Modulation of dorsal premotor cortex disrupts neuroplasticity of primary motor cortex in young and older adults

Wei-Yeh Liao[1], George M. Opie[1], Ulf Ziemann[2,3] & John G. Semmler[1]

1. Discipline of Physiology, School of Biomedicine, The University of Adelaide, Adelaide, Australia.

2. Department of Neurology & Stroke, Eberhard Karls University of Tübingen, Tübingen, Germany

3. Hertie-Institute for Clinical Brain Research, Eberhard Karls University of Tübingen, Tübingen, Germany

Running title: Premotor effects on motor cortex plasticity in young and older adults

Correspondence: Wei-Yeh Liao

School of Biomedicine

The University of Adelaide

Adelaide, South Australia 5005

Australia

Telephone: Int + 61 4 3771 9665

E-mail: wei-yeh.liao@adelaide.edu.au

Abstract

Although transcranial magnetic stimulation (TMS) research demonstrates that dorsal premotor cortex (PMd) influences neuroplasticity within primary motor cortex (M1), it is unclear how ageing modifies this communication. The present study investigated the influence of PMd on different indirect (I) wave inputs within M1 that mediate cortical plasticity in young and older adults. 15 young and 15 older participants completed two experimental sessions that examined the effects of intermittent theta burst stimulation (iTBS) to M1 when preceded by iTBS (PMd iTBS-M1 iTBS) or sham stimulation (PMd sham-M1 iTBS) to PMd. Changes in corticospinal excitability post-intervention were assessed with motor evoked potentials (MEP) recorded from right first dorsal interosseous using posterior-anterior (PA) and anterior-posterior (AP) current single-pulse TMS (PA_{1mV}; AP_{1mV}; $PA_{0.5mV}$, early I-wave; $AP_{0.5mV}$, late I-wave). Although PA_{1mV} did not change post-intervention ($P = 0.628$), PMd iTBS-M1 iTBS disrupted the expected facilitation of AP_{1mV} (to M1 iTBS) in young and older adults ($P = 0.002$). Similarly, PMd iTBS-M1 iTBS disrupted $PA_{0.5mV}$ facilitation in young and older adults ($P = 0.030$), whereas $AP_{0.5mV}$ facilitation was not affected in either group ($P = 0.218$). This suggests that while PMd specifically influences the plasticity of early I-wave circuits, this communication is preserved in older adults.

Keywords: ageing, dorsal premotor cortex, neuroplasticity, transcranial magnetic stimulation

Introduction

One of the universal effects of ageing is widespread deficits in motor function. Although these deficits occur at all levels of the motor system, the structural, functional, and biochemical changes within the brain are important (Seidler *et al.*, 2010). In particular, alterations to the ability of the brain's motor system to continuously modify its structure and function are a critical factor. Termed neuroplasticity, this process is initially mediated by changes in the strength of synaptic communication with long-term potentiation (LTP) and depression (LTD), and underpins the ability to learn new motor skills (Buonomano & Merzenich, 1998; Sanes & Donoghue, 2000). While the capacity for neuroplastic change is present across the lifespan, some studies using non-invasive brain stimulation (NIBS) show reduced plasticity in older adults (Müller-Dahlhaus *et al.*, 2008; Fathi *et al.*, 2010; Todd *et al.*, 2010; Freitas *et al.*, 2011). This reduced plasticity may contribute to the motor deficits that limit the ability of older adults to learn new motor skills that may be essential for daily life. However, the neurophysiological mechanisms underpinning these changes with advancing age remain unclear.

Transcranial magnetic stimulation (TMS) is a type of NIBS that allows investigation of specific neuronal networks within the motor system with high temporal resolution. Application of TMS over primary motor cortex (M1) produces a complex series of descending volleys within corticospinal neurons that summate at the spinal cord, resulting in a motor evoked potential (MEP) in targeted muscles (Di Lazzaro *et al.*, 1998; Rossini *et al.*, 2015). The first of these waves likely represents direct activation of corticospinal neurons, whereas subsequent waves are thought to reflect the indirect activation of interneuronal inputs to the corticospinal neurons (Di Lazzaro *et al.*, 2012; Ziemann, 2020). These responses are referred to as indirect (I) waves and are named early (I_1) or late (I_2, I_3) based on the order

of their appearance, which occurs with a periodicity of ~1.5 ms (Di Lazzaro *et al.*, 2012;

Ziemann, 2020). Early and late I-waves can be preferentially recruited by applying low-

intensity single-pulse TMS with different current directions (Sakai *et al.*, 1997; Di Lazzaro *et*

al., 2001; Ni *et al.*, 2010). For example, a posterior-anterior (PA) current (relative to the

central sulcus) preferentially recruits early I-waves, whereas an anterior-posterior (AP)

current preferentially recruits late I-waves (Sakai *et al.*, 1997; Di Lazzaro *et al.*, 2001; Ni *et*

al., 2010). Using these measures, previous work has shown that the ability to recruit late I-

waves predicts the response to plasticity-inducing TMS paradigms over M1 (Hamada *et al.*,

2013; Wiethoff *et al.*, 2014) and that the late I-waves are behaviourally relevant to the

acquisition of fine motor skills (Hamada *et al.*, 2014).

I-wave circuits are also involved in mediating the communication between other motor nodes

and M1 (Groppa *et al.*, 2012; Volz *et al.*, 2015; Spampinato *et al.*, 2020; Opie *et al.*, 2022;

Casarotto *et al.*, 2023), which form a wider network that influences M1 plasticity and

learning (Huang *et al.*, 2018; Liao *et al.*, 2022). In particular, the dorsal premotor cortex

(PMd) facilitates the planning, prediction, and correction of movements during motor

learning by updating the activity of M1 (Chouinard *et al.*, 2005; Nowak *et al.*, 2009; Parikh &

Santello, 2017). Previous studies have demonstrated that the application of repetitive TMS

(rTMS) techniques (such as theta burst stimulation; TBS) over PMd is able to modify M1

excitability, plasticity, and motor skill acquisition (Huang *et al.*, 2018; Meng *et al.*, 2020).

Furthermore, while PMd influences both early and late I-wave excitability (Liao *et al.*, 2023),

there is a stronger effect on the late I-waves (Volz *et al.*, 2015; Aberra *et al.*, 2020). Taken

together, it is likely that the influence of late I-waves on M1 plasticity reflects inputs from

PMd.

Given the role of late I-wave circuits in mediating PMd-M1 communication, changes in late

I-wave activity may affect the influence of PMd on M1 plasticity. In particular, late I-wave

activity is known to be altered with advancing age (Opie *et al.*, 2018). Age-related changes in

I-wave excitability have been investigated using the paired-pulse TMS protocol short

intracortical facilitation (SICF) (Opie *et al.*, 2018), which revealed reduced I-wave

excitability and a specific delay in the temporal characteristics of the late I-waves in older

adults (Opie *et al.*, 2018). Importantly, this delay influences NIBS-induced plasticity and is

associated with specific aspects of motor behaviour in older adults (Opie *et al.*, 2018; Opie *et*

al., 2020). In addition, it is also known that PMd-M1 effective connectivity (Ni *et al.*, 2015)

and direct PMd modulation of early I-waves within M1 is reduced in older adults (Liao *et al.*,

2023). Consequently, it is possible that the influence of PMd on M1 plasticity is altered with

advancing age, but this remains to be tested.

The purpose of the present study was, therefore, to investigate the influence of PMd on the

plasticity of early and late I-wave circuits in M1 of young and older adults. Given that

previous work has used TBS to modulate M1 plasticity in young adults (Huang *et al.*, 2018),

we applied intermittent TBS (iTBS) over PMd in young and older participants and assessed

how this influenced the neuroplastic response of M1 to iTBS. Different I-wave circuits were

assessed by varying the direction of current used to apply TMS over M1. Although we

expected iTBS over PMd to selectively modulate the plasticity of late I-wave circuits, we

hypothesised that the effect of PMd on M1 plasticity would be weaker in older adults, given

the likely alterations in late I-wave activity and PMd-M1 connectivity with advancing age.

Materials and Methods

Sample Size and Participants

15 young (mean ± standard deviation, 24.7 ± 5.0 years; range , 19-36 years) and 15 older

adults (67.2 ± 5.4 years; 61-78 years) were recruited for the study via advertisements placed

on notice boards within The University of Adelaide and the wider community, in addition to

social media platforms. Applicants for the study were excluded if they had a history of

psychiatric or neurological disease, current use of medication that affect the central nervous

system, pregnancy, metal implants, or left handedness, as assessed by a standard TMS

screening questionnaire (Rossi *et al.*, 2011). The experiment was conducted in accordance

with the Declaration of Helsinki and was approved by The University of Adelaide Human

research Ethics Committee (H-026-2008). Subjects provided written, informed consent prior

to participation.

Experimental Arrangement

All participants attended two experimental sessions where iTBS or sham iTBS was applied to

PMd, followed 30 minutes later by plasticity induction within M1 via iTBS (PMd iTBS-M1

iTBS, PMd sham-M1 iTBS). The same experimental protocol was used in both sessions (Fig.

1), with the order of intervention randomised between participants, and a washout period of at

least 1 week was used between sessions. As diurnal variations in cortisol are known to

influence the neuroplastic response to TMS (Sale *et al.*, 2008), all sessions were completed

between 11 am and 5 pm at approximately the same time of day for each participant.

During each experimental session, participants were seated in a comfortable chair with their

hands resting and relaxed. Surface electromyography (EMG) was recorded from the first

dorsal interosseous (FDI) of the right hand using two Ag-AgCl electrodes arranged in a belly-

tendon montage on the skin above the muscle, with a third electrode attached above the

styloid process of the right ulnar used to ground the electrodes. EMG signals were amplified

(300x) and filtered (band-pass 20 Hz – 1 kHz) using a CED 1902 signal conditioner

(Cambridge Electronic Design, Cambridge, UK) before being digitised at 2 kHz using a CED

1401 analogue-to-digital converter. Signal noise associated with mains power was removed

using a Humbug mains noise eliminator (Quest Scientific, North Vancouver, Canada). EMG

signals were stored on a PC for offline analysis. Real-time EMG signals were displayed on an

oscilloscope placed in front of the participant to facilitate muscle relaxation during the

experiment.

Experimental Procedures

Transcranial magnetic stimulation (TMS). A branding iron coil connected to two Magstim

200^2 magnetic stimulators (Magstim, Whitland, UK) via a BiStim unit was used to apply

TMS to left M1. The coil was held tangentially to the scalp at an angle of 45° to the sagittal

plane, inducing a PA current relative to the central sulcus. The M1 hotspot was identified as

the location producing the largest and most consistent MEPs within the relaxed FDI muscle

of the right hand (Rossini *et al.*, 2015). This location was marked on the scalp for reference

and continuously monitored throughout each experimental session. All baseline, post-PMd

iTBS, and post-M1 iTBS (5 minutes, 30 minutes) TMS was applied at a rate of 0.2 Hz, with a

10% jitter between trials to avoid anticipation of the stimulus.

Resting motor threshold (RMT) was recorded as the lowest stimulus intensity producing an

MEP amplitude ≥ 50 µV in at least 5 out of 10 trials during relaxation of the right FDI. RMT

was assessed at the beginning of each experimental session and expressed as a percentage of

maximum stimulator output (% MSO) (Rossini *et al.*, 2015). Active motor threshold (AMT)

was then assessed, defined as the lowest % MSO producing an MEP amplitude ≥ 200 µV in

at least 5 out of 10 trials during concurrent low-level activation (~10% voluntary activation)

of the right FDI (Hamada *et al.*, 2013). These measures were then repeated using the AP

current by rotating the coil 180°. Then, the stimulus intensities producing a standard MEP

amplitude approximating 1 mV (MEP_{1mV}; PA_{1mV}, AP_{1mV}), in addition to an MEP amplitude

approximating 0.5 mV ($MEP_{0.5mV}$; $PA_{0.5mV}$, $AP_{0.5mV}$), when averaged over 20 trials, were

identified. The same intensities (MEP_{1mV}, $MEP_{0.5mV}$) were then applied following PMd iTBS

and following M1 iTBS to assess changes in corticospinal excitability.

I-wave recruitment. To investigate the ability to recruit I-waves, the onset latencies of PA

(early) and AP (late) MEPs were assessed relative to the MEP onset generated by direct

activation of corticospinal neurons using a lateral-to-medial (LM) current (Hamada *et al.*,

2013). A block of 15 MEP trials in the active FDI was recorded for 110% of AMT_{PA} and

AMT_{AP}, in addition to 150% AMT_{LM} (Hamada *et al.*, 2013). If 150% AMT_{LM} exceeded 100%

MSO, 100% MSO was used, or if 150% AMT_{LM} was below 50% MSO, 50% MSO was used

(Hamada *et al.*, 2013). The difference in mean onset latencies between PA and LM (PA-LM)

and AP and LM (AP-LM) were calculated as measures of early and late I-wave recruitment

efficiency, respectively (Hamada *et al.*, 2013). In an attempt to reduce the confounding

influence of muscle contraction on neuroplasticity induction (Huang *et al.*, 2008;

Thirugnanasambandam *et al.*, 2011; Goldsworthy *et al.*, 2015), these measures were recorded

at the start and at the end of the experimental session, at least 45 minutes apart from the

plasticity induction of PMd and M1.

Theta burst stimulation (TBS). Intermittent theta burst stimulation (iTBS) was delivered over

left PMd and left M1 using a Magstim Super-rapid stimulator (Magstim, Whitland, UK),

connected to an air-cooled figure-of-eight coil. The coil was held tangentially to the scalp, at

an angle of 45° to the sagittal plane, with the handle pointing backwards and laterally,

inducing a biphasic pulse with an initial PA current followed by an AP return current (Suppa

et al., 2008). In accordance with existing literature, iTBS consisted of bursts of three pulses

given at a frequency of 50 Hz. Each burst was repeated at 5 Hz for 2 s, and repeated every 8 s

for 20 cycles, totalling 600 pulses (Huang *et al.*, 2005; Huang *et al.*, 2008; Huang *et al.*,

2018; Meng *et al.*, 2020). The location of left PMd was defined as 8% of the distance

between the nasion and inion (approximately 2.5 – 3 cm) anterior to the M1 hotspot,

consistent with previous work (Münchau *et al.*, 2002; Koch *et al.*, 2007; Huang *et al.*, 2018;

Meng *et al.*, 2020). The location of both the M1 hotspot and left PMd site were logged

relative to the MNI-ICBM152 template using Brainsight neuronavigation (Rogue Research,

Montreal, Quebec, Canada). These locations were then used to guide the assessment of RMT

(RMT$_{Rapid}$) over M1 with the Magstim Super-rapid stimulator, in addition to the application

of iTBS over left PMd and M1 at 70% RMT$_{Rapid}$.

Sham iTBS to left PMd was delivered using a sham figure-of-eight coil (replicating the coil

click), with a bar electrode connected to a constant current stimulator (Digitimer,

Hertfordshire, UK) placed underneath the coil delivering electrical stimulation (1.5 mA) to

the scalp in order to mimic the pulse sensation. Following either intervention, participants

provided answers to a visual analogue scale (VAS) questionnaire indexing the degree of

discomfort, muscle activation, and localisation of scalp sensation during PMd iTBS.

Data Analysis

Visual inspection of EMG data was completed offline, with any trials obtained from the

resting muscle having EMG activity exceeding 25 µV in the 100 ms prior to stimulus

application excluded from analysis (approximately 6.8% removed). The amplitude of MEPs

obtained from resting muscle recordings was measured peak-to-peak and expressed in mV.

The MEP onset latencies obtained from active muscle recordings was assessed with a semi-

automated process using a custom script within the Signal program (v 6.02, Cambridge

Electronic Design) and expressed in ms. MEP latency was recorded as the period from

stimulus application to the resumption of voluntary EMG activity. This was defined as the

point at which post-stimulus EMG amplitude exceeded the mean EMG amplitude recorded

within the 100 ms pre-stimulus, plus 2 standard deviations. MEP onset latencies were

averaged over individual trials within each subject and coil orientation. Within each

participant, the mean LM MEP latencies were subtracted from the mean PA and AP MEP

latencies to determine PA-LM and AP-LM MEP latency differences. Following TBS

interventions, changes in MEP latency differences were quantified by expressing the post-

intervention responses as a percentage of the baseline responses. Changes in MEP amplitude

due to PMd iTBS were quantified by expressing post-PMd iTBS responses as a percentage of

baseline MEP amplitude. For post-M1 iTBS, changes in MEP amplitude were quantified by

expressing post-M1 iTBS responses as a percentage of post-PMd iTBS responses.

Statistical Analysis

Visual inspection and Kolmogorov-Smirnov tests of the data residuals revealed non-normal,

positively-skewed distributions for all TMS data. Consequently, generalised linear mixed

models (GLMM's), which can account for non-normal distributions (Lo & Andrews, 2015;

Puri & Hinder, 2022), were used to perform all statistical analyses. Each model assessing

MEP amplitude included single trial data with repeated measures and was fitted with Gamma

distributions (Puri & Hinder, 2022), with all random subject effects included (intercepts and

slopes) (Barr *et al.*, 2013). Identity link functions were used for baseline MEP amplitude and

latency differences while log link functions were used for post-iTBS normalised MEP

amplitude and latency differences (Lo & Andrews, 2015; Puri & Hinder, 2022). To optimise

model fit, we tested different covariance structures and the structure providing the best fit

(assess with the Bayesian Schwartz Criterion; BIC) within a model that was able to converge

was used in the final model. Two-factor GLMMs were used to compare effects of session

(PMd iTBS-M1 iTBS, PMd sham-M1 iTBS) and age (young, older) at baseline in eight

separate models for $PA_{0.5mV}$, $AP_{0.5mV}$, PA_{1mV}, and AP_{1mV} stimulation intensities and MEP

amplitude. A three-factor model was used to compare the effects of session, age, and

orientation (PA, AP) on PA-LM and AP-LM latency differences at baseline.

Changes in corticospinal excitability following PMd iTBS were investigated by assessing

effects of session and age in four separate models for baseline-normalised PA_{1mV}, AP_{1mV},

$PA_{0.5mV}$, and $AP_{0.5mV}$ MEP amplitude. Changes in corticospinal excitability following PMd

iTBS-M1 iTBS and PMd sham-M1 iTBS were investigated by assessing effects of session,

time (5 minutes, 30 minutes) and age in four separate models for PA_{1mV}, AP_{1mV}, $PA_{0.5mV}$, and

$AP_{0.5mV}$ MEP amplitude normalised to the mean post-PMd iTBS MEP amplitude. As AP_{1mV}

baseline stimulation intensities varied between sessions (see Table 1), AP_{1mV} stimulation

intensities were also included in the model as a covariate to assess if varying stimulation

intensities confounded changes in post-intervention AP_{1mV} MEP amplitude. Changes in I-

wave recruitment following the intervention were investigated by assessing effects of session,

age, and coil orientation on baseline-normalised average PA-LM and AP-LM latency

differences. For all models, investigation of main effects and interactions were performed

using custom contrasts with the Bonferroni correction, and significance was set at $P < 0.05$.

Data for all models are presented as estimated marginal means (EMMs) and 95% confidence

intervals (95% CI), whereas pairwise comparisons are presented as the estimated mean

difference (EMD) and 95% CI for the estimate.

Furthermore, we used Spearman's rank order correlation analysis to assess the relationship

between different variables. Specifically, baseline MEP latency differences were correlated

with changes in corticospinal excitability immediately following PMd iTBS to investigate if

the ability to recruit I-waves is related to changes in corticospinal excitability. Baseline MEP

latency differences were also correlated with changes in corticospinal excitability during the

PMd sham-M1 iTBS session to investigate if the ability to recruit I-waves is related to

changes in corticospinal excitability following M1 iTBS. In addition, changes in corticospinal

excitability following PMd iTBS were also correlated with changes in corticospinal

excitability following M1 iTBS (during PMd iTBS-M1 iTBS) to investigate if direct PMd

modulation of M1 excitability is related to changes in M1 plasticity. Correlations are

presented as Spearman's ρ with false discovery rate-adjusted *P*-value of 0.05 following the

Benjamini-Hochberg procedure. Lastly, differences in the perception of discomfort, extent of

FDI activation, and localisation of stimulus during PMd iTBS and PMd sham were

investigated by comparing VAS responses using paired t-tests with Bonferroni correction (*P*

< 0.0167), with data presented as mean ± standard deviation.

Results

All participants completed both experimental sessions without adverse reactions. We were

unable to record PA_{1mV} in one older male participant, $AP_{0.5mV}$ in two older participants (1

female, 1 male), and AP_{1mV} in five participants (1 young female; 3 older females, 1 older

male) due to high thresholds of activation (mean RMT_{PA} = 80.0% MSO, mean RMT_{AP} =

73.0% MSO). Baseline stimulation intensities are presented in Table 1. Stimulation

intensities for AP_{1mV} differed between sessions ($F_{1,46}$ = 4.17, *P* = 0.047), with *post-hoc*

comparisons showing higher intensities for the iTBS session relative to sham session (EMD =

2.3% MSO [0.0, 4.6], *P* = 0.047). There were no other main effects or interactions for all

other baseline stimulation intensities (all *P* > 0.05).

Baseline MEP amplitude for corticospinal excitability and MEP latency differences are

shown in Table 2. For PA_{1mV} MEP amplitude, there was an interaction between session and

age ($F_{1,1121}$ = 4.194, *P* = 0.041), with *post-hoc* comparisons revealing larger MEP amplitude

for young participants relative to older participants (EMD = 0.14 mV [0.02, 0.26], *P* =

0.024). For baseline MEP latency differences, responses differed between coil orientations

($F_{1,112} = 165.20$, $P < 0.0001$), where PA-LM latencies were shorter than AP-LM latencies

(EMD = 1.95 ms [1.65, 2.25], $P < 0.0001$), as expected. There were no main effects or

interactions for all other baseline MEP amplitude or MEP latency differences (all $P > 0.05$).

Absolute MEP amplitude between sessions for young and older adults is presented in

Supplementary Materials.

Changes in corticospinal excitability following PMd iTBS

The participants' perceptions of PMd iTBS and PMd sham are shown in Table 3. While there

were no differences between sessions in the extent of discomfort ($t_{29} = 0.25$, $P = 0.804$) or

FDI activation ($t_{29} = 0.10$, $P = 0.918$) experienced by the participants, the locality of

stimulation differed ($t_{29} = 3.98$, $P = 0.004$), with the sensation of iTBS perceived as more

widespread relative to electrical scalp stimulation in sham.

Changes in MEP_{1mV} and $MEP_{0.5mV}$ measures of corticospinal excitability following PMd

iTBS are shown in Figure 2. PA_{1mV} MEP amplitude did not differ between sessions ($F_{1,1114} =$

0.90, $P = 0.343$; Fig. 2A) or age groups ($F_{1,1114} = 0.12$, $P = 0.726$), and there was no

interaction between factors ($F_{1,1114} = 2.41$, $P = 0.121$). AP_{1mV} MEP amplitude did not vary

between sessions ($F_{1,996} = 2.33$, $P = 0.127$; Fig. 2B) or age groups ($F_{1,996} = 1.31$, $P = 0.252$),

and there was no interaction between factors ($F_{1,996} = 0.51$, $P = 0.476$). In contrast, while

$PA_{0.5mV}$ MEP amplitude did not differ between age groups ($F_{1,1152} = 0.11$, $P = 0.740$),

responses varied between sessions ($F_{1,1152} = 4.23$, $P = 0.040$; Fig. 2C), with increased MEP

amplitude following PMd iTBS relative to sham (EMD = 26.3% [0.7, 51.9], $P = 0.044$).

There was no interaction between factors ($F_{1,1152} = 0.11$, $P = 0.741$). $AP_{0.5mV}$ MEP amplitude

did not vary between sessions ($F_{1,1073} = 1.04$, $P = 0.308$; Fig. 2D) or age groups ($F_{1,1073} =$

2.80, $P = 0.095$), and there was no interaction between factors ($F_{1,1073} = 1.03$, $P = 0.310$).

Changes in corticospinal excitability and I-wave recruitment following M1 iTBS.

Changes in MEP_{1mV} measures of corticospinal excitability following PMd iTBS-M1 iTBS

and PMd sham-M1 iTBS are presented in Figure 3. PA_{1mV} MEP amplitude (Fig. 3A) did not

vary between sessions ($F_{1,2234} = 2.20$, $P = 0.138$), time points ($F_{1,2234} = 0.15$, $P = 0.696$), or

age groups ($F_{1,2234} = 1.17$, $P = 0.279$), and there were no interactions between factors (all $P >$

0.05). AP_{1mV} MEP amplitude also did not differ between sessions ($F_{1,1921} = 0.98$, $P = 0.323$),

time points ($F_{1,1921} = 1.14$, $P = 0.286$), or age groups ($F_{1,1921} = 1.21$, $P = 0.272$), but there was

an interaction between session and time ($F_{1,1921} = 9.94$, $P = 0.002$; Fig 3B). *Post-hoc*

comparisons showed that MEP amplitude following PMd sham-M1 iTBS was increased at 5

minutes compared to PMd iTBS-M1 iTBS (EMD = 29.7% [6.6, 52.8], $P = 0.012$), and

compared to 30 minutes (EMD = 30.3% [7.3, 53.3], $P = 0.010$). There were no other

interactions (all $P > 0.05$). In addition, there was no effect of stimulation intensity on MEP

amplitude ($F_{1,1921} = 0.40$, $P = 0.527$).

Changes in $MEP_{0.5mV}$ measures of corticospinal excitability are presented in Figure 4. While

$PA_{0.5mV}$ MEP amplitude did not differ between time points ($F_{1,2311} = 0.03$, $P = 0.874$) or age

groups ($F_{1,2311} = 0.17$, $P = 0.678$), responses varied between sessions ($F_{1,2311} = 17.4$, $P <$

0.05), with increased MEP amplitude following PMd sham-M1 iTBS (EMD = 34.3% [17.5,

51.0], $P < 0.05$). Furthermore, there was an interaction between session, time, and age ($F_{1,2311}$

$= 4.71$, $P = 0.030$; Fig. 4A). *Post-hoc* analysis revealed increased MEP amplitude following

PMd sham-M1 iTBS compared to PMd iTBS-M1 iTBS for young adults at 30 minutes (EMD

$= 50.7\%$ [20.1, 81.3], $P = 0.001$), while this effect was observed for older adults at 5 (EMD =

43.0% [13.9, 72.0], $P = 0.004$) and 30 minutes (EMD = 32.0% [3.6, 60.4], $P = 0.027$). For

$AP_{0.5mV}$, MEP amplitude did not vary between sessions ($F_{1,2141} = 0.13$, $P = 0.723$) or age

groups ($F_{1,2141} = 3.12$, $P = 0.077$) (Fig. 4B). However, responses differed between time points

($F_{1,2141}$ = 5.91, P = 0.015; Fig. 4C), with *post-hoc* analysis revealing that MEP amplitude was

increased at 5 minutes relative to 30 minutes post-M1 iTBS (EMD = 22.0% [3.9, 40.0], P =

0.017). There were no interactions between factors (all P > 0.05).

I-wave recruitment

There was no difference between sessions ($F_{1,112}$ = 0.72, P = 0.399), coil orientations ($F_{1,112}$ =

0.09, P = 0.766), or age groups ($F_{1,112}$ = 0.38, P = 0.538), and there were no interactions

between factors (all P > 0.05).

Correlation analyses

Baseline PA-LM and AP-LM latencies were not related to changes in single-pulse measures

of corticospinal excitability (PA_{1mV}, AP_{1mV}, $PA_{0.5mV}$, $AP_{0.5mV}$) following PMd iTBS (all P >

0.05). Baseline PA-LM and AP-LM latencies were not related to changes in single-pulse

measures of corticospinal excitability following PMd sham-M1 iTBS (all P > 0.05). In

contrast, while changes in AP_{1mV} MEP amplitude following PMd iTBS were not related to

changes in AP_{1mV} responses following M1 iTBS (ρ = -0.361, P = 0.076; Fig. 5B), changes in

PA_{1mV}, $PA_{0.5mV}$, and $AP_{0.5mV}$ MEP amplitude following PMd iTBS were negatively correlated

with changes in PA_{1mV} (ρ = 0.-577, P = 0.001; Fig. 5A), $PA_{0.5mV}$ (ρ = -0.616, P = 0.0003; Fig.

5C), and $AP_{0.5mV}$ (ρ = -0.551, P = 0.002; Fig. 5D) responses following M1 iTBS,

respectively.

Discussion

In the present study, we investigated the influence of PMd on the plasticity of early and late I-

wave-generating circuits in M1 of young and older adults. This was achieved by applying

PMd iTBS as a priming intervention to modify the neuroplastic response of M1 to subsequent

iTBS (PMd iTBS-M1 iTBS, PMd sham-M1 iTBS). We measured changes in corticospinal

excitability (PA_{1mV}, AP_{1mV}, $PA_{0.5mV}$, $AP_{0.5mV}$) and I-wave recruitment (PA-LM latency, AP-

LM latency) following the intervention. The findings show that PMd iTBS specifically

modulated the excitability of the early I-wave circuits in both young and older adults.

Moreover, PMd iTBS disrupted the neuroplastic response of the early I-wave circuits to M1

iTBS in both young and older adults, whereas the neuroplastic response of the late I-wave

circuits was unaffected in both age groups.

PMd influence on corticospinal excitability in young and older adults

Previous work has reported that application of iTBS to PMd facilitates PA_{1mV} measures of

M1 corticospinal excitability in young adults by ~30%, which is thought to stem from the

induction of LTP-like effects within PMd, resulting in increased excitability within M1

(Meng *et al.*, 2020). Furthermore, we have demonstrated previously that this effect on PA_{1mV}

is preserved with ageing and extends to AP_{1mV} measures of corticospinal excitability (Liao *et*

al., 2023). The absence of any changes in PA_{1mV} or AP_{1mV} within the present study is

therefore inconsistent with these previous findings. However, inter- and intraindividual

variability in the changes in M1 excitability following TBS is well-documented (Hamada *et*

al., 2013; Corp *et al.*, 2020; Guerra *et al.*, 2020). In particular, there is some variability in the

time course of facilitation following PMd iTBS. For example, one study reported that the

facilitation of MEP amplitude only occurred at 15 minutes (Meng *et al.*, 2020), whereas we

previously demonstrated facilitation of MEP amplitude that persisted from 5 to 40 minutes

following PMd iTBS (Liao *et al.*, 2023). Consequently, our decision to record MEPs at 5

minutes post-PMd iTBS may have limited the ability to detect changes in corticospinal

excitability due to the priming intervention.

Although PA_{1mV} and AP_{1mV} MEP amplitude were not modulated following PMd iTBS,

$PA_{0.5mV}$ was facilitated (by ~30%) for both young and older adults. The conventional

interpretation of how TMS intensity and current direction influence I-wave recruitment

suggests that low-intensity PA TMS preferentially recruits early I-waves, whereas low-

intensity AP TMS preferentially recruits late I-waves (Hamada *et al.*, 2013), with either

current direction able to recruit both I-waves as the stimulation intensity is increased (Di

Lazzaro *et al.*, 2001; Di Lazzaro *et al.*, 2003). We therefore applied single-pulse TMS at

relatively lower intensities compared to MEP_{1mV} ($PA_{0.5mV}$, $AP_{0.5mV}$), where $PA_{0.5mV}$ is likely

more selective for activation of the early I-waves, while $AP_{0.5mV}$ is likely more selective for

the late I-waves (Opie *et al.*, 2022). Given that we previously reported potentiation of both

$PA_{0.5mV}$ and $AP_{0.5mV}$ (by ~50-100%) following PMd iTBS (Liao *et al.*, 2023), the increase in

$PA_{0.5mV}$ within the present study suggests that the effect of PMd iTBS on early I-wave

excitability may be immediate and more consistent. Importantly, previous work has shown

that PMd iTBS applied as it was in the current study is unlikely to have activated M1 directly.

Specifically, Huang and colleagues (2009) assessed the intensity required to activate M1

when TMS was applied over PMd, and showed that 80% of this (matching the level applied

during iTBS) applied to M1 does not influence M1 excitability (Huang *et al.*, 2009). Given

that we located PMd using similar methods (Huang *et al.*, 2009; Huang *et al.*, 2018; Meng *et*

al., 2020), it is therefore unlikely that PMd iTBS activated M1 directly in the present study.

Despite the present findings demonstrating that PMd iTBS increased early I-wave

excitability, this effect was not different between young and older adults, suggesting that the

influence of PMd on early I-wave excitability may be preserved with ageing. This contrasts

with our previous work, which specifically demonstrated weakened direct PMd modulation

of early I-waves in older adults (Liao *et al.*, 2023). Given that both studies employed the

same methods to assess changes in M1 excitability following PMd iTBS, participant factors

such as genetics, pharmacology, aerobic exercise, and diet that are known to influence

cortical plasticity (Ridding & Ziemann, 2010; Phillips, 2017) may have confounded the

present findings. As the contributions of participant characteristics on PMd-M1

communication were not examined in the present study, and the small sample sizes were not

powered for such subanalyses, it will be important to characterise their involvement in future

studies.

PMd influence on M1 plasticity in young and older adults

Previous work in young participants demonstrated that applying continuous TBS (cTBS) to

PMd disrupts the neuroplastic response of M1 to both iTBS and cTBS, assessed using PA_{1mV}

MEPs (Huang *et al.*, 2018). This demonstrated that LTP- and LTD-like effects within M1 can

be modulated by PMd cTBS, which was thought to arise from heterosynaptic metaplastic

effects, where the modulation of local synaptic plasticity within PMd affected subsequent

changes in remote synapses (that were not initially activated) within M1 (Huang *et al.*, 2018).

In the present study, we demonstrated that applying iTBS to PMd also disrupts the LTP-like

effects of M1 iTBS for AP_{1mV} measures of corticospinal excitability. However, given that

iTBS produces LTP-like effects while cTBS produces LTD-like effects, this disruption of

AP_{1mV} facilitation may stem from a different mechanism more consistent with homeostatic

metaplasticity (Müller *et al.*, 2007; Todd *et al.*, 2009; Murakami *et al.*, 2012). Importantly,

this response did not differ between young and older adults, suggesting that the influence of

PMd on the plasticity of AP circuits within M1 is maintained with age. In addition, this

difference is also unlikely to be driven by AP_{1mV} stimulation intensity differences at baseline,

as investigation of this confounding factor did not reveal any effects on post-intervention

AP_{1mV} MEP amplitude.

Furthermore, PMd iTBS disrupted the effects of M1 iTBS on $PA_{0.5mV}$ (early), but not $AP_{0.5mV}$

(late) circuits. This suggests that the influence of PMd on M1 plasticity is specific to the early

I-waves, which is unexpected given that previous work has demonstrated a stronger influence

of PMd on late I-wave circuits (Volz *et al.*, 2015; Aberra *et al.*, 2020; Liao *et al.*, 2023). One

possible explanation is that shorter AP MEP onset latencies (more consistent with early I-

wave recruitment) have been reported to predict stronger premotor-M1 functional

connectivity (Volz *et al.*, 2015) and the nature of this communication may contribute to the

influence of PMd on M1 plasticity. This appears consistent with our AP_{1mV} findings, which

may have occurred as the higher stimulus intensity required to record AP_{1mV} resulted in

mixed recruitment of early and late I-waves, but that changes in AP_{1mV} were driven

specifically by the early I-waves (Di Lazzaro *et al.*, 2001; Di Lazzaro *et al.*, 2003; Liao *et al.*,

2022). This is further complemented by the correlation analysis results demonstrating that

larger facilitation of $PA_{0.5mV}$ post-PMd iTBS is correlated with smaller facilitation of $PA_{0.5mV}$

post-M1 iTBS, suggesting that this homeostatic metaplastic effect is likely related to the early

I-wave circuits. While a similar correlation was also shown for PA_{1mV} and $AP_{0.5mV}$, PMd

iTBS-M1 iTBS did not disrupt the potentiation of these measures when compared to PMd

sham-M1 iTBS session. It is possible that the higher stimulus intensities required for PA_{1mV}

and $AP_{0.5mV}$ (relative to $PA_{0.5mV}$) may have also resulted in mixed recruitment of early and

late I-waves (Liao *et al.*, 2022). In particular, given that there is growing evidence to suggest

that PA and AP TMS can activate distinct populations of early and late I-waves (i.e., PA- and

AP-sensitive early and late I-waves) (Spampinato *et al.*, 2020; Opie & Semmler, 2021),

PA_{1mV} and $AP_{0.5mV}$ may have recruited other I-wave circuits that were less sensitive to the

modulatory effects of iTBS. However, this will need to be clarified in future research using

techniques that are more selective to these different I-waves, such as modifying the TMS

pulse width (Hannah & Rothwell, 2017). Despite this, we provide new evidence that PMd

iTBS specifically modulates M1 plasticity of early I-wave circuits recruited by AP

stimulation.

While M1 iTBS in isolation (PMd sham-M1 iTBS) potentiated $PA_{0.5mV}$ responses (compared

with PMd iTBS-M1 iTBS) in both age groups, the timing of this response varied between

groups. Whereas differences between sham and real PMd iTBS sessions were immediate for

older adults, they were only apparent after 30 minutes in young adults. Given that M1 iTBS

has not been shown to differentially modulate corticospinal excitability in young and older

adults (Di Lazzaro *et al.*, 2008; Young-Bernier *et al.*, 2014; Dickins *et al.*, 2015; Opie *et al.*,

2017), this outcome seems unlikely to reflect effects of age within M1. An alternative

explanation could be that the modulatory effects of PMd iTBS differed between groups, with

younger adults having a stronger response that was more resistant to the subsequent effects of

M1 iTBS. This is supported by the amplitude of $PA_{0.5mV}$ being reduced 5 minutes after M1

iTBS in older, but not young adults in the session involving real PMd iTBS (Fig. 4A).

Although speculative, this outcome would be consistent with our previous finding that the

influence of PMd iTBS on $PA_{0.5mV}$ is reduced in older adults (Liao *et al.*, 2023). However,

this speculation will require additional studies that more effectively characterise the time

course of facilitation in young and older adults. For example, previous work investigating the

effects of PMd cTBS on M1 neuroplastic response to iTBS or cTBS monitored changes in

corticospinal excitability for two hours following PMd cTBS (during which excitability

returned to baseline levels) before applying subsequent M1 iTBS or cTBS (Huang *et al.*,

2018).

PMd and M1 influence on I-wave recruitment in young and older adults

The ability to recruit both early and late I-waves can be investigated by comparing the

latencies evoked by PA and AP TMS to the latencies of direct corticospinal activation (PA-

LM, early; AP-LM, late) (Hamada *et al.*, 2013). The prototypical values for these measures

reveal shorter PA-LM latencies (~1.5 ms) compared to AP-LM latencies (~3 ms), providing

an index of early and late I-wave recruitment, respectively (Hamada *et al.*, 2013).

Importantly, previous studies have shown that the ability to recruit late I-waves with AP TMS

predicts the neuroplastic response of M1 to iTBS (Hamada *et al.*, 2013; Volz *et al.*, 2019),

with AP inputs thought to originate from PMd (Volz *et al.*, 2015; Aberra *et al.*, 2020). It has

also been demonstrated that AP-LM latencies can be shortened using M1 iTBS, which was

suggested to reflect the direct modulation of the late I-wave circuitry (Volz *et al.*, 2019).

Although we also assessed changes in PA-LM and AP-LM latencies following PMd sham-

M1 iTBS in the present study, the intervention failed to modulate the I-wave latencies. It is

possible that changes in AP-LM latencies occur immediately following iTBS, as the MEP

latency measures were recorded at least 45 minutes either side of PMd and M1 iTBS in order

to avoid complications involving the effects of muscle activation on neuroplasticity responses

(Huang *et al.*, 2008; Thirugnanasambandam *et al.*, 2011; Goldsworthy *et al.*, 2015).

Consequently, the effects of M1 iTBS on I-wave latencies will have to be clarified in future

studies.

Importantly, baseline I-wave recruitment was not correlated with changes in corticospinal

excitability following M1 iTBS in isolation, in contrast to previous findings (Hamada *et al.*,

2013; Volz *et al.*, 2019). While the difference between the present study and previous studies

is that we included older participants, no differences between age groups were shown for I-

wave recruitment or corticospinal excitability in the present study. The variability in the

present findings may therefore involve contributions from other factors. For example, recent

work assessing variability of M1 iTBS has suggested that the ability of iTBS to engage neural

oscillations in the β range (13-30 Hz) may be an important predictor of the neuroplastic

response to iTBS (Leodori *et al.*, 2021). Enhancing premotor-M1 communication using

cortico-cortical paired associated stimulation (ccPAS) has been recently shown to improve

the synchronisation of neural oscillations (which is thought to mediate neuronal

communication and plasticity) in the β range (Trajkovic *et al.*, 2023). Further investigation

involving these measures may therefore better characterise the variability of iTBS, and may

also have applications in understanding PMd-M1 communication.

In conclusion, the application of iTBS over PMd potentiated corticospinal excitability and

disrupted the effects of subsequent M1 iTBS. Specifically, our results show that PMd may

more consistently influence the excitability of early I-waves in young and older adults.

Importantly, we provide new evidence that PMd disrupts M1 plasticity of early I-wave

circuits in both age groups. It will therefore be useful in future studies to investigate how

PMd modulation of M1 plasticity influences different feature of motor skill learning in young

and older adults.

Funding

490 Support was provided by an Australian Research Council Discovery Projects Grant (grant

491 number DP200101009). GMO was supported by funding from the National Health and

492 Medical Research Council (APP1139723) and Australian Research Council (DE230100022).

References

Aberra AS, Wang B, Grill WM & Peterchev AV. (2020). Simulation of transcranial magnetic

stimulation in head model with morphologically-realistic cortical neurons. *Brain Stimul* **13,**

175-189.

Barr DJ, Levy R, Scheepers C & Tily HJ. (2013). Random effects structure for confirmatory hypothesis

testing: Keep it maximal. *J Mem Lang* **68,** 255-278.

Buonomano DV & Merzenich MM. (1998). Cortical plasticity: from synapses to maps. *Annu Rev*

Neurosci **21,** 149-186.

Casarotto A, Dolfini E, Cardellicchio P, Fadiga L, D'Ausilio A & Koch G. (2023). Mechanisms of

Hebbian-like plasticity in the ventral premotor – primary motor network. *J Physiol* **601,** 211-

226.

Chouinard PA, Leonard G & Paus T. (2005). Role of the primary motor and dorsal premotor cortices

in the anticipation of forces during object lifting. *J Neurosci* **25,** 2277-2284.

Corp DT, Bereznicki HGK, Clark GM, Youssef GJ, Fried PJ, Jannati A, Davies CB, Gomes-Osman J,

Stamm J, Chung SW, Bowe SJ, Rogasch NC, Fitzgerald PB, Koch G, Di Lazzaro V, Pascual-

Leone A & Enticott PG. (2020). Large-scale analysis of interindividual variability in theta-burst

stimulation data: Results from the 'Big TMS Data Collaboration'. *Brain Stimul* **13,** 1476-1488.

Di Lazzaro V, Oliviero A, Pilato F, Mazzone P, Insola A, Ranieri F & Tonali PA. (2003). Corticospinal

volleys evoked by transcranial stimulation of the brain in conscious humans. *Neurol Res* **25,**

143-150.

Di Lazzaro V, Oliviero A, Saturno E, Pilato F, Insola A, Mazzone P, Profice P, Tonali P & Rothwell J.

(2001). The effect on corticospinal volleys of reversing the direction of current induced in the

motor cortex by transcranial magnetic stimulation. *Exp Brain Res* **138,** 268-273.

Di Lazzaro V, Pilato F, Dileone M, Profice P, Oliviero A, Mazzone P, Insola A, Ranieri F, Meglio M,

Tonali PA & Rothwell JC. (2008). The physiological basis of the effects of intermittent theta

burst stimulation of the human motor cortex. *J Physiol* **586,** 3871-3879.

Di Lazzaro V, Profice P, Ranieri F, Capone F, Dileone M, Oliviero A & Pilato F. (2012). I-wave origin

and modulation. *Brain Stimul* **5,** 512-525.

Di Lazzaro V, Restuccia D, Oliviero A, Profice P, Ferrara L, Insola A, Mazzone P, Tonali P & Rothwell JC.

(1998). Effects of voluntary contraction on descending volleys evoked by transcranial

stimulation in conscious humans. *J Physiol* **508 (Pt 2),** 625-633.

Dickins DS, Sale MV & Kamke MR. (2015). Plasticity Induced by Intermittent Theta Burst Stimulation

in Bilateral Motor Cortices Is Not Altered in Older Adults. *Neural plasticity* **2015,** 323409.

Fathi D, Ueki Y, Mima T, Koganemaru S, Nagamine T, Tawfik A & Fukuyama H. (2010). Effects of aging

on the human motor cortical plasticity studied by paired associative stimulation. *Clin*

Neurophysiol **121,** 90-93.

Freitas C, Perez J, Knobel M, Tormos JM, Oberman L, Eldaief M, Bashir S, Vernet M, Peña-Gómez C &

Pascual-Leone A. (2011). Changes in cortical plasticity across the lifespan. *Front Aging*

Neurosci **3,** 5-5.

Goldsworthy MR, Müller-Dahlhaus F, Ridding MC & Ziemann U. (2015). Resistant Against De-

depression: LTD-Like Plasticity in the Human Motor Cortex Induced by Spaced cTBS. *Cereb*

Cortex **25,** 1724-1734.

Groppa S, Schlaak BH, Münchau A, Werner-Petroll N, Dünnweber J, Bäumer T, van Nuenen BFL &

Siebner HR. (2012). The human dorsal premotor cortex facilitates the excitability of

ipsilateral primary motor cortex via a short latency cortico-cortical route. *Hum Brain Mapp*

33, 419-430.

Guerra A, López-Alonso V, Cheeran B & Suppa A. (2020). Variability in non-invasive brain stimulation

studies: Reasons and results. *Neurosci Lett* **719,** 133330.

Hamada M, Galea JM, Di Lazzaro V, Mazzone P, Ziemann U & Rothwell JC. (2014). Two distinct

interneuron circuits in human motor cortex are linked to different subsets of physiological

and behavioral plasticity. *J Neurosci* **34,** 12837-12849.

Hamada M, Murase N, Hasan A, Balaratnam M & Rothwell JC. (2013). The Role of Interneuron

Networks in Driving Human Motor Cortical Plasticity. *Cereb Cortex* **23,** 1593-1605.

Hannah R & Rothwell JC. (2017). Pulse Duration as Well as Current Direction Determines the

Specificity of Transcranial Magnetic Stimulation of Motor Cortex during Contraction. *Brain*

Stimul **10,** 106-115.

Huang Y-Z, Chen R-S, Fong P-Y, Rothwell JC, Chuang W-L, Weng Y-H, Lin W-Y & Lu C-S. (2018). Inter-

cortical modulation from premotor to motor plasticity. *J Physiol* **596,** 4207-4217.

Huang Y-Z, Edwards MJ, Rounis E, Bhatia KP & Rothwell JC. (2005). Theta Burst Stimulation of the

Human Motor Cortex. *Neuron* **45,** 201-206.

Huang Y-Z, Rothwell JC, Edwards MJ & Chen R-S. (2008). Effect of Physiological Activity on an NMDA-

Dependent Form of Cortical Plasticity in Human. *Cereb Cortex* **18,** 563-570.

Huang Y-Z, Rothwell JC, Lu C-S, Wang J, Weng Y-H, Lai S-C, Chuang W-L, Hung J & Chen R-S. (2009).

The effect of continuous theta burst stimulation over premotor cortex on circuits in primary

motor cortex and spinal cord. *Clin Neurophysiol* **120,** 796-801.

Koch G, Franca M, Mochizuki H, Marconi B, Caltagirone C & Rothwell JC. (2007). Interactions

between pairs of transcranial magnetic stimuli over the human left dorsal premotor cortex

differ from those seen in primary motor cortex. *J Physiol* **578,** 551-562.

Leodori G, Fabbrini A, De Bartolo MI, Costanzo M, Asci F, Palma V, Belvisi D, Conte A & Berardelli A.

(2021). Cortical mechanisms underlying variability in intermittent theta-burst stimulation-

induced plasticity: A TMS-EEG study. *Clin Neurophysiol* **132,** 2519-2531.

Liao W-Y, Opie GM, Ziemann U & Semmler JG. (2023). Modulation of dorsal premotor cortex

differentially influences I-wave excitability in primary motor cortex of young and older

adults. *J Physiol.*

Liao W-Y, Sasaki R, Semmler JG & Opie GM. (2022). Cerebellar transcranial direct current stimulation

disrupts neuroplasticity of intracortical motor circuits. *PLoS One* **17,** e0271311.

Lo S & Andrews S. (2015). To transform or not to transform: using generalized linear mixed models

to analyse reaction time data. *Front Psychol* **6,** 1171-1171.

Meng H-J, Cao N, Zhang J & Pi Y-L. (2020). Intermittent theta burst stimulation facilitates functional

connectivity from the dorsal premotor cortex to primary motor cortex. *PeerJ* **8,** e9253-

e9253.

Müller-Dahlhaus JFM, Orekhov Y, Liu Y & Ziemann U. (2008). Interindividual variability and age-

dependency of motor cortical plasticity induced by paired associative stimulation. *Exp Brain*

Res **187,** 467-475.

Müller JF, Orekhov Y, Liu Y & Ziemann U. (2007). Homeostatic plasticity in human motor cortex

demonstrated by two consecutive sessions of paired associative stimulation. *The European*

journal of neuroscience **25,** 3461-3468.

Münchau A, Bloem BR, Irlbacher K, Trimble MR & Rothwell JC. (2002). Functional connectivity of

human premotor and motor cortex explored with repetitive transcranial magnetic

stimulation. *J Neurosci* **22,** 554-561.

Murakami T, Müller-Dahlhaus F, Lu MK & Ziemann U. (2012). Homeostatic metaplasticity of

corticospinal excitatory and intracortical inhibitory neural circuits in human motor cortex. *J*

Physiol **590,** 5765-5781.

Ni Z, Charab S, Gunraj C, Nelson AJ, Udupa K, Yeh IJ & Chen R. (2010). Transcranial Magnetic

Stimulation in Different Current Directions Activates Separate Cortical Circuits. *J*

Neurophysiol **105,** 749-756.

Ni Z, Isayama R, Castillo G, Gunraj C, Saha U & Chen R. (2015). Reduced dorsal premotor cortex and

primary motor cortex connectivity in older adults. *Neurobiol Aging* **36,** 301-303.

Nowak DA, Berner J, Herrnberger B, Kammer T, Grön G & Schönfeldt-Lecuona C. (2009). Continuous

theta-burst stimulation over the dorsal premotor cortex interferes with associative learning

during object lifting. *Cortex* **45,** 473-482.

Opie GM, Cirillo J & Semmler JG. (2018). Age-related changes in late I-waves influence motor cortex

plasticity induction in older adults. *J Physiol* **596,** 2597-2609.

Opie GM, Hand BJ & Semmler JG. (2020). Age-related changes in late synaptic inputs to corticospinal

neurons and their functional significance: A paired-pulse TMS study. *Brain Stimul* **13,** 239-

246.

Opie GM, Liao W-Y & Semmler JG. (2022). Interactions Between Cerebellum and the Intracortical

Excitatory Circuits of Motor Cortex: a Mini-Review. *Cerebellum* **21,** 159-166.

Opie GM & Semmler JG. (2021). Preferential Activation of Unique Motor Cortical Networks With

Transcranial Magnetic Stimulation: A Review of the Physiological, Functional, and Clinical

Evidence. *Neuromodulation* **24,** 813-828.

Opie GM, Vosnakis E, Ridding MC, Ziemann U & Semmler JG. (2017). Priming theta burst stimulation

enhances motor cortex plasticity in young but not old adults. *Brain Stimul* **10,** 298-304.

Parikh PJ & Santello M. (2017). Role of human premotor dorsal region in learning a conditional

visuomotor task. *J Neurophysiol* **117,** 445-456.

Phillips C. (2017). Lifestyle Modulators of Neuroplasticity: How Physical Activity, Mental

Engagement, and Diet Promote Cognitive Health during Aging. *Neural plasticity* **2017,**

3589271.

Puri R & Hinder MR. (2022). Response bias reveals the role of interhemispheric inhibitory networks

in movement preparation and execution. *Neuropsychologia* **165,** 108120.

Ridding MC & Ziemann U. (2010). Determinants of the induction of cortical plasticity by non-invasive

brain stimulation in healthy subjects. *J Physiol* **588,** 2291-2304.

Rossi S, Hallett M, Rossini PM & Pascual-Leone A. (2011). Screening questionnaire before TMS: An

update. *Clin Neurophysiol* **122,** 1686.

Rossini PM, Burke D, Chen R, Cohen LG, Daskalakis Z, Di Iorio R, Di Lazzaro V, Ferreri F, Fitzgerald PB,

George MS, Hallett M, Lefaucheur JP, Langguth B, Matsumoto H, Miniussi C, Nitsche MA,

Pascual-Leone A, Paulus W, Rossi S, Rothwell JC, Siebner HR, Ugawa Y, Walsh V & Ziemann

U. (2015). Non-invasive electrical and magnetic stimulation of the brain, spinal cord, roots

and peripheral nerves: Basic principles and procedures for routine clinical and research

application. An updated report from an I.F.C.N. Committee. *Clin Neurophysiol* **126,** 1071-

1107.

Sakai K, Ugawa Y, Terao Y, Hanajima R, Furubayashi T & Kanazawa I. (1997). Preferential activation of

different I waves by transcranial magnetic stimulation with a figure-of-eight-shaped coil. *Exp*

Brain Res **113,** 24-32.

Sale MV, Ridding MC & Nordstrom MA. (2008). Cortisol inhibits neuroplasticity induction in human

motor cortex. *J Neurosci* **28,** 8285-8293.

Sanes JN & Donoghue JP. (2000). Plasticity and primary motor cortex. *Annu Rev Neurosci* **23,** 393-

415.

Seidler RD, Bernard JA, Burutolu TB, Fling BW, Gordon MT, Gwin JT, Kwak Y & Lipps DB. (2010).

Motor control and aging: links to age-related brain structural, functional, and biochemical

effects. *Neurosci Biobehav Rev* **34,** 721-733.

Spampinato DA, Celnik PA & Rothwell JC. (2020). Cerebellar-Motor Cortex Connectivity: One or Two

Different Networks? *J Neurosci* **40,** 4230-4239.

Suppa A, Bologna M, Gilio F, Lorenzano C, Rothwell JC & Berardelli A. (2008). Preconditioning

Repetitive Transcranial Magnetic Stimulation of Premotor Cortex Can Reduce But Not

Enhance Short-Term Facilitation of Primary Motor Cortex. *J Neurophysiol* **99,** 564-570.

Thirugnanasambandam N, Sparing R, Dafotakis M, Meister IG, Paulus W, Nitsche MA & Fink GR.

(2011). Isometric contraction interferes with transcranial direct current stimulation (tDCS)

induced plasticity – evidence of state-dependent neuromodulation in human motor cortex.

Restor Neurol Neurosci **29,** 311-320.

Todd G, Flavel SC & Ridding MC. (2009). Priming theta-burst repetitive transcranial magnetic

stimulation with low- and high-frequency stimulation. *Exp Brain Res* **195,** 307-315.

Todd G, Kimber TE, Ridding MC & Semmler JG. (2010). Reduced motor cortex plasticity following

inhibitory rTMS in older adults. *Clin Neurophysiol* **121,** 441-447.

Trajkovic J, Romei V, Rushworth MFS & Sel A. (2023). Strengthening connectivity between premotor

and motor cortex increases inter-areal communication in the human brain. *bioRxiv*,

2023.2002.2015.528606.

Volz LJ, Hamada M, Michely J, Pool E-M, Nettekoven C, Rothwell JC & Grefkes Hermann C. (2019).

Modulation of I-wave generating pathways by theta-burst stimulation: a model of plasticity

induction. *J Physiol* **597,** 5963-5971.

Volz LJ, Hamada M, Rothwell JC & Grefkes C. (2015). What Makes the Muscle Twitch: Motor System

Connectivity and TMS-Induced Activity. *Cereb Cortex* **25,** 2346-2353.

Wiethoff S, Hamada M & Rothwell JC. (2014). Variability in Response to Transcranial Direct Current

Stimulation of the Motor Cortex. *Brain Stimul* **7,** 468-475.

Young-Bernier M, Tanguay AN, Davidson PS & Tremblay F. (2014). Short-latency afferent inhibition is

a poor predictor of individual susceptibility to rTMS-induced plasticity in the motor cortex of

young and older adults. *Front Aging Neurosci* **6,** 182.

Ziemann U. (2020). I-waves in motor cortex revisited. *Exp Brain Res* **238,** 1601-1610.

Figure Legends

Figure 1. (A) Subject sample and experimental setup. (B) Experimental procedure. PA, posterior-to-anterior; AP, anterior-to-posterior; LM, lateral-to-medial; RMT, resting motor threshold; AMT, active motor threshold; MEP_{1mV}, standard MEP of ~ 1mV at baseline; $MEP_{0.5mV}$, MEP of ~ 0.5 mV at baseline; PMd, dorsal premotor cortex; iTBS, intermittent theta burst stimulation; CSE, corticospinal excitability.

Figure 2. Changes in PA_{1mV} (A), AP_{1mV} (B), $PA_{0.5mV}$ (C), and $AP_{0.5mV}$ (D) measures of corticospinal excitability following PMd iTBS (grey) and sham (white) stimulation in all participants. Data show EMM (95% CI) with individual subject means. $*P < 0.05$.

Figure 3. Changes in PA_{1mV} (A) and AP_{1mV} (B) measures of corticospinal excitability following PMd iTBS-M1 iTBS (grey) and PMd sham-M1 iTBS (white) in young (no stripes) and older adults (stripes) at 5 and 30 minutes. Data show EMM (95% CI) with individual subject means. $*P < 0.05$. $\#P < 0.05$ compared to 5 minutes in same session.

Figure 4. Changes in $PA_{0.5mV}$ (A) and $AP_{0.5mV}$ (B) measures of corticospinal excitability following PMd iTBS-M1 iTBS (grey) and PMd sham-M1 iTBS (white) in young (no stripes) and older adults (stripes) at 5 and 30 minutes. (C) Changes in $AP_{0.5mV}$ following M1 iTBS in all participants (light grey) at 5 and 30 minutes. Data show EMM (95% CI) with individual subject means. $*P < 0.05$.

Figure 5. Correlation of ranked changes in post-PMd iTBS measures of corticospinal excitability (PA_{1mV}, A; AP_{1mV}, B; $PA_{0.5mV}$, C; $AP_{0.5mV}$, D) with ranked changes in post-M1 iTBS measures of corticospinal excitability.

Table 1. Baseline TMS intensities between sessions for young and older adults.

Measure	Young		Older	
	PMd iTBS-M1 iTBS	PMd sham-M1 iTBS	PMd iTBS-M1 iTBS	PMd sham-M1 iTBS
PA				
RMT_{PA} (% MSO)	47.3 [42.3, 52.3]	47.8 [42.8, 52.8]	50.4 [45.4, 55.4]	51.3 [46.3, 56.3]
AMT_{PA} (% MSO)	39.9 [36.4, 43.4]	39.3 [35.8, 42.8]	42.6 [39.1, 46.1]	43.2 [39.7, 46.7]
$1mV_{PA}$ (% MSO)	56.5 [49.9, 63.1]	57.4 [50.8, 64.0]	65.4 [58.5, 72.2]	63.3 [56.4, 70.1]
$0.5mV_{PA}$ (% MSO)	53.0 [46.1, 59.9]	53.8 [46.9, 60.7]	61.5 [54.5, 68.4]	61.3 [54.3, 68.2]
AP				
RMT_{AP} (% MSO)	61.3 [55.6, 67.0]	62.3 [56.6, 68.0]	66.3 [60.4, 72.2]	65.3 [59.4, 71.2]
AMT_{AP} (% MSO)	54.2 [49.0, 59.4]	54.1 [49.0, 59.3]	59.1 [54.0, 64.3]	57.2 [52.0, 62.4]
$1mV_{AP}$ (% MSO)	73.9 [65.6, 82.1]	73.8 [65.6, 82.0][a]	83.8 [74.6, 93.1]	79.3 [70.0, 88.5][a]
$0.5mV_{AP}$ (% MSO)	70.9 [63.5, 78.3]	71.3 [63.9, 78.7]	79.4 [71.4, 89.4]	76.4 [68.4, 84.4]
LM				
AMT_{LM} (% MSO)	45.3 [40.5, 50.2]	45.3 [40.4, 50.1]	49.9 [45.1, 54.8]	48.8 [43.9, 53.7]
TBS				
RMT_{Rapid} (% MSO)	55.7 [50.8, 60.6]	57.8 [52.9, 62.7]	57.7 [52.8, 62.6]	58.1 [53.2, 63.0]

Data show EMM [95% CI; lower, upper]. [a]$P < 0.05$ compared to iTBS session.

Table 2. Baseline responses of corticospinal excitability and I-wave recruitment between sessions.

Measure	Young		Older	
	PMd iTBS-M1 iTBS	PMd sham-M1 iTBS	PMd iTBS-M1 iTBS	PMd sham-M1 iTBS
PA				
PA-LM latency (ms)	1.39 [0.89, 1.88]	1.54 [1.04, 2.03]	1.97 [1.47, 2.46]	2.02 [1.52, 2.51]
$1mV_{PA}$ (mV)	1.03 [0.94, 1.12]	0.93 [0.85, 1.01]	0.89 [0.81, 0.97][a]	0.96 [0.87, 1.04]
$0.5mV_{PA}$ (mV)	0.53 [0.46, 0.60]	0.49 [0.42, 0.56]	0.50 [0.43, 0.57]	0.51 [0.44, 0.58]
AP				
AP-LM latency (ms)	3.49 [3.00, 3.98][b]	3.54 [3.05, 4.03][b]	3.69 [3.20, 4.18][b]	3.99 [3.50, 4.48][b]
$1mV_{AP}$ (mV)	0.97 [0.87, 1.06]	0.88 [0.79, 0.97]	1.02 [0.91, 1.14]	0.99 [0.88, 1.10]
$0.5mV_{AP}$ (mV)	0.47 [0.41, 0.53]	0.44 [0.39, 0.50]	0.45 [0.40, 0.51]	0.45 [0.39, 0.50]

Data show EMM [95% CI; lower, upper]. [a]$P < 0.05$ compared to young. [b]$P < 0.05$ compared to PA-LM latency.

Table 3. Comparison of VAS responses (mean ± STD) between sessions.

Question	PMd iTBS-M1 iTBS	PMd sham-M1 iTBS
How uncomfortable were the TMS pulses (0, not uncomfortable at all; 10, highly uncomfortable)?	2.67 ± 2.60	2.5 ± 2.79
If there were any twitches in the right hand, how strong were they (0, no twitches; 10, very strong cramp)?	0.63 ± 1.40	0.60 ± 1.13
How localised were the sensations from TMS pulses (0, highly localised; 10, widespread)?	2.03 ± 2.47	0.50 ± 1.04*

Data show mean ± standard deviation.*$P < 0.0167$ compared to iTBS.

Figure 1. (A) Subject sample and experimental setup. (B) Experimental procedure. PA, posterior-to-anterior; AP, anterior-to-posterior; LM, lateral-to-medial; RMT, resting motor threshold; AMT, active motor threshold; MEP_{1mV}, standard MEP of ~ 1mV at baseline; $MEP_{0.5mV}$, MEP of ~ 0.5 mV at baseline; PMd, dorsal premotor cortex; iTBS, intermittent theta burst stimulation; CSE, corticospinal excitability.

Figure 2. Changes in PA$_{1mV}$ (A), AP$_{1mV}$ (B), PA$_{0.5mV}$ (C), and AP$_{0.5mV}$ (D) measures of corticospinal excitability following PMd iTBS (grey) and sham (white) stimulation in all participants. Data show EMM (95% CI) with individual subject means. *$P < 0.05$.

Figure 3. Changes in PA$_{1mV}$ (A) and AP$_{1mV}$ (B) measures of corticospinal excitability following PMd iTBS-M1 iTBS (grey) and PMd sham-M1 iTBS (white) in young (no stripes) and older adults (stripes) at 5 and 30 minutes. Data show EMM (95% CI) with individual subject means. *$P < 0.05$. #$P < 0.05$ compared to 5 minutes in same session.

Figure 4. Changes in $PA_{0.5mV}$ (A) and $AP_{0.5mV}$ (B) measures of corticospinal excitability following PMd iTBS-M1 iTBS (grey) and PMd sham-M1 iTBS (white) in young (no stripes) and older adults (stripes) at 5 and 30 minutes. (C) Changes in $AP_{0.5mV}$ following M1 iTBS in all participants (light grey) at 5 and 30 minutes. Data show EMM (95% CI) with individual subject means. *$P < 0.05$.

Figure 5. Correlation of ranked changes in post-PMd iTBS measures of corticospinal excitability (PA$_{1mV}$, A; AP$_{1mV}$, B; PA$_{0.5mV}$, C; AP$_{0.5mV}$, D) with ranked changes in post-M1 iTBS measures of corticospinal excitability.