Asymmetric conformations and lipid interactions shape the ATP-coupled cycle of

a heterodimeric ABC transporter

Qingyu Tang[1], Matt Sinclair[2], Hale S. Hasdemir [2], Richard Stein[1], Erkan Karakas[1],

Emad Tajkhorshid[2], & Hassane Mchaourab[1*]

[1] Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville,

TN 37232, USA. [2]Theoretical and Computational Biophysics Group, NIH Resource for

Macromolecular Modeling and Visualization, Beckman Institute for Advanced Science

and Technology, Department of Biochemistry, and Center for Biophysics and

Quantitative Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801,

USA

*email: hassane.mchaourab@vanderbilt.edu

Abstract

To illuminate the structural origin of catalytic asymmetry of heterodimeric ABC

transporters and how it shapes the energetics of their conformational cycles, we used

cryo-electron microscopy (cryo-EM), double electron-electron resonance spectroscopy

(DEER), and molecular dynamics (MD) simulations, to capture and characterize

conformational states of the heterodimeric ABC multidrug exporter BmrCD in lipid

nanodiscs. In addition to multiple ATP- and substrate-bound inward-facing (IF)

conformations, we obtained the structure of an occluded (OC) conformation wherein the

unique extracellular domain (ECD) twists to partially open the extracellular gate. In

conjunction with DEER analysis of the populations of these conformations, the

structures reveal that ATP-powered isomerization entails changes in the relative

symmetry of the BmrC and BmrD subunits that propagates from the transmembrane

domain (TMD) to the nucleotide binding domain (NBD). The structures uncover

asymmetric substrate and Mg^{2+} binding which we hypothesize are required for triggering

ATP hydrolysis preferentially in one of the nucleotide-binding sites. MD simulations

demonstrated that multiple lipid molecules, identified from the cryo-EM density maps,

differentially bind the IF versus the OC conformation thus modulating their relative

stability. In addition to establishing how lipid interactions with BmrCD modulate the

energy landscape, our findings are framed in a distinct transport model that highlights

the role of asymmetric conformations in the ATP-coupled cycle with implications to the

mechanism of ABC transporters in general.

Introduction

ATP-binding cassette (ABC) transporters harness the energy of ATP hydrolysis to traffic diverse substrates across cell membranes[1-3]. The remarkable breadth of their substrates confers to ABC transporters critical roles in physiological processes and multidrug resistance[3-6]. Prokaryotic ABC transporters mediate nutrient uptake, antimicrobial peptide export, cell viability, virulence, and pathogenicity[7]. In humans, a number of ABC transporters of the exporter type are involved in lipid homeostasis, multidrug efflux, and the transport of immunogenic peptides. In addition, these transporters have been implicated in cardiovascular, metabolic, and neurodegenerative diseases[8-10].

Numerous studies have defined the basic molecular architectures of ABC exporters, delineated structural elements involved in their ATP-powered alternating access and revealed modes of substrate interactions[3,4,11-20]. Invariably, ABC transporters have a conserved architecture consisting of two nucleotide-binding domains (NBDs) which hydrolyze ATP to energize alternating access of two transmembrane domains (TMDs), the site of substrate recognition and translocation[3]. The NBDs are considered the hallmark of the family sharing extensive sequence similarity and architecture among all ABC transporters[1]. In contrast, at least 7 types of ABC transporters identified based on TMD folds and topologies. Among them, type IV ABC exporters, whose members include P-gp, TmrAB, TM287/288, MsbA, MRP1, McjD, and ABCB4, share a domain-swapped TMD arrangement[21].

While there is an emerging consensus on the structural elements of alternative access of type IV ABC exporters, the mechanism of ATP energy coupling has been less

characterized. One of the prominent models posits the alternating hydrolysis of ATP in

the two nucleotide binding sites. Indirect evidence, including from DEER analysis of

mammalian ABCB1 (Pgp)[22,23] was consistent with biochemical data[24-26] suggesting the

population of locally asymmetric intermediates in the substrate- and ATP-coupled cycle.

In contrast, cryo-EM structures of Pgp, in a catalytically impaired background and in the

absence of substrates, were found to be symmetric particularly at the nucleotide-binding

sites (NBSs)[20]. Subsequent DEER studies demonstrated that local asymmetry was not

observed in the absence of substrates or when bound to high affinity inhibitors[22]

suggesting a critical role for these conformations in the functional mechanism.

A remarkable variation on the theme of ATP energy conversion in ABC transporters is

the evolution of heterodimeric ABC exporters[27,28] which are distinguished by having an

intrinsically asymmetric ATP catalytic cycle resulting from the selective substitution of

one of the catalytic glutamates in the NBD[12,29,30]. Thus, they provide unique models to

investigate structural asymmetry and elucidate its consequence on the conformational

cycle with relevance to ABC exporters in general. A number of ABC heterodimers have

been investigated[12,30,31], yielding insight into their structures and the mechanism of

alternating access. TmrAB, an ABC heterodimeric transporter from *Thermus*

thermophilus, was captured in five conformations that were proposed to represent the

ATP-powered cycle[12]. However, as noted previously by us[11] and others[27,32], the

energetics of the TmrAB conformational cycle may not be reflective of the typical type IV

exporter. In particular, the dynamics of the NBDs may be constrained by strong

helix/helix interactions at the C-terminus. Moreover, the role of the lipid-transporter

interactions was not explicitly explored. Finally, the quality of the experimental density

for the peptide substrate in these structures was not amenable to detailed structure

determination.

Our model heterodimeric ABC exporter, BmrCD, is associated with antibiotic efflux in

the Gram-positive bacterium *Bacillus subtilis*[33]. Incorporated in giant unilamellar

vesicles, BmrCD can transport ethidium bromide *in vitro*, a process powered by ATP

and inhibited by orthovanadate[34]. Previous studies from our laboratory revealed that

ATP-coupled conformational changes in BmrCD entail population of asymmetric

configurations of the catalytically inequivalent NBSs[29,35]. A cryo-EM structure of an ATP-

bound inward-facing (IF) conformation of BmrCD in detergent micelles uncovered an

asymmetric arrangement in the TMD that was ostensibly stabilized by the inequivalent

binding of two molecules of the substrate Hoechst-33342 (hereafter referred to as

Hoechst, HT or H)[11].

The stability of the ATP- and Hoechst-bound IF conformation of BmrCD was

considered unusual, as most type IV ABC exporters were captured in an outward-facing

(OF) conformation under these conditions[12,20,36-39]. Along with DEER analysis

demonstrating the requirement of ATP hydrolysis for transition to the OF conformation[29],

this finding suggests a distinct energy landscape for BmrCD, particularly in comparison

to TmrAB. Another unique aspect of BmrCD is the energetic dependence of the

conformational equilibrium on interactions with the lipid environment. Although ATP

hydrolysis is stimulated by incorporation of the BmrCD into lipid bilayers, the NBD to

TMD coupling, which drives transition to the OF conformation, becomes strictly

dependent on substrate binding[29]. Even in the presence of the substrate Hoechst, only

a fraction of BmrCD reaches the OF conformation. Thus, BmrCD provides a unique

model to dissect the energetics of lipid modulation of ATP-coupled transport in ABC

transporters[29].

To address the outstanding questions of the role of asymmetry and lipid interactions

in the conformational cycle of ABC exporters, we determined structures of BmrCD in

multiple distinct conformations in lipid nanodiscs and correlated the population of these

conformations with DEER analysis under different nucleotide and substrate conditions.

This integrated analysis illuminates novel aspects of the BmrCD conformational cycle.

First, the transition entails population of structurally asymmetric intermediates.

Asymmetry propagates from the TMD (in the IF conformation) to the NBD (in the

occluded (OC) conformation) leading to differential Mg^{2+} binding at the two NBSs and

ATP hydrolysis at the consensus NBS. Second, we determined IF structures without the

substrate Hoechst, with one, or two antiparallel Hoechst molecules. Importantly, we

observed that Hoechst binding is associated with differential Mg^{2+} binding at the NBSs.

Finally, lipids shape the energy of the cycle through differential interactions with the IF

and OF conformations. Lipid molecules, visualized from cryo-EM electron density maps,

were subjected to MD simulations to identify those that are stably bound. We found that

the IF conformation of BmrCD, by virtue of exposure of critical residues to the bilayer,

interacts more extensively with lipids, thus explaining the stabilization of IF relative to

OF. These findings are integrated to propose a transport model for heterodimeric ABC

transporters with implications to type IV exporters.

Results

IF and OC conformations of BmrCD in lipid bilayers. We determined cryo-EM

structures of wild type (WT) and a catalytically impaired mutant of BmrCD in lipid

nanodiscs under conditions expected to stabilize distinct transport intermediates. The

mutant, constructed on a cysteine-less (C-less) background, contains glutamine

substitutions that impair ATP hydrolysis at D500 of BmrC and E592 of BmrD (referred to

hereafter as BmrCD*-QQ). The purified proteins were reconstituted in lipid nanodiscs of

two different compositions: phosphatidylcholine (PC)/phosphatidic acid (PA), and PC

and *E. coli* polar lipids, previously shown to stimulate ATP hydrolysis in WT and C-less

backgrounds[29]. Although we determined multiple structures of WT and QQ BmrCD in

both types of nanodiscs (Supplementary Fig. 1 and Table 1), we focus here on the best-

quality, unique multiple conformations which were obtained in the BmrCD*-QQ

background.

We report the structure of three IF conformations bound to different combinations of

ATP and Hoechst. Under conditions of excess substrate Hoechst and ATP/Mg^{2+}, an IF

conformation of BmrCD*-QQ in PC/PA lipids was determined to 3.3 Å resolution

(referred to hereafter as BmrCD_IF-2H/ATP) (Fig. 1c, Supplementary Figs. 1-5 and

Supplementary Table 1). Along with 2 Hoechst and 2 ATP molecules, 20 lipid molecules

can also be partially or fully modeled into the electron density (Supplementary Fig. 5).

Remarkably only a single Mg^{2+} ion in the consensus NBS was visible in the density

maps (see below for a detailed discussion). The BmrCD_IF-2H/ATP structure is quite

similar to the BmrCD structure in detergent micelles (PDB ID: 7M33, referred hereafter

as BmrCD_det)[11] with an RMSD of 0.7 Å across 959 Cα atoms (Supplementary Table

2).

Under similar conditions of Hoechst and ATP/Mg^{2+} but reconstituted in PC/*E. coli*

polar lipids nanodiscs (dataset 2, Fig. 1b), another IF conformation of BmrCD*-QQ was

identified with density for only one Hoechst molecule in the chamber (Fig. 1b,

Supplementary Figs. 1-5 and Supplementary Table 1). This structure, referred to as

(BmrCD_IF-H/ATP) represents an intermediate state with two ATP molecules, one Mg^{2+}

and one substrate. It appears that the energetics of BmrCD*-QQ in the PC/ *E. coli* polar

lipids mixture is favorable for an IF conformation bound to a single Hoechst. Twenty-one

lipid molecules were visualized in this structure.

The third IF conformation of BmrCD*-QQ was obtained in the presence of ATP but

absence of Hoechst in PC/PA bilayers (referred hereafter as BmrCD_IF-ATP, Fig. 1a,

Supplementary Figs. 1-5, and Supplementary Table 1). BmrCD_IF-ATP is characterized

by a large chamber open to the cytoplasm, had two well resolved ATP molecules,

although no identifiable Mg^{2+} ion, as well as 25 densities that arise from lipid molecules

(Supplementary Fig. 5). This IF structure is also very similar to BmrCD_IF-2H/ATP and

BmrCD_det, with RMSD values of 0.9 Å across 1,153 Cα atoms and 1.4 Å across 1078

Cα atoms, respectively. However, in the absence of bound Hoechst, this IF

conformation features a larger substrate binding cavity compared to the detergent and

Hoechst-bound IF conformations. The solvent-accessible volume of BmrCD_IF-ATP

substrate-binding chamber calculated by CASTp is 9,354 $Å^3$ whereas it is 7,882 $Å^3$ in

BmrCD_IF-2H/ATP and 7,538 $Å^3$ in BmrCD_det (Supplementary Fig. 6)[40]. One

possible reason for the narrowing of the chamber in the later structures is interaction

with the substrate Hoechst which can "pull in" the helices lining the chamber. However,

it should be noted that BmrCD_IF-ATP was determined in larger nanodiscs using the

MSP1D1E3 scaffold protein[41]. To exclude the possibility that the size of the nanodiscs

influence the opening of the chamber, we examined ATP-bound BmrCD in nanodiscs

formed using a smaller membrane scaffold protein MSP1D1, with a diameter of 9.5 nm

compared to MSP1D1E3 of 12 nm (Dataset 2, Supplementary Fig. 1). We identified a

very similar IF structure (BmrCD_IF-ATP2) suggesting that BmrCD conformation is not

constrained by the size of nanodiscs (Supplementary Fig. 1), a finding consistent with

previous results by other groups[13].

In addition to the IF conformations, we obtained the structure of an occluded (OC)

conformation of BmrCD*-QQ in PC/*E. coli* polar lipids (referred hereafter as

BmrCD_OC-ATP, Fig. 1d, Supplementary Figs. 1-5, 7 and Supplementary Table 1,

dataset 4). In this inward-closed structure, the NBDs are in contact with each other to

form the NBSs and the chamber was occluded to the cytoplasm. In the same dataset,

we also detected IF conformations in 2D classification and 3D models (Supplementary

Figs. 1-5, 8). The electron density maps displayed anisotropy in both OC and IF

conformations (Supplementary Fig. 4d and e). Heterogeneous refinement implemented

in CryoSPARC[42] was helpful in resolving the anisotropy problem of the maps for the IF

and OC structures.

Finally, while attempting to obtain an OF conformation in a WT background, we

determined another OC structure in PC/PA bilayers (dataset 5). BmrCD was incubated

with ATP and vanadate (Vi) to trap the high energy state (HES) populated following ATP

hydrolysis (Supplementary Figs. 1-5). The OC in the WT and QQ backgrounds are

almost identical (RMSD value of 0.6 Å over 1039 Cα atoms, Supplementary Table 2).

Although we could not conclusively establish from the density, we presume that the

likely had ADP and Vi at the consensus NBS, given a 10-minute incubation at 37°C.

The density maps of an IF conformation under these conditions was not amenable to

high resolution structure primarily due to highly flexible NBDs. Overall, we didn't

observe substantial differences for the same conformations between the two lipid

compositions or between the WT and QQ OC conformations. Moreover, the stably

bound lipid molecules, as identified by MD simulations (see below), appeared in similar

locations in all the structures regardless of lipid composition (Supplementary Fig. 9).

A sequential Hoechst binding model involving two ionic latches. The two IF

structures with one (BmrCD_IF-H/ATP) or two (BmrCD_IF-2H/ATP) Hoechst molecules

suggest a two-step binding process that entails changes in the interactions of two ionic

"latches" in the substrate binding chamber. Superposition of the two structures highlight

high similarity with a RMSD of 0.4 Å across 1,131 Cα atoms. Remarkably, in both IF

structures one Hoechst (HT1) molecule is located at the same position on the right half

of the chamber (Fig. 2). As noted previously[11], the binding of the antiparallel Hoechst

molecules is stabilized by two patches of negatively charged residues in orientations

that match the bound Hoechst. A close-up analysis shows that W290 of BmrC and

H338 of BmrD form an ionic "latch" (WH-latch) in the middle of the left part of the

chamber in BmrCD_IF-H/ATP which blocks the binding of the second Hoechst in this

structure (Fig. 2b and c, Supplementary Fig. 10). Distinct densities for the side chains of

these residues unambiguously suggest their projection into the location of the second

Hoechst. The latch is open in the fully loaded structure. The importance of W290 was

highlighted in our previous work showing that ablation of the side chain eliminates

cooperative simulation of ATP turnover by Hoechst[11].

Furthermore, examination of the chamber entrance at or around the intracellular side

of the TMD exposes another ionic "latch" consisting of R304 and H184 of BmrC (RH-

latch, Fig. 2d). This latch is open in the one-Hoechst structure with the sidechain

distance at 11 Å (Fig. 2d, middle panel), whereas it is closed in the two-Hoechst

structure with the sidechain distance at 4.4 Å (Fig. 2d, left panel). We propose that the

energetics of these latches contribute to the overall binding of Hoechst and may well

modulate the number of bound molecules. Interestingly, in the substrate-free IF

structure BmrCD_IF-ATP, the RH-latch is open with the sidechain distance of 8 Å

primed for the first Hoechst binding (Fig. 2d, right panel). The WH-latch opens for the

second Hoechst which may then triggers the closing of the RH-latch at the entrance.

Structural rearrangements underlying the IF to OC transition of BmrCD.

Superposition of BmrCD_IF-2H/ATP and BmrCD_OC-ATP, yielded an RMSD value of

6.5 Å over 1,199 Cα atoms, suggesting drastic structural rearrangements. Comparison

at the domain level highlights that the largest movements are mainly attributed to the

TMDs with RMSD values of 2.3 Å for BmrD-TMD (259 Cα atoms) and 4.8 Å for BmrC-

TMD (314 Cα atoms). By contrast, the RMSD values at the NBDs are smaller: 1.3 Å for

BmrD-NBD (190 Cα atoms) and 1.1 Å for BmrC-NBD (215 Cα atoms).

TM by TM inspection revealed that upon transition to the OC conformation, TM1 of

BmrC tilts by 12° whereas the TM2/TM3 pair and TM6 of BmrC tilt by 16°. TMD

rearrangement is coupled with the movement of the NBDs toward each other by

approximately 23 Å to form two engaged NBSs. The transition between IF and OC

closes the intracellular TMDs thereby occluding substrate entrance. Concomitantly, the

TM4/TM5 pair of BmrD also tilts about 10° to contribute to closing of the entrance (Fig.

3a, Supplementary Fig. 6). A similar conformational change from IF to OC and OF

structures also can be observed in the heterodimeric ABC transporter TmrAB[12].

As previously reported[29], and in agreement with the cryo-EM structures, DEER

measurement of distance distributions for the intracellular spin label pair $96^{BmrC}/188^{BmrD}$

(96/188) shows a predominant IF population under all conditions. A minor distance

population corresponding to the putative OF conformation is identified in the high

energy post-hydrolysis state (HES) (trapped by vanadate following ATP turnover) but

not upon AMP-PNP (a non-hydrolyzable analog of ATP) addition (Fig. 3b and c,

Supplementary Fig. 11). Notably, in the QQ background (Fig. 3d), the distance

population stabilized by ATP binding, and associated with closing of the intracellular

side, is larger compared to the HES in the WT background suggesting a stabilization of

an OC conformation by the QQ mutations. Although conformational changes were not

detected on the extracellular side in the cryo-EM structures, the distance distribution of

the extracellular pair $55^{BmrC}/146^{BmrD}$ (55/146) shows a minor OF population (Fig. 3c and

d, Supplementary Fig. 11) at longer distance compared to the predictions from the OC

structure. Attempts to stabilize and determine the structure of the OF conformation by

cryo-EM were not successful.

NBD dimer assembly and differential Mg^{2+} binding. Supplementary Fig. 12a shows

the details of how BmrCD engages bound ATP in the consensus and degenerate NBSs

in the OC conformation. At the degenerate NBS (Supplementary Fig. 12a and b), BmrC

residue Y346 (typically part of the conserved A-loop) and S565 of BmrD interacts with

the adenine ring of ATP. The phosphate groups are coordinated by residues G374,

S375, G376, and K377 from Walker A of BmrC, Q419 from the Q-loop of BmrC, and

S568 and G570 of BmrD (Supplementary Fig. 12b).

Binding of the ATP molecule at the consensus NBS shows small but distinct

differences relative to the degenerate NBS. In contrast to the equivalent residues at the

degenerate NBS, S476 and Q479 of BmrC stabilize the ribose of ATP. G478 on the

ABC signature motif of BmrC, G468, K469, S470, and S471 on Walker A and Q592 on

Walker B of BmrD, and Q511, H623 of BmrD contribute to phosphate group binding

(Supplementary Fig. 12c). Interestingly, comparison of the catalytic residues D500 in

the degenerate NBS and E592 in consensus NBS (both mutated to glutamines) in

BmrCD_IF-2H/ATP , BmrCD_IF-ATP, and BmrCD_OC-ATP, reveals that the side chain

of Q500 is oriented toward the γ-phosphate group of ATP. On the other hand, the side

chain of Q592 flips its orientation toward the γ-phosphate group of ATP in OC while its

points away in the IF (Supplementary Fig. 13). This phenomenon was also reported in

TmrAB at canonical and non-canonical NBSs, in which the side chain of the former flips

but the other does not[12].

Comparison of the NBSs in the three nanodisc structures revealed that one of the IF

conformations, obtained in the absence of Hoechst (BmrCD_IF-ATP), lacks density

corresponding to the Mg^{2+} ions, whereas the IF obtained with Hoechst has discernable

Mg^{2+} density only in the consensus NBS. The space for Mg^{2+} is occupied by S470 in the

consensus NBS and T378 in the degenerate NBS (Supplementary Fig. 14). Because

the quality of the density maps in this region for the three structures appears equivalent,

we surmise that these observations reflect the affinity for Mg^{2+}, being preferentially

stabilized at the consensus NBS prior to engaging in ATP hydrolysis.

Structural asymmetry of the NBSs. Remarkably, despite the catalytic impairment of

the BmrCD construct, we observed stark structural asymmetry in the assembly of the

two NBSs upon transition from the IF to the OC conformations. To highlight and

experimentally support the distinct arrangement of the two NBSs, we compared

experimental distance distributions in lipid bilayers to those predicted based on the cryo-

EM structures (Fig. 4, Supplementary Fig. 15). For this purpose, distance distributions

for spin label pairs monitoring the two NBSs were calculated in all three BmrCD

structures (BmrCD_IF-2H/ATP , BmrCD_IF-ATP, and BmrCD_OC-ATP using

CHARMHH-GUI server and the MDDS method[43]). These distance distributions at the

two NBSs ($348^{BmrC}/532^{BmrD}$ in the degenerate NBS (348/532) and $440^{BmrC}/441^{BmrD}$ in the

consensus NBS (440/441)) of BmrCD_IF-2H/ATP and BmrCD_IF-ATP overlap.

Predicted distances distributions are drastically shorter and manifestly asymmetric in

BmrCD_OC-ATP (Fig. 4b red dashed traces). The corresponding distance distributions

predicted by the BmrCD_det structure are shorter and symmetric (Supplementary Fig.

15)[11] suggesting that lipids may stabilize structural asymmetry.

Experimental distance distributions obtained in the QQ background in lipid bilayers

are multi-component and asymmetric, reflecting the coexistence of multiple

conformations. At the degenerate NBS, the distribution obtained in the presence of

Hoechst (Fig. 4b) overlaps the one predicted from the IF structures. In contrast, a

second major population was observed at approximately 45 Å at the consensus pair

440/441 in nanodiscs under the same conditions.

Binding of ATP and Hoechst yields an experimental distribution that is almost

superimposable to the predicted one at the degenerate NBS, although a component at

a longer distance overlapping the IF conformation persists. In contrast, at the

consensus NBS the predicted and experimental distributions deviate substantially. Two

populations are observed in the latter compared to a unimodal population at a longer

average distance in the former (Fig. 4). The prominent short component in the

experimental distribution is almost superimposable to a similar component for the same

consensus pair obtained by vanadate trapping following ATP hydrolysis in the WT

background (Supplementary Fig. 15). Notably, this component is stabilized by Hoechst

binding suggesting that it corresponds to the OF conformation. Therefore, we conclude

that the OC structure does not capture certain structural features at the consensus

NBS, consistent with the lack of opening of the extracellular side in this structure noted

in Fig. 3.

Dynamics and asymmetry of the TMD. In addition to the large ATP-dependent

structural changes underpinning the IF to OC transition, we observed changes in the

symmetry arrangement of the BmrC and BmrD TMDs. Superposition of the two halves

of the TMD around the pseudosymmetry axis in the three nanodiscs and in the previous

detergent BmrCD structures (Fig. 5, Supplementary Fig. 16) highlights outward shifts of

the TM helices in BmrD relative to BmrC[11]. These outward shifts are observed at both

the extracellular and intracellular sides in all structures but to different extents

(Supplementary Figs. 16 and 17), suggesting the TM helices on the two sides of the

membrane may undergo independent movement.

To quantitatively capture the asymmetry across conformations in lipid bilayers, we

measured distance changes between corresponding helices of BmrC and BmrD using

TM-align[44] (Fig. 5). For both IF structures, the distance changes reveal increased

asymmetry relative to the detergent structure for TM4 on the extracellular side and TM2

and TM4 on the intracellular side (Fig. 5a, b, and c) implying that the lipid environment

is more conducive for asymmetry between BmrC and BmrD.

Drastic change in asymmetry is observed upon the transition to the OC conformation.

At the extracellular side there is an increase in the asymmetry of TM1 and TM2 that

may indicate initiation of transition to an OF conformation even though the extracellular

side remains closed in the OC conformation. Remarkably, concomitant movements in

TM2, TM3, and TM6 reduce the local asymmetry which we attribute to the loss of

interaction with Hoechst (Supplementary Fig. 18). In contrast, TM4's asymmetry is

accentuated around a kink in the bilayer.

The unique ECD domain of BmrCD twists to open. BmrCD is the only structurally

characterized heterodimer that harbors an extracellular domain, the deletion of which

abolishes substrate stimulation of ATP hydrolysis[11]. Superposition of ECDs from

BmrCD_IF-2H/ATP, BmrCD_IF-ATP, and BmrCD_OC-ATP, based on BmrC alignment,

exposes a "twisting" motion of this domain in the latter. The anatomy of this twist

involves movements of the loop between residues 118 and 127, the loop between

residues 73 and 89, and the loop between residues 67 and 70 of 13 Å, 20 Å, and 10 Å,

respectively (Fig. 6a). This twist alters the electrostatics of the domain as a whole,

displacing a patch of positive charges, which could potentially enable an exit pathway

for substrate release as it is squeezed out by the transition from IF to OC. (Fig. 6b).

Meanwhile, the charge distribution of the ECD docking surface to the extracellular side

of the TMD remains invariable in all structures (Supplementary Fig. 19) implying that the

ECD movement is independent of the TMD.

We tested if the ATP-powered partial opening movement of the ECD is also observed

in the WT background. Distance distributions of three pairs $46^{BmrC}/126^{BmrD}$ (46/126),

$55^{BmrC}/126^{BmrD}$ (55/126), and $126^{BmrD}/146^{BmrD}$ (126/146), predicted by MDDS based on

the IF structures, are in agreement with the DEER data (Fig. 6c and d, Supplementary

Fig. 20). Upon transition to the HES, prominent distance changes are observed

between TM2 and the ECD (the 55/126 pair) as predicted by the OC conformation. The

structures and the DEER data suggest that part of the ECD is anchored to the TMD

specifically by its interactions with TM1 and TM2 of BmrD (the 126/146 pair) and

interactions with TM1 and/or TM5 of BmrC (Fig. 6c and d, Supplementary Fig. 21). The

relatively broader distributions in the HES (55/126) are indicative of increased dynamics

of part of this domain which correspond to the relatively poorer cryo-EM density of ECD

in the OC conformation (Supplementary Fig. 5).

Lipids preferentially bind the IF conformation. We observed substantial density

around the TMDs in all the high-resolution structures. Based on the shape and length of

these densities, we modeled phospholipid molecules in the vicinity of the transporter. To

assess the stability of the individual lipids, we monitored the minimum distance of each

to the transporter surface (Supplementary Fig. 22a and b) during MD simulations (see

methods). This analysis identifies those lipids which may be stably bound to the protein

and those that may have been captured by cryo-EM simply due to the nanodiscs

environment. In the simulation trajectories, we observed variable stability for the cryo-

EM-modeled lipids relative to their initial position and conformation (Supplementary Fig.

22). Overall, the IF conformation appears to be more hospitable to bound lipids,

including several lipids near the elbow helix of BmrD (Lipids 2 and 14 in IF) as well as

other regions of the protein such as the interface between the two arms of the TMDs

(Lipids 3 and 9 in IF) (Supplementary Fig. 22c). The OC conformation on the other hand

seems to mostly stabilize lipids around the elbow helix.

    To further characterize the stability and dynamics of the lipids during the simulations,

we binned lipid phosphorous coordinates within 20 Å of the protein for both the upper

and lower leaflets. From this dataset, we performed kernel density estimation to obtain a

density map (Fig. 7a and b) in which sites of high-density correlate with stable lipid

binding whereas smeared densities reflect lipids that are simply gliding along the

surface of the protein. For example, it appears that Lipid 13 in the IF conformation is

stably bound near its initial placement. However, the density near Lipid 13 is attributed

to a different lipid from the bulk bilayer (Supplementary Fig. 22a). Even lipids that seem

to unbind in some but not all simulation replicas, such as Lipid 9 (Supplementary Fig.

22a), appear to stay proximal to the protein in the unbound replicas.

For the OC conformation, the kernel densities highlight that Lipids 6 and 16 appear

stably bound, being positioned directly in high density wells (Fig. 7b). Lipid 4, which

fluctuates away from the protein in simulations, does occupy the same high-density

region as Lipid 6 near the elbow helix region of BmrD (Fig. 7b, Supplementary Fig. 22b

and c). While in some replicas Lipid 4 does not stay fully bound, the convergence at the

end of the simulation indicates that it binds in this region albeit less strongly than the

equivalent lipid (Lipid 2) in the IF conformation (Supplementary Fig. 22). Interestingly,

Lipids 10 and 15 are found to be stably bound according to the distance analysis but the

density maps show that, in both cases, these lipids are moving along the protein surface

and not settled in a high density well (Fig. 7b, Supplementary Fig. 22b)

To identify BmrCD's contributions to these protein-lipid interactions, we analyzed the

binding residence times of the modeled lipids at basic surface residues (Fig. 7c). These

residence times further highlight the propensity for IF to stabilize lipids over the OC

conformation. To probe the origin of this propensity, we measured the distributions of

side chain angles of basic surface residues in both conformations (Fig. 7d). For this

purpose, this angle is defined by two vectors: one from the center of mass (CoM) of the

backbone atoms of neighboring residues, and the other vector is drawn from the alpha

carbon to the CoM of the terminal charged nitrogen (NZ for lysine and NH1/NH2 for

arginine). In cases of stable lipid coordination, the available angle space which can be

explored by a given residue is reduced as is observed for several major residues

including K2 (BmrD) of the elbow helix in the IF conformation (Supplementary Fig. 22c).

Other notable residues are K182 (BmrC), which lies at the vertex of TMD, and R180

(BmrD) which sits along the interface between BmrC and BmrD near the elbow helix.

The apparent importance of the elbow helix in recruiting lipids focused our attention

on key residues which might be inducing strong local protein-lipid interactions. Several

charged residues, K2, K5, and R15 of BmrD, were *in silico* mutated to alanine in order

to assess the effects of charge interactions on bound-lipid residence time relative to WT

(Supplementary Fig. 22d). In all cases and for both conformations, we observed a

decrease in the high residence binding regime. This was coupled with a lack of

perturbation in the low residence binding as expected for surface residues.

Discussion

Conformational dynamics of BmrCD. The integrated investigation of BmrCD

presented above addresses two long-standing questions in the field- namely the origin

and role of asymmetry in the conformational cycle of ABC transporters. We present

direct experimental evidence of asymmetric substrate and Mg^{2+} binding that appear to

be coupled to the TMD and NBD structural asymmetries. Furthermore, our study

contributes structural and dynamic insights into how lipid/transporter interactions can

modulate function with implications to membrane proteins in general. Three snapshots

of the ATP-powered conformational cycle in lipid bilayers describe how ATP binding

engenders transition to an OC state in which cytoplasmic access to the chamber at the

interface of the two TMDs is closed. The structural elements that enable this transition

are conserved among ABC transporters as illustrated in the comparison between

corresponding intermediates of BmrCD and TmrAB (Supplementary Fig. 23). That a

population of the ATP-bound conformation is also inward-facing, both in detergent

micelles and lipid bilayers, suggests that energetics of BmrCD conformational

equilibrium deviate from TmrAB and MsbA, in which ATP binding triggers NBD

dimerization and extracellular gate opening[12,13,45]. Finally, the lack of an OF

conformation in the QQ background can be ascribed partially to the previously

demonstrated requirement for ATP hydrolysis to stabilize this intermediate[29]. However,

we were unable to capture this conformation in the WT background under turnover

conditions (not shown) or in the vanadate-trapped HES which may reflect the intrinsic

heterogeneity and/or instability of this conformation. This conclusion is consistent with

the relatively transient nature of this state inferred from multiple studies of ABC

transporters[18,46].

Figure 8 summarizes our BmrCD transport model, updated based on the results

presented in this study, that emphasizes the interplay between asymmetry and ligand

binding. BmrCD partitions into an ATP-powered but substrate-uncoupled cycle (Step 2).

Based on DEER data[29], an OF conformation is not populated significantly in this cycle in

lipid bilayers. Similar to Pgp and other ABC exporters, this cycle is characterized by

lower ATP turnover relative to the substrate-coupled cycle. Subsequent to Hoechst

binding, which we surmise occurs in two distinct steps as evidenced by two IF

conformations with one or two bound Hoechst, BmrCD partitions to the ATP- and

substrate-coupled cycle. Since BmrCD bound to ATP populates the IF conformation

(Step 1) and appears to bind at least one ATP at the degenerate NBS[31], we propose

that binding of the second ATP followed by sequential coordination by Mg^{2+} represent

potential rate limiting steps for triggering ATP hydrolysis (Steps 2, 3 and 4). In support

of this contention, we could only identify one Mg^{2+} in the cryo-EM structure of the ATP-

and Hoechst-bound IF conformation (Step 3, Supplementary Fig. 14). Finally, despite

the absence of the OF from the current structural records, the DEER data unequivocally

suggest the population of an OF conformation of BmrCD that has features similar to the

corresponding MsbA conformation[29]. Therefore, substrate extrusion out of the chamber

is executed at this step. We propose that the OC structure captured here represents a

return step from the OF state, since it is likely that the ATP at the consensus NBS is

already hydrolyzed (Step 4). Because previous DEER analysis indicated that population

of the OF conformation requires ATP hydrolysis[29], we expect that the return step in the

WT background to have ADP molecule in the consensus NBS.

The impact of lipid bilayers on energetics of the conformational cycle of ABC

transporters. Reconstitution of BmrCD into lipid bilayers attenuates the coupling

between ATP hydrolysis and isomerization to the OF conformation despite a 11-14-fold

enhancement of ATP hydrolysis rate[29]. It was speculated that this could reflect a

"coupling leak" wherein the dimerization of the NBDs is not efficiently coupled to the

TMD. The mechanism responsible for the loss of efficiency, however, was not

described.

The combination of cryo-EM visualization of lipids and MD simulations to triage non-

specifically bound lipids enables the formulation of a plausible hypothesis to explain this

observation. More lipid molecules stably bind to the IF conformation relative to OC.

Importantly, we observed stably-bound lipids at the interface between the two TMD

leaflets that would have to be displaced for the transition to OC or OF conformations.

For example, Lipids 3 and 9 bind to the interface of the TMDs in the IF conformation

within 400 ns and 300 ns in the MD simulations, respectively, whereas just one lipid

(Lipid 16) is stably bound at the interface in the OC conformation (Supplementary Fig.

22a, b, and c). A wide IF conformation has been reported for multiple ABC

transporters[11,12,15,31,47-50] which could imply that lipid interactions and consequent

modulation of the energetics of conformational changes may be a ubiquitous element

for these transporters. While a more detailed analysis of the role of lipids, including the

headgroups and the chain length, in the mechanism of ABC exporters is yet to be fully

investigated, the results presented here suggest a critical role for lipid-transporter

interactions.

Structural asymmetry in the transport cycle of ABC exporters. ABC heterodimers

with a catalytically impaired NBS has presented challenges to existing models derived

from investigations of canonical ABC exporters with two active NBSs. Specifically, it

raised questions regarding the role of the impaired NBS in priming the transporter and

how asymmetry is propagated between the TMD and NBD. The findings reported here

may have more general implications to type IV ABC exporters and possibly across ABC

exporters. Our previous DEER investigation of the ATP- and substrate-coupled

conformational cycle of the mammalian ABC exporter Pgp, which were partly the

impetus of the systematic study of BmrCD, revealed that asymmetric conformations at

the NBSs are differentially stabilized by substrate and inhibitors suggesting a critical role

of these conformations in the transport cycle[22,23]. Whereas BmrCD has intrinsically

catalytically inequivalent NBSs, we found that structural asymmetry, stabilized by

substrate binding in the TMD, propagates to the NBSs where it results in exclusive ATP

hydrolysis at the consensus NBS. We surmise that differential interactions of ATP and

Mg^{2+} with the NBSs, as visualized in our structures (Supplementary Fig. 12), provide a

basis for the catalytic inequivalence. Despite the QQ background, it is remarkable that

only the side chain of the consensus catalytic residue flips in order to coordinate the

nucleophilic attack on the ATP γ-phosphate. Thus, the cryo-EM structures presented

here build the notion that such asymmetry may be a conserved aspect of ABC exporter

transport cycles. Further investigations of ABC exporters in lipid bilayers without

functionally impairing mutations will be required to further illuminate this aspect of the

transport mechanism.

Methods

Cloning, expression, and purification of BmrCD. Wild type BmrCD in pET21b(+) was

used as the original template to generate the cysteine-less BmrCD (BmrCD*) and

D500Q in BmrC, E592Q in BmrD contained BmrCD* (BmrCD*-QQ) as described

previously[11,29]. Briefly, three native cysteines, C154, C256, and C351, in BmrD were

substituted with alanine using site-directed mutagenesis to create cysteine-less BmrCD.

To impair the ATP turnover of BmrCD, the catalytic residues D500 in BmrC and E592 in

BmrD were mutated into glutamine by site-directed mutagenesis.

WT BmrCD and BmrCD*-QQ plasmids were transformed into *Escherichia coli*

BL21(DE3) cells and the transporter was expressed in minimal media supplemented

with glycerol (0.5% v/v), thiamin (2.5 µg/ml), ampicillin (100 µg/ml), MgSO4 (1 mM) and

50× MEM amino acids. Cell cultures were grown at 37°C until optical density of OD600

reached approximately 1.2, and then the temperature was decreased to 25°C with 0.7

isopropyl β-d-1-thiogalactopyranoside (IPTG) added to induce BmrCD expression. After

5 h expression, the cells were harvested and frozen at -80°C. Cell pellets were

resuspended in lysis buffer (50 mM Tris-HCl, pH 8.0, 1 mM EDTA, 1 mM PMSF, and a

complete EDTA-free protease inhibitor cocktail tablet (Roche). After sonication, the

lysate was centrifuged at 9000g for 10 min at 4°C, and supernatant was used for

another round of centrifugation at 40,000 rpm for 1 h at 4°C to get the membrane

pellets. The membrane pellets were solubilized in resuspension buffer (50 mM Tris-HCl,

pH 8.0, 0.1 M NaCl, 15% (v/v) glycerol, 1 mM DTT and 1.25% w/v n-dodecyl-β-D-

maltopyranoside (β-DDM) for 1 h on ice. The sample was centrifugated at 40,000 rpm

for 1 h at 4°C, and the supernatant was mixed with the pre-equilibrated Ni-NTA beads

for about 2 h in binding buffer (50 mM Tris-HCl, pH 8.0, 0.1 M NaCl, 15% (v/v) glycerol,

0.05% β-DDM). The binding buffer supplemented with 20 mM and 250 mM imidazole

were used for resin wash and protein elution, respectively. Eluted fractions containing

BmrCD were further purified by size exclusion chromatography (SEC) using a Superdex

200 Increase 10/300 column (Cytiva) for further purification in the SEC buffer 1 (50 mM

Tris-HCl, pH 7.5, 0.15 M NaCl, 10% (v/v) glycerol, 0.05% β-DDM).

Nanodiscs preparation of BmrCD. The purified MSP and BmrCD in detergent were

reconstituted into nanodiscs as described previously[29]. Briefly, two lipid mixtures were

made. PC (L-α phosphatidylcholine) and PA (L-α phosphatidic acid) (Avanti Polar

Lipids, Alabaster, USA) were mixed as a 9:1 molar ratio or PC and *E. coli* polar lipids

(Avanti Polar Lipids, Alabaster, USA) were combined in a 1:3 molar ratio. Lipids were

then mixed with MSP and BmrCD micelles at a molar ratio of 1800:360:3:1 (β-DDM:

lipid: MSP: BmrCD). The mixtures were rotated for 30 min, and then biobeads (0.8-1

g/ml) were added to rotate overnight at 4°C to remove the detergent. The biobeads

were precipitated by centrifugation at 2000 rpm for 5 min and then removed completely

using a 0.45 μm filter. BmrCD in nanodiscs was separated by size exclusion

chromatography using a Superdex 200 Increase 10/300 column (Cytiva) with SEC

buffer 2 composed of 50 mM Tris-HCl, pH 8.0, 0.15 M NaCl (For cryo-EM) or SEC

buffer 3 composed of 50 mM Tris-HCl, pH 7.5, 0.15 M NaCl, 10% glycerol (For CW-

EPR and DEER).

Cryo-EM sample preparation and data acquisition. Reconstituted BmrCD*-QQ (~3

mg/ml) was incubated on ice for 30 min after supplementing with 10 mM ATP, 5 mM

MgCl$_2$, and with or without 1.2 mM Hoechst-33342 (Thermo Fisher Scientific). To

prepare vanadate trapped BmrCD sample, reconstituted wild type BmrCD (~3 mg/ml)

was incubated at 37°C for 15 min with 10 mM ATP, 4 mM MgSO$_4$, and 4 mM vanadate

before plunging. For cryo-EM grids preparation, 2.5 μl of ~3 mg/ml BmrCD was placed

onto the Quantifoil UltrAuFoil grids (R1.2/1.3, 300 mesh, Electron Microscopy Sciences)

which were glow discharged for 20 sec at 25 mA. Then the grids were blotted for 3 sec

at force 12 using two layers of filter papers and were plunge-frozen in liquid ethane by

Vitrobot Mark IV (Thermo Fisher) with the environmental chamber set to 4°C and 100%

humidity. Cryo-EM grids for data collection were selected after screening by TF20 or

Glacios (Thermo Fisher Scientific) .

Cryo-EM data of BmrCD_IF-2H/ATP were collected by Titan Krios G4 microscope

(Thermo Fisher Scientific) at 300 keV, equipped with Gatan BioQuantum filter, Gatan

K3 with GIF detector. Images were recorded with two exposures per hole using EPU in

super-resolution mode with a 20 eV slit width of energy filter and at a magnification of

105,000 with a pixel size of 0.818 Å/pixel. Defocus was set to vary from -0.8 to -2.2 μm.

Each image was dose fractionated to 50 frames . The total dose over the sample was

48.271 e$^-$/Å^2 and the dose was 0.965 e$^-$/Å^2/frame. 5,696 movies were collected for this

data set. Cryo-EM data of BmrCD_OC-ATP was collected using a similar condition

except with a higher magnification of 130,000 with a pixel size of 0.647 Å/pixel. The

entire dataset contains 11,829 movies.

Cryo-EM data of BmrCD_IF-ATP was imaged with a 300 keV FEI Krios G3i

microscope equipped with a Gatan K3 direct electron. Movies containing 40-50 frames

were collected at a magnification of 81,000 in super-resolution mode with a pixel size of

0.55 Å/pixel and defocus range of -0.8 to -1.6 μm using automated imaging software

EPU (Thermo Fisher Scientific).

Cryo-EM data processing. All datasets were processed in RELION 3.0[51] and

CryoSparc[42]. MotionCor2 and Gtcf were used to perform beam-induced motion

correction and CTF estimations, respectively. Images with poor quality were discarded.

For BmrCD_IF-2H/ATP dataset processing, auto picking implemented with

Laplacian-of-Gaussian method of Relion was used to pick around 3 million particles,

and extracted as 4 x 4 binned, through 6 rounds of 2D classification and one round of

3D classification using rescaled BmrCD map (EMD-23641)[11]. After 4 more rounds of 2D

classification, the particles were unbinned and one more 3D classification was carried

out to obtain a 4 Å map. Using this map as a template, we did template-based picking,

multiple rounds of 2D classification, and 2 rounds of 3D classification to obtain a class

with 133,280 particles. Then processing was implemented with CryoSparc 3.2.0[42],

performing NU-refinement and local refinement to yield a final map at 3.34 Å. To

improve the quality of the map, more rounds of local refinements were applied using

masks covering parts of the original map. Four separate masks that cover ECD, TMD,

BmrC-NBD, and BmrD-NBD were used (Supplementary Fig. 3).

For BmrCD_IF-ATP dataset processing, a similar processing procedure was

performed. Briefly, after several rounds of 2D classification and 3D classification, a

class containing 529,549 particles was transferred to CryoSparc 3.2.0[42] to do Ab-initio,

then a class was chosen to run NU-refinement and local refinement. Finally, a cryo-EM

map with 3.27 Å resolution was obtained. More rounds of local refinements for masks

on parts of the original map were also applied to improve the quality of the electron

density.

BmrCD_OC-ATP dataset processing was similar to BmrCD_IF-ATP dataset

processing except that this data set displayed anisotropy. To solve the anisotropy of the

map, we performed heterogeneous refinement. Then NU-refinement and local

refinement were performed to get the final map at 2.9 Å. More rounds of local

refinements for masks on ECD, TMD and closed NBD of the original map were also

applied to improve the quality of the electron density map. A similar processing

procedure is applied to obtain the map of BmrCD_IF-ATP2 in this dataset.

Dataset 2 and dataset 5 in Supplementary Fig. 1 were processed using the same

strategies as above (Supplementary Fig. 2).

Model building and figure preparation. The same pipeline for model building process

was performed for all the structures. The initial model was searched with dock_in_map

implemented in Phenix by separated domains from the BmrCD structure in detergent

micelles (PDB ID: 7m33)[11]. Combined with manual adjustment in coot[52] and refinement

by real-space refinement implemented in Phenix[53], a satisfactory model was built.

MolProbity implemented in Phenix[54] was used for all the structural model validation. For

BmrCD_IF-H/ADPVi in Supplementary Fig. 1, the model of BmrCD_IF-2H/ATP was

fitted to the density map in Chimera manually due to its low resolution.

Figures were prepared by UCSF Chimera[55], ChimeraX[56], and PyMol. Solvent-

accessible volume calculated by CASTp[40].

Molecular dynamics simulations

Equilibrium simulations. Initial placement of BmrCD in the membrane and

arrangement of lipids around it were done by the CHARMM-GUI webserver[43,57] using

the published structure (PDB ID: 7M33[11]) as a proxy for our new Cryo-EM models.

Partially resolved Cryo-EM lipids were classified as POPC or POPA depending on the

resolved head group atoms. Lipids with resolved Nitrogen and Phosphate atoms were

labeled as POPC, and those with resolved P atoms only were labeled as POPA. Lipid

tails without any headgroup atoms were excluded from the modeling process. Missing

atoms in the cryo-EM lipids were completed using the PSFGEN plugin in VMD[58].

Missing atoms of the protein were completed with MOE (Chemical Computing Group,

Montreal, Canada). The protonation states of the residues were assigned using

PropKa[59]. Completed cryo-EM lipids were combined with the protein, Hoechst-

33342and two ATP molecules preserving their position in the cryo-EM structure. Upon

visual inspection, any lipid tails modeled artefactually penetrating the protein core were

rotated around one of the C-C bonds to achieve a better initial position and orientation

without changing the positions of the cryo-EM resolved atoms. The protein structure

was placed into a bilayer using the CHARMM-GUI web server and cryo-EM lipids were

retained in their position[43,57]. Any bulk membrane lipids within 0.5 Å of either the protein

or the completed cryo-EM lipids were removed to prevent clashing. This model was

then re-ionized to ensure charge neutrality using the AUTOIONIZE plugin in VMD.

Parameters for the Hoechst-33342 ligand present in the open conformation are

previously reported in our past work[11]. Replica simulations were generated using the

Membrane Mixer tool in VMD[60] to better sample the initial lipid distribution for the non-

modeled lipids.

Each simulation system was energy-minimized for 10,000 steps followed by a gentle

heating scheme of incrementing the system temperature by 25 K for 100 ps at a time

until the target temperature of 310 K was achieved. Equilibration was performed using

NAMD2.14[61,62] for 7.5 ns with restraints following this scheme: 2.5 ns with protein

backbone and lipid headgroups restrained at $1 kcal/mol/Å^2$, 2.5 ns with just protein

backbone restrained at $0.5 kcal/mol/Å^2$ and a final 2.5 ns equilibration with protein α-

carbons restrained at $0.5 kcal/mol/Å^2$ to allow the lipid tails to melt and the protein side

chains to relax. Restrained equilibration was then followed by 5 ns of simulation without

restraints during which pressure coupling was turned on. To allow modeled lipid tails to

blend into the membrane, a grid-based potential was applied using the Grid Forces[63]

option of NAMD during the first phase of equilibration. Production MD was then carried

out for 500 ns in each replica for a total of 3 µs of sampling across both conformations

and all 3 replicas.

In silico **mutagenesis**. K2, K5 and R15 were sites of the BmrD elbow helix identified

as participating in highly resident lipid-binding. Each site was individually mutated to an

alanine using the Mutagenesis tool in VMD and system charge once again corrected

using the AUTOIONIZE plugin. Three replicas were generated for each mutant for a

total of 18 simulations (3 replicas x 3 mutants x 2 conformations). Each replica was

treated as before with a slow equilibration and 500 ns of production MD.

DEER sample preparation, CW-EPR and DEER spectroscopy. To prepare samples

for continuous-wave electron paramagnetic resonance spectroscopy (CW-EPR) and

double electron-electron resonance (DEER), BmrCD double-cysteine mutants eluted

from Ni-NTA affinity purification were labeled with 20-fold molar excess of (1-Oxyl-

2,2,5,5-tetramethylpyrroline-3-methyl)methanethiosulfonate (MTSSL, Enzo Life

Science) at room temperature in dark for 2 h. Followed by labeled with another 20-fold

molar excess of MTSSL at room temperature in dark for 2 h and then moved to 4 °C

overnight. The labeled BmrCD samples separated from the free label by Superdex 200

Increase 10/300 column (Cytiva) in SEC buffer 1 for CW-EPR and DEER in detergent

buffer. Reconstitution and a size size-exclusion chromatography in SEC buffer 3 were

applied to prepare nanodisc samples for CW-EPR and DEER.

Spin-labeled BmrCD mutants were incubated with nucleotides and/or Hoechst-33342

(Thermo Fisher) for CW-EPR and DEER. Different combinations were used to capture

different states of BmrCD with final concentrations of ATP, vanadate, AMP-PNP,

Hoechst-33342, and MgSO4 at 10 mM, 5 mM, 10 mM, 1 mM, 10 mM, respectively. For

high-energy post-hydrolysis state, BmrCD in detergent micelles and in nanodiscs were

incubated with ATP, vanadate and MgCl2 at 30 °C for 30 min and 37 °C for 30 min,

respectively. To trap an ATP-bound state, BmrCD in detergent micelles and in

nanodiscs were incubated with AMP-PNP and MgSO4 at 30 °C for 30 min and 37 °C for

30 min, respectively.

For CW-EPR experiments, spin-labeled BmrCD samples were loaded in 25 µl

capillaries and spectra were collected on a Bruker EMX spectrometer using 10 mW

microwave power level and a modulation amplitude of 1.6 G.

For DEER spectroscopy, a final concentration of 24% (v/v) glycerol was used as a

cryoprotectant for all DEER experiments. An Elexsys E580 pulsed EPR spectrometer

operating at Q-band frequency (33.9 GHz) with a 40W Amp-Q amplifier (Bruker) with

the dead-time free four-pulse sequence at 83 K was used[64,65]. We used the pulse

lengths of 10 ns (p/2) and 20 ns (p) for the probe pulses, and 40 ns for the pump pulse,

separately. And the frequency separation was set as 73 MHz. Raw DEER decays were

analyzed as described previously[11,29]. Briefly, a home-written software operating in the

Matlab environment was used to analyze the raw DEER. For the same spin-labeled

position under different conditions, global analysis of the DEER decays was applied. We

assumed that the distance distribution consists of a sum of Gaussians and used a

statistical criterion to determine the number and population of the distance distribution

as previously described[29].

Distance distribution predictions based on BmrCD cryo-EM structures were carried

out by the DEER Spin-Pair Distributor at the CHARMM-GUI website[43,66] with default

parameters in 10 ns MD simulations with dummy spin labels that mimics the dynamics

of MTS spin-label.

Data availability

The three cryo-EM 3D density maps of the heterodimeric ABC transporter BmrCD in

nanodiscs in Supplementary Fig. 1 have been deposited in the Electron Microscopy

Data Bank under accession codes EMD-29297, EMD-40908, EMD-29362, EMD-41004,

EMD-29087, EMD-40974 and EMD-41058 (see Supplemental Table 1). Atomic

coordinates for the atomic models have been deposited in the Protein Data Bank under

accession codes 8FMV, 8SZC, 8FPF, 8T3K, 8FHK, and 8T1P.

References

1.   Beek, J., Guskov, A. & Slotboom, D.J. Structural diversity of ABC transporters. *J*

Gen Physiol **143**, 419-35 (2014).

2.   Rees, D.C., Johnson, E. & Lewinson, O. ABC transporters: the power to change.

Nat Rev Mol Cell Biol **10**, 218-27 (2009).

3.   Locher, K.P. Mechanistic diversity in ATP-binding cassette (ABC) transporters.

Nat Struct Mol Biol **23**, 487-93 (2016).

4.   Thomas, C. & Tampe, R. Structural and Mechanistic Principles of ABC

Transporters. *Annu Rev Biochem* **89**, 605-636 (2020).

5.   Higgins, C.F. & Linton, K.J. The ATP switch model for ABC transporters. *Nat*

Struct Mol Biol **11**, 918-26 (2004).

6.   Shvarev, D., Januliene, D. & Moeller, A. Frozen motion: how cryo-EM changes

the way we look at ABC transporters. *Trends in Biochemical Sciences* **47**, 136-

148 (2022).

7.   Gebhard, S. ABC transporters of antimicrobial peptides in Firmicutes bacteria -

phylogeny, function and regulation. *Mol Microbiol* **86**, 1295-317 (2012).

8.   Ye, Z., Lu, Y. & Wu, T. The impact of ATP-binding cassette transporters on

metabolic diseases. *Nutr Metab (Lond)* **17**, 61 (2020).

9.   Schumacher, T. & Benndorf, R.A. ABC Transport Proteins in Cardiovascular

Disease-A Brief Summary. *Molecules* **22**(2017).

10.   Dean, M., Moitra, K. & Allikmets, R. The human ATP-binding cassette (ABC)

transporter superfamily. *Human Mutation* **43**, 1162-1182 (2022).

11.   Thaker, T.M. et al. Asymmetric drug binding in an ATP-loaded inward-facing

state of an ABC transporter. *Nat Chem Biol* **18**, 226-235 (2022).

12.   Hofmann, S. et al. Conformation space of a heterodimeric ABC exporter under

turnover conditions. *Nature* **571**, 580-583 (2019).

13.   Mi, W. et al. Structural basis of MsbA-mediated lipopolysaccharide transport.

Nature **549**, 233-237 (2017).

14. Alam, A., Kowal, J., Broude, E., Roninson, I. & Locher, K.P. Structural insight into

substrate and inhibitor discrimination by human P-glycoprotein. *Science* **363**,

753-+ (2019).

15. Jin, M.S., Oldham, M.L., Zhang, Q. & Chen, J. Crystal structure of the multidrug

transporter P-glycoprotein from Caenorhabditis elegans. *Nature* **490**, 566-9

(2012).

16. Oldham, M.L., Davidson, A.L. & Chen, J. Structural insights into ABC transporter

mechanism. *Current Opinion in Structural Biology* **18**, 726-733 (2008).

17. Lewinson, O., Orelle, C. & Seeger, M.A. Structures of ABC transporters: handle

with care. *FEBS Lett* **594**, 3799-3814 (2020).

18. Wang, L. et al. Characterization of the kinetic cycle of an ABC transporter by

single-molecule and cryo-EM analyses. *Elife* **9**(2020).

19. Seeger, M.A. & van Veen, H.W. Molecular basis of multidrug transport by ABC

transporters. *Biochim Biophys Acta* **1794**, 725-37 (2009).

20. Kim, Y. & Chen, J. Molecular structure of human P-glycoprotein in the ATP-

bound, outward-facing conformation. *Science* **359**, 915-919 (2018).

21. Thomas, C. et al. Structural and functional diversity calls for a new classification

of ABC transporters. *Febs Letters* **594**, 3767-3775 (2020).

22. Dastvan, R., Mishra, S., Peskova, Y.B., Nakamoto, R.K. & McHaourab, H.S.

Mechanism of allosteric modulation of P-glycoprotein by transport substrates and

inhibitors. *Science* **364**, 689-692 (2019).

23. Verhalen, B. et al. Energy transduction and alternating access of the mammalian

ABC transporter P-glycoprotein. *Nature* **543**, 738-741 (2017).

24.   Carrier, I., Julien, M. & Gros, P. Analysis of catalytic carboxylate mutants E552Q

and E1197Q suggests asymmetric ATP hydrolysis by the two nucleotide-binding

domains of P-glycoprotein. *Biochemistry* **42**, 12875-85 (2003).

25.   Tombline, G. et al. Properties of P-glycoprotein with mutations in the "catalytic

carboxylate" glutamate residues. *Journal of Biological Chemistry* **279**, 46518-

46526 (2004).

26.   Tombline, G. & Senior, A.E. The occluded nucleotide conformation of p-

glycoprotein. *J Bioenerg Biomembr* **37**, 497-500 (2005).

27.   Toth, A., Janaszkiewicz, A., Crespi, V. & Di Meo, F. On the interplay between

lipids and asymmetric dynamics of an NBS degenerate ABC transporter.

Communications Biology **6**(2023).

28.   Chen, M., Abele, R. & Tampe, R. Functional non-equivalence of ATP-binding

cassette signature motifs in the transporter associated with antigen processing

(TAP). *J Biol Chem* **279**, 46073-81 (2004).

29.   Mishra, S. et al. Conformational dynamics of the nucleotide binding domains and

the power stroke of a heterodimeric ABC transporter. *Elife* **3**, e02740 (2014).

30.   Hohl, M. et al. Structural basis for allosteric cross-talk between the asymmetric

nucleotide binding sites of a heterodimeric ABC exporter. *Proceedings of the*

National Academy of Sciences of the United States of America **111**, 11025-

11030 (2014).

31.   Hohl, M., Briand, C., Grutter, M.G. & Seeger, M.A. Crystal structure of a

heterodimeric ABC transporter in its inward-facing conformation. *Nat Struct Mol*

Biol **19**, 395-402 (2012).

32.   Stockner, T., Gradisch, R. & Schmitt, L. The role of the degenerate nucleotide

813   binding site in type I ABC exporters. *FEBS Lett* **594**, 3815-3838 (2020).

814   33.   Reilman, E., Mars, R.A., van Dijl, J.M. & Denham, E.L. The multidrug ABC

815   transporter BmrC/BmrD of Bacillus subtilis is regulated via a ribosome-mediated

816   transcriptional attenuation mechanism. *Nucleic Acids Res* **42**, 11393-407 (2014).

817   34.   Dezi, M., Di Cicco, A., Bassereau, P. & Levy, D. Detergent-mediated

818   incorporation of transmembrane proteins in giant unilamellar vesicles with

819   controlled physiological contents. *Proc Natl Acad Sci U S A* **110**, 7276-81 (2013).

820   35.   Collauto, A., Mishra, S., Litvinov, A., McHaourab, H.S. & Goldfarb, D. Direct

821   Spectroscopic Detection of ATP Turnover Reveals Mechanistic Divergence of

822   ABC Exporters. *Structure* **25**, 1264-1274 e3 (2017).

823   36.   Choudhury, H.G. et al. Structure of an antibacterial peptide ATP-binding cassette

824   transporter in a novel outward occluded state. *Proc Natl Acad Sci U S A* **111**,

825   9145-50 (2014).

826   37.   Ward, A., Reyes, C.L., Yu, J., Roth, C.B. & Chang, G. Flexibility in the ABC

827   transporter MsbA: Alternating access with a twist. *Proc Natl Acad Sci U S A* **104**,

828   19005-10 (2007).

829   38.   Dawson, R.J. & Locher, K.P. Structure of the multidrug ABC transporter Sav1866

830   from Staphylococcus aureus in complex with AMP-PNP. *FEBS Lett* **581**, 935-8

831   (2007).

832   39.   Johnson, Z.L. & Chen, J. ATP Binding Enables Substrate Release from Multidrug

833   Resistance Protein 1. *Cell* **172**, 81-89 e10 (2018).

40. Tian, W., Chen, C., Lei, X., Zhao, J. & Liang, J. CASTp 3.0: computed atlas of

surface topography of proteins. *Nucleic Acids Res* **46**, W363-W367 (2018).

41. Ritchie, T.K. et al. Chapter 11 - Reconstitution of membrane proteins in

phospholipid bilayer nanodiscs. *Methods Enzymol* **464**, 211-31 (2009).

42. Punjani, A., Rubinstein, J.L., Fleet, D.J. & Brubaker, M.A. cryoSPARC:

algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods*

14, 290-296 (2017).

43. Jo, S., Kim, T., Iyer, V.G. & Im, W. CHARMM-GUI: a web-based graphical user

interface for CHARMM. *J Comput Chem* **29**, 1859-65 (2008).

44. Zhang, Y. & Skolnick, J. TM-align: a protein structure alignment algorithm based

on the TM-score. *Nucleic Acids Res* **33**, 2302-9 (2005).

45. Dong, J., Yang, G. & McHaourab, H.S. Structural basis of energy transduction in

the transport cycle of MsbA. *Science* **308**, 1023-8 (2005).

46. Bountra, K. et al. Structural basis for antibacterial peptide self-immunity by the

bacterial ABC transporter McjD. *EMBO J* **36**, 3062-3079 (2017).

47. Johnson, Z.L. & Chen, J. Structural Basis of Substrate Recognition by the

Multidrug Resistance Protein MRP1. *Cell* **168**, 1075-1085 e9 (2017).

48. Thelot, F.A. et al. Distinct allosteric mechanisms of first-generation MsbA

inhibitors. *Science* **374**, 580-585 (2021).

49. Manolaridis, I. et al. Cryo-EM structures of a human ABCG2 mutant trapped in

ATP-bound and substrate-bound states. *Nature* **563**, 426-430 (2018).

50. Nosol, K. et al. Cryo-EM structures reveal distinct mechanisms of inhibition of the

human multidrug transporter ABCB1. *Proc Natl Acad Sci U S A* **117**, 26245-

26253 (2020).

51. Zivanov, J. et al. New tools for automated high-resolution cryo-EM structure

determination in RELION-3. *Elife* **7**(2018).

52. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development

of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).

53. Afonine, P.V. et al. Real-space refinement in PHENIX for cryo-EM and

crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).

54. Williams, C.J. et al. MolProbity: More and better reference data for improved all-

atom structure validation. *Protein Sci* **27**, 293-315 (2018).

55. Pettersen, E.F. et al. UCSF Chimera--a visualization system for exploratory

research and analysis. *J Comput Chem* **25**, 1605-12 (2004).

56. Pettersen, E.F. et al. UCSF ChimeraX: Structure visualization for researchers,

educators, and developers. *Protein Sci* **30**, 70-82 (2021).

57. Wu, E.L. et al. CHARMM-GUI Membrane Builder toward realistic biological

membrane simulations. *J Comput Chem* **35**, 1997-2004 (2014).

58. Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J Mol*

Graph **14**, 33-8, 27-8 (1996).

59. Rostkowski, M., Olsson, M.H., Sondergaard, C.R. & Jensen, J.H. Graphical

analysis of pH-dependent properties of proteins predicted using PROPKA. *BMC*

Struct Biol **11**, 6 (2011).

60. Licari, G., Dehghani-Ghahnaviyeh, S. & Tajkhorshid, E. Membrane Mixer: A

Toolkit for Efficient Shuffling of Lipids in Heterogeneous Biological Membranes. *J*

Chem Inf Model **62**, 986-996 (2022).

61. Phillips, J.C. et al. Scalable molecular dynamics with NAMD. *J Comput Chem* **26**,

1781-802 (2005).

62. Phillips, J.C. et al. Scalable molecular dynamics on CPU and GPU architectures

with NAMD. *J Chem Phys* **153**, 044130 (2020).

63. Wells, D.B., Abramkina, V. & Aksimentiev, A. Exploring transmembrane transport

through alpha-hemolysin with grid-steered molecular dynamics. *Journal of*

Chemical Physics **127**(2007).

64. Jeschke, G. & Polyhach, Y. Distance measurements on spin-labelled

biomacromolecules by pulsed electron paramagnetic resonance. *Physical*

Chemistry Chemical Physics **9**, 1895-1910 (2007).

65. Jeschke, G. DEER Distance Measurements on Proteins. *Annual Review of*

Physical Chemistry, Vol 63 **63**, 419-446 (2012).

66. Qi, Y. et al. CHARMM-GUI DEER facilitator for spin-pair distance distribution

calculations and preparation of restrained-ensemble molecular dynamics

simulations. *J Comput Chem* **41**, 415-420 (2020).

Acknowledgements

This work was supported by NIH grant GM128087 to Hassane Mchaourab. EM data

collections were conducted at the Center for Structural Biology Cryo-EM Facility at

Vanderbilt University. We acknowledge the use of the Glacios cryo-TEM, which was

acquired by NIH grant S10 OD030292-01. EM data collection for BmrCD_IF-ATP was

conducted at Case Western Reserve University by Dr. Kunpeng Li. Molecular

simulations were performed on resources provided by the XSEDE grant MCA06N060,

and supported by NIH grants P41-GM104601, R24-145965, and R01-GM123455.

Author contributions

Q.T. performed all the experiments and solved the structures. M.S. and H.H.

performed the MD simulation under the supervision of E.T. E.K. provided advice for

structure determination. R.S. helped with DEER data collection. Q.T. and H.M. analyzed

and interpreted the structures and wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

BmrCD_IF-ATP
3.3 Å

BmrCD_IF-H/ATP
3.1 Å

BmrCD_IF-2H/ATP
3.3 Å

BmrCD_OC-ATP
2.9 Å

Fig. 1 Representative BmrCD structures in lipid bilayers.

a BmrCD inward-facing (IF) conformation bound to ATP. **b** BmrCD IF bound to one

substrate Hoechst (HT) and two ATP molecules. **c** BmrCD IF conformation bound to

two Hoechsts and two ATP molecules. **d** BmrCD occluded (OC) conformation bound to

ATP. The top and bottom rows show experimental maps and deposited models,

respectively. BmrC is shown in light sea green, BmrD is in tan, Hoechst and ATP are

shown in magenta, and lipids are blue. In the models, BmrCD is shown in cartoon

representation, and all the ligands are represented by sticks. Mg^{2+} ions are omitted.

Fig. 2 Comparison of IF conformations with one or two bound Hoechst revealed that binding involve two latches in the substrate pocket.

a Side views of the electrostatic surface of the substrate Hoechst-binding pocket of

BmrCD_IF-2H/ATP (left panel) and BmrCD_IF-H/ATP (right panel). The view is the

same as the square box in Fig. 2b. The surfaces are colored from blue (positively

charged regions) to red (negatively charged regions). Neutral regions are shown as

white. BmrCD is shown in cartoon and Hoechst (HT, magenta color) is shown in stick.

The dash black lines indicate two halves of the chamber accommodating two bound

Hoechst molecules. **b** The two latches locations are shown in BmrCD_IF-2H/ATP in the

middle panel. The WH-latch is highlighted on in the left panel, and the RH-latch is

highlighted on in the right panel. **c** WH-latch analysis. The electron density maps for

WH-latches in BmrCD_IF-2H/ATP and BmrCD_IF-H/ATP are shown in left and middle

panels with the side-chain distances highlighted. The electron density maps for WH-

latches in BmrCD_IF-ATP is shown in right panel. **d** Analysis of the RH-latch in the

absence and presence of one or two Hoechst molecules. The electron density maps for

RH-latches in BmrCD_IF-2H/ATP (left panel), BmrCD_IF-H/ATP (middle panel) and

BmrCD_IF-ATP (right panel) are shown with the side-chain distance highlighted.

Fig. 3 Conformational changes underlying the IF to OC transition.

a Superposition of BmrCD_IF-2H/ATP and BmrCD_OC-ATP based on the alignment of

BmrD. BmrC and BmrD in the OC conformation are shown in light sea green and tan,

respectively while BmrCD_IF-2H/ATP is shown in grey. TMs 1,2/3 and 6 of BmrC and

4/5 of BmrD undergo the largest shifts between the two conformations. Ligands are

omitted in both structures for clarity. **b** Close up views of BmrCD highlighting spin label

pairs 55[BmrC]/146[BmrD] and 96[BmrC]/188[BmrD]. BmrC and BmrD are shown in light sea green

and tan in BmrCD_OC-ATP, respectively. TMD helices are shown as cylinders and

ECD and NBDs are shown as transparent surfaces. The spin label locations are

highlighted by the spheres. **c** and **d** DEER distance measurements for spin-label pairs

on the intracellular (96[BmrC]/188[BmrD]) and extracellular (55[BmrC]/146[BmrD]) sides in WT and

QQ backgrounds respectively. Experimental distance distributions representing the

probability of a distance $P(r)$ versus the distance (r) between spin labels, are shown in

963   solid lines: H (refers to Hoechst, black) and H-ADP-Vi (refers to ADP-vanadate

964   hereafter) (red). The simulated distance distributions derived from cryo-EM structures

965   by MDDS are shown in dashed lines: IF-2H/ATP (blue), IF-ATP (black), and OC-ATP

966   (red).

Fig. 4 DEER distance distributions for spin-labeled pairs monitoring the NBSs.

a Closed-up view of the NBDs of BmrCD cryo-EM structures (IF-H/ATP and OC-ATP)

showing the spin label pairs in the consensus and degenerate NBSs respectively. The

spin label pairs are highlighted by raspberry spheres. The side chain of general catalytic

residues and ATP are represented by sticks. **b** Experimental (solid lines) and predicted

(dashed lines) distance distributions highlighting the asymmetry of the NBSs in the OC

conformation.

Fig. 5 Asymmetric TMDs arrangements underpin the conformational cycle.

Upper panels: Distances between alpha carbons in the TMDs of BmrC and BmrD

generated by TM-align following superposition of the two TMD. The BmrCD structure in

detergent micelles is shown for reference. Lower panels: cartoon representation of the

superimposed TM helices from BmrC (colored to match the upper panels) and BmrD

(grey). A transparent surface representation of BmrD is shown as reference.

Fig. 6 Movement of the ECD opens a putative path for substrate efflux.

a Superimposition of ECD based on the alignment of BmrC in the BmrCD cryo-EM

structures highlighting the movement of the representative loops. **b** Electrostatics

potential surface of ECD in BmrCD cryo-EM structures shown from an extracellular

view. The color scale ranges from blue (positively charged regions) to red (negatively

charged regions) whereas neutral regions are shown as white. The positive charged

patch which possibly blocks the substrate exit in the IF structures is highlighted in a

black circle, a corresponding patch in BmrCD_OC-ATP with neutral charge is also

highlighted with a black circle for comparison. **c** Cartoon representation showing the

three spin label pairs between ECD and TMD. The spin label pairs are illustrated by

raspberry spheres connected by a straight line. **d** Experimental (solid lines) and

predicted (dashed lines) DEER distance distributions highlighting the movement of the

ECD.

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

Fig. 7 Stably bound lipids identified by MD simulations.

a and **b** Kernel density estimation performed on lipid phosphorus atoms within 20 Å of

BmrCD IF (BmrCD_IF-2H/ATP) and OC (BmrCD_OC-ATP) conformations. Initial

position of cryo-EM modeled lipid phosphorus atoms is marked in teal. The regions with

oval bright yellow densities highlight stable lipid binding. The regions with smeared

densities indicate nearby mobile lipids which do not exhibit stable binding. **c** Protein-lipid

residence times for surface basic residues binned into a box-and-whisker plot.

Prevalence of stable lipids in the IF conformation is highlighted by larger mean binding

residence time. **d** Angle space distribution for each surface basic residue for both

1037   conformations. Key residues such as K2, R180, and R328 have a tighter distribution in

1038   the IF and a broader more flexible distribution for the OC conformation.

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

Fig. 8 Model of BmrCD transport cycle.

BmrC and BmrD are shown in the same color as Figure 1. Number refers to the four

steps of BmrCD transport cycle: 1→2: ATP binding and the substrate uncoupled cycle;

2→3: Hoechst and the first Mg^{2+} binding; 3→4: the second Mg^{2+} binding, ATP hydrolysis

and Hoechst release; 4→1: ADP and Pi release which resets the transport cycle. Arrow

near the ECD refers the ECD twists to open/close the exit. Red box represents the cryo-

EM structures reported here. Black box are conformations deduced from cryo-EM 2D

classification or DEER data. ATP, ADP, and substrate Hoechst are represented by red

filled "A" circle, purple filled "A" circle, and blue filled "H" trapezoid, respectively.

Released phosphate and Mg^{2+} are shown in black and green dot, respectively. The two

"H" trapezoid represents the antiparallel arrangement of Hoechst in the chamber.

1060

1061 **Table 1. Conditions and Ligands of BmrCD cryo-EM structures.**

	Dataset	MSP	Lipids (No.)	Hoechst	ATP	Mg^{2+}	Construct
BmrCD_IF-2H/ATP	1	MSP1D1	PC/PA (20)	2	2	1	BmrCD*-QQ
BmrCD_IF-H/ATP	2	MSP1D1	PC/E.coli polar lipid (21)	1	2	1	
BmrCD_IF-ATP	3	MSP1D1E3	PC/PA (25)	0	2	0	
BmrCD_IF-ATP2	4	MSP1D1	PC/E.coli polar lipid (13)	0	2	0	
BmrCD_OC-ATP	4	MSP1D1	PC/E.coli polar lipid (16)	0	2	2	
BmrCD_OC-ADPVi	5	MSP1D1	PC/PA (21)	0	2	2	BmrCD-WT

1062

1063