1

Cl⁻-dependent amplification of excitatory synaptic potentials

at distal dendrites revealed by voltage imaging

Authors

Masato Morita,[1] Reo Higashi,[1] Shin-ya Kawaguchi[1]*

Affiliations

[1] Department of Biophysics, Graduate School of Science, Kyoto University, Kitashirakawa Oiwake-cho,

Sakyo-ku, Kyoto 606-8502, Japan.

* Corresponding author: kawaguchi.shinya.7m@kyoto-u.ac.jp

Abstract

Processing of synaptic signals in somatodendritic compartments determines the neuronal computation. Although amplification of excitatory signals by local voltage-dependent cation channels has been extensively studied, its spatio-temporal dynamics in elaborate dendritic branches remains obscure because of technical limitation. Using fluorescent voltage imaging throughout dendritic arborizations in hippocampal pyramidal neurons, here we demonstrate a unique Cl^--dependent remote computation mechanism equipped in distal branches. Local laser photolysis of caged-glutamate triggered excitatory postsynaptic potentials spreading along dendrites with gradual amplification toward the distal end whereas with attenuation toward the soma. *Tour-de-force* subcellular patch-clamp recordings from thin branches complemented with biophysical model simulation revealed that the asymmetric augmentation of excitation relies on the TTX-resistant Na^+ channels and Cl^--conductances accompanied with deeper dendritic resting potential. Taken together, the present study unveils cooperative voltage-dependent actions of cation and anion conductances for dendritic supralinear computation which can locally decode the spatio-temporal context of synaptic inputs.

Introduction

Neurons extend elaborate dendrites to receive synaptic inputs from other neurons. These synaptic inputs are spatially and temporally integrated in an analogue manner, and then the signals are transformed into action potentials (APs) as digital all-or-none signals at the axon initial segment. It has been shown that hippocampal and cortical pyramidal neurons exhibit several types of regenerative local spikes upon synaptic inputs in dendrites, including Na^+, NMDA, and Ca^{2+} spikes (1, 2). Na^+ and NMDA spikes are triggered by local multiple synaptic inputs in dendritic branches (3-6). On the other hand, Ca^{2+} spikes mediated by voltage-gated Ca^{2+} channels, show larger and broader waveforms, often leading to burst APs (1, 7, 8). In contrast to such a dendritic supra-linear computation mediated by active conductances, sub-threshold potential changes are thought to obey the passive membrane properties of dendrites, attenuating during the propagation as the cable theory predicts (9, 10). However, at present, it remains elusive how the subthreshold potential changes are processed in dendrite because of the difficulty to apply the patch-clamp technique to multiple points of a thin dendrite at the same time. As a corollary, mathematical model of electrical circuit assuming multiple compartments has provided basic view of signal integration in dendrites (11, 12), substantial part of which still remains to be consolidated by direct experimental analysis.

In this decade, genetically-encoded voltage indicators (GEVIs), which detect membrane potential changes as fluorescence changes, have drastically developed (13-15). This technique has enabled to fluorescently detect membrane potential changes at a tiny structure which is difficult to apply the patch-clamp technique, for example even a dendritic spine and a presynaptic bouton (16, 17). Taking advantage of the leading-edge voltage imaging technique, in this study, we studied how subthreshold synaptic potentials are processed in dendrites. Using improved version of ASAP1 (Accelerated Sensor of Action Potentials 1) (13), a fluorescent voltage imaging probe, expressed in a cultured hippocampal neuron in combination with very local glutamate or GABA stimulation on a dendrite by 405 nm spot-laser-mediated

photoactivation, we found that even small size of glutamatergic excitatory postsynaptic potentials (EPSPs) spread in dendrites with opposite modulation depending on the direction of propagation. We also identified a novel Cl⁻-dependent mechanism underlying the unique asymmetric processing of the electrical signal in dendrites.

Results

Fluorescent imaging of membrane potential changes in a cultured hippocampal neuron

In this study, for better fluorescent detection of membrane potential changes in neurons, some amino acid replacements reported in an improved version ASAP3 (18) were introduced into the original ASAP1 (13) (here we note this version ASAP3β; for details, see methods). We evaluated fluorescence change of ASAP3β upon voltage changes by expressing in HEK293T cells (fig. S1A). ASAP3β exhibited fluorescence decrease (-40.0 ± 1.1 % ΔF/F, fig. S1, B and C) in response to 100 mV depolarization of the voltage-clamped HEK293T cell from a holding potential at -70 mV, which corresponded to almost two-fold sensitivity of the original ASAP1 (-18.5 ± 1.2 % ΔF/F, fig. S1C) similarly to ASAP3 (18).

We then examined how ASAP3β detects neuronal activity characterized by rapid membrane potential changes in a cultured hippocampal neuron by expressing it with electroporation (Fig. 1A). Both the fluorescence and the membrane potential were simultaneously recorded by imaging and patch-clamp methods. ASAP3β detected spontaneous EPSPs almost perfectly without any noticeable change in time course (Fig. 1B), showing ~ 1.3 % fluorescence change for 2 mV potential change around -70 mV of membrane potential (Fig. 1C). On the other hand, more rapid spontaneous or evoked APs were detected as smaller fluorescence changes relative to the actual change of membrane potential (-8.4 ± 0.4 % ΔF/F upon 93.6 ± 2.7 mV, Fig. 1, B and C), likely because of the insufficient velocity of ASAP3β to follow a

given rapid voltage change (Fig. 1, B and C). From these observations, it was confirmed that ASAP3β

nicely detects synaptic potentials.

Direction-dependent EPSP modulation in dendrites

Taking advantage of real-time fluorescent detection of local membrane potential changes of interest

throughout dendritic arborization, we studied how synaptic potentials propagate in dendrites.

Glutamatergic EPSPs (EPSP$_{glu}$) were evoked by local photo-activation of MNI-caged-glutamate (300 μM

in the external bath) by 405 nm laser illumination (2 μm diameter) at a spine of ASAP3β-expressing

hippocampal pyramidal neuron in culture (Fig. 1D). Simultaneous recordings of membrane potentials by

current-clamp and fluorescent imaging from an ASAP3β-expressing neuron showed similar waveforms

of EPSPs characterized by 1.4 ± 0.1 % (21 events) fluorescence change at the soma upon 2 mV potential

change without temporal lag (Fig. 1D). Thus, evoked EPSP (eEPSP) measurement by ASAP3β enabled

us to quantify the spatial pattern of EPSP spreading in a neuron.

Using this method, eEPSPs were monitored as the relative ASAP fluorescence changes (ΔF/F)

throughout the dendrites of ASAP3β-expressing hippocampal neurons in the view field (Fig. 1, E and F).

To focus on the propagation of EPSPs far from activation of typical voltage-gated active conductances,

such as Na^+ and Ca^{2+} channels, the laser intensity and duration was adjusted so that the EPSP size was

around 2 - 5 mV (~ 1 - 3 % of ΔF/F in ASAP fluorescence change). In line with the prediction of the cable

theory (11), the fluorescence change for eEPSP (ΔF/F$_{eEPSP}$) attenuated as propagation toward the soma

(Fig. 1, E and G). The ΔF/F$_{eEPSP}$ decreased by 25 ± 2 % at 33.6 ± 3.1 μm away from the uncaged site and

by 59 ± 3 % at 97.7 ± 4.9 μm (Fig. 1H, black circles). Thus, the propagation of EPSPs toward the soma

attenuates almost exponentially as characterized by a length constant of ~ 125 μm. In contrast, surprisingly,

EPSPs detected as ΔF/F$_{eEPSP}$ rather amplified (in some cases to almost double, although variable), when

propagating toward the distal end of a dendritic branch (Fig. 1, E to H and fig. S1D). The $\Delta F/F_{eEPSP}$

increased by 21 ± 4 % in average at 54.7 ± 1.9 μm away from the uncaged site and by 23 ± 3 % at 110.6

± 4.5 μm (Fig. 1H, red). This is quite contrasting to the classical view of passive traveling of subthreshold

membrane potential changes in a cable, in which membrane potential change attenuates during

propagation in a dendritic tree regardless of direction. The $\Delta F/F_{eEPSP}$ amplification detected by ASAP

imaging during propagation toward the distal end was not dependent on the absolute size of $\Delta F/F_{eEPSP}$ at

the uncaged site (ranging from 1.1 % to 10.9 %) that reflects the magnitude of EPSPs (Fig. 1I and fig.

S1D). Thus, our data indicated that excitatory synaptic inputs are oppositely modulated in a dendrite

depending on the direction of propagation.

GABAergic PSPs attenuate as spread in dendrites irrespective of direction

To further study the spatial pattern of spreading of membrane potential changes in a dendrite, we examined

how GABAergic postsynaptic potentials (PSP_{GABAS}) spread in dendrites by the ASAP3β imaging coupled

with local activation of GABA_A receptors (GABA_ARs). PSP_{GABAS} were caused by local uncaging of

DPNI-caged-GABA (200-500 μM) with a spot 405 nm laser illumination (Fig. 2 and fig. S3), in a similar

manner to the local glutamate stimulation. In many cases, local GABA stimulation on a dendrite of

hippocampal neuron exhibited little fluorescence change of ASAP3β, reflecting the resting membrane

potential (Vm_{rest}) substantially close to the equilibrium potential for Cl^- (E_{Cl}). Still, some cells exhibited

clear increase of ASAP3β fluorescence upon GABA uncaging, corresponding to hyperpolarization (Fig.

2A and fig. S3A). In contrast, only very limited cases showed depolarization as observed with fluorescence

decrease upon the local GABA, which is probably caused by somehow high internal Cl^- ($[Cl^-]_{in}$) (Fig. 2B

and fig. S3B). Importantly, in both cases, PSP_{GABAS} were not amplified during the propagation toward

both directions in dendrites (Fig. 2 and fig. S3), which was a sharp contrast to the asymmetric modulation

of EPSP$_{glu}$ (see Fig. 1). This discrepancy in the modulation of membrane depolarization spreading in a

dendritic branch between glutamate- and GABA-caused ones made us to assume that the Cl$^-$ conductance

and/or E$_{Cl}$ might play a critical role in processing of local depolarization in a distal dendritic branch.

Low [Cl$^-$]$_{in}$-dependent deeper Vm$_{rest}$ in distal dendrites

We searched for a clue to test the idea that Cl$^-$ and/or E$_{Cl}$ affect the direction of modulation potential

changes in a dendrite, first by measuring Vm$_{rest}$ in dendrites of hippocampal neurons. For that challenging

task, a direct patch-clamp recording from a very thin distal dendrite (< 1 μm in a diameter) was performed

using the EGFP-fluorescence as a guide to precisely target the pipette (Fig. 3, A and B). Whole-cell

current-clamp recording showed that, surprisingly, the Vm$_{rest}$ was more negative in a dendrite (-75.5 ± 1.4

mV) than in the soma (-68.3 ± 1.4 mV, P = 0.011, Tukey test, Fig. 3C). When the dendritic Vm$_{rest}$ was

plotted against the distance from the soma, there was a tendency for Vm$_{rest}$ to become more negative in a

distal dendrite (-3.2 mV / 100 μm, Fig. 3D). In addition to the deeper Vm$_{rest}$, leak conductance per

membrane area size (measured at -70 mV under the voltage-clamp configuration) became somehow higher

in a dendrite farther away from the soma (Fig. 3E), suggesting that more leak channels are basally open

at the distal dendrite compared to the soma.

Taken the above results together, we hypothesized that the deeper Vm$_{rest}$ in a distal branch might

be related to the Cl$^-$ conductance and/or homeostasis depending on the location of dendrite. This idea was

tested using VU0463271 (10 μM), an inhibitor of KCC2, a type of Cl$^-$ transporter which predominantly

contributes to the low [Cl$^-$]$_{in}$ in mature neurons (19). Immunocytochemistry exhibited more abundant

expression of KCC2 in distal dendrites of hippocampal pyramidal neurons than in the soma (1.5-fold, P =

0.019, Tukey test, Fig. 3, F and G). KCC2 basically establishes low [Cl$^-$]$_{in}$ by excluding Cl$^-$ out of the

cytoplasm, and VU0463271 is expected to bring about higher [Cl$^-$]$_{in}$. When KCC2 was inhibited, the Vm$_{rest}$

in a dendrite became higher by ~ 6 mV, while the somatic V_{mrest} was not affected, resulting in almost

identical potentials at dendrites and soma (dendrite: -66.5 ± 2.0 mV, soma: -67.9 ± 1.9 mV, Fig. 3, C and

D). On the other hand, the tendency of higher basal leak conductance at the distal dendrite was not affected

by the disruption of low $[Cl^-]_{in}$ (Fig. 3E). These results altogether suggested that the lower $[Cl^-]_{in}$ and/or

higher Cl^- conductance in a dendrite than in the soma could be a potential cause for the deeper V_{mrest} in

dendrites.

Deep V_{mrest} via low $[Cl^-]_{in}$ and TTX-resistant Na_V in dendrites are critical for $EPSP_{glu}$ amplification

Next we attempted to examine whether the deep V_{mrest} dependent on the local Cl^- extrusion system in a

distal dendrite is involved in the augmentation of local glutamatergic depolarizing signals. Disruption of

the dendritic deeper V_{mrest} by KCC2 inhibition using VU0463271 (10 μM) clearly, but not completely,

weakened the amplification of a glutamatergic eEPSP in distal dendrites monitored by the ASAP3β

imaging ($\Delta F/F_{eEPSP}$ at an uncaged site and distal site: 3.99 ± 0.39 % and 4.20 ± 0.37 %, P = 0.061, paired

t-test; Fig. 4, A and D). Thus, we considered that the specific $[Cl^-]_{in}$ regulation and/or the resultant deeper

V_{mrest} in distal dendrites compared with the soma might play a role in the local amplification of $EPSP_{glu}$.

We next attempted to identify the remaining factor other than Cl^--related one, mediating the

glutamate-specific dendritic augmentation of depolarization. For that aim, several ion channels were

blocked by pharmacological agents. Voltage-gated Na^+ channels (Na_V) are known to be present in

dendrites of pyramidal neurons (20), contributing to backpropagation of spikes (21). First, we applied a

typical Na_V blocker, tetrodotoxin (TTX, 1 μM) to the external bath, and examined $\Delta F/F_{eEPSP}$ propagation

by ASAP3β imaging. However, the EPSP amplification in distal dendrite was not affected by TTX

($\Delta F/F_{eEPSP}$ amplification, 1.32 ± 0.05, Fig. 4D and S4A). While TTX is a typical Na_V blocker used for

inhibition of Na^+ spikes, TTX-resistant Na_V, such as Na_V1.5, 1.8 and 1.9, are also known. Among these,

at least $Na_V1.5$ and $Na_V1.9$ are reported to be expressed in hippocampal pyramidal neurons (22-24). In the

presence of lidocaine (5 mM), which broadly inhibits Na_V, the $\Delta F/F_{eEPSP}$ amplification in distal dendrites

was not significant (P = 0.25, paired t-test, Fig. 4B), although slight augmentation was still remaining

($\Delta F/F_{eEPSP}$ amplification, 1.12 ± 0.05, Fig. 4D). Thus, TTX-resistant Na_V might also partially underlie the

EPSP amplification in distal dendrites. Simultaneous inhibition of Na_V and KCC2 by lidocaine and

VU0463271 almost abolished the $\Delta F/F_{eEPSP}$ amplification ($\Delta F/F_{eEPSP}$ amplification, 1.04 ± 0.03; P = 0.016,

Dunnett's test, Fig. 4C and D).

We further examined the possible involvement of other cation channels and glutamate receptors

in the EPSP amplification in distal dendrites, although the EPSP size studied here was out of range for

these channels. Voltage-gated Ca^{2+} channels and NMDA receptors are reported to produce dendritic Ca^{2+}

and NMDA spikes upon relatively strong glutamatergic synaptic inputs, contributing to subcellular local

computation by supralinear synaptic integration (6, 25). As expected, neither external administration of

Cd^{2+} (Ca^{2+} channel blocker, 100 μM, Fig. 4D and S4B) nor D-AP5 (NMDA receptor antagonist, 100 μM,

Fig. 4D and S4C) affected the $\Delta F/F_{eEPSP}$ amplification during propagation toward distal dendrites,

suggesting that Ca^{2+} channels and NMDA receptors are not involved in the amplification of small size of

EPSP here we are focusing on. In addition, inhibition of hyperpolarization-activated cation channels (HCN

channels) by application of a blocker, ZD7288, failed to affect the $\Delta F/F_{eEPSP}$ amplification (Fig. 4D, and

fig. S4D). Taken all these results together, it is suggested that TTX-resistant Na_V are critical for the EPSP

amplification in distal dendrites, in cooperation with the low Cl^--dependent deeper V_{mrest}.

Synergistic actions of dendritic local Cl^- conductances and deeper V_{mrest} on local $EPSP_{glu}$ amplification

Finally, we attempted to systematically understand how Cl^--dependent deeper V_{mrest} and TTX-resistant

Na_V together contribute to the local augmentation of EPSP spreading in the distal dendrite. Taking the

following 2 facts into account: 1), the distal dendritic branch exhibits deeper V_{mrest} and higher basal leak

conductance (see Fig. 3); 2), the local augmentation of PSP size is specific to the depolarization mediated

by Na^+ (glutamatergic) but not by Cl^- (GABAergic), (see Fig. 1 and 2), we speculated that the Cl^-

conductance at the distal dendrite might be responsible for the higher leak conductance which could bring

the V_{mrest} closer to E_{Cl}. Among Cl^- channels, hippocampal pyramidal neurons express ClC-2 (Fig. 5A)

(26), which is hyperpolarization-activated and known to regulate Cl^- homeostasis and membrane

excitability (27, 28). Thus, as depicted in Fig. 5B, it would be possible to hypothesize that more likely

activation of ClC-2 by the deeper V_{mrest} in a distal dendrite would increase the basal Cl^- leak conductance,

which would drive the V_{mrest} closer to the E_{Cl} that is kept deep by the powerful local action of KCC2. To

test this hypothesis mechanistically, we constructed a biophysical simple model assuming

hyperpolarization-activated Cl^- conductance like ClC-2 and depolarization-activated Na^+ conductance in

dendritic branches (Fig. 5C; see Methods for detail of model). When an excitatory input (brief increase of

conductance with a reversal potential at 0 mV) was given, an EPSP-like depolarization was caused, and

importantly, the depolarization became larger during the traveling toward a region with enriched

hyperpolarization-activated Cl^- conductances (Fig. 5, C to E). This local augmentation of depolarization

in the model was suppressed either by the elimination of sensitivity of Cl^- conductances to

hyperpolarization or by the reduction of a depolarization-caused Na_V activation (Fig. 5, D and E). Even

when the synaptic input site is altered, the local EPSP amplification at distal branches was evident

although the extent of augmentation was variable (Fig. 5E), in line with the variable size of amplification

in actual data (see Fig. 1). Thus, our simple model supported the idea that the two voltage-sensitive

conductances synergistically contribute to the local EPSP augmentation at distal branches.

The above model simulation theoretically provided a prediction that the voltage-dependent fine

tuning of Cl^- conductance in dendrites is critical for the deeper V_{mrest} and local augmentation of $EPSP_{glu}$.

To evaluate the validity of this model prediction, we first inhibited Cl⁻ conductances by NPPB (50 μM),

a blocker of relatively broad types of Cl⁻ channels including ClC-2, in addition to the Na$_V$ inhibitor

lidocaine, and examined by the ASAP3β imaging how the local augmentation of eEPSP in a distal dendrite

is affected (Fig. 6A). Simultaneous inhibition of Na$_V$ and Cl⁻ channels strongly weakened the EPSP

amplification in line with the simulation results ($\Delta F/F_{eEPSP}$ amplification, 1.04 ± 0.03; $P = 0.009$, Dunnett's

test; Fig. 6C). Furthermore, conversely, aberrant increase of Cl⁻ conductance by applying GABA (50 μM),

which would disturb the voltage-sensitive fine tuning of Cl⁻ conductance and also the Na$_V$ activation

through enhanced shunting effect, completely abolished the direction-dependent augmentation of EPSP$_{glu}$

($\Delta F/F_{eEPSP}$ amplification, 0.91 ± 0.05; $P = 0.0002$, Dunnett's test; Fig. 6, B and C). Taken all these results

together, our simulation of EPSP traveling in a cable equipped with 2 positive-feedback mechanisms,

depolarization-activated Na⁺ conductance and hyperpolarization-activated Cl⁻ conductance, nicely

predicted the cooperative action of those mechanisms to locally augment the excitatory synaptic signals,

which was confirmed by the experiment.

Direction-dependent spatio-temporal integration of synaptic inputs array

Neuronal dendrites are device for spatio-temporal integration of inputs at numerous synapses. Considering

our data that glutamatergic synaptic inputs asymmetrically spread in a dendritic tree depending on the

direction of propagation, we speculated that the synaptic integration might be affected by the asymmetric

modulation of synaptic inputs. For example, apical dendrites of a hippocampal pyramidal neuron receive

inputs from entorhinal cortex (EC) directly via perforant path (or temporo-ammonic path) at distal sites,

and indirectly via the tri-synaptic circuit (mossy fibers or Schaffer collaterals) at a proximal region (29).

Such a circuit organization makes a situation in which excitatory signals arrive with a spatio-temporal

context: from distal to proximal. To test whether synaptic summation is affected by such a spatio-temporal

context of excitatory inputs, distinct temporal orders of glutamate photolysis were performed at multiple locations along a branch (10 sites, 35 ms intervals) in the presence of TTX (1 μM). We compared the effects of three different patterns of eEPSP trains: laser spots were illuminated along a branch, 1) in a sequence from the soma toward the end ("to distal", a to j, Fig. 7A), 2) in a sequence from the end of a branch toward the soma ("to soma", j to a, Fig. 7A), and 3) at "random". Fluorescent imaging with ASAP3β showed that the $\Delta F/F_{eEPSP}$ at distal region (ROI_{distal}, Fig. 7A) was larger than that either at the uncaged branch or at proximal region ($ROI_{proximal}$) in response to all three spatio-temporal patterns of stimuli (Fig. 7B). Thus, $\Delta F/F_{eEPSP}$ upon multiple stimuli on a branch is augmented during the propagation toward the distal end of dendrite, in consistent with the data for single eEPSPs. Notably, peak of $\Delta F/F_{eEPSP}$ by 10 uncagings became smaller either by lidocaine (5 mM) & VU0463271 (10 μM) or Cd^{2+} (100 μM) & D-AP5 (100 μM), compared with the control (Fig. 7C and fig. S5C), indicating several positive feedback mechanisms synergistically operate to enhance the excitatory input train during spatio-temporal summation. In addition, the location-dependent $\Delta F/F_{eEPSP}$ amplification was weakened by application of lidocaine & VU0463271 (fig. S5A), but not by Cd^{2+} & D-AP5 (fig. S5B).

Furthermore, pyramidal neuronal dendrites exhibited distinct augmentation of EPSP summation depending on the spatio-temporal order of synaptic activation. Although the $\Delta F/F_{eEPSP}$ at ROI_{distal} reached similar peaks in two conditions, "to soma" and "to distal" (Fig. 7D and fig. S5D, $P > 0.05$, paired t-test), the signal at a distal branch (ROI_{distal}) steeply summated during glutamate uncagings in a sequence of "to soma" compared to "to distal" (Fig. 7D, $P < 0.01$, two-way ANOVA), but not at a proximal branch ($ROI_{proximal}$, fig. S5E). The ratio of the eEPSP train trace (at ROI_{distal}) for "to soma" divided by that for "to distal", reflecting the preference of "to soma" direction for effective summation of excitatory inputs, was kept higher than 1 during the EPSP array (Fig. 7E). Importantly, such a context-dependent augmented summation was clearly weakened by lidocaine & VU0463271 (Fig. 7D, $P > 0.05$, two-way ANOVA; and

Fig. 7E, P < 0.01, two-way ANOVA). Thus, the Na$_V$- and low [Cl$^-$]$_{in}$-dependent ΔF/F$_{eEPSP}$ amplification

seems to make it possible for a distal branch to decode the spatio-temporal context of synaptic inputs,

resulting in local computation in a distinct manner.

Discussion

In this study, using voltage imaging with a genetically-encoded probe ASAP3β combined with local

glutamate or GABA uncaging, subcellular patch-clamp recordings, and biophysical model simulation, we

explored how subthreshold size of glutamatergic and GABAergic synaptic inputs propagated in

hippocampal neuronal dendrites. Imaging analysis of membrane potential changes at multiple dendritic

branches demonstrated direction-dependent asymmetric propagation of excitatory synaptic inputs in a

dendrite, that is, EPSPs attenuated during propagation toward the soma, while amplified toward the distal

end. Moreover, the EPSP amplification depended on synergistic actions of TTX-resistant Na$_V$ and low

[Cl$^-$]$_{in}$-related Cl$^-$ conductance. Thus, the data presented in this study showed asymmetric modulation of

glutamatergic synaptic inputs in contrast to the conventional view established from the classical cable

theory: subthreshold membrane potential changes gradually attenuate during spreading in dendrites.

Together, our findings provide a novel concept of local modulation of synaptic integration relying on a

unique Cl$^-$-dependent mechanism in a single neuron.

To examine the spatio-temporal propagation pattern of synaptic inputs, we used fluorescent

voltage imaging with a GEVI. Patch-clamp electrophysiology has been a powerful technique to precisely

record electrical signals at a given position in a neuron, although it requires a sophisticated technique. On

the other hand, GEVIs enable us to measure membrane potential changes much more easily even from

multiple subcellular regions at the same time, that is very difficult to perform by conventional patch-clamp

recordings. In this decade, several types of GEVI, including ASAP (13), have been developed, such as

ArcLight (30), Ace2N-mNeon (14), Voltron (31), and so on (15, 32). Taking advantage of this merit,

recent studies applied GEVIs to various preparations *in vivo* and *in vitro,* demonstrating neuronal

excitability changes dependent on behavioral states in animals (32). Moreover, GEVIs could be non-

invasive compared with whole-cell patch-clamp recordings which is typically accompanied with washout

of cytoplasmic components. As demonstrated in this study, $[Cl^-]_{in}$ is critical for the local EPSP

amplification (see Fig. 3-7), and we speculate that monitoring of membrane potential without intracellular

dialysis must have been essential for the detection of such cellular functions. In addition, the EPSP

amplification demonstrated here operates even for small EPSPs (locally 2 ~ 5 mV and hence ~ 1 mV or

smaller at the soma) in a very local branch of distal dendrites specifically, which cannot be recognized by

the somatic recordings. Therefore, the lack of useful technique like voltage imaging based on GEVIs must

have veiled the Cl^--dependent asymmetric EPSP amplification at a distal dendrite.

Previous studies have suggested local augmentation of synaptic inputs by active conductances via

voltage-gated cation channels such as Na^+ and/or Ca^{2+} channels and NMDA receptors (4, 6, 25, 33).

Notably, such augmentation of local excitatory inputs emerges when the input size is relatively large (~ 5

mV or larger when recorded at the soma and hence ~ 20 mV at the stimulated site). On the other hand, as

shown in this study, eEPSPs caused here by spot laser illumination on a spine, was not so large in

amplitude: ~ 3 % $\Delta F/F$ in average, corresponding to ~ 5 mV depolarization at the stimulated site, and the

local EPSP amplification relied on the TTX-resistant Na^+ channels. Interestingly, among three subtypes

of such Na^+ channels ($Na_V1.5$, 1.8 and 1.9), $Na_V1.9$ is especially activated at more negative membrane

potential (-70 ~ -60 mV) (34, 35). Taking it into consideration that activations of typical Na_V and Ca^{2+}

channels are triggered at -40 ~ -50 mV, these channels are unlikely to be involved in the amplification of

small size of EPSPs. Thus, distinct subtypes of Nav with different activation profile might provide neurons

wide range of locally computation.

In contrast to the conventional view of dendritic computation focusing on cation channels, this

study identified a unique mechanism of local augmentation of excitatory inputs in a distal dendrite: low

$[Cl^-]_{in}$- and Cl^- conductance-dependent enhancement of depolarization in distal dendrites (see Fig. 4-6).

Our biophysical model simulation indicated that a type of hyperpolarization-activated Cl^- channel, ClC-2,

could constitute a kind of positive-feedback loop to stably keep the membrane potential negative as close

to the E_{Cl}. Indeed, direct subcellular patch-clamp recordings from dendritic branches demonstrated the

deeper Vm_{rest} dependent on low $[Cl^-]_{in}$ and higher leak conductances at distal branches in a distance-

dependent manner (see Fig. 3). Such a positive-feedback hyperpolarizing system is switched-off upon a

depolarization, resulting in indirect enhancement of depolarizing responses (see Fig. 5). The specific

subtype of Cl^- channels in the asymmetric EPSP modulation, here we could not determine because of the

lack of specific blockers, is an important issue to be addressed in a future study.

In addition to the involvement of Cl^- conductance, E_{Cl} may also be altered at distal dendrites

contributing to deeper Vm_{rest} there because of the lower $[Cl^-]_{in}$ which is established by KCC2 activity, as

reported previously (36-39). More abundant expression of KCC2 in spines and vicinity of spines (37, 40)

and/or more efficient Cl^- extrusion by KCC2 might operate at distal dendrites based on the distinct

surface/volume ratio, as shown by Báldi et al. (41). Furthermore, regulation of KCC2 by phosphorylation

and dephosphorylation in distal dendrites (36, 42, 43) may also give rise to diverse modes of Cl^-

homeostasis, potentially impacting the local processing of synaptic potentials in a branch. As shown in

Fig. 1 and 2, in contrast to the asymmetric modulation of $EPSP_{glu}$s, depolarizing PSP_{GABA}s symmetrically

attenuated during propagation in dendrites. Depolarizing PSP_{GABA}s would have been caused by unusually

high $[Cl^-]_{in}$, in a manner similar to the situation with VU0463271 (see Fig. 3), so that the E_{Cl} should be

higher and then the deep V_{mrest} at a distal dendrite would also be disturbed, precluding one of the two

critical positive-feedback mechanisms for the local amplification of depolarization.

As demonstrated in this study (see Fig. 7), hippocampal and cortical pyramidal neurons exhibit

supralinear synaptic integration upon multiple inputs in local dendrites dependent on active conductances

(5, 6), whereas other types such as dentate gyrus granule cells and cerebellar interneurons show linear or

sublinear summation (44-46). Accordingly, among hippocampal culture preparation consisting of several

types of neurons, pyramidal cells, but not apparent granule cells, selectively exhibited the local

asymmetric EPSP modulation (see Fig. 1). Thus, distinct neuronal types seem to be equipped with specific

machinery for local dendritic computation. Asymmetric propagation of EPSPs may play a role in

associative induction of synaptic plasticity such as long-term potentiation or depression. EPSPs caused at

a dendrite sequentially in the direction toward the soma, likely cause an AP firing in cortical pyramidal

neurons (47). Furthermore, the long-term plasticity at excitatory synapses tends to be induced by dendritic

spike firing in a distal thin branch caused by clustered inputs, rather than distributed ones (48, 49). Thus,

asymmetric modulation of EPSPs revealed here might associate a given pattern of EPSPs arriving at a

distal dendrite, contributing to the local dendritic spikes and long-term plasticity.

Methods

Plasmid construction

Plasmids were constructed by standard molecular biology techniques. The coding DNA for ASAP1 was

obtained from Addgene (#52519), and some modifications for higher voltage sensitivity (D145Δ, L146G,

S147T, N149R, S150G, and H151D) in an improved version ASAP3 (18) were introduced by PCR-

mediated mutations. Modified-ASAP (ASAP3β) or original ASAP1 cDNAs was inserted into the vector

made from pAAV-MSC by replacement of CMV promotor and beta-globin intron region by CA promotor

and intron region from pCAGGS, followed by insertion of WPRE sequence before hGH pA region.

HEK293T cells

HEK293T cells were cultured at 37 °C in humidified air containing 5 % CO_2 in DMEM (Nakalai Tesque,

Kyoto, Japan) supplemented with 10 % (v/v) FBS (NICHIREI BIOSCIENCES, Tokyo, Japan) and

penicillin (100 U/mL)-streptomycin (0.1 mg/mL, Nakalai Tesque). For testing voltage-dependent

fluorescence changes, HEK293T cells were transfected with original ASAP1 or modified-ASAP

(ASAP3β) using Lipofectamine 3000 (1 μg DNA, 2 μL P3000 reagent, 1.5 μL Lipofectamine for 35 mm

dish; Thermo Fisher Scientific, Waltham, MA, USA).

Hippocampal neuron culture and transfection

The animals were maintained and treated according to the NIH guide for the care and use of laboratory

animals, and to the ethical guidelines on animal experimentation of Kyoto University. Hippocampi were

dissected out from male and female newborn Wistar rats and incubated in Ca^{2+} and Mg^{2+}-free Hank's

balanced salt solution (CMF-HBSS) containing 0.1 % trypsin for 5 min at 37 °C. Neurons were dissociated

by trituration with a fire-polished Pasteur pipette and seeded on poly-D-lysine (Sigma-Aldrich, St. Louis,

MO, USA)-coated glasses in Neurobasal plus medium containing 2 % B27 plus supplement (Thermo

Fisher Scientific) and penicillin (100 U/mL)-streptomycin (0.1 mg/mL, Nakalai Tesque). Some cells were

transfected with plasmids by means of electroporation with NEPA21 (Nepa Gene, Chiba, Japan). 1×10^6

cells were mixed with plasmids (6 μg ASAP3β plasmid or 3 μg pCAsalEGFP (50)) in 100 μL CMF-

HBSS. Square electric pulses were applied at 300 V (pulse length, 0.5 ms; two pulses; interval, 50 ms;

voltage decay rate, 10 %), followed by additional pulses at 20 V (pulse length, 50 ms; five pulses; interval,

50 ms; voltage decay rate, 40 %; polarity exchanged pulse). Transfected cells were then cultured on the

glasses together with the non-treated ones. The cells were cultured at 37 °C in humidified air containing

5 % CO_2. One day after seeding, 75 % of the medium was replaced with fresh one. Since then, half of the

medium was replaced every 4 days.

Electrophysiology

Whole-cell voltage-clamp recording from an ASAP-expressing HEK293T cell was performed with an

amplifier (EPC10; HEKA Elektronik GmbH, Reutlingen, Germany) in an extracellular solution containing

(in mM) 145 NaCl, 5 KOH, 2 $CaCl_2$, 1 $MgCl_2$, 10 Hepes and 10 glucose (pH 7.3) at room temperature

(20-24 °C). Patch pipettes (3-5 MΩ) were filled with an internal solution containing (in mM) 147 CsCl, 5

EGTA, 10 Hepes, 15 CsOH, 2 ATP, 0.2 GTP (pH 7.3). The membrane potential (a holding potential of -

70 mV) was changed by step voltage depolarizations and hyperpolarizations to a voltage (-100 mV to 50

mV) for 30 ms.

In recordings from hippocampal neurons in culture for 2-3 weeks, we used an extracellular solution

containing (in mM) 120 NaCl, 5 KOH, 2 $CaCl_2$, 1 $MgCl_2$, 10 Hepes and 10 glucose (pH 7.3). Patch pipettes

(3-5 MΩ, somatic recordings; 20-25 MΩ, dendritic recordings) were filled with an internal solution

containing (in mM) 120 K-gluconate, 7 KCl, 5 EGTA, 10 Hepes, 2 ATP, 0.2 GTP (pH 7.3 adjusted with

KOH). In some experiments, TTX (1 μM, FUJIFILM Wako Pure Chemical Corporation, Tokyo, Japan),

lidocaine (5 mM, Nakalai Tesque), D-AP5 (100 μM, Tocris Bioscience, Bio-Techne SRL, Minneapolis,

MN, USA), Cd^{2+} (100 μM, FUJIFILM Wako Pure Chemical Corporation), ZD7288 (100 μM, Tocris

bioscience), VU0463271 (10 μM, Tocris Bioscience) and NPPB (50 μM, Tocris Bioscience) were used

to inhibit Na^+ channels, NMDA receptors, Ca^{2+} channels, HCN channels, IRK channels, KCC2 and Cl^-

channels, respectively. GABA (50 μM, Nakalai Tesque) was used to activate $GABA_A$ receptors.

Fluorescent imaging

Fluorescence imaging was performed using an inverted microscope IX71 (Olympus, Tokyo, Japan) through a 40× 0.95-NA objective (Olympus) for HEK293T cells, or a 60× 1.42-NA oil-immersion objective (Olympus) for hippocampal neurons. Fluorescent excitation was delivered using a white LED (SOLA; Lumencor, OR, USA) with 15 mW/mm^2 through a 450-480 nm filter. Fluorescent images were obtained with a Zyla4.2 sCMOS camera (Andor Technology Ltd, Belfast, UK) at 1500 Hz (128 × 128 pixels, binning 2 × 2) for HEK293T cells, at 400 Hz (512 × 512 pixels, binning 2 × 2) or at 200 Hz (1392 × 1040 pixels, binning 2 × 2) for hippocampal neurons and analyzed with SOLIS (Andor) or Image J software (NIH, Bethesda, MD, USA).

The LED illumination in our self-customized fluorescent imaging system is designed to be assembled around the center of view field so that high fluorescent signal is attained as possible even for a very fast image acquisition. As a result, the absolute fluorescent signals around the edge of view field tended to be smaller, potentially giving rise to a tendency for the signal to noise ratio to become lower at those dimmer regions. However, we confirmed that, as shown in fig. S2, spatial pattern of $\Delta F/F_{eEPSP}$ signals in dendrites was little affected by the relative brightness of fluorescence altered by moving the microscope.

Glutamate or GABA uncaging

For glutamate or GABA uncaging, MNI-caged-L-glutamate (300 μM, Tocris Bioscience) or DPNI-caged-GABA (200-500 μM, Tocris Bioscience) was applied to the external solution. Uncaging was caused by 405 nm laser illumination (Single line laser; Rapp OptoElectronic GmbH, Wedel, Germany) with 4-6 mW laser power (51, 52). Glutamate uncaging was performed by laser illumination at a 2 μm diameter spot

(0.2 ms duration; or 1 ms in the presence of external GABA) around 200 μm from the soma, while GABA

was uncaged by a spot or a line scan (0.5-5 ms duration). The location of laser illumination was controlled

by a localized photomanipulation system based on Galvano mirrors (UGA-42 Firefly; Rapp

OptoElectronic) implemented in SysCon software (Rapp OptoElectronic).

Immunocytochemistry

Cultured neurons were fixed with 4 % paraformaldehyde in phosphate buffered saline, permeabilized with

0.5 % Tween 20, then blocked with 1 % goat serum, and finally labeled with primary and secondary

antibodies. The following antibodies were used: rabbit polyclonal antibodies (pAb) against KCC2 (1:1000,

Sigma-Aldrich), ClC-2 (1:5000, Alomone Labs, Jerusalem, Israel), chicken pAb against GFP (1:2000,

Sigma-Aldrich), Alexa 568-conjugated pAb against rabbit IgG (1:400, Thermo Fisher Scientific) and

Alexa 488-conjugated pAb against chicken IgG (1:400, Thermo Fisher Scientific). Fluorescent images

were recorded with a confocal laser microscope (FV1000 imaging system, Olympus, Japan), and analyzed

using ImageJ software (NIH). The neuronal dendrites were defined as the area positive for GFP.

Simulation of simple dendrite model with voltage-dependent Na$^+$ and Cl$^-$ channels

Simple model of a dendrite equipped with the depolarization-activated Na$_V$ and the hyperpolarization-

activated Cl$^-$ channels was constructed and simulated using the *NEURON* simulator (53) version 8.2

operating under the Python programming language. Here we assumed a multi-compartment model

consisting of a soma and 18 dendritic sections with single branch point where 3 daughter branches

stemmed out (see Fig. 5C), one of which had 120 μm and the other two 90 μm in length, yielding total

arborization length of 500 or 470 μm long. The diameter of a dendritic branch was assumed to be 2.5 μm

at the most proximal one, narrowing down towards the distal end until it reaches 0.5 μm at 200 μm away

from the soma. Cytosolic resistivity was here set at 250 Ω·cm, and membrane capacitance 0.8 mF/cm^2.

Membrane of individual compartments were equipped with TTX-resistant voltage-gated Na$^+$ channel

Na$_V$1.9, hyperpolarization-activated Cl$^-$ channel like ClC-2, Na$^+$/K$^+$ pump, KCC2, and linear leak

conductances for Na$^+$, K$^+$, and Cl$^-$ (for detailed parameters of each components, see table S1).

Individual components in each compartment at the membrane potential (V_m) were expressed in

the model based on the equilibrium potentials for Na$^+$ (E_{Na}), K$^+$ (E_K), and Cl$^-$ (E_{Cl}) by implementing the

equations which were already published in previous studies, as described below.

For TTX-resistant persistent Na channels, Na$_V$1.9 model (54) was here used:

450
$$I_{Nav} = \overline{g_{Nav}} \times m_{Nav} \times h_{Nav} \times (V_m - E_{Na})$$

451 , where $\overline{g_{Nav}}$, m_{Nav}, h_{Nav} are the maximal conductance, voltage-dependent activation and inactivation

452 probability factors, respectively. Voltage-dependent changes of m_{Nav} and h_{Nav} were defined by the rate

453 constants (α and β) as described by the following equations:

454
$$\frac{dm_{Nav}}{dt} = \alpha_{m_{Nav}} \times (1 - m_{Nav}) - \beta_{m_{Nav}} \times m_{Nav}$$

455
$$\frac{dh_{Nav}}{dt} = \alpha_{h_{Nav}} \times (1 - h_{Nav}) - \beta_{h_{Nav}} \times h_{Nav}$$

456 , where forward (α) and reverse (β) rate constants for m_{Nav} and h_{Nav} were assumed as below:

457
$$\alpha_{m_{Nav}} = \frac{1.032}{1 + \exp\left(\frac{V_m + 6.99}{-14.87115}\right)}$$

458
$$\beta_{m_{Nav}} = \frac{5.79}{1 + \exp\left(\frac{V_m + 130.4}{22.9}\right)}$$

459
$$\alpha_{h_{Nav}} = \frac{0.06435}{1 + \exp\left(\frac{V_m + 73.26415}{3.71928}\right)}$$

460
$$\beta_{h_{Nav}} = \frac{0.13496}{1 + \exp\left(\frac{V_m + 10.27853}{-9.09334}\right)}$$

For hyperpolarization-activated Cl⁻ component, a modified biophysical model for ClC-2 (55) was

incorporated:

$$I_{Cl} = \bar{g}_{Cl} \times m_{Cl} \times (V_m - E_{Cl})$$

$$\frac{dm_{Cl}}{dt} = 0.1 \times (m_{Cl,ss} - m_{Cl})$$

, where $\bar{g}_{Cl}$ is the maximal conductance, m_{Cl} is the relative open probability factor of the channel, and its

steady state at V_m, $m_{Cl,ss}$ is defined as:

$$m_{Cl,ss} = \frac{1}{1 + \exp\left(\frac{V_m - E_{Cl} + V_h}{V_s}\right)}$$

, where V_h and V_s are parameters to define the voltage-sensitivity of the channel activation.

For KCC2, K⁺ and Cl⁻ fluxes were calculated as a product of the parameter for its anti-porting efficiency

(U_{KCC2}) and the difference between their equilibrium potentials as demonstrated in Gentiletti et al. (56):

$$I_{K(KCC2)} = U_{KCC2} \ln\left(\frac{[K^+]_{in}[Cl^-]_{in}}{[K^+]_{out}[Cl^-]_{out}}\right)$$

$$I_{Cl(KCC2)} = -I_{K(KCC2)}$$

Na⁺/K⁺ pump activity was calculated based on the formulations in Takeuchi et al. (57), as listed below,

$$I_{NaK} = I_{NaKmax}(k_1 \times p_{E1Na} \times y - 0.04 \times p_{E2Na} \times (1 - y))$$

$$I_{K(pump)} = -2I_{NaK}$$

$$I_{Na(pump)} = 3I_{NaK}$$

State transitions of the pump were defined by rate constants as below:

$$\frac{dy}{dt} = \alpha_y(1 - y) - \beta_y y$$

$$\alpha_y = 0.04 \times p_{E2Na} + 0.165 \times p_{E2K}$$

$$\beta_y = \frac{0.37}{1 + \frac{0.094}{[ATP]_{in}}} \times p_{E1Na} + 0.01 \times p_{E1K}$$

$$[Na^+]_{eff} = [Na^+]_{out} \exp\left(-\frac{0.82 \times 96.4853 \times V_m}{8.314463 \times T}\right)$$

$$p_{E1Na} = \left(1 + \left(\frac{4.05}{[\text{Na}^+]_{in}}\right)^{1.06} \left(1 + \frac{[K^+]_{in}}{32.88}\right)^{1.12}\right)^{-1}$$

$$p_{E1K} = \left(1 + \left(\frac{32.88}{[\text{K}^+]_{in}}\right)^{1.12} \left(1 + \frac{[\text{Na}^+]_{in}}{4.05}\right)^{1.06}\right)^{-1}$$

$$p_{E2Na} = \left(1 + \left(\frac{69.8}{[\text{Na}^+]_{eff}}\right)^{1.06} \left(1 + \frac{[K^+]_{out}}{0.258}\right)^{1.12}\right)^{-1}$$

$$p_{E2K} = \left(1 + \left(\frac{0.258}{[\text{K}^+]_{out}}\right)^{1.12} \left(1 + \frac{[\text{Na}^+]_{eff}}{69.8}\right)^{1.06}\right)^{-1}$$

In our model, calculation of the Na^+/K^+ pump activity was specifically implemented in Neuron simulator, by modifying the program code presented in Botta et al. (58).

Notably, here we assumed the hyperpolarization-activated Cl^- channels only at distal compartments (see Fig. 5C). An EPSP was represented as a double exponential conductance change (reversal potential at 0 mV) at a single compartment (#6-8 in Fig. 5C), defined by the rise and decay time constants of 10 and 80 ms, respectively, and the peak conductance was optimized to cause similar size of EPSP (~ 12 mV) at the stimulated compartment. The model was simulated in *Neuron* platform with backward Euler method (dt = 0.025 ms). In some simulations to evaluate the role of the voltage-dependent Na^+ and Cl^- conductances in the model, the maximal conductance of Na_V was halved and/or the Cl^- conductances at distal branches were set constant.

Data analysis of ASAP fluorescence change

Dendritic regions were selected for analysis as ROIs in ImageJ, and F_{dend} was calculated as the mean intensity of fluorescence in each ROI which was subtracted by background noise defined as the mean of fluorescence intensity at the area nearby each ROI. During image acquisition of ASAP-expressing cells,

fluorescence gradually decreased due to photobleaching, which was defined as F_{base} obtained by double

exponential curve fitting of the total fluorescence change as follows:

$$F_{base} = A + B \cdot exp \{\alpha \cdot (-t / \tau_1) + (1 - \alpha) \cdot (-t/\tau_2)\}$$

All above parameters (A, B, α, τ_1 and τ_2) were determined by the least square method from the time course

of fluorescence change before the uncaging. The relative change of fluorescence ($\Delta F/F$, %) was

$$\Delta F/F = (F_{dend} / F_{base} - 1) \cdot 100$$

In this study, relative fluorescence change of ASAP is represented as $-\Delta F/F$ because of the negative

relation of fluorescence against membrane potential change. $-\Delta F/F$ traces in individual ROIs were

obtained by averages of 3-7 trials. Adjacent area to the laser spot was excluded from analysis because of

an artifact by laser illumination. Distance between two sites of interest was measured using ImageJ (NIH).

Data fitting for $\Delta F/F$ against the distance from the uncaged site was performed by Igor Pro8 (WaveMetrics,

Lake Oswego, OR, USA) or Microsoft Excel.

Statistics analysis

Statistical significance of differences between groups was tested by two-tailed paired Student's t-test,

Dunnett's test, Tukey test or two-way ANOVA, and $P < 0.05$ was considered as significant. Throughout

the project, data are presented as mean $\pm$ standard error of the mean (SEM) unless otherwise stated. In all

figures, symbols with error bars indicate mean $\pm$ SEM; asterisk, $P < 0.05$; double asterisk, $P < 0.01$; triple

asterisk, $P < 0.001$. These analyses were performed by Microsoft Excel and Igor Pro8 (WaveMetrics).

References

1. G. J. Stuart, N. Spruston, Dendritic integration: 60 years of progress. *Nat Neurosci*. **18**, 1713–1721 (2015).

2. M. J. Higley, B. L. Sabatini, Calcium Signaling in Dendrites and Spines: Practical and Functional Considerations. *Neuron*. **59**, 902–913 (2008).

3. G. Ariav, A. Polsky, J. Schiller, Submillisecond Precision of the Input-Output Transformation Function Mediated by Fast Sodium Dendritic Spikes in Basal Dendrites of CA1 Pyramidal Neurons. *J. Neurosci.* **23**, 7750–7758 (2003).

4. A. Losonczy, J. C. Magee, Integrative Properties of Radial Oblique Dendrites in Hippocampal CA1 Pyramidal Neurons. *Neuron*. **50**, 291–307 (2006).

5. T. Branco, M. Häusser, Synaptic Integration Gradients in Single Cortical Pyramidal Cell Dendrites. *Neuron*. **69**, 885–892 (2011).

6. J. K. Makara, J. C. Magee, Variable Dendritic Integration in Hippocampal CA3 Pyramidal Neurons. *Neuron*. **80**, 1438–1450 (2013).

7. N. L. Golding, H. Jung, T. Mickus, N. Spruston, Dendritic Calcium Spike Initiation and Repolarization Are Controlled by Distinct Potassium Channel Subtypes in CA1 Pyramidal Neurons. *J. Neurosci.* **19**, 8789–8798 (1999).

8. S. R. Williams, G. J. Stuart, Mechanisms and consequences of action potential burst firing in rat neocortical pyramidal neurons. *The Journal of Physiology*. **521**, 467–482 (1999).

9. N. L. Golding, T. J. Mickus, Y. Katz, W. L. Kath, N. Spruston, Factors mediating powerful voltage attenuation along CA1 pyramidal neuron dendrites: Voltage attenuation in CA1 dendrites. *The Journal of Physiology*. **568**, 69–82 (2005).

10. T. Nevian, M. E. Larkum, A. Polsky, J. Schiller, Properties of basal dendrites of layer 5 pyramidal neurons: a direct patch-clamp recording study. *Nat Neurosci*. **10**, 206–214 (2007).

11. W. Rall, Distinguishing theoretical synaptic potentials computed for different soma-dendritic distributions of synaptic input. *Journal of Neurophysiology*. **30**, 1138–1168 (1967).

12. A. Tran-Van-Minh, R. D. Cazé, T. Abrahamsson, L. Cathala, B. S. Gutkin, D. A. DiGregorio, Contribution of sublinear and supralinear dendritic integration to neuronal computations. *Front. Cell. Neurosci.* **9** (2015), doi:10.3389/fncel.2015.00067.

13. F. St-Pierre, J. D. Marshall, Y. Yang, Y. Gong, M. J. Schnitzer, M. Z. Lin, High-fidelity optical reporting of neuronal electrical activity with an ultrafast fluorescent voltage sensor. *Nat Neurosci*. **17**, 884–889 (2014).

14. Y. Gong, C. Huang, J. Z. Li, B. F. Grewe, Y. Zhang, S. Eismann, M. J. Schnitzer, High-speed recording of neural spikes in awake mice and flies with a fluorescent voltage sensor. *Science*. **350**, 1361–1366 (2015).

15. Y. Xu, P. Zou, A. E. Cohen, Voltage imaging with genetically encoded indicators. *Current Opinion in Chemical Biology*. **39**, 1–10 (2017).

16. T. Kwon, M. Sakamoto, D. S. Peterka, R. Yuste, Attenuation of Synaptic Potentials in Dendritic Spines. *Cell Reports*. **20**, 1100–1110 (2017).

17. M. B. Hoppa, G. Gouzer, M. Armbruster, T. A. Ryan, Control and Plasticity of the Presynaptic Action Potential Waveform at Small CNS Nerve Terminals. *Neuron*. **84**, 778–789 (2014).

18. V. Villette, M. Chavarha, I. K. Dimov, J. Bradley, L. Pradhan, B. Mathieu, S. W. Evans, S. Chamberland, D. Shi, R. Yang, B. B. Kim, A. Ayon, A. Jalil, F. St-Pierre, M. J. Schnitzer, G. Bi, K. Toth, J. Ding, S. Dieudonné, M. Z. Lin, Ultrafast Two-Photon Imaging of a High-Gain Voltage Indicator in Awake Behaving Mice. *Cell*. **179**, 1590-1608.e23 (2019).

19. R. Liu, J. Wang, S. Liang, G. Zhang, X. Yang, Role of NKCC1 and KCC2 in Epilepsy: From Expression to Function. *Front. Neurol.* **10**, 1407 (2020).

20. S. Remy, H. Beck, Y. Yaari, Plasticity of voltage-gated ion channels in pyramidal cell dendrites. *Current Opinion in Neurobiology*. **20**, 503–509 (2010).

21. G. J. Stuart, B. Sakmann, Active propagation of somatic action potentials into neocortical pyramidal cell dendrites. *Nature*. **367**, 69–72 (1994).

22. R. Blum, K. W. Kafitz, A. Konnerth, Neurotrophin-evoked depolarization requires the sodium channel NaV1.9. *Nature*. **419**, 687–693 (2002).

23. S.-Y. Jeong, J. Goto, H. Hashida, T. Suzuki, K. Ogata, N. Masuda, M. Hirai, K. Isahara, Y. Uchiyama, I. Kanazawa, Identification of a Novel Human Voltage-Gated Sodium Channel α Subunit Gene, SCN12A. *Biochemical and Biophysical Research Communications*. **267**, 262–270 (2000).

24. Y. Zhou, Z. Luo, F. Bi, J. Li, B. Xiao, Expression of Nav1.5 in the pathogenesis of temporal lobe epilepsy. *Cell Mol Biol (Noisy-le-grand)*. **65**, 58–62 (2019).

25. S. Yang, V. Emiliani, C.-M. Tang, The kinetics of multibranch integration on the dendritic arbor of CA1 pyramidal neurons. *Front. Cell. Neurosci.* **8** (2014), doi:10.3389/fncel.2014.00127.

26. A. Sík, R. L. Smith, T. F. Freund, Distribution of chloride channel-2-immunoreactive neuronal and astrocytic processes in the hippocampus. *Neuroscience*. **101**, 51–65 (2000).

27. K. Staley, R. Smith, J. Schaack, C. Wilcox, T. J. Jentsch, Alteration of GABAA Receptor Function Following Gene Transfer of the CLC-2 Chloride Channel. *Neuron*. **17**, 543–551 (1996).

28. I. Rinke, J. Artmann, V. Stein, ClC-2 Voltage-Gated Channels Constitute Part of the Background Conductance and Assist Chloride Extrusion. *J. Neurosci.* **30**, 4776–4786 (2010).

29. J. Stepan, J. Dine, M. Eder, Functional optical probing of the hippocampal trisynaptic circuit in vitro: network dynamics, filter properties, and polysynaptic induction of CA1 LTP. *Frontiers in Neuroscience*. **9** (2015), doi:10.3389/fnins.2015.00160.

30. L. Jin, Z. Han, J. Platisa, J. R. A. Wooltorton, L. B. Cohen, V. A. Pieribone, Single Action Potentials and Subthreshold Electrical Events Imaged in Neurons with a Fluorescent Protein Voltage Probe. *Neuron*. **75**, 779–785 (2012).

31. A. S. Abdelfattah, T. Kawashima, A. Singh, O. Novak, H. Liu, Y. Shuai, Y.-C. Huang, L. Campagnola, S. C. Seeman, J. Yu, J. Zheng, J. B. Grimm, R. Patel, J. Friedrich, B. D. Mensh, L. Paninski, J. J. Macklin, G. J. Murphy, K. Podgorski, B.-J. Lin, T.-W. Chen, G. C. Turner, Z. Liu, M. Koyama, K. Svoboda, M. B. Ahrens, L. D. Lavis, E. R. Schreiter, Bright and photostable chemigenetic indicators for extended in vivo voltage imaging. *Science*. **365**, 699–704 (2019).

32. Y. Adam, J. J. Kim, S. Lou, Y. Zhao, M. E. Xie, D. Brinks, H. Wu, M. A. Mostajo-Radji, S. Kheifets, V. Parot, S. Chettih, K. J. Williams, B. Gmeiner, S. L. Farhi, L. Madisen, E. K. Buchanan, I. Kinsella, D. Zhou, L. Paninski, C. D. Harvey, H. Zeng, P. Arlotta, R. E. Campbell, A. E. Cohen, Voltage imaging and optogenetics reveal behaviour-dependent changes in hippocampal dynamics. *Nature*. **569**, 413–417 (2019).

33. A. Losonczy, J. K. Makara, J. C. Magee, Compartmentalized dendritic plasticity and input feature storage in neurons. *Nature*. **452**, 436–441 (2008).

34. M. Stevens, S. Peigneur, J. Tytgat, Neurotoxins and Their Binding Areas on Voltage-Gated Sodium Channels. *Front. Pharmacol.* **2** (2011), doi:10.3389/fphar.2011.00071.

35. S. D. Dib-Hajj, J. A. Black, S. G. Waxman, NaV1.9: a sodium channel linked to human pain. *Nat Rev Neurosci*. **16**, 511–519 (2015).

36. I. Chamma, M. Heubl, Q. Chevy, M. Renner, I. Moutkine, E. Eugène, J. C. Poncer, S. Lévi, Activity-Dependent Regulation of the K/Cl Transporter KCC2 Membrane Diffusion, Clustering, and Function in Hippocampal Neurons. *J. Neurosci.* **33**, 15488–15503 (2013).

37. G. Gauvain, I. Chamma, Q. Chevy, C. Cabezas, T. Irinopoulou, N. Bodrug, M. Carnaud, S. Lévi, J. C. Poncer, The neuronal K-Cl cotransporter KCC2 influences postsynaptic AMPA receptor content and lateral diffusion in dendritic spines. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 15474–15479 (2011).

38. S. Khirug, K. Huttu, A. Ludwig, S. Smirnov, J. Voipio, C. Rivera, K. Kaila, L. Khiroug, Distinct properties of functional KCC2 expression in immature mouse hippocampal neurons in culture and in acute slices: Regulation of KCC2 in immature hippocampal neurons. *European Journal of Neuroscience*. **21**, 899–904 (2005).

39. S. Khirug, J. Yamada, R. Afzalov, J. Voipio, L. Khiroug, K. Kaila, GABAergic Depolarization of the
Axon Initial Segment in Cortical Principal Neurons Is Caused by the Na–K–2Cl Cotransporter NKCC1.
J. Neurosci. **28**, 4635–4639 (2008).

40. I. Chamma, Q. Chevy, J. C. Poncer, S. Lévi, Role of the neuronal K-Cl co-transporter KCC2 in
inhibitory and excitatory neurotransmission. *Front. Cell. Neurosci.* **6** (2012),
doi:10.3389/fncel.2012.00005.

41. R. Báldi, C. Varga, G. Tamás, Differential distribution of KCC2 along the axo-somato-dendritic axis
of hippocampal principal cells: KCC2 distribution along hippocampal principal cells. *European Journal
of Neuroscience*. **32**, 1319–1325 (2010).

42. J. Rinehart, Y. D. Maksimova, J. E. Tanis, K. L. Stone, C. A. Hodson, J. Zhang, M. Risinger, W. Pan,
D. Wu, C. M. Colangelo, B. Forbush, C. H. Joiner, E. E. Gulcicek, P. G. Gallagher, R. P. Lifton, Sites of
Regulated Phosphorylation that Control K-Cl Cotransporter Activity. *Cell*. **138**, 525–536 (2009).

43. E. Côme, X. Marques, J. C. Poncer, S. Lévi, KCC2 membrane diffusion tunes neuronal chloride
homeostasis. *Neuropharmacology*. **169**, 107571 (2020).

44. R. Krueppel, S. Remy, H. Beck, Dendritic Integration in Hippocampal Dentate Granule Cells. *Neuron*.
71, 512–528 (2011).

45. T. Abrahamsson, L. Cathala, K. Matsui, R. Shigemoto, D. A. DiGregorio, Thin Dendrites of Cerebellar
Interneurons Confer Sublinear Synaptic Integration and a Gradient of Short-Term Plasticity. *Neuron*. **73**,
1159–1172 (2012).

46. A. Tran-Van-Minh, T. Abrahamsson, L. Cathala, D. A. DiGregorio, Differential Dendritic Integration
of Synaptic Potentials and Calcium in Cerebellar Interneurons. *Neuron*. **91**, 837–850 (2016).

47. T. Branco, B. A. Clark, M. Häusser, Dendritic Discrimination of Temporal Input Sequences in Cortical
Neurons. *Science*. **329**, 1671–1675 (2010).

48. Á. Magó, J. P. Weber, B. B. Ujfalussy, J. K. Makara, Synaptic Plasticity Depends on the Fine-Scale
Input Pattern in Thin Dendrites of CA1 Pyramidal Neurons. *J. Neurosci.* **40**, 2593–2605 (2020).

49. J. P. Weber, B. K. Andrásfalvy, M. Polito, Á. Magó, B. B. Ujfalussy, J. K. Makara, Location-
dependent synaptic plasticity rules by dendritic spine cooperativity. *Nat Commun*. **7**, 11380 (2016).

50. K. Horikawa, M. Takeichi, Requirement of the juxtamembrane domain of the cadherin cytoplasmic
tail for morphogenetic cell rearrangement during myotome development. *Journal of Cell Biology*. **155**,
1297–1306 (2001).

51. L. C. Gomez, S. Kawaguchi, T. Collin, A. Jalil, M. del P. Gomez, E. Nasi, A. Marty, I. Llano, Influence
of spatially segregated IP3-producing pathways on spike generation and transmitter release in Purkinje
cell axons. *Proceedings of the National Academy of Sciences*. **117**, 11097–11108 (2020).

52. J. Zorrilla de San Martin, F. F. Trigo, S. Kawaguchi, Axonal GABAA receptors depolarize presynaptic
terminals and facilitate transmitter release in cerebellar Purkinje cells. *The Journal of Physiology*. **595**,
7477–7493 (2017).

53. M. L. Hines, N. T. Carnevale, The NEURON Simulation Environment. *Neural Computation*. **9**, 1179–
1209 (1997).

54. R. I. Herzog, T. R. Cummins, S. G. Waxman, Persistent TTX-Resistant Na+ Current Affects Resting
Potential and Response to Depolarization in Simulated Spinal Sensory Neurons. *Journal of*
Neurophysiology. **86**, 1351–1364 (2001).

55. S. Ratté, S. A. Prescott, ClC-2 Channels Regulate Neuronal Excitability, Not Intracellular Chloride
Levels. *J. Neurosci.* **31**, 15838–15843 (2011).

56. D. Gentiletti, M. de Curtis, V. Gnatkovsky, P. Suffczynski, Focal seizures are organized by feedback
between neural activity and ion concentration changes. *eLife*. **11**, e68541 (2022).

57. A. Takeuchi, S. Tatsumi, N. Sarai, K. Terashima, S. Matsuoka, A. Noma, Ionic Mechanisms of Cardiac
Cell Swelling Induced by Blocking Na+/K+ Pump As Revealed by Experiments and Simulation. *Journal*
of General Physiology. **128**, 495–507 (2006).

58. P. Botta, F. M. S. de Souza, T. Sangrey, E. De Schutter, C. F. Valenzuela, Alcohol Excites Cerebellar
Golgi Cells by Inhibiting the Na+/K+ ATPase. *Neuropsychopharmacol*. **35**, 1984–1996 (2010).

Acknowledgments

729 We thank to Drs. M. Midorikawa and T. Inoshita for critical reading of the manuscript and helpful

730 comments.

Funding:

JSPS/MEXT KAKENHI Grant Numbers JP22H02721 (SK)

JSPS/MEXT KAKENHI Grant Numbers JP22K19360 (SK)

the Takeda Science Foundation (SK)

the Naito Foundation (SK)

Grant-in-Aid for JSPS Fellows Grant Numbers JP22J14296 (MM)

Grant-in-Aid for JSPS Fellows Grant Numbers JP22KJ1851 (MM)

Author contributions: MM performed most imaging and patch-clamp experiments and data analysis; RH performed model simulation and analysis; SK conceptualized and administrated the study; SK and MM interpreted data and wrote the manuscript. All authors approved the final version.

Competing interests: The authors declare no competing interests.

Data and materials availability: All plasmids used in this study except for those available at Addgene can be requested to the corresponding author. All data and any additional information required to reanalyze the data reported in this paper will be shared by the corresponding author upon request. Program codes for simulating the simple biophysical multi-compartment model will be available at the website of the corresponding author.

Fig. 1. Direction-dependent EPSP modulation in dendrites by voltage imaging

(A) Fluorescent image of a cultured hippocampal neuron expressing ASAP3β under the whole-cell patch clamp recording.

(B) Representative spontaneous EPSP (left) and action potential (right) in a cultured hippocampal neuron, recorded by ASAP imaging (black) and current-clamp recording (red).

(C) Relative fluorescence changes ($\Delta F/F$) for EPSPs (red, 29 events, 5 cells) and action potentials (blue, 12 events, 3 cells) plotted against the size of membrane potential changes (ΔV_m).

(D) Top, single-photon (405 nm) glutamate uncaging around a spine of an ASAP3β-expressing neuron under the whole-cell patch-clamp recording. Bottom, somatic voltage change upon glutamate uncaging recorded by current-clamp (red) or fluorescence imaging (black).

(E) Top, image of an ASAP3β expressing neuron. Yellow rectangular area is expanded below, showing branches used for voltage imaging of local glutamate-evoked EPSPs. Magenta point indicates the location of 405 nm spot laser illumination.

(F) $\Delta F/F_{eEPSP}$ traces at six ROIs (1-6) indicated in E.

(G) Top, Schematic drawing of classification of dendrites (black, dendrite on the way to soma; red, distal branches; blue, bifurcated branches on the way to soma). Bottom, $\Delta F/F_{eEPSP}$ sizes for individual eEPSPs (normalized to that at ROI 3) in a cell (show in E) are plotted against the distance from the uncaged site.

(H) Relative $\Delta F/F_{eEPSP}$ against the distance from the uncaged site (21 cells, red: 102, black: 85, blue: 226 regions based on the categorization shown in G). Square plots indicate averages ($\pm$ SEM).

(I) Absolute $\Delta F/F_{eEPSP}$ sizes at three categories of location: distal branch, uncaged site, and proximal branch (n = 21 cells, **: $P < 0.01$, ***: $P < 0.001$, paired t-test). Square plots indicate averages ($\pm$ SEM). See also fig. S1 and S2.

Figure 2. Symmetric attenuative propagation of GABAergic PSPs

(A) and (B) Left, dendritic segments of interest used for voltage imaging upon local GABA-evoked hyperpolarizing (A) or depolarizing (B) postsynaptic potentials (PSP$_{GABA}$). Magenta point or line indicates the location of GABA uncaging. Middle, $\Delta F/F$ for PSP$_{GABA}$ traces at five ROIs (4-8 for A, 2-6 for B) indicated in the left image. Right, relative $\Delta F/F$ for PSP$_{GABA}$ is plotted against the distance from the uncaged site (6 cells, red: 25, black: 18 regions for A, 8 cells, red: 33, black: 20 regions for B). Square plots indicate averages (± SEM). See also fig. S3.

Figure 3. [Cl$^-$]$_{in}$-dependent more negative membrane potential in dendrites

(A) Image of direct patch-clamp recording from an EGFP-expressing dendrite (right) and magnified view of the patched branch in yellow (left).

(B) V_m trace recorded from a dendritic branch (shown in A).

(C) Somatic (blue, n = 19 cells) and dendritic (green, n = 11 cells) Vm_{rest} without or with 10 μM VU0463271 (soma: yellow, n = 9 cells; dendrite: red, n = 11 cells). Tukey test, *: P < 0.05, **: P < 0.01, n.s.: not significant.

(D) and (E) Vm_{rest} (D) and leak conductance (E) plotted against the distance from soma. VU : in the presence of VU0463271.

(F) Images of immunostained KCC2 (magenta, top) in a GFP-labeled hippocampal neuron (green, middle). Three rectangular areas (a - c) are expanded below with color-merged.

(G) Ratio of fluorescent intensity of KCC2 to GFP in soma, proximal and distal dendrites (n = 9 cells, *: P < 0.05, n.s.: not significant, Tukey test). Square plots indicate averages (± SEM).

Fig. 4. Contribution of low [Cl⁻]$_{in}$ and TTX-resistant Na⁺ channels to EPSP amplification

(A-C) Left, Images for dendritic segment of interest (ROIs 1-5) and $\Delta F/F_{eEPSP}$ traces measured in the presence of VU0463271 (10 µM, A), lidocaine (5 mM, B) or both (C). Magenta points indicate the locations of glutamate uncaging. Middle, relative $\Delta F/F_{eEPSP}$ plotted against the distance from the uncaged site (17 cells, red: 44, black: 52 regions for A; 17 cells, red: 55, black: 56 regions for B; 15 cells, red: 38, black: 44 regions for C). Square plots indicate averages (± SEM). Right, absolute $\Delta F/F_{eEPSP}$ sizes in distal region of dendrites and uncaged sites (A, n = 17 cells; B, n = 17 cells; C, n = 15 cells). Paired t-test, n.s.: not significant. Square plots indicate averages (± SEM).

(D) Statistical summary of $\Delta F/F_{eEPSP}$ amplification in distal branches with or without VU0463271 (10 µM), lidocaine (5 mM), TTX (1 µM), Cd^{2+} (100 µM), D-AP5 (100 µM), ZD7288 (100 µM). Dunnett's test, *: $P < 0.05$, n.s.: not significant. Error bars are the SEM. See also fig. S4.

Figure 5. Role of Cl⁻- and Na⁺ channels in low Vmrest and local EPSP amplification

(A) Immunostained ClC-2 image in a hippocampal neuron.

(B) Hypothetical model for mechanisms of EPSP amplification in distal dendrites.

(C) Geometric design of a multi-compartment model equipped with the deep V_m-activated Cl⁻ conductance and the low-threshold Na$_V$ channels. Dendrites are composed of 18 compartments, and an excitatory synaptic input was given at the compartment #6, #7, or #8. See also table S1.

(D) Simulated EPSP traces (caused at #7) at five representative compartments (#1,4,7,9,12) indicated in C, with or without the 50 % inhibition of Na$_V$ and/or the abolishment of deeper V_m-activated Cl⁻ conductance change.

(E) Relative amplitude of simulated EPSPs caused at #6 (right), #7 (middle), or #8 (left) plotted against the distance from the input site in 4 conditions of the model.

Figure 6. Validation of a model prediction: critical role of Cl⁻ conductance in EPSP amplification
(A) and (B) Left, stimulated dendritic segment in the presence of NPPB (50 μM) & lidocaine (5 mM, A), or GABA (50 μM, B). Magenta points indicate the locations of glutamate uncaging. Middle, $\Delta F/F_{eEPSP}$ traces at five ROIs (1-5) indicated in the left image. Right, relative $\Delta F/F_{eEPSP}$ plotted against the distance from the uncaged site (13 cells, red: 37, black: 34 regions for A; 10 cells, red: 29, black: 31 regions for B). Square plots indicate averages (± SEM).
(C) Statistical summary of $\Delta F/F_{eEPSP}$ amplification in distal region with or without NPPB & lidocaine, and GABA (**: P < 0.01, ***: P < 0.001, Dunnett's test). Error bars are the SEM.

Figure 7. Dynamic spatio-temporal integration of synaptic inputs

(A) Left, image of an ASAP3β-expressing neuron, with yellow rectangular area expanded in right, showing 10 sites of local glutamate uncaging (magenta, 35 ms intervals) in three patterns: to distal (a to j), to soma (j to a) or random.

(B) Averaged $\Delta F/F_{eEPSP}$ traces (± SEM, 17 cells) at three ROIs (uncaged branch: black, proximal: yellow, distal: red as indicated in A right) upon three patterns of glutamate uncagings. Magenta lines indicate spot laser illumination.

(C) $\Delta F/F_{eEPSP}$ traces (mean ± SEM) at ROI$_{distal}$ in three conditions: control (red, 17 cells), with lidocaine (5 mM) & VU0463271 (10 μM) (blue, 13 cells), and with Cd^{2+} (100 μM) & D-AP5 (100 μM) (green, 14 cells).

(D) Comparison of synaptic integration at ROI$_{distal}$ upon three patterns of glutamate inputs in the absence (ctrl, top) or presence of lidocaine and VU0463271 (bottom). Two-way ANOVA, **: $P < 0.01$, ***: $P < 0.001$, n.s.: not significant.

(E) Ratio of $\Delta F/F_{eEPSP}$ signals (mean ± SEM) upon opposite directions of glutamate uncagings ("to soma" relative to "to distal") at ROI$_{distal}$ in the absence (ctrl) or presence of lidocaine & VU0463271. Two-way ANOVA, **: $P < 0.01$. See also fig. S5.