Deciphering deep-sea chemosynthetic symbiosis by single-nucleus RNA-sequencing

Hao Wang[1,2,3,4#], Kai He[5#], Huan Zhang[1#], Quanyong Zhang[6#], Lei Cao[1], Jing Li[1,7], Zhaoshan Zhong[1], Hao Chen[1], Li Zhou[1], Chao Lian[1], Minxiao Wang[1], Kai Chen[6], Pei-Yuan Qian[3,4*], and Chaolun Li[1,7,8*]

[1]Center of Deep-Sea Research, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China

[2]Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China

[3]Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou), Nansha, Guangzhou, China

[4]Department of Ocean Science, Hong Kong University of Science and Technology, Hong Kong S.A.R, China.

[5]Key Laboratory of Conservation and Application in Biodiversity of South China, School of Life Sciences, Guangzhou University, Guangzhou, Guangdong 510006, China

[6]State Key Laboratory of Primate Biomedical Research, Institute of Primate Translational Medicine, Kunming University of Science and Technology, Kunming, China

[7]South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, China

[8]University of Chinese Academy of Sciences, Beijing, China

Correspondence authors: Prof. Pei-yuan Qian and Prof. Chaolun Li

Email addresses: boqianpy@ust.hk and lcl@qdio.ac.cn

#These authors contributed equally to this work

The authors declare no conflict of interest.

Abstract:

Bathymodioline mussels dominate deep-sea methane seep and hydrothermal vent habitats and obtain nutrients and energy primarily through chemosynthetic endosymbiotic bacteria in the bacteriocytes of their gill. However, the molecular mechanisms that orchestrate mussel host-symbiont interactions remain unclear. Here, we constructed a comprehensive cell atlas of the gill in the mussel *Gigantidas platifrons* from the South China Sea methane seeps (1100m depth) using single-nucleus RNA sequencing (snRNA-seq) and whole-mount *in situ* hybridisation. We identified 13 types of cells, including three previously unknown ones, and uncovered unknown tissue heterogeneity. Every cell type has a designated function in supporting the gill's structure and function, creating an optimal environment for chemosynthesis, and effectively acquiring nutrients from the endosymbiotic bacteria. Analysis of snRNA-seq of *in situ* transplanted mus-sels clearly showed the shifts in cell state in response to environmental oscillations. Our find-ings provide insight into principles of host-symbiont interaction and the bivalves' environmental adaption mechanisms.

Introduction:

Mutualistic interactions between multicellular animals and their microbiota play a fundamental role in animals' adaptation, ecology, and evolution [1,2]. By associating with symbionts, host animals benefit from the metabolic capabilities of their symbionts and gain fitness advantages that allow them to thrive in habitats they could not live in on their own[1,3]. Prime examples of such symbioses are bathymodioline mussels and gammaproteobacteria endosymbionts[4]. The bathymodioline mussels occur worldwide at deep-sea chemosynthetic ecosystems, such as cold-seeps, hydrothermal vents, and whale falls[4]. The host mussel acquired sulfur-oxidising (SOX) and/or methane-oxidising (MOX) symbiont through horizontal transmission at their early life stage[3]. Their symbionts, which are hosted in a specialised gill epithelial cell, namely, the bacteriocytes[5,6], utilise the chemical energy from the reduced chemical compounds such as CH_4 and H_2S released from cold seeps or hydrothermal vents to fix carbon and turn into carbon source for the host mussel[7,8]. The ecological success of the bathymodioline symbioses is apparent: the bathymodiolin mussels are often among the most dominant species in the deep-sea chemosynthetic ecosystems[9,10]. Thus, it is critical to know how the bathymodioline mussels interact with the symbionts and maintain the stability and efficiency of the symbiosis.

The gill structure of bathymodiolin mussels has undergone remarkable adaptations at the molecular, cellular, and tissue levels to support their deep-sea chemosynthetic lifestyle[11,12]. Compared to shallow water mussels[13], the gill filaments of bathymodiolin mussels have enlarged surfaces allowing them to hold more symbionts per filament (Supplementary Fig. S1). This adaption requires not only novel cellular and molecular mechanisms to maintain and support the enlarged gill filament structure, but also a strong ciliary ventilation system to pump vent/seep fluid to fuel the symbiotic bacteria. Previous studies based on whole-genome sequencing and bulk RNA-seq projects have shown that genes in the categories of nutrient transporters, lysosomal proteins, and immune receptors are either expanded in the host mussel's genome or up-regulated in the gill[12, 14-17], suggesting that these genes involved in host-symbiont interaction. Though providing deep molecular insights, these studies mainly used homogenised tissues that average genes' expression levels amongst different cell types and eliminate cell and gene spatial distribution information[18-21]. In addition, the broad expression and function of po-

tentially 'symbiosis-related' proteins also greatly limited data interpretation. Therefore, a sys-

temic atlas of gill cell types and the descriptions of cell-type-specific gene expressional profiles

are warranted to a better understanding of the host-symbiont interaction and environmental

adaptation mechanisms of the bathymodioline symbiosis.

In recent years, single cell/nucleus RNA-sequencing (sc/sn RNA-seq) technologies have be-

come one of the preferred methods for investigating the composition of complex tissue at the

transcriptional level[22]. snRNA-seq has several advantages, such as compatibility with frozen

samples, elimination of dissociation-induced transcriptional stress responses, and reduced dis-

sociation bias[23]. These advantages are significant for deep-sea and other ecological studies be-

cause cell and molecular biology facilities are not available in the field. To examine the cellular

and molecular mechanisms of the environmental adaptations and host-symbiont interactions in

bathymodioline mussels, we conducted a single nucleus RNA-sequencing-based transcriptomic

study. We analysed the gill symbiosis in the dominate deep-sea mussels inhabiting the F-site

cold-seep *Gigantidas platifrons*[24], which hosts a single a MOX endosymbiont population that

consist of only one 16S rRNA phylotype. Our work provides a proof-of-principle for environ-

mental adaptation mechanisms study in non-traditional but ecologically important organisms

with single-nucleus RNA-sequencing technologies.

Results and discussion

1. *G. platifrons* deep-sea *in situ* transplant experiment and single-cell transcriptomic sequencing:

We conducted a *G. platifrons* deep-sea *in situ* transplant experiment at the "F-Site" Cold-seep (~1117 m depth) and retrieved three groups of samples (Fig. 1A), as follows: the 'Fanmao' (meaning prosperous) group, which comprised the mussels collected from methane-rich (~40 µM) site of the cold-seep, where the animals thrived; the 'starvation' group, which comprised the mussels collected from the methane-rich site and then moved to a methane-limited (~0.054 µM) starvation site for 14 days before retrieval; and the 'reconstitution' group, which comprised the methane-rich site mussels moved to the 'Starvation' site for 11 days and then moved back to the methane-rich site for another 3 days.

We next performed single nucleus RNA-sequencing (snRNA-seq) of the gill posterior tip of the three groups of samples using the microwell-based BD Rhapsody platform[™] Single-Cell Analysis System (Fig. 1A). After quality control, we obtained 9717 (Fanmao), 21614 (Starvation), and 28928 (Reconstitution) high-quality single nuclei transcriptomic data.

2. Cell atlas of *G. platifrons* gill

To unravel the intricate cellular composition, we utilised a reciprocal PCA (RPCA) strategy to project cells in the three samples onto the same space based on conserved expressed genes among them[25]. This strategy maximises the number of cells per cluster regardless of the organism's state and therefore maximises genes per cluster[26].

Through the implementation of Seurat, we revealed 14 cell clusters, each was associated with a set of marker genes (Figs. 1B- D) [27,28]. Given the limited availability of canonical marker genes for *G. platifrons* and molluscs in general, we undertook a meticulous approach to characterize each cell cluster. This involved: 1) examining the cluster marker genes' functions, 2) identifying the expression pattern of cluster marker gene using whole-mount *in situ* hybridisation (WISH) and/or double-fluorescent *in situ* hybridisation (FISH) analyses, and 3) conducting scanning electron microscopy (SEM) and transmission electron microscopy (TEM) analyses. We successfully identified and characterized 13 of the 14 cell clusters, which could be categorized into

four major groups, including 1) the supportive cells, 2) the ciliary cells, 3) the proliferation cells,

and 4) the bacteriocytes. One cell cluster remained ambiguous. To assess the robustness of

each cell cluster, we employed a bootstrap sampling and clustering algorithm examining the

similarity among clusters and obtained strong support for all clusters through a combined

analysis of the three samples (Supplementary Fig. S2A). Furthermore, when examining each

sample individually, we found that the majority of clusters demonstrated robust support, with

the exception of the three ciliary cell clusters, which showed overlaps of assignment probabili-

ties among them (Supplementary Fig. S2 B-D). These three cell types are derived from a same

precursor, and exhibited relatively lower numbers of nuclei, resulting in a reduced availability of

genes per cell type, which is particularly true for the food grove ciliary cell (Supplementary Ta-

ble S1). By integrating the spatial and functional annotations of these cell clusters, we gained

insight into their collective efforts in supporting the symbiotic relationship and maintaining the

high-efficiency chemosynthetic system within the gill tissue.

Supportive cells: We have identified four supportive cell types, namely, inter lamina cells

basal membrane cells (BMC)1, BMC2, and mucus cells (Figs 2A, B and C).

The inter lamina cells, marked by high expression of fibrillar-forming collagens, were the

cells located between the two layers of basal membranes (Fig 2D and E). These cells were pre-

viously identified as amoeboid cells[13], and their function has not been explored. The inter lam-

ina cells were densely distributed around the food groove and at the rim of the gill filament.

While on the middle part of the gill filament, the inter lamina cells distributed in parallel rows. It

might help to connect the two sheets of basal membranes and maintain the spatial integrity of

the gill filament. In addition, we observed the enrichment of ribosomal proteins involved in

protein synthesis (Supplementary Table S2), indicating a high metabolic rate[29] of inter lamina

cells. This implies that the inter lamina cells may process the nutrients acquired from the sym-

biont.

We have also identified two types of BMC populations which have not been recognised be-

fore. The BMC1 (Fig. 2F and G) expressed genes encoding extracellular matrix and adhesive pro-

teins (Fig. 2B and Supplementary Table S2), such as SCO-spondin (Bpl_scaf_27638-4.18),

Zonadhesin (Bpl_scaf_1371-2.4), Cartilage matrix proteins (Bpl_scaf_16371-0.14 and

Bpl_scaf_3387-5.47), and Collagen alpha-1(XXI) chain protein (Bpl_scaf_51350-0.19). BMC2 (Fig.

2H and I) was distinctive from BMC1 in the context of expression (Fig 2A) and marker genes (Fig

2B). Genes encoding hemicentins (Bpl_scaf_38731-0.50 and Bpl_scaf_7293-0.14), which are

known to stabilise the cells' contact with the basal membrane[30], were highly expressed in BMC2.

Thus, these two cell types are likely to help building the basal lamina, and stabilising the epithe-

lial-derived cells, such as bacteriocytes and intercalary cells, on the surface of the basal mem-

brane.

Mucus cells are specialised secretory cells with intracellular mucus vacuoles[13]. In shallow-

water filter-feeding bivalves, mucus cells secret mucus, which cooperates with the ciliary venti-

lation system to capture, process, and transport food particles to their mouths, known as filter

feeding mechanism[31-34]. Deep-sea mussels may not necessarily retain this function since they

do not normally acquire food resources such as phytoplankton and planktonic bacteria but ob-

tain most of nutrient through their chemosynthetic symbiont. Thus, it was hypothesised that

mucus cells are involved in other biological functions such as immune responses to pathogens[18].

Herein, genes encoding proteins with microbe-binding functionalities were enriched in mucus

cells (Fig. 2B and Supplementary Table S2), such as C-type lectins (Bpl_scaf_58694-0.22 and

Bpl_scaf_38133-0.8), C1q domain-containing protein (Bpl_scaf_55216-2.11), and Fibrinogen C

domain-containing protein (Bpl_scaf_18519-2.15)[35,36]. The expression of similar immune genes

was upregulated in a shallow water mussel *Mytilus galloprovincialis* when challenged by a

pathogenic bacteria[37]. Our WISH and double-FISH analyses showed that mucus cells were em-

bedded within the outer rim cilia and scattered on the gill lamella alongside the bacteriocytes

(Figs. 2J and 2K). These data collectively suggest that mucus cells may help mussel maintain the

immune homeostasis of gill. Interestingly, WISH and double-FISH analyses showed that mucus

cells were also distributed alongside the gill lamella's food groove and the inner edge, where

the density of bacteriocytes is low (Fig. 2J). Because the food groove is the main entry of food

practical to the labial palps and mouth[38], this distribution pattern implies that mucus cells may

be also involved in capturing planktonic bacteria and sending them to the mouth.

Ciliary and smooth muscle cells: A remarkable feature of bathymodioline mussel's gill is its

ciliary ventilation system, which constantly agitates the water and provides the symbiont with

the necessary gas[39]. We identified four types of ciliary cells (Figs. 3A and 3B): apical ridge ciliary

cells (ARCCs), food groove ciliary cells (FGCCs), lateral ciliary cells (LCCs), and intercalary cells

(ICs), as well as a newly identified type of smooth muscle cells. All ciliary cells were marked by

canonical cilium genes, such as genes encoding flagellar proteins, ciliary motor kinesin proteins,

ciliary dynein proteins, and ciliary microtubules and were clearly distinguishable by specifically

expressed genes (Supplementary Table S3 and Supplementary Fig. S3). ARCCs were character-

ized by expression of Tubulin alpha-1A chain Bpl_3489-0.37 (Fig. 3C) and Bpl_scaf_20631-1.16,

which encodes homeobox Dlx6a-like protein (Fig. 3D) which is a marker of the apical ectoder-

mal ridge[40]. FGCCs, which could be ladled by expression of marker gene Bpl_scaf_5544-0.0 (Fig.

3E), expressed genes encoding primary cilia development regulator Tubby-related protein 3[41]

(Bpl_scaf_24834-2.3) and primary ciliary cell structural protein TOG array regulator of axonemal

microtubules protein 1[42,43] (Bpl_scaf_55620-1.3), suggesting the FGCCs are the sensory ciliary

cells that gather information from the surrounding environment. Both ARCCs and FGCCs were

located around the ventral tip of the gill filament (Fig. 3F). LCCs were distributed as two parallel

rows along the gill lamella's outer rim and ciliary disks' outer rim (Figs. 3G and 3H) and had

highly expressed genes involved in cilium structure, cilium movement and ATP synthases (Fig.

3B and Supplementary Table S3), indicating that LCCs may have a strong ability to beat their

cilia. Interestingly, we identified a group of previously unreported smooth muscle cells (SMCs)

co-localised with the LCCs as showed by WISH (Fig. 3I and Supplementary Fig. S4). SMCs

strongly expressed several Low-density lipoprotein receptors (LDL receptor) and LDLR-

associated proteins[44] (Fig. 3B). In addition, these cells also expressed the angiotensin-

converting enzyme-like protein[45] (Bpl_scaf-29477-5.10) and the 'molecular spring' titin-like pro-

tein[46] (Bpl_scaf_56354-7.8) (Supplementary Table S3). The expression of these genes could be

commonly found in human vascular smooth muscle cells[47-49]. Collectively, we suspect that SMCs

are involved in lateral cilium movement and the gill slice contraction.

ICs are the specialised ciliary cells surrounding the bacteriocytes (Figs. 3B and 3J). WISH

analysis showed that the expressions of IC markers had apparent spatial variations (Fig. 3J). The

expression of ICs marker rootletin[50-52] (Bpl_scarf_22362-6.9) was considerably higher in the ICs

close to the frontal edge of the gill filament, and the expression gradually decreased along with

the direction of inter lamina water flow (Fig. 3K and Supplementary Fig. S5), implying that ICs

ventilate the water flow and the mucus through the gill filaments. Furthermore, compared with

the other three types of ciliary cells, the ICs expressed several genes encoding transcription fac-

tors involved in determining cell fate (Fig. 3B and Supplementary Table S3), such as transcrip-

tion factors Sox 8 (Bpl_scarf_40595-5.30) and Wnt pathway cell polarity regulator secreted friz-

zled-related protein 5 (Bpl_scaf_57424-6.1)[53], suggesting that the ICs might also play regulatory

roles[54-56].

Proliferation cells: It has long been known that the bathymodioline mussel gill has three

types of proliferation cells that are conserved throughout all filibranchia bivalves: the budding

zone at the posterior end of the gill where new gill filaments are continuously formed, and the

dorsal and ventral ends of each gill filaments[57,58]. These "cambial-zone"-like cell populations

could continuously proliferate throughout the whole life span of the mussel[58]. Our snRNA-seq

data recognised three types of proliferation cells, which is consistent with previous findings (Fig.

4A, Fig. 4B and Supplementary Table S4). The gill posterior end budding zone cells (PEBZCs) are

located on the first few freshly proliferated filaments of the posterior tip of gill (Figs. 4C and 4D).

The PBEZCs marker (Bpl_scaf_61993-0.4) gradually disappeared at around the 11th -12th row

of the gill filament (Fig. 4D), suggesting the maturation of the gill filaments, which is similar to

the developmental pattern reported in another deep-sea mussel *Bathymodiolus azoricus* [59]. The

dorsal end proliferation cells (DEPCs), which expressed the hallmarks genes of muscular tissue

(Fig. 4B), as well as cell proliferation and differentiation regulators, were the proliferation cells

in connective tissue at the dorsal end of the gill slice (Fig. 4A, 4E). The ventral end proliferation

cells (VEPCs) were two symmetrical triangle-shaped cell clusters of small symbiont-free cells

(Fig. 4F). In VEPCs, genes encoding ribosomal proteins, chromatin proteins, RNA and DNA bind-

ing proteins, and cell proliferation markers were all up-regulated (Fig. 4B and Supplementary

Table S4), indicating that VEPCs are meristem-like cells[59,60].

It has been hypothesised that new proliferation cells will be colonised by symbionts, serving

as the vital mechanism of bacteriocyte recruitment[59]. We then determined which proliferation

cell type gives rise to the mature cells, especially the bacteriocytes, which has little available

information regarding their precursor and transmission mode in bathymodioline mussels [59,61].

Slingshot trajectory analysis suggested that the PEBZCs might be the origin of all gill epithelial

cells, including the other two proliferation cells (VEPC and DEPC) and bacteriocytes (Supple-

mentary Fig. S6). The sole exception was BMC2, which may be derived from VEPC rather than

PEBZC. This result is consistent with previous studies which suggested new gill filaments of the

filibranch mussels are formed in the gill's posterior budding zones[59]. The colonisation by the

symbiont might play a crucial role in determining the fate of the bacteriocytes. Noticeably, in *G.*

platifrons, only the pillar-shaped first row of gill filament, comprised of small meristem-like cells,

was symbiont-free (Figs. 4D and D'), whereas all the other gill filaments were colonised by sym-

bionts. This pattern of symbiosis establishment is different from that of *B. azoricus*, in which

PEBZCs are symbiont free and are gradually colonised by symbionts released from the bacterio-

cytes on the adjacent mature gill filaments after maturation[59]. On mature gill filaments, the

DEPCs and VEPCs are seemingly the sources of new cells that sustain the growth of the gill fila-

ment from both dorsal and ventral directions, respectively[57]. Interestingly, comparable active

ventral and dorsal end proliferation zones have also been identified in the symbiotic mussel *B.*

azoricus, whereas they are absent in the shallow-water mussel *Mytilus edulis*[62]. This contrast

further suggests the potential involvement of DEPCs and VEPCs in the establishment of symbio-

sis.

3. Bacteriocytes and host-symbiont interaction:

We conducted the whole-mount FISH using a bacterial 16S rRNA probe for symbiont to de-

termine the spatial distribution of bacteriocytes, and their positions relative to the other cell

types on the gill filament (Figs. 5A, B). Bacteriocytes covered the majority of the surface of the

gill filament, except the ventral tip, the ciliary disk, and the frontal edge (lateral ciliary). The

bacteriocytes were surrounded by intercalary cells with microvilli and cilia on the surface (Fig.

5C).

The gene expression profile of bacteriocytes aligned well with the morphological analysis,

which suggested that the bacteriocytes have structural, metabolic, and regulatory adaptions to

cope with the symbiont (Figs. 5D, 5F and 5I). It has been hypothesised that the bacteriocytes

extract nutrients from the symbiont through two pathways: lysing the symbiont through the

lysosome (the 'farming' pathway) or directly utilising the nutrient produced by the symbiont

(the 'milking' pathway)[63,64]. Our TEM observations clearly detected intracellular vacuole and

lysosome system of the bacteriocyte that harbour, transport, and digest the symbionts (Fig. 5D),

which is consistent with previous studies[18,65]. The snRNA data showed bacteriocytes expressed

cellular membrane synthesis enzyme (phosphoethanolamine N-methyltransferase,

Bpl_scaf_15282-3.10) and a series of lysosomal proteins such as lysosomal proteases (cathep-

sins; Bpl_scaf_61711-5.12, Bpl_scaf_46838-6.40 and Bpl_scaf_59648-4.5; lysosomal alpha-

glucosidase-like), protease regulators 56 (TMPRSS15 proteins; Bpl_scaf_52188-1.15 and

Bpl_scaf_15410-0.9), and lysosomal traffic regulator proteins (rabenosyn-5, Bpl_scaf_54816-0.3

and Bpl_scaf_52809-1.6, lysosomal-trafficking regulator-like isoform X1). Among the proteases,

cathepsins were thought to be evolutionary conserved molecular tools that host utilized to con-

trol the residence of their symbiont microbes[66]. They were also highly expressed in symbiotic

tissue of other deep-sea chemosynthetic animals, such as vesicomyid clams and vestimen-

tiferan tubeworms[67-69]. Bacteriocytes also expressed genes encoding cellular vesicle transports

(kinesins, Bpl_scaf_14819-0.11, Bpl_scaf_54265-1.13, and Bpl_scaf_4784-1.40)[53], potential

amino acid transporter[57] (Bpl_scaf_36159-5.5), and genes involved in intracellular vesicle trans-

port, such as the FYVE and coiled-coil domain-containing protein 1[58] (Bpl_scaf_33726-5.7). In

addition to form and mobilise early endosomes, the protein products of these genes could

transport symbiont-secreted nutrient vesicles to the host (Fig. 5F and Supplementary Table S5),

supporting the "milking" pathway.

The symbiont of *G. platifrons* belongs to type I methanotrophy, of which the core metabolic

function is linked with the development of intracytoplasmic membranes leading to a high

lipid/biomass content[70,71]. Recent lipid biomarker analyses showed that the gill of *G. platifrons*

contains a high amount of bacterial lipids, which are directly utilised by the host to synthesise

most of its lipid contents[67]. Downstream of the bacteriocyte's metabolic cascade, genes encod-

ing proteins that may be involved in fatty acid/lipid metabolism, such as perilipin2

(Bpl_scaf_27158-3.8), which is the critical protein to form intracellular lipid droplets[72], and a

variety of fatty acid metabolism enzymes (acetyl carboxylase 2, Bpl_scaf_55250-5.11[73]; fatty

acid desaturase 1-like isoform X1, Bpl_scaf_35916-0.6[74,75]; long-chain fatty acid-ligase ACSBG2-

like isoform X1, Bpl_scaf_28862-1.5 [76,77]), were up-regulated, suggesting that the fatty acid

could be a major form of nutrients passing from the symbiont to the host mussel.

Additionally, bacteriocytes expressed several solute carriers, including sodium/ascorbate co-

transporter (Fig. 5I, solute carrier family 23 members 1-like, Bpl_scaf_32311-1.19), so-

dium/potassium/calcium exchanger (sodium potassium calcium exchanger 3-like,

Bpl_scaf_14503-1.28), sodium/chloride ion cotransporter (solute carrier family 12 member 3-

like isoform X1, Bpl_scaf_44604-3.7), ferrous iron transporter (solute carrier family 40 member

1-like, Bpl_scaf_63447-0.12) and zinc transporter (zinc transporter ZIP14-like, Bpl_scaf_44428-

5.3). The solute carriers are a large family of ATP-dependent transporters that shuttle a variety

of small molecules across the cellular membrane[78]. In several model symbiotic systems, solute

carriers play a vital role in host-symbiont interaction by either providing the symbiont with sub-

strates[79,80] or transporting symbiont-produced nutrients to the host [81-83]. Previous studies dem-

onstrated that solute carrier genes were expanded in deep-sea chemosynthetic animals' ge-

nome[84] and highly expressed in symbiotic organs[85] including bathymodioline mussel's gill[12]. In

bathymodioline mussels, previous bulk RNA-seq studies detected up-regulated expression of a

large variety of solute carriers in the gill, which is consistent with the present study, suggesting

that solute carriers may play crucial roles in shuttling nutrients in and out of bacteriocytes and

in maintaining the suitable intracellular micro-environment (such as the SLC23A1 and SLC39A14)

for the symbiont[86,87].

4. Cell-type-specific response to environmental stresses:

To examine cell-type-specific acclimatisation to environmental changes the expression pro-

files of differentially expressed genes (DEGs) were compared between the three states (Fanmao,

starved and reconstituted animals) of samples collected in our *in situ* transplantation experi-

ment (full lists of DEGs are shown in Supplementary Table S6-S18). Notably, for each state three

mussels were processed but pooled for nuclei extraction before snRNA sequencing (see

Method for detail). We tested our hypothesis here that at the starvation site, a relatively low

concentration of methane shall upset symbiont metabolism and thus substantially affect sym-

biont-hosting bacteriocytes by assessing the transcriptional changes per cell type. We calcu-

lated the distances between the three sets of samples on the UMAP plot (Fig. 6A) estimating

the centroid coordinates per sample per cell type, and calculating the Euclid distance between

the centroid coordinates of each pair of states per cell type (Supplementary Table S19). The im-

pact of starvation was variable across cell types, as reflected by the cross-state distances (Fig.

6B, green bar). Starvation resulted in the most significant transcriptional changes in bacterio-

cytes reflected by a large Fanmao-vs-starvation distance (2.3), followed by VEPC and inter lam-

ina cells (2.1 and 1.4, respectively). On the other hand, starvation had little impact on the tran-

scriptions of ciliary cells and most supportive cells such as the food grove ciliary cell, BMC2, and

mucus cells (distances <0.5). Fig 6B also shows that after reconstitution (moving the mussels

back to the methane-rich site Fanmao for 3 days), the expressional profile of bacteriocytes rap-

idly changed back, reflected by a large starvation-vs-reconstitution distance and much smaller

Fanmao-vs-reconstitution distance (2.3 vs. 0.8; Suppl. Tab. S19). This result coincided with and

was supported by our pseudo-time analysis for bacteriocytes (Fig. 6C), showing that bacterio-

cytes in the reconstitution are in intermediate and transitional states between Fanmao and

starvation.

For the bacteriocytes population, we conducted cell trajectory pseudo-timing and detailed

DEG analysis to interpret the mechanism of host-symbiont interactions. It is important to

acknowledge that our sampling strategy might have limitations, as more closely spaced time

points could enhance the confidence of trajectory reconstruction. The branched heatmap

showed both up- and down-regulated genes during starvation and reconstitution compared

with the Fanmao state (Fig. 6D and supplementary Fig. S7). The Kyoto Encyclopedia of Genes

and Genomes (KEGG) pathway enrichment analysis of the bacteriocytes' DEGs provides an

overall view of the pathways enriched in each environmental state (Supplementary Fig. S8 A-C).

The genes encoding ribosomal proteins are highly expressed in the bacteriocytes under the

methane-rich 'Fanmao' state (Supplementary Fig. S8A), suggesting an active protein synthesis

and cellular metabolism [88]. Moreover, organic ion transporters (A BCB P-glycol,

Bpl_scaf_18613-16.10; canalicular multispecific organic anion transporter 2, Bpl_scaf_52110-

4.43; sodium- and chloride-dependent glycine transporter 1-like, Bpl_scaf_13642-5.8), which

may be involved in transporting symbiont-produced nutrients, are also highly expressed (Sup-

plementary Fig. S9).

In starved *G. platifrons*, negative regulators of cell proliferation, such as autocrine prolifera-

tion repressors and receptor-type tyrosine phosphatase beta, were up-regulated in bacterio-

cytes, suggesting repression of cell growth. We also observed the enrichment of genes in the

apoptosis pathway (Supplementary Fig. S8B), which is attributed to the upregulation of cas-

pases (Bpl_scaf_25165-3.8 and Bpl_scaf_24225-0.12) and cathepsins (Bpl_scaf_64706-0.16,

Bpl_scaf_61711-5.12) which can trigger caspases-dependent cell death, suggesting cellular

stress condition. Correspondingly, a gene encoding baculoviral IAP repeat-containing protein

(Bpl_scaf_6172-0.3), which can bind caspase and inhibit apoptosis, was up-regulated in bacte-

riocytes (Fig. 6E). In *Drosophila*, the baculoviral IAP proteins are important in animal's response

to cellular stress and promote cell survival[89,90]. Similarly, the gene encoding MAP3K14

(Bpl_scaf_60908-4.14) and potential E3 ubiquitin ligase RNF213 (Bpl_scaf_25983-1.26,

Bpl_scaf_1894-1.45, Supplementary Fig. S10), which were apoptosis suppressor or regulator,

were up-regulated[91]. These results suggest that starved bacteriocytes were carrying out cell-

type-specific adjustments to cope with stresses. The E3 ubiquitin ligases could also work as in-

tracellular immune sensors of bacterial lipopolysaccharides[92]. Thus, the encoded protein may

be able to active downstream immunological toolkit to digest the symbiont population for nu-

trients.

As mentioned above, bacteriocytes obtain nutrients from endosymbionts. KEGG pathway

analyses suggested that phagosome, lysosome-related pathways were upregulated in the star-

vation state, indicating that bacteriocytes were actively digesting the endosymbionts in the

starved *G. platifrons*. Interestingly, protein synthesis activity was more activated in inter lamina

cells and VEPCs as supported by the up-regulation of genes encoding ribosomal proteins in both

cell populations (Supplementary Tables S6 and S17). This was contrary to the situation in bacte-

riocytes and may be a consequence of the high activity of "farming" in bacteriocytes.

We anticipated that after moving the starved *G. platifrons* back to Fanmao site, bacterio-

cytes and their endosymbionts would "reconstitute" leading to the partly restored "farming"

and "milking" pathways. This was confirmed by higher expression of fatty acid metabolic genes

in the mussels in the reconstitution state than those in the starvation state, such as long-chain-

fatty-acid-ligase ACSBG2-like (Bpl_scaf_28862-1.5), elongation of very long-chain fatty acids 7-

like (Bpl_scaf_5959-1.9), and fatty acid desaturase 1-like (Bpl_scaf_35916-0.6). These findings

were also consistent with the result of KEGG analyses. The mitochondrial trifunctional enzyme

(Bpl_scaf_42376-0.5) was also highly expressed, suggesting a high level of energy-producing

activity. The KEGG analyses showed that glutamatergic synthase was enriched in the reconstitu-

tion state in comparison with those in both the Fanmao and starvation states. In the insect

aphid—Buchnera endosymbiosis model, the host-produced glutamate could be transported and

directly utilised by the symbiont[93]. Similar glutamate-based host—symbiont metabolic interac-

tion mechanisms were also proposed in the deep-sea mussel *Bathymodiolus thermophilus* and

B. azoricus[64,94]. As the pseudotime analyses showed above (Fig. 6C), reconstitution was the in-

termediate state between Fanmao and starvation. Metabolic and gene regulatory functions of

bacteriocytes were still different from that in Fanmao. The KEGG analyses suggested regulatory

(Rap1 signalling, retrograde endocannabinoid signalling, chemokine signalling, glucagon signal-

ling, thyroid hormone synthesis, mRNA surveillance pathway, etc.) and metabolic pathways

were enriched in the reconstitution state (fatty acid metabolism, salivary secretion, aldosterone

synthesis and section, insulin secretion and pancreatic section) (Supplementary Fig. S8C). For

example, we detected up-regulation of the genes encoding carbonic anhydrases

(Bpl_scaf_33596-7.3 and Bpl_scaf_48274-0.3), electroneutral sodium bicarbonate exchanger 1

(Bpl_scaf_61230-5.11), and globin-like proteins (Bpl_scaf_50392-1.14 and Bpl_scaf_24370-0.5),

which could provide the symbiont with carbon dioxide and oxygen necessary for the symbiont's

chemosynthetic metabolism[84,95].

5. Summary and outlook:

Using the deep-sea mussel *G. platifrons* as a model organism, we demonstrated the power

of integrating snRNA-seq and WISH data in unravelling the mechanism behind animal-microbe

symbiosis. The robustness of this strategy showed stable and highly distinguishable expression

patterns of each cell type regardless of the different environmental states. We successfully pro-

filed the specific roles of different types of cells, including the previously unknown cell types, in

maintaining the structure and function of the gill. The supportive cells that located in between
(inter lamina cells) and on the basal membrane's surface (BMC1 and BMC2) helped maintain
the anatomical structure of the basal membrane. Ciliary and smooth muscle cells involved in gill
slice contraction and cilium beat allowing the gathering of material and information from the
surrounding environment. Proliferation cells (PBEZCs, DEPCs and VEPCs) gave rise to new cells,
including bacteriocytes which obtained nutrients from endosymbionts using intracellular vacu-
oles and lysosomes.

The snRNA analyses also revealed that different cell types collaborated to support bacterio-
cytes' functionality. The PBEZCs gave rise to new bacteriocytes that allowed symbiont colonisa-
tion. Bacteriocytes attached to the basal membrane and were stabilised by supportive cells.
Mucus cells co-localise with bacteriocytes and help to maintain immune homeostasis. The beat-
ing ciliary cells controlled the water flow, providing bacteriocytes with necessary inorganic sub-
stances from the environment. This new information on cell-cell interaction certainly advanced
our overall understanding of how endosymbiotic microbes and host cells communicate and col-
laborate, which cannot be easily achieved through other methods.

Moreover, the analysis of snRNA data from bacteriocytes has provided insight into the mo-
lecular mechanisms employed by the host to maintain and regulate the symbiosis. Notably, the
bacteriocytes-enriched transcripts involved in harbouring, digesting symbionts, and transport-
ing nutrients produced by the symbionts were identify and characterised. Our *in situ* transplant
experiments by moving mussels between methane-rich and methane-limited sites also pro-
vided clues of cell-type specific responses to environmental change. Under a methane-limited
environment, the staved mussels more actively consumed endosymbionts through the "farm-
ing" pathway. After being moved back to the methane-rich environment, mussels produced
more glutamates which sustained the regrowth of symbionts. These preliminary findings
showed that the deep-sea mussels were able to control their endosymbionts using a set of
genes in response to environmental change. Due to the limitation of remotely operated vehicle
(ROV) dives and sampling capacity, we had to have pooled samples of each state for nuclei ex-
traction and sequencing. Thus, the cells per cluster could be considered as technical than bio-
logical replicates. Although this sampling process strategy has been broadly used in

snRNA/scRNA sequencing[96], we recognise the possible violation of assumptions in p-value cal-

culation for DEGs between the three states.

Overall, the single-cell spatial and functional atlas developed in the present work will deci-

pher some common principles of symbiosis and environmental adaption mechanisms of ani-

mals. The workflow developed in the present study could provide insightful references for re-

searchers focusing on the mechanistic study of the biological adaptation of biologically and

ecologically important non-model animals.

Acknowledgements: This study was supported by the Science and Technology Innovation Pro-

ject of Laoshan Laboratory (Project Number No. LSKJ202203104), the National Natural Science

Foundation of China (Grant No. 42030407), Southern Marine Science and Engineering Guang-

dong Laboratory (Guangzhou)(HJ202101, SMSEGL20SC01), Major Project of Basic and Applied

Basic Research of Guangdong Province (2019B030302004), the Research Grants Council of

Hong Kong (C6026-19G-A and 16101822) and the Guangdong Natural Science Funds for Distin-

guished Young Scholar (2022B1515020033). We appreciate all the assistance provided by the

crew of *R/V* 'Kexue' and the operation team of ROV 'Faxian'.

Author contributions: W.H., K.C., P.Y.Q. and L.C.L designed the study. Z.H., L. Cao, Z.Z.S., L.

Chao and W.M.X conducted the *in situ* transplant experiments and collected the samples; Z.Q.Y.

and W.H. constructed the snRNA-seq library. H.K. performed the bioinformatics analysis. W.H.,

Z.H., L. Cao, J.L., H.C. and L.Z. performed the experiments. This paper was written by W.H., H.K.,

K.C., P.Y.Q. and L.C.L. All authors revised the manuscript. All authors read, approved, and con-

tributed to the final manuscript.

Competing interests: The authors declare no competing interests.

Figure legends

Figure 1: Identification of 14 cell types in the gill of deep-sea symbiotic mussel *Gigantidas plati-frons*. A: Overall experimental scheme describing the deep-sea *in situ* transplant experiment, the sample preparation procedures and the single-cell analysis and validation pipeline. Three *G. platifrons* samples were included in the present study: 'Fanmao,' Starvation and Reconstitution. The cell nucleus was extracted from each sample, which included a pool of gill posterior tip of three mussels. The snRNA-seq libraries were constructed according to the BD Rhapsody single-nuclei 3′ protocol. Cell population-specific markers were validated by WISH and ISH. B: The image shows the posterior end of the gill of *G. platifrons*. C: UMAP representation of *G. plati-frons* gill single cells. Cell clusters are coloured and distinctively labelled. D: Heat map profile of markers in each cluster. The colour gradient represents the expression level of every single cell.

Figure 2: Supportive cell populations of *G. platifrons* gill. A: UMAP representation of the four supportive cell populations. B: Expression profiles of the cell markers that are specific or en-riched in the supportive cell populations. The sizes of the circles represent the percentages of cells in those clusters that expressed a specific gene. Genes shown in red were validated by WISH or double FISH. C and C': Schematics demonstrating the overall structural (panel C) and supportive cell distribution (C'). D, F, H, and J: WISH characterisation of the selected representa-tive cell population markers. E, G, I, and K: Double FISH characterisation of the selected repre-sentative cell population markers. The white arrowheads in G indicate the BMC1 cells locates at the otter rim of gill slice. Scale bar: 50μm.

Figure 3: Ciliary cell populations of *G. platifrons* gill. A: UMAP representation of the four ciliary cell populations and potential smooth muscle cell population. B: Expression profiles of the cell markers that are specific or enriched in the ciliary cell populations. The sizes of the circles rep-resent the percentages of cells in those clusters that expressed a specific gene. The genes shown in red were validated by WISH or double FISH. C-E, G and I-J: WISH characterisation of the selected representative cell population markers. F and H: SEM analysis of the ciliary cells of *G. platifrons* gill. K: Schematic of the water flow agitated by different ciliary cell types. The

coluor of arrowheads corresponds to water flow potentially influenced by specific types of cilia,
as indicated by their colour code in Figure 3A.

Figure 4: Proliferation cell populations of *G. platifrons* gill. A: UMAP representation of the three
proliferation cell populations. B: Expression profiles of the cell markers that are specific or en-
riched in the supportive cell populations. The sizes of the circles represent the percentages of
cells in those clusters that expressed a specific gene. Genes shown in red were validated by
WISH or double FISH. C and D: Photographic and schematic analyses of the spatial position of
the three Proliferation cell populations. E, F and J: FISH and WISH characterisation of the se-
lected population markers. The marker genes confirmed by ISH or WISH in the current study are
indicated in red. Scale bar: 50 μm.

Figure 5: Characterisation of the bacteriocytes of *G. platifrons*. A: Schematic of the overall
structure of *G. platifrons* gill filaments. B: Whole-mount FISH analyses of the overall distribution
of bacteriocytes on the *G. platifrons* gill filament. C: SEM analysis of the bacteriocytes. D: TEM
analysis of a bacteriocyte. E: UMAP representation of *G. platifrons* bacteriocytes. F: Expression
profiles of the cell markers that are specific or enriched in the bacteriocytes. The sizes of the
circles represent the percentages of cells in those clusters that expressed a specific gene. Dou-
ble FISH validated the genes shown in red. I: Schematic of the host-symbiont interaction based
on the single-cell transcriptome of *G. platifrons* bacteriocytes. The marker genes confirmed by
ISH in the current study are indicated in red. Scale bar in panel G and H: 25 μm.

Figure 6: Analysis of cell population-specific DEGs. A: UMAP representation of the impact of
deep-sea *in situ* transplant treatments on the gene expression pattern of each cell population.
The cells from different treatments were labelled with different colours. The dashed line encir-
cled bacteriocyte populations have a considerably altered expression profile. B: Histogram of
cross-state distances between the centroids of the Fanmao, Starvation and Reconstitution
groups per cell type on UMAP. The black dashed lines indicate the bacteriocyte populations
whose expression profile was remarkably altered. C: Visualization of bacteriocytes onto the

pseudo time map using monocle. The black lines indicate the main path of the pseudotime or-
dering of the cells. D: Bifurcation of selected gene expression along two branches in response to
environmental perturbation. Genes are clustered hierarchically into two groups, illustrating up-
(cluster 1) and down- (cluster2) regulated genes in the starvation state compared with Fanmao.
Genes in red colour were discussed in Section 4. The heat map showing the gene expression
profiles of all bacteriocytes' DEGs is shown in supplementary Fig. S7; E: Proposed model for the
molecular mechanisms of host-symbiont interactions in response to environmental changes.
500
501

Materials and methods:

Deep-sea *in situ* transplant experiment and sample collection: *In situ* transplant experiment
was conducted at the 'F-site' cold seep during the R/V 'Kexue' 2020 South China Sea cold seep
cruise. The overall design of the *in situ* transplant experiment and environmental states are
shown in Figure 1. First, mussels in the methane-rich 'Fanmao' site (meaning 'prosperous' site,
22° 06′ 55.425″ N,119° 17′ 08.287″ E, depth 1117m) were scoped into three nylon bags
with approximately 10 mussels in each bag. Then, two bags of mussels were transplanted to the
low methane Starvation site (Figure 1, 22° 07′ 00″ N, 119° 17′ 07.02″ E, depth 1147.42m).
After 11 days of transplantation, one bag of mussels in the Starvation site was moved back to
the 'Fanmao' site. On the 14th day of the transplant, three bags of mussels: one bag from the
Fanmao site (designated as the 'Fanmao' sample), one bag from the starvation site (designated
as the 'Starvation' sample) and one bag of mussels which were first transplanted to the Starva-
tion site for 11 days and moved back to the Fanmao site for 3 days (designated as the 'Reconsti-
tution' sample) were all retrieved by ROV Fanxian. The mussels were kept in a hydraulic pres-
sure-sealed biobox during the ascending of the ROV. The biobox is made of heat-insulated ma-
terial, which will prohibit heat exchange with warm surface water. The samples were immedi-
ately processed once onboard R/V Kexue. For snRNA-seq, the posterior end tip of the mussel's
gill was dissected (Figure 1C), snap-frozen with liquid nitrogen and then stored at –80 °C until
use. For WISH, ISH and FISH, the gills of the mussels were first fixed with 4% paraformaldehyde
(PFA, prepared with autoclaved 0.22 μM membrane filtered *in situ* seawater) at 4 °C overnight.

Then, the gill was washed with ice-cold 1× PBS three times, dehydrated and stored in 100%

methanol at −20 °C.

Single-nucleus RNA-sequencing of *G. platifrons*: The gill nucleus was extracted using the Nuclei

PURE prep nuclei isolation kit (Sigma-Aldrich). For each sample, posterior end tips from 3 indi-

vidual mussels were randomly selected and pooled together. The posterior tips were homoge-

nised in 10 mL of ice-cold lysis solution (Nuclei PURE lysis buffer with 0.1% Triton X100 and 1

mM DTT) on ice. The cell nuclei were then separated by sucrose gradient centrifugation accord-

ing to the manufacturer's protocol. The cell nuclei pellets were washed by re-suspending in ice-

cold DPBS–BSA solution (1× DPBS, 0.04% nuclease-free BSA, 0.01% RNase inhibitor; Takara).

The nucleus was spun down by centrifugation at 500×g for 5 min at 4 °C. This step was repeated

to remove the contaminants from the cell plasma. Finally, the nucleus was re-suspended in the

DPBS–BSA solution. The concentration of the cell nucleus was counted by Cell Countess II. Sin-

gle-nucleus RNA-seq libraries were then constructed with the BD Rhapsody™ single-cell analysis

system using the BD Rhapsody WTA Amplification Kit according to the manufacturer's protocol.

The library was subjected to 150bp paired-end sequencing using the Illumina HiSeq 4000 plat-

form. The clean reads of three datasets were submitted to the National Centre for Biotechnol-

ogy Information Sequence Read Archive database (Bioproject: PRJNA779258).

Bioinformatics: The three raw datasets (Fanmao, Starvation, and Reconstitution) were proc-

essed individually following the BD single-cell genomics analysis setup user guide (Doc ID:

47383 Rev. 8.0) and BD single-cell genomics bioinformatics handbook (Doc ID: 54169 Rev. 7.0).

This process involved the preparation of gene names, alignment of data to the genome, and

generation of an expression matrix for each dataset. A *G. platifrons* genome reference v1.0

(available at the Dryad Digital Repository; http://dx.doi.org/10.5061/dryad.h9942) was utilized.

The genome was indexed and constructed using STAR 2.5.2b[97]. Subsequently, the sequencing

data was mapped to the indexed genome using the BD Rhapsody single-nucleus pipeline v1.9,

employing default parameters.

Each data matrix was converted to a SingleCellExperiment data format, and empty barcodes

were removed using the emptyDrops function of DropletUtils v3.14[98]. Then, we converted and

processed the data in Seurat v3[28]. We removed cells that had <100 or > 2500 genes and <100

or > 6000 unique molecular identifiers (UMIs). We also removed genes that had <10 UMIs in

each data matrix. Then, we log-normalised the data and used DoubletFinder v2.0[99] to remove

potential doublet, assuming a 7.5% doublet formation rate. The numbers of retained nuclei

were 9717, 21614 and 28928 for Fanmao, Starvation and Reconstitution data, respectively

(Supplementary Table S20). We used the top 3000 highly variable genes for principal compo-

nent analysis (PCA) and a reciprocal PCA approach to integrating the three datasets[100]. The first

40 principal components (PCs) were used for uniform manifold approximation and projection

(UMAP) dimensional reduction and following clustering (Supplementary Fig. S11). We em-

ployed an empirical parameter - a resolution of 0.2 - in the FindClusters function, utilizing the

original Louvain algorithm. To identify unique marker genes associated with each cluster, we

utilized the FindAllMarkers function from the Seurat package. This analysis employed the Wil-

coxon rank sum test, focusing on genes that demonstrated a minimum 0.3-fold difference be-

tween two groups of cells. The annotation of cell types relied on the reference of previously

published marker genes. In instances where clusters exhibited marker genes that couldn't be

associated with known cell types, we pursued the validation process through whole-mount *in*

situ hybridization (WISH; elaborated below), discerning cell types based on gene expression

patterns and morphological characteristics. When WISH displayed uniform expression of

marker genes from different clusters within the same cell type, we consolidated those clusters

into a single cell type. We assigned cells to 14 reliable cell types (Supplementary Fig. S11). Sup-

plementary data S1 presents the counting matrix and Supplementary data S1 presents the av-

erage expression of each gene per cell type. To evaluate the stability of each cell type, we im-

plemented a bootstrap sampling and clustering strategy comprising 100 iterations using cells

combined from all three samples and individual samples[101]. The determination of marker genes

per cell type followed the previously described methodology. The identified marker genes were

subsequently utilized in KEGG enrichment analysis, employing ClusterProfiler v4.2[102].

Cell trajectory analysis: We conducted slingshot trajectory analyses using Slingshot[103] v2.2.1

for a subset of clusters (intercalary cells, dorsal-end proliferation cells,ventral-end proliferation

cells, mucus cells, basal membrane cells 1, basal membrane cells 2 and bacteriocytes) to ex-

plore the developmental trajectory of cells, assuming that all these cells were developed from

the same precursor ('posterior-end budding zone'). We performed PCA for the subsampled data

set and conducted dimensional reduction using phateR v1.0.7 for slingshot analyses. We used

the Potential of Heat-diffusion for Affinity-based Trajectory Embedding (PHATE) because it

could better reveal developmental branches than other tools[104].

We also examined the effect of deep-sea transplant experiments on shaping gene expression

patterns by comparing the expression levels amongst the three different states of a given clus-

ter. We conducted Monocle analyses using Monocle2 and Monocle 3 in R environment[105,106].

This comparison was done for the bacteriocytes. The biased number of cells per state could af-

fect the results of the dimensional reduction and calculation of marker genes, and the se-

quenced nucleus per state was unbalanced; therefore, we first downsampled the cells per clus-

ter per state to to a maximum of 1000 nuclei per cell type. Thereafter, we performed PCA and

dimensional reduction using UMAP and PHATE, calculated marker genes and conducted sling-

shot trajectory and KEGG enrichment analyses as mentioned above for each cell type.

Phylogenetic estimation: Phylogenetic estimation was conducted for E3 ubiquitin ligase

RNF213 genes. We downloaded RNF213 genes from GenBank for representative vertebrate and

invertebrate species across the tree of life. The amino acid sequences were aligned with the *G.*

platifrons sequences annotated as E3 ubiquitin ligase (Bpl_scaf_25983-1.26 and Bpl_scaf_1894-

1.45) using MAFFT v7.450. The alignment was used to estimate a maximum-likelihood best

gene tree and to calculate bootstrap values on each node using RAxML v8.2.

***G. platifrons* gill fixation and storage:** The gill tissues of *G. platifrons* were dissected within

minutes after the ROV, and samples were retrieved on board R/V Kexue. The gill tissues were

briefly washed with ice-cold filtered and autoclaved *in situ* seawater (FAISW) and then fixed in 4%

PFA prepared in FAISW at 4 °C overnight. The gill tissues were washed three times with ice-cold

PBST, dehydrated in 100% methanol and stored at -20 °C until use.

Synthesise probes for mRNA *in situ* hybridisations: For WISH and double FISH analyses, the

DNA fragments (~1000bp) of the targeted genes were first PCR amplified with gene-specific

primers (GSPs) pairs using *G. platifrons* gill cDNA as template (the sequences of targeted genes

and gene-specific primers were provided in Supplementary Data S3). The amplified fragments

were ligated into the pMD18-simpleT vector (Takara) and transformed into *E. coli*. Individual

colonies were picked up, and their plasmids were sequenced to confirm the inserts. The tem-

plates for in vitro mRNA transcription were amplified using T7 forward GSP (sense probe con-

trol) or Sp6 reversed GSPs (antisense probe) combined with either forward or reversed gene-

specific primer. Labelled probes and control probes were generated using digoxigenin (DIG)-12-

UTP (Roche) or fluorescein-12-UTP (Roche) according to the protocol described by Thisse[107]

with Sp6 and T7 RNA polymerase, respectively.

Paraffin embedding and double fluorescent *in situ* hybridisation: The methanal-dehydrated

gill slices were incubated in 100% ethanol, a 1:1 mixture of 100% ethanol and xylene and Xylene

twice for 1 h each at RT. The samples were embedded by incubating in Paraplast Plus (Sigma-

Aldrich) for 2 h at 65 °C and then cooled down to RT. Sections with 5 µM thickness were cut us-

ing a microtome (Leica).

For double FISH, sections were dewaxed by incubating in xylene twice, a 1:1 mixture of 100%

ethanol and xylene, 100% ethanol twice, 95% ethanol, 85% ethanol and 75% ethanol for 15 min

each at RT. The sections were washed with PBST three times for 10 min each and then perme-

abilised by 2 µg/mL proteinase K (NEB) in PBST for 15 min at RT. Post-digestion fixation was

conducted by incubating the sections in 4% PFA in PBST for 30 min at RT. The sections were

washed three times with PBST for 15 min each. Pre-hybridisation was conducted by incubating

the sections in HM for 1 h at 55 °C. Then, the *in situ* hybridisation was performed by incubating

the sections in ~0.5 ng/µL fluorescein (Roche)-labelled probed prepared in fresh HM overnight

at 55 °C. The sections were washed three times with 2×SSC for 15 min each at 55 °C, cooled

down to room temperature and washed three times with PBST.

A second-round hybridisation was conducted on the DIG-labelled oligonucleotides to label the

symbiont. The slices were hybridised for 1 h at 46 °C with 100 ng DIG-labelled *G. platifrons*

symbiont-specific probe in FISH buffer (0.9 M NaCl, 0.02 M Tris-HCl, 0.01% sodium dodecyl sul-

phate [SDS] and 30% formamide). The slices were then washed with FISH washing buffer (0.1 M

NaCl, 0.02 M Tris HCl, 0.01% SDS and 5 mM EDTA) three times at 5 min each at 48 °C. The slices

were washed with PBST three times and then blocked with blocking buffer (2.5% sheep serum

and 2% BSA in sheep serum) for 1 h at room temperature.

The slices were then incubated with 1:1000 diluted anti-fluorescein–peroxidase (POD) (Roche)

overnight at 4 °C, then washed six times with PBST for 15 min each and three times with TNT

buffer (100 mM Tris-HCl, pH 7.5; 100 mM NaCl; 0.1% Tween 20) for 15 min each. Afterwards,

the fluorescent signal of the *G. platifrons* gene expression pattern was developed by the TSA

fluorescein kit (Akoya Biosciences) according to the manufacturer's protocol. The slices were

washed three times, and the remaining POD activity was quenched by incubation in 1% hydro-

gen peroxide solution for 1 h at room temperature. Then, the slices were washed three times

with PBS, blocked with blocking buffer for 30 min at room temperature, incubated with 1:2500

diluted anti-DIG–POD (Roche) for 2 h at RT and washed with PBST for six times and TNT for

three times. The symbiont FISH signal was developed using the TSA Cy3 kit (Akoya Biosciences).

Finally, the slices were washed with PBST, stained with DAPI, and mounted with ProLong Dia-

mond Antifade Mountant (Thermo Fisher).

Whole-mount *in situ* hybridisation: For WISH, the connective region at the end of the W-

shaped gill filament was cut off, and each gill slice was carefully peeled off with fine-tip tweez-

ers. We dissected gill tissues from 5 individual mussels and pooled all the gill slices together.

The gill slices were then rehydrated in 75%, 50% and 25% methanol-PBST (1×PBS with 0.1%

Tween 20) for 15 min each, followed by 3×5 min PBST washes. The gill slices were then perme-

abilised with 2 μg/mL proteinase K in PBST for 30 min at 37 °C. Post-digestion fixation was con-

ducted by fixing the gill slices with 4% PFA in PBST for 30 min at room temperature (RT). After

3×5 min PBST wash to remove the residual fixative, the gill slices were pre-hybridised with a

hybridisation mix (HM; containing 50% formamide, 5×saline–sodium citrate (SSC), 0.1% Tween

20, 10 µg/mL heparin, 500 µg/mL yeast tRNA) for 1 h at 65 °C. For each hybridisation, 5-10 gill

slices were added to 400 µL of fresh HM containing ~0.5 ng/µL DIG-labelled probe. Hybridisa-

tion was conducted in a 55 °C shaking water bath overnight. Post-hybridisation washes were

performed according to the following steps: the gill slices were first washed 3×15 min with hy-

bridisation washing buffer (50% formamide, 5×SSC, 0.1% Tween 20), followed by 3×15 min

2×SSC with 0.1% Tween 20 and 3×15 min 0.2×SSC with 0.1% Tween 20. The washings were also

conducted in a shaking water batch, and all the washing buffers were pre-heated to 55 °C. The

samples were washed three times with PBST and then blocked in blocking buffer (2.5% sheep

serum, 2% BSA in PBST) for 1 h at RT. Each sample was incubated with 1:10,000 diluted anti-

DIG-AP antibody (Roche) at 4 °C overnight. The samples were incubated with the antibody, then

washed for 6×15 min with PBST, followed by washing in 3×15 min alkaline Tris buffer (100 mM

Tris-HCl, pH 9.5; 100 mM NaCl; 50 mM $MgCl_2$). The samples were incubated in nitro blue tetra-

zolium/5-Bromo-4-chloro-3-indolyl phosphate staining solution (Sangon). After the desired ex-

pression pattern was revealed, the staining reaction was stopped by 3×15 min PBST–EDTA wash

(PBST, 1 mM ETDA). The gill slices were cleared by incubating in 100% glycerol overnight at 4 °C

and then mounted on glass slides. The results of control hybridisations (with sense probes)

were provided in Supplementary Fig. S13-S16. WISH analyses were repeated with another

batch of *G. platifrons* gill slices samples collected during the *R/V* 'Kexue' 2017 South China Sea

cold seep cruise to confirm the consistency in expression patterns.

Microscopy imaging: All the WISH samples and whole-mount 16S FISH images were observed

and imaged with a Nikon Eclipse Ni microscope with a DS-Ri2 camera. The double FISH slides

were imaged with a Zeiss LSM710 confocal microscope.

Electron microscopy analysis: The gill slices of the *G. platifrons* were dissected and fixed in

electron microscopy fixative (2.5% glutaraldehyde and 2% PFA) at 4 °C. For SEM analysis, the

samples were dehydrated in a graded ethanol series and then dried at the critical point. The

samples were then coated with gold (sputter/carbon Thread, EM ACE200) and observed under

a scanning electron microscope (VEGA3, Tescan). For TEM analysis, the samples were rinsed

with double distilled water, post-fixed with 1% osmium tetroxide and then washed with double

distilled water. The samples were then rinsed, dehydrated and embedded in Ep812 resin. Ultra-

thin sections were obtained with an ultramicrotome (70 nm thickness, Reichert-Jung Ultracut E).

The sections were then double-stained with lead citrate and uranyl acetate. The cells were ob-

served under a transmission electron microscope (JEM1200, Jeol) operated under 100 kV.

References

Kremer, N. *et al.* Initial symbiont contact orchestrates host-organ-wide transcriptional changes that prime tissue colonization. *Cell Host & Microbe.* **14**, 183-194 (2013). https://doi.org:10.1016/j.chom.2013.07.006

Bang, C. *et al.* Metaorganisms in extreme environments: do microbes play a role in organismal adaptation? *Zoology (Jena)* (2018). https://doi.org:10.1016/j.zool.2018.02.004

Franke, M., Geier, B., Hammel, J. U., Dubilier, N. & Leisch, N. Coming together-symbiont acquisition and early development in deep-sea bathymodioline mussels. *Proc. R. Soc.* **288**, 20211044 (2021). https://doi.org:10.1098/rspb.2021.1044

Dubilier, N., Bergin, C. & Lott, C. Symbiotic diversity in marine animals: the art of harnessing chemosynthesis. *Nat. Rev. Microbiol.* **6**, 725 (2008). https://doi.org:10.1038/nrmicro1992

Sogin, E. M., Kleiner, M., Borowski, C., Gruber-Vodicka, H. R. & Dubilier, N. Life in the dark: phylogenetic and physiological diversity of chemosynthetic symbioses. *Annu. Rev. Microbiol.* **75**, 695-718 (2021). https://doi.org:10.1146/annurev-micro-051021-123130

Xu, T., Feng, D., Tao, J. & Qiu, J.-W. A new species of deep-sea mussel (Bivalvia: Mytilidae: *Gigantidas*) from the South China Sea: Morphology, phylogenetic position, and gill-associated microbes. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **146**, 79-90 (2019). https://doi.org:https://doi.org/10.1016/j.dsr.2019.03.001

DeChaine, E. G. & Cavanaugh, C. M. Symbioses of methanotrophs and deep-sea mussels (Mytilidae: Bathymodiolinae). *Prog. Mol. Subcell. Biol.* **41**, 227-249 (2006). https://doi.org:10.1007/3-540-28221-1_11

Fujiwara, Y. *et al.* Phylogenetic characterization of endosymbionts in three hydrothermal vent mussels influence on host distributions. *Mar. Ecol. Prog. Ser.* **208**, 147-155 (2000). https://doi:10.3354/meps208147

Kiel, S. The vent and seep biota : aspects from microbes to ecosystems. (Springer, 2010). https://doi.org/10.1007/978-90-481-9572-5

Vrijenhoek, R. C. Genetic diversity and connectivity of deep-sea hydrothermal vent metapopulations. *Mol. Ecol.* **19**, 4391-4411 (2010). https://doi.org:10.1111/j.1365-294X.2010.04789.x

Halary, S., Riou, V., Gaill, F., Boudier, T. & Duperron, S. 3D FISH for the quantification of methane- and sulphur-oxidizing endosymbionts in bacteriocytes of the hydrothermal vent mussel *Bathymodiolus azoricus*. *ISME J* **2**, 284-292 (2008). https://doi.org:10.1038/ismej.2008.3

Zheng, P. *et al.* Insights into deep-sea adaptations and host-symbiont interactions: A comparative transcriptome study on Bathymodiolus mussels and their coastal relatives. *Mol. Ecol.* **26**, 5133-5148 (2017). https://doi.org:10.1111/mec.14160

Fiala-Médioni, A., Métivier, C., Herry, A. & Le Pennec, M. *Ultrastructure of the gill of the hydrothermal-vent mytilid Bathymodiolus sp.. Mar. Biol.* **92**, 65–72 (1986). https://doi.org/10.1007/BF00392747

Wong, Y. H. *et al.* High-throughput transcriptome sequencing of the cold seep mussel Bathymodiolus platifrons. *Sci. Rep.* **5**, 16597 (2015). https://doi.org:10.1038/srep16597

15   Sun, J. *et al.* Adaptation to deep-sea chemosynthetic environments as revealed by mussel
genomes. *Nat. Ecol. Evol.* **1**, 121 (2017). https://doi.org:10.1038/s41559-017-0121
16   Bettencourt, R. *et al.* An Insightful Model to Study Innate Immunity and Stress Response
in Deep‐Sea Vent Animals: Profiling the Mussel *Bathymodiolus azoricus. Organismal and*
Molecular Malacology, **8**, (2017). https://doi.org:10.5772/68034.
17   Barros, I. *et al.* Post-capture immune gene expression studies in the deep-sea
hydrothermal vent mussel *Bathymodiolus azoricus* acclimatized to atmospheric pressure.
Fish Shellfish Immunol **42**, 159-170 (2015). https://doi.org:10.1016/j.fsi.2014.10.018
18   Wang, H. *et al.* Molecular analyses of the gill symbiosis of the bathymodiolin mussel
Gigantidas platifrons. iScience **24**, 101894 (2021).
https://doi.org:10.1016/j.isci.2020.101894
19   Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. RNA imaging. Spatially
resolved, highly multiplexed RNA profiling in single cells. *Science* **348**, 6090 (2015).
https://doi.org:10.1126/science.aaa6090
20   Hwang, B., Lee, J. H. & Bang, D. Single-cell RNA sequencing technologies and
bioinformatics pipelines. *Exp. Mol. Med.* **50**, 1-14 (2018). https://doi.org:10.1038/s12276-
018-0071-8
21   Saliba, A. E., Westermann, A. J., Gorski, S. A. & Vogel, J. Single-cell RNA-seq: advances and
future challenges. *Nucleic Acids Res* **42**, 8845-8860 (2014).
https://doi.org:10.1093/nar/gku555
22   Chen, X., Teichmann, S. A. & Meyer, K. B. From tissues to cell types and back: single-cell
gene expression analysis of tissue architecture. *Annu. Rev. Biomed. Data Sci.* **1**, 29-51
(2018). https://doi.org:10.1146/annurev-biodatasci-080917-013452
23   Wu, H., Kirita, Y., Donnelly, E. L. & Humphreys, B. D. Advantages of single-nucleus over
single-cell RNA sequencing of adult kidney: rare cell types and novel cell states revealed in
fibrosis. *J. Am. Soc. Nephrol.* **30**, 23-32 (2019). https://doi.org:10.1681/ASN.2018090912
24   Feng, D. *et al.* Using Bathymodiolus tissue stable carbon, nitrogen and sulfur isotopes to
infer biogeochemical process at a cold seep in the South China Sea. *Deep Sea Res. Part I*
Oceanogr. Res. **104**, 52-59 (2015).
https://doi.org:https://doi.org/10.1016/j.dsr.2015.06.011
25   Kharchenko, P. V., Silberstein, L. & Scadden, D. T. Bayesian approach to single-cell
differential expression analysis. *Nat. Methods.* **11**, 740-742 (2014).
https://doi.org:10.1038/nmeth.2967
26   Elyanow, R., Dumitrascu, B., Engelhardt, B. E. & Raphael, B. J. netNMF-sc: leveraging gene-
gene interactions for imputation and dimensionality reduction in single-cell expression
analysis. *Genome. Res.* **30**, 195-204 (2020). https://doi.org:10.1101/gr.251603.119
27   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
(2021). https://doi.org:10.1016/j.cell.2021.04.048
28   Stuart, T. *et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902 (2019).
https://doi.org:10.1016/j.cell.2019.05.031
29   Petibon, C., Malik Ghulam, M., Catala, M. & Abou Elela, S. Regulation of ribosomal protein
genes: An ordered anarchy. *WIREs RNA* **12**, e1632 (2021).
https://doi.org:https://doi.org/10.1002/wrna.1632

30 Welcker, D. *et al.* Hemicentin-1 is an essential extracellular matrix component of the
dermal–epidermal and myotendinous junctions. *Sci. Rep.* **11**, 17926 (2021).
https://doi.org:10.1038/s41598-021-96824-4
31 Dufour, S. C. & Beninger, P. G. A functional interpretation of cilia and mucocyte
distributions on the abfrontal surface of bivalve gills. *Mar. Biol.* **138**, 295-309 (2001).
https://doi.org:10.1007/s002270000466
32 Dufour, S. C. Gill anatomy and the evolution of symbiosis in the bivalve family Thyasiridae.
Biol. Bull. **208**, 200-212 (2005). https://doi.org:10.2307/3593152
33 Gómez-Mendikute, A., Elizondo, M., Venier, P. & Cajaraville, M. P. Characterization of
mussel gill cells in vivo and in vitro. *Cell Tissue Res.* **321**, 131-140 (2005).
https://doi.org:10.1007/s00441-005-1093-9
34 Beninger, P. G. & Dufour, S. Mucocyte distribution and relationship to particle transport
on the pseudolamellibranch gill of *Crassostrea virginica* (Bivalvia:Ostreidae). *Mar. Ecol.*
Progr. Ser. **137**, 133-138 (1996). https://doi.org:10.3354/meps137133
35 Gerdol, M. *et al.* The C1q domain containing proteins of the Mediterranean mussel
Mytilus galloprovincialis: A widespread and diverse family of immune-related molecules.
Dev. Comp. Immunol. **35**, 635-643 (2011).
https://doi.org:https://doi.org/10.1016/j.dci.2011.01.018
36 Wang, W., Song, X., Wang, L. & Song, L. Pathogen-derived carbohydrate recognition in
molluscs immune defense. *Int. J. Mol. Sci.* **19**, 721 (2018).
https://doi.org:10.3390/ijms19030721
37 Saco, A., Rey-Campos, M., Novoa, B. & Figueras, A. Transcriptomic response of mussel
gills after a *Vibrio splendidus* infection demonstrates their role in the immune response.
Front. Immunol. **11**, 615580 (2020). https://doi.org:10.3389/fimmu.2020.615580
38 Richoux, N. B. & Thompson, R. J. Regulation of particle transport within the ventral groove
of the mussel (*Mytilus edulis*) gill in response to environmental conditions. *J. Exp. Mar.*
Biol. Ecol. **260**, 199-215 (2001). https://doi.org:10.1016/s0022-0981(01)00254-4
39 Riisgård, H. U., Egede, P. P. & Barreiro Saavedra, I. Feeding behaviour of the mussel,
Mytilus edulis: New observations, with a minireview of current knowledge. *J. Mar. Biol.*
2011, 312459 (2011). https://doi.org:10.1155/2011/312459
40 Heude, É., Shaikho, S. & Ekker, M. The dlx5a/dlx6a genes play essential roles in the early
development of zebrafish median fin and pectoral structures. *PLOS ONE* **9**, e98505 (2014).
https://doi.org:10.1371/journal.pone.0098505
41 Han, S. *et al.* TULP3 is required for localization of membrane-associated proteins ARL13B
and INPP5E to primary cilia. *Biochem. Biophys. Res. Commun.* **509**, 227-234 (2019).
https://doi.org:10.1016/j.bbrc.2018.12.109
42 Louka, P. *et al.* Proteins that control the geometry of microtubules at the ends of cilia. *J.*
Cell Biol. **217**, 4298-4313 (2018). https://doi.org:10.1083/jcb.201804141
43 Das, A., Dickinson, D. J., Wood, C. C., Goldstein, B. & Slep, K. C. Crescerin uses a TOG
domain array to regulate microtubules in the primary cilium. *Mol. Biol. Cell.* **26**, 4248-
4264 (2015). https://doi.org:10.1091/mbc.E15-08-0603
44 Llorente-Cortes, V., Martinez-Gonzalez, J. & Badimon, L. LDL receptor-related protein
mediates uptake of aggregated LDL in human vascular smooth muscle cells. *Arterioscler.*
Thromb. Vasc. Biol. **20**, 1572-1579 (2000). https://doi.org:10.1161/01.atv.20.6.1572

45 Chen, X. *et al.* Angiotensin-converting enzyme in smooth muscle cells promotes
atherosclerosis-brief report. *Arterioscler. Thromb. Vasc. Biol.* **36**, 1085-1089 (2016).
https://doi.org:10.1161/ATVBAHA.115.307038
46 Linke, W. A. & Grützner, A. Pulling single molecules of titin by AFM—recent advances and
physiological implications. *Pflug. Arch. Eur.* **456**, 101-115 (2008).
https://doi.org:10.1007/s00424-007-0389-x
47 St Paul, A., Corbett, C. B., Okune, R. & Autieri, M. V. Angiotensin II, Hypercholesterolemia,
and vascular smooth muscle cells: A perfect trio for vascular pathology. *Int. J. Mol. Sci.* **21**
4525 (2020). https://doi.org:10.3390/ijms21124525
48 Ytrehus, K., Ludvigsen, S., Mancusi, C., Gerdts, E. & de Simone, G. Heart angiotensin-
converting enzyme and angiotensin-converting enzyme 2 gene expression associated with
male sex and salt-sensitive hypertension in the Dahl Rat. *Front. physiol.* **12** (2021).
https://doi.org:10.3389/fphys.2021.663819
49 Keller, T. C., 3rd *et al.* Role of titin in nonmuscle and smooth muscle cells. *Adv. Exp. Med.*
Biol. **481**, 265-277 (2000). https://doi.org:10.1007/978-1-4615-4267-4_16
50 Chen, J. V. *et al.* Rootletin organizes the ciliary rootlet to achieve neuron sensory function
in *Drosophila*. *J. Cell. Biol.* **211**, 435-453 (2015). https://doi.org:10.1083/jcb.201502032
51 Styczynska-Soczka, K. & Jarman, A. P. The *Drosophila* homologue of rootletin is required
for mechanosensory function and ciliary rootlet formation in chordotonal sensory
neurons. *Cilia* **4**, 9 (2015). https://doi.org:10.1186/s13630-015-0018-9
52 Mohan, S., Timbers, T. A., Kennedy, J., Blacque, O. E. & Leroux, M. R. Striated rootlet and
nonfilamentous forms of rootletin maintain ciliary function. *Curr. Biol.* **23**, 2016-2022
(2013). https://doi.org:10.1016/j.cub.2013.08.033
53 Jones, S. E. & Jomary, C. Secreted Frizzled-related proteins: searching for relationships
and patterns. *Bioessays* **24**, 811-820 (2002). https://doi.org:10.1002/bies.10136
54 Gillis, W. Q., Bowerman, B. A. & Schneider, S. Q. The evolution of protostome GATA
factors: molecular phylogenetics, synteny, and intron/exon structure reveal orthologous
relationships. *BMC Evol. Biol.* **8**, 112-112 (2008). https://doi.org:10.1186/1471-2148-8-
112
55 Zhang, Z., Liu, L., Twumasi-Boateng, K., Block, D. H. S. & Shapira, M. FOS-1 functions as a
transcriptional activator downstream of the *C. elegans* JNK homolog KGB-1. *Cell. Signal.*
30, 1-8 (2017). https://doi.org:https://doi.org/10.1016/j.cellsig.2016.11.010
56 Phochanukul, N. & Russell, S. No backbone but lots of Sox: Invertebrate Sox genes. *Int. J.*
Biochem. Cell Biol. **42**, 453-464 (2010).
https://doi.org:https://doi.org/10.1016/j.biocel.2009.06.013
57 Cannuel, R., Beninger, P. G., McCombie, H. & Boudry, P. Gill Development and its
functional and evolutionary implications in the blue mussel *Mytilus edulis* (Bivalvia:
Mytilidae). *Biol. Bull.* **217**, 173-188 (2009). https://doi.org:10.1086/BBLv217n2p173
58 Leibson, N. L. & Movchan, O. T. Cambial zones in gills of Bivalvia. *Mar. Biol.* **31**, 175-180
(1975). https://doi.org:10.1007/BF00391629
59 Wentrup, C., Wendeberg, A., Schimak, M., Borowski, C. & Dubilier, N. Forever competent:
deep-sea bivalves are colonized by their chemosynthetic symbionts throughout their
lifetime. *Environ. Microbiol.* **16**, 3699-3713 (2014). https://doi.org:10.1111/1462-
2920.12597

60   Mohieldin, A. M. *et al.* Proteomic identification reveals the role of ciliary extracellular-like
vesicle in cardiovascular function. *Adv. Sci.* **7**, 1903140 (2020).
https://doi.org:https://doi.org/10.1002/advs.201903140
61   Neumann, D. & Kappes, H. On the growth of bivalve gills initiated from a lobule-producing
budding zone. *Biol. Bull.* **205**, 73-82 (2003). https://doi.org:10.2307/1543447
62   Piquet, B. et al. Regionalized cell proliferation in the symbiont-bearing gill of the
hydrothermal vent mussel Bathymodiolus azoricus. Symbiosis 82, 225-233 (2020).
https://doi.org/10.1007/s13199-020-00720-w
63   Streams, M. E., Fisher, C. R. & Fiala-Médioni, A. Methanotrophic symbiont location and
fate of carbon incorporated from methane in a hydrocarbon seep mussel. *Mar. Biol.* **129**,
465-476 (1997). https://doi.org:10.1007/s002270050187
64   Ponnudurai, R. *et al.* Metabolic and physiological interdependencies in the *Bathymodiolus*
azoricus symbiosis. *ISME J* **11**, 463-477 (2017). https://doi.org:10.1038/ismej.2016.124
65   Barry, J. P. *et al.* Methane‐based symbiosis in a mussel, *Bathymodiolus platifrons*, from
cold seeps in Sagami Bay, Japan. *Invertebr. Biol.* **121**, 47-54 (2002).
https://doi.org:doi:10.1111/j.1744-7410.2002.tb00128.x
66   Renoz, F., Noel, C., Errachid, A., Foray, V. & Hance, T. Infection dynamic of symbiotic
bacteria in the pea *Aphid acyrthosiphon pisum* gut and host immune response at the early
steps in the infection process. *Plos One* **10,** 0122099 (2015).
https://doi.org:ARTNe012209910.1371/journal.pone.
67   Guan, H. *et al.* Lipid Biomarker Patterns Reflect Nutritional Strategies of Seep-Dwelling
Bivalves From the South China Sea. *Front. Mar. Sci.* **9,** 3398 (2022).
https://doi.org:10.3389/fmars.2022.831286
68   Li, Y. *et al.* Genomic adaptations to chemosymbiosis in the deep-sea seep-dwelling
tubeworm *Lamellibrachia luymesi*. *BMC Biol.* **17**, 91 (2019).
https://doi.org:10.1186/s12915-019-0713-x
69   Sun, Y. *et al.* Genomic signatures supporting the symbiosis and formation of chitinous
tube in the deep-sea tubeworm *Paraescarpia echinospica*. *Mol. Biol. Evol.* **38**, 4116-4134
(2021). https://doi.org:10.1093/molbev/msab203
70   Takishita, K. *et al.* Genomic evidence that methanotrophic endosymbionts likely provide
deep-sea bathymodiolus mussels with a sterol intermediate in cholesterol biosynthesis.
Genome Biol. Evol. **9**, 1148-1160 (2017). https://doi.org:10.1093/gbe/evx082
71   Demidenko, A., Akberdin, I. R., Allemann, M., Allen, E. E. & Kalyuzhnaya, M. G. Fatty acid
biosynthesis pathways in *Methylomicrobium buryatense* 5G(B1). *Front. Microbiol.* **7**, 2167
(2016). https://doi.org:10.3389/fmicb.2016.02167
72   Brasaemle, D. L. *et al.* Adipose differentiation-related protein is an ubiquitously expressed
lipid storage droplet-associated protein. *J. Lipid Res.* **38**, 2249-2263 (1997).
https://doi.org/10.1016/S0022-2275(20)34939-7
73   Cheng, D. *et al.* Expression, purification, and characterization of human and rat acetyl
coenzyme A carboxylase (ACC) isozymes. *Protein Expr. Purif.* **51**, 11-21 (2007).
https://doi.org:10.1016/j.pep.2006.06.005
74   Monroig, Ó. & Kabeya, N. Desaturases and elongases involved in polyunsaturated fatty
acid biosynthesis in aquatic invertebrates: a comprehensive review. *Fish. Sci.* **84**, 911-928
(2018). https://doi.org:10.1007/s12562-018-1254-x

75   Kabeya, N. et al. Unique fatty acid desaturase capacities uncovered in Hediste diversicolor
illustrate the roles of aquatic invertebrates in trophic upgrading. *Philos. Trans. R. Soc.* **375**,
20190654 (2020). https://doi.org:10.1098/rstb.2019.0654

76   Soupene, E. & Kuypers, F. A. Mammalian long-chain acyl-CoA synthetases. *Exp. Biol. Med.*
(Maywood). **233**, 507-521 (2008). https://doi.org:10.3181/0710-MR-287

77   Alves-Bezerra, M. *et al.* Long-chain acyl-CoA synthetase 2 knockdown leads to decreased
fatty acid oxidation in fat body and reduced reproductive capacity in the insect *Rhodnius*
prolixus. Biochim. Biophys. Acta. **1861**, 650-662 (2016).
https://doi.org:10.1016/j.bbalip.2016.04.007

78   Hoglund, P. J., Nordstrom, K. J. V., Schioth, H. B. & Fredriksson, R. The solute carrier
families have a remarkably long evolutionary history with the majority of the human
families present before divergence of bilaterian species. *Mol. Biol. Evol.* **28**, 1531-1541
(2011). https://doi.org:10.1093/molbev/msq350

79   Mohamed, A. R. *et al.* Dual RNA-sequencing analyses of a coral and its native symbiont
during the establishment of symbiosis. *Mol. Ecol.* **29**, 3921-3937 (2020).
https://doi.org:10.1111/mec.15612

80   Bertucci, A., Foret, S., Ball, E. E. & Miller, D. J. Transcriptomic differences between day and
night in *Acropora millepora* provide new insights into metabolite exchange and light-
enhanced calcification in corals. *Mol. Ecol.* **24**, 4489-4504 (2015).
https://doi.org:10.1111/mec.13328

81   Hamada, M. *et al.* Metabolic co-dependence drives the evolutionarily ancient Hydra-
Chlorella symbiosis. *Elife* **7,** e35122 (2018). https://doi.org:10.7554/eLife.35122

82   Duncan, R. P., Feng, H., Nguyen, D. M. & Wilson, A. C. Gene family expansions in aphids
maintained by endosymbiotic and nonsymbiotic traits. *Genome Biol. Evol.* **8**, 753-764
(2016). https://doi.org:10.1093/gbe/evw020

83   Feng, H. L. *et al.* Trading amino acids at the aphid-Buchnera symbiotic interface. *P. Natl.*
Acad. Sci. USA. **116**, 16003-16011 (2019). https://doi.org:10.1073/pnas.1906223116

84   Ip, J. C. *et al.* Host-endosymbiont genome integration in a deep-sea chemosymbiotic clam.
Mol. Biol. Evol. **38**, 502-518 (2021). https://doi.org:10.1093/molbev/msaa241

85   Hongo, Y. *et al.* Expression of genes involved in the uptake of inorganic carbon in the gill
of a deep-sea vesicomyid clam harboring intracellular thioautotrophic bacteria. *Gene* **585**,
228-240 (2016). https://doi.org:https://doi.org/10.1016/j.gene.2016.03.033

86   Sotiriou, S. *et al.* Ascorbic-acid transporter Slc23a1 is essential for vitamin C transport into
the brain and for perinatal survival. *Nat. Med.* **8**, 514-517 (2002).
https://doi.org:10.1038/0502-514

87   Aydemir, T. B. et al. Zinc transporter ZIP14 functions in hepatic zinc, iron and glucose
homeostasis during the innate immune response (endotoxemia). *PLoS One* **7**, e48679
(2012). https://doi.org:10.1371/journal.pone.0048679

88   Turi, Z., Lacey, M., Mistrik, M. & Moudry, P. Impaired ribosome biogenesis: mechanisms
and relevance to cancer and aging. Aging (Albany NY) **11**, 2512-2540 (2019).
https://doi.org:10.18632/aging.101922

89   Hay, B. A. Understanding IAP function and regulation: a view from Drosophila. Cell. Death.
Differ. **7**, 1045-1056 (2000). https://doi.org:10.1038/sj.cdd.4400765

90   Dubrez-Daloz, L., Dupoux, A. & Cartier, J. IAPS : More than just inhibitors of apoptosis
proteins. Cell Cycle **7**, 1036-1046 (2008). https://doi.org:10.4161/cc.7.8.5783
91   Pflug, K. M. & Sitcheran, R. Targeting NF-κB-inducing kinase (NIK) in immunity,
inflammation, and cancer. *Int. J. Mol. Sci.* **21**, 8470 (2020).
92   Otten, E. G. *et al.* Ubiquitylation of lipopolysaccharide by RNF213 during bacterial
infection. *Nature* **594**, 111-116 (2021). https://doi.org:10.1038/s41586-021-03566-4
93   Price, D. R. *et al.* Aphid amino acid transporter regulates glutamine supply to intracellular
bacterial symbionts. *Proc. Natl. Acad. Sci. USA.* **111**, 320-325 (2014).
https://doi.org:10.1073/pnas.1306068111
94   Ponnudurai, R. *et al.* Comparative proteomics of related symbiotic mussel species reveals
high variability of host–symbiont interactions. *ISME J.* **14**, 649-656 (2020).
https://doi.org:10.1038/s41396-019-0517-6
95   Tashian, R. E. The carbonic anhydrases: widening perspectives on their evolution,
expression and function. *Bioessays* **10**, 186-192 (1989).
https://doi.org:10.1002/bies.950100603
96   Hicks, S. C., Townes, F. W., Teng, M. & Irizarry, R. A. Missing data and technical variability
in single-cell RNA-sequencing experiments. *Biostatistics* **19**, 562-578 (2017).
https://doi.org:10.1093/biostatistics/kxx053
97   Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15-21 (2013).
https://doi.org:10.1093/bioinformatics/bts635
98   Lun, A. T. L. *et al.* EmptyDrops: distinguishing cells from empty droplets in droplet-based
single-cell RNA sequencing data. *Genome Biol.* **20**, 63 (2019).
https://doi.org:10.1186/s13059-019-1662-y
99   McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet detection in single-
cell RNA sequencing data using artificial nearest neighbors. *Cell Systems* **8**, 329-337
(2019). https://doi.org:https://doi.org/10.1016/j.cels.2019.03.003
100   Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e1821
(2019). https://doi.org:https://doi.org/10.1016/j.cell.2019.05.031
101   Singh, P. & Zhai, Y. Deciphering Hematopoiesis at single cell level through the lens of
reduced dimensions. *bioRxiv*, 2022.2006.2007.495099 (2022).
https://doi.org:10.1101/2022.06.07.495099
102   Wu, T. et al. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
The Innovation **2**, 100141 (2021).
https://doi.org:https://doi.org/10.1016/j.xinn.2021.100141
103   Street, K. et al. Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genom.* **19**, 477 (2018). https://doi.org:10.1186/s12864-018-4772-0
104   Moon, K. R. et al. Visualizing structure and transitions in high-dimensional biological data.
Nat. Biotechnol. **37**, 1482-1492 (2019). https://doi.org:10.1038/s41587-019-0336-3
105   Qiu, X. et al. Reversed graph embedding resolves complex single-cell trajectories. Nat
Methods 14, 979-982 (2017). https://doi.org/10.1038/nmeth.4402
106   Cao, J. et al. The single-cell transcriptional landscape of mammalian organogenesis.
Nature 566, 496-502 (2019). https://doi.org/10.1038/s41586-019-0969-x
107   Thisse, C. & Thisse, B. High-resolution in situ hybridization to whole-mount zebrafish
embryos. Nat. Protoc. 3, 59-69 (2008). https://doi.org:10.1038/nprot.2007.514

B

DAPI | Symbiont | Merge

Ciliary disk

Ciliary disk | Food grove | VEPCs | B'

C

Bacteriocyte

IC | IC | IC | IC

IC | Bacteriocyte

10μm

D

Bacteriocyte | IC | IC

Symbiont

N | N | N

lysosomes

Basal membrane

Interlaminar lumen

5μm

E

UMAP_1 | UMAP_2

Bacteriocytes

F

		Average Expression
Ambiguous		
Bacteriocytes		
Apical ridge ciliary cell		
Food grove ciliary cell		
Lateral cilia cell		
Smooth muscle cell		
Intercalary cell		
Inter lamina cell		
Basal membrane cell 1		
Basal membrane cell 2		
Mucus cell		
Posterior end budding zone cell		
Dorsal end proliferation cell		
Ventral end proliferation cell		

Percent Expressed: 25 50 75

Membrane associated
Lysosome associated
Cellular metabolism
Host-symbiont interaction
Pro and Anti- Apoptosis

Chitin synthase 1(6901-7.5)
NMT3(15282-3.10)
KIF-16B(54265-1.13)
KLC-like(4784-1.40)
CTSL1-like(61711-5.12)
CTSL1-like(46838-6.40)
GAA(59648-4.5)
CLCN7(7842-1.11)
Rabenosyn-5(54816-0.3)
LYST(52809-1.6)
DMBT1(52188-1.15)
TMPRSS15(154410-0.9)
SLC7A9(36159-5.5)
FYCO1-like(33726-5.7)
PLIN2-like(27158-3.8)
ACC1(55250-5.11)
PLCB4-like(35916-0.6)
ACSBG2-like(28862-1.5)
SLC23A1-like(32311-1.19)
SLC24A3-like(14503-1.28)
SLC12A3-like(44604-3.7)
SLC40A1-like(63447-0.12)
SLCA39A14(44428-5.3)
CHDH(34315-4.10)
RBM25(42979-0.15)
HIP1(41172-3.3)
BIRC3(6172-0.3)
BIRC7(32300-1.15)

G CTSL1-like(46838-6.40)
Symbiont DAPI

Frontal

Abfrontal

H PLCB4-like(35916-0.6)
Symbiont DAPI

Frontal

Abfrontal

I

Membrane
Chitin synthase 1
NMT-3

Vesicle transport
Kinesin
kinesin light chain-like
FYVE and coiled-coil
domain-containing 1-like

Rabenosyn-5

lysosomal-trafficking
regulator-like

H(+)Cl(-) exchange
transporter 7-like

Symbiont

Metabolite exchange
b(0,+)-type amino
acid transporter 1-like

Globin
Carbonic anhydrase

Cellular metabolism
Perilipin 2-like
Acetyl-carboxylase 2
Fatty acid desaturase 1-like
long-chain-fatty-acid ligase
ACSBG2-like

Lysosome
Cathepsin L1-like
Lysosomal alpha-glucosidase-like
TMPRSS15

Transporters
Vitamin C
Na⁺ Zn²⁺ → SLC23A1
→ SLC24A3
Fe
Mg²⁺ → SLC12A3
Cl⁻
Ca²⁺ K⁺ → SLC40A1
→ SLC39A14
Glycose and sugars
CO² O²

Anti-apoptosis
BIRC3

Pro-apoptosis
RNA-binding 25-like
huntingtin-interacting 1-like