Deciphering deep-sea chemosynthetic symbiosis by single-nucleus RNA-sequencing

Hao Wang[1,2,3#], Kai He[4#], Huan Zhang[1#], Quanyong Zhang[5#], Lei Cao[1], Jing Li[1,6], Zhaoshan Zhong[1], Hao Chen[1], Li Zhou[1], Chao Lian[1], Minxiao Wang[1], Kai Chen[5], Pei-Yuan Qian[2,3*], and Chaolun Li[1,6,7*]

[1]Center of Deep-Sea Research, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China

[2]Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou), Nansha, Guangzhou, China

[3]Department of Ocean Science, Hong Kong University of Science and Technology, Hong Kong S.A.R, China.

[4]Key Laboratory of Conservation and Application in Biodiversity of South China, School of Life Sciences, Guangzhou University, Guangzhou, Guangdong 510006, China

[5]State Key Laboratory of Primate Biomedical Research, Institute of Primate Translational Medicine, Kunming University of Science and Technology, Kunming, China

[6]South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, China

[7]University of Chinese Academy of Sciences, Beijing, China

Correspondence authors: Prof. Pei-yuan Qian and Prof. Chaolun Li

Email addresses: boqianpy@ust.hk and lcl@qdio.ac.cn

#These authors contributed equally to this work

The authors declare no conflict of interest.

Abstract:

Bathymodioline mussels dominate deep-sea methane seep and hydrothermal vent habitats and

obtain nutrients and energy primarily through chemosynthetic endosymbiotic bacteria in the

bacteriocytes of their gill. However, the molecular mechanisms that orchestrate mussel host-

symbiont interactions remain unclear. Here, we constructed a comprehensive cell atlas of the

gill in the mussel *Gigantidas platifrons* from the South China Sea methane seeps (1100m depth)

using single-nucleus RNA sequencing (snRNA-seq) and whole-mount *in situ* hybridisation. We

identified 13 types of cells, including three previously unknown ones, uncovered unknown

tissue heterogeneity. Every cell type has a designated function in supporting the gill's structure

and function, creating an optimal environment for chemosynthesis, and effectively acquiring

nutrients from the endosymbiotic bacteria. Analysis of snRNA-seq of *in situ* transplanted

mussels clearly showed the shifts in cell state in response to environmental oscillations. Our

findings provide insight into principles of host-symbiont interaction and the bivalves'

environmental adaption mechanisms.

Introduction:

Mutualistic interactions between multicellular animals and their microbiota play a fundamental role in animals' adaptation, ecology, and evolution [1,2]. By associating with symbionts, host animals benefit from the metabolic capabilities of their symbionts and gain fitness advantages that allow them to thrive in habitats they could not live in on their own[1,3]. Prime examples of such symbioses are bathymodioline mussels and gammaproteobacteria endosymbionts[4]. The bathymodioline mussels occur worldwide at deep-sea chemosynthetic ecosystems, such as cold-seeps, hydrothermal vents, and whale falls[4]. The host mussel acquired sulfur-oxidising (SOX) and/or methane-oxidising (MOX) symbiont through horizontal transmission at their early life stage[3]. Their symbionts, which are hosted in a specialised gill epithelial cell, namely, the bacteriocytes[5,6], utilise the chemical energy from the reduced chemical compounds such as CH_4 and H_2S released from cold seeps or hydrothermal vents to fix carbon and turn into carbon source for the host mussel[7,8]. The ecological success of the bathymodioline symbioses is apparent: the bathymodiolin mussels are often among the most dominant species in the deep-sea chemosynthetic ecosystems[9,10]. Thus, it is critical to know how the bathymodioline mussels interact with the symbionts and maintain the stability and efficiency of the symbiosis.

The gill of bathymodiolin mussels has adapted to deep-sea chemosynthetic life at the levels of molecule, cell and tissue[11,12]. Compared to shallow water mussels[13], the gill filaments of bathymodiolin mussels have enlarged surfaces allowing them to hold more symbionts per filament (Supplementary Fig. S1). This adaption requires not only novel cellular and molecular mechanisms to maintain and support the enlarged gill filament structure but also a strong ciliary ventilation system to pump vent/seep fluid to fuel the symbiotic bacteria. Previous studies based on whole-genome sequencing and bulk RNA-seq projects have shown that genes in the categories of nutrient transporters, lysosomal proteins, and immune receptors are either expanded in the host mussel's genome or up-regulated in the gill[12, 14-17], suggesting that these genes involved in host-symbiont interaction. Though providing deep molecular insights, these studies mainly used homogenised tissues that average genes' expression levels amongst different cell types and eliminate cell and gene spatial distribution information[18-21]. In addition,

the broad expression and function of potentially 'symbiosis-related' proteins also greatly

limited data interpretation. Therefore, a systemic atlas of gill cell types and the descriptions of

cell-type-specific gene expressional profiles are warranted to a better understanding of the

host-symbiont interaction and environmental adaptation mechanisms of the bathymodioline

symbiosis.

In recent years, single cell/nucleus RNA-sequencing (sc/sn RNA-seq) technologies have

become one of the preferred methods for investigating the composition of complex tissue at

the transcriptional level[22]. snRNA-seq has several advantages, such as compatibility with frozen

samples, elimination of dissociation-induced transcriptional stress responses, and reduced

dissociation bias[23]. These advantages are significant for deep-sea and other ecological studies

because cell and molecular biology facilities are not available in the field. To examine the

cellular and molecular mechanisms of the environmental adaptations and host-symbiont

interactions in bathymodioline mussels, we conducted a single nucleus RNA-sequencing-based

transcriptomic study. We analysed the gill symbiosis using deep-sea mussels *Gigantidas*

platifrons[24]. Our work provides a proof-of-principle for environmental adaptation mechanisms

study in non-traditional but ecologically important organisms with single-nucleus RNA-

sequencing technologies.

Results and discussion

1. *G. platifrons* deep-sea *in situ* transplant experiment and single-cell transcriptomic analysis:

We conducted a *G. platifrons* deep-sea *in situ* transplant experiment at the "F-Site" Cold-seep (~1117 m depth) and retrieved three groups of samples (Fig. 1A), as follows: the 'Fanmao' (meaning prosperous) group, which comprised the mussels collected from methane-rich (~40 µM) site of the cold-seep, where the animals thrived; the 'starvation' group, which comprised the mussels collected from the methane-rich site and then moved to a methane-limited (~0.054 µM) starvation site for 14 days before retrieval; and the 'reconstitution' group, which comprised the methane-rich site mussels moved to the 'Starvation' site for 11 days and then moved back to the methane-rich site for another 3 days.

We next performed single nucleus RNA-sequencing (snRNA-seq) of the gill posterior tip of the three groups of samples using the microwell-based BD Rhapsody platform™ Single-Cell Analysis System (Fig. 1A). After quality control, we obtained 9717 (Fanmao), 21614 (Starvation), and 28928 (Reconstitution) high-quality single nuclei transcriptomic data.

To minimise the dropout effect in snRNAseq, we utilised a reciprocal PCA (RPCA) strategy to project cells in the three samples onto the same space based on conserved expressed genes among them[25]. This strategy maximises the number of cells per cluster regardless of the organism's state and therefore maximises genes per cluster[26]. We revealed 14 cell clusters and their corresponding marker genes (Figs. 1B- D) with Seurat[27,28]. Because there are few available canonical marker genes per cell type for *G. platifrons* and molluscs, we carefully characterised each cell cluster by 1) examining the cluster marker genes' functions, 2) identifying the expression pattern of cluster marker gene using whole-mount *in situ* hybridisation (WISH) and/or double-fluorescent *in situ* hybridisation (FISH) analyses, 3) conducting scanning electron microscopy (SEM) and transmission electron microscopy (TEM) analyses, and 4) evaluating the stability per cluster using a bootstrapping algorithm. We then successfully identified 13 of the 14 cell clusters, which could be generally divided into four groups, including 1) the supportive cells, 2) the ciliary cells, 3) the proliferation cells, and 4) the bacteriocytes. One cell cluster remained ambiguous. Notably, a bootstrapping strategy suggested that all clusters were highly supported when using all three samples (Supplementary Fig. S2A). When using each sample

individually, most clusters were also well supported except of the three ciliary cells

(Supplementary Fig. S2 B-D). This could be due to that these cells were derived from the same

precursor cells, and at least partially because they were represented by fewer numbers of

nuclei and thus fewer available genes, especially in the food groove ciliary cell (Supplementary

Table S1). The spatial and functional annotation of the cell clusters revealed that the cell

clusters worked in close cooperation to support the symbiont and maintain a high-efficiency

chemosynthetic system.

2. Cell atlas of *G. platifrons gill*

Supportive cells: We have identified four supportive cell types, namely, inter lamina cells

basal membrane cells (BMC)1, BMC2, and mucus cells (Figs 2A and 2B).

The inter lamina cells, marked by high expression of fibrillar-forming collagens, were the

cells located between the two layers of basal membranes. These cells were previously

identified as amoeboid cells[13], and their function has not been explored. The inter lamina cells

were densely distributed around the food groove and at the rim of the gill filament. While on

the middle part of the gill filament, the inter lamina cells distributed in parallel rows. It might

help to connect the two sheets of basal membranes and maintain the spatial integrity of the gill

filament. In addition, we observed the enrichment of ribosomal proteins involved in protein

synthesis (Supplementary Table S2), indicating a high metabolic rate[29] of inter lamina cells. This

implies that the inter lamina cells may process the nutrients acquired from the symbiont.

We have also identified two types of BMC populations which have not been recognised

before. The BMC1 expressed genes encoding extracellular matrix and adhesive proteins (Fig. 2B

and Supplementary Table S2), such as SCO-spondin (Bpl_scaf_27638-4.18), Zonadhesin

(Bpl_scaf_1371-2.4), Cartilage matrix proteins (Bpl_scaf_16371-0.14 and Bpl_scaf_3387-5.47),

and Collagen alpha-1(XXI) chain protein (Bpl_scaf_51350-0.19). BMC2 (Fig. 2B and

Supplementary Table S2) was distinctive from BMC1 in the context of expression (Fig 2A) and

marker genes (Fig 2B). Genes encoding hemicentins (Bpl_scaf_38731-0.50 and Bpl_scaf_7293-

0.14), which are known to stabilise the cells' contact with the basal membrane[30], were highly

expressed in BMC2. Thus, these two cell types are likely to help building the basal lamina, and

stabilising the epithelial-derived cells, such as bacteriocytes and intercalary cells, on the surface

of the basal membrane.

Mucus cells are specialised secretory cells with intracellular mucus vacuoles[13]. In shallow-

water filter-feeding bivalves, mucus cells secret mucus, which cooperates with the ciliary

ventilation system to capture, process, and transport food particles to their mouths, known as

filter feeding mechanism[31-34]. Deep-sea mussels may not necessarily retain this function since

they do not normally acquire food resources such as phytoplankton and planktonic bacteria but

obtain most of nutrient through their chemosynthetic symbiont. Thus, it was hypothesised that

mucus cells are involved in other biological functions such as immune responses to pathogens[18].

Herein, genes encoding proteins with microbe-binding functionalities were enriched in mucus

cells (Fig. 2B and Supplementary Table S2), such as C-type lectins (Bpl_scaf_58694-0.22 and

Bpl_scaf_38133-0.8), C1q domain-containing protein (Bpl_scaf_55216-2.11), and Fibrinogen C

domain-containing protein (Bpl_scaf_18519-2.15)[35,36]. The expression of similar immune genes

was upregulated in a shallow water mussel *Mytilus galloprovincialis* when challenged by a

pathogenic bacteria[37]. Our WISH and double-FISH analyses showed that mucus cells were

embedded within the outer rim cilia and scattered on the gill lamella alongside the

bacteriocytes (Figs. 2G and 2K). These data collectively suggest that mucus cells may help

mussel maintain the immune homeostasis of gill. Interestingly, WISH and double-FISH analyses

showed that mucus cells were also distributed alongside the gill lamella's food groove and the

inner edge, where the density of bacteriocytes is low (Fig. 2G). Because the food groove is the

main entry of food practical to the labial palps and mouth[38], this distribution pattern implies

that mucus cells may be also involved in capturing planktonic bacteria and sending them to the

mouth.

Ciliary and smooth muscle cells: A remarkable feature of bathymodioline mussel's gill is its

ciliary ventilation system, which constantly agitates the water and provides the symbiont with

the necessary gas[39]. We identified four types of ciliary cells (Figs. 3A and 3B): apical ridge ciliary

cells (ARCCs), food groove ciliary cells (FGCCs), lateral ciliary cells (LCCs), and intercalary cells

(ICs), as well as a newly identified type of smooth muscle cells. All ciliary cells were marked by

canonical cilium genes, such as genes encoding flagellar proteins, ciliary motor kinesin proteins,

ciliary dynein proteins, and ciliary microtubules and were clearly distinguishable by specifically

expressed genes (Supplementary Table S3 and Supplementary Fig. S3). ARCCs were

characterized by expression of Bpl_scaf_20631-1.16, which encodes homeobox Dlx6a-like

protein (Fig 3D and 3F) which is a marker of the apical ectodermal ridge[40]. FGCCs expressed

genes encoding primary cilia development regulator Tubby-related protein 3[41]

(Bpl_scaf_24834-2.3) and primary ciliary cell structural protein TOG array regulator of axonemal

microtubules protein 1[42,43] (Bpl_scaf_55620-1.3), suggesting the FGCCs are the sensory ciliary

cells that gather information from the surrounding environment. Both ARCCs and FGCCs were

located around the ventral tip of the gill filament. LCCs were distributed as two parallel rows

along the gill lamella's outer rim and ciliary disks' outer rim (Figs. 3G and 3H) and had highly

expressed genes involved in cilium structure, cilium movement and ATP synthases (Fig. 3B and

Supplementary Table S3), indicating that LCCs may have a strong ability to beat their cilia.

Interestingly, we identified a group of previously unreported smooth muscle cells (SMCs) co-

localised with the LCCs as showed by WISH (Fig. 3I and Supplementary Fig. S4). SMCs strongly

expressed several Low-density lipoprotein receptors (LDL receptor) and LDLR-associated

proteins[44] (Fig. 3B). In addition, these cells also expressed the angiotensin-converting enzyme-

like protein[45] (Bpl_scaf-29477-5.10) and the 'molecular spring' titin-like protein[46]

(Bpl_scaf_56354-7.8) (Supplementary Table S3). The expression of these genes could be

commonly found in human vascular smooth muscle cells[47-49]. Collectively, we suspect that SMCs

are involved in lateral cilium movement and the gill slice contraction.

ICs are the specialised ciliary cells surrounding the bacteriocytes (Figs. 3B and 3J). WISH

analysis showed that the expressions of IC markers had apparent spatial variations (Fig. 3J). The

expression of ICs marker rootletin[50-52] (Bpl_scarf_22362-6.9) was considerably higher in the ICs

close to the frontal edge of the gill filament, and the expression gradually decreased along with

the direction of inter lamina water flow (Fig. 3K and Supplementary Fig. S5), implying that ICs

ventilate the water flow and the mucus through the gill filaments. Furthermore, compared with

the other three types of ciliary cells, the ICs expressed several genes encoding transcription

factors involved in determining cell fate (Fig. 3B and Supplementary Table S3), such as

transcription factors Sox 8 (Bpl_scarf_40595-5.30) and Wnt pathway cell polarity regulator

secreted frizzled-related protein 5 (Bpl_scaf_57424-6.1)[53], suggesting that the ICs might also

play regulatory roles[54-56].

Proliferation cells: It has long been known that the bathymodioline mussel gill has three

types of proliferation cells that are conserved throughout all filibranchia bivalves: the budding

zone at the posterior end of the gill where new gill filaments are continuously formed, and the

dorsal and ventral ends of each gill filaments[57,58]. These "cambial-zone"-like cell populations

could continuously proliferate throughout the whole life span of the mussel[58]. Our snRNA-seq

data recognised three types of proliferation cells, which is consistent with previous findings (Fig.

4A, Fig. 4B and Supplementary Table S4). The gill posterior end budding zone cells (PEBZCs) are

located on the first few freshly proliferated filaments of the posterior tip of gill (Figs. 4C and 4D).

The PBEZCs marker gradually disappeared at around the 11th -12th role of the gill filament,

suggesting the maturation of the gill filaments, which is similar to the developmental pattern

reported in another deep-sea mussel *Bathymodiolus azoricus* [59]. The dorsal end proliferation

cells (DEPCs), which expressed the hallmarks genes of muscular tissue, as well as cell

proliferation and differentiation regulators, were the proliferation cells in connective tissue at

the dorsal end of the gill slice (Fig. 4A, 4E). The ventral end proliferation cells (VEPCs) were two

symmetrical triangle-shaped cell clusters of small symbiont-free cells (Fig. 4F). In VEPCs, genes

encoding ribosomal proteins, chromatin proteins, RNA and DNA binding proteins, and cell

proliferation markers were all up-regulated (Fig. 4B and Supplementary Table S4), indicating

that VEPCs are meristem-like cells[59,60].

It has been hypothesised that new proliferation cells will be colonised by symbionts, serving

as the vital mechanism of bacteriocyte recruitment[59]. We then determined which proliferation

cell type gives rise to the mature cells, especially the bacteriocytes, which has little available

information regarding their precursor and transmission mode in bathymodioline mussels [59,61].

Slingshot trajectory analysis suggested that the PEBZCs might be the origin of all gill epithelial

cells, including the other two proliferation cells (VEPC and DEPC) and bacteriocytes

(Supplementary Fig. S6). The sole exception was BMC2, which may be derived from VEPC

rather than PEBZC. This result is consistent with previous studies which suggested new gill

filaments of the filibranch mussels are formed in the gill's posterior budding zones[59]. The

colonisation by the symbiont might play a crucial role in determining the fate of the

bacteriocytes. Noticeably, in *G. platifrons*, only the pillar-shaped first row of gill filament,

comprised of small meristem-like cells, was symbiont-free (Figs. 4D and D'), whereas all the

other gill filaments were colonised by symbionts. This pattern of symbiosis establishment is

different from that of *B. azoricus*, in which PEBZCs are symbiont free and are gradually

colonised by symbionts released from the bacteriocytes on the adjacent mature gill filaments

after maturation[59]. On mature gill filaments, the DEPCs and VEPCs are seemingly the sources of

new cells that sustain the growth of the gill filament from both dorsal and ventral directions,

respectively[57].

3. Bacteriocytes and host-symbiont interaction:

We conducted the whole-mount FISH using a bacterial 16S rRNA probe for symbiont to

determine the spatial distribution of bacteriocytes, and their positions relative to the other cell

types on the gill filament (Figs. 5A, B). Bacteriocytes covered the majority of the surface of the

gill filament, except the ventral tip, the ciliary disk, and the frontal edge (lateral ciliary). The

bacteriocytes were surrounded by intercalary cells with microvilli and cilia on the surface (Fig.

5C). Noticeably, the bacteriocytes' size and FISH signal intensity, which reflected the symbiont

number in each bacteriocyte, gradually decreased from the heavily ciliated frontal edge to the

inner edge, demonstrating that the symbiont abundance in each bacteriocyte was affected by

the environmental gradient[11].

The gene expression profile of bacteriocytes aligned well with the morphological analysis,

which suggested that the bacteriocytes have structural, metabolic, and regulatory adaptions to

cope with the symbiont (Figs. 5D, 5F and 5I). It has been hypothesised that the bacteriocytes

extract nutrients from the symbiont through two pathways: lysing the symbiont through the

lysosome (the 'farming' pathway) or directly utilising the nutrient produced by the symbiont

(the 'milking' pathway)[62,63]. Our TEM observations clearly detected intracellular vacuole and

lysosome system of the bacteriocyte that harbour, transport, and digest the symbionts (Fig. 5D),

which is consistent with previous studies[18,64]. The snRNA data showed bacteriocytes expressed

cellular membrane synthesis enzyme (phosphoethanolamine N-methyltransferase,

Bpl_scaf_15282-3.10) and a series of lysosomal proteins such as lysosomal proteases

(cathepsins; Bpl_scaf_61711-5.12, Bpl_scaf_46838-6.40 and Bpl_scaf_59648-4.5; lysosomal

alpha-glucosidase-like), protease regulators 56 (TMPRSS15 proteins; Bpl_scaf_52188-1.15 and

Bpl_scaf_15410-0.9), and lysosomal traffic regulator proteins (rabenosyn-5, Bpl_scaf_54816-0.3

and Bpl_scaf_52809-1.6, lysosomal-trafficking regulator-like isoform X1). Among the proteases,

cathepsins were thought to be evolutionary conserved molecular tools that host utilized to

control the residence of their symbiont microbes[65]. They were also highly expressed in

symbiotic tissue of other deep-sea chemosynthetic animals, such as vesicomyid clams and

vestimentiferan tubeworms[66-68]. Bacteriocytes also expressed genes encoding cellular vesicle

transports (kinesins, Bpl_scaf_14819-0.11, Bpl_scaf_54265-1.13, and Bpl_scaf_4784-1.40)[53],

potential amino acid transporter[57] (Bpl_scaf_36159-5.5), and genes involved in intracellular

vesicle transport, such as the FYVE and coiled-coil domain-containing protein 1[58]

(Bpl_scaf_33726-5.7). In addition to form and mobilise early endosomes, the protein products

of these genes could transport symbiont-secreted nutrient vesicles to the host (Fig. 5F and

Supplementary Table S5), supporting the "milking" pathway.

The symbiont of *G. platifrons* belongs to type I methanotrophy, of which the core metabolic

function is linked with the development of intracytoplasmic membranes leading to a high

lipid/biomass content[69,70]. Recent lipid biomarker analyses showed that the gill of *G. platifrons*

contains a high amount of bacterial lipids, which are directly utilised by the host to synthesise

most of its lipid contents[66]. Downstream of the bacteriocyte's metabolic cascade, genes

encoding proteins that may be involved in fatty acid/lipid metabolism, such as perilipin2

(Bpl_scaf_27158-3.8), which is the critical protein to form intracellular lipid droplets[71], and a

variety of fatty acid metabolism enzymes (acetyl carboxylase 2, Bpl_scaf_55250-5.11[72]; fatty

acid desaturase 1-like isoform X1, Bpl_scaf_35916-0.6[73,74]; long-chain fatty acid-ligase ACSBG2-

like isoform X1, Bpl_scaf_28862-1.5 [75,76]), were up-regulated, suggesting that the fatty acid

could be a major form of nutrients passing from the symbiont to the host mussel.

Additionally, bacteriocytes expressed several solute carriers, including sodium/ascorbate

cotransporter (solute carrier family 23 members 1-like, Bpl_scaf_32311-1.19),

sodium/potassium/calcium exchanger (sodium potassium calcium exchanger 3-like,

Bpl_scaf_14503-1.28), sodium/chloride ion cotransporter (solute carrier family 12 member 3-

like isoform X1, Bpl_scaf_44604-3.7), ferrous iron transporter (solute carrier family 40 member

1-like, Bpl_scaf_63447-0.12) and zinc transporter (zinc transporter ZIP14-like, Bpl_scaf_44428-

5.3). The solute carriers are a large family of ATP-dependent transporters that shuttle a variety

of small molecules across the cellular membrane[77]. In several model symbiotic systems, solute

carriers play a vital role in host-symbiont interaction by either providing the symbiont with

substrates[78,79] or transporting symbiont-produced nutrients to the host [80-82]. Previous studies

demonstrated that solute carrier genes were expanded in deep-sea chemosynthetic animals'

genome[83] and highly expressed in symbiotic organs[84] including bathymodioline mussel's gill[12].

In bathymodioline mussels, previous bulk RNA-seq studies detected up-regulated expression of

a large variety of solute carriers in the gill, which is consistent with the present study,

suggesting that solute carriers may play crucial roles in shuttling nutrients in and out of

bacteriocytes and in maintaining the suitable intracellular micro-environment for the

symbiont[85,86].

4. Cell-type-specific response to environmental stresses:

To examine cell-type-specific acclimatisation to environmental changes the expression

profiles of differentially expressed genes (DEGs) were compared between the three states

(Fanmao, starved and reconstituted animals) of samples collected in our *in situ* transplantation

experiment (full lists of DEGs are shown in Supplementary Table S6-S18). Notably, for each

state three mussels were processed but pooled for nuclei extraction before snRNA sequencing

(see Method for detail). We tested our hypothesis here that at the starvation site, a relatively

low concentration of methane shall upset symbiont metabolism and thus substantially affect

symbiont-hosting bacteriocytes by assessing the transcriptional changes per cell type. We

calculated the distances between the three sets of samples on the UMAP plot (Fig. 6A)

estimating the centroid coordinates per sample per cell type, and calculating the Euclid distance

between the centroid coordinates of each pair of states per cell type. The impact of starvation

was variable across cell types, as reflected by the cross-state distances (Fig. 6B, green bar).

Starvation resulted in the most significant transcriptional changes in bacteriocytes reflected by

a large Fanmao-vs-starvation distance, followed by VEPC and inter lamina cells. On the other

hand, starvation had little impact on the transcriptions of ciliary cells and most supportive cells

such as the food groove ciliary cell, BMC2, and mucus cells. Fig 6B also shows that after

reconstitution (moving the mussels back to the methane-rich site Fanmao for 3 days), the

expressional profile of bacteriocytes rapidly changed back, reflected by a large starvation-vs-

reconstitution distance and much smaller Fanmao-vs-reconstitution distance. This result

coincided with and was supported by our pseudo-time analysis for bacteriocytes (Fig. 6C),

showing that bacteriocytes in the reconstitution are in intermediate and transitional states

between Fanmao and starvation.

For the bacteriocytes population, we conducted cell trajectory pseudo-timing and detailed

DEG analysis to interpret the mechanism of host-symbiont interactions. The branched heatmap

showed distinct gene expression patterns amongst the three treatments (Fig. 6D and

supplementary Fig. S7). The Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway

enrichment analysis of the bacteriocytes' DEGs provides an overall view of the pathways

enriched in each environmental state (Supplementary Fig. S8 A-C). The genes encoding

ribosomal proteins are highly expressed in the bacteriocytes under the methane-rich 'Fanmao'

state (Supplementary Fig. S8A), suggesting an active protein synthesis and cellular metabolism

[87]. Moreover, organic ion transporters (A BCB P-glycol, Bpl_scaf_18613-16.10; canalicular

multispecific organic anion transporter 2, Bpl_scaf_52110-4.43; sodium- and chloride-

dependent glycine transporter 1-like, Bpl_scaf_13642-5.8), which may be involved in

transporting symbiont-produced nutrients, are also highly expressed.

In starved *G. platifrons*, negative regulators of cell proliferation, such as autocrine

proliferation repressors and receptor-type tyrosine phosphatase beta, were up-regulated in

bacteriocytes, suggesting repression of cell growth. We also observed the enrichment of genes

in the apoptosis pathway (Supplementary Fig. S8B), which is attributed to the upregulation of

caspases (Bpl_scaf_25165-3.8 and Bpl_scaf_24225-0.12) and cathepsins (Bpl_scaf_64706-0.16,

Bpl_scaf_61711-5.12) which can trigger caspases-dependent cell death, suggesting cellular

stress condition. Correspondingly, a gene encoding baculoviral IAP repeat-containing protein

(Bpl_scaf_6172-0.3), which can bind caspase and inhibit apoptosis, was up-regulated in

bacteriocytes (Fig. 6E). In *Drosophila*, the baculoviral IAP proteins are important in animal's

response to cellular stress and promote cell survival[88,89]. Similarly, the gene encoding MAP3K14

(Bpl_scaf_60908-4.14) and potential E3 ubiquitin ligase RNF213 (Bpl_scaf_25983-1.26,

Bpl_scaf_1894-1.45, Supplementary Fig. S9), which were apoptosis suppressor or regulator,

were up-regulated[90]. These results suggest that starved bacteriocytes were carrying out cell-

type-specific adjustments to cope with stresses. The E3 ubiquitin ligases could also work as

intracellular immune sensors of bacterial lipopolysaccharides[91]. Thus, the encoded protein may

be able to active downstream immunological toolkit to digest the symbiont population for

nutrients.

As mentioned above, bacteriocytes obtain nutrients from endosymbionts. KEGG pathway

analyses suggested that phagosome, lysosome-related pathways were upregulated in the

starvation state, indicating that bacteriocytes were actively digesting the endosymbionts in the

starved *G. platifrons*. Interestingly, protein synthesis activity was more activated in inter lamina

cells and VEPCs as supported by the up-regulation of genes encoding ribosomal proteins in both

cell populations (Supplementary Tables S6 and S17). This was contrary to the situation in

bacteriocytes and may be a consequence of the high activity of "farming" in bacteriocytes.

We anticipated that after moving the starved *G. platifrons* back to Fanmao site,

bacteriocytes and their endosymbionts would "reconstitute" leading to the partly restored

"farming" and "milking" pathways. This was confirmed by higher expression of fatty acid

metabolic genes in the mussels in the reconstitution state than those in the starvation state,

such as long-chain-fatty-acid-ligase ACSBG2-like (Bpl_scaf_28862-1.5), elongation of very long-

chain fatty acids 7-like (Bpl_scaf_5959-1.9), and fatty acid desaturase 1-like (Bpl_scaf_35916-

0.6). These findings were also consistent with the result of KEGG analyses. The mitochondrial

trifunctional enzyme (Bpl_scaf_28862-1.5) was also highly expressed, suggesting a high level of

energy-producing activity. The KEGG analyses showed that glutamatergic synthase was

enriched in the reconstitution state in comparison with those in both the Fanmao and

starvation states. In the insect aphid–Buchnera endosymbiosis model, the host-produced

glutamate could be transported and directly utilised by the symbiont[92,93]. Similar glutamate-

based host–symbiont metabolic interaction mechanisms were also proposed in the deep-sea

mussel *Bathymodiolus thermophilus* and *B. azoricus*[63,94]. As the pseudotime analyses showed

above (Fig. 6C), reconstitution was the intermediate state between Fanmao and starvation.

Metabolic and gene regulatory functions of bacteriocytes were still different from that in

Fanmao. The KEGG analyses suggested regulatory (Rap1 signalling, retrograde endocannabinoid

signalling, chemokine signalling, glucagon signalling, thyroid hormone synthesis, mRNA

surveillance pathway, etc.) and metabolic pathways were enriched in the reconstitution state

(fatty acid metabolism, salivary secretion, aldosterone synthesis and section, insulin secretion

and pancreatic section) (Supplementary Fig. S8C). For example, we detected up-regulation of

the genes encoding carbonic anhydrases (Bpl_scaf_33596-7.3 and Bpl_scaf_48274-0.3),

electroneutral sodium bicarbonate exchanger 1 (Bpl_scaf_61230-5.11), and globin-like proteins

(Bpl_scaf_50392-1.14 and Bpl_scaf_24370-0.5), which could provide the symbiont with carbon

dioxide and oxygen necessary for the symbiont's chemosynthetic metabolism[83-94].

5. Summary and outlook:

Using the deep-sea mussel *G. platifrons* as a model organism, we demonstrated the power

of integrating snRNA-seq and WISH data in unravelling the mechanism behind animal-microbe

symbiosis. The robustness of this strategy showed stable and highly distinguishable expression

patterns of each cell type regardless of the different environmental states. We successfully

profiled the specific roles of different types of cells, including the previously unknown cell types,

in maintaining the structure and function of the gill. The supportive cells that located in

between (inter lamina cells) and on the basal membrane's surface (BMC1 and BMC2) helped

maintain the anatomical structure of the basal membrane. Ciliary and smooth muscle cells

involved in gill slice contraction and cilium beat allowing the gathering of material and

information from the surrounding environment. Proliferation cells (PBEZCs, DEPCs and VEPCs)

gave rise to new cells, including bacteriocytes which obtained nutrients from endosymbionts

using intracellular vacuoles and lysosomes.

The snRNA analyses also revealed that different cell types collaborated to support

bacteriocytes' functionality. The PBEZCs gave rise to new bacteriocytes that allowed symbiont

colonisation. Bacteriocytes attached to the basal membrane and were stabilised by supportive

cells. Mucus cells co-localise with bacteriocytes and help to maintain immune homeostasis. The

beating ciliary cells controlled the water flow, providing bacteriocytes with necessary inorganic

substances from the environment. This new information on cell-cell interaction certainly

advanced our overall understanding of how endosymbiotic microbes and host cells

communicate and collaborate, which cannot be easily achieved through other methods.

Moreover, the analysis of snRNA data from bacteriocytes has provided insight into the

molecular mechanisms employed by the host to maitain and regulate the symbiosis. Notably,

the bacteriocytes-enriched transcripts involved in harbouring, digesting symbionts, and

transporting nutrients produced by the symbionts were identify and characterised. Our *in situ*

transplant experiments by moving mussels between methane-rich and methane-limited sites

also provided clues of cell-type specific responses to environmental change. Under a methane-

limited environment, the staved mussels more actively consumed endosymbionts through the

"farming" pathway. After being moved back to the methane-rich environment, mussels

produced more glutamates which sustained the regrowth of symbionts. These preliminary

findings showed that the deep-sea mussels were able to control their endosymbionts using a

set of genes in response to environmental change. Due to the limitation of remotely operated

vehicle (ROV) dives and sampling capacity, we had to have pooled samples of each state for

nuclei extraction and sequencing. Thus, the cells per cluster could be considered as technical

than biological replicates. Although this sampling process strategy has been broadly used in

snRNA/scRNA sequencing[95], we recognise the possible violation of assumptions in p-value

calculation for DEGs between the three states.

Overall, the single-cell spatial and functional atlas developed in the present work will

decipher some common principles of symbiosis and environmental adaption mechanisms of

animals. The workflow developed in the present study could provide insightful references for

researchers focusing on the mechanistic study of the biological adaptation of biologically and

ecologically important non-model animals.

Acknowledgements: This study was supported by the Science and Technology Innovation

Project of Laoshan Laboratory (Project Number No. LSKJ202203104), the National Natural

Science Foundation of China (Grant No. 42030407), Southern Marine Science and Engineering

Guangdong Laboratory (Guangzhou)(HJ202101, SMSEGL20SC01), Major Project of Basic and

Applied Basic Research of Guangdong Province (2019B030302004), the Research Grants Council

of Hong Kong (C6026-19G-A and 16101822) and the Guangdong Natural Science Funds for

Distinguished Young Scholar (2022B1515020033). We appreciate all the assistance provided by

the crew of *R/V* 'Kexue' and the operation team of ROV 'Faxian'.

Author contributions: W.H., K.C., P.Y.Q. and L.C.L designed the study. Z.H., L. Cao, Z.Z.S., L.

Chao and W.M.X conducted the *in situ* transplant experiments and collected the samples; Z.Q.Y.

and W.H. constructed the snRNA-seq library. H.K. performed the bioinformatics analysis. W.H.,

Z.H., L. Cao, J.L., H.C. and L.Z. performed the experiments. This paper was written by W.H., H.K.,

K.C., P.Y.Q. and L.C.L. All authors revised the manuscript. All authors read, approved, and

contributed to the final manuscript.

Competing interests: The authors declare no competing interests.

Figure legends

Figure 1: Identification of 14 cell types in the gill of deep-sea symbiotic mussel *Gigantidas platifrons*. A: Overall experimental scheme describing the deep-sea *in situ* transplant experiment, the sample preparation procedures and the single-cell analysis and validation pipeline. Three *G. platifrons* samples were included in the present study: 'Fanmao,' Starvation and Reconstitution. The cell nucleus was extracted from each sample, which included a pool of gill posterior tip of three mussels. The snRNA-seq libraries were constructed according to the BD Rhapsody single-nuclei 3′ protocol. Cell population-specific markers were validated by WISH and ISH. B: The image shows the posterior end of the gill of *G. platifrons*. C: UMAP representation of *G. platifrons* gill single cells. Cell clusters are coloured and distinctively labelled. D: Heat map profile of markers in each cluster. The colour gradient represents the expression level of every single cell.

Figure 2: Supportive cell populations of *G. platifrons* gill. A: UMAP representation of the four supportive cell populations. B: Expression profiles of the cell markers that are specific or enriched in the supportive cell populations. The sizes of the circles represent the percentages of cells in those clusters that expressed a specific gene. Genes shown in red were validated by WISH or double FISH. C and C': Schematics demonstrating the overall structural (panel C) and supportive cell distribution (C'). D-F: WISH characterisation of the selected representative cell population markers. H-K: Double FISH characterisation of the selected representative cell population markers. The marker genes confirmed by WISH and ISH in the current study are indicated in red. Scale bar: 50µm.

Figure 3: Ciliary cell populations of *G. platifrons* gill. A: UMAP representation of the four ciliary cell populations and potential smooth muscle cell population. B: Expression profiles of the cell markers that are specific or enriched in the ciliary cell populations. The sizes of the circles represent the percentages of cells in those clusters that expressed a specific gene. The genes shown in red were validated by WISH or double FISH. C-E, G and I-J: WISH characterisation of the selected representative cell population markers. F and H: SEM analysis of the ciliary cells of

G. platifrons gill. K: Schematic of the water flow agitated by different ciliary cells. The marker

genes confirmed by WISH in the current study are indicated in red.

Figure 4: Proliferation cell populations of *G. platifrons* gill. A: UMAP representation of the three

proliferation cell populations. B: Expression profiles of the cell markers that are specific or

enriched in the supportive cell populations. The sizes of the circles represent the percentages of

cells in those clusters that expressed a specific gene. Genes shown in red were validated by

WISH or double FISH. C and D: Photographic and schematic analyses of the spatial position of

the three Proliferation cell populations. E, F and J: FISH and WISH characterisation of the

selected population markers. The marker genes confirmed by ISH or WISH in the current study

are indicated in red. Scale bar: 50 μm.

Figure 5: Characterisation of the bacteriocytes of *G. platifrons*. A: Schematic of the overall

structure of *G. platifrons* gill filaments. B: Whole-mount FISH analyses of the overall distribution

of bacteriocytes on the *G. platifrons* gill filament. C: SEM analysis of the bacteriocytes. D: TEM

analysis of a bacteriocyte. E: UMAP representation of *G. platifrons* bacteriocytes. F: Expression

profiles of the cell markers that are specific or enriched in the bacteriocytes. The sizes of the

circles represent the percentages of cells in those clusters that expressed a specific gene.

Double FISH validated the genes shown in red. I: Schematic of the host-symbiont interaction

based on the single-cell transcriptome of *G. platifrons* bacteriocytes. The marker genes

confirmed by ISH in the current study are indicated in red.

Figure 6: Analysis of cell population-specific DEGs. A: UMAP representation of the impact of

deep-sea *in situ* transplant treatments on the gene expression pattern of each cell population.

The cells from different treatments were labelled with different colours. The dashed line

encircled bacteriocyte populations have a considerably altered expression profile. B: Histogram

of cross-state distances between the centroids of the Fanmao, Starvation and Reconstitution

groups per cell type on UMAP. The black dashed lines indicate the bacteriocyte populations

whose expression profile was remarkably altered. C: Visualization of bacteriocytes onto the

pseudo time map using monocle. The black lines indicate the main path of the pseudotime

ordering of the cells. D: Bifurcation of selected gene expression along two branches in response

to environmental perturbation. Genes are clustered hierarchically into two groups, illustrating

up- (cluster 1) and down- (cluster2) regulated genes in the starvation state compared with

Fanmao. Genes in red colour were discussed in Section 4. The heat map showing the gene

expression profiles of all bacteriocytes' DEGs is shown in supplementary Fig. S7; E: Proposed

model for the molecular mechanisms of host-symbiont interactions in response to

environmental changes.

Materials and methods:

Deep-sea *in situ* transplant experiment and sample collection: *In situ* transplant experiment

was conducted at the 'F-site' cold seep during the R/V 'Kexue' 2020 South China Sea cold seep

cruise. The overall design of the *in situ* transplant experiment and environmental states are

shown in Figure 1. First, mussels in the methane-rich 'Fanmao' site (meaning 'prosperous' site,

22° 06′ 55.425″ N,119° 17′ 08.287″ E, depth 1117m) were scoped into three nylon bags

with approximately 10 mussels in each bag. Then, two bags of mussels were transplanted to the

low methane Starvation site (Figure 1, 22° 07′ 00″ N, 119° 17′ 07.02″ E, depth 1147.42m).

After 11 days of transplantation, one bag of mussels in the Starvation site was moved back to

the 'Fanmao' site. On the 14th day of the transplant, three bags of mussels: one bag from the

Fanmao site (designated as the 'Fanmao' sample), one bag from the starvation site (designated

as the 'Starvation' sample) and one bag of mussels which were first transplanted to the

Starvation site for 11 days and moved back to the Fanmao site for 3 days (designated as the

'Reconstitution' sample) were all retrieved by ROV Fanxian. The mussels were kept in a

hydraulic pressure-sealed biobox during the ascending of the ROV. The biobox is made of heat-

insulated material, which will prohibit heat exchange with warm surface water. The samples

were immediately processed once onboard R/V Kexue. For snRNA-seq, the posterior end tip of

the mussel's gill was dissected (Figure 1C), snap-frozen with liquid nitrogen and then stored at

−80 °C until use. For WISH, ISH and FISH, the gills of the mussels were first fixed with 4%

paraformaldehyde (PFA, prepared with autoclaved 0.22 μM membrane filtered *in situ* seawater)

at 4 °C overnight. Then, the gill was washed with ice-cold 1× PBS three times, dehydrated and

stored in 100% methanol at −20 °C.

Single-nucleus RNA-sequencing of *G. platifrons*: The gill nucleus was extracted using the Nuclei

PURE prep nuclei isolation kit (Sigma-Aldrich). For each sample, posterior end tips from 3

individual mussels were randomly selected and pooled together. The posterior tips were

homogenised in 10 mL of ice-cold lysis solution (Nuclei PURE lysis buffer with 0.1% Triton X100

and 1 mM DTT) on ice. The cell nuclei were then separated by sucrose gradient centrifugation

according to the manufacturer's protocol. The cell nuclei pellets were washed by re-suspending

in ice-cold DPBS–BSA solution (1× DPBS, 0.04% nuclease-free BSA, 0.01% RNase inhibitor;

Takara). The nucleus was spun down by centrifugation at 500×g for 5 min at 4 °C. This step was

repeated to remove the contaminants from the cell plasma. Finally, the nucleus was re-

suspended in the DPBS–BSA solution. The concentration of the cell nucleus was counted by Cell

Countess II. Single-nucleus RNA-seq libraries were then constructed with the BD Rhapsody™

single-cell analysis system using the BD Rhapsody WTA Amplification Kit according to the

manufacturer's protocol. The libraries were then sequenced with the Illumina HiSeq 4000

platform. The clean reads of three datasets were submitted to the National Centre for

Biotechnology Information Sequence Read Archive database (Bioproject: PRJNA779258).

Bioinformatics: Three raw datasets (Fanmao, Starvation and Reconstitution) were processed

individually by following the BD single-cell genomics analysis setup user guide (Doc ID: 47383

Rev. 8.0) and BD single-cell genomics bioinformatics handbook (Doc ID: 54169 Rev. 7.0). In brief,

a *G. platifrons* genome reference v1.0 (the genome could be downloaded at Dryad Digital

Repository http://dx.doi.org/10.5061/dryad.h9942)[15] was indexed and built using STAR 2.5.2b.

Gene names were prepared according to the BD-single cell guideline. The sequencing data were

then mapped to the indexed genome using the BD Rhapsody single-nucleus pipeline v1.9 with

default parameters to generate an expression matrix per dataset.

Each data matrix was converted to a SingleCellExperiment data format, and empty barcodes

were removed using the emptyDrops function of DropletUtils v3.14[96]. Then, we converted and

processed the data in Seurat v3[28]. We removed cells that had <100 or > 2500 genes and <100

or > 6000 unique molecular identifiers (UMIs). We also removed genes that had <10 UMIs in

each data matrix. Then, we log-normalised the data and used DoubletFinder v2.0[97] to remove

potential doublet, assuming a 7.5% doublet formation rate. The numbers of retained nuclei

were 9717, 21614 and 28928 for Fanmao, Starvation and Reconstitution data, respectively

(Supplementary Table S19). We used the top 3000 highly variable genes for principal

component analysis (PCA) and a reciprocal PCA approach to integrating the three datasets[98].

The first 40 principal components (PCs) were used for uniform manifold approximation and

projection (UMAP) dimensional reduction and following clustering (Supplementary Fig. S10).

We used an empirical parameter, a resolution of 0.2, in the FindClusters function. The average

expression of each gene per cluster were also provided as supplementary data S1. Based on the

WISH results (see below), we manually combined the clusters with similar gene expressions and

WISH patterns. Eventually, we assigned all the cells to 14 reliable clusters (Supplementary Fig.

S11). We adopted a bootstrap sampling strategy to evaluate the stability for each cluster[99]. We

ran bootstrapping with 100 iterations using cells in the three samples together and in each

sample individually. We determined marker genes per cluster using the FindAllMarkers function

based on the Wilcoxon rank sum test. The marker genes were used for KEGG enrichment using

ClusterProfiler v4.2[100].

Cell trajectory analysis: We conducted slingshot trajectory analyses using Slingshot[101] v2.2.1

for a subset of clusters ('intercalary cells,' 'dorsal-end proliferation cells,' 'ventral-end

proliferation cells,' 'mucus cells,' 'basal membrane cells 1,' 'basal membrane cells 2' and

'bacteriocytes') to explore the developmental trajectory of cells, assuming that all these cells

were developed from the same precursor ('posterior-end budding zone'). We performed PCA

for the subsampled data set and conducted dimensional reduction using phateR v1.0.7 for

slingshot analyses. We used the Potential of Heat-diffusion for Affinity-based Trajectory

Embedding (PHATE) because it could better reveal developmental branches than other tools[102].

We also examined the effect of deep-sea transplant experiments on shaping gene expression

patterns by comparing the expression levels amongst the three different states of a given

cluster. This comparison was done for the bacteriocytes. The biased number of cells per state

could affect the results of the dimensional reduction and calculation of marker genes, and the

sequenced nucleus per state was unbalanced; therefore, we first downsampled the cells per

cluster per state to about the same number. Thereafter, we performed PCA and dimensional

reduction using UMAP and PHATE, calculated marker genes and conducted slingshot trajectory

and KEGG enrichment analyses as mentioned above for each cell type.

Phylogenetic estimation: Phylogenetic estimation was conducted for E3 ubiquitin ligase

RNF213 genes. We downloaded RNF213 genes from GenBank for representative vertebrate and

invertebrate species across the tree of life. The amino acid sequences were aligned with the *G.*

platifrons sequences annotated as E3 ubiquitin ligase (Bpl_scaf_25983-1.26 and Bpl_scaf_1894-

1.45) using MAFFT v7.450. The alignment was used to estimate a maximum-likelihood best

gene tree and to calculate bootstrap values on each node using RAxML v8.2.

***G. platifrons* gill fixation and storage:** The gill tissues of *G. platifrons* were dissected within

minutes after the ROV, and samples were retrieved on board R/V Kexue. The gill tissues were

briefly washed with ice-cold filtered and autoclaved *in situ* seawater (FAISW) and then fixed in 4%

PFA prepared in FAISW at 4 °C overnight. The gill tissues were washed three times with ice-cold

PBST, dehydrated in 100% methanol and stored at -20 °C until use.

Whole-mount *in situ* hybridisation: For WISH and double FISH analyses, the DNA fragments

(~1000bp) of the targeted genes were first PCR amplified with gene-specific primers (GSPs)

pairs using *G. platifrons* gill cDNA as template (the sequences of targeted genes and gene-

specific primers were provided in Supplementary Data S2). The amplified fragments were

ligated into the pMD18-simpleT vector (Takara) and transformed into *E. coli*. Individual colonies

were picked up, and their plasmids were sequenced to confirm the inserts. The templates for in

vitro mRNA transcription were amplified using T7 forward GSP (sense probe control) or Sp6

reversed GSPs (antisense probe) combined with either forward or reversed gene-specific primer.

Labelled probes and control probes were generated using digoxigenin (DIG)-12-UTP (Roche) or

fluorescein-12-UTP (Roche) according to the protocol described by Thisse[103] with Sp6 and T7

RNA polymerase, respectively.

For WISH, the connective region at the end of the W-shaped gill filament was cut off, and each

gill slice was carefully peeled off with fine-tip tweezers. We dissected gill tissues from 5

individual mussels and pooled all the gill slices together. The gill slices were then rehydrated in

75%, 50% and 25% methanol-PBST (1×PBS with 0.1% Tween 20) for 15 min each, followed by

3×5 min PBST washes. The gill slices were then permeabilised with 2 µg/mL proteinase K in

PBST for 30 min at 37 °C. Post-digestion fixation was conducted by fixing the gill slices with 4%

PFA in PBST for 30 min at room temperature (RT). After 3×5 min PBST wash to remove the

residual fixative, the gill slices were pre-hybridised with a hybridisation mix (HM; containing 50%

formamide, 5×saline–sodium citrate (SSC), 0.1% Tween 20, 10 µg/mL heparin, 500 µg/mL yeast

tRNA) for 1 h at 65 °C. For each hybridisation, 5-10 gill slices were added to 400 µL of fresh HM

containing ~0.5 ng/µL DIG-labelled probe. Hybridisation was conducted in a 55 °C shaking water

bath overnight. Post-hybridisation washes were performed according to the following steps:

the gill slices were first washed 3×15 min with hybridisation washing buffer (50% formamide,

5×SSC, 0.1% Tween 20), followed by 3×15 min 2×SSC with 0.1% Tween 20 and 3×15 min 0.2×SSC

with 0.1% Tween 20. The washings were also conducted in a shaking water batch, and all the

washing buffers were pre-heated to 55 °C. The samples were washed three times with PBST

and then blocked in blocking buffer (2.5% sheep serum, 2% BSA in PBST) for 1 h at RT. Each

sample was incubated with 1:10,000 diluted anti-DIG-AP antibody (Roche) at 4 °C overnight.

The samples were incubated with the antibody, then washed for 6×15 min with PBST, followed

by washing in 3×15 min alkaline Tris buffer (100 mM Tris-HCl, pH 9.5; 100 mM NaCl; 50 mM

MgCl$_2$). The samples were incubated in nitro blue tetrazolium/5-Bromo-4-chloro-3-indolyl

phosphate staining solution (Sangon). After the desired expression pattern was revealed, the

staining reaction was stopped by 3×15 min PBST–EDTA wash (PBST, 1 mM ETDA). The gill slices

were cleared by incubating in 100% glycerol overnight at 4 °C and then mounted on glass slides.

The results of control hybridisations (with sense probes) were provided in Supplementary Fig.

S12-S15. WISH analyses were repeated with another batch of *G. platifrons* gill slices samples

collected during the *R/V* 'Kexue' 2017 South China Sea cold seep cruise to confirm the

consistency in expression patterns.

Paraffin embedding and double fluorescent *in situ* hybridisation: The methanal-dehydrated

gill slices were incubated in 100% ethanol, a 1:1 mixture of 100% ethanol and xylene and Xylene

twice for 1 h each at RT. The samples were embedded by incubating in Paraplast Plus (Sigma-

Aldrich) for 2 h at 65 °C and then cooled down to RT. Sections with 5 μM thickness were cut

using a microtome (Leica).

For double FISH, sections were dewaxed by incubating in xylene twice, a 1:1 mixture of 100%

ethanol and xylene, 100% ethanol twice, 95% ethanol, 85% ethanol and 75% ethanol for 15 min

each at RT. The sections were washed with PBST three times for 10 min each and then

permeabilised by 2 μg/mL proteinase K (NEB) in PBST for 15 min at RT. Post-digestion fixation

was conducted by incubating the sections in 4% PFA in PBST for 30 min at RT. The sections were

washed three times with PBST for 15 min each. Pre-hybridisation was conducted by incubating

the sections in HM for 1 h at 55 °C. Then, the *in situ* hybridisation was performed by incubating

the sections in ~0.5 ng/μL fluorescein (Roche)-labelled probed prepared in fresh HM overnight

at 55 °C. The sections were washed three times with 2×SSC for 15 min each at 55 °C, cooled

down to room temperature and washed three times with PBST.

A second-round hybridisation was conducted on the DIG-labelled oligonucleotides to label the

symbiont. The slices were hybridised for 1 h at 46 °C with 100 ng DIG-labelled *G. platifrons*

symbiont-specific probe in FISH buffer (0.9 M NaCl, 0.02 M Tris-HCl, 0.01% sodium dodecyl

sulphate [SDS] and 30% formamide). The slices were then washed with FISH washing buffer (0.1

M NaCl, 0.02 M Tris HCl, 0.01% SDS and 5 mM EDTA) three times at 5 min each at 48 °C. The

slices were washed with PBST three times and then blocked with blocking buffer (2.5% sheep

serum and 2% BSA in sheep serum) for 1 h at room temperature.

The slices were then incubated with 1:1000 diluted anti-fluorescein–peroxidase (POD) (Roche)

overnight at 4 °C, then washed six times with PBST for 15 min each and three times with TNT

buffer (100 mM Tris-HCl, pH 7.5; 100 mM NaCl; 0.1% Tween 20) for 15 min each. Afterwards,

the fluorescent signal of the *G. platifrons* gene expression pattern was developed by the TSA

fluorescein kit (Akoya Biosciences) according to the manufacturer's protocol. The slices were

washed three times, and the remaining POD activity was quenched by incubation in 1%

hydrogen peroxide solution for 1 h at room temperature. Then, the slices were washed three

times with PBS, blocked with blocking buffer for 30 min at room temperature, incubated with

1:2500 diluted anti-DIG–POD (Roche) for 2 h at RT and washed with PBST for six times and TNT

for three times. The symbiont FISH signal was developed using the TSA Cy3 kit (Akoya

Biosciences). Finally, the slices were washed with PBST, stained with DAPI, and mounted with

ProLong Diamond Antifade Mountant (Thermo Fisher).

Microscopy imaging: All the WISH samples and whole-mount 16S FISH images were observed

and imaged with a Nikon Eclipse Ni microscope with a DS-Ri2 camera. The double FISH slides

were imaged with a Zeiss LSM710 confocal microscope.

Electron microscopy analysis: The gill slices of the *G. platifrons* were dissected and fixed in

electron microscopy fixative (2.5% glutaraldehyde and 2% PFA) at 4 °C. For SEM analysis, the

samples were dehydrated in a graded ethanol series and then dried at the critical point. The

samples were then coated with gold (sputter/carbon Thread, EM ACE200) and observed under

a scanning electron microscope (VEGA3, Tescan). For TEM analysis, the samples were rinsed

with double distilled water, post-fixed with 1% osmium tetroxide and then washed with double

distilled water. The samples were then rinsed, dehydrated and embedded in Ep812 resin. Ultra-

thin sections were obtained with an ultramicrotome (70 nm thickness, Reichert-Jung Ultracut E).

The sections were then double-stained with lead citrate and uranyl acetate. The cells were

observed under a transmission electron microscope (JEM1200, Jeol) operated under 100 kV.

References

680		**References**
681	1	Kremer, N. *et al.* Initial symbiont contact orchestrates host-organ-wide transcriptional
682		changes that prime tissue colonization. *Cell Host & Microbe.* **14**, 183-194 (2013).
683		https://doi.org:10.1016/j.chom.2013.07.006
684	2	Bang, C. *et al.* Metaorganisms in extreme environments: do microbes play a role in
685		organismal adaptation? *Zoology (Jena)* (2018). https://doi.org:10.1016/j.zool.2018.02.004
686	3	Franke, M., Geier, B., Hammel, J. U., Dubilier, N. & Leisch, N. Coming together-symbiont
687		acquisition and early development in deep-sea bathymodioline mussels. *Proc. R. Soc.* **288**,
688		20211044 (2021). https://doi.org:10.1098/rspb.2021.1044
689	4	Dubilier, N., Bergin, C. & Lott, C. Symbiotic diversity in marine animals: the art of
690		harnessing chemosynthesis. *Nat. Rev. Microbiol.* **6**, 725 (2008).
691		https://doi.org:10.1038/nrmicro1992
692	5	Sogin, E. M., Kleiner, M., Borowski, C., Gruber-Vodicka, H. R. & Dubilier, N. Life in the dark:
693		phylogenetic and physiological diversity of chemosynthetic symbioses. *Annu. Rev.*
694		*Microbiol.* **75**, 695-718 (2021). https://doi.org:10.1146/annurev-micro-051021-123130
695	6	Xu, T., Feng, D., Tao, J. & Qiu, J.-W. A new species of deep-sea mussel (Bivalvia: Mytilidae:
696		*Gigantidas*) from the South China Sea: Morphology, phylogenetic position, and gill-
697		associated microbes. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **146**, 79-90 (2019).
698		https://doi.org:https://doi.org/10.1016/j.dsr.2019.03.001
699	7	DeChaine, E. G. & Cavanaugh, C. M. Symbioses of methanotrophs and deep-sea mussels
700		(Mytilidae: Bathymodiolinae). *Prog. Mol. Subcell. Biol.* **41**, 227-249 (2006).
701		https://doi.org:10.1007/3-540-28221-1_11
702	8	Fujiwara, Y. *et al.* Phylogenetic characterization of endosymbionts in three hydrothermal
703		vent mussels influence on host distributions. *Mar. Ecol. Prog. Ser.* **208**, 147-155 (2000).
704		https://doi:10.3354/meps208147
705	9	Kiel, S. The vent and seep biota : aspects from microbes to ecosystems. (Springer, 2010).
706		https://doi.org/10.1007/978-90-481-9572-5
707	10	Vrijenhoek, R. C. Genetic diversity and connectivity of deep-sea hydrothermal vent
708		metapopulations. *Mol. Ecol.* **19**, 4391-4411 (2010). https://doi.org:10.1111/j.1365-
709		294X.2010.04789.x
710	11	Halary, S., Riou, V., Gaill, F., Boudier, T. & Duperron, S. 3D FISH for the quantification of
711		methane- and sulphur-oxidizing endosymbionts in bacteriocytes of the hydrothermal vent
712		mussel *Bathymodiolus azoricus*. *ISME J* **2**, 284-292 (2008).
713		https://doi.org:10.1038/ismej.2008.3
714	12	Zheng, P. *et al.* Insights into deep-sea adaptations and host-symbiont interactions: A
715		comparative transcriptome study on Bathymodiolus mussels and their coastal relatives.
716		*Mol. Ecol.* **26**, 5133-5148 (2017). https://doi.org:10.1111/mec.14160
717	13	Fiala-Médioni, A., Métivier, C., Herry, A. & Le Pennec, M. *Ultrastructure of the gill of the*
718		*hydrothermal-vent mytilid Bathymodiolus sp.*. *Mar. Biol.* **92**, 65–72 (1986).
719		https://doi.org/10.1007/BF00392747
720	14	Wong, Y. H. *et al.* High-throughput transcriptome sequencing of the cold seep mussel
721		Bathymodiolus platifrons. *Sci. Rep.* **5**, 16597 (2015). https://doi.org:10.1038/srep16597

15 Sun, J. *et al.* Adaptation to deep-sea chemosynthetic environments as revealed by mussel
genomes. *Nat. Ecol. Evol.* **1**, 121 (2017). https://doi.org:10.1038/s41559-017-0121
16 Bettencourt, R. *et al.* An Insightful Model to Study Innate Immunity and Stress Response
in Deep‐Sea Vent Animals: Profiling the Mussel *Bathymodiolus azoricus*. *Organismal and*
Molecular Malacology, **8**, (2017). https://doi.org:10.5772/68034.
17 Barros, I. *et al.* Post-capture immune gene expression studies in the deep-sea
hydrothermal vent mussel *Bathymodiolus azoricus* acclimatized to atmospheric pressure.
Fish Shellfish Immunol **42**, 159-170 (2015). https://doi.org:10.1016/j.fsi.2014.10.018
18 Wang, H. *et al.* Molecular analyses of the gill symbiosis of the bathymodiolin mussel
Gigantidas platifrons. *iScience* **24**, 101894 (2021).
https://doi.org:10.1016/j.isci.2020.101894
19 Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. RNA imaging. Spatially
resolved, highly multiplexed RNA profiling in single cells. *Science* **348**, 6090 (2015).
https://doi.org:10.1126/science.aaa6090
20 Hwang, B., Lee, J. H. & Bang, D. Single-cell RNA sequencing technologies and
bioinformatics pipelines. *Exp. Mol. Med.* **50**, 1-14 (2018). https://doi.org:10.1038/s12276-
018-0071-8
21 Saliba, A. E., Westermann, A. J., Gorski, S. A. & Vogel, J. Single-cell RNA-seq: advances and
future challenges. *Nucleic Acids Res* **42**, 8845-8860 (2014).
https://doi.org:10.1093/nar/gku555
22 Chen, X., Teichmann, S. A. & Meyer, K. B. From tissues to cell types and back: single-cell
gene expression analysis of tissue architecture. *Annu. Rev. Biomed. Data Sci.* **1**, 29-51
(2018). https://doi.org:10.1146/annurev-biodatasci-080917-013452
23 Wu, H., Kirita, Y., Donnelly, E. L. & Humphreys, B. D. Advantages of single-nucleus over
single-cell RNA sequencing of adult kidney: rare cell types and novel cell states revealed in
fibrosis. *J. Am. Soc. Nephrol.* **30**, 23-32 (2019). https://doi.org:10.1681/ASN.2018090912
24 Feng, D. *et al.* Using Bathymodiolus tissue stable carbon, nitrogen and sulfur isotopes to
infer biogeochemical process at a cold seep in the South China Sea. *Deep Sea Res. Part I*
Oceanogr. Res. **104**, 52-59 (2015).
https://doi.org:https://doi.org/10.1016/j.dsr.2015.06.011
25 Kharchenko, P. V., Silberstein, L. & Scadden, D. T. Bayesian approach to single-cell
differential expression analysis. *Nat. Methods.* **11**, 740-742 (2014).
https://doi.org:10.1038/nmeth.2967
26 Elyanow, R., Dumitrascu, B., Engelhardt, B. E. & Raphael, B. J. netNMF-sc: leveraging gene-
gene interactions for imputation and dimensionality reduction in single-cell expression
analysis. *Genome. Res.* **30**, 195-204 (2020). https://doi.org:10.1101/gr.251603.119
27 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
(2021). https://doi.org:10.1016/j.cell.2021.04.048
28 Stuart, T. *et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902 (2019).
https://doi.org:10.1016/j.cell.2019.05.031
29 Petibon, C., Malik Ghulam, M., Catala, M. & Abou Elela, S. Regulation of ribosomal protein
genes: An ordered anarchy. *WIREs RNA* **12**, e1632 (2021).
https://doi.org:https://doi.org/10.1002/wrna.1632

30 Welcker, D. *et al.* Hemicentin-1 is an essential extracellular matrix component of the
dermal–epidermal and myotendinous junctions. *Sci. Rep.* **11**, 17926 (2021).
https://doi.org:10.1038/s41598-021-96824-4
31 Dufour, S. C. & Beninger, P. G. A functional interpretation of cilia and mucocyte
distributions on the abfrontal surface of bivalve gills. *Mar. Biol.* **138**, 295-309 (2001).
https://doi.org:10.1007/s002270000466
32 Dufour, S. C. Gill anatomy and the evolution of symbiosis in the bivalve family Thyasiridae.
Biol. Bull. **208**, 200-212 (2005). https://doi.org:10.2307/3593152
33 Gómez-Mendikute, A., Elizondo, M., Venier, P. & Cajaraville, M. P. Characterization of
mussel gill cells in vivo and in vitro. *Cell Tissue Res.* **321**, 131-140 (2005).
https://doi.org:10.1007/s00441-005-1093-9
34 Beninger, P. G. & Dufour, S. Mucocyte distribution and relationship to particle transport
on the pseudolamellibranch gill of *Crassostrea virginica* (Bivalvia:Ostreidae). *Mar. Ecol.*
Progr. Ser. **137**, 133-138 (1996). https://doi.org:10.3354/meps137133
35 Gerdol, M. *et al.* The C1q domain containing proteins of the Mediterranean mussel
Mytilus galloprovincialis: A widespread and diverse family of immune-related molecules.
Dev. Comp. Immunol. **35**, 635-643 (2011).
https://doi.org:https://doi.org/10.1016/j.dci.2011.01.018
36 Wang, W., Song, X., Wang, L. & Song, L. Pathogen-derived carbohydrate recognition in
molluscs immune defense. *Int. J. Mol. Sci.* **19**, 721 (2018).
https://doi.org:10.3390/ijms19030721
37 Saco, A., Rey-Campos, M., Novoa, B. & Figueras, A. Transcriptomic response of mussel
gills after a *Vibrio splendidus* infection demonstrates their role in the immune response.
Front. Immunol. **11**, 615580 (2020). https://doi.org:10.3389/fimmu.2020.615580
38 Richoux, N. B. & Thompson, R. J. Regulation of particle transport within the ventral groove
of the mussel (*Mytilus edulis*) gill in response to environmental conditions. *J. Exp. Mar.*
Biol. Ecol. **260**, 199-215 (2001). https://doi.org:10.1016/s0022-0981(01)00254-4
39 Riisgård, H. U., Egede, P. P. & Barreiro Saavedra, I. Feeding behaviour of the mussel,
Mytilus edulis: New observations, with a minireview of current knowledge. *J. Mar. Biol.*
2011, 312459 (2011). https://doi.org:10.1155/2011/312459
40 Heude, É., Shaikho, S. & Ekker, M. The dlx5a/dlx6a genes play essential roles in the early
development of zebrafish median fin and pectoral structures. *PLOS ONE* **9**, e98505 (2014).
https://doi.org:10.1371/journal.pone.0098505
41 Han, S. *et al.* TULP3 is required for localization of membrane-associated proteins ARL13B
and INPP5E to primary cilia. *Biochem. Biophys. Res. Commun.* **509**, 227-234 (2019).
https://doi.org:10.1016/j.bbrc.2018.12.109
42 Louka, P. *et al.* Proteins that control the geometry of microtubules at the ends of cilia. *J.*
Cell Biol. **217**, 4298-4313 (2018). https://doi.org:10.1083/jcb.201804141
43 Das, A., Dickinson, D. J., Wood, C. C., Goldstein, B. & Slep, K. C. Crescerin uses a TOG
domain array to regulate microtubules in the primary cilium. *Mol. Biol. Cell.* **26**, 4248-
4264 (2015). https://doi.org:10.1091/mbc.E15-08-0603
44 Llorente-Cortes, V., Martinez-Gonzalez, J. & Badimon, L. LDL receptor-related protein
mediates uptake of aggregated LDL in human vascular smooth muscle cells. *Arterioscler.*
Thromb. Vasc. Biol. **20**, 1572-1579 (2000). https://doi.org:10.1161/01.atv.20.6.1572

45 Chen, X. *et al*. Angiotensin-converting enzyme in smooth muscle cells promotes
atherosclerosis-brief report. *Arterioscler. Thromb. Vasc. Biol.* **36**, 1085-1089 (2016).
https://doi.org:10.1161/ATVBAHA.115.307038
46 Linke, W. A. & Grützner, A. Pulling single molecules of titin by AFM—recent advances and
physiological implications. *Pflug. Arch. Eur.* **456**, 101-115 (2008).
https://doi.org:10.1007/s00424-007-0389-x
47 St Paul, A., Corbett, C. B., Okune, R. & Autieri, M. V. Angiotensin II, Hypercholesterolemia,
and vascular smooth muscle cells: A perfect trio for vascular pathology. *Int. J. Mol. Sci.* **21**
4525 (2020). https://doi.org:10.3390/ijms21124525
48 Ytrehus, K., Ludvigsen, S., Mancusi, C., Gerdts, E. & de Simone, G. Heart angiotensin-
converting enzyme and angiotensin-converting enzyme 2 gene expression associated with
male sex and salt-sensitive hypertension in the Dahl Rat. *Front. physiol.* **12** (2021).
https://doi.org:10.3389/fphys.2021.663819
49 Keller, T. C., 3rd *et al*. Role of titin in nonmuscle and smooth muscle cells. *Adv. Exp. Med.*
Biol. **481**, 265-277 (2000). https://doi.org:10.1007/978-1-4615-4267-4_16
50 Chen, J. V. *et al*. Rootletin organizes the ciliary rootlet to achieve neuron sensory function
in *Drosophila*. *J. Cell. Biol.* **211**, 435-453 (2015). https://doi.org:10.1083/jcb.201502032
51 Styczynska-Soczka, K. & Jarman, A. P. The *Drosophila* homologue of rootletin is required
for mechanosensory function and ciliary rootlet formation in chordotonal sensory
neurons. *Cilia* **4**, 9 (2015). https://doi.org:10.1186/s13630-015-0018-9
52 Mohan, S., Timbers, T. A., Kennedy, J., Blacque, O. E. & Leroux, M. R. Striated rootlet and
nonfilamentous forms of rootletin maintain ciliary function. *Curr. Biol.* **23**, 2016-2022
(2013). https://doi.org:10.1016/j.cub.2013.08.033
53 Jones, S. E. & Jomary, C. Secreted Frizzled-related proteins: searching for relationships
and patterns. *Bioessays* **24**, 811-820 (2002). https://doi.org:10.1002/bies.10136
54 Gillis, W. Q., Bowerman, B. A. & Schneider, S. Q. The evolution of protostome GATA
factors: molecular phylogenetics, synteny, and intron/exon structure reveal orthologous
relationships. *BMC Evol. Biol.* **8**, 112-112 (2008). https://doi.org:10.1186/1471-2148-8-
112
55 Zhang, Z., Liu, L., Twumasi-Boateng, K., Block, D. H. S. & Shapira, M. FOS-1 functions as a
transcriptional activator downstream of the *C. elegans* JNK homolog KGB-1. *Cell. Signal.*
30, 1-8 (2017). https://doi.org:https://doi.org/10.1016/j.cellsig.2016.11.010
56 Phochanukul, N. & Russell, S. No backbone but lots of Sox: Invertebrate Sox genes. *Int. J.*
Biochem. Cell Biol. **42**, 453-464 (2010).
https://doi.org:https://doi.org/10.1016/j.biocel.2009.06.013
57 Cannuel, R., Beninger, P. G., McCombie, H. & Boudry, P. Gill Development and its
functional and evolutionary implications in the blue mussel *Mytilus edulis* (Bivalvia:
Mytilidae). *Biol. Bull.* **217**, 173-188 (2009). https://doi.org:10.1086/BBLv217n2p173
58 Leibson, N. L. & Movchan, O. T. Cambial zones in gills of Bivalvia. *Mar. Biol.* **31**, 175-180
(1975). https://doi.org:10.1007/BF00391629
59 Wentrup, C., Wendeberg, A., Schimak, M., Borowski, C. & Dubilier, N. Forever competent:
deep-sea bivalves are colonized by their chemosynthetic symbionts throughout their
lifetime. *Environ. Microbiol.* **16**, 3699-3713 (2014). https://doi.org:10.1111/1462-
2920.12597

60 Mohieldin, A. M. *et al.* Proteomic identification reveals the role of ciliary extracellular-like
vesicle in cardiovascular function. *Adv. Sci.* **7**, 1903140 (2020).
https://doi.org:https://doi.org/10.1002/advs.201903140
61 Neumann, D. & Kappes, H. On the growth of bivalve gills initiated from a lobule-producing
budding zone. *Biol. Bull.* **205**, 73-82 (2003). https://doi.org:10.2307/1543447
62 Streams, M. E., Fisher, C. R. & Fiala-Médioni, A. Methanotrophic symbiont location and
fate of carbon incorporated from methane in a hydrocarbon seep mussel. *Mar. Biol.* **129**,
465-476 (1997). https://doi.org:10.1007/s002270050187
63 Ponnudurai, R. *et al.* Metabolic and physiological interdependencies in the *Bathymodiolus*
azoricus symbiosis. *ISME J* **11**, 463-477 (2017). https://doi.org:10.1038/ismej.2016.124
64 Barry, J. P. *et al.* Methane‐based symbiosis in a mussel, *Bathymodiolus platifrons*, from
cold seeps in Sagami Bay, Japan. *Invertebr. Biol.* **121**, 47-54 (2002).
https://doi.org:doi:10.1111/j.1744-7410.2002.tb00128.x
65 Renoz, F., Noel, C., Errachid, A., Foray, V. & Hance, T. Infection dynamic of symbiotic
bacteria in the pea *Aphid acyrthosiphon pisum* gut and host immune response at the early
steps in the infection process. *Plos One* **10,** 0122099 (2015).
https://doi.org:ARTNe012209910.1371/journal.pone.
66 Guan, H. *et al.* Lipid Biomarker Patterns Reflect Nutritional Strategies of Seep-Dwelling
Bivalves From the South China Sea. *Front. Mar. Sci.* **9,** 3398 (2022).
https://doi.org:10.3389/fmars.2022.831286
67 Li, Y. *et al.* Genomic adaptations to chemosymbiosis in the deep-sea seep-dwelling
tubeworm *Lamellibrachia luymesi*. *BMC Biol.* **17**, 91 (2019).
https://doi.org:10.1186/s12915-019-0713-x
68 Sun, Y. *et al.* Genomic signatures supporting the symbiosis and formation of chitinous
tube in the deep-sea tubeworm *Paraescarpia echinospica*. *Mol. Biol. Evol.* **38**, 4116-4134
(2021). https://doi.org:10.1093/molbev/msab203
69 Takishita, K. *et al.* Genomic evidence that methanotrophic endosymbionts likely provide
deep-sea bathymodiolus mussels with a sterol intermediate in cholesterol biosynthesis.
Genome Biol. Evol. **9**, 1148-1160 (2017). https://doi.org:10.1093/gbe/evx082
70 Demidenko, A., Akberdin, I. R., Allemann, M., Allen, E. E. & Kalyuzhnaya, M. G. Fatty acid
biosynthesis pathways in *Methylomicrobium buryatense* 5G(B1). *Front. Microbiol.* **7**, 2167
(2016). https://doi.org:10.3389/fmicb.2016.02167
71 Brasaemle, D. L. *et al.* Adipose differentiation-related protein is an ubiquitously expressed
lipid storage droplet-associated protein. *J. Lipid Res.* **38**, 2249-2263 (1997).
https://doi.org/10.1016/S0022-2275(20)34939-7
72 Cheng, D. *et al.* Expression, purification, and characterization of human and rat acetyl
coenzyme A carboxylase (ACC) isozymes. *Protein Expr. Purif.* **51**, 11-21 (2007).
https://doi.org:10.1016/j.pep.2006.06.005
73 Monroig, Ó. & Kabeya, N. Desaturases and elongases involved in polyunsaturated fatty
acid biosynthesis in aquatic invertebrates: a comprehensive review. *Fish. Sci.* **84**, 911-928
(2018). https://doi.org:10.1007/s12562-018-1254-x
74 Kabeya, N. et al. Unique fatty acid desaturase capacities uncovered in Hediste diversicolor
illustrate the roles of aquatic invertebrates in trophic upgrading. *Philos. Trans. R. Soc.* **375**,
20190654 (2020). https://doi.org:10.1098/rstb.2019.0654

75 Soupene, E. & Kuypers, F. A. Mammalian long-chain acyl-CoA synthetases. *Exp. Biol. Med.*
(Maywood). **233**, 507-521 (2008). https://doi.org:10.3181/0710-MR-287
76 Alves-Bezerra, M. *et al.* Long-chain acyl-CoA synthetase 2 knockdown leads to decreased
fatty acid oxidation in fat body and reduced reproductive capacity in the insect *Rhodnius*
prolixus. Biochim. Biophys. Acta. **1861**, 650-662 (2016).
https://doi.org:10.1016/j.bbalip.2016.04.007
77 Hoglund, P. J., Nordstrom, K. J. V., Schioth, H. B. & Fredriksson, R. The solute carrier
families have a remarkably long evolutionary history with the majority of the human
families present before divergence of bilaterian species. *Mol. Biol. Evol.* **28**, 1531-1541
(2011). https://doi.org:10.1093/molbev/msq350
78 Mohamed, A. R. *et al.* Dual RNA-sequencing analyses of a coral and its native symbiont
during the establishment of symbiosis. *Mol. Ecol.* **29**, 3921-3937 (2020).
https://doi.org:10.1111/mec.15612
79 Bertucci, A., Foret, S., Ball, E. E. & Miller, D. J. Transcriptomic differences between day and
night in *Acropora millepora* provide new insights into metabolite exchange and light-
enhanced calcification in corals. *Mol. Ecol.* **24**, 4489-4504 (2015).
https://doi.org:10.1111/mec.13328
80 Hamada, M. *et al.* Metabolic co-dependence drives the evolutionarily ancient Hydra-
Chlorella symbiosis. *Elife* **7**, e35122 (2018). https://doi.org:10.7554/eLife.35122
81 Duncan, R. P., Feng, H., Nguyen, D. M. & Wilson, A. C. Gene family expansions in aphids
maintained by endosymbiotic and nonsymbiotic traits. *Genome Biol. Evol.* **8**, 753-764
(2016). https://doi.org:10.1093/gbe/evw020
82 Feng, H. L. *et al.* Trading amino acids at the aphid-Buchnera symbiotic interface. *P. Natl.*
Acad. Sci. USA. **116**, 16003-16011 (2019). https://doi.org:10.1073/pnas.1906223116
83 Ip, J. C. *et al.* Host-endosymbiont genome integration in a deep-sea chemosymbiotic clam.
Mol. Biol. Evol. **38**, 502-518 (2021). https://doi.org:10.1093/molbev/msaa241
84 Hongo, Y. *et al.* Expression of genes involved in the uptake of inorganic carbon in the gill
of a deep-sea vesicomyid clam harboring intracellular thioautotrophic bacteria. *Gene* **585**,
228-240 (2016). https://doi.org:https://doi.org/10.1016/j.gene.2016.03.033
85 Sotiriou, S. *et al.* Ascorbic-acid transporter Slc23a1 is essential for vitamin C transport into
the brain and for perinatal survival. *Nat. Med.* **8**, 514-517 (2002).
https://doi.org:10.1038/0502-514
86 Aydemir, T. B. et al. Zinc transporter ZIP14 functions in hepatic zinc, iron and glucose
homeostasis during the innate immune response (endotoxemia). *PLoS One* **7**, e48679
(2012). https://doi.org:10.1371/journal.pone.0048679
87 Turi, Z., Lacey, M., Mistrik, M. & Moudry, P. Impaired ribosome biogenesis: mechanisms
and relevance to cancer and aging. Aging (Albany NY) **11**, 2512-2540 (2019).
https://doi.org:10.18632/aging.101922
88 Hay, B. A. Understanding IAP function and regulation: a view from Drosophila. Cell. Death.
Differ. **7**, 1045-1056 (2000). https://doi.org:10.1038/sj.cdd.4400765
89 Dubrez-Daloz, L., Dupoux, A. & Cartier, J. IAPS : More than just inhibitors of apoptosis
proteins. Cell Cycle **7**, 1036-1046 (2008). https://doi.org:10.4161/cc.7.8.5783
90 Pflug, K. M. & Sitcheran, R. Targeting NF-κB-inducing kinase (NIK) in immunity,
inflammation, and cancer. *Int. J. Mol. Sci.* **21**, 8470 (2020).

91 Otten, E. G. *et al.* Ubiquitylation of lipopolysaccharide by RNF213 during bacterial
infection. *Nature* **594**, 111-116 (2021). https://doi.org:10.1038/s41586-021-03566-4
92 Tashian, R. E. The carbonic anhydrases: widening perspectives on their evolution,
expression and function. *Bioessays* **10**, 186-192 (1989).
https://doi.org:10.1002/bies.950100603
93 Price, D. R. *et al.* Aphid amino acid transporter regulates glutamine supply to intracellular
bacterial symbionts. *Proc. Natl. Acad. Sci. USA.* **111**, 320-325 (2014).
https://doi.org:10.1073/pnas.1306068111
94 Ponnudurai, R. *et al.* Comparative proteomics of related symbiotic mussel species reveals
high variability of host–symbiont interactions. *ISME J.* **14**, 649-656 (2020).
https://doi.org:10.1038/s41396-019-0517-6
95 Hicks, S. C., Townes, F. W., Teng, M. & Irizarry, R. A. Missing data and technical variability
in single-cell RNA-sequencing experiments. *Biostatistics* **19**, 562-578 (2017).
https://doi.org:10.1093/biostatistics/kxx053
96 Lun, A. T. L. *et al.* EmptyDrops: distinguishing cells from empty droplets in droplet-based
single-cell RNA sequencing data. *Genome Biol.* **20**, 63 (2019).
https://doi.org:10.1186/s13059-019-1662-y
97 McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet detection in single-
cell RNA sequencing data using artificial nearest neighbors. *Cell Systems* **8**, 329-337
(2019). https://doi.org:https://doi.org/10.1016/j.cels.2019.03.003
98 Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e1821
(2019). https://doi.org:https://doi.org/10.1016/j.cell.2019.05.031
99 Singh, P. & Zhai, Y. Deciphering Hematopoiesis at single cell level through the lens of
reduced dimensions. *bioRxiv*, 2022.2006.2007.495099 (2022).
https://doi.org:10.1101/2022.06.07.495099
100 Wu, T. et al. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
The Innovation **2**, 100141 (2021).
https://doi.org:https://doi.org/10.1016/j.xinn.2021.100141
101 Street, K. et al. Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genom.* **19**, 477 (2018). https://doi.org:10.1186/s12864-018-4772-0
102 Moon, K. R. et al. Visualizing structure and transitions in high-dimensional biological data.
Nat. Biotechnol. **37**, 1482-1492 (2019). https://doi.org:10.1038/s41587-019-0336-3
103 Thisse, C. & Thisse, B. High-resolution *in situ* hybridization to whole-mount zebrafish
embryos. *Nat. Protoc.* **3**, 59-69 (2008). https://doi.org:10.1038/nprot.2007.514

A

In situ experiment

Sample dissection, on-board treatment and storage

Nuclei extraction and gradient centrifugation

"Fanmao" Site
CH₄:~44.23μM
DO :~3.01mg/L

Fanmao

Transplant

Starvation Site

3 days of Recovery

Reconstitution

CH₄:~0.054μM
DO :~3.20mg/L

11 days of CH₄ starvation

14 days of CH₄ starvation

Starvation

Liquid nitrogen and store in -80℃

4% PFA fixation and store in MeOH at -20℃

BD Rhapsody snRNA-seq

Whole mount *in situ* hybridizaiton (WISH) and ISH and IHC on paraffin slices with Fanmao mussels

Illumina sequencing

Cell cluster specific environmental responds

Cell cluster specific DEG analysis

Starvation
Reconstitution
Fanmao

Cell clusters validation and spatial mapping using WISH and ISH

Bacteriocytes
Budding zone
Intercalary cells
Food groove
Ciliary cells
......

Integration and cell clustering

Bioinformatics pipelines

snRNA-Seq Pipeline
Seurat
Monocle

B

1cm

C

Intercalary cell

Food groove ciliary cells

Bacteriocyte

Apical ridge ciliary cell

Posterior end budding zone cell

Lateral ciliary cell

Smooth muscle

Dorsal end proliferation cell

Inter lamina cell

Ventral end proliferation cell

Basal membrane cell 1

Ambiguous

Mucus cell

Basal membrane cell 2

UMAP_2

UMAP_1

D

Posterior end budding zone cell
Ventral end proliferation cell
Inter lamina cell
Bacteriocyte
Lateral ciliary cell
Apical ridge ciliary cell
Intercalary cell
Mucus cell
Ambiguous
Basal membrane cell 1
Basal membrane cell 2
Dorsal end proliferation cell
Smooth musle proliferation cell
Food groove ciliary cell