Ripply1 and Gsc collectively suppress anterior endoderm differentiation from prechordal plate progenitors

Tao Cheng[1,2#], Xiang Liu[2#], Yang Dong[1,2#], Yi-Meng Tian[2], Yan-Yi Xing[1,2], Chen-Yi Chen[2], Cong Liu[2], Yun-Fei Li[2], Ying Huang[2], Ding-Hao Zhuo[2], Xiao Xu[1], Jing-Yun Luan[2], Xin-Xin Fu[2], Zi-Xin Jin[2], Jing Mo[3], Xiang Xu[2], Hong-Qing Liang[1,2]*, Peng-Fei Xu[1,2]*

[1] Women's Hospital, Zhejiang University School of Medicine, Hangzhou, China.

[2] Institute of Genetics and Department of Human Genetics, Zhejiang University School of Medicine, Hangzhou, China.

[3] Department of Immunology, Guizhou Medical University, Guiyang 550004, China.

*Corresponding author: lianghongqing@zju.edu.cn, pengfei_xu@zju.edu.cn

[#]Tao Cheng, Xiang Liu and Yang Dong contributed equally to this work and should be regarded as joint first authors.

Abstract:

During gastrulation, the mesendoderm is firstly specified by morphogens such as Nodal, and then segregates into endoderm and mesoderm in a Nodal concentration-dependent manner. However, the mechanism underlying the segregation and crosstalk of different sub-groups within the meso- and endoderm lineages remains unclear. Here, taking zebrafish prechordal plate (PP) and anterior endoderm (Endo) as research model, using single-cell multi-omics and live imaging analyses, we show that anterior Endo progenitors originate directly from PP progenitors. A single-cell transcriptomic trajectory analysis of wild-type, *ndr1* knockdown and *lft1* knockout Nodal explants confirms the diversification of anterior Endo fate from PP progenitors. Gene Ontology (GO) enrichment analysis indentifies that the change of chromatin organization potentiates the segregation of endodermal cell fate from PP progenitors. Single-cell ATAC & RNA sequencing further reveals that two transcriptional regulators, *gsc* and *ripply1*, exhibit varied activation patterns in PP and Endo lineages at both the chromatin and RNA expression levels. We further demonstrate that Ripply1 functions coordinately with Gsc to repress endodermal cell fate by directly binding to the *cis*-elements of *sox32* and *sox17*. Modulating the expression levels of these regulators tilts the cell fate decision between the PP and Endo lineages.

Introduction:

During gastrulation, instructed by genetic and epigenetic signals, naïve cells from blastula progressively acquire distinct cell fates and are allocated to specific domains within the embryo[1,2]. Nodal, a member of the transforming growth factor β (TGF-β) superfamily, acts as a morphogen[1,3-5] and plays highly conserved roles in mesendoderm induction and patterning[6-8].

It is widely accepted that Nodal induces endodermal and mesodermal cell fates via its concentration gradient[8-10]. Higher Nodal signaling activity bias cells toward endoderm, while lower levels are thought to promote mesoderm induction[11,12]. Mesodermal and endodermal common progenitors are induced by Nodal and mixed together at the onset of gastrulation[1], while it remains elusive how the common progenitors of mesendoderm took on distinct lineages. Several recent studies reported that multiple important signaling pathways interacted with

Nodal to drive mesendoderm separation[2,13-15]. Notably, lateral endoderm specification may

comply with a stochastic cell fate switching model in zebrafish[16].

The specification of dorsal mesendoderm requires the highest level of Nodal signaling, as

evidenced by that this cell population was first to be depleted when Nodal signaling was

gradually inhibited[11]. Both the PP and anterior Endo originate from anterior mesendoderm

(derived from the dorsal mesendoderm), and the commitment of these cell fates is regulated by

similar levels of Nodal signaling. However, the mechanisms governing the separation of cell

fates within the anterior mesendoderm remain unclear. Previous studies show that Nodal

signaling enhances the duration of cell-cell contact in PP cells, creating a positive feedback

loop between cell-cell contact duration and cell fate specification[17]. However, what factors

downstream of Nodal pathway play major roles in segregating these two cell fates, when and

how PP and Endo cells can be distinguished transcriptionally are the questions remaining to be

investigated.

Previous research has shown that Gsc, whose expression can be induced by Nodal, acts as

a repressor to suppress anterior endoderm specification by binding to the promoter region of

sox17[18]. However, investigations into loss-of-function of Gsc in zebrafish have not yielded

conclusive evidence supporting its significant roles in mesoderm and/or endoderm

specification[19,20]. These results imply that there might be additional, unidentified transcriptional

repressors that work in collaboration with Gsc and possess a redundant function in suppressing

the specification of endoderm.

In this study, we constructed a single-cell transcriptional trajectory to delineate the cell

state segregation between PP and anterior Endo in zebrafish embryos and Nodal-injected

zebrafish explants[21,22]. Single-cell transcriptional trajectory and live imaging analyses

demonstrated that anterior Endo cells originated and were specified from PP progenitors.

Interestingly, this lineage separation can be traced back to as early as the onset of gastrulation.

To achieve a more comprehensive understanding, we integrated single-cell datasets from wild-

type, *ndr1* (Nodal ligand) knockdown and *lft1* (Nodal inhibitor) knockout Nodal explants at

shield stage to construct an extended single-cell trajectory, which fully captured the branching

event between PP and anterior Endo fates and unveiled a potential involvement of epigenetic

regulation in this process. This finding was further supported by the experimental evidence that

perturbation of the chromatin remodeler *srcap* could affect the cell specification of PP and

anterior Endo. Furthermore, combining the multi-omic analysis and gain-/loss-of-function

experiments, we found that a transcriptional repressor, Ripply1, cooperated with Gsc to

supresse endoderm specification by directly binding to the regulatory elements of *sox17* and

sox32.

Results:

Anterior Endo originates from PP progenitors in zebrafish.

Both anterior Endo and PP are derived from embryonic shield and require high Nodal signaling

to be specified[11,18,23]. To investigate how these two cell lineages are further segregated, we

collected the single-cell RNA-seq datasets from the published studies[24,25], identified the

subpopulations representing the anterior Endo and PP cells, and reconstructed their lineage

branching tree (Figures 1A-1C and S1A). We found that these two lineages started to be

separated at shield stage, and their common progenitors highly expressed PP marker genes, like

gsc, frzb, etc. (Figures 1C and S1A). Then, we further explored the dynamic molecular

signatures of these two lineages during their fate determination. Pseudotime analysis revealed

that PP and Endo exhibited distinct differentiation states by the end of gastrulation (Figure S1B).

Of note, compared to *sox32*, *gsc* displayed earlier transcriptional differences in these two

lineages (Figure S1C). The Uniform Manifold Approximation and Projection (UMAP)

analysis[26] and PCA analysis showed that the common progenitors of PP and Endo clustered

closely to PP and away from Endo (Figures S1D and S1E), suggesting that anterior Endo

lineage is branched out from PP lineage. The cell lineage prediction analysis[27] also revealed

that the progenitor cells resembled more of the PP lineage (Figures 1D, 1E and S1F), and similar

result was obtained using another single-cell RNA-seq dataset (Figures S1G-S1I). Therefore,

both UMAP and the cell lineage prediction analysis suggest that anterior Endo and PP originate

from the same progenitor with transcriptome profile more similar to PP lineage. Anterior Endo

becomes transcriptionally distinguishable from PP as early as at the shield stage.

Cells from hatching gland (derived from PP) and pharyngeal pouch (derived from anterior

Endo) are distant on the transcriptome-based phylogenetic tree of zebrafish embryos[25]. Thus, the resolution of this dataset may not be sufficient to identify the exact lineage separation point between PP and Endo. To investigate the key molecules that regulate the PP and Endo separation, we employed our previously published Nodal induced embryonic explant system (referred to Nodal explant hereafter), which encompasses the axial mesendoderm, along with the PP and anterior Endo lineages[22]. Notably, the single-cell trajectory tree of Nodal explants showed a closer transcriptomic relationship between the PP and anterior Endo (Figure 1F). Thus, we reconciled the single-cell transcriptome dynamics to look for the branching point of PP and anterior Endo in the Nodal explants (Figures 1G, 1H, S2A and S2B). Consistent with the observations in wild-type embryos, the common progenitors of these two lineages highly expressed the PP marker gene (*gsc*), and the segregation between PP and Endo had already occurred by 6 hpf (hours post fertilization) (Figures 1G, 1H and S2B). Cell lineage prediction analysis on the Nodal explant datasets further indicated that the common progenitor cells were transcriptionally similar to PP (Figures 1I and S2I), which aligned with our previous observations in wild-type embryos (Figures 1D, 1E, S1H and S1I).

To explore the molecular signatures during the specification of PP and anterior Endo, we performed differential expression analysis for these two lineages in both wild-type embryos and Nodal explants. 18 genes were highly enriched along the trajectory of Endo specification in both wild-type embryos and Nodal explants (Figure S2C). These genes enriched GO terms such as "gland development", "negative regulation of transcription, DNA-templated" and "negative regulation of RNA biosynthetic process" (Figure S2D). While for PP, 49 highly expressed genes were identified (Figure S2C), which enriched GO terms including "proteolysis involved in cellular protein catabolic process", "cellular protein catabolic process" , "protein catabolic process", etc. (Figure S2D). Cell fate specification requires the activation of a series of genes, and these gene sets can be regarded as gene modules, which are probably responsible for specific lineage formation[28]. We identified two gene modules associated with Endo and PP lineages respectively (Figure S2E). Gene Ontology enrichment analyses[29] showed that terms like "endoderm development" and "cell motility" were enriched for Endo (gene module 1), while "embryo development ending in birth or egg hatching" was enriched for PP (gene module

2) (Figures S2F-S2H). These enriched GO terms were highly correlated with the developmental

processes and functions of these two cell types.

To further validate this observation and investigate the dynamics of lineage segregation

between anterior Endo and PP during development, we performed live imaging analysis on

embryos of Tg(*gsc*:EGFP;*sox17*:DsRed) (Figures 2A and 2B). Notably, we observed a gradual

emergence of *sox17*-positive cells from a subset of the *gsc*-positive cells (Figures 2A-2D, S3A-

S3D and Movies S2, S3). And this cell fate transition was also evident in Nodal explants

constructed from Tg(*gsc*:EGFP;*sox17*:DsRed) embryos (Figures S3E-S3H and Movie S1).

Furthermore, we generated *sox17*:Cre and *gsc*:loxP-STOP-loxP-mCherry constructs (Figure

S4D) and co-injected them with transposase mRNA into one-cell stage embryos (Figure S4A).

Interestingly, distinct mCherry-positive cells were observed in the injected embryos, providing

clear evidence that *gsc*-positive cells can give rise to *sox17*-positive cells (Figures S4B and

S4C).

We co-stained *gsc* and *sox17* in embryos and in Nodal explants respectively at different

time points during gastrulation (Figures 2E and S3I). By assessing the expression levels of

sox17 and *gsc* in *sox17*-positive anterior mesendoderm cells, we observed a positive correlation

between *sox17* and *gsc* expression at the onset of gastrulation. However, with advancing

development, *sox17* and *gsc* expression exhibited a negative correaltion in both embryos and

Nodal explants (Figures 2F and S3J). This suggests that the increase of *sox17* expression during

anterior Endo differentiation is accompanied by the simultaneous reduction of *gsc* expression.

These findings align with our live imaging experiments (Figures 2A, S3B and S3F). In summary,

combing single-cell tanscriptomic trajectory analyses and live imaging experiments using

embryos and explants, we demonstrated that anterior Endo cells were derived from PP

progenitor cells in zebrafish.

Nodal-Lefty regulatory loop is needed for PP and anterior Endo fate specification

In zebrafish, the PP and anterior Endo lineages are physically in close proximity and direct

contact[17]. During development, interactions through cell-cell contacts between neighboring cell

types often play a crucial role in promoting cell fate segregation[17,30]. We hypothesized that

communications between PP and anterior Endo cells may influence each other's cell fate

specification. We employed LIANA[31], a ligand-receptor analysis framework, to investigate the

molecular signals involved in these interactions using single-cell RNA-seq datasets. Our

analysis revealed strong interactions between PP and Endo cells in both wild-type embryos and

Nodal explants (Figures S5C-S5F). Interestingly, key molecular factors in the Nodal-Lefty

regulatory loops (Figure 3D), such as *ndr1*, *lft1* and *lft2*, were significantly enriched when

setting PP as source cells and Endo as target cells (Figures 3A, S5A, S5B and S5G-S5I), while

Nodal-Lefty interactions were reduced when setting the cell types conversely (Figures 3B and

S5G-S5I). In addition to Nodal-Lefty signaling, Fgf and Wnt signalings were also significantly

enriched in cell-cell communications between these two cell types in both embryos and Nodal

explants (Figures 3A, 3B and S5G-S5I). This observation is consistent with previous findings

that Fgf signaling interacts with Nodal to regulate mesoderm and endoderm separation[13,32].

The results presented above suggest that differences in the Nodal-Lefty signaling may exist

between PP and anterior Endo cells. It is widely accepted that Nodal-Lefty functions as

reaction-diffusion system to facilitate Nodal gradient formation and cell fate determination[33-37]

(Figure 3D). As zebrafish possess redundant Nodal (Ndr1, Ndr2 and Ndr3) and Lefty (Lefty1

and Lefty2) paralogs, we firstly investigated the perturbation effects of individual Nodal or

Lefty paralog on Nodal activity and its gradient. We generated an ectopic Nodal gradient in

zebrafish animal pole cells by injection of *ndr2* mRNA at 128-cell stage in wild-type embryos,

ndr1 morphants and *lft1* mutants (Figures 3E, S6A-S6F and S6H-S6M). By immunostaining

and quantification of the pSmad2 signaling, we observed a significant enhancement of Nodal

activity in *lft1* mutants, while its level decreased upon *ndr1* knockdown (Figures 3E, 3F and

S6A-S6M).

To further determine how Nodal-Lefty is involved in the regulation of PP and anterior

Endo separation, we performed single-cell transcriptomic sequencing for Nodal explants

constructed from *ndr1* morphants and *lft1* mutants at shield stage (Figure 3G). We integrated

these two single-cell RNA-seq datasets with the dataset of wild-type Nodal explants[22]. After

applying unsupervised clustering, we identified a total of 9 different cell clusters (Figures 3H,

S7A and S7B). Among these clusters, two mesodermal clusters named mesoderm_PP and

mesoderm_NT (Figure 3H) were obtained. Mesoderm_PP cells transcriptionally resembled the progenitors of PP evidenced by the high expression of *gsc* and *chrd*, while mesoderm_NT cells were the progenitors of notochord characterized by the high expression of *noto* and *tbxta* (Figures S7A and S7B). We observed that a small cluster of *sox32* and *sox17* positive anterior Endo cells was close to the PP cells on the UMAP (Figure 3H), further supporting the transcriptional similarity between anterior Endo and the PP lineage. We then calculated the proportions of PP and Endo cells in the three datasets, and found that the proportions of both cell types increased in *lft1*-mutant explants, while decreased in *ndr1*-morphant explants compared to wild-type explants (Figures 3I and S7C).

To better estimate Nodal activity during PP and Endo cell fate segregation, we defined a Nodal score by regressing out the expression level of all the Nodal downstream targets, which were differentially expressed upon Nodal treatment[22] and also possessed pSmad2 binding peaks near their genetic loci[22,38] (**see Methods**). A total of 29 Nodal downstream targets, whose expression exhibited a high correlation with Nodal activity, were used to calculate Nodal score (Figure 3J). We found Nodal score was dramatically decreased in *ndr1* knockdown samples and slightly increased in *lft1* mutant samples compared to that in wild-type samples (Figures 3K and S7D), which is consistent with the results of pSmad2 immunostaining (Figures 3E and 3F). We then examined the levels of Nodal score across all cell types in Nodal explants at 6 hpf, and found that axial mesendodermal cells, including PP, notochord and anterior Endo, displayed the highest Nodal scores (Figure S7E), which was consistent with previous knowledge that Nodal activity was the highest in the dorsal margin of zebrafish embryos at the onset of gastrulation[1,39,40]. A further comparsion of Nodel scores between PP and Endo revealed that PP had a sightly higher Nodal score than Endo (Figure S7E). These analyses underscored the importance of Nodal signaling levels in PP and anterior Endo lineage specification.

A meticulous single-cell trajectory tree reveals the role of chromatin remodeling in PP and anterior Endo segregation.

To thoroughly investigate the dynamic transcriptome changes during PP and anterior Endo separation, we reconstructed a single-cell trajectory tree using all PP and anterior Endo cells

from three Nodal explant datasets at 6 hpf which was a crucial time point previously identified for the separation of these two lineages (Figure 4A). This trajectory tree captured a continuous cell-state-transition process from PP progenitors to anterior Endo, and a key branching point in this process was identified (Figures 4A and S7F). To elucidate the molecular dynamics in this branching process, we performed differential expression analysis for these two lineages around the branching point (Figure 4B). Three clusters of genes were identified, which were highly expressed in pre-branch (progenitors), left-branch (Endo) and right-branch (PP) respectively. GO enrichment analysis using these three gene sets identified terms related to mesendodermal cell fate specification (Figure 4B). For instance, "regulation of endodermal cell fate specification" was observed in Endo lineage, while "negative regulation of transcription by RNA polymerase II" was found in PP lineage. Interestingly, "chromatin organization" was significantly enriched in the pre-branch cells, and maintained high enrichment during PP specification, but was down-regulated upon endodermal lineage formation (Figure 4B). Furthermore, the GO term "chromatin organization" was also enriched in the differentially expressed genes of the PP cells when comparing *lft1* mutant or *ndr1* knockdown samples to wild-type samples (Figures 4C-4G and S7G).

These analyses unveiled a potential role of chromatin regulators in the regulation of PP and anterior Endo segregation, prompting us to identify the specific factors involved in this process. Among the candidates within GO term of "chromatin organization" (Figure 4B-4E), we identified a key member of SWI/SNF complex, *srcap*, which was differentially expressed between PP and Endo cells in both embryos and Nodal explants (Figures 4J, 4K and S7H). Previous studies have demonstrated that the SWI/SNF nucleosome remodeling complex is necessary for TGF-β-induced transcription of numerous target genes[41,42]. To investigate the role of the SWI/SNF complex in the separation of PP and Endo cell fates , we treated embryos with AU15330 (Figure 4H), a known degrader of SWI/SNF ATPase components[43]. Subsequently, we assessed the cell fate specification of PP and anterior Endo lineages by HCR co-staining for *gsc*, *frzb* and *sox32*. We found that the numbers of anterior Endo cells were significantly increased in the embryos treated with AU15330 compared to DMSO-treated embryos (Figures 4H and 4I). Similar effects were observed in *srcap* morphants, where the cell number of anterior

Endo increased compared to wild-type embryos (Figures 4L, 4M, and S7I).

Collectively, these findings underscore the importance of SWI/SNF complexes in ensuring

accurate cell fate segregation within the anterior mesendoderm.

Integrative analysis of single-nucleus RNA-Seq and ATAC-Seq reveals distinct chromatin

states underlying differential expression profiling in PP and Endo cells.

To delve deeper into the involvement of chromatin states in the separation of PP and Endo, we

conducted single-cell assays for transposase-accessible chromatin (ATAC) and nuclear RNA

sequencing in zebrafish embryos at 6 hpf (Figure 5A). Recognizing that accurate cell clustering

cannot solely rely on chromatin accessibility dataset, we integrated this dataset with RNA

expression data to achieve precise cell type identification. The resulting cells were clustered

into 10 different groups on the UMAP (Figures 5B, S8A, and S8B). Consistent with the earlier

single-cell RNA atlas of Nodal explants, Endo was found to locate closely to the axis mesoderm

(PP and notochord progenitors) on the joint UMAP (Figure 5B).

Using the marker gene *gsc*, we identified PP progenitor cells within the axis mesoderm,

and then Endo and PP progenitor cells were selected for downstream analysis (Figure 5C). A

small cluster of cells expressing *sox32*, *sox17* and *foxj1a*, which were located distantly from PP

and Endo, were classified as dorsal forerunner cells (DFCs) and excluded from further analysis

(Figures 5C and S8C).

Our data unveiled differential chromatin accessibility between PP and Endo cells (Figures

5G, S8K, and S8L). Correlation analysis revealed a strong correspondence between chromatin

accessibility and RNA expression levels for marker genes in these two cell types (Figure S8D).

To identify master regulators of PP and Endo separation, we performed differential analysis for

both chromatin accessibility and gene expression (Figures 5D and S8E). A total of 170 up-

regulated genes in PP, including *gsc* and *ripply1*, and 261 up-regulated genes in Endo, such as

sox32 and *sox17*, were identified. Most of these genes also exhibited differential chromatin

opening states in PP and anterior Endo lineages (Figures 5E-5G and S8F-S8I).

Previous study reported that Nodal signaling can boost chromatin accessibility both *in vivo*

and *ex vivo*[44,45] . And we observed that PP displayed a higher Nodal levels compared to Endo

cells (Figures S2J and S7E). In this ATAC and RNA-seq dataset, we also observed a higher Nodal score in PP cells (Figures 5H and 5I). Thus, it is possible that the differences in Nodal activity may lead to differential chromatin accessibility states and gene expression level in PP and Endo.

To further determine how Nodal siganling levels may differentially influence the gene expression dynamics, we first explored the expression profiles of key markers of anterior mesendoderm in the bulk RNA-seq data of Nodal explants injected with different dosages of Nodal mRNA[22]. We observed that the highest Nodal dosages did not result in further elevation of the expression of the endodermal marker *sox17* (Figure 5J), while higher concentrations of Nodal promoted the expression of *gsc* and *ripply1* (Figure 5J). Importantly, these patterns of differential marker expression in response to Nodal/Activin concentration were not only observed in our Nodal explant model but also validated in a human embryonic stem cell system (Figure 5K). These findings suggest that Nodal signaling levels play a conserved role in determining the cell fate specification, which influences the differentiation towards either Endo or PP lineages. To further validate this finding, we injected different dosages of Nodal mRNA into one zebrafish animal pole blastomere at the 128-cell stage, and performed HCR co-staining for *sox32* and *frzb* at 6 hpf (Figure S9A). By quantifying the cell numbers of Endo and the area sizes of PP cluster, we found that moderate concentration of Nodal levels promoted Endo cell specification (Figures S9B-S9E).

Based on our aforementioned findings, it becomes evident that subtle variation in Nodal signaling levels may regulate PP and Endo gene expression. This variation is likely regulated by Nodal-Lefty regulatory loops, and contributes to the establishment of distinct chromatin states and transcriptional response of key transcriptional regulators within these two lineages.

Gsc and Ripply1 collaborate to suppress the specification of anterior Endo in zebrafish.

It has been reported that endoderm differentiation could be suppressed by upregulation of *gsc*[18], while the role of *ripply1* in endoderm development remains unknown, though it has been characterized as a transcriptional repressor [46,47]. Moreover, both Gsc and Ripply1 are capable of inducing a secondary axis formation when overexpressed in the ventral side of zebrafish

embryos[19,22]. Combining these findings from literature with our observation of high expression

of *ripply1* in PP of Nodal explants and zebrafish embryos at 6 hpf (Figures 5D and S8I), we

hypothesized that Ripply1 may also play a role in suppressing Endo specification. To test this

hypothesis, we generated mutant lines for *gsc* and *ripply1* in zebrafish (Figure 6B). In order to

comprehensively analyze the regulatory role of Ripply1 in cooperation with Gsc in the

specification of cell fates between PP and Endo cells, we conducted a self-cross of the

Tg(*gsc*$^{+/-}$;*ripply1*$^{+/-}$) lines (Figure 6A). Subsequently, we collected the descendant embryos to

perform HCR co-staining for *frzb* and *sox32,* along with genotyping (Figure 6A). From the

descendant embryos, we obtained a total of 9 genotypes with different phenotypes displaying

variable defects in PP and anterior Endo separation (Figures 6C and S10A). Interestingly, we

observed an increase in the number of anterior Endo cells with the loss of wild-type alleles of

gsc and *ripply1* (Figures 6D, S10B, and S10C). To further investigate the mechanisms of

Ripply1 suppressing anterior Endo specification in zebrafish, we injected *ripply1* mRNA at 1-

cell stage in zebrafish embryos and found that Endo cell number was dramatically decreased as

shown by whole-mount *in situ* hybridization (WISH) of *sox32* (Figure 6F). Additionally, we

constructed a plasmid that contained *ripply1* cDNA downstream of the *sox17* promoter.

Injection of this plasmid at 1-cell stage also resulted in a decrease in Endo specification shown

by EGFP expression (Figure 6E). Taken together, these results demonstrated that Ripply1 acted

as an Endo repressor in zebrafish.

It is well established that Gsc functions as an Endo suppressor by directly binding to the

promoter region of *sox17*[18]. However, the mechanism by which Ripply1 suppresses Endo

specification remains unknown. To address this question, we injected HA-tagged *ripply1*

mRNA into zebrafish embryos for conducting CUT&Tag experiments (Figures 6G, S11A and

S11B). The CUT&Tag sequencing dataset was analysed following the standard process

(Figures S11C-S11H), and 72,683 peaks were enriched in HA-*ripply1* injected samples (Figure

6H). Genes nearby the differentially enriched peaks were used to perform GO enrichment

analysis (Figure 6I). Notably, we observed significant enrichment of terms related to "muscle

structure development", "skeletal system development" and "connective tissue development"

(Figure 6I), which were consistent with the well-established functions of Ripply1 reported in

the published studies[46-48]. Additionally, we also observed significant enrichment of GO terms

such as "regulation of cell differentiation", "negative regulation of signaling" and "negative

regulation of transcription by RNA polymerase II" (Figure 6I), further supporting the

transcriptional repressive function of Ripply1. More interestingly, Ripply1 binding peaks were

found to be enriched in the immediate upstream region of the gene body of *sox17* and *sox32*

(Figures 6J and 6K), suggesting a direct function of Ripply1 in the transcriptional regulation of

these two endodermal markers.

In summary, our data proposes a model within PP and anterior Endo lineage separation.

Variations of Nodal activities promote the establishment of distinct chromatin states in PP and

anterior Endo lineages, which is facilitated by SWI/SNF complexes. These differential

chromatin accessibility patterns, in turn, drive the differential expression of key regulators such

as *gsc* and *ripply1* in PP and Endo lineages. Collectively, these regulators collaborate to finely

regulate the segregation of mesendoderm cell fates (Figure 7).

Discussion:

In this study, we delved into the intricacies of specifying and segregating PP and anterior Endo

lineages in zebrafish through single-cell multi-omics and live imaging analyses. Our findings

revealed the origin of anterior Endo from PP progenitors and unveiled a correlation between

subtle differences of Nodal signaling levels and lineage commitment. These signaling

variations were identified as potential regulators influencing the expression of cell-fate-specific

genes at both epigenetic and transcriptional levels, with the collaboration of SWI/SNF

complexes. Through gain-/loss-of-function studies, we demonstrated the collaborative

suppression of Gsc and Ripply1 in anterior Endo specification.

While the induction of mesendodermal cell fates has been extensively studied[9,12,49,50], the

mechanisms underlying the separation of meso- and endo-lineages from common progenitors

remain a subject of ongoing research. The widely accepted notion posits that high Nodal

signaling levels promote endoderm specification, while lower levels induce mesoderm[11-13].

These quantitative effects of Nodal signaling lead to differential activation of transcriptional

targets which have differential sensitivity to Nodal activation[2,38]. However, recent studies

challenge this perspective, proposing that Nodal signal level may not strictly determine cell fate

separation of mesendoderm. Instead, Nodal signaling provides competency for mesendodermal

progenitors to stochastically switch and commit to lineage-specific fates in zebrafish[16].

Moreover, the duration of Nodal signaling has been implicated in specifying different cell fates

of mesendoderm, with prolonged signaling promoting prechordal plate specification and

suppressing endoderm induction[18]. Besides, other important signaling pathways in early

embryonic development also interact with Nodal signaling to regulate cell fate separation[13,32].

It has been well established that Fgf signaling plays a crucial role in regulating cell fate

switching between endoderm and mesoderm in lateral margin of zebrafish embryo. In our

current work, we preliminarily explored whether Fgf siganling plays roles in regulating cell

fate segregation between PP and anterior Endo cells (Figure S12). We observed that several

genes related to Fgf siganling were highly enriched in PP and Endo cells (Figure S12B), and

inhibition of Fgf signaling could obviously increase the cell number of anterior Endo (Figures

S12A and S12C). The findings above suggest that Fgf signaling inhibits the specification of

anterior Endo from PP progenitors in zebrafish, which is a similar process to the suppression

observed during lateral endoderm specification. However, further investigation is required to

understand the specific interactions between Fgf signaling[51] and factors such as Gsc, Ripply1,

Sox32[52,53] and the SWI/SNF complexes, as well as their roles in regulating the fate

diversification of PP and Endo lineages.

Our study contributed a new layer of understanding to the regulation of mesendodermal

cell fate segregation by Nodal signaling. We proposed that Nodal signaling may influence

chromatin remodeling factors, thereby affecting the chromatin accessibility states of key

regulators such as *gsc* and *ripply1*. It is worth noting that *gsc* and *ripply1* have been identified

as direct targets of Nodal signaling[22]. This suggests that the transcription of these genes may

directly and sensitively respond to variations in Nodal signaling levels. Based on this

understanding, we hypothesize that Nodal signaling can achieve more precise and robust

control over the fate segregation of closely related cell types by introducing an additional layer

of epigenetic regulatory mechanisms. Additionally, our findings underscored the pivotal role of

chromatin openness in morphogen interpretation during development, emphasizing the need to

consider chromatin structure when studying how morphogen gradients regulate patterning

formation across different species.

In our work, combining single-cell multi-omics analyses with perturbation experiments,

we revealed that chromatin remodeling together with Nodal morphogen promoted the PP and

Endo separation in zebrafish. However, which and how chromatin remodeling factors respond

to Nodal siganling to facilitate fate diversification remains largely unknown. Our current study

identified a key member of SWI/SNF complexes, *srcap*, which is differentially expressed in PP

and Endo cells. What triggers the transcriptional difference of *srcap* in these two lineages is

still a open question. One of the hypotheses is that the transcription of *srcap* is very sensitive

to Nodal signaling (Nodal concentration or Nodal duration), and a little variance in Nodal

signaling can drive differential activation of *srcap*. Another speculation is that the activation of

srcap needs the assistance of pioneer factors, like forkhead family[41,44], whose binding motifs

are particularly accessible in PP cells compared to Endo cells (Figures S8K and S8L). All these

hypotheses need to be further determined in future works.

Despite these insights, several questions remain unanswered. Firstly, we discovered a

potential role of Nodal activity levels in the separation of PP and Endo cells. However, further

investigation is required to understand how Nodal activity levels and signaling duration are

integrated to regulate this process. Secondly, although our study determined the cooperative

role of Gsc and Ripply1 in suppressing endoderm specification, the potential involvement of

other unidentified transcriptional repressors, as suggested in previous work[18], warrants further

investigation. Notably, even in double homozygous mutants of *gsc* and *ripply1*, the

specification of PP clusters was still detected, and the enrichment of another transcriptional

repressor, *osr1*, in PP cells (Figure S8J) suggested a complex network of cooperative endoderm

repressors in preventing cells from differentiating into Endo. Thus, it would be worth studying

how Gsc, Ripply1, Osr1 and other potential factors cooperatively regulate PP and Endo

specification in the future. Lastly, the anterior endoderm and ventral-lateral endoderm were

mixed together in our single-cell ATAC-seq dataset, which brought in interferences when we

interpreted the mechanisms of the separation between PP and anterior Endo. More replicates of

embryonic single-cell multi-omics or performing single-cell multi-omics on Nodal explants

may help to address this issue in future studies.

Methods:

Animal ethics

Zebrafish strains were conducted following standard procedures, and experimental procedures

were approved by the Institutional Review Board of Zhejiang University. The protocol number

is ZJU20220375.

Generation of Nodal explants

Nodal explant was generated as our previous work described[54]. 10 pg *ndr2* mRNA was injected into

one blastomere of zebrafish embryonic animal pole at the 128-cell stage. About half of the animal

pole region of the embryo was cut off from the blastula at the 1k-cell stage, and was cultured in a

Petri dish coated with 1.5% agarose gel filled with Dulbecco's modified Eagle's medium/nutrient

mixture F-12 (DMEM/F-12) with 10 mM HEPES, 1x minimum essential medium (MEM)

containing non-essential amino acids and supplemented with 7 mM $CaCl_2$, 1 mM sodium pyruvate,

50 μg/ml gentamycin, 100 μM 2-mercaptoethanol, $1\times$ antibiotic-antimycotic (15240062, Thermo

Fisher) and 10% serum replacement. For generating *ndr1*-morphant Nodal explants, *ndr1*

morpholino (0.5 mM, 2 nL) was injected into yolk of wild-type zebrafish embryos at 1-cell stage;

while for generating *lft1*-mutant Nodal explants, the embryos of *lft1* mutants were used to construct

Nodal explants. The sequence of *ndr1* morpholino was 5' - ATGTCAAATCAAGGTAATAATCCAC

- 3'.

Live imaging

Tg(*gsc*:EGFP;*sox17*:dsRed) embryos and Nodal explants generated from those embryos were

used to perform live imaging analysis. 0.5% low-melt agarose was used to mount the embryos

or Nodal explants in glass-bottom dishes (Cellvis, D35-20-0-N). The embryos or explants were

live-imaged by confocal laser scanning microscopy (OLYMPUS FV12-IXCOV) using a 20X

objective lens or a 40X oil immersion lens.

***In situ* hybridization chain reaction (HCR)**

HCR co-staining of *frzb*, *sox32* and *gsc* was performed as previous studies described[21]. The

probes of those genes were ordered and produced from Molecular Technologies

(http://www.moleculartechnologies.org/). Embryos were fixed in DEPC-treated PBS with

4%(w/v) paraformaldehyde at 4°C overnight. The embryos were incubated in 500 μL 30%

formamide probe hybridization with 2 pmol of each HCR probe set at 37°C overnight. Next

day, 30% formamide probe wash buffer was used to stop hybridization by repeated washing of

the embryos at 37°C. Fluorescent signals were generated and amplified by probes that bound

to fluorescent HCR amplifiers in an amplification buffer overnight at room temperature. To halt

this process, the samples were washed several times using 5× SSC with 0.001% Tween 20. The

reaction buffers, fluorescent hairpins and probes were manufactured by Molecular

Technologies. OLYMPUS FV12-IXCOV confocal microscope was used to photograph those

HCR co-staining samples.

Double-color RNA-fluorescence *in situ* hybridization (FISH)

Double-color FISH of *gsc* and *sox17* was conducted according to the protocol from manufactory

(http://www.pinpoease.com/, GD Pinpoease Biotech Co., Ltd.). Wild-type embryos or Nodal

explants (6 hpf, 7 hpf and 8 hpf) were fixed in DEPC-treated PBS with 4%(w/v) paraformaldehyde

at 4°C overnight. Pre-A solution was used to inhibit the peroxidase activity. Target RNA molecules

were exposed through protease treatment, followed by hybridization with probes at 4°C for 2 hours.

The fluorescent signal was amplified through sequential reactions 1, 2, and 3. Lastly, a tyramide

fluorescent substrate (OpalTM520, Akoya Biosciences) was used to incubate the embryos, enabling

fluorescent labeling of the target RNA through the Tyramide Signal Amplification (TSA) assay.

Immunostaining and quantification of pSmad2

Samples were fixed at 4% paraformaldehyde overnight for pSmad2/3 immunostaining. The

experiment was performed as previously described[54]. Confocal laser scanning microscopy

(OLYMPUS FV12-IXCOV) was used to scan the immunostained samples. Imaris (Version 9.7)

software was used to quantify the signaling activity of pSmad2 (488 nm), RFP (561 nm) and DAPI

(405 nm). Firstly, the raw image data was loaded in Imaris and visualized in a three-dimensional

(3D) view. The Spot plugin from Imaris was used to detect spots with pSmad2 signal. The diameter

of spot detection was set to 5-6 μm for GFP channel. The spots were automatically identified with

default parameters. And then, as the identification of some spots was inaccurate, we manually added

or removed some spots. Lastly, the coordinates and signal intensities of each spot were exported for

next-step analyses. To make all images comparable, we used DAPI signal as an internal control to

normalize the signal intensity in other channels. When plotting a pSmad2 signal intensity gradient

along a distance, the cell with the highest pSmad2 signal was assigned as the start point, and then

the distance of other cells was calculated by comparing them with the start point cell. The fitting

curve was plotted by ggplot2[55] in R (https://www.r-project.org).

Whole-mount *in situ* hybridization (WISH)

The processes of WISH and the construction of *sox32* probes were described in our previous work[54].

Embryos were fixed in DEPC-treated PBS with 4%(w/v) paraformaldehyde at 4°C overnight. To

enable long-term storage, the embryos were dehydrated using 100% methanol. For WISH

experiments, probes were labeled with digoxigenin-11-UTP (Roche Diagnostics), and the substrate

was NBT/BCIP.

Activation of Activin/Nodal signaling in human embryonic stem cell system

Human embryonic stem cell (hESC) lines were used between passage 45-50. Cells were seeded

equally on matrigel-coated plate and cultured in mTeSR medium. Before the experiment, each group

was pretreated in N2B27 medium for 6 hours. Then the control group was maintained in N2B27,

while the experiment groups were cultured in N2B27 medium supplemented with different

concentrations of Activn-A (0, 10, 40, 70 and 100 ng/mL). The cellular RNA of each group was

extracted with Trizol after 12 hours of treatment.

Overexpression of *gsc* and *ripply1* in zebrafish embryos

The mRNA of *gsc* and *ripply1* (synthesized *in vitro*) and *sox17* promotor-driven *ripply1* plasmid

were co-injected into zebrafish embryos at 1-cell stage. The ddH$_2$O was injected as control. The

embryos were imaged or fixed at 8 hpf. The *sox32* probe was used for WISH.

Generating heterozygous embryos of *gsc* and *ripply1*

The *ripply1* mutants were obtained from Dr. Ming Shao's lab at Shandong University. We

knockout *gsc* in *ripply1*-mutant embryos by CRISPR-Cas9 system. F0 embryos were raised up,

and then crossed with wild-type (AB). The descendant embryos (F1) were raised up and

genotyped. The adult zebrafish of the target genotype were selected.

Quantifying the cell number of anterior Endo

Imaris (version 9.7) was used to identify the Endo cells based on the HCR images. Briefly, the raw

image data was imported into Imaris, and the object was visualized in a 3D view. The Spot plugin

from Imaris was used to detect each cell. The diameter of spot detection was set to 10-12 μm for

GFP channel (staining for *sox32* or *sox17*). The candidate spots were selected by setting an

appropriate quality threshold. Finally, the coordinates and signal intensities of each spot were

exported from Imaris and used for further analysis.

CUT&Tag sample preparation

HA-*ripply1* sequence with homologous sequences was generated through two PCR steps using

two separate sets of primers (Table S1, reverse primer was common, *ripply1*-F1 primer:

tcaaggcctctcgagcctctagaATGTACCCATACGATGTTCCAGATTACGCT; *ripply1*-F2 primer:

ACGATGTTCCAGATTACGCTGGCAGCATGAATTCTGTGTGCTTTGCCACT; *ripply1*-R

primer: TACGACTCACTATAGTTCTAGAtcagttgaaagctgtgaagtga). The details of generating

mRNA *in vitro* was described in our previous protocol[21]. Briefly, the HA-*ripply1* PCR products

were inserted into pCS^{2+} plasmid through homologous recombination using ClonExpress II One

Step Cloning Kit. The plasmid containing HA-*ripply1* was linearized using *Not1* cleavage, and

HA-*ripply1* mRNA were generated through mRNA *in vitro* synthesis using mMESSAGE

mMACHINETM SP6 transcription kit. Wild-type embryos were injected with HA-*ripply1*

mRNA or ddH$_2$O at one-cell stage and harvested at 6 hpf. Yolk was removed through pipetting,

and the cells were collected. The cell viability was assessed by Trypan blue staining and the

cell number was then counted.

CUT&Tag experiment

CUT&Tag was performed on samples of wildtype, HA-*ripply1*-injected and HA-ripply1-

injected without primary antibody using <u>Hyperactive Universal CUT&Tag Assay Kit for</u>

<u>Illumina Pro</u> (Vazyme, TD904), and each sample had two replicates. Briefly, around 100,000

cells were collected and immobilized on Concanavalin A-coat Magnetic Beads. Cells were then

incubated with primary antibody (HA-Tag Rabbit mAb, Cell Signaling Technology, 3724S,

1:50 dilution) at 4°C for 12 h-16 h. Tethered cells were washed thoroughly to remove excessive

primary antibody and incubated with secondary antibody (Goat Anti-Rabbit IgG H&L, Vazyme,

Ab207-01, 1:100 dilution) for 60 minutes at room temperature. Then samples were washed

thoroughly to remove unbound secondary antibody and incubated with pA/G-Tnp for 1 hour at

room temperature. DNA was then fragmentated through Trueprep Tagment Buffer L (TTBL).

Fragmented DNA was extracted using DNA extract beads and amplified through PCR. PCR

products were purified using VAHTS DNA CLEAN Beads (Vazyme, N411) and then sent to

Novogene Co., Ltd. (Beijing, China) for sequencing. Paired-end sequencing at 150 bp read

length was performed on an NovaSeq 6000 System.

CUT&Tag analysis

Sequencing data of CUT&Tag samples were first treated by Cutadapt v4.5[56] to remove adapters

and trim reads using the following parameters "*-m 18 -q 30,30 --max-n=0.05 -e 0.2 -n 2*". The

trimmed reads were then aligned to zebrafish genome (danRer11) through Bowtie2 v2.5.2[57]

using the following parameters "*--end-to-end --very-sensitive --no-mixed --no-discordant --*

phred33 -I 10 -X 700". Samtools v1.18[58] and BEDtools v2.31.0[59] were then used for post-

alignment processing to covert file formats in preparation for the peak calling and visualization.

BigWig tracks were generated for visualization of chromatin landscapes in interested regions

using the Integrative Genomics Viewer (IGV)[60]. To assess the reproducibility between

replicates, the genome was spilt into 500 bp bins, and a Pearson correlation of the $\log_2$-

transformed read counts in each bin was calculated between datasets.

SEACR[61] was used to call "non stringent" peaks for wild-type and HA-*ripply1*-injected samples

using merged HA-*ripply1*-injected samples without primary antibody as control. Additionally,

top 1% of enrich regions were selected as peaks by AUC without controls. Peaks were annotated

583 by  ChIPpeakAnno  v3.34.1[62]  using  UCSC  annotation  from

584 TxDb.Drerio.UCSC.danRer11.refGene v3.4.6[63]. Significantly enriched peaks in HA-*ripply1*-

585 injected samples were identified using DESeq2 v1.40.2[64], and enriched GO terms were

586 identified using clusterProfiler v4.9.0.2[65]. Signal intensity heatmaps were generated using

587 deeptools v3.5.4[66]. The peaks of HA-*ripply1*-injected rep1 were used as the reference to

588 generate  computeMatrix  using  the  *reference-point*  model  (*--skipZeros --*

589 *afterRegionStartLength 3000 --beforeRegionStartLength 3000 --referencePoint center*), and

590 *PlotHeatmap* function was used to plot signal intensity.

592 **Processing and analyzing single-cell RNA-seq data**

593 Illumina sequencing reads were aligned to the reference genome (GRCz11) of zebrafish through

594 10× Genomics CellRanger pipeline (version 3.0.2) with default parameters. Nodal explants

595 generated from *lft1* mutants and *ndr1* morphants yielded 5,419 and 5,195 cells respectively. The

596 expression matrix of each sample was obtained via CellRanger pipeline analysis and used for further

597 analysis. The single-cell RNA-seq data of wild-type Nodal explants was reused from our previous

598 study[54].

600 **Processing and analyzing single-cell multi-omics data**

601 Illumina sequencing reads were aligned to the reference genome (GRCz11) of zebrafish using 10×

602 Genomics CellRanger-arc pipeline (version 2.0.0) with default parameters. Expression matrix and

603 fragments files were obtained after running the pipeline. 4,879 cells were obtained. Signac[67] was

604 used for downstream analyses with default parameters. Low-quality cells were filtered out with

605 nCount_ATAC < 100000 & nCount_RNA < 50000 & nCount_ATAC > 1000 & nCount_RNA >

606 1000 & nucleosome_signal < 2 & TSS.enrichment > 1 & percent.mt < 4. And then, gene expression

607 data and DNA accessibility data were processed respectively by Signac with default parameters.

608 The RNA expression level was utilized to identify cluster markers for cell annotation. Gene activity

609 matrix was obtained by assessing the chromatin accessibility associated with each gene through

610 *GeneActivity* function.

612 **Differential expression analysis of PP cells between wild-type, *lft1*-mutant and *ndr1*-morphant**

613 **Nodal explants**

The three sing-cell RNA-seq datasets of Nodal explants constructed from wild-type embryos, *lft1*

mutants and *ndr1* morphants were integrated by Seurat[68]. To investigate the functional and

regulatory effects of Nodal signals on cell differentiation, we compared the wild-type mesoderm PP

cells to the mesoderm PP cells in *lft1* mutants and *ndr1* morphants respectively, and differentially-

expressed (DE) genes were identified through Seurat v4.0.2 (https://github.com/satijalab/seurat)

with a Bonferroni adjusted p-value < 0.05. As it is known that EVL is not affected by Nodal

signaling[54,69], DE genes identified in EVL were regarded as background noises and were removed

from the mesoderm PP DE genes. The mesoderm PP DE genes positively correlated with Nodal

concentration were then selected, constructing two gene sets: upregulated DE genes in *lft1* mutants

and downregulated DE genes in *ndr1* morphants. GO enrichment analysis was performed on these

two   gene   sets   respectively   using   clusterProfilter   v4.2.2   (https://github.com/YuLab-

SMU/clusterProfiler). The enriched GO terms in "biological process" subontology were identified

using a Benjamini-Hochberg adjusted p-value < 0.05.

Cell-cell communication analysis by LIANA

R package, LIANA[31] v0.1.5 (https://github.com/saezlab/liana/), was used for the investigation of

ligand-receptor interactions among different cell types. The ligand-receptor analysis was performed

against the ligand-receptor reference achieved from Omnipath resource using 5 different methods

implemented in LIANA, including Natmi, Connectome, SingleCellSingalR, iTALK and

CellPhoneDBv2. The consensus rank aggregated from the results of these methods was used for the

identification of enriched ligand-receptor interactions.

Inferring cell-cell communication by CellChat

CellChat[70] was employed to systematically analyze cell-cell communication based on prior

known zebrafish ligand-receptor interaction database CellChatDB. As some known interactions

were missing in the database, we manually added some interactions of interest into the database.

The expression data of scRNA-seq was preprocessed to identify over-expressed ligands and

receptors for each group. CellChat inferred communication probability between two interacting

cell groups based on the average gene expression of a ligand in one cell group and the average

gene expression of a receptor in another cell group. The communication probabilities of

signaling pathways were then calculated by summarizing the communication probabilities of

all ligands-receptors interactions associated with each pathway.

Gene set enrichment analysis

The wild-type mesoderm PP gene expression was compared to the mesoderm PP gene expression

of *lft1* mutants and *ndr1* morphants respectively using Seurat v4.0.2

(https://github.com/satijalab/seurat). The genes expressed in more than 10% cells in either of the

two compared populations were retained. A ranked list was formed on the retained genes using

sign($\log_2$FC) * (-$\log_{10}$PValue) as the ranking statistic. The mapping between GO terms and zebrafish

genes was achieved through org.Dr.eg.db v3.14.0 (https://bioconductor.org/packages/org.Dr.eg.db/).

GSEA was performed on the selected GO terms by clusterProfiler v4.2.2

(https://github.com/YuLab-SMU/clusterProfiler), using fgsea (https://github.com/ctlab/fgsea) with

10^5 interactions. R package, enrichplot v1.14.2 (https://github.com/YuLab-SMU/enrichplot), was

then used to visualize the GSEA results.

Single-cell RNA-seq trajectory analysis

Three different methods were used to perform single-cell RNA-seq trajectory analysis. The

developmental trajectory tree of zebrafish embryos and Nodal explants was constructed by URD[71]

with default parameters as described in previous studies[71]. The branchpoint preference plot of PP

and Endo was constructed by URD with default parameters. Firstly, *branchpointPreferenceLayout*

function was used to define the preference layout for the branchpoint. And then stage, pseudotime

information and gene expression were plotted on the branchpoint using *plotBranchpoint* function.

The expression matrix of PP and Endo cells was also used to construct trajectory tree by monocle2[28]

and monocle3[72] with default parameters. And differential expression analysis was performed at the

branching point of PP and Endo separation on trajectory tree constructed by *BEAM* function in

monocle2. Modules of co-regulated genes along PP or Endo differentiation lineage were identified

by *find_gene_modules* function in monocle3 with default parameters. The cells contributed to the

preference layout for the Endo and PP brachpoint from URD analysis were also selected as inputs

for Palantir[27]. We used Harmony[73] to calculate the augmented affinity matrix for data of all time

points. Data was visualized using force directed layouts. Palantir was used for downstream analysis,

and a *nanog^{high}* cell was used as the start cell. A *frzb^{high}* cell (PP) and a *sox17^{high}* cell (Endo) were

selected as the terminal cells. Endo and PP cell differential trajectory was detected by

palantir.core.run_palantir function from Palantir. And finally, gene expression trend along Palantir

inferred pseudotime was calculated by *palantir.presults.compute_gene_trends* function.

Calculating Nodal score in single-cell RNA-seq data

Nodal functions through activating a set of key downstream genes in a concentration-dependent

manner. Our previous study identified 105 Nodal immediate target genes by analyzing bulk RNA-

seq datasets of Nodal mRNA injected explants[22]. We overlapped these 105 genes with 61 Nodal

direct targets, which were previously identified through ChIP-seq analysis of pSmad2[38]. This

analysis allowed us to pinpoint 29 Nodal downstream genes whose expression was highly sensitive

to Nodal activity. To quantify Nodal activity, we defined a Nodal score based on its downstream

transcriptional response. This score was calculated by regressing out the expression levels of these

29 Nodal downstream genes. To accomplish this, the *AddModuleScore* function in Seurat was

employed.

Data availability

The data generated in this study is deposited in the Gene Expression Omnibus (GEO). The single-

cell RNA sequencing and single-cell multi-omics data is under accession number GSE223636, and

the CUT&Tag data is under the accession number GSE249292.

Code availability

Any custom code and data are available from the authors upon request. All analyses are based on

previously published code and software.

Additional information

Supplemental Information: Supplementary Figures 1-12, Supplementary Table S1, Supplementary

Data 1-8 and Supplementary Movies 1-3.

Refference:

1.   Schier, A.F., and Talbot, W.S. (2005). Molecular genetics of axis formation in zebrafish.
Annual review of genetics *39*, 561-613. 10.1146/annurev.genet.37.110801.143752.

2.   Jones, W.D., and Mullins, M.C. (2022). Cell signaling pathways controlling an axis
organizing center in the zebrafish. Current topics in developmental biology *150*, 149-209.
10.1016/bs.ctdb.2022.03.005.

3.   Gurdon, J.B., and Bourillot, P.Y. (2001). Morphogen gradient interpretation. Nature *413*,
797-803. 10.1038/35101500.

4.   Xu, P.F., Houssin, N., Ferri-Lagneau, K.F., Thisse, B., and Thisse, C. (2014). Construction of
a vertebrate embryo from two opposing morphogen gradients. Science (New York, N.Y.)
344, 87-89. 10.1126/science.1248252.

5.   Xu, P.F., Borges, R.M., Fillatre, J., de Oliveira-Melo, M., Cheng, T., Thisse, B., and Thisse, C.
(2021). Construction of a mammalian embryo model from stem cells organized by a
morphogen signalling centre. Nature communications *12*, 3277. 10.1038/s41467-021-
23653-4.

6.   Chen, Y., and Schier, A.F. (2001). The zebrafish Nodal signal Squint functions as a
morphogen. Nature *411*, 607-610. 10.1038/35079121.

7.   Schier, A.F. (2009). Nodal morphogens. Cold Spring Harbor perspectives in biology *1*,
a003459. 10.1101/cshperspect.a003459.

8.   Hill, C.S. (2018). Spatial and temporal control of NODAL signaling. Current opinion in cell
biology *51*, 50-57. 10.1016/j.ceb.2017.10.005.

9.   Schier, A.F. (2003). Nodal signaling in vertebrate development. Annual review of cell and
developmental biology *19*, 589-621. 10.1146/annurev.cellbio.19.041603.094522.

10.   Feldman, B., Gates, M.A., Egan, E.S., Dougan, S.T., Rennebeck, G., Sirotkin, H.I., Schier, A.F.,
and Talbot, W.S. (1998). Zebrafish organizer development and germ-layer formation
require nodal-related signals. Nature *395*, 181-185. 10.1038/26013.

11.   Thisse, B., Wright, C.V., and Thisse, C. (2000). Activin- and Nodal-related factors control
antero-posterior patterning of the zebrafish embryo. Nature *403*, 425-428.
10.1038/35000200.

12.   Dougan, S.T., Warga, R.M., Kane, D.A., Schier, A.F., and Talbot, W.S. (2003). The role of the
zebrafish nodal-related genes squint and cyclops in patterning of mesendoderm.
Development (Cambridge, England) *130*, 1837-1851. 10.1242/dev.00400.

13.   van Boxtel, A.L., Economou, A.D., Heliot, C., and Hill, C.S. (2018). Long-Range Signaling
Activation and Local Inhibition Separate the Mesoderm and Endoderm Lineages.
Developmental cell *44*, 179-191.e175. 10.1016/j.devcel.2017.11.021.

14.   Martyn, I., Kanno, T.Y., Ruzo, A., Siggia, E.D., and Brivanlou, A.H. (2018). Self-organization
of a human organizer by combined Wnt and Nodal signalling. Nature *558*, 132-135.
10.1038/s41586-018-0150-y.

15.   Poulain, M., Fürthauer, M., Thisse, B., Thisse, C., and Lepage, T. (2006). Zebrafish endoderm
formation is regulated by combinatorial Nodal, FGF and BMP signalling. Development
(Cambridge, England) *133*, 2189-2200. 10.1242/dev.02387.

16.   Economou, A.D., Guglielmi, L., East, P., and Hill, C.S. (2022). Nodal signaling establishes a
competency window for stochastic cell fate switching. Developmental cell *57*, 2604-2622
e2605. 10.1016/j.devcel.2022.11.008.

17.   Barone, V., Lang, M., Krens, S.F.G., Pradhan, S.J., Shamipour, S., Sako, K., Sikora, M., Guet,
C.C., and Heisenberg, C.P. (2017). An Effective Feedback Loop between Cell-Cell Contact
Duration and Morphogen Signaling Determines Cell Fate. Developmental cell *43*, 198-
211.e112. 10.1016/j.devcel.2017.09.014.

18.   Sako, K., Pradhan, S.J., Barone, V., Inglés-Prieto, Á., Müller, P., Ruprecht, V., Čapek, D.,
Galande, S., Janovjak, H., and Heisenberg, C.P. (2016). Optogenetic Control of Nodal
Signaling Reveals a Temporal Pattern of Nodal Signaling Regulating Cell Fate Specification
during Gastrulation. Cell reports *16*, 866-877. 10.1016/j.celrep.2016.06.036.

19.   Dixon Fox, M., and Bruce, A.E. (2009). Short- and long-range functions of Goosecoid in
zebrafish axis formation are independent of Chordin, Noggin 1 and Follistatin-like 1b.
Development (Cambridge, England) *136*, 1675-1685. 10.1242/dev.031161.

20.   Seiliez, I., Thisse, B., and Thisse, C. (2006). FoxA3 and goosecoid promote anterior neural
fate through inhibition of Wnt8a activity before the onset of gastrulation. Developmental
biology *290*, 152-163. 10.1016/j.ydbio.2005.11.021.

21.   Cheng, T., Xing, Y.Y., Dong, Y., and Xu, P.F. (2023). Protocol for generation and assessment
of head-like structure in zebrafish. STAR Protoc *4*, 102553. 10.1016/j.xpro.2023.102553.

22.   Cheng, T., Xing, Y.Y., Liu, C., Li, Y.F., Huang, Y., Liu, X., Zhang, Y.J., Zhao, G.Q., Dong, Y., Fu,
X.X., et al. (2023). Nodal coordinates the anterior-posterior patterning of germ layers and
induces head formation in zebrafish explants. Cell reports *42*, 112351.
10.1016/j.celrep.2023.112351.

23.   Warga, R.M., and Nüsslein-Volhard, C. (1999). Origin and development of the zebrafish
endoderm. Development (Cambridge, England) *126*, 827-838.

24.   Wagner, D.E., Weinreb, C., and Collins, Z.M. (2018). Single-cell mapping of gene
expression landscapes and lineage in the zebrafish embryo. Science (New York, N.Y.) *360*,
981-987. 10.1126/science.aar4362.

25.   Farrell, J.A., and Wang, Y. (2018). Single-cell reconstruction of developmental trajectories
during zebrafish embryogenesis. Science (New York, N.Y.) *360*. 10.1126/science.aar3131.

26.   Becht, E., and McInnes, L. (2018). Dimensionality reduction for visualizing single-cell data
using UMAP. Nature biotechnology. 10.1038/nbt.4314.

27.   Setty, M., Kiseliovas, V., Levine, J., Gayoso, A., Mazutis, L., and Pe'er, D. (2019).
Characterization of cell fate probabilities in single-cell data with Palantir. Nature
biotechnology *37*, 451-460. 10.1038/s41587-019-0068-4.

28.   Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).
Reversed graph embedding resolves complex single-cell trajectories. Nature methods *14*,
979-982. 10.1038/nmeth.4402.

29.   Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C.,
and Chanda, S.K. (2019). Metascape provides a biologist-oriented resource for the
analysis of systems-level datasets. Nature communications *10*, 1523. 10.1038/s41467-
019-09234-6.

30.   Xia, P., Gutl, D., Zheden, V., and Heisenberg, C.P. (2019). Lateral Inhibition in Cell
Specification Mediated by Mechanical Signals Modulating TAZ Activity. Cell *176*, 1379-
1392 e1314. 10.1016/j.cell.2019.01.019.

31.   Dimitrov, D., Turei, D., Garrido-Rodriguez, M., Burmedi, P.L., Nagai, J.S., Boys, C., Ramirez
Flores, R.O., Kim, H., Szalai, B., Costa, I.G., et al. (2022). Comparison of methods and

resources for cell-cell communication inference from single-cell RNA-Seq data. Nature
communications *13*, 3224. 10.1038/s41467-022-30755-0.

32. Mathieu, J., Griffin, K., Herbomel, P., Dickmeis, T., Strahle, U., Kimelman, D., Rosa, F.M., and
Peyrieras, N. (2004). Nodal and Fgf pathways interact through a positive regulatory loop
and synergize to maintain mesodermal cell populations. Development (Cambridge,
England) *131*, 629-641. 10.1242/dev.00964.

33. Müller, P., Rogers, K.W., Jordan, B.M., Lee, J.S., Robson, D., Ramanathan, S., and Schier, A.F.
(2012). Differential diffusivity of Nodal and Lefty underlies a reaction-diffusion patterning
system. Science (New York, N.Y.) *336*, 721-724. 10.1126/science.1221920.

34. Chen, Y., and Schier, A.F. (2002). Lefty proteins are long-range inhibitors of squint-
mediated nodal signaling. Current biology : CB *12*, 2124-2128. 10.1016/s0960-
9822(02)01362-3.

35. Sekine, R., Shibata, T., and Ebisuya, M. (2018). Synthetic mammalian pattern formation
driven by differential diffusivity of Nodal and Lefty. Nature communications *9*, 5456.
10.1038/s41467-018-07847-x.

36. Kuhn, T., Landge, A.N., Morsdorf, D., Cossmann, J., Gerstenecker, J., Capek, D., Muller, P.,
and Gebhardt, J.C.M. (2022). Single-molecule tracking of Nodal and Lefty in live zebrafish
embryos supports hindered diffusion model. Nature communications *13*, 6101.
10.1038/s41467-022-33704-z.

37. Liu, L., Nemashkalo, A., Rezende, L., Jung, J.Y., Chhabra, S., Guerra, M.C., Heemskerk, I.,
and Warmflash, A. (2022). Nodal is a short-range morphogen with activity that spreads
through a relay mechanism in human gastruloids. Nature communications *13*, 497.
10.1038/s41467-022-28149-3.

38. Dubrulle, J., Jordan, B.M., Akhmetova, L., Farrell, J.A., Kim, S.H., Solnica-Krezel, L., and
Schier, A.F. (2015). Response to Nodal morphogen gradient is determined by the kinetics
of target gene induction. eLife *4*. 10.7554/eLife.05042.

39. Xing, C., Shen, W., Gong, B., Li, Y., Yan, L., and Meng, A. (2022). Maternal Factors and
Nodal Autoregulation Orchestrate Nodal Gene Expression for Embryonic Mesendoderm
Induction in the Zebrafish. Front Cell Dev Biol *10*, 887987. 10.3389/fcell.2022.887987.

40. Pinheiro, D., Kardos, R., Hannezo, E., and Heisenberg, C.P. (2022). Morphogen gradient
orchestrates pattern-preserving tissue morphogenesis via motility-driven unjamming.
Nature Physics. 10.1038/s41567-022-01787-6.

41. Coda, D.M., Gaarenstroom, T., East, P., Patel, H., Miller, D.S., Lobley, A., Matthews, N.,
Stewart, A., and Hill, C.S. (2017). Distinct modes of SMAD2 chromatin binding and
remodeling shape the transcriptional response to NODAL/Activin signaling. eLife *6*.
10.7554/eLife.22474.

42. Ross, S., Cheung, E., Petrakis, T.G., Howell, M., Kraus, W.L., and Hill, C.S. (2006). Smads
orchestrate specific histone modifications and chromatin remodeling to activate
transcription. EMBO J *25*, 4490-4502. 10.1038/sj.emboj.7601332.

43. Xiao, L., Parolia, A., Qiao, Y., Bawa, P., Eyunni, S., Mannan, R., Carson, S.E., Chang, Y., Wang,
X., Zhang, Y., et al. (2022). Targeting SWI/SNF ATPases in enhancer-addicted prostate
cancer. Nature *601*, 434-439. 10.1038/s41586-021-04246-z.

44. Coda, D.M., Patel, H., Gori, I., Gaarenstroom, T.E., Song, O.R., Howell, M., and Hill, C.S.
(2022). A network of transcription factors governs the dynamics of NODAL/Activin

transcriptional responses. Journal of cell science *135*. 10.1242/jcs.259972.

45. Akhmetova, L. (2016). Transcriptional Regulation of Nodal Target Genes in Early Zebrafish Development. Doctor (Harvard University).

46. Kawamura, A., Koshida, S., Hijikata, H., Ohbayashi, A., Kondoh, H., and Takada, S. (2005). Groucho-associated transcriptional repressor ripply1 is required for proper transition from the presomitic mesoderm to somites. Developmental cell *9*, 735-744. 10.1016/j.devcel.2005.09.021.

47. Yabe, T., Uriu, K., and Takada, S. (2023). Ripply suppresses Tbx6 to induce dynamic-to-static conversion in somite segmentation. Nature communications *14*, 2115. 10.1038/s41467-023-37745-w.

48. Windner, S.E., Doris, R.A., Ferguson, C.M., Nelson, A.C., Valentin, G., Tan, H., Oates, A.C., Wardle, F.C., and Devoto, S.H. (2015). Tbx6, Mesp-b and Ripply1 regulate the onset of skeletal myogenesis in zebrafish. Development (Cambridge, England) *142*, 1159-1168. 10.1242/dev.113431.

49. Shen, M.M. (2007). Nodal signaling: developmental roles and regulation. Development (Cambridge, England) *134*, 1023-1034. 10.1242/dev.000166.

50. Takenaga, M., Fukumoto, M., and Hori, Y. (2007). Regulated Nodal signaling promotes differentiation of the definitive endoderm and mesoderm from ES cells. Journal of cell science *120*, 2078-2090. 10.1242/jcs.004127.

51. Mizoguchi, T., Izawa, T., Kuroiwa, A., and Kikuchi, Y. (2006). Fgf signaling negatively regulates Nodal-dependent endoderm induction in zebrafish. Developmental biology *300*, 612-622. 10.1016/j.ydbio.2006.08.073.

52. Kikuchi, Y., Agathon, A., Alexander, J., Thisse, C., Waldron, S., Yelon, D., Thisse, B., and Stainier, D.Y. (2001). casanova encodes a novel Sox-related protein necessary and sufficient for early endoderm formation in zebrafish. Genes & development *15*, 1493-1505. 10.1101/gad.892301.

53. Aoki, T.O., David, N.B., Minchiotti, G., Saint-Etienne, L., Dickmeis, T., Persico, G.M., Strahle, U., Mourrain, P., and Rosa, F.M. (2002). Molecular integration of casanova in the Nodal signalling pathway controlling endoderm formation. Development (Cambridge, England) *129*, 275-286. 10.1242/dev.129.2.275.

54. Cheng, T., Xing, Y.-Y., Li, Y.-F., Liu, C., Huang, Y., Zhang, Y.-J., Megason, S.G., and Xu, P.-F. (2021). Single cell response landscape of graded Nodal signaling in zebrafish explants. bioRxiv. 10.1101/2021.04.25.441305.

55. Wickham, H. (2016). ggplot2 : Elegant Graphics for Data Analysis. Use R!,. 2nd ed. Springer International Publishing : Imprint: Springer,.

56. Kechin, A., Boyarskikh, U., Kel, A., and Filipenko, M. (2017). cutPrimers: A New Tool for Accurate Cutting of Primers from Reads of Targeted Next Generation Sequencing. J Comput Biol *24*, 1138-1143. 10.1089/cmb.2017.0096.

57. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nature methods *9*, 357-359. 10.1038/nmeth.1923.

58. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics (Oxford, England) *25*, 2078-2079. 10.1093/bioinformatics/btp352.

880   59.   Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
881   genomic features. Bioinformatics (Oxford, England) *26*, 841-842.
882   10.1093/bioinformatics/btq033.

883   60.   Robinson, J.T., Thorvaldsdottir, H., Turner, D., and Mesirov, J.P. (2023). igv.js: an
884   embeddable JavaScript implementation of the Integrative Genomics Viewer (IGV).
885   Bioinformatics (Oxford, England) *39*. 10.1093/bioinformatics/btac830.

886   61.   Meers, M.P., Tenenbaum, D., and Henikoff, S. (2019). Peak calling by Sparse Enrichment
887   Analysis for CUT&RUN chromatin profiling. Epigenetics Chromatin *12*, 42.
888   10.1186/s13072-019-0287-4.

889   62.   Zhu, L.J. (2013). Integrative analysis of ChIP-chip and ChIP-seq dataset. Methods in
890   molecular biology (Clifton, N.J.) *1067*, 105-124. 10.1007/978-1-62703-607-8_8.

891   63.   Team, B.C.M., B.P. (2019). TxDb.Drerio.UCSC.danRer11.refGene: Annotation package for
892   TxDb object(s).

893   64.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
894   dispersion for RNA-seq data with DESeq2. Genome biology *15*, 550. 10.1186/s13059-
895   014-0550-8.

896   65.   Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et al.
897   (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
898   Innovation (Camb) *2*, 100141. 10.1016/j.xinn.2021.100141.

899   66.   Ramirez, F., Ryan, D.P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dundar,
900   F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing
901   data analysis. Nucleic acids research *44*, W160-165. 10.1093/nar/gkw257.

902   67.   Stuart, T., Srivastava, A., Madad, S., Lareau, C.A., and Satija, R. (2021). Single-cell chromatin
903   state analysis with Signac. Nature methods *18*, 1333-1341. 10.1038/s41592-021-01282-
904   5.

905   68.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
906   cell transcriptomic data across different conditions, technologies, and species. Nature
907   biotechnology *36*, 411-420. 10.1038/nbt.4096.

908   69.   Sagerström, C.G., Gammill, L.S., Veale, R., and Sive, H. (2005). Specification of the
909   enveloping layer and lack of autoneuralization in zebrafish embryonic explants.
910   Developmental dynamics : an official publication of the American Association of
911   Anatomists *232*, 85-97. 10.1002/dvdy.20198.

912   70.   Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.H., Myung, P., Plikus,
913   M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using CellChat.
914   Nature communications *12*, 1088. 10.1038/s41467-021-21246-9.

915   71.   Farrell, J.A., Wang, Y., Riesenfeld, S.J., Shekhar, K., Regev, A., and Schier, A.F. (2018). Single-
916   cell reconstruction of developmental trajectories during zebrafish embryogenesis. Science
917   (New York, N.Y.) *360*. 10.1126/science.aar3131.

918   72.   Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos, S.,
919   Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional landscape of
920   mammalian organogenesis. Nature *566*, 496-502. 10.1038/s41586-019-0969-x.

921   73.   Nowotschin, S., Setty, M., Kuo, Y.Y., Liu, V., Garg, V., Sharma, R., Simon, C.S., Saiz, N.,
922   Gardner, R., Boutet, S.C., et al. (2019). The emergent landscape of the mouse gut
923   endoderm at single-cell resolution. Nature *569*, 361-367. 10.1038/s41586-019-1127-1.

74. Pijuan-Sala, B., Griffiths, J.A., Guibentif, C., Hiscock, T.W., Jawaid, W., Calero-Nieto, F.J., Mulas, C., Ibarra-Soria, X., Tyser, R.C.V., Ho, D.L.L., et al. (2019). A single-cell molecular map of mouse gastrulation and early organogenesis. Nature *566*, 490-495. 10.1038/s41586-019-0933-9.

75. Sagerstrom, C.G., Gammill, L.S., Veale, R., and Sive, H. (2005). Specification of the enveloping layer and lack of autoneuralization in zebrafish embryonic explants. Developmental dynamics : an official publication of the American Association of Anatomists *232*, 85-97. 10.1002/dvdy.20198.

Figures:

Figure 1. Single-cell trajectory analysis for studying PP and anterior Endo specification.

(A) URD differentiation tree of zebrafish embryos from blastula stage to somitogenesis stage[74]. Cells are colored according to developmental stages. Gray dotted frame indicates the ectoderm lineages and black dotted frame indicates the mesendoderm lineages. Green line shows the

pharyngeal endoderm lineage (anterior endoderm, roman number I), red line shows the

prechordal plate lineage (roman number II) and yellow line indicates their progenitors. (B)

Simplified schematic showing the patterns of endoderm and axial mesoderm of zebrafish

embryos at 75% epiboly. Dotted frame indicates the enlarged view of dorsal anterior region,

showing that anterior Endo sporadically locates near PP. (C) URD branchpoint plots of the PP

and anterior Endo development from scRNA-seq data of zebrafish embryos[74], showing

pseudotime (x-axis) and random walk visitation preference from pharyngeal endoderm to

prechordal plate domains (y-axis). Cells are colored by developmental stages (left) and

expression of Endo markers (*sox32*, *sox17* and *pax1b*; right top) and PP markers (*gsc*, *frzb* and

he1a; right bottom). (D) Force-directed layout of PP and anterior Endo cells of zebrafish

embryos from blastula stage to somitogenesis stage. Cells are colored by Palantir[27] pseudotime

(top left), differentiation potential (top right) and branch probabilities of PP (bottom left) and

anterior Endo (bottom right). (E) Branch probabilities of five cells randomly selected from

progenitor cells. Bars are colored by cell types. (F) URD differentiation tree of Nodal explants

from blastula stage to the end of gastrulation. Gray dotted frame indicates the ectoderm lineages

and black dotted frame indicates the mesendoderm lineages. The green, red and yellow dots

indicate the Endo cells (anterior Endo), PP cells and the progenitor cells respectively, and the

gray dots indicate other cells. (G-H) URD branchpoint plots of the PP and Endo development

from scRNA-seq data of Nodal explants showing pseudotime (y-axis) and random walk

visitation preference from PP to Endo domains (x-axis). Cells are colored by developmental

stages (G) and expression of Endo markers (*sox32*, *sox17* and *nkx2.7*; top) and PP markers (*gsc*,

frzb and *he1a*; bottom) (H). (I) Force-directed layout of PP and anterior Endo cells of Nodal

explants from blastula stage to the end of gastrulation. Cells are colored by the levels of

Palantir[27] pseudotime (top left), differentiation potential (top right) and branch probabilities of

PP (bottom left) and anterior Endo (bottom right).

Figure 2. Live imaging analyses for Tg(*gsc*:EGFP;*sox17*:DsRed) embryos. (A) Time series

of 3D reconstruction of a representative Tg(*gsc*:EGFP;*sox17*:DsRed) embryo. (B) Schematic

diagram indicating the focused region in the Tg(*gsc*:EGFP;*sox17*:DsRed) embryo. (C and D)

Temporal profiles of *sox17* (RFP) & *gsc* (GFP) (C) and the ratio between RFP intensity and

GFP intensity (D) in the highlighted cell from (A). (E) Double-color RNA-fluorescence *in situ*

hybridization (FISH) of *sox17* and *gsc* in wild-type embryos at 6 hpf (top) and 7.5 hpf (bottom).

(N >= 30, n = 5/5, 6 hpf; N >= 30, n = 5/5, 7.5 hpf; N: embryos were used in the experiment;

n: embryos were imaged, expression observed/total imaged) (F) Scatter plots and inferred linear

regression comparing the relative intensities of *sox17* and *gsc* in wild-type embryos at 6 hpf

(top) and 7.5 hpf (bottom). Each FISH experiment was performed for at least 3 independent

replicates (technical replicates). Scale bar: 30 μm (A) and 50 μm (E).

Figure 3 Nodal-Lefty regulatory loops are involved in cell fate separation between anterior Endo and PP. (A and B) Dot plots visualizing ligand-receptor interactions between PP and Endo. The interactions from PP (ligand) to Endo (receptor) and from Endo (ligand) to PP (receptor) are plotted respectively in A and B. (C) Schematic diagram illustrating PP (red) and Endo (green) in zebrafish embryos at 6 hpf. (D) Simplified schematic showing the interaction network between Nodal and Lefty. (E) 3D reconstruction of representative confocal-scanned images showing pSmad2 levels in zebrafish embryonic animal pole stimulated by Nodal injection at 128-cell stage (WT, N >= 10, n = 3/4; *ndr1*kd, N >= 10, n = 3/4; *lft1*ko, N >= 10, n = 4/4; N: embryos were used in the experiment; n: embryos were imaged, observed/total imaged). (F) Comparisons of pSmad2 levels in wild-type embryos, *lft1* mutants and *ndr1* morphants at 6 hpf (quantified from E). Statistical differences between two samples were evaluated by Student's t-test. *indicates P-value < 0.05, **indicates P-value < 0.01 and ***indicates P-value < 0.001. (G) Schematic diagram showing the experimental workflow of single-cell RNA sequencing of Nodal explants constructed from *lft1* mutants and *ndr1* morphants at 6 hpf. (H) Overall UMAP plot of integrated single-cell datasets of Nodal explants generated from wild-type embryos (left), *ndr1* morphants (middle) and *lft1* mutants (right). (I) Histograms showing the ratios of the cell proportions of endoderm against EVL (top) and

prechordal plate (bottom) against EVL. Per indicates percentage. As EVL cells can be specified

spontaneously and are Nodal-independent[75], here the proportions of prechordal plate and

endoderm were adjusted by using the proportions of EVL cells as an internal control. (J) Venn

plot showing the selection of Nodal direct target genes used for defining Nodal score (**see**

Methods). (K) Dot plot visualizing Nodal score in wild-type, *ndr1*-morphant and *lft1*-mutant

Nodal explants. Dot size indicates the percentage of cells within the cell group; the color

indicates the average Nodal score levels. Scale bar: 50 µm (E).

1006

1007

1008

Figure 4. Single-cell trajectory tree analyses of PP and anterior Endo reveal that chromatin organization is involved in the separation of these two lineages. (A) Pseudotime trajectory analysis of PP and anterior Endo using integrated single-cell datasets of wild-type (left), *ndr1*-morphant (middle) and *lft1*-mutant (right) Nodal explants. Cells are colored by cell types (top) and pseudotime levels (bottom). (B) Heatmaps representing clusters of genes that co-vary along the pseudotime during the separation of PP and anterior Endo from common progenitors. Top GO terms enriched by the genes in each cluster are listed with their corresponding adjusted P-values. Representative GO terms of each cluster are highlighted in different colors. (C) Diagram of gene set selection for GO analysis. As a control of batch effects,

EVL differentially expressed (DE) genes are removed from mesoderm PP DE genes. (D and E) GO enrichment analysis of mesoderm PP up-regulated genes in *lft1* mutants (D) and down-regulated genes in *ndr1* morphants (E). Redundancy of enriched GO terms are removed, and each GO term contains at least 10 transcripts. Note that only the top 16 enriched GO terms identified in *lft1* mutant are plotted for a visualization purpose. (F and G) GSEA of "chromatin organization" term on PP cells from *lft1* mutants (F) and *ndr1* morphants (G) respectively. P-adjust indicates Benjamini-Hochberg adjusted p-value. (H-M) SWI/SNF complexes involved in regulating cell fate separation between the PP and anterior Endo. (H) HCR co-staining of *gsc*, *frzb* and *sox32* in wild-type embryos (top, N >= 40, n = 5/5, N: embryos were used in the experiment, n: embryos were imaged, expression observed/total imaged), as well as those treated with 1 μM (middle, N >= 40, n = 4/5, N: embryos were used in the experiment, n: embryos were imaged, expression observed/total imaged) and 5 μM (bottom, n = 5/6, expression observed/total imaged) AU15330. Regions framed by dotted white lines were used to quantify the cell number of anterior Endo. (I) Box plot showing the cell numbers of anterior Endo in (H). (J and K) Volinplot indicating the expression levels of *srcap* in PP and Endo cells within both embryos (J) and Nodal explants (K). (L) HCR co-staining of *gsc*, *frzb* and *sox32* in wild-type embryos and embryos with *srcap* knockdown. Embryos at 7.5 hpf (Ctrl, N >= 30, n = 6/6; *srcap* MO, N >= 30, n = 4/5; N: embryos were used in the experiment; n: embryos were imaged, expression observed/total imaged) were evaluated. (M) Box plot showing the cell numbers of anterior Endo in (L). Statistical differences between two samples were evaluated by Student's t-test (I and M). *indicates P-value < 0.05; NS indicates P-value >=0.05. Each HCR experiment was performed for at least 3 independent replicates (technical replicates). Scale bar: 50 μm (H and L).

1043

Figure 5. Investigating the mechanisms of PP and Endo separation through integrative

single-cell RNA-seq and ATAC-seq analysis. (A) Schematic diagram showing the

experimental workflow of single-cell multi-omics of 6 hpf zebrafish embryos. (B) UMAPs

based on scATAC-Seq (left), scRNA-Seq (middle) and co-embedding scRNA-seq and scATAC-

seq (right) datasets. Cells are colored by different cell types. (C) UMAP plot of the PP and Endo

cells. Cells are colored by cell types (left) and the expression of *gsc* (middle) and *sox32* (right).

(D) Volcano plot showing the differentially expressed genes in PP and Endo based on RNA

levels of single-cell multi-omics datasets. (E-F) Track plot showing chromatin accessibility of

each cell type on the gene locus of *gsc* (E) and *ripply1* (F). (G) Heatmap showing the chromatin

accessibility level of genes with differential chromatin accessibility in PP and Endo. Red and

green box indicates genes with higher chromatin accessbility levels in PP and Endo cells

respectively. (H and I) Dot plot (H) and ridge plot (I) showing the level of Nodal score in PP

and Endo. (J) Box plots showing the expression profiles of *sox17* (Endo), *gsc* (PP), *ripply1* (PP)

and *krt4* (EVL) in our previous bulk RNA-seq datasets of Nodal explants. (K) Box plots (left)

and bar plot (right) showing the expression levels of *SOX17* and *GSC* in the human embryonic

stem cells (hESCs) treated with different dosages of Activin protein (**see Methods**). The values

were calculated as the mean ± SEM (K).

1061

1062

Figure 6. Exploring the mechanisms underlying the role of Ripply1 in the regulation of cell fate separation between PP and anterior Endo. (A) Schematic diagram illustrating the workflow for investigating the role of *gsc* and *ripply1* in mesendoderm cell fate separation through loss-of-function studies. (B) Schematic presentation of *gsc* and *ripply1* mutations and the sequencing results of each genotype. The mutations are a 10-bp deletion and a 1-bp insertion in *gsc* and *ripply1* coding sequences respectively. Texts in red indicate coding sequences. Sanger sequencing results display at the bottom, showcasing wild-type (top), heterozygous mutant (middle) and homozygous mutant (bottom) embryos of *gsc* and *ripply1* mutations. (C) HCR co-staining of *sox32* and *frzb* in the embryos of Tg(*gsc*[+/+];*ripply1*[+/+]) (WT),

Tg(*gsc$^{+/+}$;ripply1$^{-/-}$*) (*ripply1* mutant) and Tg(*gsc$^{-/-}$;ripply1$^{-/-}$*) (double mutant) at 8 hpf (Here

embryos of three different genotypes are shown. Other genotypes are shown in Figure S10A.)

[Tg(*gsc$^{+/+}$;ripply1$^{+/+}$*), n = 4/5; Tg(*gsc$^{+/+}$;ripply1$^{-/-}$*), n = 4/6; Tg(*gsc$^{-/-}$;ripply1$^{-/-}$*), n = 5/5;

expression observed/total imaged]. (D) Box plot showing the quantification of the numbers of

anterior Endo cells identified in (C). (E and F) Assessing the role of *ripply1* in anterior Endo

specification. EGFP driven by *sox17* promotor (E) and WISH of *sox32* (F) are used to indicate

the Endo cells. Embryos with *gsc* overexpression are used as a positive control. The statistics

of observed/total sampled embryos were shown in bottom left of the panels (E and F). (G)

Schematic diagram showing the strategy of conducting CUT&Tag experiment for *ripply1*. (H)

Metaplots and heatmaps for CUT&Tag signals over R1_rep1 binding peaks (n=72,683) in wild-

type and HA-*ripply1*-injected embryos. Two replicates were performed. (I) Dot plot showing

the enriched GO terms using the genes that annotated from the differentially enriched peaks. (J

and K) Screenshot of R1_con (technical control, without primary antibody, **see Methods**), WT

(ddH$_2$O-injected) and R1 (HA-*ripply1*-injected) CUT&Tag peaks around *sox17* (J) and *sox32*

(K) loci. Each HCR and ISH experiment was performed for at least 3 independent replicates

(technical replicates). Statistical differences between two samples were evaluated by Student's

t-test (D). Scale bar: 50 µm (C) and 200 µm (E and F).

1089

1090

Figure 7. Model underlying the separation of PP and anterior Endo. An outline depicts PP and anterior Endo specification. Nodal signaling drives a differential chromatin states between PP and anterior Endo through epigenetic regulators. And then, key transcriptional factors, such as *gsc* and *ripply1*, are differentially expressed in these two lineages and suppress anterior Endo specification by directly binding to the *cis*-elements of *sox17* and *sox32*.

Acknowledgments:

We would like to thank Dr. Alex Schier for his kind gift of the *lft1* mutant. We thank Dr. Ming Shao from Shandong University for his kind gift of the *ripply1* mutant. We thank Dr. Peng Xia from Zhejiang University for his kind gift of the Tg(*gsc*:EGFP) embryos. We also thank Dr. Nai-He Jing at University of Chinese Academy of Sciences, Dr. Jun Ma, Dr. Xiao-hang Yang, Dr. Min-Xin Guan, Dr. Feng He and the members of Laboratory of Development and Organogenesis (LDO) at Zhejiang University for helpful suggestions and discussions. We thank Mr. Guang-Xu Zhang from Biostar Technology for the technical help on constructing 10x single-cell multi-omics sequencing library. We thank Shuang-Shuang Liu from the Imaging Platform and Ying-Niang Li from the zebrafish core facility at Zhejiang University School of

1107 Medicine for their technical support. This work was supported by grants from the Chinese

1108 National Key Research and Development Project (2022YFA1103100, 2019YFA0802402) and

1109 the National Scientific Foundation of China (32050109, 32300688, 32300677).

1110

1111 **Conflict of Interest:**
1112 The authors declare no competing interests.
1113

Author Contributions

PFX, TC and HQL conceived and designed the research; TC, XL, YD, YMT, YYX, CYC, CL,

YFL, YH, DHZ, XX, XXF, ZXJ, JYL and JM performed research; TC and YD analyzed the

single-cell RNA-seq and single-cell multi-omics data; YD performed the CUT&Tag

experiment and analyzed the data; TC, YD, HQL and PFX wrote the manuscript; all authors

reviewed and approved the manuscript. PFX and HQL supervised this study. Fundings for the

study were provided by PFX, TC and YD.

1121

1122