1 # Spatial autocorrelation of the environment influences the patterns and genetics of local adaptation

4 *Tom R. Booker[1]*

5    1.  Department of Forest and Conservation Science, Faculty of Forestry, University of British
6         Columbia

8 Correspondence to: thomas.booker@ubc.ca

## Abstract

Environmental heterogeneity can lead to spatially varying selection, which can, in turn, lead to local adaptation. Population genetic models have shown that the pattern of environmental variation in space can strongly influence the evolution of local adaptation. In particular, when environmental variation is highly autocorrelated in space local adaptation will more readily evolve. Despite this long-held prediction, the evolutionary genetic consequences of different patterns of environmental variation have not been thoroughly explored. In this study, simulations are used to model local adaptation to different patterns of environmental variation. The simulations confirm that local adaptation is expected to increase with the degree of spatial autocorrelation in the selective environment, but also show that highly heterogeneous environments are more likely to exhibit high variation in local adaptation, a result not previously described. Spatial autocorrelation in the environment also influences the evolution and genetic architecture of local adaptation, with different combinations of allele frequency and effect size arising under different patterns of environmental variation. These differences influence the ability to characterise the genetic basis of local adaptation in different environments. Finally, I analyse a large-scale provenance trial conducted on lodgepole pine and find suggestive evidence that spatially autocorrelated environmental variation leads to stronger local adaptation in natural populations of lodgepole pine. Overall, this work emphasizes the profound importance that the spatial pattern of selection can have on the evolution of local adaptation and how spatial autocorrelation should be considered when formulating hypotheses in ecological and genetic studies.

## Lay Summary

Many species exhibit local adaptation to environmental variation across their ranges. Theoretical population genetics predicts that the evolution of local adaptation and patterns of genetic variation underlying it will be influenced by the spatial pattern of variation across a species' range. However, this prediction has not been thoroughly explored for cases of complex heterogeneous landscapes. In this paper, I analyse simulations and empirical data to characterise the effects that the spatial pattern of environmental variation can have on the evolution of local adaptation and the genetics underlying it. From these analyses, I show that the pattern of environmental variation influences the average level of local adaptation, variation in local adaptation as well as the genetics underlying this important phenomenon.

## Introduction

Local adaptation is an important phenomenon in the natural world. Along with phenotypic plasticity, local adaptation can dictate the extent of environmental heterogeneity that a species can tolerate, shape its geographic range (Kirkpatrick & Barton, 1997), and help predict how it will respond to changing environments (Rellstab et al., 2021). In forest trees, for example, local adaptation is widely observed (Leites & Benito Garzón, 2023) and is central to plans for adapting forestry practice in light of climate change (O'Neill & Gómez-Pineda, 2021; Ying & Yanchuk, 2006). Local adaptation can be defined as a kind of genotype-by-environment interaction for fitness, where individuals have higher chances of survival and/or reproduction when they are reared at home as opposed to away, though several other definitions are used in the literature (Blanquart et al., 2013; Kawecki & Ebert, 2004). Local adaptation is a property of a particular population at a particular point in time rather than a property of a species as a whole. For example, populations at range edges are often expected to be maladapted to their conditions, while populations in the core of a range may be well adapted (Angert et al., 2020). Locally adapted populations may harbour genetic variation that could help buffer susceptible ones against the detrimental effects of climate change (Aitken & Whitlock, 2013), which are already wreaking havoc on important species around the world (Hartmann et al., 2022). A deep understanding of local adaptation, the agents that have given rise to it and the genetics that underpin this phenomenon is thus important for our understanding of biodiversity and for species management and conservation in the Anthropocene (Aitken & Whitlock, 2013; Exposito-Alonso, 2023; Wadgymar et al., 2022).

The ultimate cause of local adaptation is variation in the environment. Whether it is biotic (e.g. disease/parasite prevalence or intraspecific competition) or abiotic (e.g. climate, geology or photoperiod), variation in the environment may lead to spatially varying selection pressures where phenotypic optima differ over a landscape. Such variation in selection across space has been well documented (e.g. Siepielski et al., 2013) and there are, of course, myriad aspects of the environment that could conceivably induce spatially varying selection, many of which would be highly inter-correlated. However, while there is an infinite number of ways to describe the environment, most of these may be functionally disconnected from a species' biology. A recent review by Wadgymer et al (2022) highlighted a critical gap in our knowledge of local adaptation - that the aspects of environmental variation that have given rise to local adaptation (what they term the 'agents of selection') are unknown in most cases. For example, in the absence of experimental evidence many genetic studies have assumed that various climatic measures recorded in databases such as WorldClim correspond to agents of selection and search for the genetic basis of local adaptation using those data (Lasky et al., 2023). However, that a particular aspect of environmental variation could conceivably induce spatially varying selection is not a guarantee that it will have led to local adaptation.

Population genetic studies have revealed numerous factors that can influence the evolution of local adaptation in a particular location, the most important being the strength of selection and rates of gene flow. The strength of natural selection is of foremost importance because larger fitness consequences for deviating from the optimal phenotype in a particular location can potentially lead to greater evolutionary change (Falconer & MacKay, 1995). The rate of gene flow is important because migration into a region experiencing idiosyncratic selection can overwhelm that selection, preventing regional trait differences (i.e. local adaptation) from accumulating (Nagylaki, 1975; Slatkin, 1978; Wright, 1931; Yeaman, 2015). In discrete space models, the evolution of local adaptation in a particular location depends on the ratio of gene flow from dissimilar environments ($m$) to the strength of selection ($s$), $m/s$ (Slatkin, 1978; Wright, 1931; Yeaman, 2015). In models of continuous space, patterns of dispersal relative to the strength of selection can be used to determine the minimum size a region experiencing idiosyncratic selection needs to be for locally adaptive differences to accumulate, the so-called

"characteristic length" of a cline. Specifically, when dispersal is modelled as a diffusion process, the
characteristic length is the standard deviation of dispersal distances ($\sigma$) relative to the square root of the
strength of selection (i.e. $\sigma/\sqrt{s}$). Such characteristic lengths can be described for individual alleles
(Nagylaki, 1975; Slatkin, 1973) or for stabilising selection acting on polygenic traits (Barton, 1999;
Slatkin, 1978), see reviews by Felsenstein, (1976) and Lenormand (2002). Furthermore, the mean values
of polygenic traits will more closely track changes in phenotypic optima over space if those changes are
small (Barton, 1999; Slatkin, 1978). Because natural populations often inhabit large spatial ranges
encompassing complex patterns of environmental variation, relative rates of gene flow among regions
of high environmental similarity or dissimilarity will vary across the landscape, influencing the evolution
of local adaptation.
100
That the spatial pattern of environmental heterogeneity will influence the evolution of local adaptation
has been recognised since at least the 1960s (Antonovics, 1971; Antonovics & Bradshaw, 1970; Forester
et al., 2016; Hadfield, 2016; Levins, 1966; Schiffers et al., 2014). Indeed, the overall level of local
adaptation a species exhibits can be strongly affected by the pattern of environmental variation over
space (Forester et al., 2016; Gilbert & Whitlock, 2017; Hadfield, 2016; Schiffers et al., 2014) and several
studies have framed this concept in terms of the spatial autocorrelation of the environment (Hadfield,
2016; Urban, 2011). Spatial autocorrelation describes the similarity of observations from nearby
locations and can be quantified, for example, using Moran's I (Moran, 1950). Consider the maps of
environmental heterogeneity shown in Figure 1A. When the environment that gives rise to spatially
varying selection across a species' range exhibits high spatial autocorrelation (e.g. the right-hand map in
Figure 1A), selection pressures may be similar over large areas and changes in environment over space
will tend to be gradual. On the other hand, when the environment exhibits weak autocorrelation (e.g.
the left-hand map in Figure 1A), regions experiencing idiosyncratic selection will be comparatively small
and selection pressures may change rapidly over space. Of course, other factors such as variation in
population density, the magnitude of environmental variation and the scale of dispersal will also affect
the outcomes of spatially varying selection. All else being equal, though, a species with restricted
migration will tend to evolve the strongest local adaptation if agents of selection exhibit high levels of
spatial autocorrelation.
119
It is likely that the spatial pattern of selection across a species' range affects the genetic architecture
underlying local adaptation. In simple two-patch models, the relative balance of selection and migration
influences the number of alleles underlying local adaptation, their effect sizes and rates of allelic
turnover (Yeaman & Whitlock, 2011). In continuous, linearly varying landscapes, frequencies of alleles
contributing to locally adaptive traits are expected to vary in relation to their effect sizes and the
proximity of local trait means to phenotypic optima (Polechová & Barton, 2015). However, such models
may not fully predict the patterns of genetic variation expected in complex landscapes where the
relative balance of selection and migration vary across space. Previous studies have examined the
effects of landscape structure on the genetics of local adaptation in single locus models (Forester et al.,
2016) or in models of population expansion (Gilbert & Whitlock, 2017; Schiffers et al., 2014), but it is
unclear how the polygenic architecture of local adaptation will be influenced by the spatial pattern of
environmental variation.
132
In this paper, I examine patterns and the genetic bases of local adaptation in complex landscapes.
Following previous studies, I cast spatial patterns of environmental variation across a species' range in
terms of spatial autocorrelation. With population genetic simulations, I examine the patterns and
genetic architectures of local adaptation that evolve in environments that vary in their degree of spatial
autocorrelation. These simulations show ways that the pattern of environmental variation across a

species' range can influence the genetic variation underlying local adaptation. Finally, I analyse empirical
data from a large-scale experiment in lodgepole pine and find evidence suggesting a link between spatial
autocorrelation in climatic/environmental variation with the strength of local adaptation in a natural
system. Taken together, the results of this study highlight the importance of considering the spatial
pattern of environmental variation in studies of local adaptation.
143
## Results and Discussion

### Simulating local adaptation to spatially heterogeneous environments

To further understand the effects that the spatial pattern of environmental variation can have on
patterns of local adaptation, I constructed a simulation model of spatially varying selection. I used
forward-in-time population genetic simulations in SLiM (4.1; Haller & Messer, 2023) modelling a 2-
dimensional stepping-stone metapopulation of 196 demes (i.e. a 14x14 grid). Migration was restricted
to adjacent demes with rates of gene flow that resulted in pronounced population structure with clear
isolation-by-distance (Figure S1). Spatially varying selection was modelled as stabilising selection, where
each deme ($d$) had a particular phenotypic optimum ($\theta_d$), i.e. an individual in deme $d$ had higher fitness
if its phenotype was close to $\theta_d$. The strength of stabilising selection was set such that an individual with
the optimal phenotype for the deme with the most negative optimum translocated into the deme with
the most positive optimum would experience a 50% reduction in fitness (strong selection) or a 25%
reduction in fitness (moderate selection). I used a quantitative trait model to study local adaptation,
because it is thought that the traits involved in local adaptation are generally polygenic (Savolainen et
al., 2013). Additionally, there is evidence that the genetic basis of local adaptation can involve alleles
that have spatially antagonistic fitness effects as well as conditionally neutral effects (Anderson et al.,
2013) and both kinds of effects can arise in a quantitative trait model given variation in genetic
backgrounds and environments. I constructed a set of 200 maps of normally distributed environmental
variation that varied in the degree of spatial autocorrelation (three examples are shown in Figure 1A). I
quantified spatial autocorrelation in the environment using Moran's I, which varied from 0.05 (weak
autocorrelation) to 0.95 (strong autocorrelation) in the maps I constructed. The maps of environmental
variation were used to specify phenotypic optima in individual simulations.
166

**Figure 1** The pattern of environmental heterogeneity influences the outcomes of spatially varying selection. A) Three examples of environmental heterogeneity with similar distributions of phenotypic optima ranging from low autocorrelation on the left to high autocorrelation on the right. B) The pattern of local adaptation that evolved on the landscapes shown in panel A. C) The average local adaptation that arises as a function of Moran's I across 200 maps of environmental variation. C) The coefficient of variation for local adaptation across the 200 maps. The simulation results shown are for cases with mean $F_{ST}$ ~ 2% and moderate stabilising selection.

The effects of environmental structure on patterns of local adaptation that evolved in simulations were profound. I measured local adaptation in simulated populations by comparing an individual's fitness at "home" versus "away" following the method outlined by (Blanquart et al., 2013). Using this method, the observed local adaptation for each deme ($LA$) in the metapopulation was computed (e.g. Figure 1B). As expected, the mean local adaptation across populations ($\overline{LA}$) increased with the degree of spatial

autocorrelation in the environment (Figure 1C), consistent with Hadfield (2016). This result held over
various levels of gene flow and strengths of stabilising selection (Figure S2). Varying the rate of gene
flow and/or the strength of selection had an effect on the level of local adaptation that arose in a given
case, but a pattern of increasing $\overline{LA}$ with Moran's I was always observed (Figure S2). In natural
populations, it is likely that multiple aspects of environmental variation will induce spatially varying
selection. In cases where simulated populations had multiple traits subject to selection due to different
aspects of environmental variation, the trait corresponding to the more autocorrelated environment
exhibited greater local adaptation (Figure S3).
189
The spatial pattern of environmental variation did not just affect the average level of local adaptation,
though, it also had a large influence on the variation in local adaptation across the landscape (Figure 1B).
The coefficient of variation in local adaptation $[CV(LA)]$ across the landscape decreased rapidly with
increasing autocorrelation (Figure 1D, S2B). When environmental variation was weakly autocorrelated,
the $CV(LA)$ was as much as 30x higher than for more highly autocorrelated environments (Figure 1D).
Variation in the degree of local adaptation across a species range is understudied in the population
genetics literature but has important implications (see below).
197
At a finer scale, environmental variation in the immediate vicinity of a particular deme predicted its level
of local adaptation and genetic variation, as predicted by theory (Barton, 1999; Guillaume & Whitlock,
2007; Slatkin, 1978). Demes that were surrounded by populations with highly similar phenotypic optima
evolved greater local adaptation than demes bordering more dissimilar environments (Figure S4A). This
was particularly evident when the overall landscape was weakly autocorrelated (Figure S4A),
presumably because in highly autocorrelated landscapes most demes are surrounded by similar
environments. Furthermore, additive genetic variance ($V_A$) for the trait under selection was highest in
demes surrounded by dissimilar environments (Figure S4B), suggesting that gene flow among locally
divergent populations has an effect of increasing genetic variability. Such a positive correlation between
$V_A$ and local environmental heterogeneity has been reported in lodgepole pine (Yeaman & Jarvis, 2006).
The magnitude of a species' response to selection on a trait is expected to be proportional to $V_A$
(Falconer & MacKay, 1995), thus the pattern of environmental variation that local adaptation evolves
under may influence how a species responds to changing environments.
211
Environmental structure and the genetic architecture of local adaptation
The results so far demonstrate that the structure of the environment can have a clear impact on the
patterns of local adaptation that evolve, but does it influence the genetic basis of that adaptation?
Under a model of spatially varying stabilising selection, each polymorphism that affects the phenotypes
under selection will influence local adaptation, but the extent of this will depend on its effect size,
where it is present and its allele frequencies. For each polymorphism in a simulation, I quantified the
contribution it makes to mean local adaptation ($\overline{LA}$) as follows. I shuffled the presence/absence of a
particular polymorphism across the landscape, effectively erasing its contribution to local adaptation.
Average local adaptation was then recalculated without the contribution of the focal polymorphism
($\overline{LA_\iota}$). The relative contribution of the focal polymorphism to mean local adaptation across the
landscape was then calculated as $LA_{Rel,l} = (1 - \overline{LA_\iota}/\overline{LA})$. For example, a polymorphism with $LA_{Rel} \approx$
1.0 would be the basis of all local adaptation, while one with $LA_{Rel} \approx 0.0$ would have no effect. Note,
$LA_{Rel}$ is not strictly a proportion (see Methods for details).
225

226
**Figure 2** The genetic architecture of local adaptation is influenced by the structure of the environment.
A) The proportion of total local adaptation explained by alleles that individually explain different
amounts of local adaptation ($LA_{Rel}$) varies as a function of environmental autocorrelation. Lines
represent LOESS curves fit with a span parameter of 1.5. B) The mean allele frequencies compared to
the squared effect sizes of polymorphisms that underly local adaptation differ depending on the pattern
of the environment. The contour lines indicate regions with high densities of points. High
autocorrelation refers to data from maps with the 50 highest values of Moran's I. Low autocorrelation
refers to data from maps with the 50 lowest values of Moran's I. Results in both panels come from
simulations with $F_{ST}$ = 0.02 and moderate stabilising selection.
236
The distribution of locally adaptive effects varied in relation to the pattern of environmental variation
(Figure 2A, S5). In environments exhibiting weak autocorrelation, polymorphisms that individually made
a large contribution to local adaptation across the species' range ($LA_{Rel} > 0.10$) were largely absent
and polymorphisms that made intermediate ($0.01 < LA_{Rel} < 0.10$) and small contributions ($LA_{Rel} <$
$0.01$) explained most of the local adaptation that evolved (Figure 2A). In environments that were more
highly autocorrelated, polymorphisms with $LA_{Rel} > 0.10$ made a substantial contribution to local
adaptation alongside those with intermediate and small effects, particularly under strong stabilising
selection (Figure S5). These general patterns held over different levels of gene flow (Figure S5).
245

Patterns of genetic variation underlying locally adaptive polymorphisms varied depending on the degree
of spatial autocorrelation in the environment. In general, locally adaptive polymorphisms with similar
$LA_{Rel}$ tended to have smaller phenotypic effects but larger allele frequencies in highly versus weakly
autocorrelated environments (Figure 2B). In highly autocorrelated environments, alleles may readily
spread among populations facing similar environmental challenges. However, when alleles with large
phenotypic effects spread among neighbouring demes, they may cause individuals to overshoot their
respective phenotypic optima, so the alleles that are maintained may tend to have smaller phenotypic
effects. In weakly autocorrelated environments, on the other hand, genes flowing from one location to
another have a much greater chance of encountering highly divergent environments, preventing locally
adaptive alleles from spreading across wide regions. In such cases, phenotypic effects may need to be
large for locally adaptive mutations to withstand the swamping effects of gene flow. These general
patterns were observed with both strong and moderate selection (Figure 2B, S6A) as well as over
varying levels of gene flow (Figure S6A). Indeed, the patterns of allele frequency versus phenotypic
effect still held when looking at absolute effects on local adaptation, though they were much less
pronounced (Figure S6B).
261
Characterising the genetic basis of local adaptation is important, and researchers generally attempt to
do so using one of two strategies; by comparing phenotypic variation for traits important for local
adaptation to genetic variation (i.e. genome-wide association studies, GWAS) or environmental variation
to genetic variation (i.e. genotype-environment association analysis, GEA analysis)(reviewed in Lasky et
al. 2023). Since patterns of genetic variation underlying local adaptation can differ depending on the
pattern of environmental variation, statistical power to identify the genetic basis of local adaptation will
likely vary for different aspects of the environment. To demonstrate this, I performed GWAS on
phenotypes for 1,000 randomly chosen individuals from the simulations and corrected for population
structure using the kinship matrix. Figure S7 shows that the -$\log_{10}$($p$-values) for alleles that contribute
similar levels of local adaptation tend to be smaller (i.e. there is less power) for high versus low
autocorrelation environments. Some of this difference may partially be due to the population structure
correction procedure (see below), but it demonstrates that the pattern of environmental variation that
gave rise to local adaptation can affect the ability to study the genetics of that adaptation.
275
There are numerous factors that may interact with the pattern of spatially varying selection to shape the
genetics of local adaptation that I did not explore here. The degree of genetic redundancy in relevant
traits, distribution of phenotypic effect sizes, mutation rates and patterns of dispersal can all influence
the genetics of local adaptation (e.g. Láruson et al., 2020; Yeaman, 2013; Yeaman & Whitlock, 2011).
Follow up studies looking at how such factors influence the genetic basis of local adaptation in
differently structured environments are needed. However, the results from the simulations should
provide researchers seeking to characterise the genetic basis of local adaptation with useful intuition.
283
The evolution of local adaptation: maladaptation and allelic turnover
In heterogeneous environments certain polymorphisms may have a net effect of reducing local
adaptation across a species' range. All populations will harbour such locally maladaptive alleles, because
any new mutation that increases distance between an individual's phenotype and the local optimum will
reduce local adaptation even if only by a small amount. By summing the effects of all polymorphisms
with $LA_{Rel} < 0$ across a simulation, I obtained a measure of the cumulative local maladaptation across a
meta-population. Note that the cumulative local maladaptation is analogous to "migration load". All
simulations exhibited some degree of maladaptation regardless of the level of autocorrelation, but the
cumulative effects of maladaptive alleles were always higher under weakly versus highly autocorrelated

environments (Figure 3A, S8B). Increasing the rate of gene flow increased the degree of maladaptation
and increasing the strength of selection decreased it (Figure S8B).
295
Regardless of the pattern of environmental variation in a simulation, levels of local adaptation had been
maintained at a steady state for many generations before they were sampled (Figure S9). However, the
average age of alleles underlying local adaptation increased with increasing autocorrelation in the
environment (Figure 3B). Furthermore, weighing the average allele age within a simulation by effect
size, the increase in allele age with spatial autocorrelation was even more pronounced (Figure 3B). Thus
large effect alleles, in particular, are maintained for longer times in more highly autocorrelated
landscapes. Taken together, these results demonstrate that the rate of allelic turnover is greater for
more weakly autocorrelated environments.

304
**Figure 3** Species-wide maladaptation and the age of locally adaptive alleles are influenced by the pattern
of environmental variation. A) Cumulative local maladaptation, the summed effects of all polymorphism
that have a net negative effect on local adaptation across a simulated species' range, decreases with
increasing autocorrelation. Points represent individual simulations and lines represent LOESS curves fit
with a span parameter of 1.5. B) Alleles underlying local adaptation tend to be older when the
environment is more highly autocorrelated. Lines represent LOESS curves fit to the data treating all
polymorphisms equally, or by giving higher weight to polymorphisms with greater effect size.
312
Local adaptation, environmental autocorrelation, provenance trials and lodgepole pine
The simulation results clearly show how the pattern of environmental variation across a species range
may influence the evolution of local adaptation and the genetics underlying it. In natural populations, if

the relative strength of selection acting on different aspects of environmental variation is unknown, we
should perhaps predict that local adaptation will be strongest when the environment is highly
autocorrelated. Despite such strong predictions, though, empirical evidence that patterns of local
adaptation coincide with spatially autocorrelated features of the environment is lacking (Siepielski et al.,
2013). Spatial autocorrelation in patterns of biotic interactions can explain a large proportion of
variation in trait differentiation among populations in several species (Urban, 2011), but such variation is
not necessarily locally adapted. Local adaptation has been demonstrated in many forest tree species
using provenance trials (Leites & Benito Garzón, 2023), but not to test the prediction that the pattern of
environmental variation influences local adaptation.
325
Provenance trials involve planting multiple populations of a species in numerous common gardens to
assess how "transfer distance", the distance between home and the common garden, affects
productivity (reviewed in Wadgymar et al., 2022). However, the structure of provenance trials is such
that the methods I used to quantify local adaptation in the simulations above are not necessarily
applicable. To demonstrate how provenance trial data could be used to quantify local adaptation, I
conducted *in silico* provenance trials on the simulations (e.g. Figure 4A). As expected, the slope of fitness
on transfer distance in provenance trials is increasingly negative with increasing autocorrelation (Figure
4B, S10) and strongly negatively correlated with mean local adaptation ($\overline{LA}$) in simulations (Figure 4C,
S10). Provenance trials may, thus, contain information that is useful for assessing whether the pattern of
environmental across a landscape is important in shaping local adaptation.
336

338
**Figure 4** Comparing the results from a simulated provenance trial to measures of local adaptation. A) A
map of a provenance trial conducted on a simulated population showing the locations of planting sites
and provenances. B) Linear regressions of relative fitness on environmental transfer distance for
landscapes with differing levels of environmental autocorrelation. C) The slope of relative fitness on
transfer distance compared to the mean local adaptation ($\overline{LA}$) across simulated meta-populations.
Spearman's ρ and its *p*-value are shown inset in the panel C.

345
The Illingworth trial is an exceptionally large provenance trial established by the Ministry of Forestry in
British Columbia, Canada in the 1970s to establish seed-transfer guidelines for the lodgepole pine (*Pinus*
*contorta*) (Illingworth, 1978). Seeds were collected from 140 provenances from Northwestern North
America and seedlings were planted in a set of 62 sites distributed across British Columbia (Figure 5A).
Phenotypic data has been recorded for around 60,000 individual trees since the Illingworth trial began
and previous studies have used this data to demonstrate clear local adaptation in lodgepole pine
(Mahony et al., 2020; Wang et al., 2006). Given the geographic breadth of the Illingworth trial (Figure
5A), it represents a suitable dataset to test the prediction that spatial patterns of environmental
variation influence the evolution of local adaptation.

355
I analyzed data from the Illingworth trial using a mixed-modelling approach. Different aspects of climatic
variation across the sites in the Illingworth trial are highly intercorrelated (Figure S11A), so I used
principal components analysis to identify independent axes of climatic/environmental variation in the
dataset. I restricted the analysis to the first 6 principal components (PCs), as each one explained at least
1% of the variation in the data and combined they explained 95% of the variation (Figure S11B). I then
regressed tree diameter at breast height and survival measured after 20 years on transfer distance
between planting site and provenances in PC-space (Figure 5B-C) (see Methods for details). For both
diameter at breast height and survival, there were significant negative relationships predicting
phenotypic variation from climatic PCs (Figure 5B-C). If diameter at breast height and/or survival are
considered proxied for fitness, then the results shown in Figures 5B-C indicates local adaptation along
several dimensions of climatic variability.

367
Patterns in the Illingworth trial data suggest that local adaptation in lodgepole pine is strongest when
climatic/environmental variation is highly spatially autocorrelated (Figure 5C-D). For both phenotypes,
PC1 was the strongest climatic predictor (Figure 5B-C). PC1 captures climatic/environmental differences
between coastal and inland locations (Figure S12). Spatial autocorrelation for PC1 was also among the
highest in the dataset (Figure 5D-E). For survival, there was a negative relationship between the
regression slope and Moran's I (Figure 5E), which is qualitatively similar to the analysis of simulated
provenance trials (Figure 4B). However, I did not conduct a formal statistical test of the relationship

between spatial autocorrelation and local adaptation because only 4 PCs exhibited statistically
significant evidence for local adaptation so such an analysis would be underpowered. While this means
the results are merely suggestive rather than concrete, they are in line with the prediction that spatial
autocorrelation in the abiotic environment predicts the strength of local adaptation in natural
populations.

380
**Figure 5** Analysis of local adaptation in lodgepole pine from the Illingworth provenance trial. Panel A)
shows the map of provenances and planting sites in the Illingworth trials across the Northwest of North
America. B) Fitted relationships between tree diameter at breast height (DBH) and transfer distance for
8 principal components. C) Survival probability as a function of transfer function for 8 environmental
principal components. D) Linear regression coefficients for the relationships shown in B compared to
degree of autocorrelation in the environment. E) Logistic regression coefficients for the relationships
shown in C compared to the degree of spatial autocorrelation in the environment. Statistical significance
was assessed at α = 0.05 after correcting for multiple comparisons using the Dunn-Šidak method.

## Population structure and the genetic basis of local adaptation
Knowing the genetic basis of local adaptation in natural systems would give us a better understanding of
evolution, but may also be informative for conservation and management (Grummer et al., 2022). Many
studies have used methods that associate environmental/phenotypic variation with genotypes or allele
frequencies to characterise the genetic basis of local adaptation (Lasky et al., 2023). A large proportion
of species exhibit a specific pattern of population structure termed "isolation-by-distance" (IBD) where
genetic distance is positively correlated with geographic distance or measures of resistance based on
features of the landscape (Jenkins et al., 2010). IBD can arise with restricted migration, when rates of
gene flow are highest among parts of a species' range that are in close proximity, though it can also
reflect demographic histories such as past population expansion (Slatkin, 1993). Greater levels of local
adaptation may evolve across a species' range if the spatial pattern of environmental variation aligns

with the opportunity for migration (Figure 1). Thus, stronger local adaptation is expected to arise when
selection is co-linear or co-autocorrelated with patterns of population structure.

The relationship between population structure and the structure of the environment likely impacts our
ability to study the genetic basis of local adaptation. It is well established that a pattern of IBD can
confound the search for genes involved in local adaptation (Meirmans, 2012). Indeed characterising the
genetic basis of local adaptation when the agents of selection are co-linear or co-autocorrelated with
patterns of gene flow, termed "isolation by environment", is particularly challenging (Wang & Bradburd,
2014). Many studies have analyzed genotype-environment associations (GEA) to characterize the
genetic basis of local adaptation. Such association methods often treat population structure as a
nuisance variable and various approaches are taken to correct for it. This is done for the statistical
necessity of establishing a suitable null model (Meirmans, 2012). For example, latent factor mixed
models (LFMMs) are widely used to conduct GEA analyses that correct for population structure (Caye et
al., 2019; Frichot et al., 2013). However, Lotterhos (2023) recently found that the sensitivity of the
LFMM method declined with increasing correlation between the environment and major axes of
population structure. This all suggests that characterising the genetic basis of local adaptation is
particularly difficult in the exact cases where local adaptation is expected to be strongest, when
selection pressures and population structure are highly co-autocorrelated over space. Careful sampling
strategies may alter the power of association methods (Lotterhos & Whitlock, 2015; Meirmans, 2015;
Wang & Bradburd, 2014), but such strategies require *a priori* hypotheses about locally adapted traits,
the agents of selection (e.g. Kreiner et al., 2022) and/or the genes involved (e.g. Fournier-Level et al.,
2011).

Implications for conservation management
A result from the simulations that was particularly striking is the heterogeneity in local adaptation that
can arise under different patterns of environmental variation. The only difference between the
simulations shown in Figure 1 is the pattern of environmental variation, yet the average level of local
adaptation increased by a factor of roughly 5x and the coefficient of variation in local adaptation
decreased by 30x when comparing the cases with the highest and lowest spatial autocorrelation in the
environment. High heterogeneity in local adaptation across a species' range may impact conservation
interventions and population genetic analyses.

Any practical conservation intervention that uses average patterns of local adaptation to project
performance under changing climates should carefully consider heterogeneity in local adaptation. In
forestry, for example, seed-transfer guidelines based on average transfer functions from provenance
trials may give a misleading picture of performance for some provenances if there is high heterogeneity
in local adaptation. The applicability of a single transfer function would vary depending on how
heterogeneous local adaptation is among the populations in question.
439
In recent years, population genetic analyses have been developed to identify parts of a species range
that are particularly vulnerable to climate change (i.e. genomic offset; Rellstab et al., 2021). Such
methods analyse present-day relationships between allele frequency and the environment to predict
how species will fare given predicted patterns of environmental change. If the agents of local adaptation
are highly spatially autocorrelated, neutral population structure may be partially aligned with gradients
of selection, which could explain why the use of "adaptive" genetic markers and randomly chosen
markers seem to perform equally well in some offset analyses (Fitzpatrick et al., 2021; Láruson et al.,
2022; Lind et al., 2023). Violating the assumption of homogeneous local adaptation in offset analyses,
for example, would likely introduce noise into predictions but could potentially lead to spurious results.
449
Thinking about environmental structure when building hypotheses about local adaptation
Detailed prediction of the environmental variation that is relevant to patterns of local adaptation
requires an understanding of a species' life history and physiology. The fundamental factor underlying
the evolution of local adaptation is the relative balance of selection and dispersal (see Introduction).
However, the specific pattern of environmental variation across a landscape influences whether
dispersing individuals are likely to encounter environments similar to those of their parents. Unlike
dispersal or the strength of selection, which are hard to quantify, spatial autocorrelation in the
environment is readily measurable. When seeking to characterise the genetic basis of local adaptation,
studies comparing different aspects of environmental variation should consider the pattern of such
variation when forming their hypotheses. For example, before comparing GEA results for the different
*bioclimatic* variables from WorldClim, researchers could examine how these variables are distributed
over space to form *a priori* hypotheses about factors underlying local adaptation. Of course, an aspect of
the environment may be highly autocorrelated in space, but if the variation it exhibits does not
correspond to varying selection pressures, then it is unlikely to be directly related to local adaptation.
That local adaptation is predicted to be stronger with increasing autocorrelation in the environment
does not imply that strong local adaptation cannot arise in highly heterogeneous environments or with
little spatial autocorrelation. There are numerous examples of local adaptation to environmental
heterogeneity that is not smoothly distributed in space. For example, heavy metal concentrations in
mine tailings impose selection that is so strong it overwhelms the effects of gene flow (Jain & Bradshaw,
1966). It must be kept in mind that the patchy distributions of environmental variation across a
landscape, as may be the case for heavy-metal rich soils, may be more or less autocorrelated from the
perspective of a given species depending on its dispersal behaviour. In my simulations, I matched the
granularity of dispersal with that of the environment (Figure 1A). For real species, considering
environmental variation at a scale relevant to how species disperse is critical and recent population
genetic advances may make estimating dispersal in natural populations much less time-intensive than
previously (Bradburd & Ralph, 2019; Smith et al., 2023). Comparing patterns of species dispersal with
patterns of variation in environmental variation that is plausibly relevant to selection may help identify
the drivers of local adaptation in natural populations.

### Closing remarks

While it has been a long-standing expectation that the pattern of environmental variation (and
particularly spatial autocorrelation) will influence the evolution of local adaptation (e.g. Hadfield, 2016;
Levins, 1966), the simulation results and analysis of the lodgepole pine data should serve to emphasise
how important the spatial pattern of climatic/environmental variation can be. The spatial pattern of
environmental variation that a natural population has experienced will have likely shaped the evolution,
current patterns and genetic underpinnings of local adaptation. Thus, it likely also influences how
populations will respond to changes in the future.

Materials and Methods

Simulating spatially varying selection

To explore the effects of landscape structure on the outcomes of spatially varying selection, I
constructed set of maps that exhibited varying degrees of spatial autocorrelation. Maps of normally
distributed environmental heterogeneity were constructed using the midpoint displacement algorithm
as implemented in the *NLMpy* package (Etherington et al., 2015). I simulated a 14x14 cell grid (i.e.
landscape), specifying the desired level of autocorrelation to achieve a set of 200 maps, spanning the
range of Moran's I values from 0.05 to 0.95 (i.e. Moran's I varied in increments of 0.0045). Simulated
maps were rejected if the mean value across the landscape was less than 0.4 or greater than 0.6. This
ensured that the the approximately normal distributions of environmental values across the landscape
were roughly equivalent across maps.

Using SLiM v4.1 (Haller & Messer, 2023), I modelled a 2-dimensional stepping-stone meta-populations
with 196 demes (i.e. a 14x14 grid). Each deme contained 100 diploid individuals for total meta-
population size of 19,600. Migration occurred between adjacent demes in the four cardinal directions
except for populations at the range edge where migrants only moved back into demes they were
connected to. Migration rates were set at 0.07, 0.035 or 0.0175, leading to population-wide neutral $F_{ST}$
values of 0.02, 0.05 and 0.10, respectively (Figure S1A). Each diploid individual had a 10Mbp long
genome that recombined at a constant rate of $r$=1 x $10^{-7}$. When modelling a single trait, mutational
effects were distributed as $N(0,1)$ and occurred at random along the sequence at a rate of $\mu = 10^{-10}$,
corresponding to a mutational variance of 0.001 for the trait subject to stabilising selection. When
modelling two traits, the mutation rate was the same, but effects were modelled as multivariate normal
with means of 0, variances of 1 and covariances of 0 (i.e. mutational effects for the two traits were
independent). A diploid individual's phenotype for a given trait was the additive combination of the
effects on that trait for the alleles the individual possessed (i.e. mutations were semidominant).
Spatially varying stabilising selection was modelled using the maps of environmental heterogeneity to
specify the distribution of phenotypic optima across the landscape. An individual's relative fitness $W_i$
was calculated using the standard expression for Gaussian stabilising selection (Walsh & Lynch, 2018):

$W_i = \exp\left[-\left(\frac{(\alpha_{i,d}-\theta_d)^2}{2V_s}\right)\right]$ Equation 1

where $V_S$ is the variance of the Gaussian fitness function, $\alpha_i$ is the phenotype of the $i^{th}$ individual in deme
d, and $\theta_d$ is the phenotypic optimum of deme d. When modelling stabilising selection in cases with two
traits, an individual's fitness was calculated as follows:

$W_i = \exp\left[-\left(\frac{(\alpha_{i,1,d}-\theta_{d,1})^2+(\alpha_{i,2,d}-\theta_{d,2})^2}{2V_s}\right)/2\right],$    Equation 2

where $\alpha_{i,1,d}$ and $\alpha_{i,2,d}$ are the values for traits 1 and 2 for individual i in deme d, respectively, and $\theta_{d,1}$ and
$\theta_{d,2}$ are the phenotypic optima for traits 1 and 2, respectively. In effect, an individual's relative fitness in
this 2-trait model is the average of the marginal finesses for each trait.

To achieve an equilibrium of migration, selection and drift, meta-populations evolved for 100,401
generations. Initially, meta-populations evolved under stabilising selection with an optimum of 0 in all
demes. After 400 generations, the landscape was altered to one of the 200 maps of environmental
heterogeneity and kept in that state for a further 100,000 generations. At the end of the simulation,
phenotypes of each individual in each deme were recorded as well as the genealogical history of the
meta-population stored as a tree-sequence. PySlim, tskit and msprime packages (Baumdicker et al.,
2022; Haller et al., 2019) were used to work with the output tree-sequence files. To calculate Weir and
Cockerham's $F_{ST}$, neutral mutations were added to the simulated population using PySlim at a rate of $10^{-8}$/bp.
533    

 Analyzing simulated data

Local adaptation was quantified for each deme using the "home-versus-away" (HA) method outlined by
Blanquart et al. (2013). Specifically, each individual's fitness was quantified in its home deme and every
other possible location on the landscape. The mean local adaptation was calculated in each deme as the
mean difference in fitness between home and away conditions across all individuals. For each deme $d$
not on the edge of the simulated landscape, I quantified local heterogeneity in the landscape as the
mean sum of squares between the focal deme's environment and that of the four adjacent demes (in
the cardinal directions). For each deme, across the $n$ polymorphisms that affected the phenotype
additive genetic variance for the trait was calculated as $V_{A,d} = \sum_{i=1}^{n} p_{i,d}(1 - p_{i,d})\gamma_i^2$, where $p_{i,d}$ is the
allele frequency of SNP $i$ in deme $d$ and $\gamma_i$ is the phenotypic effect of SNP $i$.

The contribution of individual SNPs to local adaptation was quantified as follows. For each
polymorphism that affected the trait(s) under selection, the presence/absence of the allele in different
haplotypes in different demes can be represented as a vector of 1s and 0s. By shuffling this vector, the
contribution of this polymorphism to local adaptation is effectively erased, while keeping its
contribution to additive genetic variance across the species' range constant. For polymorphism $l$, I
recomputed all phenotypes for all individuals after shuffling allele frequencies and re-quantified local
adaptation as $\overline{LA_l}$. The relative contribution of the focal polymorphism to local adaptation is calculated
as:

$$LA_{Rel,l} = (1 - \overline{LA_l}/\overline{LA}). \hspace{3cm} \text{Equation 3}$$

Note that $LA_{Rel}$ is not strictly a proportion, as epistasis for fitness that arises in models of stabilising
selection means that the $\sum_{i=1}^{n} LA_{Rel,i} \neq 1$ for the $n$ SNPs that affect phenotypes. Furthermore, alleles
that have a net negative effect on local adaptation (i.e. they are locally maladaptive) will have negative
$LA_{Rel}$ values. Indeed, the total amount of local maladaptation in a meta-population was calculated as
the additive combination of all polymorphisms with negative $LA_{Rel}$.

Provenance trials were conducted on simulated data by sampling a set of 50 "planting sites" and a set of
100 "provenances". The relative fitness of each provenance was computed in each of the 50 planting
sites. The absolute difference in phenotypic optimum for each provenance and each planting site was
used as environmental distance. Using *lme4* in R, I fitted a linear mixed model regressing relative fitness
on environmental distance with provenance as a random effect, with slopes and intercepts varying
across provenances.

I combined results across the 50 simulations with the lowest levels of spatial autocorrelation (weak
autocorrelation) and the 50 simulations with the highest autocorrelation (high autocorrelation) and
examined the relationship between allele frequency and phenotypic effect sizes for the alleles
underlying local adaptation.

For each simulation, I randomly sampled 1,000 individuals from the landscape and I recording their trait
values as well as neutral and phenotype affecting polymorphisms. I applied LD-pruning (with a threshold
of $r^2 < 0.2$) to the neutral SNPs and used these data to infer the kinship matrix using PLINK (v2; Chang et
al., 2015). I performed association studies on the phenotype affecting SNPs from individual simulations
with GEMMA (v0.98.5; Zhou & Stephens, 2012), using the inferred kinship matrix as a random effect to
account for population structure.

### Analysis of data from the Illingworth trial

ClimateNA (Wang et al., 2016) was used to extract climatic data for each location in the Illingworth Trial. Across the locations in the Illingworth trial, many aspects of climatic/environmental variation are highly inter-correlated (Figure S11B). Because such inter-correlation would make it difficult to tease apart the effects of individual aspects of climatic/environmental variation on local adaptation, I conducted a principal components analysis (PCA) to separate the variation onto independent axes. I restricted the analysis to the first 6 principal components as these explained 95% of climatic variation. Diameter at breast height (DBM) and tree height exhibit a strong positive correlation (Pearson's $r$ = 0.9), so analyses were restricted solely to DBM. Trees that were dead or dying after 20 years were given a survival score of 0, living trees were scored a 1.

Phenotype and survival data after 20 years for individual trees from the Illingworth trial were analyzed using mixed models. Mean normalised DBH was modelled as a normally distributed variable using the *lme4* package and survival using a generalised linear mixed model with a "logit" link function using the *glmer* package. The normalised Euclidean distance between each individual's provenance and planting site (i.e. transfer distance) in PC-space was used as a predictor in the model. Provenance, planting site and planting block within sites were included as having random effects on the slope and intercept of the relationship between phenotype and transfer distance.

Moran's I was calculated for each principal component of climatic variation across provenances (using the *ape* package) incorporating a pairwise Haversine distance matrix as weights in the calculation.

### Data accessibility

All the code used to perform, analyse, and plot the results of simulations is available at https://github.com/TBooker/LocalAdaptationArchitechture. R scripts to analyse and plot the results of the Illingworth trial data are available at https://github.com/TBooker/LocalAdaptationArchitechture, but the raw data files were used by permission of the BC Ministry of Forestry.

### Acknowledgements

Thanks to the Aitken and Whitlock labs, Anna Bazzicalupo, Katie Lotterhos, Ailene Macpherson, Greg O'Neil, Monty Slatkin, Wouter van der Wijl and Sam Yeaman for discussions and/or comments on the manuscript. Thanks to Hazel Booker for late night chat sessions. The data from the Illingworth trial was compliments of Nick Ukrainetz, the lodgepole pine breeder with the Forest Improvement and Research Management branch of the BC Ministry of Forests. TRB was supported by a Bioinformatics Postdoctoral Fellowship from the Biodiversity Research Centre at the University of British Columbia.

## Literature cited

Aitken, S. N., & Whitlock, M. C. (2013). Assisted Gene Flow to Facilitate Local Adaptation to Climate Change. *Annual Review of Ecology, Evolution, and Systematics*, *44*(1), 367–388.

Anderson, J. T., Lee, C.-R., Rushworth, C. A., Colautti, R. I., & Mitchell-Olds, T. (2013). Genetic trade-offs and conditional neutrality contribute to local adaptation. *Molecular Ecology*, *22*(3), 699–708.

Angert, A. L., Bontrager, M. G., & Ågren, J. (2020). What Do We Really Know About Adaptation at Range Edges? *Annual Review of Ecology, Evolution, and Systematics*, *51*(1), 341–361.

Antonovics, J. (1971). The Effects of a Heterogeneous Environment on the Genetics of Natural Populations: The realization that environments differ has had a profound effect on our views of the origin and role of genetic variability in populations. *American Scientist*, *59*(5), 593–599.

Antonovics, J., & Bradshaw, A. D. (1970). Evolution in closely adjacent plant populations VIII. Clinal patterns at a mine boundary. *Heredity*, *25*(3), 349–362.

Barton, N. H. (1999). Clines in polygenic traits. *Genetical Research*, *74*(3), 223–236.

Baumdicker, F., Bisschop, G., Goldstein, D., Gower, G., Ragsdale, A. P., Tsambos, G., Zhu, S., Eldon, B., Ellerman, E. C., Galloway, J. G., Gladstein, A. L., Gorjanc, G., Guo, B., Jeffery, B., Kretzschumar, W. W., Lohse, K., Matschiner, M., Nelson, D., Pope, N. S., … Kelleher, J. (2022). Efficient ancestry and mutation simulation with msprime 1.0. *Genetics*, *220*(3). https://doi.org/10.1093/genetics/iyab229

Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to measuring local adaptation. *Ecology Letters*, *16*(9), 1195–1205.

Bradburd, G. S., & Ralph, P. L. (2019). Spatial Population Genetics: It's About Time. *Annual Review of Ecology, Evolution, and Systematics*, *50*(1), 427–449.

Caye, K., Jumentier, B., Lepeule, J., & François, O. (2019). LFMM 2: Fast and accurate inference of gene-environment associations in genome-wide studies. *Molecular Biology and Evolution*, *36*(4), 852–860.

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1), 7.

Etherington, T. R., Holland, E. P., & O'Sullivan, D. (2015). NLM py: a python software package for the creation of neutral landscape models within a general numerical framework. *Methods in Ecology and Evolution / British Ecological Society*, *6*(2), 164–168.

Exposito-Alonso, M. (2023). Understanding local plant extinctions before it is too late: bridging evolutionary genomics with global ecology. *The New Phytologist*, *237*(6), 2005–2011.

Falconer, D. S., & MacKay, T. F. C. (1995). *Introduction to Quantitative Genetics* (4th ed.). Longman.

Felsenstein, J. (1976). The theoretical population genetics of variable selection and migration. *Annual Review of Genetics*, *10*, 253–280.

Fitzpatrick, M. C., Chhatre, V. E., Soolanayakanahally, R. Y., & Keller, S. R. (2021). Experimental support
for genomic prediction of climate maladaptation using the machine learning approach Gradient
Forests. *Molecular Ecology Resources*, *21*(8), 2749–2765.

Forester, B. R., Jones, M. R., Joost, S., Landguth, E. L., & Lasky, J. R. (2016). Detecting spatial genetic
signatures of local adaptation in heterogeneous landscapes. *Molecular Ecology*, *25*(1), 104–120.

Fournier-Level, A., Korte, A., Cooper, M. D., Nordborg, M., Schmitt, J., & Wilczek, A. M. (2011). A map of
local adaptation in Arabidopsis thaliana. *Science*, *334*(6052), 86–89.

Frichot, E., Schoville, S. D., Bouchard, G., & François, O. (2013). Testing for associations between loci and
environmental gradients using latent factor mixed models. *Molecular Biology and Evolution*,
*30*(7), 1687–1699.

Gilbert, K. J., & Whitlock, M. C. (2017). The genetics of adaptation to discrete heterogeneous
environments: frequent mutation or large-effect alleles can allow range expansion. *Journal of
Evolutionary Biology*, *30*(3), 591–602.

Grummer, J. A., Booker, T. R., Matthey-Doret, R., Nietlisbach, P., Thomaz, A. T., & Whitlock, M. C. (2022).
The immediate costs and long-term benefits of assisted gene flow in large populations.
*Conservation Biology: The Journal of the Society for Conservation Biology*, *36*(4), e13911.

Guillaume, F., & Whitlock, M. C. (2007). Effects of migration on the genetic covariance matrix. *Evolution;
International Journal of Organic Evolution*, *61*(10), 2398–2409.

Hadfield, J. D. (2016). The spatial scale of local adaptation in a stochastic environment. *Ecology Letters*,
*19*(7), 780–788.

Haller, B. C., Galloway, J., Kelleher, J., Messer, P. W., & Ralph, P. L. (2019). Tree-sequence recording in
SLiM opens new horizons for forward-time simulation of whole genomes. *Molecular Ecology
Resources*, *19*(2), 552–566.

Haller, B. C., & Messer, P. W. (2023). SLiM 4: Multispecies Eco-Evolutionary Modeling. *The American
Naturalist*, *201*(5), E127–E139.

Hartmann, H., Bastos, A., Das, A. J., Esquivel-Muelbert, A., Hammond, W. M., Martínez-Vilalta, J.,
McDowell, N. G., Powers, J. S., Pugh, T. A. M., Ruthrof, K. X., & Allen, C. D. (2022). Climate
Change Risks to Global Forest Health: Emergence of Unexpected Events of Elevated Tree
Mortality Worldwide. *Annual Review of Plant Biology*, *73*, 673–702.

Illingworth, K. (1978). Study of lodgepole pine genotype–environment interaction in B.C. *Proceedings
International Union of Forestry Research Organizations (IUFRO) Joint Meeting of Working
Parties: Douglas-Fir Provenances, Lodgepole Pine Provenances, Sitka Spruce Provenances, and
Abies Provenances. Vancouver, British Columbia, Canada.*, 151–158.

Jain, S. K., & Bradshaw, A. D. (1966). Evolutionary divergence among adjacent plant populations I. The
evidence and its theoretical analysis. *Heredity*, *21*(3), 407–441.

Jenkins, D. G., Carey, M., Czerniewska, J., Fletcher, J., Hether, T., Jones, A., Knight, S., Knox, J., Long, T.,
Mannino, M., McGuire, M., Riffle, A., Segelsky, S., Shappell, L., Sterner, A., Strickler, T., & Tursi,

R. (2010). A meta-analysis of isolation by distance: relic or reference standard for landscape
genetics? *Ecography*, *33*, 315–320.

Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*(12), 1225–
1241.

Kirkpatrick, M., & Barton, N. H. (1997). Evolution of a species' range. *The American Naturalist*, *150*(1), 1–
23.

Kreiner, J. M., Caballero, A., Wright, S. I., & Stinchcombe, J. R. (2022). Selective ancestral sorting and de
novo evolution in the agricultural invasion of Amaranthus tuberculatus. *Evolution; International*
*Journal of Organic Evolution*, *76*(1), 70–85.

Láruson, Á. J., Yeaman, S., & Lotterhos, K. E. (2020). The importance of genetic redundancy in evolution.
*Trends in Ecology & Evolution*, *35*(9), 809–822.

Láruson, Á. J., Fitzpatrick, M. C., Keller, S. R., Haller, B. C., & Lotterhos, K. E. (2022). Seeing the forest for
the trees: Assessing genetic offset predictions from gradient forest. *Evolutionary Applications*,
*15*(3), 403–416.

Lasky, J. R., Josephs, E. B., & Morris, G. P. (2023). Genotype-environment associations to reveal the
molecular basis of environmental adaptation. *The Plant Cell*, *35*(1), 125–138.

Leites, L., & Benito Garzón, M. (2023). Forest tree species adaptation to climate across biomes: Building
on the legacy of ecological genetics to anticipate responses to climate change. *Global Change*
*Biology*. https://doi.org/10.1111/gcb.16711

Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, *17*(4),
183–189.

Levins, R. (1966). The Strategy of Model Building in Population Biology. *American Scientist*, *54*(4), 421–
431.

Lind, B. M., Candido-Ribeiro, R., Singh, P., Lu, M., Vidakovic, D. O., Booker, T. R., Whitlock, M. C.,
Yeaman, S., Isabel, N., & Aitken, S. N. (2023). How useful is genomic data for predicting
maladaptation to future climate? In *bioRxiv* (p. 2023.02.10.528022).
https://doi.org/10.1101/2023.02.10.528022

Lotterhos, K. E. (2023). The paradox of adaptive trait clines with nonclinal patterns in the underlying
genes. *Proceedings of the National Academy of Sciences of the United States of America*,
*120*(12), e2220313120.

Lotterhos, K. E., & Whitlock, M. C. (2015). The relative power of genome scans to detect local adaptation
depends on sampling design and statistical method. *Molecular Ecology*, *24*(5), 1031–1046.

Mahony, C. R., MacLachlan, I. R., Lind, B. M., Yoder, J. B., Wang, T., & Aitken, S. N. (2020). Evaluating
genomic data for management of local adaptation in a changing climate: A lodgepole pine case
study. *Evolutionary Applications*, *13*(1), 116–131.

Meirmans, P. G. (2012). The trouble with isolation by distance. *Molecular Ecology*, *21*(12), 2839–2846.

Meirmans, P. G. (2015). Seven common mistakes in population genetics and how to avoid them.
724            *Molecular Ecology*, *24*(13), 3223–3231.

725    Moran, P. A. P. (1950). Notes on continuous stochastic phenomena. *Biometrika*, *37*(1–2), 17–23.

726    Nagylaki, T. (1975). Conditions for the existence of clines. *Genetics*, *80*(3), 595–615.

727    O'Neill, G. A., & Gómez-Pineda, E. (2021). Local*was*best: sourcing tree seed for future climates. *Canadian
728            Journal of Forest Research*, *51*(10), 1432–1439.

729    Polechová, J., & Barton, N. H. (2015). Limits to adaptation along environmental gradients. *Proceedings of
730            the National Academy of Sciences of the United States of America*, *112*(20), 6401–6406.

731    Rellstab, C., Dauphin, B., & Exposito-Alonso, M. (2021). Prospects and limitations of genomic offset in
732            conservation management. *Evolutionary Applications*, *14*(5), 1202–1212.

733    Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation. *Nature Reviews.
734            Genetics*, *14*(11), 807–820.

735    Schiffers, K., Schurr, F. M., Travis, J. M. J., Duputié, A., Eckhart, V. M., Lavergne, S., McInerny, G., Moore,
736            K. A., Pearman, P. B., Thuiller, W., Wüest, R. O., & Holt, R. D. (2014). Landscape structure and
737            genetic architecture jointly impact rates of niche evolution. *Ecography*, *37*(12), 1218–1229.

738    Siepielski, A. M., Gotanda, K. M., Morrissey, M. B., Diamond, S. E., DiBattista, J. D., & Carlson, S. M.
739            (2013). The spatial patterns of directional phenotypic selection. *Ecology Letters*, *16*(11), 1382–
740            1392.

741    Slatkin, M. (1973). Gene flow and selection in a cline. *Genetics*, *75*(4), 733–756.

742    Slatkin, M. (1978). Spatial patterns in the distributions of polygenic characters. *Journal of Theoretical
743            Biology*, *70*(2), 213–228.

744    Slatkin, M. (1993). Isolation by distance in equilibrium and non-equilibrium populations. *Evolution;
745            International Journal of Organic Evolution*, *47*(1), 264.

746    Smith, C. C. R., Tittes, S., Ralph, P. L., & Kern, A. D. (2023). Dispersal inference from population genetic
747            variation using a convolutional neural network. In *bioRxiv* (p. 2022.08.25.505329).
748            https://doi.org/10.1101/2022.08.25.505329

749    Urban, M. C. (2011). The evolution of species interactions across natural landscapes. *Ecology Letters*,
750            *14*(7), 723–732.

751    Wadgymar, S. M., DeMarche, M. L., Josephs, E. B., Sheth, S. N., & Anderson, J. T. (2022). Local
752            Adaptation: Causal Agents of Selection and Adaptive Trait Divergence. *Annual Review of
753            Ecology, Evolution, and Systematics*, *53*(1), 87–111.

754    Walsh, B., & Lynch, M. (2018). *Evolution and selection of quantitative traits*. Oxford University Press.

755    Wang, I. J., & Bradburd, G. S. (2014). Isolation by environment. *Molecular Ecology*, *23*(23), 5649–5662.

756    Wang, T., Hamann, A., Yanchuk, A., O'neill, G. A., & Aitken, S. N. (2006). Use of response functions in
757            selecting lodgepole pine populations for future climates. *Global Change Biology*, *12*(12), 2404–
758            2416.

759     Wang, T., Hamann, A., Spittlehouse, D., & Carroll, C. (2016). Locally downscaled and spatially
760            customizable climate data for historical and future periods for north America. *PloS One*, *11*(6),
761            e0156720.

762     Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*(3), 290–290.

763     Yeaman, S. (2013). Genomic rearrangements and the evolution of clusters of locally adaptive loci.
764            *Proceedings of the National Academy of Sciences of the United States of America*, *110*(19),
765            E1743-51.

766     Yeaman, S. (2015). Local Adaptation by Alleles of Small Effect. *The American Naturalist*, *186 Suppl 1*, S74-
767            89.

768     Yeaman, S., & Jarvis, A. (2006). Regional heterogeneity and gene flow maintain variance in a quantitative
769            trait within populations of lodgepole pine. *Proceedings. Biological Sciences*, *273*(1594), 1587–
770            1593.

771     Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation under migration-selection
772            balance. *Evolution; International Journal of Organic Evolution*, *65*(7), 1897–1911.

773     Ying, C. C., & Yanchuk, A. D. (2006). The development of British Columbia's tree seed transfer guidelines:
774            Purpose, concept, methodology, and implementation. *Forest Ecology and Management*, *227*(1),
775            1–13.

776     Zhou, X., & Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association studies.
777            *Nature Genetics*, *44*(7), 821–824.
778

**Supplementary Figure 1** Overall $F_{ST}$ (panel A) and isolation by distance (panel B) in simulated populations. Note the varying y-axes in panel B. In the main text, $F_{ST}$ is used to refer to the panels of individual graphs. In panel A values from 200 independent simulations were used to construct the boxplot and in panel B individual simulations are shown as lines. Weir and Cockerham's method for calculating $F_{ST}$, as implemented in the *sci-kit-allel* Python package, was used.

787  **Supplementary Figure 2** The average extent of local adaptation (panel A) and coefficient of variation in
788  local adaptation (panel B) as a function of spatial autocorrelation in the environment from simulated
789  datasets. The upper cell of each column is included in Figure 1 of the main text.

**Supplementary Figure 3** Comparison of local adaptation that evolves for two traits subject to spatially
varying selection. Selection on each trait was dictated by distinct maps of environmental
variation/phenotypic optima. The environment that exhibited the greater degree of spatial
autocorrelation (as measured by Moran's I) was designated "Environment 2". The 1:1 line is shown for
reference.

**Supplementary Figure 4** The effect of spatial autocorrelation in the environment on the correlations of
local environmental heterogeneity with local adaptation and additive genetic variance. A) The partial
correlation between local adaptation and local environmental heterogeneity, controlling for additive
genetic variance. B) The partial correlation between additive genetic variance and environmental
heterogeneity, controlling for local adaptation. Statistical significance was assessed after correcting for
multiple comparisons. The solid black line indicates the statistical null expectation of 0.

**Supplementary Figure 5** The distribution of locally adaptive effects as a function of spatial
autocorrelation in the environment. The area shown was calculated across 200 independent simulations
and smoothed using a LOESS regression with span 1.5. The upper right cell is included in Figure 2 of the
main text.

**Supplementary Figure 6** The relationship between allele frequency and the squared phenotypic effect size for polymorphisms that contribute varying degrees of local adaptation in either relative (panel A) or absolute terms (panel B).

814
815 **Supplementary Figure 7** Results from a GWAS on 1,000 randomly chosen individuals from either high or
816 low autocorrelation environments. Each panel compares the relative density of -log10($p$-values) from a
817 GWAS conducted on data from the 50 maps with the highest or lowest levels of spatial autocorrelation.
818

819    A                                                                   B

820    **Supplementary Figure 8** A) Cumulative local maladaptation as a function of spatial autocorrelation in
821    the environment across all parameter combinations. B) The average age of locally adaptive alleles in
822    meta-populations subject to spatially varying selection. The lines represent LOESS regression curves
823    calculated with span parameters of 1.5.
824

826 **Supplementary Figure 9** Establishment of local adaptation in the simulations. Panel A) shows the
827 average level of local adaptation across all demes. Panel B) shows the coefficient of variation in local
828 adaptation across demes. Panel C) shows the Kendall's tau rank correlation between phenotypes and
829 local optima.

830
831

**Supplementary Figure 10** Comparing local adaptation summary statistics to results of linear models applied to simulated provenance trials. Panel A compared the slopes of the relationship between relative fitness and transfer distance in simulated provenance trials to home-versus-away measure of local adaptation described by Blanquart et al., (2013). Panel B compares the variance in slopes across provenances to the variance in local adaptation across all populations. In both panels, each point summarises analyses from a single simulation. The Spearman correlation coefficient and the associated *p*-value are shown within each cell.

840
841

A)                                                                            B)

**Supplementary Figure 11** A) Percent variance explained by the principal component analysis conducted
on climatic/environmental variation in the Illingworth trial data. B) The correlation matrix for the 28
climatic/environmental variables for planting sites and provenances in the Illingworth trial. A key to the
abbreviations for the 25 annual climatic variables from ClimateBC along can be obtained from
https://climatebc.ca/Help2. Additionally, latitude (lat), longitude (Long) and elevation (Elev) are
included.

**Supplementary Figure 12** The spatial pattern of loadings onto the first 6 principal components of
environmental/climatic variation across provenances and planting sites in the Illingworth Trial. The first
6 principal components explained a total of 95% of the variation in the data.