Genotype imputation accuracy and the quality metrics of the minor ancestry in

multi-ancestry reference panels

Mingyang Shi,[1] Chizu Tanikawa,[2] Hans Markus Munter,[3] Masato Akiyama,[4] Satoshi

Koyama,[5] Kohei Tomizuka,[6] Koichi Matsuda,[2] Gregory Mark Lathrop,[3] Chikashi Terao,[6]

Masaru Koido,[1] and Yoichiro Kamatani[1]

[1] Laboratory of Complex Trait Genomics, Department of Computational Biology and

Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo,

Japan.

[2] Laboratory of Clinical Genome Sequencing, Department of Computational Biology and

Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo,

Japan.

[3] Department of Human Genetics, School of Biomedical Sciences, Faculty of Medicine

and Health Sciences, McGill University, Canada.

[4] Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu

University, Japan.

[5] Cardiovascular Disease Initiative, The Broad Institute of MIT and Harvard, Cambridge,

Massachusetts, US.

[6] Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative

Medical Sciences, Japan.

Correspondence: M.K. (mkoido@edu.k.u-tokyo.ac.jp) and Y.K.

(kamatani.yoichiro@edu.k.u-tokyo.ac.jp)

24   Tel: +81 (0)3-5449-5286

25   Fax: None

Abstract

Large-scale imputation reference panels are now available and have contributed to

efficient genome-wide association studies through genotype imputation. However, it is

still under debate whether large-size multi-ancestry or small-size population-specific

reference panels are the optimal choices for under-represented populations. We

imputed genotypes of East Asian (EAS; 180k Japanese) subjects using the Trans-

Omics for Precision Medicine (TOPMed) reference panel and found that the standard

imputation quality metric (Rsq) substantially overestimated the dosage r^2 (squared

correlation between imputed dosage and true genotype). Variance component analysis

of Rsq revealed that the increased imputed-genotype certainty (dosages closer to 0, 1,

or 2) caused upward bias, indicating some systemic bias in the imputation. Through

systematic simulations using different template switching rates (θ value) in the hidden

Markov model, we uncovered that the lower θ value increased the imputed-genotype

certainty and Rsq; however, dosage r^2 was insensitive to the θ value, thereby causing a

deviation. In simulated reference panels with different sizes and ancestral diversities,

the θ value estimates from Minimac decreased with the size of a single ancestry and

increased with the ancestral diversity. Thus, Rsq could overestimate or underestimate

dosage r^2 for a subpopulation in the multi-ancestry panel and the deviation represents

different imputed-dosage distributions. Finally, despite the impact of θ value, distant

ancestries in the reference panel contributed only a few additional variants passing a

predefined Rsq threshold. We conclude that the θ value has a substantial impact on the

imputed dosage and the imputation quality metric value.

Introduction

Genotype imputation is a cost-efficient technique to expand the number of markers in

genetic studies. Li and Stephen's hidden Markov model (HMM) and the pre-phasing-

imputation pipeline are employed by the popular tools Minimac, IMPUTE, and BEAGLE,

with slightly different implementations but similar accuracy [1–5].

The imputation accuracy (*e.g.*, the squared correlation between imputed dosage and

true genotype; dosage r^2) [6], the scores generated by software without the true

genotype (Rsq and INFO) [3,7,8], and the number of variants passing a predetermined

Rsq threshold (high-Rsq variants) are widely used quality metrics to evaluate imputation

performance [9]. The performance primarily depends on the reference panel size and

genetic similarity between the reference panel and target sample [10]. In studies using

the International HapMap Project (HapMap) and the 1000 Genomes Project (1KGP)

[11,12], pooling multiple ancestries together to maximize the panel size typically

improved the imputation performance compared to using only the matched ancestry

[8,13,14]. Deelen et al. combined the population-specific whole genome sequencing

(WGS) dataset of the Genome of the Netherlands (GoNL) with the 1KGP and showed

higher dosage r^2 than when using either GoNL or 1KGP alone [15]. Huang et al.

combined the UK10K with the 1KGP to generate more variants with high INFO [9].

Moreover, larger multi-ancestry reference panels, such as the Haplotype Reference

Consortium (HRC) and the Trans-Omics for Precision Medicine (TOPMed) [16,17],

greatly improved the imputation accuracy and increased the number of high-Rsq

variants in European (EUR), African (AFR), and Admixed American (AMR) [10,18].

However, constructing large reference panels by combining diverse ancestries is not

always beneficial. Small-size population-specific panels, such as Norwegian and

Estonian panels [19,20], achieve similar performance to the HRC panel in EUR.

Moreover, even though the HRC panel includes all 1KGP samples, the imputation

accuracy in non-EUR populations can be inferior to that of the 1KGP panel alone

[10,21]. Bai et al. found that adding 27 samples from a different ancestry to the Han

Chinese (1KGP-CHB, size = 103) resulted in better imputation accuracy than using the

1KGP-CHB or 1KGP panel [22]. In studies of Asian, combining population-specific WGS

datasets with the 1KGP had improved the imputation accuracy in some instances [23–

25], but not in others [26–28]. So far, the reason for these discrepancies is unclear.

Two sets of parameters are used in the HMM-based imputation algorithms. Transition

probability is governed by the template switching rate (θ) between adjacent markers,

which models the recombination and relatedness, and emission probability is

determined by the error rate (ε) for each marker, which models genotyping error, gene

conversion, and recurrent mutation [3,7]. These parameters are crucial to the HMM for

calculating matching probabilities between the template and target during the imputation

process, but dosage r^2 is robust to these parameters [3,5]. Many studies have

suggested different Rsq or INFO thresholds to achieve a similar dosage r^2 [29,30],

making it difficult to ascertain the relationship between the imputation process, Rsq, and

dosage r^2.

In this study, we imputed the East Asian (EAS) using the TOPMed reference panel

(EAS comprised 1.22% of the total samples) and found that Rsq substantially

overestimated dosage r^2. We introduced novel variance component analysis for Rsq

and analytically investigated why Rsq was overestimated and characterized the

relationship between the template switching rate used in the HMM, quality metrics, and

imputed dosage. Further, we evaluated the θ value, Rsq, dosage r^2, the deviation

between them, and the number of high-Rsq variants from matched or distant ancestry in

cases where the target ancestry was the major or minor component of the multi-

ancestry panels.

Material and methods

Subjects, genotyping, and quality control (QC)

Cohort specifications of Biobank Japan (BBJ-180k), genotyping, and QC are provided in

Supplementary Method 1. For imputation using the TOPMed panel, we lifted the

coordinates of the genotyping array from hg19 to hg38 using LiftOver [31]. Palindromic

variants (reference/alternative (ref/alt) alleles were G/C or A/T) and variants not in the

TOPMed freeze5b (https://bravo.sph.umich.edu/freeze5/hg38/) were excluded. The

ref/alt alleles were swapped and/or reverse-complemented according to the hg38

reference sequence. A total of 515,587 autosomal variants remained.

Pre-phasing, reference panel construction, and imputation

EAGLE v2.4.1 was used for pre-phasing without an external reference [32]. The 1KGP

reference panel was downloaded from https://genome.sph.umich.edu/wiki/Minimac3.

Methods used to construct the population-specific panels (named the BBJ1k and

JEWEL3k) are provided in **Supplementary Method 2**. We used an in-house server to

perform imputation using the 1KGP, BBJ1k, and JEWEL3k panels. Minimac3 v2.0.3

was used to estimate the HMM parameters and prepare m3vcf files.[3] Minimac4 v1.0.2

was used for imputation [33]. Imputation using the TOPMed and HRC panel were

performed using the TOPMed and Michigan imputation servers [3,17], respectively.

Evaluation of the imputation performance using different reference panels

Rsq from the Minimac4's info file was extracted for the subsequent analyses. Variants

with Rsq < 0.3 were removed [34]. Imputation performance was empirically evaluated

using 993 samples with WGS (named WGS$_{993}$). QC and processing are elaborated in

Supplementary Method 3. Coverage of a reference panel was the fraction of variants

in WGS$_{993}$ that could be imputed with Rsq ≥ 0.3. Dosage r^2 was the squared Pearson

correlation coefficient between the imputed dosage and true genotype (encoded as 0, 1,

or 2). Minor allele frequency (MAF) and minor allele count (MAC) of WGS_{993} were used

to stratify variants into bins.

Quantification of the deviation between Rsq and dosage r^2

We modeled the relationship between the imputed allelic dosage ($\boldsymbol{y} = (y_1, \dots, y_n)$, $y_i \in$

$[0,1]$) and the true allele ($\boldsymbol{x} = (x_1, \dots, x_n)$, $x_i \in \{0,1\}$) using the simple linear regression

formula:

143
$$\boldsymbol{y} = \beta_0 + \boldsymbol{x}\beta_{imp} + \boldsymbol{\varepsilon}$$

(Equation 1)

where β_{imp} is a scalar of the regression coefficient with $\hat{\beta}_{imp} = Cov(\boldsymbol{x}, \boldsymbol{y})/Var(\boldsymbol{x})$, β_0 is

the intercept, and $\boldsymbol{\varepsilon} = (\epsilon_1, \dots, \epsilon_n)$ is an error term. The Minimac's EmpRsq metric, which

is the squared Pearson correlation coefficient between $\boldsymbol{x}$ and $\boldsymbol{y}$

(https://genome.sph.umich.edu/wiki/Minimac3_Info_File), equals the ratio between the

regression sum of squares (SS_{reg}) and the total sum of squares (SS_{tot}) in this simple

linear regression. Then we have:

151
$$EmpRsq = \frac{Cov(\boldsymbol{x}, \boldsymbol{y})^2}{Var(\boldsymbol{x})Var(\boldsymbol{y})} = \frac{SS_{reg}}{SS_{tot}}$$

152 (Equation 2)

153 where $Cov(\boldsymbol{x}, \boldsymbol{y})$, $Var(\boldsymbol{x})$, and $Var(\boldsymbol{y})$ are the covariance between $\boldsymbol{x}$ and $\boldsymbol{y}$, the variance

154 of $\boldsymbol{x}$, and the variance of $\boldsymbol{y}$, respectively.

156   Rsq is the ratio between Var($\boldsymbol{y}$) and $p(1-p)$, where p is the alternative allele frequency

157   (AAF) in the imputed dataset [3,7]. Then,

$$Rsq = \frac{Var(\boldsymbol{y})}{p(1-p)} = \frac{SS_{tot}/n}{p(1-p)} = \frac{SS_{reg}/n}{p(1-p)} + \frac{SS_{res}/n}{p(1-p)}$$

159   (Equation 3)

160   where SS_{res} is the residual sum of squares that follows $SS_{tot} = SS_{reg} + SS_{res}$, and n is

161   the number of imputed haplotypes. Hence, Rsq comprises two parts: regression-related

162   and residual-related. We define $\frac{SS_{res}/n}{p(1-p)}$ as <u>M</u>AF-<u>A</u>djusted-<u>R</u>esidual-<u>E</u>rror (MARE). Then:

$$Rsq = \frac{SS_{reg}/n}{p(1-p)} + MARE = \frac{EmpRsq \times SS_{tot}/n}{p(1-p)} + MARE = \frac{EmpRsq \times Var(\boldsymbol{y})}{p(1-p)} + MARE$$

164   (Equation 4)

165   By assuming the equal AAF in $\boldsymbol{x}$ and $\boldsymbol{y}$, *i.e.*, $Var(\boldsymbol{x}) = p(1-p)$, Rsq could be further

166   treated as:

$$Rsq = \frac{Var(\boldsymbol{y})Cov(\boldsymbol{x},\boldsymbol{y})^2}{Var(\boldsymbol{x})Var(\boldsymbol{x})Var(\boldsymbol{y})} + MARE = \frac{Cov(\boldsymbol{x},\boldsymbol{y})^2}{Var(\boldsymbol{x})^2} + MARE = \hat{\beta}_{imp}^2 + MARE$$

168   (Equation 5)

169   Finally, *MARE* and $\hat{\beta}_{imp}$ could be obtained from Rsq and EmpRsq:

$$MARE = Rsq - \frac{Var(\boldsymbol{y})EmpRsq}{p(1-p)} = Rsq - Rsq \times EmpRsq = (1 - EmpRsq)Rsq$$

171   (Equation 6)

$$\hat{\beta}_{imp} = \sqrt{Rsq - (1 - EmpRsq)Rsq} = \sqrt{EmpRsq \times Rsq}$$

173   (Equation 7)

In Equation 7, $\hat{\beta}_{imp}$ is negative if $\boldsymbol{x}$ and $\boldsymbol{y}$ are negatively correlated. We do not consider

that situation. Hence, each combination of Rsq and EmpRsq indicates specific values of

MARE and $\hat{\beta}_{imp}$. Hereafter, we refer to $\hat{\beta}_{imp}$ as the β_{imp} metric and *MARE* as the MARE

metric.

Because the WGS dataset is unphased diploid data, dosage r^2 calculated from that will

be slightly different from EmpRsq. **Supplementary Note 1** provides the detailed

methods to calculate MARE and β_{imp} from haploid or diploid data, and the concordance

between values obtained using Equations 6–7 and calculated from the imputed dosage

in both haploid and diploid cases.

Imputation using the simulated 1KGP reference panels with different θ values

The 1KGP vcf files were downloaded from the Minimac3 website (see above).

Parameters were estimated using Minimac3. The θ value was extracted (denoted as

"Recom" in the m3vcf file) [3], scaled by 21 folds manually (0.01–100), and replaced in

the original file. A brief explanation of the θ value is provided in **Supplementary Note 2**.

The array data of WGS$_{993}$, a subset of BBJ-180k, was used as the target sample. The

imputation procedure was the same as described above.

Evaluation of the imputation performance using the simulated reference panels

We obtained the imputed allelic dosage (LooDosage) from Minimac4's empiricalDose

file (by turning the "--meta" option on) [35]. It was derived from the leave-one-out (LOO)

method by hiding the markers on the array during the imputation. Rsq, EmpRsq, MARE,

and β_{imp} were calculated from the LooDosage and array data. MAF was obtained from

the array data.

Simulation of reference panels and estimating the θ value

We sampled subsets from the 1KGP and JEWEL3k, shuffled the sample order 10 times,

and created new vcf files using bcftools v1.14 (https://samtools.github.io/bcftools/) [36].

We estimated the parameters using Minimac3. **Supplementary Method 4** provides the

detailed methods. The total θ value along chr19 (by summing the θ values between

adjacent markers) was used to evaluate the reference panel size and ancestral

diversity's impact. A discussion of the θ value qualification is provided in

Supplementary Note 3.

EUR-EAS reference panel simulation and imputation

We randomly sampled 403 individuals (named 1KGP-EUR$_{403}$) from the 1KGP-EUR and

combined them with the 1KGP-EAS and 6 subsets (size = 500, 1,000, 1,500, 2,000,

2,500, and 3,256) of the 3,256 JPT WGS samples (named as JPT$_{3256}$) in JEWEL3k.

The remaining 100 individuals in the 1KGP-EUR were used as the target sample. We

extracted the 10,375 polymorphic variants (chr19) on the Illumina Global Screening

Array v3.0 (https://support.illumina.com/content/dam/illumina-

support/documents/downloads/productfiles/global-screening-array-24/v3-0/infinium-

global-screening-array-24-v3-0-a1-b151-rsids.zip) to simulate the genotyping array.

Parameter estimation and imputation were the same as above.

JPT-1KGP reference panel simulation and imputation

221 We sampled 7 subsets (size = 100, 500, 1,000, 1,500, 2,000, 2,500, and 3,256) from the

222 JPT_{3256} and then combined the JPT_{3256} with 6 subsets (1KGP-JPT and 1–5 ancestries)

223 of the 1KGP. WGS_{993} was used as the target. The other processing methods were the

224 same as above.

Number of confident alleles and high-Rsq variants

Haploid dosage (HDS) is the imputed alternative allele's allelic dosage at the haploid

level, which could be obtained by the "--format HDS" option in Minimac4. Briefly, it is in

a per-variant and per-individual manner and a higher value represents an imputed

alternative allele is more certain. We quantified the θ value's impact on the imputed

dosages by the number of confident alleles (HDS > 0.9). Rsq was used to judge how

many high-Rsq (Rsq > 0.7) variants could be passed to the downstream analyses.

Furthermore, we determined how many confident alleles and high-Rsq variants could be

obtained only from the additional distant ancestries in the multi-ancestry reference

panel. We defined ancestry-specific variants as follows: EUR-only variants only existed

in the $1KGP\text{-}EUR_{403}$, and it has no non-EUR variants. JPT_{3256}-only and 1KGP-EAS-only

variants only existed in the JPT_{3256} and 1KGP-EAS, respectively. Non-EAS variants

were not found in the JPT_{3256} or 1KGP-EAS.

Replication of the TOPMed imputation pipeline

We followed the TOPMed imputation pipeline (Accessed on Oct. 15, 2022,

https://topmedimpute.readthedocs.io/) and used the 1KGP and JEWEL3k reference

panels. Specifically, the HapMap2 genetic map was used as a reference for the θ value

instead of estimating it using Minimac3. WGS_{993} was used as the target. As the θ value

was not outputted by default, we modified the Minimac4's source code to obtain the

transformed θ value (**Supplementary Method 5**).

Results

BBJ imputation using the four reference panels

The BBJ-180k was imputed using the TOPMed, 1KGP, BBJ1k, and JEWEL3k reference

panels (**Figure 1A**). Characteristics of each panel and the target sample are listed in

Table 1. We categorized the imputed variants by MAF and Rsq in each imputed

dataset. With more EAS samples in the panel, more variants with low MAF passed each

Rsq threshold (**Figure 2A** and **Supplementary Table 1**). In addition to the absolute

number, unique variants were imputed from each panel (**Figure 2B**). These results

reproduced the benefits of using large and different reference panels [19].

We then used WGS$_{993}$ to empirically evaluate the imputation performance. MAF and

MAC of WGS$_{993}$ were used to categorize the variants into six bins: common (MAF ≥

5%), low-frequency (5% > MAF ≥ 1% and 1% > MAF ≥ 0.5%), rare (0.5% > MAF >

0.1%), doubleton (MAC = 2), and singleton (MAC = 1). For common single nucleotide

polymorphisms (SNVs), the TOPMed imputation result showed the highest coverage

(0.950), whereas BBJ1k (0.890) and JEWEL3k (0.907) were lower than 1KGP (0.948).

This loss of variants was possibly due to the additional QC steps in combining the

1KGP with Japanese WGS (**Supplementary Method 2**). As MAF decreased, the

coverage also decreased; however, more EAS samples in the reference panel mitigated

this decrease, as expected (**Figure 2C**).

The 155,297 shared SNVs (on chr19) imputed by the four panels were used to compare

imputation accuracy. The mean dosage r^2 using the TOPMed panel was between those

of 1KGP and BBJ1k (**Figure 2D**). However, Rsq was substantially upward biased only

in the TOPMed imputation result (**Figure 2E**). Using all SNVs and short insertions and

deletions (indels) with Rsq ≥ 0.3, although with higher coverage, the TOPMed

imputation result showed an inferior dosage r^2 than 1KGP (**Supplementary Figure 1**).

Quantifying the deviation between Rsq and dosage r^2

The deviation between Rsq and dosage r^2 was persistent in all MAF bins only when

using the TOPMed panel (**Supplementary Figure 2**), which indicated a potential

systematic bias from the imputation pipeline or the reference panel. To justify our

observation, we analytically derived the relationship between Rsq and dosage r^2

(**Methods**). Two novel metrics, MARE and β_{imp}, were introduced to quantify the

deviation. MARE, a MAF-adjusted form of residual error, takes a value between 0 and

1, and increases with SS_{res}. β_{imp} describes the distinguishability between the mean

imputed dosage of each true genotype group. Rsq is the ratio between observed and

expected variance and dosage r^2 shows the correlation. Under the assumption of "well-

calibration" (the posterior allele probability from imputation equals the expected true

allele dose), Rsq equals dosage r^2 [10]. Our analytical derivation of the relationship

between Rsq and dosage r^2 did not assume the "well-calibration" (**Supplementary**

Figure 3–7 and **Supplementary Note 1**). Any two of Rsq, dosage r^2, MARE, and β_{imp}

could entirely quantify the relationship between the imputed dosage and true genotype

and determine the other two metrics (Equations 6–7), whereas Rsq or dosage r^2 alone

could not.

To exhibit the relationships, we plotted the theoretical values of MARE and β_{imp} on the

coordinates of Rsq and dosage r^2 (**Figure 3A** and **3F**). It clearly showed that the

overestimated Rsq was accompanied by a higher MARE (**Figure 3A**). We used

rs142572000 as an example (**Figure 3B–E**). In the TOPMed imputation result, imputed

genotypes were more certain (defined as the imputed dosage closer to 0, 1, or 2)

(**Figure 3B**) [37], compared to the other three panels (**Figure 3C–E**). The high certainty

increased Var(**y**) and Rsq. In **Supplementary Note 4**, the positive relationship between

imputed-genotype certainty and Rsq is demonstrated. As discussed further below, high

certainty or Rsq did not mean the imputation is more accurate. As shown in **Figure 3B**,

many heterozygotes were incorrectly imputed with a dosage of approximately 0 in the

TOPMed imputation, causing higher MARE and Rsq, and an even low dosage r^2.

Variants with dosage r^2 < Rsq showed a lower β_{imp} (**Figure 3F**). We used rs671 as

another example (**Figure 3G–J**). Except for the TOPMed imputation result, imputed

dosages were shrunk to the AAF in EAS (0.247) (**Figure 3H–J**), causing uncertainty in

the imputed genotype, and decreased β_{imp}, MARE, and Rsq. However, imputed

dosages were still highly correlated with the true genotypes (**Figure 3F**), causing

dosage r^2 > Rsq. Taking the two examples together, imputation results of a similar Rsq

or dosage r^2 may have a great difference in the imputed dosage, thus, more

comprehensive evaluations are necessary and the deviation between Rsq and dosage

r^2 should not be ignored.

We categorized MARE into Rsq bins and β_{imp} into dosage r^2 bins to compare them

between reference panels and to the expected values when assuming Rsq equals

dosage r^2. The mean MARE of the TOPMed result was above the expected for Rsq bins

0.35–0.9, and the mean β_{imp} of the 1KGP result was below those of the other panels

(**Supplementary Figure 8**). These results suggested that the TOPMed result was more

certain and might contain more wrongly imputed genotypes (**Figure 3B** and **3G**), while

the 1KGP result might possess higher shrinkage in the imputed dosage, as shown

above (**Figure 3C** and **3H**).

Template switching rate impacts the deviation between Rsq and dosage r^2

The high certainty suggested an overconfident matching between the reference panel

and target sample. We investigated how the template switching rate (θ) used in the

HMM affects the imputed-allele certainty using the 1KGP reference panel and 21

scalings of the θ value (0.01–100-fold of that estimated by Minimac3; **Methods**; **Figure**

1B). Low θ values caused Rsq > EmpRsq (EmpRsq is the alternate of dosage r^2;

Methods) (**Supplementary Figure 9A–C**), while high θ values caused Rsq < EmpRsq

(**Supplementary Figure 9E–G**). Using rs10410162 as an example, Rsq and MARE

increased as the θ value decreased (**Figure 4A**), with increasing certainty of the

imputed alleles (**Supplementary Figure 10A–C**) and the deviation toward Rsq >

EmpRsq (**Figure 4B**). As the θ value increased, the imputed allelic dosages were

shrunk to the AAF (0.351) (**Supplementary Figure 10E–G)**, resulting in a more drastic

decrease in Rsq than EmpRsq (**Figure 4A**), thereby causing EmpRsq > Rsq. Notably,

EmpRsq and β_{imp} were roughly maintained unless the θ value was scaled up by 2-fold

or higher (**Figure 4A** and **4C**), suggesting EmpRsq was insensitive to altering the θ

value, particularly the downscaling. In contrast, Rsq was sensitive to the θ value and

consistently increased with downscaling of the θ value, leading to the deviation between

Rsq and EmpRsq.

We also showed that low θ values increased the imputed-allele certainty using a

randomly selected target haplotype (**Figure 4D** and **Supplementary Figure 11).** High

certainty might cause some variants to be wrongly imputed as the opposite allele

(**Supplementary Figure 12**). An explanation is provided in **Supplementary Note 5**.

Such imputed-dosage properties were also revealed by the mean MARE and β_{imp}

across all variants on chr19 (**Supplementary Figure 13**). Taken together, the simulated

imputation using an extremely low θ value mimics the high certainty and Rsq

overestimation in the TOPMed imputation result in EAS (**Figure 3**).

Template switching rate impacts the imputation performance

Using the modified 1KGP panels, although the mean EmpRsq was insensitive to the

scaling of the θ value (maximum difference of 0.034 for the θ value scaling between

0.01 and 2-fold; **Figure 5**), low θ value increased Rsq (**Figure 5**). We evaluated the

number of confident alleles (HDS > 0.9) and high-Rsq variants (Rsq > 0.7) (**Methods**).

Downscaling the θ value increased the number of confident alleles and high-Rsq

variants (**Supplementary Table 2**). When the θ value was 0.5-fold, the number of

confident alleles and high-Rsq variants increased by 5.46% and 8.89%, respectively,

and if the θ value was 2-fold, these numbers decreased by 11.8% and 13.8%,

respectively. These results indicated that the θ value shaped the imputed allelic dosage

and changed Rsq and the number of high-Rsq variants while leaving EmpRsq almost

the same.

Reference panel and θ estimates

We evaluated how Minimac3's parameter estimation changed with the composition of

the reference panel (**Supplementary Method 4**, **Supplementary Figure 14**, and

Supplementary Note 6). The θ estimates decreased with the sample size of a single

ancestry and increased with ancestral diversity when size was fixed (**Figure 6A** and

6B). On pooling samples from different ancestries together at a small panel size, the θ

estimates still decreased as the panel size increased (**Figure 6C**); however, at a larger

panel size, it increased with simultaneously increasing the panel size and ancestral

diversity (**Figure 6D**), suggesting that the θ estimates were in a trade-off between the

panel size and ancestral diversity (**Figure 6E**). **Supplementary Note 6** explains these

effects.

Fitness of the θ value, deviation, and imputation performance

To elucidate the multi-ancestry reference panel's impact on the imputation result, we

simulated two scenarios using the multi-ancestry reference panels: the target sample

was from the (1) minor and (2) major ancestry (**Figure 1C**).

Scenario 1: The target sample was from the minor ancestry

We simulated 8 EUR-EAS reference panels and used 100 EUR samples as the target

(**Methods**). As the panel size increased and θ value decreased (**Table 2**), Rsq and

MARE were upwardly biased, as expected (**Supplementary Figure 15** and **S16**). The

mean EmpRsq decreased consistently with the addition of EAS samples to the

reference panel (**Table 2**). However, only a marginal difference was observed

(maximum difference of 0.020).

There were 114,606 EUR-only and 696,017 non-EUR variants (**Methods**). As the panel

size increased and θ value decreased, the number of confident alleles increased from

84,797 to 101,116 and from 0 to 458 for EUR-only and non-EUR variants, respectively

(**Supplementary Table 3**). The number of high-Rsq variants increased from 163,468 to

200,161, 18,646 to 27,941, and 0 to 521 for all, EUR-only, and non-EUR variants,

respectively (**Supplementary Table 3**). These results revealed that a lower θ value

increases the number of confident alleles and high-Rsq variants, without improving the

EmpRsq (**Supplementary Table 3**). Meanwhile, even if the reference panel comprised

90.3% EAS samples, non-EUR variants only comprised 0.26% of the total variants in

the imputation result (when setting a cutoff of Rsq > 0.7). The majority of variants

gaining from the larger panel when imputing the under-represented ancestry was

because of the lower θ value.

Scenario 2: The target sample was from the major ancestry

We simulated 13 JPT-1KGP reference panels and used WGS_{993} as the target

(**Methods**). When only JPT samples were in the panel, the mean EmpRsq and Rsq

increased with the panel size, as expected (**Supplementary Figure 17**). When

combined with the 1KGP subsets, the mean EmpRsq was highest when using

JPT_{3256}+1KGP-EAS for variants with MAF ≥ 1% and JPT_{3256}+1KGP-JPT for variants

with 1% > MAF ≥ 0.5% (**Table 3**). Adding other ancestries decreased the mean

EmpRsq marginally (maximum difference < 0.01 for all MAF categories). The mean Rsq

was the highest when using JPT_{3256} and decreased with the addition of more

ancestries, with maximum differences of 0.007, 0.024, and 0.048 for variants with MAF

≥ 5%, 5% > MAF ≥ 1%, and 1% > MAF ≥ 0.5%, respectively (**Table 3**). None of the

imputation results showed a noticeable deviation in the MARE and $β_{imp}$. However, the

MARE and $β_{imp}$ of the JPT_{3256}, JPT_{3256}+1KGP-JPT, and JPT_{3256}+1KGP-EAS results

were closer to the expected values (**Supplementary Figure 18**), while adding other

ancestries made MARE and $β_{imp}$ correspond to the condition of using a higher θ value.

There were 74,490 JPT_{3256}-only, 7,627 1KGP-EAS-only, and 495,489 non-EAS variants

(**Methods**). From JPT_{3256} to JPT_{3256}+1KGP, the number of confident alleles decreased

from 24,599 to 23,666, increased from 0 to 131, and increased from 0 to 254 for these

three groups of variants (**Supplementary Table 4**). The number of high-Rsq variants

decreased from 274,343 to 264,221 (a decrease of 3.69%) when using JPT$_{3256}$ or

JPT$_{3256}$+1KGP (**Supplementary Table 4**). Only 265 non-EAS variants reached an

Rsq > 0.7 when using JPT$_{3256}$+1KGP, which was 0.10% of the total variants. Hence,

combining with the 1KGP caused fewer confident alleles and high-Rsq variants to

remain in the imputation results, as expected from using higher θ values. Both

scenarios suggest that the distant ancestry in the reference panel affects the θ

estimates and the number of high-Rsq variants. However, the expanded haplotypes and

variant sets from distant ancestries provided only a few additional variants.

EAS imputation using the public reference panels

We used the TOPMed imputation pipeline (**Methods**) and observed an upward

deviation of Rsq, particularly when using the 1KGP panel (**Supplementary Figure 19**).

The θ value transformed from the genetic map was 0.15- and 0.27-fold of that estimated

by Minimac3 when using the 1KGP and JEWEL3k panels, respectively

(**Supplementary Table 5**).

As the θ estimates decreased with the size of the major ancestry (**Figure 6C**), it would

be underestimated for the minor ancestry in the multi-ancestry panel regardless of using

the genetic map or estimating it by the Mimimac3. One example was the HRC panel,

which did not use the genetic map but induced Rsq overestimation in the BBJ-180k

(**Supplementary Figure 20**). Thus, caution may be required when imputing the under-

represented population using large ancestry-imbalanced panels under the current

framework.

Discussion

The deviation between Rsq and dosage r^2 in the imputation result is raised from the

imputed dosage, and has been widely observed using different reference panels and

software [9,15,38]. One reason was that the θ value used in the HMM does not fit the

panel size, ancestral components in the reference panel, and the target population.

When using the multi-ancestry reference panel, distant ancestries affect the θ estimates

(in Minimac3/4), then the imputed dosage and Rsq. Our simulations indicated that the

lower θ value used by the larger panel could increase the imputed-genotype certainty

and the number of high-Rsq variants but not dosage r^2. It may cause confusing

benchmarking results and increase the chance of false positives in association tests.

Ferwerda et al. reported height and body mass index association signals in an ethnically

diverse cohort only when imputing against the TOPMed panel [39]. Bai et al. reported

the highest Rsq but lowest dosage r^2 when imputing the Han Chinese using the HRC

panel, compared to the 1KGP and a population-specific panel [22], similar to what we

had observed in Japanese. In addition to simply comparing dosage r^2 or Rsq, we

provided the changes in those metrics, the θ estimates, imputed-dosage certainty, and

the number of high-Rsq variants passing to downstream analyses. We provided the

script (https://github.com/shimaomao26/impumetric) to allow users to check their results

using the leave-one-out imputation of Minimac4 (external WGS not required) or

additional WGS data.

The TOPMed imputation pipeline used the θ value transformed from the HapMap2

genetic map by assuming 0.01 centimorgan (cM) corresponds 1% switching rate. The

switching rate would thereby be fixed given the variant's base pair position and cM. Two

reasons may explain why it works well. First, the genetic map did not significantly

change the imputation accuracy, as verified in a Finnish study [40]. We had also shown

that dosage r^2 was insensitive to the θ value. Second, the θ value was only

underestimated for the EAS (size = 1,184), but the sizes of EUR, AFR, and AMR (size =

17,085–47,159) might fit this value. Previous studies reported that Rsq in the TOPMed

imputation result was sometimes misleadingly high in AFR [38] and even EUR [41]. A

recent paper reported that the TOPMed panel was more robust to the low-density

genotyping array than the HRC and 1KGP panels [18]. These results also suggested

that a fixed low θ value might be used by the TOPMed pipeline, as we have inferred.

Further studies are warranted to comprehensively elucidate the impacts on genetic

studies.

Our results suggest that avoiding ancestral diversity is best when more than 3,000 WGS

samples are available to construct a JPT reference panel. Increased diversity would

then only have a marginal impact on dosage r^2 but would affect the individual's imputed

dosage and cause fewer variants to pass a predetermined Rsq filter. Zhang et al. and

Cong et al. similarly observed that adding the 1KGP to about 3,000 Chinese WGS

samples would neither benefit nor harm dosage r^2 [26,27]. We focused on explaining

the reason in this work. Further studies are warranted to take advantage of combining

population-specific and public WGS datasets while avoiding our identified problems.

Our study has several limitations. First, our simulations of the reference panel were

study-specific. As discussed, the θ estimates decreased with the panel size and

increased with the ancestral diversity. However, a larger panel typically increases the

diversity simultaneously. Therefore, the θ value and imputation result need to be

discussed case by case. This limitation also implies that the general experience may not

work well for a new reference panel and target dataset. Second, in the simulations of

multi-ancestry reference panels (scenarios 1 and 2), WGS datasets were merged using

IMPUTE2 [9]. We did not investigate the impact of IMPUTE2 but treated the merged

dataset similarly to the WGS dataset. It might underestimate the number of high-Rsq

variants from the distant ancestry (**Supplementary Note 7)**. Our results using the 1KGP

(not modified by IMPUTE2) also revealed that only a limited number of confident alleles

and high-Rsq variants in the imputation result were from the distant ancestry

(**Supplementary Note 7** and **Supplementary Table 2**). Therefore, the conclusion that

distant ancestry in the reference panel affects the θ estimates rather than providing

high-Rsq variants would be valid. However, further studies should be conducted to

determine the impact of the panel-merging method and the net gain of variants from

distant ancestries.

In summary, we explained that the HMM parameter could be a potential reason for

inaccurate Rsq and inferior dosage r^2 when using large multi-ancestry reference panels.

This is also the first study of the relationship between the template switching rate,

imputed-genotype certainty, Rsq, and dosage r^2. We envision that our methods and

conclusions could provide insights into benchmarking studies, construction of reference

panels, and development of imputation algorithms and pipelines in the future.

Funds

This research was supported by the Ministry of Education, Culture, Sports, Sciences

and Technology (MEXT) of Japanese government and the Japan Agency for Medical

Research and Development (AMED) under grant numbers JP18km0605001 (the

BioBank Japan project) and JP19km0405215 (to C.T., K.M., and Y.K.).

Acknowledgments

We thank the staff of the BBJ for collecting and managing samples and clinical

information. We acknowledged the Human Genome Center, the Institute of Medical

Science, the University of Tokyo (http://sc.hgc.jp/shirokane.html) for providing the

super-computing resources.

Web resources

Impumetric, https://github.com/shimaomao26/impumetric

TOPMed imputation server, https://imputation.biodatacatalyst.nhlbi.nih.gov/#!

Michigan Imputation server, https://imputationserver.sph.umich.edu/index.html

Minimac3, https://genome.sph.umich.edu/wiki/Minimac3

Minimac4, https://genome.sph.umich.edu/wiki/Minimac4

EAGLE, https://alkesgroup.broadinstitute.org/Eagle/

Bcftools, https://samtools.github.io/bcftools/

LiftOver, https://genome.ucsc.edu/cgi-bin/hgLiftOver

Data and code availability

We provided the tool to replicate the metrics used in this work

(https://github.com/shimaomao26/impumetric). Other scripts were also deposited there.

Genotype data for BBJ and the TOPMed imputation results were deposited at NBDC

Human Database (research ID: hum0014 and hum0311, respectively).

Appendix

Imputation quality metrics

Rsq: The standard quality metric in Minimac. The ratio between observed variance and

expected variance of imputed allelic dosage.

INFO: The standard quality metric in IMPUTE. The ratio between observed information

and complete information of imputed genotype distribution.

Dosage r^2: The squared Pearson correlation between imputed dosage and true

genotype.

EmpRsq: The squared Pearson correlation between imputed allelic dosage and true

allele dose. Provided by Minimac and is only available for the genotyped markers on

array.

MARE: The minor allele frequency adjusted residual error in linear regression between

imputed dosage and true genotype.

β_{imp}: The regression coefficient between imputed dosage and true genotype.

References

1. Li N, Stephens M. Modeling linkage disequilibrium and identifying recombination

hotspots using single-nucleotide polymorphism data. Genetics 2003; 165:2213–2233

2. Howie B, Fuchsberger C, Stephens M, et al. Fast and accurate genotype imputation

in genome-wide association studies through pre-phasing. Nat. Genet. 2012; 44:955–

959

3. Das S, Forer L, Schönherr S, et al. Next-generation genotype imputation service and

methods. Nat. Genet. 2016; 48:1284–1287

4. Howie BN, Donnelly P, Marchini J. A Flexible and Accurate Genotype Imputation

Method for the Next Generation of Genome-Wide Association Studies. PLOS Genet.

2009; 5:e1000529

5. Browning BL, Browning SR. Genotype Imputation with Millions of Reference

Samples. Am. J. Hum. Genet. 2016; 98:116–126

6. Li Y, Sidore C, Kang HM, et al. Low-coverage sequencing: Implications for design of

complex trait association studies. Genome Res. 2011; 21:940–951

7. Li Y, Willer CJ, Ding J, et al. MaCH: using sequence and genotype data to estimate

haplotypes and unobserved genotypes. Genet. Epidemiol. 2010; 34:816–834

8. Marchini J, Howie B. Genotype imputation for genome-wide association studies. Nat.

Rev. Genet. 2010; 11:499–511

9. Huang J, Howie B, McCarthy S, et al. Improved imputation of low-frequency and rare

variants using the UK10K haplotype reference panel. Nat. Commun. 2015; 6:8111

10. Das S, Abecasis GR, Browning BL. Genotype Imputation from Large Reference

Panels. Annu. Rev. Genomics Hum. Genet. 2018; 19:73–96

11. Frazer KA, Ballinger DG, Cox DR, et al. A second generation human haplotype map

of over 3.1 million SNPs. Nature 2007; 449:851–861

12. Auton A, Abecasis GR, Altshuler DM, et al. A global reference for human genetic

variation. Nature 2015; 526:68–74

13. Huang L, Li Y, Singleton AB, et al. Genotype-imputation accuracy across worldwide

human populations. Am. J. Hum. Genet. 2008; 84:235–250

14. Liu Q, Cirulli ET, Han Y, et al. Systematic assessment of imputation performance

using the 1000 Genomes reference panels. Brief. Bioinform. 2015; 16:549–562

15. Deelen P, Menelaou A, Van Leeuwen EM, et al. Improved imputation quality of low-

frequency and rare variants in European samples using the 'Genome of the

Netherlands'. Eur. J. Hum. Genet. 2014; 22:1321–1326

16. McCarthy S, Das S, Kretzschmar W, et al. A reference panel of 64,976 haplotypes

for genotype imputation. Nat. Genet. 2016; 48:1279–1283

17. Taliun D, Harris DN, Kessler MD, et al. Sequencing of 53,831 diverse genomes from

the NHLBI TOPMed Program. Nature 2021; 590:290–299

18. Hanks SC, Forer L, Schönherr S, et al. Extent to which array genotyping and

imputation with large reference panels approximate deep whole-genome sequencing.

Am. J. Hum. Genet. 2022; 109:1653–1666

19. Zhou W, Fritsche LG, Das S, et al. Improving power of association tests using

multiple sets of imputed genotypes from distributed reference panels. Genet. Epidemiol.

2017; 41:744–755

20. Mitt M, Kals M, Pärn K, et al. Improved imputation accuracy of rare and low-

frequency variants using population-specific high-coverage WGS-based imputation

reference panel. Eur. J. Hum. Genet. 2017; 25:869–876

21. Lin Y, Liu L, Yang S, et al. Genotype imputation for Han Chinese population using

Haplotype Reference Consortium as reference. Hum. Genet. 2018; 137:431–436

22. Bai WY, Zhu XW, Cong PK, et al. Genotype imputation and reference panel: a

systematic evaluation on haplotype size and diversity. Brief. Bioinform. 2020; 21:1806–

1817

23. Akiyama M, Ishigaki K, Sakaue S, et al. Characterizing rare and low-frequency

height-associated variants in the Japanese population. Nat. Commun. 2019; 10:4393

24. Jeon S, Bhak Y, Choi Y, et al. Korean Genome Project: 1094 Korean personal

genomes with clinical information. Sci. Adv. 2020; 6:eaaz7835

25. Yoo S-K, Kim C-U, Kim HL, et al. NARD: whole-genome reference panel of 1779

Northeast Asians improves imputation accuracy of rare and low-frequency variants.

Genome Med. 2019; 11:64

26. Cong P-K, Bai W-Y, Li J-C, et al. Genomic analyses of 10,376 individuals in the

Westlake BioBank for Chinese (WBBC) pilot project. Nat. Commun. 2022; 13:2939

27. Zhang P, Luo H, Li Y, et al. NyuWa Genome resource: A deep whole-genome

sequencing-based variation profile and reference panel for the Chinese population. Cell

Rep. 2021; 37:110017

28. Kawai Y, Mimori T, Kojima K, et al. Japonica array: improved genotype imputation

by designing a population-specific SNP array with 1070 Japanese individuals. J. Hum.

Genet. 2015; 60:581–587

29. Pistis G, Porcu E, Vrieze SI, et al. Rare variant genotype imputation with thousands

of study-specific whole-genome sequences: implications for cost-effective study

designs. Eur. J. Hum. Genet. 2015; 23:975–983

30. Roshyara NR, Horn K, Kirsten H, et al. Comparing performance of modern genotype

imputation methods in different ethnicities. Sci. Rep. 2016; 6:34386

31. Kent WJ, Sugnet CW, Furey TS, et al. The Human Genome Browser at UCSC.

Genome Res. 2002; 12:996–1006

32. Loh P-R, Danecek P, Palamara PF, et al. Reference-based phasing using the

Haplotype Reference Consortium panel. Nat. Genet. 2016; 48:1443–1448

33. Das S. Next Generation of Genotype Imputation Methods. 2017;

34. Liu EY, Buyske S, Aragaki AK, et al. Genotype Imputation of Metabochip SNPs

Using a Study-Specific Reference Panel of ~4,000 haplotypes in African Americans

From the women's health initiative. Genet. Epidemiol. 2012; 36:107–117

35. Yu K, Das S, LeFaive J, et al. Meta-imputation: An efficient method to combine

genotype data after imputation with multiple reference panels. Am. J. Hum. Genet.

2022; 109:1007–1015

36. Li H. A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data.

Bioinformatics 2011; 27:2987–2993

37. Marchini J, Howie B, Myers S, et al. A new multipoint method for genome-wide

association studies by imputation of genotypes. Nat. Genet. 2007; 39:906–913

38. Sun Q, Yang Y, Rosen JD, et al. MagicalRsq: Machine-learning-based genotype

imputation quality calibration. Am. J. Hum. Genet. 2022; 109:1986–1997

39. Ferwerda B, Abdellaoui A, Nieuwdorp M, et al. A Genetic Map of the Modern Urban

Society of Amsterdam. Front. Genet. 2021; 12:

40. Hassan S, Surakka I, Taskinen M-R, et al. High-resolution population-specific

recombination rates and their effect on phasing and genotype imputation. Eur. J. Hum.

Genet. 2021; 29:615–624

41. Sun Q, Liu W, Rosen JD, et al. Leveraging TOPMed imputation server and

constructing a cohort-specific imputation reference panel to enhance genotype

imputation among cystic fibrosis patients. Hum. Genet. Genomics Adv. 2022; 3:100090

Figures

Figure 1. Overview of this study. (A) Imputation using the TOPMed, 1KGP, BBJ1k (1,037 Japanese (JPT) WGS + 1KGP), and JEWEL3k (3,256 JPT WGS + 1KGP) reference panels. The 180k East Asian (EAS) samples in the Biobank Japan (BBJ) first cohort were imputed against the four panels. Samples with whole genome sequencing (WGS_{993}) were used to empirically evaluate the imputation performance, including a count of imputed variants and estimated (Rsq) and true (dosage r^2) imputation accuracy. The deviation between Rsq and dosage r^2 was examined and quantified analytically. Two novel metrics, MAF-adjusted-residual-error (MARE) and β_{imp}, were introduced to indicate the different imputed-dosage distributions underlying the

deviation. (B) Simulations to evaluate the relationship between the hidden Markov

model (HMM) parameter, Rsq, dosage r^2, the deviation between Rsq and dosage r^2,

and imputed dosage. The reference panel and target sample were fixed to the 1KGP

and WGS$_{993}$, and 21 scaled template switching rates (the θ value) were used for

running imputations. Evaluations were the same as in (A). (C) Simulations to evaluate

the relationship between the reference panel, HMM parameter, imputation metrics, and

imputation result. Two scenarios were simulated to represent the imputations of the

major and minor ancestry in a multi-ancestry panel. The 3,256 JPT WGS and 1KGP

were used to simulate reference panels with different sizes and ancestral diversities.

European (EUR) and JPT samples were used as the target. Imputation results were

evaluated the same as in (B). In addition, we evaluated how the Minimac3's parameter

estimates change with the panel composition.

Figure 2. Imputation results using the TOPMed, 1KGP, BBJ1k, and JEWEL3k

reference panels. (A) Count of imputed variants using the four panels, stratified by

minor allele frequency (MAF) and Rsq in each imputation result. (B) Count of unique

and shared variants (with Rsq ≥ 0.3) using the TOPMed, 1KGP, and JEWEL3k panels.

(C) Fraction of SNV and indel in WGS_{993} that could be imputed with Rsq ≥ 0.3

(coverage) using each panel, stratified by MAF of WGS_{993}. (D) The mean dosage r^2 and

Rsq, stratified by MAF of WGS_{993}. (E) Comparison of dosage r^2 and Rsq in the

imputation results using the four panels. The scatter represents the variants, with the

blue lines showing the density. The red line shows that Rsq equals dosage r^2, and the

histograms on the side show the distribution. In (C), (D), and (E), variants on

chromosome 19 were used. In (D) and (E), overlapping variants imputed by the four

panels (Rsq ≥ 0.3 using all panels) were used.

Figure 3. MARE, β_{imp}, Rsq, dosage r², and imputed dosage of rs142572000 and rs671. (A) The plot shows Rsq and dosage r² of rs142572000 and rs671, and the theoretical value of MARE on the Rsq ~ dosage r² plot. (B–E) The plot shows the imputed dosage of rs142572000 in the imputation result using the (B) TOPMed, (C) 1KGP, (D) BBJ1k, and (E) JEWEL3k reference panels. (F) The plot shows the theoretical value of β_{imp} on the Rsq ~ dosage r² plot. (G–J) The plot shows the imputed dosage of rs671 in the imputation result using the (G) TOPMed, (H) 1KGP, (I) BBJ1k, and (J) JEWEL3k panels. In (A) and (F), the red line shows that Rsq equals dosage r², and the color keys show the theoretical MARE or β_{imp} values. The dashed lines represent the contours of MARE or β_{imp}, and adjacent contours have a value difference of 0.05. The red dashed arrow highlights the difference in MARE and β_{imp} (crossing how many contour lines) for variants showing a similar Rsq but different dosage r²

(rs142572000), or similar dosage r^2 but different Rsq (rs671). In (B–E) and (G–J), the

strip plot shows the individual and the violin plot shows the distribution. The red line

shows the linear regression line between the imputed dosage and true genotype. In (B–

E), MARE is 0.473, 0.274, 0.328, and 0.187, and β_{imp} is 0.336, 0.097, 0.439, and 0.717.

In (G–J), MARE is 0.139, 0.072, 0.071, and 0.085, and β_{imp} is 0.843, 0.565, 0.694, and

0.675.

Figure 4. Changes in the metrics value and the imputed allelic dosage with

different scalings of the θ value. (A) Rs10410162 is used as an example to visualize

the changes in Rsq, EmpRsq, MARE, and β_{imp} (y-axis) with the 21 scalings of the θ

value (x-axis). To have a better illustration, the x-axis is in the minus log scale. The

scalings of 100, 10, 2, 1, 0.5, 0.1, and 0.01-fold are circled, marked with vertical dashed

lines, and labeled. The other scaling folds are shown as the scatters, with the color key

denoting the fold in a log scale. (B–C) Rs10410162 is used as an example to visualize

the deviation between Rsq, EmpRsq, and MARE (B) or β_{imp} (C). The red line shows that

Rsq equals dosage r^2, and the color keys show the theoretical MARE or β_{imp} values.

The dashed lines represent the contours of MARE or β_{imp}, and adjacent contours have a

value difference of 0.05. The 21 scatters are the same as (A). (D) The plot shows the

imputed allelic dosage of a randomly selected target haplotype (chr19) with the 7

scalings of the θ value. The scatters show variants. The boxes show the median, upper

(75%), and lower (25%) quartiles. The whiskers show the 1.5-fold of the interquartile

range (IQR) extended from the upper or lower quartile if a value exceeds them;

otherwise, they show the maximum or minimum value. Only variants with imputed allelic

dosage < 0.5 and 0.3 < EmpRsq < 0.8 when using the 1-fold θ value are shown.

Figure 5. Rsq and EmpRsq using the 1KGP panel with different scalings of the θ

value. The plot shows the mean Rsq and EmpRsq of each imputation result, stratified

by the MAF of the array. The scalings of 100, 10, 2, 1, 0.5, 0.1, and 0.01-fold are

marked with vertical dashed lines and labeled.

Figure 6. Total θ value of the simulated reference panels. The Supplementary

Figure hows the total θ value of simulated reference panels, estimated by Minimac3.

(A) JPT-only panels with different sample sizes. (B) Fixed-size panels (504) with 1–5

levels of ancestral diversity, as shown in the legend. (C) Adding large-size (504–3,760)

EAS samples to a small-size (403) EUR panel. (D) Adding 1KGP samples to a large-

size (3,256) JPT panel. (E) Simultaneously increasing the size and ancestral diversity

using 1–5 ancestries in the 1KGP. In each plot, the scatters show the values of 10 runs.

The boxes show the median, upper (75%), and lower (25%) quartiles. The whiskers

show the 1.5-fold interquartile range (IQR) extended from the upper or lower quartile if a

value exceeds them; otherwise, they show the maximum or minimum value. In (C),

751 EURn403 represents the 403 EUR. In (C) and (D), 500–3256JPT represents the

752 number of JPT samples in the panel.

754 **Tables**

756 **Table 1. Characteristics of the reference panels and target samples used in this**

757 **study, and the number of variants imputed.**

Reference panel	Reference panel size	Size of East Asian subjects	Percentage of EAS subjects	Sequencing depth	Target sample size	Target sample selection criteria	Number of imputed variants on autosomes
1KGP	2,504	504	20.13%	4–8 ×	180,882	All EAS in BBJ -180k passed QC	47,109,465
BBJ1k	3,541	1,541	43.52%	1KGP + 30 ×	179,947	Same as above. Exclude samples in the reference panel	59,387,070
JEWEL3k	5,760	3,760	65.28%	1KGP + 15–30 ×	177,930	Same as above. Exclude samples in the reference panel	69,076,783
TOPMed	97,256	1,184	1.22%	38 ×	180,882	All EAS in BBJ-180k passed QC	291,214,407

Table 2. Reference panel description, the total θ value estimated by Minimac3, the mean EmpRsq, and the mean Rsq using the simulated EUR-EAS reference panels with different sizes and EUR proportions, and 100 EUR as the target sample.

Reference panel	Reference panel size	EUR (%)	Total θ value	MAF ≥ 5%		5% ≥ MAF > 1%	
				EmpRsq	Rsq	EmpRsq	Rsq
1KGP-EURn403	403	100	740.0	0.849	0.797	0.75	0.679
1KGP-EURn403+1KGP-EAS	907	44.43	722.7	0.841	0.795	0.743	0.7
1KGP-EURn403+1KGP-EAS+500JPT	1,407	28.64	595.9	0.839	0.814	0.744	0.736
1KGP-EURn403+1KGP-EAS+1000JPT	1,907	21.13	534.9	0.836	0.822	0.74	0.752
1KGP-EURn403+1KGP-EAS+1500JPT	2,407	16.74	466.0	0.834	0.831	0.739	0.767
1KGP-EURn403+1KGP-EAS+2000JPT	2,907	13.86	437.5	0.833	0.835	0.737	0.774
1KGP-EURn403+1KGP-EAS+2500JPT	3,407	11.83	408.4	0.831	0.839	0.734	0.78
1KGP-EURn403+1KGP-EAS+3256JPT	4,163	9.68	364.1	0.829	0.844	0.732	0.788

EURn403 represents the 403 EUR; 1KGP-EAS represents the 504 EAS; 500–3256JPT represents the number of JPT samples in the reference panel. Minor allele frequency (MAF) is determined by the genotyping array.

765 **Table 3. Reference panel description, the total θ value estimated by Minimac3, the**

766 **mean EmpRsq, and the mean Rsq using the simulated JPT-1KGP reference**

767 **panels with different sizes and ancestral diversity, and WGS$_{993}$ as the target**

768 **sample.**

Reference panel	Reference panel size	EAS (%)	Total θ value	MAF ≥ 5%		5% > MAF ≥ 1%		1% > MAF ≥ 0.5%	
				EmpRsq	Rsq	EmpRsq	Rsq	EmpRsq	Rsq
3256JPT	3,256	100	326.9	0.959	0.949	0.921	0.91	0.861	0.841
3256JPT+1KGP-JPT	3,360	100	336.2	0.959	0.949	0.921	0.909	0.861	0.839
3256JPT+1KGP-EAS	3,760	100	334.2	0.959	0.948	0.923	0.908	0.859	0.834
3256JPT+1KGP-EAS+EUR	4,263	88.2	377.4	0.959	0.945	0.921	0.899	0.855	0.816
3256JPT+1KGP-EAS+EUR+AFR	4,924	76.36	395.7	0.958	0.944	0.92	0.892	0.857	0.8
3256JPT+1KGP-EAS+EUR+AFR+AMR	5,217	72.07	426.4	0.958	0.942	0.919	0.887	0.857	0.793
3256JPT+1KGP-All	5,760	65.28	422.4	0.958	0.942	0.919	0.886	0.854	0.793

769 The 3,256 WGS samples of BBJ are included in all panels. 1KGP- followed by the

770 ancestry represents the 1KGP subset. Minor allele frequency (MAF) is determined

771 by the genotyping array.