Turing pattern formation in reaction-cross-diffusion systems with a bilayer geometry

Antoine DIEZ[1], Andrew L. KRAUSE[2], Philip K. MAINI[3], Eamonn A. GAFFNEY[3,*], AND Sungrim SEIRIN-LEE[1,4,*]

[1]Institute for the Advanced Study of Human Biology (ASHBi)
Kyoto University Institute for Advanced Study,
Kyoto University,
Yoshida-Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan.

[2]Mathematical Sciences Department,
Durham University,
Upper Mountjoy Campus, Stockton Rd,
Durham DH1 3LE, United Kingdom

[3]Wolfson Centre For Mathematical Biology,
Mathematical Institute, Andrew Wiles Building,
University of Oxford,
Radcliffe Observatory Quarter, Woodstock Road,
Oxford, OX2 6GG, United Kingdom

[4]Department of Mathematical Medicine,
Graduate School of Medicine,
Kyoto University,
Yoshida-Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan

* Corresponding authors
gaffney@maths.ox.ac.uk
lee.seirin.2c@kyoto-u.ac.jp

Abstract

Conditions for self-organisation via Turing's mechanism in biological systems represented by reaction-diffusion or reaction-cross-diffusion models have been extensively studied. Nonetheless, the impact of tissue stratification in such systems is under-explored, despite its ubiquity in the context of a thin epithelium overlying connective tissue, for instance the epidermis and underlying dermal mesenchyme of embryonic skin. In particular, each layer can be subject to extensively different biochemical reactions and transport processes, with chemotaxis – a special case of cross-diffusion – often present in the mesenchyme, contrasting the solely molecular transport typically

found in the epidermal layer. We study Turing patterning conditions for a class of reaction-cross-diffusion systems in bilayered regions, with a thin upper layer and coupled by a linear transport law. In particular, the role of differential transport through the interface is explored together with the presence of asymmetry between the homogeneous equilibria of the two layers. A linear stability analysis is carried out around a spatially homogeneous equilibrium state in the asymptotic limit of weak and strong coupling strengths, where quantitative approximations of the bifurcation curve can be computed. Our theoretical findings, for an arbitrary number of reacting species, reveal quantitative Turing conditions, highlighting when the coupling mechanism between the layered regions can either trigger patterning or stabilize a homogeneous equilibrium regardless of the independent patterning state of each layer. We support our theoretical results through direct numerical simulations, and provide an open source code to explore such systems further.

Keywords: Turing instabilities, stratified systems, skin patterns, interface, chemotaxis

1 Introduction

1.1 Biological motivation and study objectives

Among many other pioneering works, Turing [60] introduced a new, mathematical, way to understand symmetry-breaking phenomena in biology. Over the intervening decades, his concept of a diffusion driven instability in reaction-diffusion systems has been extensively studied in the mathematical literature and confronted with biological observations and experiments [31, 4, 44, 45]. Despite extensive evidence that Turing patterning mechanisms can explain numerous complex phenomena observed in nature, a core challenge is modernizing Turing's ideas to accommodate advances in our understanding of the living world that have emerged since his study. In its most classical version, Turing's theory predicts that a homogeneous equilibrium state between two species can be broken by the sole effect of the increased diffusion of one species [45, Chapter 2], as in the renowned local activation/long-range inhibition paradigm advocated in [23, 40]. Many recent works have refined this theory and proposed more realistic biological scenarios, in particular regarding the geometry of the domain [32], the complexity of the signalling network [34] or the inclusion of other influences, such as cross-diffusion [18, 37], mechanical forces or active cell transport phenomena [46, 38, 61]. Contemporary perspectives on Turing systems are summarized in [32].

One refinement of Turing's ideas that has been under-explored is the bilayer structure of many biological pattern forming systems. For instance, layered development has been indicated as relevant for the morphogenesis of the plant shoot apical meristem [20], cell-membrane Turing self-organisation modulated by signalling molecules in the cytosol [35], cell-cell membrane communication processes [59], and propagation problems in population ecology [21, 56, 7]. Outside biology, bilayer systems have been studied in chemistry in the context of the so-called CIMA and CDIMA experiments [5] which have motivated many analytical and experimental works on the patterning mechanisms of bilayer systems [62, 63, 64, 6, 9].

In particular, an important biological system that especially motivates this work is the embryonic skin, which consists of a thin epidermal layer superimposed on the extracellular matrix of the dermal mesenchyme. While the former is a thin layer mostly composed of tightly packed cells with limited movement, the latter can be much deeper and is composed of a network of collagen fibers supporting motile cells. During embryonic development, the interplay between these two layers gives rise to various repeating anatomical patterns such as hair follicles [25] (Fig. 1a), feather placodes [28] or finger prints [24]. These studies have identified complex signalling networks of diffusive molecules produced in each layer and which interact with the motile mesenchymal cells in order to initiate local cell clustering. Although mathematical reaction-diffusion-chemotaxis models [3, 49, 19] have been able to accurately reproduce the biological observations, they have focused on a simplified mono-layer geometry, i.e. a single domain in which all the species interact. However, biological experiments in [28] have stressed the importance of the bilayer structure on pattern formation by considering chimera skins composed of an epidermis from one species and a dermis from another. Depending on the species considered, the patterning ability may be conserved, or not, as well as the periodic structure of the patterning. This demonstrates that the coupling between the two layers is in itself a crucial component for symmetry-breaking.

Hence, on noting that cross-diffusion encompasses chemotaxis as a special case, the focus of this study will be the mathematical derivation of patterning conditions for bilayered reaction-cross diffusion systems, coupled via linear transport between the layers. In particular, our objective will be to determine when the bilayer structure, with a thin upper layer represented via a one dimensional domain, is predicted to enhance self-organisation or stabilise the homogeneous equilibrium of the system.

1.2 Related works and previous theoretical results

Bilayer systems similar to, but simpler than, the ones considered in the present article have been analytically and computationally studied, in particular in [33] and in [9]. The former article [33] is motivated by patterns formed by bacteria growing on an agar substrate and thus considers a pure reaction-diffusion system with a passive bulk (i.e. with only diffusion). Depending on the thickness of the layers, several asymptotic instability conditions and reduced dispersion relations are derived and studied numerically. The latter article [9] provides a detailed analysis of a coupled reaction-diffusion system of two exactly identical two-component layers. In this setting, the linear stability analysis of the coupled system is amenable to block-matrix computations which reduce the problem to a classical 2D eigenvalue problem and thus allows for a detailed bifurcation analysis.

Much earlier works were also motivated by the role of reaction-diffusion mechanisms in skin patterning problems. In particular, in [48], a very similar example based on a two-component reaction-diffusion system is studied numerically with the aim of distinguishing the mechanisms leading to spot and stripe patterns. The authors assume equal reaction terms in both layers but possibly different diffusion coefficients and a nonlinear transport law between the two layers. Patterning is assumed to

be driven by the epidermis where the homogeneous state is unstable. Related works [58, 11, 46] have also considered mechano-chemical models of pattern formation in the skin, with an overview presented in [45, Chapter 6].

Other closely related theoretical works [54, 53, 35, 36, 26] in the literature are motivated by the cell membrane-cytosol system and thus focus on a spherical geometry and most often a passive bulk. Although in this context the curvature of the domain itself may have an important effect on pattern formation, we restrict ourselves to a planar geometry which is more relevant for the example of skin patterning. We also generalize the system to consider chemotaxis and other nonlinear reaction-diffusion effects present in both layers. Another class of related models recently studied in the literature are the so-called compartmental models [51, 50] where two or more reaction-diffusion compartments are spatially coupled through a passive diffusive bulk.

Although all these works study patterning conditions for various types of coupling between reaction-diffusion systems, their modelling frameworks are quite different from the bilayer structure that we consider here. In this article, we will consider a more general cross-diffusion framework (which includes chemotaxis models) with two non-necessarily identical active layers and an arbitrary number of interacting species. We give a set of quantitative conditions for (non)-patterning in the asymptotic limit of small and large couplings and provide several examples of patterning scenarios, theoretically, and with numerical evidence for reaction-diffusion and chemotaxis systems. In particular we consider the general scenario of pattern formation driven by an individual layer or their mutual coupling. Our analysis is based on classical linear stability analysis in the context of multi-component reaction-cross-diffusion systems. Since explicit analytical results cannot typically be obtained in this situation, we derive patterning conditions via quantitative approximations of the bifurcation curve depending on the coupling strength. In the weak coupling case, related techniques have been used in a different context for the study of weakly coupled oscillators and reaction-diffusion networks [15, 16].

The present article is structured as follows. The modelling framework is described in Section 2 and the main contributions are summarized in Section 2.4. As a starting point of the analysis, the dispersion relations and bifurcation conditions are written in full generality in Section 3. We then split the analysis into two asymptotic cases, first the weak coupling case in Section 4 and secondly the large coupling case in Section 5. Phenomena in the intermediate coupling case are briefly described in Section 6 before the conclusions and discussion in Section 7. The supplementary material contains a description of the numerical methods (Appendix A) and the list and description of the supplementary videos (Appendix B).

2   Models and basic properties

2.1   1D surface - 2D bulk space model

We first consider a suitably non-dimensionalised bilayer system $\Omega_S \cup \Omega_B$ where the thin upper layer of length $L > 0$, $\Omega_S = [0, L]$ – referred to as the surface – is taken to be infinitesimally thin compared

to the lower layer $\Omega_B = [0, L] \times [0, H]$ – referred to as the bulk – which has a depth $H > 0$ (Fig. 1b). We consider n species in the surface and m species in the bulk. In both layers we assume interactions of these species according to a system of reaction-cross-diffusion equations. Their concentrations are denoted by[1] $\boldsymbol{u_S}(x) \in \mathbb{R}^n$ in the surface at the location $x \in \Omega_S$ and $\boldsymbol{u_B}(x, y) \in \mathbb{R}^m$ in the bulk at the location $(x, y) \in \Omega_B$. We will always consider $m \geq n$ and we assume that the n species in the surface can diffuse to the bulk and vice-versa for the first n components of the bulk species $\boldsymbol{u_B}$. There are $m - n$ species which do not diffuse to the surface and remain in the bulk. The mathematical analysis would hold similarly in the reverse case but we choose $m \geq n$ for both parsimony and also noting the biological case of skin patterning where an extra chemotaxis component would be included in the bulk domain. Then, the general form of the model is given by

$$\partial_t \boldsymbol{u_S}(x) = \boldsymbol{f_S}(\boldsymbol{u_S}) + \eta \boldsymbol{A}\big(\tilde{\boldsymbol{u}}_{\boldsymbol{B}}(x, H) - \boldsymbol{u_S}(x)\big) + \nabla \cdot \big(\boldsymbol{D_S}(\boldsymbol{u_S})\nabla \boldsymbol{u_S}\big), \tag{1}$$

$$\partial_t \boldsymbol{u_B}(x, y) = \boldsymbol{f_B}(\boldsymbol{u_B}) + \nabla \cdot \big(\boldsymbol{D_B}(\boldsymbol{u_B})\nabla \boldsymbol{u_B}\big), \tag{2}$$

where the matrix $\boldsymbol{A} \in \mathbb{R}^{n \times n}$ specifies the exchange rates of the different species between the two layers and $\tilde{\boldsymbol{u}}_{\boldsymbol{B}} \in \mathbb{R}^n$ denotes the vector constructed by taking the first n components of $\boldsymbol{u_B} \in \mathbb{R}^m$. The parameter $\eta \geq 0$ represents the strength of the coupling between the two layers. The reaction functions $\boldsymbol{f_S}$ and $\boldsymbol{f_B}$ are arbitrary and the cross-diffusion matrices $\boldsymbol{D_S}$ and $\boldsymbol{D_B}$ are positive definite.

On the lateral sides and the bottom side of the bulk, we assume zero-flux boundary conditions. At $y = H$, we consider the following linear transport law for the bulk species:

$$\boldsymbol{D_B}\frac{\partial \boldsymbol{u_B}}{\partial y}(x, H) = \eta \tilde{\boldsymbol{B}}\big(\tilde{\boldsymbol{u}}_{\boldsymbol{S}}(x) - \boldsymbol{u_B}(x, H)\big), \tag{3}$$

with $\tilde{\boldsymbol{B}} \in \mathbb{R}^{m \times m}$ denoting the matrix constructed on padding $\boldsymbol{B} \in \mathbb{R}^{n \times n}$ by zeros, where $\boldsymbol{B}$ specifies linear transport from the bulk to the surface. Similarly, $\tilde{\boldsymbol{u}}_{\boldsymbol{S}} \in \mathbb{R}^m$ denotes the vector constructed from $\boldsymbol{u_S} \in \mathbb{R}^n$ by adding $m - n$ rows with zero components. Furthermore, note that $\boldsymbol{B} - \boldsymbol{A}$ corresponds to interfacial sources/sinks of species, which are allowed for generality, but these will often be zero in applications.

Note that as in earlier works [48], one could also consider nonlinear transport laws but for simplicity and because we will only carry out a linear stability analysis that would also linearize this part of the equation, we will only work with linear transport between the layers.

As motivated in the introduction, we focus on a model representing the bilayer structure of embryonic skin [25, 28, 24], with bulk chemotaxis where cells restricted to the bulk are chemo-attracted by n species, which diffuse between the two layers (Fig. 1). In this case $m = n + 1$ and $\boldsymbol{u_B} = (\tilde{\boldsymbol{u}}_{\boldsymbol{B}}, c)$ where $c(t, x, y)$ denotes the cell concentration. Assuming a classical reaction-diffusion interaction for

---

[1] We suppress the dependence on time, t, for notational convenience.

the chemical species, the equation in the bulk reduces to

$$\partial_t \tilde{\boldsymbol{u}}_{\boldsymbol{B}} = \tilde{\boldsymbol{f}}_{\boldsymbol{B}}(\tilde{\boldsymbol{u}}_{\boldsymbol{B}}, c) + \tilde{\boldsymbol{D}}_{\boldsymbol{B}} \nabla^2 \boldsymbol{u}_{\boldsymbol{B}} \tag{4}$$

$$\partial_t c = -\nabla \cdot \left(h(c) \nabla k(\tilde{\boldsymbol{u}}_{\boldsymbol{B}}) \right) + r(c) + d_c \nabla^2 c, \tag{5}$$

where $\tilde{\boldsymbol{D}}_{\boldsymbol{B}} \in \mathbb{R}^{n \times n}$ is a positive definite diagonal matrix and $d_c > 0$. The representative examples for $h(c)$, $r(c)$ and $k(\tilde{\boldsymbol{u}})$ are given by

$$h(c) = c, \quad r(c) = r_0 c(c^* - c),$$

with r_0, c^* positive constants, and $k(\tilde{\boldsymbol{u}}) = u_1$, where u_1 is the first component of $\tilde{\boldsymbol{u}} \in \mathbb{R}^n$. The boundary condition at the interface can be rewritten

$$\tilde{\boldsymbol{D}}_{\boldsymbol{B}} \frac{\partial \tilde{\boldsymbol{u}}_{\boldsymbol{B}}}{\partial y} = \eta \boldsymbol{B}(\boldsymbol{u}_{\boldsymbol{S}} - \tilde{\boldsymbol{u}}_{\boldsymbol{B}}), \quad \frac{\partial c}{\partial y} = 0.$$

This corresponds to Eqs. (2), (3) with

$$\boldsymbol{D}_{\boldsymbol{B}}(\tilde{\boldsymbol{u}}, c) = \begin{pmatrix} \tilde{\boldsymbol{D}}_{\boldsymbol{B}} & 0 \\ -h(c)\nabla k(\tilde{\boldsymbol{u}})^{\mathrm{T}} & d_c \end{pmatrix}, \quad \boldsymbol{f}_{\boldsymbol{B}}(\tilde{\boldsymbol{u}}, c) = \left(\tilde{\boldsymbol{f}}_{\boldsymbol{B}}(\tilde{\boldsymbol{u}}, c)^{\mathrm{T}}, r(c) \right)^{\mathrm{T}},$$

where the superscript T denotes the transpose.

2.2 1D surface - 1D thin bulk space model

In the embryonic skin example described in the introduction, local cell clustering is typically observed only at the interface between the epidermal and dermal layers (Fig. 1a) and the whole process takes place on a spatial range which does not exceed the diameter of a few cells [25, 28]. Thus, we will also consider the case of an infinitesimally thin bulk (Fig. 1c).

Under an appropriate rescaling of the coupling intensity η, the 1D-2D model (1)-(3) can be reduced to two coupled one-dimensional equations. In particular, let us consider a bulk concentration $\boldsymbol{u}_{\boldsymbol{B}}$ and a bulk depth of $\varepsilon > 0$, with $\varepsilon \ll 1$ corresponding to the additional assumption that the bulk depth is much smaller than any other lengthscale in the 1D-2D model of the previous subsection. We then introduce the rescaled concentration $\boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon}(x, Y) = \boldsymbol{u}_{\boldsymbol{B}}(x, y)$ where $Y = \varepsilon^{-1} y \in [0, 1]$. With this change of variable, Eq. (2) becomes

$$\partial_t \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon}(x) = \boldsymbol{f}_{\boldsymbol{B}}(\boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon}) + \partial_x \left(\boldsymbol{D}_{\boldsymbol{B}} \, \partial_x \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon} \right) + \frac{1}{\varepsilon^2} \partial_Y \left(\boldsymbol{D}_{\boldsymbol{B}} \, \partial_Y \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon} \right).$$

We also rewrite $\boldsymbol{B} = \varepsilon \hat{\boldsymbol{B}}$, with the components of $\hat{\boldsymbol{B}}$ scaling as $O(1)$, to ensure there is a balance between the inter-layer flux and reaction terms of the rescaled bulk equations, noting this scaling will also be inherited by $\tilde{\boldsymbol{B}} = \varepsilon \hat{\tilde{\boldsymbol{B}}}$. In particular, for other scalings, after the limit of infinitesimal bulk thickness has been taken, only one of the reaction term or the inter-layer flux will appear in the

Figure 1: Embryonic skin patterning in bilayer tissue and domain geometry. (a) An example of skin patterning: the emergence of primordia of hair follicles at the interface between the epidermis and the dermis of a mouse embryo at 15.5 days post coitum. The formation of these condensates of mesenchymal cells can be explained by a complex network of reaction-diffusion and chemotaxis interactions between the two layers (see [25, 28]). Original picture from the e-Mouse Atlas Project under a CC BY 3.0 licence [2]. (b) Mathematical model in a 1D-2D geometry. The epidermis (later referred to as the surface) is modelled by a one-dimensional layer while the dermis (later referred to as the bulk) is two-dimensional to take into account its depth. The arrow symbolizes the boundary condition along the outward normal of the domain. (c) Mathematical model in the 1D-1D geometry where the two layers are taken to be one-dimensional.

leading order dominant balance. If the reaction terms dominate, then the layers are uncoupled and the patterning is based on the individual layer dynamics at leading order, while if the inter-layer transport dominates then there is no chemotaxis or interaction between the signalling molecules, which again is not the system that is to be considered, as biological mechanisms of self-organisation are lost. Thus, with the scaling $\boldsymbol{B} = \varepsilon \hat{\boldsymbol{B}}$, we expand $\boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon}$ in powers of ε,

$$\boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon} = \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,0} + \varepsilon \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,1} + \varepsilon^2 \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,2} + \dots,$$

to obtain, at order ε^{-2} and ε^{-1},

$$\partial_Y \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,0} = \partial_Y \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,1} = \boldsymbol{0}.$$

The boundary condition Eq. (3) becomes

$$\boldsymbol{D}_{\boldsymbol{B}} \partial_Y \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,2} = \eta \widehat{\tilde{\boldsymbol{B}}} \left(\tilde{\boldsymbol{u}}_{\boldsymbol{S}} - \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,0} \right).$$

Moreover, it follows that at leading order,

$$\partial_t \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,0}(x) = \boldsymbol{f}_{\boldsymbol{B}}(\boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,0}) + \partial_x \left(\boldsymbol{D}_{\boldsymbol{B}} \, \partial_x \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,0} \right) + \partial_Y \left(\boldsymbol{D}_{\boldsymbol{B}} \, \partial_Y \boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,2} \right).$$

Noting $\boldsymbol{u}_{\boldsymbol{B}}^{\varepsilon,0}, \tilde{\boldsymbol{u}}_{\boldsymbol{B}}^{\varepsilon,0}$ are independent of Y, integrating the equation in Y between 0 and 1 and, for notational simplicity, dropping the hat of $\widehat{\tilde{\boldsymbol{B}}}$ and the superscript label "$\varepsilon,0$", reveals

$$\partial_t \boldsymbol{u}_{\boldsymbol{S}} = \boldsymbol{f}_{\boldsymbol{S}}(\boldsymbol{u}_{\boldsymbol{S}}) + \eta \boldsymbol{A}(\tilde{\boldsymbol{u}}_{\boldsymbol{B}} - \boldsymbol{u}_{\boldsymbol{S}}) + \nabla \cdot \left(\boldsymbol{D}_{\boldsymbol{S}}(\boldsymbol{u}_{\boldsymbol{S}}) \nabla \boldsymbol{u}_{\boldsymbol{S}} \right) \tag{6}$$

$$\partial_t \boldsymbol{u}_{\boldsymbol{B}} = \boldsymbol{f}_{\boldsymbol{B}}(\boldsymbol{u}_{\boldsymbol{B}}) + \eta \tilde{\boldsymbol{B}}(\tilde{\boldsymbol{u}}_{\boldsymbol{S}} - \boldsymbol{u}_{\boldsymbol{B}}) + \nabla \cdot \left(\boldsymbol{D}_{\boldsymbol{B}}(\boldsymbol{u}_{\boldsymbol{B}}) \nabla \boldsymbol{u}_{\boldsymbol{B}} \right). \tag{7}$$

Hence one may also observe that once the infinitesimal bulk thickness limit has been taken, and with the constraint $\boldsymbol{A} = \boldsymbol{B}$, (where we recall that $\boldsymbol{B}$ is the $n \times n$ block of $\tilde{\boldsymbol{B}}$ corresponding to species that can transport between the upper and lower regions), then there are no additional sources or sinks at the layer interface for the leading order equations. So the 1D-1D model is in some sense a special case of the 1D-2D model given in Eqs. (1)-(3), though they may not admit the same spatially-homogeneous equilibria.

2.3 Equilibria structure

Throughout this article, we assume that there exists a constant equilibrium point $\boldsymbol{w}^* = (\boldsymbol{u}_{\boldsymbol{S}}^*, \boldsymbol{u}_{\boldsymbol{B}}^*)$ such that in the 1D-1D case

$$\boldsymbol{f}_{\boldsymbol{S}}(\boldsymbol{u}_{\boldsymbol{S}}^*) + \eta \boldsymbol{A}(\tilde{\boldsymbol{u}}_{\boldsymbol{B}}^* - \boldsymbol{u}_{\boldsymbol{S}}^*) = 0, \tag{8}$$

$$\boldsymbol{f}_{\boldsymbol{B}}(\boldsymbol{u}_{\boldsymbol{B}}^*) + \eta \tilde{\boldsymbol{B}}(\tilde{\boldsymbol{u}}_{\boldsymbol{S}}^* - \boldsymbol{u}_{\boldsymbol{B}}^*) = 0. \tag{9}$$

Note that this equilibrium point $\boldsymbol{w}^* \equiv \boldsymbol{w}^*(\eta)$ may depend on η whenever $\boldsymbol{u}_S^* \neq \tilde{\boldsymbol{u}}_B^*$. In the 1D-2D case, the boundary condition at the interface imposes $\boldsymbol{u}_S^* = \tilde{\boldsymbol{u}}_B^*$ and $\boldsymbol{w}^*$ is thus independent of the coupling strength η and only depends on the reaction terms, which must satisfy

$$\boldsymbol{f}_S(\boldsymbol{u}_S^*) = 0, \quad \boldsymbol{f}_B(\boldsymbol{u}_B^*) = 0.$$

Note that when $m > n$, the $m - n$ pure bulk species concentrations may act as free parameters in the second relation. For instance in the chemotaxis case, the equilibrium cell concentration c^* can act as a parameter of the system, specified by a growth term of the form $r(c) = c(c^* - c)$. As we are going to show in the following, the ability of the 1D-1D system to have different equilibrium concentrations in the surface and the bulk can have a strong influence on pattern formation.

An important point to note is that the existence or uniqueness of such an equilibrium point in the 1D-1D case cannot be guaranteed by the sole intrinsic properties of the reaction terms, due to the dependence on η. For a sufficiently small coupling strength η, we will thus adopt a perturbative approach. We always assume that at $\eta = 0$, the uncoupled systems have a homogeneous equilibrium point. Then, by the implicit function theorem, for η sufficiently small, it is possible to find a curve $\eta \mapsto \boldsymbol{w}^*(\eta)$ which satisfies the relations Eqs. (8)-(9), provided that the Jacobian matrices of $\boldsymbol{f}_S$ and $\boldsymbol{f}_B$ evaluated, respectively, at $\boldsymbol{u}_S^*(0)$ and $\boldsymbol{u}_B^*(0)$ are invertible (see Section 4.3 for more details).

2.4 Objectives and contributions

The goal of this study is to understand how Turing self-organisation is affected by the coupling in a bilayer geometry and to derive mathematical conditions for the linear instability of a homogeneous equilibrium in order to generate patterns. Thus we consider the fundamental mechanism of Turing pattern formation theory, namely diffusion-driven instability. We will focus our analysis on the two asymptotic regimes $\eta \ll 1$ and $\eta \gg 1$ and we will obtain explicit quantitative patterning conditions that can be numerically computed. We will always assume that the two layers independently can generate Turing patterns in general, though not necessarily with the parameters investigated, so that our main objective will be to determine analytically the cases where the coupling enhances or diminishes this patterning ability. Before proceeding to the mathematical analysis, we briefly summarize our main contributions below:

- **Weak coupling case ($\eta \ll 1$) for the 1D-1D system:** Turing patterns can be formed by coupling two independently non-patterning layers provided that the exchange rate of one species is large enough (Section 4.2, Fig. 2b, Video 1) or that the equilibria of the two uncoupled layers are different (Section 4.3, Fig. 3).

- **Weak coupling case ($\eta \ll 1$) for the 1D-2D system:** we extend the 1D-1D results by studying (asymptotically) the influence of the bulk depth (Section 4.4.1). We also show different patterning scenarios in Section 4.4.3 (Videos 2-3-4).

- **Strong coupling case ($\eta \gg 1$) for the 1D-1D system:** we show that the coupled system reduces to one single cross-diffusion equation (Section 5.1) and find a simple criterion to determine whether the coupling stabilizes the homogeneous state (Section 5.2.1, Video 5) or enhances patterning (Section 5.2.2, Video 6).

- **Intermediate coupling case:** In Section 6, we briefly comment on the wide variety of patterning scenarios, and the difficulty in studying them in any generality, beyond the asymptotic regimes above.

3 Dispersion relations and bifurcation points

As a starting point, we derive the dispersion and bifurcation relations associated with a weak linear perturbation of the homogeneous state. In addition, in the 1D-2D case, the change of variable $y' = H - y$ is convenient, so as to locate the interface at $y' = 0$, though the prime is dropped below.

3.1 1D-1D

For the 1D-1D system (6)-(7), taking into account the Neumann boundary conditions on the lateral sides, we consider a linear perturbation of the equilibrium point of the form

$$
\begin{aligned}
\boldsymbol{u_S} &= \boldsymbol{u_S^*} + \varepsilon \mathrm{e}^{\lambda t} \boldsymbol{u_{S,q}} \cos(k_q x), \\
\boldsymbol{u_B} &= \boldsymbol{u_B^*} + \varepsilon \mathrm{e}^{\lambda t} \boldsymbol{u_{B,q}} \cos(k_q x),
\end{aligned}
\tag{10}
$$

where q is an integer and $k_q = q\pi/L$. Linearizing Eqs. (6)-(7) around this equilibrium point shows that at order ε, the perturbation vector $\boldsymbol{w}_q = (\boldsymbol{u_{S,q}}, \boldsymbol{u_{B,q}})^{\mathrm{T}}$ belongs to the kernel of the following matrix

$$
\boldsymbol{M} = \begin{pmatrix} \lambda \boldsymbol{I} + \boldsymbol{P_0} & \boldsymbol{0} \\ \boldsymbol{0} & \lambda \boldsymbol{I} + \boldsymbol{Q_0} \end{pmatrix} + \eta \boldsymbol{E}.
\tag{11}
$$

Here $\boldsymbol{I}$ is the identity matrix, $\xi = k_q^2$ is treated as a continuous variable, $\boldsymbol{P_0} = \xi \boldsymbol{D_S} - \boldsymbol{J_S}$, $\boldsymbol{Q_0} = \xi \boldsymbol{D_B} - \boldsymbol{J_B}$, where $\boldsymbol{J_B}$ and $\boldsymbol{J_S}$ are the Jacobian matrices of $\boldsymbol{f_S}$ and $\boldsymbol{f_B}$ with respect to $\boldsymbol{u_S}$ and $\boldsymbol{u_B}$, respectively and,

$$
\boldsymbol{E} = \begin{pmatrix} \boldsymbol{A} & -\boldsymbol{A} & \boldsymbol{0} \\ -\boldsymbol{B} & \boldsymbol{B} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{0} & \boldsymbol{0} \end{pmatrix}.
$$

Unless specified otherwise, all expressions are evaluated at the equilibrium point $\boldsymbol{w}^*$. Note again that $\boldsymbol{w}^*$ may depend on η if the surface and bulk equilibrium concentrations are not equal.

The dispersion relation is given by the determinant of $\boldsymbol{M}$ being zero, i.e.

$$
|\boldsymbol{M}| = 0,
$$

which is a polynomial relation in λ and ξ but possibly non polynomial in η due to the unknown dependence of $\boldsymbol{w^*}$ on η. For a given set of parameters and a given η, it is nevertheless possible to determine the stability of each mode ξ by applying the Routh-Hurwitz criterion to the polynomial in λ of degree $(m+n)$.

3.2   1D-2D

In the 1D-2D case, the amplitude of the perturbation in the bulk depends on the y-variable so that the linearized equations become

$$\lambda \boldsymbol{u_{S,q}} = (\boldsymbol{J_S} - k_q^2 \boldsymbol{D_S})\boldsymbol{u_{S,q}} + \eta \boldsymbol{A}(\tilde{\boldsymbol{u}}_{\boldsymbol{B},q}(y=0) - \boldsymbol{u_{S,q}})$$
$$\lambda \boldsymbol{u_{B,q}}, = \boldsymbol{J_B}\boldsymbol{u_{B,q}} + \boldsymbol{D_B}(-k_q^2 + \partial_{yy}^2)\boldsymbol{u_{B,q}}.$$

The second equation can be rewritten

$$\partial_{yy}^2 \boldsymbol{u_{B,q}} = \boldsymbol{M_B^2}\boldsymbol{u_{B,q}}, \tag{12}$$

where the matrix $\boldsymbol{M_B}$ is a square root of

$$\boldsymbol{M_B^2} = \boldsymbol{D_B^{-1}}(\lambda \boldsymbol{I} + \boldsymbol{Q_0}).$$

Note that in principle the square root of a matrix is not unique and may not even exist. The square root of a diagonal matrix can be constructed by taking the principal square root of the diagonal elements. This construction readily extends to diagonalizable matrices which form a dense open set. However, as we shall see, this particular convention does not play any role in the following. Solving Eq. (12) in y, with $y = 0$ corresponding to the interface, and no-flux conditions at $y = H$, gives

$$\boldsymbol{u_{B,q}} = \cosh((H-y)\boldsymbol{M_B})\boldsymbol{v_{B,q}},$$

for some vector $\boldsymbol{v_{B,q}}$, with the interfacial conditions still to be imposed. These, in turn, give

$$\boldsymbol{D_B}\boldsymbol{M_B}\sinh(H\boldsymbol{M_B})\boldsymbol{v_{B,q}} = \eta \tilde{\boldsymbol{B}}\big(\tilde{\boldsymbol{u}}_{\boldsymbol{S}} - \cosh(H\boldsymbol{M_B})\boldsymbol{v_{B,q}}\big).$$

In the previous expressions, the hyperbolic sine and cosine of a matrix are defined by their power series expansion, namely for a matrix $\boldsymbol{M}$,

$$\cosh(H\boldsymbol{M}) := \sum_{k=0}^{+\infty} \frac{H^{2k}}{(2k)!}\boldsymbol{M}^{2k},$$
$$\boldsymbol{M}\sinh(H\boldsymbol{M}) := \sum_{k=1}^{+\infty} \frac{H^{2k-1}}{(2k-1)!}\boldsymbol{M}^{2k}.$$

₂₄₀ Note in particular that these are functions of $\boldsymbol{M}^2$, so that the final results do not depend on taking a
₂₄₁ matrix square root, nor the choice of the square root.

₂₄₂   Finally, we conclude that $\boldsymbol{w}_q(0) := (\boldsymbol{u}_{S,q}, \boldsymbol{v}_{B,q})$ belongs to the kernel of the following matrix

$$\boldsymbol{M} = \begin{pmatrix} \lambda\boldsymbol{I} + \boldsymbol{P_0} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{D_B}\boldsymbol{M_B}\sinh(H\boldsymbol{M_B}) \end{pmatrix} + \eta\boldsymbol{E}, \tag{13}$$

₂₄₃ with

$$\boldsymbol{E} = \begin{pmatrix} \boldsymbol{A} & -\boldsymbol{A}[\cosh(H\boldsymbol{M_B})]_n & \boldsymbol{0} \\ -\boldsymbol{B} & & \tilde{\boldsymbol{B}}\cosh(H\boldsymbol{M_B}) \\ \boldsymbol{0} & & \end{pmatrix},$$

₂₄₄ where $[\cosh(H\boldsymbol{M_B})]_n$ denotes the first n rows of $\cosh(H\boldsymbol{M_B})$. Note that, unlike $\lambda\boldsymbol{I} + \boldsymbol{P_0}$, the matrix
₂₄₅ $\boldsymbol{M_B}$ does not depend linearly on λ or ξ due to the square root and the hyperbolic sine and cosine.
₂₄₆ Consequently, the dispersion relation

$$|\boldsymbol{M}| = 0,$$

₂₄₇ is a transcendental relation in all variables. In particular, there is no standard criterion, like the
₂₄₈ Routh-Hurwitz criterion in the polynomial case, to determine the stability of a given mode. Note also
₂₄₉ that this dispersion relation is a straightforward extension of the model studied in [33] in a 2D-2D
₂₅₀ reaction-diffusion scenario with a passive bulk layer.

₂₅₁   As an example, in the chemotaxis case given by Eqs. (4)-(5), one can check that

$$\boldsymbol{M}_{\boldsymbol{B}}^2 = \begin{pmatrix} \boldsymbol{R} & \boldsymbol{w}_1 \\ \boldsymbol{w}_2^{\mathrm{T}} & \omega \end{pmatrix},$$

with

$$\boldsymbol{R} = \boldsymbol{D_B}^{-1}\left(\lambda\boldsymbol{I} - \boldsymbol{J_B} + k_q^2\boldsymbol{D_B}\right)$$

$$\boldsymbol{w}_1 = -\boldsymbol{D_B}^{-1}\frac{\partial\boldsymbol{f_B}}{\partial c}$$

$$\boldsymbol{w}_2 = d_c^{-1}h(c^*)(\boldsymbol{R}^{\mathrm{T}} - k_q^2\boldsymbol{I})(\nabla_{\boldsymbol{u_B}}k)(\boldsymbol{u}_{\boldsymbol{B}}^*)$$

$$\omega = d_c^{-1}\left(\lambda + d_c k_q^2 + h(c^*)(\nabla_{\boldsymbol{u_B}}k)(\boldsymbol{u}_{\boldsymbol{B}}^*)\cdot\boldsymbol{w}_1 - r_c\right),$$

₂₅₂ where $r_c = \partial r/\partial c$, evaluated at the equilibrium point.

₂₅₃ 3.3 Bifurcation points

₂₅₄ In the following, all the parameters are assumed to be fixed except for one, generically denoted by
₂₅₅ δ which will be taken as a bifurcation parameter. Classically, δ is the diffusion coefficient of one of
₂₅₆ the species, for instance the inhibitor diffusion in a classical two-species reaction-diffusion system [45,
₂₅₇ Chapter 2]. Note however that in the following, the parameter δ could be any parameter of the model,

such as the chemotaxis strength. Highlighting the dependence with respect to the various parameters, the dispersion relation reads

$$|\boldsymbol{M}|(\eta, \boldsymbol{w^*}, \lambda, \xi, \delta) = 0, \tag{14}$$

where $\boldsymbol{M}$ is given by Eq. (11) or Eq. (13).

Following the classical Turing theory for marginal instability, for a given η we are interested in the critical value δ_c of the bifurcation parameter at which $\lambda = 0$ is a solution of the dispersion relation, while all the other solutions have a negative real part (i.e. a Turing bifurcation). In this article, we will only consider the case of a Turing bifurcation and leave the case of Hopf (or Wave) bifurcations (i.e. associated with $\lambda = i\rho$, ρ real) for future work. Thus, for a given η and a given equilibrium $\boldsymbol{w^*}$, the critical values ξ_c and δ_c satisfy the following relation:

$$a_0(\eta, \boldsymbol{w^*}, \xi_c, \delta_c) := |\boldsymbol{M}|(\eta, \boldsymbol{w^*}, 0, \xi_c, \delta_c) = 0, \tag{15}$$

and our goal is to study the dependence of this solution on η and $\boldsymbol{w^*}$. Moreover, since all the other solutions of the dispersion relation are assumed to have a non positive real part, it follows that $a_0 \geq 0$ in a ξ neighbourhood of ξ_c, which imposes the second relation

$$\left.\frac{\partial a_0}{\partial \xi}\right|_{\xi=\xi_c} = 0. \tag{16}$$

In the following, it will be useful to expand a_0 as

$$a_0(\eta, \boldsymbol{w^*}, \xi, \delta) = \sum_{k=0}^{n+m} \eta^k H_k(\boldsymbol{w^*}, \xi, \delta). \tag{17}$$

The first term is computed by setting $\eta = 0$, in which case Eq. (15) reduces to a block-diagonal determinant:

$$H_0(\boldsymbol{w^*}, \xi, \delta) = |\boldsymbol{P_0}||\boldsymbol{W_0}|, \tag{18}$$

where $\boldsymbol{P_0} = \xi \boldsymbol{D_S} - \boldsymbol{J_S}$ and $\boldsymbol{W_0} = \boldsymbol{Q_0} = \xi \boldsymbol{D_B} - \boldsymbol{J_B}$ in the 1D-1D case. For the 1D-2D case, $\boldsymbol{P_0}$ has the same definition but $\boldsymbol{W_0} := \boldsymbol{D_B}\sqrt{\boldsymbol{R_0}}\sinh(H\sqrt{\boldsymbol{R_0}})$ with $\boldsymbol{R_0} = \boldsymbol{D_B^{-1}}(\xi \boldsymbol{D_B} - \boldsymbol{J_B})$ and $\sqrt{\boldsymbol{R_0}}$ denoting any square root of $\boldsymbol{R_0}$. For the second term, we recall Jacobi's formula for the differential of the determinant function,

$$\frac{\mathrm{d}}{\mathrm{d}h}|\boldsymbol{A}(h)| = \mathrm{co}(\boldsymbol{A}(h)) \cdot \boldsymbol{A'}(h),$$

for any matrix-valued differentiable curve $h \in \mathbb{R} \mapsto \boldsymbol{A}(h)$, with derivative at h denoted by $\boldsymbol{A'}(h)$, where $\boldsymbol{A} \cdot \boldsymbol{B} := \mathrm{Tr}(\boldsymbol{A^T B})$ denotes the usual matrix dot product and $\mathrm{co}(\boldsymbol{A})$ is the comatrix of the matrix $\boldsymbol{A}$ (which is the transpose of the adjugate matrix).

Consequently, we obtain that

$$H_1(\boldsymbol{w^*}, \xi, \delta) = \left.\frac{\partial a_0}{\partial \eta}\right|_{\eta=0} = |\boldsymbol{W_0}|\,\mathrm{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A} + |\boldsymbol{P_0}|\,\mathrm{co}(\boldsymbol{W_0}) \cdot \tilde{\boldsymbol{B}}, \tag{19}$$

in the 1D-1D case and

$$H_1(\boldsymbol{w^*}, \xi, \delta) = \left. \frac{\partial a_0}{\partial \eta} \right|_{\eta=0} = |\boldsymbol{W_0}| \operatorname{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A} + |\boldsymbol{P_0}| \operatorname{co}(\boldsymbol{W_0}) \cdot \tilde{\boldsymbol{B}} \cosh(H\sqrt{\boldsymbol{R_0}}), \qquad (20)$$

in the 1D-2D case.

Remark 1. *The expansion* (17) *will be useful to study the small coupling case $\eta \ll 1$ in the next section. Following the same methodology, it would possible to compute the full Taylor expansion of a_0 using the Faa di Bruno formula, although, due the algebraic complexity, it does not seem possible to obtain exploitable analytical results beyond the first order.*

4 Weak coupling case for the 1D-1D and 1D-2D models

When $\eta = 0$, the surface and bulk systems are decoupled and we assume that at least one of them has a bifurcation parameter with the ability to produce Turing patterns. Classically, it is possible to compute, explicitly or numerically, the critical value of this bifurcation parameter, $\delta_c(0)$, as well as the critical wave number $\xi_c(0)$ (see for instance [45, Section 2.3] as well as Section 4.2.2 in the case of a two-species reaction-diffusion system). The goal of this section is to study how this critical parameter changes when the two systems are coupled.

4.1 General formula

We first give a simple criterion for the existence of a critical bifurcation parameter function $\eta \mapsto \delta_c(\eta)$ and $\eta \mapsto \xi_c(\eta)$ in a neighbourhood of $\eta = 0$. Since these critical values are defined by Eqs. (15)-(16), by the implicit function theorem and the expansion given in Eq. (17), this reduces to proving that

$$\det \begin{pmatrix} \partial_\xi H_0 & \partial_\delta H_0 \\ \partial^2_{\xi\xi} H_0 & \partial^2_{\xi\delta} H_0 \end{pmatrix} \neq 0,$$

where all the partial derivatives are evaluated at $(\xi_c(0), \delta_c(0))$. We will call surface-driven (resp. bulk-driven) symmetry-breaking the case where δ is the diffusion coefficient of a surface (resp. bulk) species and thus at $\delta_c(0)$, $|\boldsymbol{P_0}| = \partial_\xi |\boldsymbol{P_0}| = 0$ and $|\boldsymbol{Q_0}| \neq 0$ (resp. with $\boldsymbol{Q_0}$ and $\boldsymbol{P_0}$ switched). Without loss of generality, let us therefore assume a surface-driven symmetry breaking (again, otherwise $\boldsymbol{P_0}$ and $\boldsymbol{Q_0}$ are simply switched). In this case, using Eq. (18), the above condition thus reduces to

$$\partial^2_{\xi\xi} |\boldsymbol{P_0}| \, \partial_\delta |\boldsymbol{P_0}| \neq 0. \qquad (21)$$

This condition is fulfilled, for instance, for a classical two-species activator-inhibitor system away from higher codimension points, as it is just a transversality condition. We will assume Eq. (21) holds in the following.

In order to know if the coupling increases or decreases the ability to form patterns, we want to

compute the derivative of δ_c at $\eta = 0$ and thus obtain a first order approximation of the bifurcation curve $\eta \mapsto \delta_c(\eta)$. To do so, since

$$a_0(\eta, \boldsymbol{w}^*(\eta), \xi_c(\eta), \delta_c(\eta)) = 0, \quad \partial_\xi a_0(\eta, \boldsymbol{w}^*(\eta), \xi_c(\eta), \delta_c(\eta)) = 0,$$

taking the derivative of these relations with respect to η (and using $'$ to indicate derivatives with respect to η), we find that $\delta_c'(0)$ and $\xi_c'(0)$ satisfy the following system of coupled partial differential equations

$$\partial_\eta a_0 + \boldsymbol{w}^{*\prime}(0) \cdot \nabla_{\boldsymbol{w}^*} a_0 + \delta_c'(0) \partial_\delta a_0 = 0,$$
$$\partial_{\eta\xi}^2 a_0 + \boldsymbol{w}^{*\prime}(0) \cdot \nabla_{\boldsymbol{w}^*} \partial_\xi a_0 + \xi_c'(0) \partial_{\xi\xi}^2 a_0 + \delta_c'(0) \partial_{\xi\delta}^2 a_0 = 0.$$

Using the expansion in Eq. (17), this reduces to

$$H_1 + \boldsymbol{w}^{*\prime}(0) \cdot \nabla_{\boldsymbol{w}^*} H_0 + \delta_c'(0) \partial_\delta H_0 = 0,$$
$$\partial_\xi H_1 + \boldsymbol{w}^{*\prime}(0) \cdot \nabla_{\boldsymbol{w}^*} \partial_\xi H_0 + \xi_c'(0) \partial_{\xi\xi}^2 H_0 + \delta_c'(0) \partial_{\xi\delta}^2 H_0 = 0.$$

In particular, we deduce the general formula

$$\delta_c'(0) = -\frac{H_1 + \boldsymbol{w}^{*\prime}(0) \cdot \nabla_{\boldsymbol{w}^*} H_0}{\partial_\delta H_0}, \tag{22}$$

where H_1 and H_0 are evaluated at $(\boldsymbol{w}^*(0), \xi_c(0), \delta_c(0))$. In the case of a surface-driven symmetry breaking and in the 1D-1D case, using the formula Eqs. (18)-(19),

$$\delta_c'(0) = -\frac{\mathrm{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A} + \boldsymbol{u_S^{*\prime}}(0) \cdot \nabla_{\boldsymbol{u}^*} |\boldsymbol{P_0}|}{\partial_\delta |\boldsymbol{P_0}|}, \tag{23}$$

and, for bulk-driven symmetry breaking, $(\boldsymbol{P_0}, \boldsymbol{A}, \boldsymbol{u_S^*}(0))$ is replaced by $(\boldsymbol{W_0}, \tilde{\boldsymbol{B}}, \tilde{\boldsymbol{u}_B^*}(0))$. In the 1D-2D case, we recall that the two layers must share the same equilibria, so $\boldsymbol{w}^{*\prime}(0) = 0$ and, in the surface-driven symmetry breaking situation, the formula Eq. (22) thus simplifies to

$$\delta_c'(0) = -\frac{\mathrm{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A}}{\partial_\delta |\boldsymbol{P_0}|}. \tag{24}$$

Finally, in the 1D-2D bulk-driven symmetry breaking situation, we need to take into account the additional hyperbolic cosine term in Eq. (20), and finally obtain

$$\delta_c'(0) = -\frac{\mathrm{co}(\boldsymbol{W_0}) \cdot \tilde{\boldsymbol{B}} \cosh(H\sqrt{\boldsymbol{R_0}})}{\partial_\delta |\boldsymbol{W_0}|}. \tag{25}$$

One surprising consequence of the formula Eq. (23) (and similarly for its variants Eqs. (24)-(25)) is that at first order when $\eta \ll 1$, the function $\eta \mapsto \delta_c(\eta)$ does not depend on the bulk (resp. surface)

parameters. When $\boldsymbol{u_S^*}'(0) = 0$, it is an intrinsic property of the surface (resp. bulk) system. When $\boldsymbol{u_S^*}'(0) \neq 0$, as explained in Section 4.3, there is an additional contribution, but this only depends on the difference between the equilibrium concentrations of the bulk and the surface.

We recall that at $(\xi_c(0), \delta_c(0))$, $|\boldsymbol{P_0}| = \partial_\xi |\boldsymbol{P_0}| = 0$ and $|\boldsymbol{P_0}| \geq 0$ elsewhere for surface driven symmetry breaking. Moreover, patterning occurs in the surface system as soon as there exist parameters (ξ, δ) such that $|\boldsymbol{P_0}|(\xi, \delta) < 0$. Consequently, when $\partial_\delta |\boldsymbol{P_0}| < 0$, patterning occurs in the surface layer independently for $\delta > \delta_c(0)$ and when $\delta < \delta_c(0)$ if $\partial_\delta |\boldsymbol{P_0}| > 0$. The second case is typically encountered when δ is the diffusion parameter of the cells in a chemotaxis system, whereas the first case corresponds, for instance, to a classical activator-inhibitor system with δ the diffusion parameter of the inhibitor. Following this observation, one can conclude that the coupling enhances patterning when $\delta_c'(0)$ and $\partial_\delta |\boldsymbol{P_0}|$ have the same sign. For instance, when $\partial_\delta |\boldsymbol{P_0}| < 0$, this would imply that $\delta_c(\eta) < \delta_c(0)$ for η small enough. Due to Formula (23), this occurs if and only if

$$\mathrm{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A} + \boldsymbol{u_S^*}'(0) \cdot \nabla_{\boldsymbol{u^*}} |\boldsymbol{P_0}| < 0. \tag{26}$$

The increased or decreased ability to form Turing patterns thus depends on the balance between the two terms in Eq. (26). The second term may be non-zero only in the case where the two layers have different equilibrium concentrations. Note that this can only happen in the 1D-1D case. The contribution of this term is difficult to estimate in full generality, so we will give some examples in Section 4.3. We now focus on the contribution of the first term, theoretically and for various examples.

4.2   Global equilibrium

4.2.1   Theoretical considerations with equal surface and bulk equilibrium concentrations

Let us assume that $\boldsymbol{u_S^*}(0) = \boldsymbol{\tilde{u}_B^*}(0)$ and hence focus on the contribution of the first term in (26). That is, we want to compute the sign of

$$I_1 := \mathrm{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A}. \tag{27}$$

Note that in the 1D-1D model, the surface and bulk are exchangeable so the analysis in this section can be immediately translated to 1D-1D bulk-driven symmetry breaking. Moreover, owing to Eq. (24), the analysis in this section also applies to surface-driven symmetry breaking in the 1D-2D case. The bulk-driven symmetry breaking situation in the 1D-2D model will be discussed in Section 4.4.1.

Let us first consider the case $\boldsymbol{A} = \boldsymbol{I}$. The quantity (27) is thus equal to the sum of the n principal minors associated with the principal submatrices of order $(n-1)$. It can be shown that this quantity is equal to the product of the $(n-1)$ non-zero eigenvalues of $\boldsymbol{P_0}$ [41, Section 7.1]. By definition, at $(\xi_c(0), \delta_c(0))$, all these eigenvalues have a positive real part and thus we conclude that $I_1 \geq 0$ and thus $\partial_\delta |\boldsymbol{P_0}|$ and $\delta_c'(0)$ have opposite signs. This implies that the coupling always degrades the ability to form patterns. In order to enhance patterning, one needs to choose a matrix $\boldsymbol{A}$ whose elements corresponding to negative cofactors are non-negative and large.

For pure reaction-diffusion systems – that is $\boldsymbol{D_S}$ is diagonal with nonnegative coefficients – such

cofactors always exist since

$$\partial_\xi |\boldsymbol{P_0}| = \text{co}(\boldsymbol{P_0}) \cdot \boldsymbol{D_S} = 0,$$

which implies that at least one diagonal cofactor is negative. In particular, it is important to note that since the first order approximation $\delta_c'(0)$ is directly proportional to the norm of $\boldsymbol{A}$, it can be made as small as possible by an appropriate scaling of the exchange rates. In theory, this fact thus allows the possibility that patterning can be enhanced by the layer-coupling given an arbitrarily small diffusion of the reactive species. Compared to the Turing theory for reaction-diffusion systems, the differential transport (i.e. the fact that one of the interacting species has a higher exchange rate through the interface) can compensate for, or even replace, the usually theoretically necessary differential diffusion for the classical version of the Turing mechanism, which has long been a subject of debate in real-world biological systems [4, 40, 37, 61].

In the following section, we will illustrate these phenomena through various examples in the 1D-1D situation.

4.2.2 Two-species 1D-1D reaction-diffusion systems

For this example, we consider only two reacting species with concentrations

$$\boldsymbol{u_B} = (u_B, v_B), \quad \boldsymbol{u_S} = (u_S, v_S),$$

which share the same equilibrium. Thus we can, without loss of generality, consider only the surface-driven symmetry breaking situation. In the rest of this article, the subscript S will refer to surface quantities and the subscript B refers to bulk quantities. In many cases, one of the two layers plays no role so we will omit this index when no confusion is possible. In the present example, in the surface-driven symmetry breaking situation, the bulk system plays no role at first order and we will thus denote for simplicity $(u_S, v_S) \equiv (u, v)$. Similarly, the surface reaction function is denoted by $\boldsymbol{f_S} \equiv (f, g)$ and u and v subscripts will denote partial differentiation.

As in the classical Turing theory, let us assume that the reaction functions satisfy the Turing conditions [45, Eqs. (2.31)]. The bifurcation parameter $\delta \equiv d_v$ is the diffusion parameter of the v-species (the inhibitor) while the diffusion coefficient of the u-species (the activator) is fixed to $d_u = 1$. Moreover, we assume that $\boldsymbol{A}$ is diagonal with non-negative diagonal coefficients denoted by α, β. In this case,

$$I_1 = \text{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A} = \alpha(\xi\delta - g_v) + \beta(\xi - f_u),$$

and

$$\partial_\delta |\boldsymbol{P_0}| = \xi(\xi - f_u).$$

We recall [45, Eq. (2.25)] that the critical diffusion and critical wave number are linked by the relation

$$\xi_c(0) = \frac{\delta_c(0)f_u + g_v}{2\delta_c(0)},$$

so that

$$\xi_c(0)\delta_c(0) - g_v = \frac{\delta_c(0)f_u - g_v}{2}, \quad \xi_c(0) - f_u = \frac{-\delta_c(0)f_u + g_v}{2\delta_c(0)}.$$

In particular, when $f_u > 0$ and $g_v < 0$, it follows that $\partial_\delta|\boldsymbol{P_0}| < 0$ and Eq. (23) simplifies to

$$\delta_c'(0) = \frac{\alpha\delta_c(0)}{\xi_c(0)}\left(1 - \frac{\beta/\alpha}{\delta_c(0)}\right). \tag{28}$$

In conclusion, $\delta_c'(0) > 0$ if and only if $\delta_c(0) > \frac{\beta}{\alpha}$. Since the classical Turing instability requires that the inhibitor diffusion coefficient is larger than the activator diffusion coefficient, and thus $\delta_c(0) \geq 1$, this condition for $\delta_c(0)$ is fulfilled when $\beta < \alpha$. In other words, the prospect of patterning is reduced if the exchange rate of the activator is higher than the exchange rate of the inhibitor. This is always the case when $\alpha = \beta$ provided $\delta_c(0) > 1$. In order to enhance patterning it is sufficient to take $\beta > \alpha\delta_c(0)$, which can be understood as off-setting a small inhibitor diffusion with a high exchange rate. Note that the quantity $\xi_c(0) - f_u$ is the negative cofactor in the general case outlined in Section 4.2.1. This result is illustrated in Fig. 2 and Video 1 for the following Schnakenberg system [57]:

$$f(u,v) = s(a - u + u^2v), \tag{29a}$$
$$g(u,v) = s(b - u^2v), \tag{29b}$$

where $0 < a < b$ and $s > 0$.

4.2.3 Chemotaxis systems

Let us consider now the bulk-driven symmetry breaking scenario with a population of chemotactic cells in the bulk and two chemical species with equal equilibrium concentrations in the two layers. Let us first consider the simpler case where only one chemical species, the chemo-attractant, diffuses between the two layers. Similarly as before, we omit the B indexing since the surface does not play any role. The chemo-attractant concentration is denoted by u and the attracted species with concentration c is assumed to satisfy Eq. (5) and to produce the chemo-attractant. Both species are assumed to have constant diffusion coefficients, respectively denoted by d_u and d_c. A typical example is the linear Keller-Segel model

$$f(u,c) = ac - bu, \quad r(c) = c(c^* - c)$$

where $a, b, c^* > 0$ so that $f_u < 0, f_c > 0$ and $r_c < 0$ (when evaluated at the equilibrium point c^*).

In this case, the bifurcation parameter $\delta \equiv d_u$ can be taken as the diffusion coefficient of the chemo-attractant. For this type of model, patterning occurs for small values of δ as we now show. The linearized matrix of the bulk system reads,

$$\boldsymbol{W_0}(\xi,\delta) := \xi\boldsymbol{D_B} - \boldsymbol{J_B} = \begin{pmatrix} \xi\delta - f_u & -f_c \\ -\xi h k_u & \xi d_c - r_c \end{pmatrix},$$

Figure 2: Coupling two identical Schnakenberg systems with $a = 02305$, $b = 0.7695$ and $s = 1$. The bifurcation parameter δ is the diffusion coefficient of the v-species. The instability region in the (η, δ) plane is depicted in grey and is computed numerically by applying the Routh-Hurwitz criterion to the fourth order polynomial Eq. (14) for the modes $k_q = q\pi/L$ with $L = 1000$ and $q \in \{0, \ldots, 1500\}$. The thin solid line at the boundary of the instability region is computed by solving numerically Eqs. (15)-(16). The dashed horizontal line indicates the critical bifurcation parameter $\delta_c(0)$ at $\eta = 0$. The thick solid line is the first-order approximation $\delta_c(\eta) = \delta_c(0) + \eta\delta'_c(0)$ computed using Eq. (28). (a) When $\alpha = \beta = 1$, the coupling degrades the ability to form Turing patterns in the sense that a higher diffusion coefficient is needed (i.e. the instability region is reduced when η is increased). (b) When $\alpha = 1$ and $\beta = 30$, the coupling enlarges the parameter space in which Turing patterns are formed. See also Video 1. In both cases the bulk layer is in a non-patterning state with diffusion coefficients $d_{u_B} = 1$ and $d_{v_B} = 15 < \delta_c(0) \simeq 17.3$.

396 and its determinant is given by

$$|\boldsymbol{W_0}|(\xi, \delta) = \xi^2 \delta d_c - (d_c f_u + \delta r_c + h f_c k_u)\xi + f_u r_c$$

Since $f_u, r_c < 0$, and thus $f_u r_c > 0$, a necessary condition for this polynomial (in ξ) to take negative values for $\xi \geq 0$ is that its derivative in 0 is nonpositive, that is $\delta < (h f_c k_u - d_c|f_u|)/|r_c|$. Then, the minimal value of this polynomial is given by $-(d_c f_u + \delta r_c + h f_c k_u)^2/(4\delta d_c) + f_u r_c$. This function of δ is monotonically increasing between 0 and $(h f_c k_u - d_c|f_u|)/|r_c|$ and is negative for δ smaller than a certain value which defines the critical value $\delta_c(0)$ below which patterning occurs. Moreover, a direct computation shows that

$$\partial_\delta |\boldsymbol{W_0}| = \xi(\xi d_c - r_c) > 0,$$
$$\mathrm{co}(\boldsymbol{W_0}) \cdot \tilde{\boldsymbol{B}} = \alpha(\xi d_c - r_c),$$

397 with diagonal matrix $\tilde{\boldsymbol{B}} = \mathrm{diag}(\alpha, 0)$. Consequently, Eq. (23) simplifies to

$$\delta'_c(0) = -\frac{\alpha}{\xi_c(0)} < 0$$

and hence $\partial_\delta|\boldsymbol{W_0}|\delta_c'(0) < 0$. Thus, in the case of chemotaxis, coupling reduces the ability of the system to form patterns.

Let us now consider the more complex case of chemotaxis coupled to a two-species reaction-diffusion system in the form of Eqs. (4)-(5) with $\tilde{\boldsymbol{f}}_{\boldsymbol{B}}(u,v) = (f(u,v), g(u,v))$. As a three-component reaction-cross-diffusion system, this case is more difficult and explicit computations become more complex. In particular, the full dispersion relation for the bulk reads

$$|\lambda\boldsymbol{I} + \boldsymbol{W_0}| = 0,$$

where

$$\boldsymbol{W_0} = \begin{pmatrix} \xi d_u - f_u & -f_v & -f_c \\ -g_u & \xi\delta - g_v & -g_c \\ -\xi h k_u & -\xi h k_v & \xi d_c - r_c \end{pmatrix}.$$

Thus

$$\partial_\delta|\boldsymbol{W_0}| = \xi\Big((\xi d_u - f_u)(\xi d_c - r_c) - \xi h k_u f_c\Big),$$

and when $\boldsymbol{B}$ is diagonal with diagonal coefficients α, β,

$$\mathrm{co}(\boldsymbol{W_0}) \cdot \tilde{\boldsymbol{B}} = \alpha\Big((\xi\delta - g_v)(\xi d_c - r_c) - \xi h k_v g_c\Big) + \beta\Big((\xi d_u - f_u)(\xi d_c - r_c) - \xi h k_u f_c\Big).$$

Assessing the sign of this expression in a general setting is difficult due to the number of degrees of freedom. However, one scenario where the equilibria of the reacting and diffusing species have the same values in the bulk and surface is where the cells are slave, with no feedback on the signalling molecules, so that $f_c = g_c = 0$. Further, the requirement of stability to homogeneous perturbations, i.e. when $\xi = 0$, gives $r_c < 0$, $f_u + g_v < 0$, and $f_u g_v - g_u f_v > 0$, with the latter two constituting standard Turing conditions. With the chemoattractant species, v, as the activator, and the assumption that u, v are a Turing pair in the absence of chemotaxis, so that u is the inhibitor, entails we additionally have $g_v > 0 > f_u$ and hence $g_u f_v < f_u g_v < 0$. Thus the requirement that $|\boldsymbol{W_0}| = 0$ at the critical point gives

$$(\xi d_u - f_u)(\xi\delta - g_v)(\xi d_c - r_c) - f_v g_u(\xi d_c - r_c) = 0,$$

and hence the coefficient of α in $\mathrm{co}(\boldsymbol{W_0}) \cdot \tilde{\boldsymbol{B}}$ is given by

$$(\xi\delta - g_v)(\xi d_c - r_c) = f_v g_u \left(\frac{\xi d_c + |r_c|}{\xi d_u + |f_u|}\right) < 0.$$

In contrast the coefficient of β is given by $(\xi d_u + |f_u|)(\xi d_c + |r_c|) > 0$. Hence, whether the bilayer enhances patterning depends on the between-layer transport of the inhibitor, u here, relative to that of the activator, v here, with sufficiently low relative activator between-layer transport acting to increase the ability to form patterns. We also note that the chemotactic parameters occur only in the grouping $(\xi d_c + |r_c|)$, which factors out of both the coefficients of α and β so that, as might be expected

421 given that the cells are slave to the signalling molecules, the chemotatic properties of the cells do not
422 influence these observations.

4.3 Asymmetric equilibria in the 1D-1D model

424 Let us now focus on the contribution of the second term in Eq. (26),

$$I_2 := \boldsymbol{u_S^*}'(0) \cdot \nabla_{\boldsymbol{u}^*} |\boldsymbol{P_0}|, \tag{30}$$

425 for cases where the equilibria in the uncoupled bulk and surface layers are different but the coupled
426 system has a homogeneous equilibrium (and thus necessarily in a 1D-1D model). Without loss of
427 generality, we consider the surface-driven patterning situation. Crucially, we need to compute the
428 derivative $\boldsymbol{u_S^*}'(0)$ of the modified equilibrium concentration induced by the coupling. This follows by
429 a simple application of the implicit function theorem to the function

$$\boldsymbol{f} : (\eta, \boldsymbol{u_S}, \boldsymbol{u_B}) \mapsto \Big(\boldsymbol{f_S}(\boldsymbol{u_B}) + \eta A(\boldsymbol{u_B} - \boldsymbol{u_S}), \boldsymbol{f_B}(\boldsymbol{u_B}) + \eta\tilde{B}(\tilde{\boldsymbol{u}}_S - \boldsymbol{u_B})\Big).$$

430 The determinant of the Jacobian matrix $\boldsymbol{J_f}$ of $\boldsymbol{f}$ with respect to $(\boldsymbol{u_S}, \boldsymbol{u_B})$ at $\eta = 0$ is given by

$$|\boldsymbol{J_f}| = |\boldsymbol{J_S}||\boldsymbol{J_B}|.$$

431 Since we only consider diffusion-driven instabilities, both Jacobian matrices on the right-hand side are
432 invertible (as they must have eigenvalues with strictly negative real parts) and hence so is $\boldsymbol{J_f}$. Hence,
433 applying the implicit function theorem, there exist smooth functions $\eta \mapsto \boldsymbol{u_S^*}(\eta)$ and $\eta \mapsto \boldsymbol{u_B^*}(\eta)$ close
434 to $\eta = 0$. Next, we can compute the derivative of these functions at $\eta = 0$ by simply differentiating
435 with respect to η the relation

$$\boldsymbol{f}\big(\eta, \boldsymbol{u_S^*}(\eta), \boldsymbol{u_B^*}(\eta)\big) = 0.$$

Their explicit expression can be found by solving the linear system:

$$\boldsymbol{u_S^*}'(0) = \boldsymbol{J_S^{-1}} A\big(\boldsymbol{u_S^*}(0) - \tilde{\boldsymbol{u}}_B^*(0)\big),$$
$$\boldsymbol{u_B^*}'(0) = \boldsymbol{J_B^{-1}} \tilde{B}\big(\boldsymbol{u_B^*}(0) - \tilde{\boldsymbol{u}}_S^*(0)\big).$$

436 Consequently, the first-order contribution due to $\boldsymbol{u_S^*}'(0)$ depends only on the difference between the
437 equilibrium concentrations and the reaction term of the surface but it does not involve the reaction
438 term of the bulk layer.

439 Note that using Jacobi's formula again, it follows that, for a given vector $\boldsymbol{p} = (p_1, \ldots, p_n)^{\mathrm{T}}$,

$$\boldsymbol{p} \cdot \nabla_{\boldsymbol{u}^*} |\boldsymbol{P_0}| = \mathrm{co}(\boldsymbol{P_0}) \cdot (\boldsymbol{p} \cdot \nabla_{\boldsymbol{u}^*})\boldsymbol{P_0},$$

440 where we use the operator $\boldsymbol{p} \cdot \nabla_{\boldsymbol{u}^*} := \sum_{k=1}^{n} p_k \partial_{u_k^*}$.
441 Consequently, similarly to the previous situation, we can also rewrite the second term in Eq. (26)

as an inner product with the comatrix of $\boldsymbol{P_0}$:

$$I_2 = \boldsymbol{u_S^{*}}'(0) \cdot \nabla_{\boldsymbol{u^*}}|\boldsymbol{P_0}| = \mathrm{co}(\boldsymbol{P_0}) \cdot \left(\boldsymbol{J_S^{-1}}\boldsymbol{A}(\boldsymbol{u_S^*}(0) - \boldsymbol{u_B^*}(0)) \cdot \nabla_{\boldsymbol{u^*}}\right)\boldsymbol{P_0}.$$

Thus, this contribution will dominate when the uncoupled equilibrium concentrations in the two layers are sufficiently different. This case is illustrated in Fig. 3. The situation is simpler in the strong coupling case, as explained in the forthcoming Section 5, but nonetheless progress can be made in simple examples, as we now show.

For example, we consider the Schnakenberg kinetics of Eq. (29), with $s = 1$, a coupling matrix $\boldsymbol{A} = \boldsymbol{B} = \mathrm{diag}(\alpha, 0)$ and, to force a difference in the surface and bulk concentrations, we take the equivalent parameter for a in the bulk to be $a_B \neq a$, with all parameters positive. The equilibria are given by the solution of the four simultaneous algebraic equations governing $u_S^*, v_S^*, u_B^*, v_B^*$, namely

$$u_S^* = a + b + \frac{(a_B - a)\alpha\eta}{1 + 2\alpha\eta}, \quad u_B^* = a_B + b + \frac{(a - a_B)\alpha\eta}{1 + 2\alpha\eta}, \quad v_S^* = \frac{b}{u_S^{*2}}, \quad v_B^* = \frac{b}{u_B^{*2}}.$$

Thus

$$u_S^{*'} = \frac{\alpha(a_B - a)}{(1 + 2\alpha\eta)^2}, \quad v_S^{*'} = -\frac{2b}{u_S^{*3}}u_S^{*'}.$$

Hence, explicitly evaluating Eq. (30) with d_u, d_v the diffusion coefficients of the respective species, we have

$$I_2 = u_S^{*'}\left(\frac{\partial|\boldsymbol{P_0}|}{\partial u_S^*} - \frac{2b}{u_S^{*3}}\frac{\partial|\boldsymbol{P_0}|}{\partial v_S^*}\right) = u_S^{*'}\left(2\xi(d_u u_S^* - d_v v_S^*) + 2u_s^* + \frac{4\xi b d_v}{u_S^{*2}}\right)$$

$$= u_S^{*'}\left(2\xi\left(d_u u_S^* + \frac{b d_v}{u_S^{*2}}\right) + 2u_S^*\right),$$

with the final equality arising from noting that $v_S^* = b/u_S^{*2}$. Hence, for this case we have

$$\mathrm{co}(\boldsymbol{P_0}) \cdot \boldsymbol{A} + \boldsymbol{u_S^{*}}'(0) \cdot \nabla_{\boldsymbol{u^*}}|\boldsymbol{P_0}| = \alpha(u_S^{*2} + \xi d_v) + \alpha(a_B - a)\left(2\xi\left(d_u u_S^* + \frac{b d_v}{u_S^{*2}}\right) + 2u_S^*\right),$$

where we recall that this expression is evaluated at the critical point $\eta = 0, \xi = \xi_c(0), d_v = \delta_c(0)$ and $u_S^*(0) = a + b$. Thus, when $a_B > a$, this expression is always positive and the layering acts to inhibit pattern, whilst for $a > a_B$ sufficiently large, we have that the difference between the resulting surface and bulk equilibrium concentration levels will enhance prospective patterning.

(a)   (b)

Figure 3: Coupling two different Schnakenberg systems with parameters $\boldsymbol{A} = \boldsymbol{B} = \boldsymbol{I}$, $a = 0.15$, $b = 0.2$, $s = 0.5$, $d_{u_B} = 1$, $d_{v_B} = 15 < \delta_c[\boldsymbol{f_B}] \simeq 25.7$ for the bulk (non-patterning state) and $a = 0.2305$, $b = 0.7695$, $s = 2$ for the surface. The instability region in grey and its boundary are computed numerically as in Fig. 2. The coupling enhances patterning for small and large η but reduces patterning for intermediate values. The first-order approximation at $\eta = 0$ is computed using Eq. (23). The asymptotic value is indicated by the dotted line and is computed using Eq. (35). (a) $\eta \in (0, 5)$. (b) $\eta \in (0, 200)$.

4.4 The case of the 1D-2D model

4.4.1 Large and small depth

In the 1D-2D case, the boundary condition Eq. (3) imposes a global homogeneous equilibrium identical in both layers. Thus, for bulk-driven patterns, using (20), the formula Eq. (22) reduces to

$$\delta'_c(0) = \frac{\text{co}\left(\boldsymbol{D_B}\sqrt{\boldsymbol{R_0}}\sinh(H\sqrt{\boldsymbol{R_0}})\right) \cdot \tilde{\boldsymbol{B}}\cosh(H\sqrt{\boldsymbol{R_0}})}{\partial_\delta|\boldsymbol{D_B}\sqrt{\boldsymbol{R_0}}\sinh(H\sqrt{\boldsymbol{R_0}})|}, \tag{31}$$

where $\boldsymbol{R_0} = \boldsymbol{D_B^{-1}}\boldsymbol{Q_0}$, $\boldsymbol{Q_0} = \xi\boldsymbol{D_B} - \boldsymbol{J_B}$ and the notation $\sqrt{\boldsymbol{R_0}}$ denotes any square root of $\boldsymbol{R_0}$. We recall that

$$\sqrt{\boldsymbol{R_0}}\sinh(H\sqrt{\boldsymbol{R_0}}) = \sum_{k=1}^{+\infty}\frac{H^{2k-1}}{(2k-1)!}\boldsymbol{R_0^k} =: F_1(H, \boldsymbol{R_0}), \quad \cosh(H\sqrt{\boldsymbol{R_0}}) = \sum_{k=0}^{+\infty}\frac{H^{2k}}{(2k)!}\boldsymbol{R_0^k} =: F_2(H, \boldsymbol{R_0}),$$

and thus the expression in Eq. (31) is a function of $\boldsymbol{R_0}$ only. When $H \to 0$, one can compute the following equivalent

$$\delta'_c(0) \sim \frac{1}{H}\frac{\text{co}(\boldsymbol{Q_0}) \cdot \tilde{\boldsymbol{B}}}{\partial_\delta|\boldsymbol{Q_0}|},$$

that is, up to a positive factor H^{-1}, this is the same as in the 1D-1D case. This scaling is expected and consistent with the analysis for the reduction from 2D to 1D obtained when $H \to 0$.

When H is large, we also expect that the bulk will pattern on its own without influence of the

surface, that is $\delta_c'(0) \to 0$. Using Jacobi's formula and the series expansion in Eq. (17) we note that

$$\partial_\delta |\boldsymbol{D_B}\sqrt{\boldsymbol{R_0}}\sinh(H\sqrt{\boldsymbol{R_0}})| = \mathrm{co}\left(\boldsymbol{D_B}F_1(H,\boldsymbol{R_0})\right)\cdot\left(\partial_\delta\boldsymbol{D_B}F_1(H,\boldsymbol{R_0}) + H\boldsymbol{D_B}\partial_\delta\boldsymbol{R_0}\tilde{F}_2(H,\boldsymbol{R_0})\right),$$

where

$$\tilde{F}_2(H,\boldsymbol{R_0}) := \sum_{k=0}^{+\infty}\frac{k+1}{2k+1}\frac{H^{2k}}{(2k)!}\boldsymbol{R_0^k}.$$

We can expect that, generically when H is large, the denominator in Eq. (31) will behave as H multiplied by a quantity of the same order of magnitude as the numerator $\mathrm{co}\left(\boldsymbol{D_B}F_1(H,\boldsymbol{R_0})\right)\cdot\tilde{\boldsymbol{B}}F_2(H,\boldsymbol{R_0})$. With this formal argument we thus expect that $\delta_c'(0)$ generically decreases as H^{-1}.

4.4.2 Two-species case

For a two-species system like the one in Section 4.2.2, we can make the above computation exact for any H. This follows from the fact that

$$\boldsymbol{R_0} = \begin{pmatrix} \xi - d_u^{-1}f_u & -d_u^{-1}f_u \\ -d_v^{-1}g_u & \xi - d_v^{-1}g_v \end{pmatrix}$$

is nilpotent, satisfying $\boldsymbol{R_0^2} = \boldsymbol{0}$ (for example by noting $\mathrm{Tr}(\boldsymbol{R_0}) = \det(\boldsymbol{R_0}) = 0$ at the critical bifurcation point $\xi = \xi_c(0)$, $d_v = \delta_c(0)$, and then using the Cayley-Hamilton theorem). Further note that, in this case, the matrix $\boldsymbol{R_0}$ has no square root. Thus it follows that,

$$F_1(H,\boldsymbol{R_0}) = H\boldsymbol{R_0}, \quad F_2(H,\boldsymbol{R_0}) = \boldsymbol{I} + \frac{H^2}{2}\boldsymbol{R_0},$$

and Eq. (31) simplifies to

$$\delta_c'(0) = \frac{1}{H}\frac{\mathrm{co}(\boldsymbol{D_B}\boldsymbol{R_0})\cdot(\boldsymbol{B} + \frac{H^2}{2}\boldsymbol{B}\boldsymbol{R_0})}{\partial_\delta|\boldsymbol{D_B}\boldsymbol{R_0}|}.$$

We note that since $|\boldsymbol{R_0}| = 0$,

$$\mathrm{co}(\boldsymbol{D_B}\boldsymbol{R_0})\cdot\boldsymbol{B}\boldsymbol{R_0} = \frac{\mathrm{d}}{\mathrm{d}h}\det(\boldsymbol{D_B}\boldsymbol{R_0} + h\boldsymbol{B}\boldsymbol{R_0})\Big|_{h=0} = \left(\mathrm{co}(\boldsymbol{D_B})\cdot\boldsymbol{B}\right)|\boldsymbol{R_0}| = 0.$$

Moreover, $\boldsymbol{D_B}\boldsymbol{R_0} = \boldsymbol{Q_0}$ so it follows that

$$\delta_c'(0) = \frac{1}{H}\frac{\mathrm{co}(\boldsymbol{Q_0})\cdot\boldsymbol{B}}{\partial_\delta|\boldsymbol{Q_0}|},$$

and this exact formula remains valid for all H.

Remark 2. *It should be noted that the eigenvalues of $\boldsymbol{R_0}$ at $\xi = \xi_c(0)$ and $\delta = \delta_c(0)$ are the solutions of*

$$|(\xi_c(0) - \lambda)\boldsymbol{D_B} - \boldsymbol{J_B}| = |\boldsymbol{Q_0}|\left(\xi_c(0) - \lambda, \delta_c(0)\right) = 0,$$

so $\lambda = 0$ *is always an eigenvalue of* $\boldsymbol{R_0}$*. In the two-species case, this is the only eigenvalue since the polynomial* $\xi \mapsto |\boldsymbol{Q_0}|\big(\xi, \delta_c(0)\big)$ *is of degree 2, vanishes at* $\xi_c(0)$ *and is non-negative in a neighbourhood of this root. Consequently,* $\boldsymbol{R_0}$ *is a nilpotent matrix, as computed before. The same situation could possibly happen for an arbitrary, but even, number of species, although it would require some fine tuning of the parameters, in which case the functions* F_1 *and* F_2 *would actually be polynomials in* $\boldsymbol{R_0}$*. However, for an odd number of species, the polynomial* $\xi \mapsto |\boldsymbol{Q_0}|\big(\xi, \delta_c(0)\big)$ *is an even-degree polynomial and thus it must have at least one other root* $\xi_0 < \xi_c(0)$*. Consequently, the matrix* $\boldsymbol{R_0}$ *always has a positive eigenvalue* $\lambda = \xi_c(0) - \xi_0 > 0$ *so the functions* F_1 *and* F_2 *cannot be polynomials. This observation suggests that, in layered systems, 2-species models may not be representative of higher species models since the impact of nilpotency in 2-species models is a specific feature that may reflect a genuinely different behaviour compared to higher species models.*

4.4.3 Patterning dynamics

Since Eq. (23) involves the characteristics of the patterning layer only, the critical bifurcation curve in the surface-patterning case for the 1D-2D system has the same first-order approximation as in the 1D-1D system. However, although the 1D-2D and 1D-1D systems look similar from the point of view of patterning conditions, the structure of the patterns is different depending on whether symmetry-breaking is driven by the surface or the bulk. As may be expected, in the former case, patterning preferentially occurs at the interface and is not observed to extend into the depth of a non-patterning bulk system, at least for the cases we will consider below. For bulk-driven self-organisation, patterns can be observed throughout the bulk layer, with the potential of creating surface concentration inhomogeneities when these bulk patterns are formed close to the interface.

These phenomena are illustrated in Fig. 4 and Videos 2-3 for a reaction-diffusion-chemotaxis model (4)-(5) using the following piece-wise linear reaction functions: in the surface the reaction term is given by $\boldsymbol{f_S}(u, v) = (f(u, v), g(u, v))$ with

$$f(u, v) = \phi(a_0 + a_1 u - bv) - a_2 u, \tag{32a}$$

$$g(u, v) = \phi(b_0 + c_1 v + du) - c_2 v. \tag{32b}$$

Here signalling molecules are produced by a fixed density of surface cells in an autocatlytic feedback response, with the continuous piece-wise linear function $\phi(\zeta) = \max(0, \min(\zeta, M))$ for a given $M > 0$, and parameters $a_0, a_1, b, a_2, b_0, c_1, d, c_2 \geq 0$. In this definition the value M is chosen sufficiently large so that the reaction function behaves as the identity function around the equilibrium value (obtained by solving the linear system, with ϕ taken as the identity). The threshold values 0 and M are added to ensure the positivity of the solutions and to prevent blow-up. This model has been introduced in the seminal work [30]. In the bulk we consider the same system but where the chemical species are produced by the chemotactic cells without feedback, so that $\boldsymbol{f_B}(u, v, c) = (f(u, v, c), g(u, v, c), r(c))$

with

$$f(u, v, c) = \phi(a_0 + a_1 c - bv) - a_2 u, \tag{33a}$$

$$g(u, v, c) = \phi(b_0 + c_1 c + dc) - c_2 v, \tag{33b}$$

$$r(c) = r_0 c(c^* - c). \tag{33c}$$

The parameters in both systems and the equilibrium cell concentration c^* are chosen so that $\boldsymbol{u}_{\boldsymbol{S}}^*(0) = \tilde{\boldsymbol{u}}_{\boldsymbol{B}}^*(0)$. For a given set of parameters, the critical diffusion of the v-species in the surface and the bulk are, respectively, denoted in the caption of the figures by $d_{v_S}^c$ and $d_{v_B}^c$.

In this situation, since one layer is in a patterning state, spot patterns appear simultaneously throughout the patterning layer and propagate through the interface to the non-patterning layer. The amplitude of the patterns created in the non-patterning layer depends on the coupling strength η, and it has been confirmed numerically that they are directly proportional (results not shown). In particular, since we consider the case of small η in all of the numerical experiments presented in this section, the amplitude of the patterns in the non-patterning layer is typically much lower than in the patterning layer. For instance in Fig. 4b and Fig. 5, the amplitude of the surface patterns is about one percent or less of the value of the homogeneous steady state, but it would be proportionally larger for larger values of the coupling strength η (in the cases shown, we took $\eta = 0.01$). The same is true for the bulk amplitude in Fig. 4a.

The patterning dynamics may be different when the patterns are created by coupling two non-patterning layers with sufficiently large and different exchange rates, i.e. by taking $\boldsymbol{A}$ or $\boldsymbol{B}$ sufficiently different from the identity, similarly to the system in Section 4.2.2 and Fig. 2b. In particular, Fig. 5 shows the result for the piece-wise linear system (32)-(33) with $\boldsymbol{A} = \boldsymbol{I}$ and $\boldsymbol{B} = \text{diag}(1, \beta_B)$. Starting from a random perturbation of the homogeneous equilibrium, when the exchange rate β_B of the inhibitor v is sufficiently large, patterns appear locally at the interface and then propagate throughout the bulk and the surface (see also Video 4).

The pattern propagation in the bulk (Fig. 5) can be understood as a multi-stability phenomenon: although linearly stable, the homogeneous state in the bulk is only one stable equilibrium of the PDE, and patterns in the uncoupled system can be obtained by choosing an initial condition very far from the homogeneous equilibrium. In the simulation, the initial condition is a small random perturbation of the homogeneous state and the bulk independently thus does not typically escape from this linearly stable state. However, when coupling is added, the interface plays the role of an active boundary able to locally drive the system far from the homogeneous equilibrium before spreading throughout the domain, in a similar fashion as the models introduced by [39, 47].

(a) Surface-driven patterning   (b) Bulk-driven patterning

Figure 4: Surface and bulk patterning with $\eta = 0.01$ in the pseudo-linear 1D-2D system (32)-(33) computed in a 128×128 grid at time $t = 500$ after a steady state is reached. Surface parameters: $a_0 = 1$, $a_1 = 2$, $b = 2$, $a_2 = 2$, $b_0 = 0.25$, $c_1 = 1$, $d = 1$, $c_2 = 2$. Bulk parameters: $r_0 = 0.2$, $a_0 = 3$, $a_1 = 8$, $b = 2$, $a_2 = 7$, $b_0 = 0.75$, $c_1 = 0.5$, $c_2 = 1.5$, $d = 1$, $c^* = 0.25$, $\chi = 50$. (a) Surface-driven patterning: $d_{u_S} = 1$, $d_{v_S} = 6.41 > d_{v_S}^c \simeq 5.85$, $d_{u_B} = 10$, $d_c = 10$, $d_{v_B} = 6 < d_{v_B}^c \simeq 8$. (b) Bulk-driven patterning: $d_{u_S} = 1$, $d_{v_S} = 2.91 < d_{v_S}^c$, $d_{u_B} = 10$, $d_c = 10$, $d_{v_B} = 60 > d_{v_B}^c$. Note that for both plots the original y-coordinate is used, with the no-flux condition at $y = 0$ and the surface at $y = H = 50$. A brief description of the numerical method used can be found in Appendix A as well as a link to the numerical code freely available online.

5   Strong coupling in the 1D-1D case

5.1   Asymptotic reduction

When $\eta \geq 1$ and for sufficiently nice $\boldsymbol{A}$ and $\boldsymbol{B}$, the concentrations $\tilde{\boldsymbol{u}}_B$ and $\boldsymbol{u_S}$ are expected to converge towards a common value $\boldsymbol{u}$. For simplicity, we consider here only the case $m = n$, but the extension to the general case would be straightforward. Thus, splitting $\boldsymbol{u_S}$ and $\boldsymbol{u_B}$ as

$$\boldsymbol{u_S} = \boldsymbol{u_p} + \boldsymbol{u_m}, \quad \boldsymbol{u_B} = \boldsymbol{u_p} - \boldsymbol{u_m},$$

where

$$\boldsymbol{u_p} = \frac{\boldsymbol{u_S} + \boldsymbol{u_B}}{2}, \quad \boldsymbol{u_m} = \frac{\boldsymbol{u_S} - \boldsymbol{u_B}}{2},$$

Figure 5: Wave pattern with $\eta = 0.01$ computed on a 512×512 grid until a steady state is reached. Surface parameters: $d_{u_S} = 1$, $d_{v_S} = 2.91 < d_{v_S}^c$. Bulk parameters: $\chi = 120$, $d_{u_B} = 25$, $d_c = 25$, $d_{v_B} = 35 < d_{v_B}^c \simeq 37.5$, $\beta_B = 180$. The other parameters are the same as in Fig. 4.

provided that $\boldsymbol{u_S}$ and $\boldsymbol{u_B}$ remain uniformly bounded, the half difference $\boldsymbol{u_m}$ satisfies

$$\partial_t \boldsymbol{u_m} = -\eta(\boldsymbol{A} + \boldsymbol{B})\boldsymbol{u_m} + \mathcal{O}(1).$$

As a consequence, if $\boldsymbol{A}$ and $\boldsymbol{B}$ are such that their sum $\boldsymbol{A} + \boldsymbol{B}$ has only non-negative eigenvalues (for instance when $\boldsymbol{A}$ and $\boldsymbol{B}$ are both symmetric and positive definite), it follows that $\boldsymbol{u_m} = \mathcal{O}(\eta^{-1})$. Up to an error of order η^{-1}, the concentrations $\boldsymbol{u_S}$ and $\boldsymbol{u_B}$ are thus equal to their average $\boldsymbol{u_p}$. The dynamics of $\boldsymbol{u_p}$ can be found by multiplying Eq. (6) by $\boldsymbol{A}^{-1}$ and Eq. (7) by $\boldsymbol{B}^{-1}$ and summing the two resulting equations so that the exchange term cancels out. Again, up to an error of order η^{-1}, the behaviour of $\boldsymbol{u_p}$ is given by the following equation

$$(\boldsymbol{A}^{-1} + \boldsymbol{B}^{-1})\partial_t \boldsymbol{u} = \boldsymbol{A}^{-1}\boldsymbol{f_S}(\boldsymbol{u}) + \boldsymbol{B}^{-1}\boldsymbol{f_B}(\boldsymbol{u}) + \nabla \cdot \big((\boldsymbol{A}^{-1}\boldsymbol{D_S} + \boldsymbol{B}^{-1}\boldsymbol{D_B})\nabla\boldsymbol{u}\big).$$

For instance, in the simple case where $\boldsymbol{A} = \boldsymbol{B}$, corresponding to no sinks or sources at the interface, $\boldsymbol{u}$ satisfies

$$\partial_t \boldsymbol{u} = \frac{\boldsymbol{f_S}(\boldsymbol{u}) + \boldsymbol{f_B}(\boldsymbol{u})}{2} + \nabla \cdot \left(\frac{\boldsymbol{D_S} + \boldsymbol{D_B}}{2}\nabla\boldsymbol{u}\right). \tag{34}$$

A further case with a simple reduction occurs when $\boldsymbol{f_S} = \boldsymbol{f_B}$ so that the reaction term remains unchanged while the diffusion matrix becomes a linear combination of the surface and bulk diffusion matrices. In addition, for a mass conserving system with $\boldsymbol{f_S} = -\boldsymbol{f_B}$, the reduced equation Eq. (34) simply becomes a pure (cross-)diffusion. However, in full generality, the behaviour of the solution does not readily simplify, since the properties of the new reaction term cannot be expected to immediately follow from the properties of the surface and bulk reaction terms.

5.2 Patterning conditions for two-species systems

In the case of two coupled two-species reaction-diffusion systems with $d_{u_S} = d_{u_B} = 1$, the critical bifurcation parameter of the inhibitor in the surface can be simply computed from Eq. (34) and is given by

$$\delta_c(\infty) := \lim_{\eta \to +\infty} \delta_c(\eta) = 2\delta_c \left[\frac{\boldsymbol{f_S} + \boldsymbol{f_B}}{2} \right] - d_{v_B}, \tag{35}$$

where we denote by $\delta_c[\boldsymbol{f}]$ the critical inhibitor diffusion of a reaction-diffusion system with reaction term $\boldsymbol{f} = (f, g)$ and activator diffusion $d_u = 1$. Note that there is a balance between the diffusion coefficients of the two layers. In particular, and as in the small coupling case (Section 4.2.1), the critical diffusion of the v-species in the surface in the coupled system can be made as small as desired, provided that the diffusion in bulk d_{v_B} is large enough.

For later convenience, we recall the following formula corresponding to a standard reaction-diffusion system with $\boldsymbol{f} = (f, g)$ [45, Eq. (2.27)]:

$$\delta_c[\boldsymbol{f}] = \left(\frac{\sqrt{f_u g_v - f_v g_u} + \sqrt{-f_v g_u}}{f_u} \right)^2. \tag{36}$$

5.2.1 Stabilization by coupling

When the two coupled systems are identical, $\boldsymbol{f_S} = \boldsymbol{f_B}$, with a common critical bifurcation parameter for the inihibitor diffusion denoted by $\delta_c(0)$, then, asymptotically when η is large,

$$\delta_c(\infty) = 2\delta_c(0) - d_{v_B}, \tag{37}$$

and in particular, if the bulk is in a non-patterning state, i.e. $d_{v_B} < \delta_c(0)$, then $\delta_c(\infty) > \delta_c(0)$. As might be expected, and similarly to the small coupling case (Section 4.2), this means that a large coupling has a stabilizing effect: a larger diffusion of the inhibitor in the surface is needed to de-stabilize the homogeneous state. This observation is illustrated in Fig. 6a.

More generally, the analysis presented in Section 4 reveals that if a layer is in a patterning state, then for small coupling, the coupled system remains in a patterning state. This can be seen as a consequence of the fact that the derivative $\delta_c'(0)$ of the critical parameter is an intrinsic property of the considered layer. Consequently, if δ is chosen above the critical value $\delta_c(0)$ and $\delta_c'(0) < +\infty$, then the coupled system must remain in a patterning state for η sufficiently small (when $\delta_c'(0) > 0$ and δ is close to $\delta_c(0)$, a first order approximation of this value would be $\eta = (\delta - \delta_c(0))/\delta_c'(0)$). Therefore, if one wants to use the coupling to stabilize a homogeneous state, it is necessary to consider a sufficiently large coupling strength. When the two layers are independently in a patterning state, i. e. when $d_{v_B} > \delta_c[\boldsymbol{f_B}]$ and $d_{v_S} > \delta_c[\boldsymbol{f_S}]$, an asymptotically non-patterning state for the coupled system corresponds to the case where

$$d_{v_S} < \delta_c(\infty) = 2\delta_c \left[\frac{\boldsymbol{f_S} + \boldsymbol{f_B}}{2} \right] - d_{v_B}.$$

It is possible to find such parameters (d_{v_S}, d_{v_B}) if and only if

$$\delta_c \left[\frac{\boldsymbol{f_S} + \boldsymbol{f_B}}{2} \right] > \frac{\delta_c[\boldsymbol{f_B}] + \delta_c[\boldsymbol{f_S}]}{2}. \tag{38}$$

This strict midpoint-concavity property requires that $\boldsymbol{f_B} \neq \boldsymbol{f_S}$. Checking this property for two arbitrary reaction functions heavily depends on the form of these functions. When the reaction function depends linearly on its parameters, using the explicit formula Eq. (36) for the critical diffusion, checking Eq. (38) reduces to checking the convexity properties of this several-variable function (note that midpoint convexity is equivalent to convexity for continuous functions). For instance, it is straightforward to select reaction functions which satisfy this concavity property: one can consider a family of (pseudo)-linear models of the form (32) parametrised by a single parameter $p > 1$ such that the Jacobian matrix of the equilibrium system is given by

$$J_p = \begin{pmatrix} 1 & -1 \\ p & 1 - p \end{pmatrix}. \tag{39}$$

In this case

$$\delta_c = 1 + p + 2\sqrt{p}$$

is a concave function of p. This situation is illustrated in Fig. 6b and Video 5.

5.2.2 Strong coupling patterns

Reciprocally, if $d_{v_B} < \delta_c[\boldsymbol{f_B}]$ and $d_{v_S} < \delta_c[\boldsymbol{f_S}]$ (i.e. the two layers are independently in a non-patterning state), then the homogeneous state of the coupled system, with η large, is unstable whenever $\delta_c(\infty) < d_{v_S} < \delta_c[\boldsymbol{f_S}]$. Using Eq. (35), such a value of d_{v_S} will exist if and only if the following midpoint-convexity condition is satisfied:

$$\delta_c \left[\frac{\boldsymbol{f_S} + \boldsymbol{f_B}}{2} \right] < \frac{\delta_c[\boldsymbol{f_B}] + \delta_c[\boldsymbol{f_S}]}{2}. \tag{40}$$

Once again, it is necessary to consider two different systems in the surface and bulk layers. Note that in contrast to the small coupling case (Section 4.3), the patterning condition (40) is typically more straightforward to ascertain, as it reduces to computing the convexity properties of a function. For instance, the condition (40) is always satisfied for the Schnakenberg (29) [57] and Gierer-Meinhardt [23], [45, Eq. (2.8)] systems, as it can be checked using the formula (36). An example is shown in Fig. 3b and Video 6.

6 Beyond the asymptotic cases

For intermediate values of η many different scenarios are possible. An important point to note is that the Turing conditions for each layer at $\eta = 0$ do not ensure that the coupled system at $\eta > 0$ is stable

(a)   (b)

Figure 6: Large coupling asymptotics. The dotted horizontal line indicates the asymptotic value $\delta_c(\infty)$ computed using Eq. (35). (a) Coupling two identical Schnakenberg systems with the same parameters as in Fig. 2. The asymptotic value of the critical bifurcation parameter is larger than $\delta_c(0)$. (b) Coupling two pseudo-linear systems with Jacobians given by Eq. (39) with $p = 2$ for the bulk and $p = 200$ for the surface. The bulk is in a patterning state with diffusion coefficients $d_{u_B} = 1$ and $d_{v_B} = 10$. Consequently, for small η the coupled system is in a patterning state for all values of d_{v_S}. However, since $\delta_c(\infty) > \delta_c(0)$, for η sufficiently large and $d_{v_S} \in (\delta_c(0), \delta_c(\infty))$, the coupled system is in a non-patterning state and thus the coupling stabilizes the homogeneous equilibrium. See also Video 5.

without diffusion (i.e. that $\xi = 0$ is a stable mode). For instance, taking two identical 1D-1D layers with Jacobian matrix $\boldsymbol{J}$ and $\boldsymbol{A} = \boldsymbol{B}$, as in Eq. (11), the stability of the mode $\xi = 0$ can be checked by computing the eigenvalues of the block matrix

$$\begin{pmatrix} \boldsymbol{J} - \eta\boldsymbol{A} & \eta\boldsymbol{A} \\ \eta\boldsymbol{A} & \boldsymbol{J} - \eta\boldsymbol{A} \end{pmatrix}.$$

Using the formula for block-diagonal determinants, the eigenvalues of this matrix are the eigenvalues of the matrix $\boldsymbol{J}$ and the eigenvalues of the matrix $\boldsymbol{J} - 2\eta\boldsymbol{A}$. If $\boldsymbol{A} \neq \boldsymbol{I}$, then the latter may have non-negative eigenvalues, indicating that the mode $\xi = 0$ is unstable. For instance for $\boldsymbol{A} = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix}$, the mode $\xi = 0$ is unstable for all η between the roots of the polynomial (in η)

$$4\alpha\beta\eta^2 - 2(\alpha g_v + \beta f_u)\eta + |\boldsymbol{J}|, \tag{41}$$

as soon as this polynomial has two real roots. This phenomenon is illustrated on Fig. 7.

Figure 7: Coupling two identical Schnakenberg systems with $a = 0.2305$, $b = 0.7695$, $s = 2$, $d_{v_B} = 1$ and $\boldsymbol{A} = \boldsymbol{B}$ diagonal with coefficients $\alpha = 1$ and $\beta = 40$. The behaviour of the boundary of the instability region (in grey) near $\eta = 0$ and for large η is well-predicted respectively by Eq. (28) (solid thick line) and Eq. (35) (dotted horizontal line). For η between the roots of the polynomial Eq. (41) (depicted by the randomly dotted band), the 0-th mode is unstable, which indicates a non-Turing type instability.

7 Conclusion and discussion

During development, many organs are characterised by a bilayer geometry, such as the skin and internal epithelium. However, the precise effect of this bilayer coupling on self-organisation dynamics is under-explored. In this study, we have investigated patterning conditions for bilayer reaction-cross-diffusion systems with weak and strong coupling.

Our analysis provides a quantitative description of this coupling effect and shows in particular that not only can spatial patterns emerge from the coupling of two non-patterning layers, but also that the coupling of two independent patterning layers can stabilize a homogeneous equilibrium and thus diminish patterning. Furthermore, the reaction-cross-diffusion systems investigated in this study show that the classical paradigm of local activation and long-range inhibition can be weakened by considering alternative mechanisms specific to the bilayer geometry. In particular, we found that different transport rates between the two layers, or the asymmetry of the equilibrium states of the model components between the two layers, can have a critical influence on pattern formation. In the asymptotic regimes of weak and strong coupling, we proved that these constraints are necessary to induce patterning between two non-patterning layers and we highlighted several explicit examples where they are sufficient. Furthermore, we also classified the cases where the coupling has a negative effect on patterning when one or both layers is independently in a patterning state. We observed an additional complication that the layered coupling can disrupt the stability of the homogeneous steady state that is required for Turing's mechanism, further highlighting the need to consider the coupled system rather than layers in isolation. Further behaviours observed in these bilayer systems include a wave of patterning induction, contrasting the behaviour of the classical Turing system, which patterns via simultaneous self-organisation across the domain. In particular, the combined wave and

patterning dynamics bears some resemblance to cell density waves seen in feather patterning. We note that feather placode formation is also typically accompanied by an additional signalling-molecule wave [28], so this phenomenological observation needs more careful study.

More generally, our analysis covers the cases of 1D-1D and 1D-2D systems and these findings offer the prospect of improving our understanding of patterning mechanisms under the control of signalling molecules, and responses to them, in organs incorporating a bilayer structure in their development, such as the skin. Other organs that exhibit an analogous bilayering alongside spatial patterning in development include the villi of the gut, and the branches of the lung and various models for these systems are reviewed in [1, 42].

In addition to investigating this application to biological systems in detail, several other extensions of this work could be considered. First of all, the models studied in this paper are one-dimensional along the x-axis, as Turing patterning conditions are independent of the dimension, modulo geometric effects on wavemode selection [32]. However, in order to quantitatively study the patterning dynamics or to develop the analysis of pattern types, the models should be extended to a 2D surface and a 2D or 3D bulk. This configuration would also be more realistic from the perspective of prospective applications, such as feather array patterning [28]. From a mathematical perspective, and following the seminal developments of Shaw et. al [58] and Ermentrout [17], one should not only study the emergence of patterning but also the self-organized pattern shapes, requiring weakly-nonlinear bifurcation analysis. Weakly nonlinear analysis in the 1D-1D setting has been done in some special cases, e.g. [9, 8], so extending the present theory to this case would be challenging but plausible. In the 1D-2D setting, however, substantial new mathematics would need to be developed to perform such analysis given the complexity of the eigenfunctions in this setting. Another mathematical direction meriting further investigation is a consideration of Hopf/Wave bifurcations in these coupled systems, whereby linear instability leads to spatiotemporally oscillating solutions. Such instabilities are not possible in the classical two-species case, but exist in three-species models or models incorporating inertial (hyperbolic) effects [32, 55]. Regarding potential biological applications, see, for example, [10].

Furthermore, when the bulk has a non-zero depth along the y-axis, another generalisation could consider when the two layers do not admit the same equilibrium state. This would lead to a non-homogeneous equilibrium state, constant along the x-axis of the interface, but possessing a gradient along the y-axis. Studying the linear stability of such a system would require perturbing about a heterogeneous steady state and, for instance, may further localise patterning to near the interface, in contrast to the situation obtained in Fig. 5. We remark that even in simple 1D scenarios, spatial heterogeneity often requires the study of specific asymptotic regimes to make progress [22]. Another direction is to consider the influence of the interface shape and mechanical structure on patterning. For instance, a fundamental question is whether a local deformation of the interface could have an influence on patterning, noting that the self-organisation is associated with a local mechanical compression and deformation of the epithelium [25, 28], as well as motivating earlier and more recent theoretical work [11, 13].

In summary, we have derived conditions for self organisation for a coupled bilayer system, with

reaction-cross diffusion dynamics in each layer, under a number of limiting cases, such as thin layers and weak or strong coupling. The exploration of these conditions has revealed how the bilayering influences pattern formation, and can act to enhance or inhibit the prospect of patterning. In particular, our study emphasises that a detailed comparison of theory with observation for developmental periodic-structure formation, or chemical patterning in layered reaction diffusion systems [14], has to accommodate the prospect that considering the layers in isolation is insufficient to determine the presence, or absence, of self-organisation.

Acknowledgments

The authors are grateful to Denis Headon for his advice and for the useful discussions regarding the biology of the skin. AD and SSL acknowledge the hospitality of the Wolfson Centre for Mathematical Biology of the University of Oxford (UK) where part of this research was carried out. This work is partially funded by the following grants: KAKENHI Transformative Research A (Grant number: JP22H05110) and KAKENHI Fostering Joint International Research (Grant number: JP17KK0094).

Competing Interests

The authors declare no competing interests.

A   Numerical methods

The code used to produce all the simulations and plots presented in this article is freely accessible at

$$\texttt{https://github.com/antoinediez/BilayerReactionCrossDiffusion}$$

It is written in the `Julia` programming language and is based on the following open source libraries.

- All the plots and videos are produced using the `Makie.jl` data visualization ecosystem developed by [12].

- The bifurcation curves and instability regions in the (η, δ) plane for the 1D-1D case are computed using the Routh-Hurwitz criterion and the explicit equations Eqs. (15)-(16). Most of the computations are done symbolically using the `Symbolics.jl` package developed by [27] and the numerical solution uses the nonlinear solver `NLsolve.jl` developed by [43].

- The simulations of the 1D-1D and 1D-2D systems are implemented using a classical method of lines with an upwind scheme for chemotaxis [29] and an implicit adaptive time-stepping using the ODE solver `DifferentialEquations.jl` developed by [52].

B Supplementary videos

This article is supplemented with the following illustrative videos which show example simulations of 1D-1D and 1D-2D reaction-diffusion-chemotaxis systems. These videos can be accessed online along with the numerical code to reproduce them at

https://github.com/antoinediez/BilayerReactionCrossDiffusion

Video 1. Example of patterning with small coupling for the system described in Fig. 2b with $\eta = 0.01$ and $d_{v_S} = \delta_c(0) + \frac{1}{2}\delta_c'(0) < \delta_c(0)$.

Video 2. Example of surface-driven patterning for the 1D-2D system described in Fig. 4a.

Video 3. Example of bulk-driven patterning for the 1D-2D system described in Fig. 4b.

Video 4. Wave patterning in the 1D-2D system described in Fig. 5.

Video 5. Example of stabilization of a homogeneous equilibrium using a large coupling in the system described in Fig. 6b with $\eta = 200$ and $d_{v_S} = 232 > \delta_c(0)$.

Video 6. Example of patterning with large coupling for the system described in Fig. 3b with $\eta = 100$ and $d_{v_S} = 15$.

References

[1] A. A. Almet, P. K. Maini, D. E. Moulton, and H. M. Byrne. "Modeling Perspectives on the Intestinal Crypt, a Canonical System for Growth, Mechanics, and Remodeling". *Curr. Opin. Biomed.* 15 (2020), pp. 32–39. DOI: 10.1016/j.cobme.2019.12.012.

[2] C. Armit, L. Richardson, S. Venkataraman, L. Graham, N. Burton, B. Hill, Y. Yang, and R. A. Baldock. "eMouseAtlas: An Atlas-Based Resource for Understanding Mammalian Embryogenesis". *Developmental Biology* 423.1 (2017), pp. 1–11. DOI: 10.1016/j.ydbio.2017.01.023.

[3] R. Bailleul, C. Curantz, C. Desmarquet-Trin Dinh, M. Hidalgo, J. Touboul, and M. Manceau. "Symmetry Breaking in the Embryonic Skin Triggers Directional and Sequential Plumage Patterning". *PLoS Biol.* 17.10 (2019). Ed. by C. S. Hill, e3000448. DOI: 10.1371/journal.pbio.3000448.

[4] J. Bard and I. Lauder. "How Well Does Turing's Theory of Morphogenesis Work?" *J. Theoret. Biol.* 45.2 (1974), pp. 501–531. DOI: 10.1016/0022-5193(74)90128-3.

[5] I. Berenstein, M. Dolnik, L. Yang, A. M. Zhabotinsky, and I. R. Epstein. "Turing Pattern Formation in a Two-Layer System: Superposition and Superlattice Patterns". *Phys. Rev. E* 70.4 (2004), p. 046219. DOI: 10.1103/PhysRevE.70.046219.

[6]   M. Bestehorn. "Two-Layer Model Showing a Variety of Pattern Types near Nonequilibrium Phase Transitions". *Physical Review E* 53.5 (1996), pp. 4842–4846. DOI: `10.1103/PhysRevE.53.4842`.

[7]   R. S. Cantrell and C. Cosner. *Spatial Ecology via Reaction-Diffusion Equations.* Wiley Series in Mathematical and Computational Biology. Chichester, West Sussex, England ; Hoboken, NJ: J. Wiley, 2003.

[8]   J. K. Castelino, D. J. Ratliff, A. M. Rucklidge, P. Subramanian, and C. M. Topaz. "Spatiotemporal Chaos and Quasipatterns in Coupled Reaction–Diffusion Systems". *Phys. D* 409 (2020), p. 132475. DOI: `10.1016/j.physd.2020.132475`.

[9]   A. J. Catllá, A. McNamara, and C. M. Topaz. "Instabilities and Patterns in Coupled Reaction-Diffusion Layers". *Phys. Rev. E* 85.2 (2012), p. 026215. DOI: `10.1103/PhysRevE.85.026215`.

[10]   J. C. Cavallo, S. Scholpp, and M. B. Flegg. "Delay-Driven Oscillations via Axin2 Feedback in the Wnt/ β -Catenin Signalling Pathway". *J. Theoret. Biol.* 507 (2020), p. 110458. DOI: `10.1016/j.jtbi.2020.110458`.

[11]   G. C. Cruywagen, P. K. Maini, and J. D. Murray. "Sequential Pattern Formation in a Model for Skin Morphogenesis". *Math. Med. Biol.* 9.4 (1992), pp. 227–248. DOI: `10.1093/imammb/9.4.227`.

[12]   S. Danisch and J. Krumbiegel. "Makie.Jl: Flexible High-Performance Data Visualization for Julia". *Journal of Open Source Software* 6.65 (2021), p. 3349. DOI: `10.21105/joss.03349`.

[13]   L. M. De Oliveira Vilaca, M. C. Milinkovitch, and R. Ruiz-Baier. "Numerical Approximation of a 3D Mechanochemical Interface Model for Skin Patterning". *J. Comput. Phys.* 384 (2019), pp. 383–404. DOI: `10.1016/j.jcp.2019.01.023`.

[14]   B. Dúzs, P. De Kepper, and I. Szalai. "Turing Patterns and Waves in Closed Two-Layer Gel Reactors". *ACS Omega* 4.2 (2019), pp. 3213–3219. DOI: `10.1021/acsomega.8b02997`.

[15]   S.-I. Ei. "The Motion of Weakly Interacting Pulses in Reaction-Diffusion Systems". *J. Dynam. Differential Equations* 14.1 (2002), pp. 85–137. DOI: `10.1023/A:1012980128575`.

[16]   S.-I. Ei and K. Ohgane. "A New Treatment for Periodic Solutions and Coupled Oscillators". *Kyushu J. Math.* 65.2 (2011), pp. 197–217. DOI: `10.2206/kyushujm.65.197`.

[17]   B. Ermentrout. "Stripes or Spots? Nonlinear Effects in Bifurcation of Reaction—Diffusion Equations on the Square". *Proc. R. Soc. Lond. A* 434.1891 (1991), pp. 413–417. DOI: `10.1098/rspa.1991.0100`.

[18]   D. Fanelli, C. Cianci, and F. Di Patti. "Turing Instabilities in Reaction-Diffusion Systems with Cross Diffusion". *The European Physical Journal B* 86.4 (2013), p. 142. DOI: `10.1140/epjb/e2013-30649-7`.

[19]   C. Fraga Delfino Kunz, A. Gerisch, J. Glover, D. Headon, K. J. Painter, and F. Matth. "Novel Aspects in Pattern Formation Arise From Coupling Turing Reaction-Diffusion and Chemotaxis". *Preprint* (2023).

[20] H. Fujita and M. Kawaguchi. "Pattern Formation by Two-Layer Turing System with Complementarysynthesis". *Journal of Theoretical Biology* 322 (2013), pp. 33–45. DOI: 10.1016/j.jtbi.2013.01.008.

[21] E. F. Fussell, A. L. Krause, and R. A. Van Gorder. "Hybrid Approach to Modeling Spatial Dynamics of Systems with Generalist Predators". *J. Theoret. Biol.* 462 (2019), pp. 26–47. DOI: 10.1016/j.jtbi.2018.10.054.

[22] E. A. Gaffney, A. L. Krause, P. K. Maini, C. Wang, and Mathematical Sciences Department, Durham University, Upper Mountjoy Campus, Stockton Rd, Durham DH1 3LE, United Kingdom. "Spatial Heterogeneity Localizes Turing Patterns in Reaction-Cross-Diffusion Systems". *Discrete Contin. Dyn. Syst. Ser. B* In press (2023). DOI: 10.3934/dcdsb.2023053.

[23] A. Gierer and H. Meinhardt. "A Theory of Biological Pattern Formation". *Kybernetik* 12.1 (1972), pp. 30–39. DOI: 10.1007/BF00289234.

[24] J. D. Glover, Z. R. Sudderick, B. B.-J. Shih, C. Batho-Samblas, L. Charlton, A. L. Krause, C. Anderson, J. Riddell, A. Balic, J. Li, V. Klika, T. E. Woolley, E. A. Gaffney, A. Corsinotti, R. A. Anderson, L. J. Johnston, S. J. Brown, S. Wang, Y. Chen, M. L. Crichton, and D. J. Headon. "The Developmental Basis of Fingerprint Pattern Formation and Variation". *Cell* (2023), S0092867423000454. DOI: 10.1016/j.cell.2023.01.015.

[25] J. D. Glover, K. L. Wells, F. Matthäus, K. J. Painter, W. Ho, J. Riddell, J. A. Johansson, M. J. Ford, C. A. B. Jahoda, V. Klika, R. L. Mort, and D. J. Headon. "Hierarchical Patterning Modes Orchestrate Hair Follicle Morphogenesis". *PLoS Biol.* 15.7 (2017). Ed. by C. Hill, e2002117. DOI: 10.1371/journal.pbio.2002117.

[26] D. Gomez, S. Iyaniwura, F. Paquin-Lefebvre, and M. J. Ward. "Pattern Forming Systems Coupling Linear Bulk Diffusion to Dynamically Active Membranes or Cells". *Phil. Trans. R. Soc. A.* 379.2213 (2021), p. 20200276. DOI: 10.1098/rsta.2020.0276.

[27] S. Gowda, Y. Ma, A. Cheli, M. Gwóźdź, V. B. Shah, A. Edelman, and C. Rackauckas. "High-Performance Symbolic-Numerics via Multiple Dispatch". *ACM Communications in Computer Algebra* 55.3 (2021), pp. 92–96. DOI: 10.1145/3511528.3511535.

[28] W. K. W. Ho, L. Freem, D. Zhao, K. J. Painter, T. E. Woolley, E. A. Gaffney, M. J. McGrew, A. Tzika, M. C. Milinkovitch, P. Schneider, A. Drusko, F. Matthäus, J. D. Glover, K. L. Wells, J. A. Johansson, M. G. Davey, H. M. Sang, M. Clinton, and D. J. Headon. "Feather Arrays Are Patterned by Interacting Signalling and Cell Density Waves". *PLoS Biol* 17.2 (2019). Ed. by G. S. Barsh, e3000132. DOI: 10.1371/journal.pbio.3000132.

[29] W. Hundsdorfer and J. Verwer. *Numerical Solution of Time-Dependent Advection-Diffusion-Reaction Equations.* Vol. 33. Springer Series in Computational Mathematics. Berlin, Heidelberg: Springer Berlin Heidelberg, 2003. DOI: 10.1007/978-3-662-09017-6.

[30] S. Kondo and R. Asai. "A Reaction–Diffusion Wave on the Skin of the Marine Angelfish Pomacanthus". *Nature* 376.6543 (1995), pp. 765–768. DOI: 10.1038/376765a0.

[31] S. Kondo and T. Miura. "Reaction-Diffusion Model as a Framework for Understanding Biological Pattern Formation". *Science* 329.5999 (2010), pp. 1616–1620. DOI: 10.1126/science.1179047.

[32] A. L. Krause, E. A. Gaffney, P. K. Maini, and V. Klika. "Modern Perspectives on Near-Equilibrium Analysis of Turing Systems". *Phil. Trans. R. Soc. A.* 379.2213 (2021), p. 20200268. DOI: 10.1098/rsta.2020.0268.

[33] A. L. Krause, V. Klika, J. Halatek, P. K. Grant, T. E. Woolley, N. Dalchau, and E. A. Gaffney. "Turing Patterning in Stratified Domains". *Bull. Math. Biol.* 82.10 (2020), p. 136. DOI: 10.1007/s11538-020-00809-9.

[34] A. N. Landge, B. M. Jordan, X. Diego, and P. Müller. "Pattern Formation Mechanisms of Self-Organizing Reaction-Diffusion Systems". *Developmental Biology* 460.1 (2020), pp. 2–11. DOI: 10.1016/j.ydbio.2019.10.031.

[35] H. Levine and W.-J. Rappel. "Membrane-Bound Turing Patterns". *Phys. Rev. E* 72.6 (2005), p. 061912. DOI: 10.1103/PhysRevE.72.061912.

[36] A. Madzvamuse, A. H. W. Chung, and C. Venkataraman. "Stability Analysis and Simulations of Coupled Bulk-Surface Reaction–Diffusion Systems". *Proc. R. Soc. A.* 471.2175 (2015), p. 20140546. DOI: 10.1098/rspa.2014.0546.

[37] A. Madzvamuse, H. S. Ndakwo, and R. Barreira. "Cross-Diffusion-Driven Instability for Reaction-Diffusion Systems: Analysis and Simulations". *Journal of Mathematical Biology* 70.4 (2015), pp. 709–743. DOI: 10.1007/s00285-014-0779-6.

[38] P. K. Maini, M. R. Myerscough, K. H. Winter, and J. D. Murray. "Bifurcating Spatially Heterogeneous Solutions in a Chemotaxis Model for Biological Pattern Generation". *Bull. Math. Biol.* 53.5 (1991), pp. 701–719. DOI: 10.1007/BF02461550.

[39] P. Maini and M. Myerscough. "Boundary-Driven Instability". *Applied Mathematics Letters* 10.1 (1997), pp. 1–4. DOI: 10.1016/S0893-9659(96)00101-2.

[40] H. Meinhardt and A. Gierer. "Pattern Formation by Local Self-Activation and Lateral Inhibition". *Bioessays* 22.8 (2000), pp. 753–760. DOI: 10.1002/1521-1878(200008)22:8<753::AID-BIES9>3.0.CO;2-Z.

[41] C. D. Meyer. *Matrix Analysis and Applied Linear Algebra*. Society for Industrial and Applied Mathematics, 2001.

[42] T. Miura. "Models of Lung Branching Morphogenesis". *J. Biochem.* 157.3 (2015), pp. 121–127. DOI: 10.1093/jb/mvu087.

[43] P. K. Mogensen, K. Carlsson, S. Villemot, S. Lyon, M. Gomez, C. Rackauckas, T. Holy, D. Widmann, T. Kelman, D. Karrasch, A. Levitt, A. N. Riseth, C. Lucibello, C. Kwon, D. Barton, J. TagBot, M. Baran, M. Lubin, S. Choudhury, S. Byrne, S. Christ, T. Arakaki, T. A. Bojesen, Benneti, and M. R. G. Macedo. *JuliaNLSolvers/NLsolve.Jl: V4.5.1*. Zenodo. 2020. DOI: 10.5281/ZENODO.4404703.

[44] J. D. Murray. *Mathematical Biology: I. An Introduction*. Third. Vol. 17. Interdisciplinary Applied Mathematics. New York, NY: Springer New York, 2002. DOI: `10.1007/b98868`.

[45] J. D. Murray. *Mathematical Biology: II: Spatial Models and Biomedical Applications*. Third. Vol. 18. Interdisciplinary Applied Mathematics. New York, NY: Springer New York, 2003. DOI: `10.1007/b98869`.

[46] J. Murray, P. Maini, and R. Tranquillo. "Mechanochemical Models for Generating Biological Pattern and Form in Development". *Phys. Rep.* 171.2 (1988), pp. 59–84. DOI: `10.1016/0370-1573(88)90003-8`.

[47] M. Myerscough. "Pattern Formation in a Generalized Chemotactic Model". *Bulletin of Mathematical Biology* 60.1 (1998), pp. 1–26. DOI: `10.1006/bulm.1997.0010`.

[48] B. N. Nagorcka and J. R. Mooney. "From Stripes to Spots: Prepatterns Which Can Be Produced in the Skin by a Reaction-Diffusion System". *Math. Med. Biol.* 9.4 (1992), pp. 249–267. DOI: `10.1093/imammb/9.4.249`.

[49] K. J. Painter, W. Ho, and D. J. Headon. "A Chemotaxis Model of Feather Primordia Pattern Formation during Avian Development". *J. Theoret. Biol.* 437 (2018), pp. 225–238. DOI: `10.1016/j.jtbi.2017.10.026`.

[50] M. Pelz and M. J. Ward. "Symmetry-Breaking Bifurcations for Compartmental Reaction Kinetics Coupled by Two Bulk Diffusing Species with Comparable Diffusivities in 2-D". *arXiv:2301.06040* (2023).

[51] M. Pelz and M. J. Ward. "The Emergence of Spatial Patterns for Compartmental Reaction Kinetics Coupled by Two Bulk Diffusing Species with Comparable Diffusivities". *Phil. Trans. R. Soc. A.* 381.2245 (2023), p. 20220089. DOI: `10.1098/rsta.2022.0089`.

[52] C. Rackauckas and Q. Nie. "DifferentialEquations.Jl – A Performant and Feature-Rich Ecosystem for Solving Differential Equations in Julia". *Journal of Open Research Software* 5.1 (2017), p. 15. DOI: `10.5334/jors.151`.

[53] A. Rätz. "Turing-Type Instabilities in Bulk–Surface Reaction–Diffusion Systems". *Journal of Computational and Applied Mathematics* 289 (2015), pp. 142–152. DOI: `10.1016/j.cam.2015.02.050`.

[54] A. Rätz and M. Röger. "Symmetry Breaking in a Bulk–Surface Reaction–Diffusion Model for Signalling Networks". *Nonlinearity* 27.8 (2014), pp. 1805–1827. DOI: `10.1088/0951-7715/27/8/1805`.

[55] J. S. Ritchie, A. L. Krause, and R. A. Van Gorder. "Turing and Wave Instabilities in Hyperbolic Reaction–Diffusion Systems: The Role of Second-Order Time Derivatives and Cross-Diffusion Terms on Pattern Formation". *Ann. Physics* 444 (2022), p. 169033. DOI: `10.1016/j.aop.2022.169033`.

[56] L. Roques and O. Bonnefon. "Modelling Population Dynamics in Realistic Landscapes with Linear Elements: A Mechanistic-Statistical Reaction-Diffusion Approach". *PLoS One* 11.3 (2016). Ed. by Z. Jin, e0151217. DOI: 10.1371/journal.pone.0151217.

[57] J. Schnakenberg. "Simple Chemical Reaction Systems with Limit Cycle Behaviour". *J. Theoret. Biol.* 81.3 (1979), pp. 389–400. DOI: 10.1016/0022-5193(79)90042-0.

[58] L. J. Shaw and J. D. Murray. "Analysis of a Model for Complex Skin Patterns". *SIAM Journal on Applied Mathematics* 50.2 (1990), pp. 628–648. DOI: 10.1137/0150037.

[59] K. Sugimura and H. Kori. "A Reduced Cell-Based Phase Model for Tissue Polarity Alignment through Global Anisotropic Cues". *Scientific Reports* 7.1 (2017), p. 17466. DOI: 10.1038/s41598-017-17611-8.

[60] A. Turing. "The Chemical Basis of Morphogenesis". *Phil. Trans. R. Soc. Lond. B* 237.641 (1952), pp. 37–72. DOI: 10.1098/rstb.1952.0012.

[61] F. Veerman, M. Mercker, and A. Marciniak-Czochra. "Beyond Turing: Far-from-Equilibrium Patterns and Mechano-Chemical Feedback". *Phil. Trans. R. Soc. A.* 379.2213 (2021), p. 20200278. DOI: 10.1098/rsta.2020.0278.

[62] L. Yang, M. Dolnik, A. M. Zhabotinsky, and I. R. Epstein. "Spatial Resonances and Superposition Patterns in a Reaction-Diffusion Model with Interacting Turing Modes". *Phys. Rev. Lett.* 88.20 (2002), p. 208303. DOI: 10.1103/PhysRevLett.88.208303.

[63] L. Yang and I. R. Epstein. "Oscillatory Turing Patterns in Reaction-Diffusion Systems with Two Coupled Layers". *Phys. Rev. Lett.* 90.17 (2003), p. 178303. DOI: 10.1103/PhysRevLett.90.178303.

[64] L. Yang and I. R. Epstein. "Symmetric, Asymmetric, and Antiphase Turing Patterns in a Model System with Two Identical Coupled Layers". *Phys. Rev. E* 69.2 (2004), p. 026211. DOI: 10.1103/PhysRevE.69.026211.