Title: Invasion of spontaneous germinal centers by naive B cells is rapid and persistent

Authors: T. van den Broek,[1] K. Oleinika,[1] S. Rahmayanti,[1] C. Castrillon,[1] C.E. van der Poel,[1] M.C. Carroll,[1]*

Affiliations:

[1] Program in Cellular and Molecular Medicine, Boston Children's Hospital, Harvard Medical School, Boston; MA 02115, USA.

*Corresponding author. Email: Michael.carroll@childrens.harvard.edu

Abstract: In autoreactive germinal centers (GC) initiated by a single rogue B cell clone, wild-type B cells expand and give rise to clones that target other autoantigens, known as epitope spreading. The chronic, progressive nature of epitope spreading calls for early interventions, but the kinetics and molecular requirements for wild-type B cell invasion and participation in GC remain largely unknown. With parabiosis and adoptive transfer approaches in a murine model of systemic lupus erythematosus, we demonstrate that wild-type B cells join existing GCs rapidly, clonally expand, persist, and contribute to autoantibody production and diversification. The invasion of autoreactive GCs required TLR7, B cell receptor specificity, antigen presentation, and type I interferon signaling. The adoptive transfer model provides a novel tool for identifying early events in the breaking of B cell tolerance in autoimmunity.

One Sentence Summary: The autoreactive germinal center is an open structure that is susceptible to rapid and persistent naive B cell invasion, with clonal expansion and auto-antibody induction and diversification.

Main Text:

INTRODUCTION

Systemic lupus erythematosus (SLE) is a prototypical autoimmune disease characterized by a diverse and evolving repertoire of autoantibodies, which are often present prior to the clinical manifestation of disease and drive multi-organ involvement (*1, 2*). In the healthy individual, the mature B cell repertoire contains autoreactive B cell clones, but several tolerance checkpoints prevent them from being activated and producing auto-antibodies (*3, 4*). However, after the initial breakdown of self-tolerance to a single nuclear self-antigen, such as small nuclear ribonucleoproteins (RNPs), there is a diversification of antibody self-specificity, called epitope spreading, that leads to overt disease (*2, 5*).

Germinal centers (GC) are the primary sites for generating high-affinity antibodies and subsequent epitope spreading (*6*). As such, early interventions in the GC response could lead to better therapies. We have previously shown that GCs consist mainly of WT B cells that undergo clonal selection and maturation, resulting in epitope spreading (*7*). What is more, WT B cells eventually become independent of the autoreactive B cells, being able to sustain the GC response in their absence. For that study, we used a mixed bone marrow chimera model, where wild-type (WT) B cells co-exist with transgenic B cells specific for ribonuclear complexes (derived from 564Igi mouse bone marrow) (*7, 8*). However, this model lacks a way to synchronize B cell entry into the GC, so the chronicity of WT B cell engagement precludes the specific evaluation of

early mature B cell activation, GC entry, and autoantibody production. Furthermore, apart from TLR7 (*7, 8*), the molecular requirements for WT B cell participation in autoreactive GCs are unknown.

Here, by synchronizing WT B cell entry, via parabiosis and adoptive transfer experiments, we pinpoint when and how naïve WT B cells gain initial access to the autoreactive germinal center. These data indicate that polyclonal naïve B cells can rapidly enter, clonally expand, and persist within spontaneous GC. Notably, via adoptive transfer of WT B cells, we identified crucial factors involved in their invasion of spontaneous GC and hereby provide a tool to further study the requirements for GC entry.

RESULTS

WT B cells enter and dominate mature autoreactive GCs

We used a parabiosis model to characterize the overall dynamics of WT B cells with synchronized access to spontaneous mature GCs. Specifically, WT mice were surgically bound to 564Igi or WT mice (control parabiosis) (Fig.1A), and the B and T cell exchange between mice was assessed by periodic blood draws and tracked using allelic variants of the pan-leucocyte marker CD45 (Fig.1B, C). In the 564Igi:WT parabiont, B and T cell exchange in the peripheral blood reached 50% at 2 weeks following parabiosis (Fig.1D, Suppl.Fig.1A). Peripheral blood T cell frequencies equilibrated thereafter at comparable exchange rates between the 564Igi:WT and control WT:WT parabiosis (Suppl.Fig.1A,B), indicating a successful surgery. However, WT B cells began to dominate the circulation of the 564Igi:WT parabiont by two weeks and increased over time (Fig.1D, gray line), thereby reducing the proportion of circulating 564Igi B cells (Fig.1E). We looked beyond the circulation and found that, at 8-9 weeks post-surgery, WT B

cells were significantly enriched within the splenic GC compartment (phenotypically GL7+CD38-) of the 564Igi partner; in contrast, this pattern was not observed for 564Igi B cells in the WT parabiont (Fig.1F).

To address whether the increased competitiveness of WT B cells in 564Igi parabionts was due to the relative abundance of B cells in WT mice, we performed WT parabiosis with 564Igi heterozygous mice (50% WT, 50% 564Igi B cells), which have at least 2-fold more B cells than 564Igi homozygous mice. However, the circulatory B cell exchange kinetics in parabionts of WT mice with either 564Igi heterozygous or 564Igi homozygous mice were comparable (Suppl.Fig.1C-F), indicating that it was not the increased number of B cells that made the WT cells more competitive. WT B cell dominance of the splenic GCs in the 564Igi parabiont was also dependent on Toll-like receptor 7 (TLR7) expression (Fig.1G), consistent with our earlier observations of TLR7-dependence in WT B cell participation in the GC response in the 564Igi:WT bone marrow chimera system (*7*). Consistently, in the mesenteric lymph node (mesLN), which is primarily targeted at microbial antigens, WT B cell GC residence was TLR7 independent (Fig.1H).

We next assessed the reactivity of WT-derived serum antibodies to nucleolar autoantigens. Due to allotype differences in the heavy chain locus of 564Igi and WT B cells, the source of IgG2 antibodies can be distinguished by two isoforms, i.e. IgG2a (564Igi) and IgG2c (WT) respectively. We found increased reactivity of IgG2c antibodies against nucleolar auto-antigens in the sera of 564Igi:WT parabionts in contrast to control parabiosis (WT:WT)(Suppl.Fig.1G). Together these results mirror the findings from the earlier mixed bone marrow chimera model, where WT B cells outcompete the spontaneous GC-initiating 564Igi

clone and generate autoantibodies but with the advantage to observe the WT B cell ability to dominate mature GCs in the 564Igi mice.

Enrichment of WT B cells in GCs is persistent

To examine the dynamics of WT B cell invasion and their persistence in autoreactive GCs without a continuous WT B cell source, parabiotic mice were separated at 2 weeks following surgery (Fig.2A). This time point was chosen as this is when parabionts reached adequate cross-circulation, as indicated by T cell equilibration. Following separation, the abundance of WT B and T cells gradually decreased in the circulation of 564Igi recipients, displaying a similar trend to control separation parabionts (Fig.2B,C, Suppl.Fig.2A,B). Of note, WT B cells in the 564Igi parabiont only constituted approximately 20% of peripheral blood B cells 6-7 weeks following separation. By contrast, in the spleen, WT B cells were dominant. Specifically, WT B cells constituted more than 80% of all splenic GC B cells 6-7 weeks after separation in the 564Igi mouse (Fig.2D). WT B cells also dominated the GC response in the axillary/brachial and mesenteric LNs (Suppl.Fig.2C). Importantly, splenic GC enrichment was not apparent at the time of parabiotic separation (Suppl.Fig.2D) but was present for up to a year post-separation (Suppl.Fig.2E). Together, these results show that WT B cells can invade and persist in the autoreactive GC without a continuous source.

WT B cells clonally expand in spontaneous GCs

To determine if the WT B cells underwent clonal selection and expansion in 564Igi GCs, Aicda-Cre[ERT2] Confetti reporter mice were attached to 564Igi mice for two weeks, then they were separated and the 564Igi ex-parabiont was treated with tamoxifen. Upon tamoxifen induction,

Aid-expressing (GC) B cells and their progeny express 1 of 10 different color combinations (*9*), allowing us to visualize the clonal dynamics of these B cells in the spontaneous GCs (*9–11*). At several time points post-separation, spleens were harvested, and their sections analyzed for Confetti-expressing B cells within GCs (Fig.3A). The possibilities are that, upon activation and AID expression, (1) many individual B cells of different clonalities are equally capable of expanding, and as such, they undergo minimal clonal selection and maintain the original color diversity; (2) fate-mapped B cells do not persist in the GC and are replaced by newly recruited B cells that do not show any of the Confetti colors (dark clones), thus diminishing fate-mapped B cells over time; or (3) specific clones are selected and expand, giving rise to the dominance of a single or few colors (referred to as pauci-clonality). The latter is observed in foreign antigen responses and also in 564Igi:WT bone marrow chimeras (*7, 9*). Indeed, we also observed that fate-mapped B cells in the GCs of 564Igi parabionts gradually lost color diversity, indicating active participation and clonal selection of these B cells in the spontaneous GCs (Fig.3B-D). Despite the overall proportion of WT B cells increasing over time after separation in our experiment above (Fig.2D, Suppl.Fig.2D), the total number of fate-mapped B cells per GC decreased from day 14 to 30 post-tamoxifen (Fig.3E), suggesting that non-fate-mapped cells (dark clones), likely naïve B cells, are the source of this B cell enrichment. Together, these results show that after a transient WT B cell source, WT B cells clonally expand in the autoreactive GC and contribute, in part, to WT B cell enrichment in the GC.

WT B cells enter the pre-existing autoreactive GCs within days

Having observed the engraftment of WT B cells in the autoimmune environment as early as 2 weeks following parabiosis surgery, an adoptive transfer approach was established to further

characterize and identify the kinetics of WT B cell participation in autoreactive GCs. WT B cells were adoptively transferred to 564Igi recipients, and spleens were harvested at varying periods post-transfer (Fig.4A). As early as day 4 post-adoptive transfer, WT B cells participated in the GC response. By day 7 post-transfer, the proportion of transferred WT B cells within the total GC B cell pool was significantly increased over the proportion of WT B cells within the total B cell compartment of the spleen (Fig.4B,C). Early enrichment of donor B cells was specific to the autoreactive GC, as it was not seen in the mesenteric lymph node and Peyer's patches of the 564Igi recipient or in immunized WT mice (Suppl.Fig.3A-D). Of note, we saw little engraftment of WT B cells transferred to unimmunized WT hosts (Suppl.Fig.3A-D). WT B cells were frequently found in 564Igi GCs, with most GCs in the 564Igi recipient mice populated by WT B cells (Fig.4D). This was consistent among 564Igi recipients despite differences in the number of white pulp areas and GCs (Fig.4E). A range of WT B cell residency patterns within GCs – from dominant to sparse – was observed (Fig.4F), but there were no GCs composed solely of donor B cells at day 7 post-adoptive transfer. Thus, we conclude that, within four days post-transfer, donor B cells joined existing GCs. WT B cell entry of GCs may be dependent on the size of the GC. Conversely, the size of the GC could be impacted by WT B cell residency. However, the GC size was comparable between GCs with both donor and recipient residents and GCs with 564Igi B cells only (Fig.4G). Together, these data show that, like foreign-antigen-driven GCs, autoreactive GCs are open structures to which B cells can be recruited rapidly (*12*).

WT B cells actively participate in autoreactive GCs and can differentiate into autoantibody-producing cells

We next aimed to confirm that adoptively transferred WT B cells actively engaged in the GC response in contrast to transient occupancy (*13*). To address this, we used B cells from S1PR2-Cre[ERT2]TdTomato reporter animals; TAM administration permanently labels *S1pr2*-expressing GC and activated B cells and their progeny with tdTomato (*14*). We transferred S1pr2 reporter B cells to 564Igi Aicda-Cre[ERT2]EYFP recipients. Notably, S1pr2-TdTomato-expressing donor B cells (WT) co-localized in 564Igi GCs with Aicda-Cre[ERT2]EYFP -expressing 564Igi B cells (Suppl.Fig.3E). Interestingly, donor-derived antibodies (IgG2c) specific for nuclear antigens (ANA), neutrophil cytoplasmic antigens (ANCA) and ssDNA were present as early as 7 days post-adoptive transfer (Suppl.Fig.3F-H). After transfer, the relatively quick rise of WT-derived plasma cells before the peak of donor GC B cells suggests that both extra-follicular and GC-derived antibody-producing plasma cells are generated (Suppl.Fig.3I).

To demonstrate that GC-derived B cells possessed the ability to produce autoantibody at this early time-point, S1pr2-TdTomato reporter B cells (WT B cell repertoire) were transferred into tamoxifen-treated 564Igi recipients. TdTomato-expressing (S1pr2$^+$) B cells were then sorted and cultured, then the antibody reactivity of GC-experienced WT B cells was measured in the cell culture supernatant (Suppl.Fig.3J). Self-reactive antibody against ssDNA was detected in sorted TdTomato-expressing (S1pr2$^+$) B cells. We further observed that sorted donor B cells that did not express the S1pr2 reporter also secreted ssDNA antibody, although at a lower level, suggesting a GC-independent source of auto-antibody production or incomplete labeling of B cells in the GC in this narrow timeframe of tamoxifen exposure (Suppl.Fig.3K). Together, these results show that donor-derived WT B cells can enter the autoreactive GC and subsequently differentiate into autoantibody-producing cells.

GC invasion by WT B cells is dependent on BCR specificity and MHC expression and can be regulated by TLR7 and IFNAR expression

The adoptive transfer approach allowed us to further examine the molecular requirements for B cell entry into spontaneous GCs and to identify possible targets for intervention. The role of BCR specificity for entry into GC was assessed by the transfer of B1-8 transgenic (Tg) B cells into 564Igi recipients. In the B1-8 Tg mouse strain, the pairing of the Ig heavy chain knock-in with Ig lambda (Igλ) light chain confers specificity to the hapten 4-hydroxy-3-nitrophenylacetate (NP); thus approximately 5% of the peripheral B cells are NP-specific (Fig.5A) (*15*). In contrast to observations with WT B cells, the proportion of Igλ$^+$ Tg B cells (NP-specific) within the GC population was decreased relative to naïve B1-8-derived B cells in both spleen and mesLN (Fig.5B, Suppl.Fig.4A). Thus, NP-specific B cells did not participate in the spontaneous GC response, indicating the essential role of antigen recognition in this process. NP-specific Igλ+ B cells did however readily enter GCs in WT mice immunized with haptenated NP-CGG in a similar adoptive transfer scheme (Suppl.Fig.4B), as previously published (*16*). In contrast to the results with NP-specific B cells, we found that Sle1yaa-derived B cells not only readily enter the GC when transferred into 564Igi recipients, but their capacity to invade GCs was superior to the B cells derived from WT mice (Fig.5C). Sle1yaa mice have the lupus erythematosus susceptibility locus as well as a Y chromosome translocation that increases the expression of TLR7 approximately 2-fold, which suggests a role for increased TLR7 expression in enhancing GC entry (*17*).

To determine the requirement for antigen presentation in the participation of WT B cells in spontaneous GCs, MHCII KO B cells were adoptively transferred into 564Igi recipients (Fig.5C). Notably, MHC II–deficient B cells failed to participate in spontaneous GCs, even when

MHCII expression was halved (MHCII haploinsufficiency) (Suppl.Fig.4C). MHCII–deficient B cells were able to enter the B cell follicle but not the GC area (Suppl. Fig.4D). B cells similarly failed to engage in the GC response upon adoptive transfer when they lacked the expression of TLR7 or type I interferon receptor (IFNAR) signaling (Fig.5C). Diminished expression of IFNAR (IFNAR haploinsufficiency) did, however, result in partial recovery of B cell participation in the GC response (Suppl.Fig.4C). Of note, both TLR7-KO and IFNAR-deficient B cells competed similar to WT B cells for entry into the GC of the mesLN (Suppl.Fig.4E), suggesting selective TLR7 and IFNAR dependence towards the autoreactive GCs. Together, these data demonstrate that WT B cell entry and participation in spontaneous GCs is an active process dependent on BCR specificity and antigen presentation and can be titrated by the expression level of TLR7 and IFNAR.

DISCUSSION

In a foreign antigen response, established germinal centers are dynamic, open structures that harbor B cells with different specificities and allow naïve B cells to enter and participate in the GC process (*12*, *15*, *16*, *18–22*). We found that, within the 564Igi mice, WT B cells were able to enter and participate in the autoimmune GC response, eventually acquiring a specificity for ssDNA and also displaying a wide array of class-switched autoantibody reactivities, suggesting possible recruitment of B cells with other specificities. Our observations are reminiscent of the breach of B cell tolerance observed in tonsil biopsies of SLE patients, where despite autoreactive B cells being excluded from the GCs of healthy individuals, in SLE patients, autoreactive B cells readily engaged in the GC response (*23*).

Progressive GC enrichment of WT B cells is evident in the parabiosis model, yet there was a decrease in the number of fate-mapped B cells after temporary parabiosis. This suggests that WT naive B cells (dark cells) entered the GC, as is observed in long-lasting GCs primed by respiratory viruses (*20*). In the adoptive transfer model, from 10 days post-transfer onwards, the proportion of WT B cells decreased and persisted at a lower level, supporting that persistent WT B cell GC enrichment may be dependent on a continuous source of naïve B cells. WT B cells did increasingly invade the GCs of the mesenteric lymph node (mesLN) and Peyer's patches (PP) and persisted at 8 weeks post-transfer. This could suggest that WT B cells initially favor entry into the autoreactive GC, but at a later stage, the majority of naïve B cells preferentially enter and persist in the mesLN and PP, as they are likely more reactive than 564Igi B cells to the microbial antigens found in gut-related lymph nodes.

In murine infection and vaccination response models, the entry of naive B cells into GCs depends on specific BCR and MHCII expression, as in our autoimmune models (*12*, *16*, *20*, *22*, *24*, *25*). In contrast, we show that TLR7 and IFNAR expression altered the threshold for naive B cell entry in the autoreactive GC but minimally affected entry in the mesenteric lymph node. This difference is of interest as it could potentially be utilized to specifically address the autoreactive GC process, thus could aid in the identification of early intervention strategies that could halt or slow disease progression. TLR7 signaling is important in the establishment of autoimmune diseases, such as SLE, by inducing extrafollicular antibody-forming cells (*26*, *27*). The dual function of TLR7 might highlight the requirements for the initiation of disease by autoreactive B cells (GC-independent) and, at least in part, the propagation of disease by recruitment of new B cells (GC-dependent). IFNAR signals play an important role in the regulation of TLR7 responses in B cells and could contribute to further GC-dependent

propagation of disease, as B cell–intrinsic type 1 IFN receptor signals are not required for SLE development but do enhance disease (*28–30*). Lack of GC-specific IFNAR signaling indeed reduces the size and frequency of the GC as well as auto-antibody production, and our data suggest that this might be due to reduced B cell entry to the GC (*31*, *32*).

Our findings not only give mechanistic insight into the kinetics and requirements for a break of tolerance of naïve B cells but also underscore the potential therapeutic targets in autoimmunity. The chronic and progressive nature of WT B cells joining the GC response calls for early interventions. Total blockade of GC responses might, however, exacerbate autoimmune disease (*33*, *34*), and warrants selective GC checkpoints such as TLR7 and IFNAR to regulate autoreactive from foreign GC responses, but could conflict with responses to viral infection (*35*, *36*). We suggest that adoptive transfer in 564Igi mice can be utilized more widely as a novel tool to address the early events and molecular mechanisms facilitating the break in tolerance of naive B cells.

MATERIALS AND METHODS

Study design

We performed parabiosis and adoptive transfers to assess the kinetics of WT B cell participation and break of tolerance in the autoreactive GC. The details of the biological replicates in each group and the number of repetitions per experiment are indicated in the respective figure legends. Experimental end-points were established before any *in vivo* experiments for animal welfare considerations. All animals used for comparison within each experiment were age- and

sex-matched. Mice in different groups were randomly assigned, outliers were not excluded, and studies were unblinded.

Mice

This study was conducted in accordance with the NIH Guide for the Care and Use of Laboratory Animals. Animal experiments were performed as specified in protocols approved by the local IACUC of Harvard Medical School (protocol number IS00000111). All surgeries and cell transfers were performed under isoflurane anesthesia, and every effort was made to minimize suffering. 564Igi mice were provided by Theresa Imanishi-Kari (Tufts University) and maintained in-house, and different congenic backgrounds (CD45.1 and CD45.1.2) and a 564Igi AicdaCreERT2-YFP were generated. AicdaCreERT2-Confetti, B1-8, and S1PR2CreERT2-TdTomato donors were provided by Gabriel D. Victora (Rockefeller University). B6 (C57BL/6J), B6 CD45.1 (B6.SJL-Ptprca Pepcb/BoyJ), TLR7 KO (B6.129S1-Tlr7 tm1Flv /J), I-AB-flox (MHCII-fl) (B6.129X1-H2-Ab1tm1Koni/J), Sle1Yaa (B6.Cg-Sle1NZM2410/Aeg Yaa/DcrJ), MB1-Cre (B6.C(Cg)-Cd79atm1(cre)Reth/EhobJ) were from Jackson Laboratories and maintained in house. IFNaR fl/fl (Ifnar1tm1Uka) mice were provided by Ulrich Kalinke. MB1-Cre mice were crossed with I-AB-flox and with IFNaR fl/fl to generate B cell-specific expression. AicdaCreERT2 flox-stop-flox- EYFP mice were from Claude-Agnes Reynaud and Jean-Claude Weill (Institut Necker) and crossed with 564Igi mice. To generate 564IgiHet, MHCII+/-, and IFNaR +/- mice, corresponding homozygous mice were crossed with B6, B6 CD45.1, or MB1-Cre and maintained in-house.

Experimental models: Organisms/strains	Source	Identifier
Aid-CreERT2 EYFP: AicdaCreERT2 flox-stop-flox-	Dogan et al., 2009	Ref (*13*)

EYFP		
564Igi	Berland et al., 2006	Ref (*8*)
Aid-CreERT2 Confetti: AicdaCreERT2-Rosa26Confetti	Tas et al, 2016	Ref (*9*)
C57BL/6j (CD45.2)	The Jackson Laboratory	000664
B1-8 (PA-GFP)	Victora et al, 2010	Ref (*37*)
S1PR2-CreERT2-TdTomato	Shinnakasu et al., 2016	Ref (*14*)
B6.SJL-Ptprca Pepcb/BoyJ (CD45.1)	The Jackson Laboratory	002014
TLR7 KO: B6.129S1-Tlr7 tm1Flv /J	The Jackson Laboratory	008380
I-AB-flox: B6.129X1-H2-Ab1tm1Koni/J	The Jackson Laboratory	013181
Sle1Yaa: B6.Cg-Sle1NZM2410/Aeg Yaa/DcrJ	The Jackson Laboratory	021569
MB1-Cre: B6.C(Cg)-Cd79atm1(cre)Reth/EhobJ	The Jackson Laboratory	020505
Ifnar1 tm1Uka	Le Bon et al. 2006	Ref (*38*)

Mice were bred and maintained in an AAALAC-accredited facility at Harvard Medical School under specific pathogen-free (SPF) conditions. Both female and male mice were used in the adoptive transfer studies. Only females are suited for parabiosis experiments. Mice were on a standard 12hr light/dark cycle and were kept on a high-fat diet for 1-2 weeks before and during the parabiosis experiments and on a regular low-fat diet for adoptive transfer studies. Mice were 7–9 weeks old at the start of experiments. Mice were age-matched when comparing different strains (max 1 week between experimental and control groups).

Antibodies and staining reagents

Immunogen	Fluorophore	Clone	Source	Identifier	RRID

B220	FITC	RA3-6B2	Biolegend	103206	AB_312991
B220	PerCP-Cy5.5	RA3-6B2	Biolegend	103236	AB_893354
B220	APC	RA3-6B2	Biolegend	103212	AB_312997
B220	BV510	RA3-6B2	Biolegend	103247	AB_2561394
CD4	BV510	RM4-5	Biolegend	100553	AB_2561388
CD21/CD35	Pacific Blue	7E9	Biolegend	123414	AB_2085158
CD38	PE-Cy7	90	Biolegend	102718	AB_2275531
CD45.1	FITC	A20	Biolegend	110706	AB_313495
CD45.1	PE	A20	Biolegend	110708	AB_313497
CD45.1	APC	A20	Biolegend	110713	AB_313502
CD45.1	PE-Cy7	A20	Biolegend	110730	AB_1134168
CD45.2	FITC	104	Biolegend	109805	AB_313442
CD45.2	APC	104	Biolegend	109814	AB_389211
CD138	APC	281-1	Biolegend	142506	AB_10962911
CD138	BV510	281-2	Biolegend	142521	AB_2562727
CD169	FITC	3D6.112	Biolegend	142405	AB_2563106
CD169	PE	3D6.112	Biolegend	142403	AB_10915470
GL7	Pacific Blue	GL7	Biolegend	142403	AB_2563292
GL7	PE	GL7	Biolegend	144607	AB_2562925
GL7	PerCP-Cy5.5	GL7	Biolegend	144610	AB_2562979
IgMa	PE	MA-69	Biolegend	408608	AB_940545
IgMb	FITC	AF6-78	Biolegend	406206	AB_315039
IgD	Pacific Blue	11-26c.2a	Biolegend	405712	AB_1937244

Rabbit anti-Goat IgG (H+L)	AF488	polyclonal	ThermoFischer Scientific	A-11078	AB_2534122
Streptavidin	PE	N/A	Biolegend	405204	-
Streptavidin	PE-Cy7	N/A	Biolegend	405206	-
Fixable Viability Dye	eFluor780	N/A	Invitrogen	65-0865-14	-

In-house generated anti-idiotypic Ab, clone 9D11, conjugated to Alexa Fluor 647 or streptavidin.

Genotyping and FACS-typing mice

Genotyping was performed using the primers and reaction conditions indicated in Table S1. For

FACStyping, 564Igi mice were bled retro-orbitally. Using heparinized capillary tubes,

approximately 60–100 µl blood was drawn into Eppendorf tubes containing 30 µl acid-citrate-

dextrose solution. Following collection, the stabilized blood was briefly spun down, then under-

layered with 1 ml of Lymphocyte Separation Medium, and spun for 25 min at 1,600 RPM at RT.

The mononuclear cell layer was aspirated and transferred into 1 ml ice-cold FACS buffer, mixed,

then pelleted at 200 g for 5 min. Cells were resuspended in FACSbuffer and processed for flow

cytometric analysis as described further below. FACStyping of 564Igi mice was performed using

B220, anti-IgMa, anti-IgMb, and 9D11. 564Igi mice were genotyped by ddPCR using primers

(Table S1) to assess the expression of heavy and light chains.

Parabiosis surgery

Mice were co-housed for 1-2 weeks before parabiosis surgery, and their flanks were shaved a day before. Surgeries were performed on heating pads under aseptic conditions with controlled isoflurane anesthesia. Mirror-image incisions at the left and right flanks of the mouse were made through the skin. Elbow and knee joints from each parabiont were sutured together (Ethilon 4-0), followed by a continuous suture to join the skin together (Ethilon 4-0). Mice in each pair were injected subcutaneously with buprenorphine twice a day for 3 days or buprenorphine SR once with daily meloxicam to manage pain, supplemented with 0.9% (w/v) sodium chloride for hydration. Control parabiosis surgeries consisted of joining wild-type mice differing in congenic marker (CD45.1 and CD45.2). In addition, the side of the surgery was alternated between similar parabiosis pairings. Mice were housed in alpha-dry bedding and received sulfamethoxazole/trimethoprim in their drinking water for 10 days. For overall health and maintenance behavior, mice were monitored daily for the first week and analyzed for weight loss, grooming, stress, and pain response, followed by visual monitoring 2-3 times a week.

The parabiosis separation surgery was performed after 2 weeks of pairing and followed the same setup, analgesia, and monitoring as the initial parabiosis surgery. The sutures joining the knee and elbow joints were removed and the skin of each mouse was closed with a continuous suture (Ethilon 4-0). For blood collection mice were bled retro-orbitally and via the submandibular route. Using heparinized capillary tubes, approximately 60–100 µl blood was drawn into Eppendorf tubes containing 30 µl acid-citrate-dextrose solution.

B cell isolation and adoptive transfer

Spleens were passed through a 70-um cell strainer, and splenocytes were resuspended in RBC lysis buffer (155 mM NH_4Cl, 12 mM $NaHCO_3$, 0.1 mM EDTA), incubated for 2–3 minutes then spun down as before and resuspended in MACS buffer. Splenocytes were incubated with 100ul

of biotin-antibody cocktail for 1×10^8 cells in 1ml of MACS buffer (2% fetal calf serum, 1mM EDTA) for 15min on ice, followed by 100ul of Streptavidin nanobeads for 1×10^8 cells in 1ml of MACS buffer for 15min on ice (MojoSort Mouse pan B cell isolation kit II, Cat.no. 480088, Biolegend). The cell suspension was diluted and put on a pre-rinsed LS Column (Cat. No. 130-042-401, Milteny Biotec), and the column was washed three times. The flow-through was collected on ice, counted and diluted until 10×10^6 cells per 100ul of HBSS and kept on ice. Purity was assessed by FACS and showed consistent >96% B cell (B220+) purity. The adoptive transfer was performed following appropriate dilution by retroorbital injection on anesthetized mice. Adoptive transfer following immunization was done after 7 days of NP-CGG immunization and 11 days after the first SRBC immunization. For the adoptive transfer of NP-specific B cells, mice were immunized with 100ug of NP(24)-CGG (Cat. N5055C-5, Biosearch technologies) emulsified in alum (Cat. 77161, Thermo Fisher Scientific). The following day, isolated B cells from B1-8 mice were adoptively transferred (10×10^6 B cells/recipient); mice were sacrificed on day 7 post-transfer.

564Igi recipients were placed on a tamoxifen diet (TD130860, Envigo) for 3 weeks before B cell transfer and maintained on a tamoxifen diet until the end of the experiment. B cells were isolated from S1pr2-TdTomato reporter mice and adoptively transferred to 564Igi animals (10×10^6 B cells/recipient). On day 7 following adoptive transfer, 564Igi recipient animals were sacrificed. Following RBC lysis, splenocytes were stained with antibodies against IgD, CD138, B220, GL7, CD45.1, and viability dye. Viable CD45.1-B220+CD138-GL7-IgD-S1pr2+ and CD45.1-B220+CD138-GL7-IgD-S1pr2- B cells were sorted. Viable CD45.1+ B220+CD138-GL7-IgD- B cells were sorted as positive controls for anti-ssDNA autoantibody production. Sorting was performed on the SH-800Z cell sorter (Sony Biotechnology). 5000 B cells were then

cultured in the presence of 50,000 sorted B220- cells and with 1ug/mL R848 (tlrl-r848, InVivoGen) and anti-CD40 (clone FGK4.5, BE0016-2, BioXCell) for 6 days. Anti-ssDNA autoantibody production was then assessed by ELISA.

Immunization

To generate (transient) foreign-Ag-elicited GCs, mice were immunized intraperitoneally with 100 µg of chicken gamma globulin (Rockland Immunochemicals) precipitated in an equal volume of Imject Alum (ThermoScientific). To generate chronic GCs, mice were immunized with sheep red blood cells (SRBC) twice. For SRBC immunizations, 10–15 mL of Sheep blood in Alsevers (Colorado Serum Company) was washed two times in HBSS before counting SRBC using a Neubauer hemacytometer. Mice received a single injection of 2E8 SRBC in HBSS i.p followed by a boost of 2E8 SRBC 7 days later.

Confetti analysis

Analysis was performed as described in the manuscript of S.Degn et al. (*7*). Briefly, acquired stacks were rendered in Fiji (ImageJ). Counting was performed on representative z-plane from stacks acquired at 5-µm steps through the GC, with multiple GCs per mouse. Raw counts were converted to relative frequencies for each observable color to simplify analysis. Clonal dominance was calculated as the frequency of the most dominant clone/color at a given time point. The clonal divergence score is calculated at a given time point by comparing the observed distribution of colors at time t to the expected frequencies resulting from random recombination, as reported previously (*9*). Expected frequencies for each of 10 possible colors were derived using explant data from day 3 post-tamoxifen treatment, where color distribution most closely

mimics random recombination and before any selection events. Descriptors of GC behavior and clonality were graphed using GraphPad Prism 8.

Flow cytometry

Spleens and LNs were harvested into ice-cold FACS buffer and mechanically dissociated using pestles in 1.5 ml Eppendorf tubes. Samples were filtered through 70-µm cell strainers (Corning) and spun down at ~200 g for 5 min. LN samples were resuspended in FACS buffer. Spleen samples were resuspended in RBC lysis buffer (155 mM NH4Cl, 12 mM NaHCO3, 0.1 mM EDTA), incubated for 2–3 minutes then spun down as before and resuspended in FACS buffer. Samples were added to wells of 96-well round-bottom plates, spun down, and resuspended in 100 µl staining mix (appropriate Ab and viability dye cocktail in FACS buffer). Staining was performed for 30 min on ice in the dark, followed by the addition of 150 µl of wash buffer. Plates were subsequently centrifuged at 200 g for 5 min, and supernatants were flicked out of the plates. For two-step staining procedures, secondary staining mix was added (appropriate secondary Ab mix and viability dye in FACS buffer) and the process was repeated. When appropriate, cells were permeabilized for 30 min on ice and subsequently washed with perm/wash buffer. Appropriate intracellular Ab staining mix in perm/wash buffer was added for 30-45min on ice and subsequently washed. Following the last wash, samples were resuspended in 200 µl FACS buffer and transferred to FACS tubes. Flow cytometric analyses were performed on a FASCSCanto2, with 8-color and 10-parameter analytical capabilities.

Immunofluorescence confocal microscopy

Freshly harvested tissue was put in 4% PFA for several hours on ice and subsequently embedded in OCT (TissueTek) and immediately frozen at −80°C. Tissue blocks were equilibrated at cutting

temperature (−16 to −20°C depending on tissue type), and 10-15 µm thick sections were cut on a cryostat. Tissue sections were mounted on SuperFrost+ slides (Fisher Scientific) and fixed using ice-cold acetone. Slides were then rinsed with PBS and incubated with block/perm buffer (PBS, 2% FBS, 0.1% NaN3, and 0.1% Triton-X100) for 30 min. This was followed by incubation with primary Ab mixture in staining buffer (PBS, 2% FBS, 0.1% NaN3), overnight at 4°C. For two-step staining procedures, the slides were washed 3 times with PBS, 0.01% Tween-20. Then secondary Ab mixture in staining buffer was added and incubated for 2 hrs at RT. At the end of either one- or two-step staining procedures, slides were washed once with staining buffer for 5 min, then 3 times for 5 min with PBS, 0.01% Tween-20. Slides were spot-dried, then mounted in Fluoro-Gel (Electron Microscopy Sciences), and coverslipped. Imaging was performed using a Fluoview FV1000 inverted Olympus IX 81 confocal microscope, equipped with 6 laser lines (405, 457, 488, 515, 559, 635 nms) and 4 fluorescence + 1 transmission detectors (PMTs).

For each mouse, 2 non-consecutive sections of the spleen (separated by at least 50mM during cryosectioning) were stained and imaged on the FV3000R resonant scanning confocal microscope (Olympus). Images were acquired at 30x magnification with at least 4 individual fields of view per section to cover the entire area. To visualize entire spleens for downstream analysis, manual tiling of individual fields of view was carried out for each section. To determine the frequency of GCs with 564Igi and WT, 564Igi only or WT only 'residents', each section was manually examined and assigned to one of the groups. In parallel, GCs and white pulp areas were enumerated. GC area was determined using the 'measure' function in Fiji. For each mouse, the plotted values represent the average of the two non-consecutive splenic sections.

Additional procedures

Recombination of the Rosa26Confetti allele in AicdaCreERT2 mice was induced by a single gavage of 10 or 15 mg of tamoxifen (Sigma) dissolved in sunflower oil at 20 or 30 mg/ml. For the transfer of S1PR2CreERT2 TdTomato B cells into 564Igi AicdaCreERT2 eYFP mice, tamoxifen gavage was performed on day 3 and day 5 post-adoptive transfer; mice were sac'ed on day 7 post-adoptive transfer. For the transfer of S1PR2CreERT2 TdTomato B cells into 564IgiHomo (CD45.1) mice for anti-ssDNA IgG analysis, a tamoxifen-containing diet was started 21 days prior and continued during adoptive transfer.

Multiphoton imaging

Spleens were harvested at different times post-tamoxifen. Thick transversal sections (~1 mm) were manually cut from spleens using surgical scissors. Spleen sections were placed in PBS in vacuum-grease chambers on microscope slides. Chambers were coverslipped and kept on ice before imaging. All imaging was performed on an upright Olympus FV1200 MPE multiphoton system microscope fitted with either a 20× 0.95NA Plan water-immersion objective or a 25× 1.05NA Plan IR optimized water-immersion objective, a MaiTai HP DeepSee Ti-Sapphire laser (Spectraphysics), and 4 non-descanned detectors (2 GaAsP and 2 regular PMTs). Imaging of Confetti alleles was performed using $\lambda=940$ nm excitation. Fluorescence emission was collected in three channels, using the following filter sets: a pair of CFP (480/40 nm) and YFP (525/50 nm) filters, separated by a 505 nm dichroic mirror, for CFP/GFP/YFP detection, and a third filter (605/70 nm) for RFP detection.

Autoantibody array

MILLIPLEX MAP Human Autoimmune Autoantibody Panel (HAIAB-10K, Millipore Sigma) was used following manufacturer's instructions. Due to supply chain issues, ribosomal P antigen was not included in the original kit, and the assay was performed against 19 autoantigens instead of the original 20. Serum was collected from whole blood by allowing clotting at room temperature for 30 minutes, followed by 2 centrifugation steps at 4°C: 1000g for 10 minutes, and 2700g for 2 minutes, collecting the supernatant after each step. BPI (Surmonics, Cat. A192), Ro60 (Surmonics, Cat.A174), and APRIL (Peprotech, Cat. 310-10C) were added to the assay. For anti-IgG2c (Cat 1079-09, Southern Biotech) and IgG-PE (Cat 1030-09, Southern Biotech), the amount of protein used for coating the beads was 0.005 nmoles; for the other proteins BPI, APRIL, Ro60 the concentration was 0.5 nmoles. For all beads, the calculations were made to have at least 400 beads of each protein (region IDs) at the start of each assay. Serum from 3 parabiont pairs of each type was included in the autoantigen array.

Autoantibody measurement

Anti-single-stranded DNA autoantibody measurement by ELISA. ssDNA salmon DNA 20ul/ml (D9156, Sigma) was diluted in 5% DMSO in carbonate/bicarbonate coating buffer and incubated at 95C for 10 minutes. It was then coated on a 96-well ELISA plate and incubated overnight. The plate was washed x3 with wash buffer (0.01% Tween20 in PBS), then blocking buffer (5%BSA, 1%FCS in PBS) was added at 37C for 1h. The plate was washed x3 with wash buffer and sera (diluted 1:20 in 10% blocking buffer in PBS (dilution buffer)), added to the plate, and incubated at 4C overnight. Cell culture supernatants were added undiluted. The plate was washed x3, goat anti-mouse IgG2c-HRP (Southern Biotech, 1079-05) was diluted in dilution buffer and the plate was incubated at 37C for

1h. After washing the plate x3, TMB substrate was added for 15-20 minutes and the reaction was stopped with H2SO4. The 450nm absorbance was measured on the X Mark microplate spectrophotometer (Biorad). C11 (564Igi antibody derived from a hybridoma) was used to standardize absolute levels of anti-ssDNA antibody.

Hep-2 Immunofluorescence assay

ANA (Hep-2) slides (AN-1012, MBL International) and anti-neutrophil cytoplasmic antibody (ANCA) slides (29419, Biorad) were removed from pouches and allowed to equilibrate at room temperature. Sera were diluted (1:20) in PBS, and 30uL was added to each well for 1h at room temperature. Slides were washed twice with blocking buffer (0.1% Tween20, 0.5% BSA in PBS). Secondary antibody goat anti-mouse IgG2c-AF488 (1079-30, Southern Biotech) and (CTB)- AlexaFluor555 (C34776, Thermo Fisher Scientific) were diluted in blocking buffer, added to the slides, and incubated for 1h at room temperature. Slides were washed twice with blocking buffer and mounted with DAPI Fluoromount-G mounting medium (0100-20, Southern Biotech).

Image analysis

Images were quantified using CellProfiler . For HEp-2 slides, quantifications were done based on the DAPI (nucleus) and Phalloidin staining. Cell outlines were detected using the phalloidin staining and as a function of the detected nuclei. A cytoplasmic mask was generated by subtracting the nucleus mask from the cell mask. Each bar represents a mean of individual objects (for instance a mean of MFI from 20 nuclei).

Statistical analysis

Statistical tests are described in the figure legends. Data were analyzed using GraphPad Prism software (GraphPad 8, San Diego, CA). No statistical methods were used to predetermine the sample size before the experiments.

Supplementary Materials

Figs. S1 to S4

References and Notes

1. P. E. Lipsky, Systemic lupus erythematosus: an autoimmune disease of B cell hyperactivity. *Nat. Immunol.* **2**, 764–766 (2001).

2. M. R. Arbuckle, M. T. McClain, M. V. Rubertone, R. H. Scofield, G. J. Dennis, J. A. James, J. B. Harley, Development of Autoantibodies before the Clinical Onset of Systemic Lupus Erythematosus. *N. Engl. J. Med.* **349**, 1526–1533 (2003).

3. H. Wardemann, S. Yurasov, A. Schaefer, J. W. Young, E. Meffre, M. C. Nussenzweig, Predominant Autoantibody Production by Early Human B Cell Precursors. *Science*. **301**, 1374–1377 (2003).

4. K. Koelsch, N.-Y. Zheng, Q. Zhang, A. Duty, C. Helms, M. D. Mathias, M. Jared, K. Smith, J. D. Capra, P. C. Wilson, Mature B cells class switched to IgD are autoreactive in healthy individuals. *J. Clin. Invest.* **117**, 1558–1565 (2007).

5. U. S. Deshmukh, F. Gaskin, J. E. Lewis, C. C. Kannapell, S. M. Fu, Mechanisms of Autoantibody Diversification to SLE-Related Autoantigens. *Ann. N. Y. Acad. Sci.* **987**, 91–98 (2003).

6. G. D. Victora, M. C. Nussenzweig, Germinal Centers. *Annu. Rev. Immunol.* **30**, 429–457 (2012).

7. S. E. Degn, C. E. van der Poel, D. J. Firl, B. Ayoglu, F. A. Al Qureshah, G. Bajic, L. Mesin, C.-A. Reynaud, J.-C. Weill, P. J. Utz, G. D. Victora, M. C. Carroll, Clonal Evolution of Autoreactive Germinal Centers. *Cell*. **170**, 913-926.e19 (2017).

8. R. Berland, L. Fernandez, E. Kari, J.-H. Han, I. Lomakin, S. Akira, H. H. Wortis, J. F. Kearney, A. A. Ucci, T. Imanishi-Kari, Toll-like Receptor 7-Dependent Loss of B Cell Tolerance in Pathogenic Autoantibody Knockin Mice. *Immunity*. **25**, 429–440 (2006).

9. J. M. J. Tas, L. Mesin, G. Pasqual, S. Targ, J. T. Jacobsen, Y. M. Mano, C. S. Chen, J.-C. Weill, C.-A. Reynaud, E. P. Browne, M. Meyer-Hermann, G. D. Victora, Visualizing antibody affinity maturation in germinal centers. *Science*. **351**, 1048–1054 (2016).

10. H. J. Snippert, L. G. van der Flier, T. Sato, J. H. van Es, M. van den Born, C. Kroon-Veenboer, N. Barker, A. M. Klein, J. van Rheenen, B. D. Simons, H. Clevers, Intestinal Crypt Homeostasis Results from Neutral Competition between Symmetrically Dividing Lgr5 Stem Cells. *Cell*. **143**, 134–144 (2010).

11. J. Livet, T. A. Weissman, H. Kang, R. W. Draft, J. Lu, R. A. Bennis, J. R. Sanes, J. W. Lichtman, Transgenic strategies for combinatorial expression of fluorescent proteins in the nervous system. *Nature*. **450**, 56–62 (2007).

12. T. A. Schwickert, R. L. Lindquist, G. Shakhar, G. Livshits, D. Skokos, M. H. Kosco-Vilbois, M. L. Dustin, M. C. Nussenzweig, In vivo imaging of germinal centres reveals a dynamic open structure. *Nature*. **446**, 83–87 (2007).

13. I. Dogan, B. Bertocci, V. Vilmont, F. Delbos, J. Mégret, S. Storck, C.-A. Reynaud, J.-C. Weill, Multiple layers of B cell memory with different effector functions. *Nat. Immunol.* **10**, 1292–1299 (2009).

14. R. Shinnakasu, T. Inoue, K. Kometani, S. Moriyama, Y. Adachi, M. Nakayama, Y. Takahashi, H. Fukuyama, T. Okada, T. Kurosaki, Regulated selection of germinal-center cells into the memory B cell compartment. *Nat. Immunol.* **17**, 861–869 (2016).

15. T.-A. Y. Shih, M. Roederer, M. C. Nussenzweig, Role of antigen receptor affinity in T cell–independent antibody responses in vivo. *Nat. Immunol.* **3**, 399–406 (2002).

16. T. A. Schwickert, B. Alabyev, T. Manser, M. C. Nussenzweig, Germinal center reutilization by newly activated B cells. *J. Exp. Med.* **206**, 2907–2914 (2009).

17. A. Fairhurst, S. Hwang, A. Wang, X. Tian, C. Boudreaux, X. J. Zhou, J. Casco, Q. Li, J. E. Connolly, E. K. Wakeland, *Yaa* autoimmune phenotypes are conferred by overexpression of TLR7. *Eur. J. Immunol.* **38**, 1971–1978 (2008).

18. T. A. Schwickert, G. D. Victora, D. R. Fooksman, A. O. Kamphorst, M. R. Mugnier, A. D. Gitlin, M. L. Dustin, M. C. Nussenzweig, A dynamic T cell–limited checkpoint regulates affinity-dependent B cell entry into the germinal center. *J. Exp. Med.* **208**, 1243–1252 (2011).

19. C. D. C. Allen, T. Okada, H. L. Tang, J. G. Cyster, Imaging of Germinal Center Selection Events During Affinity Maturation. *Science*. **315**, 528–531 (2007).

20. R. V. H. De Carvalho, J. Ersching, A. Barbulescu, A. Hobbs, T. B. R. Castro, L. Mesin, J. T. Jacobsen, B. K. Phillips, H.-H. Hoffmann, R. Parsa, M. C. C. Canesso, C. R. Nowosad, A. Feng, S. R. Leist, R. S. Baric, E. Yang, P. J. Utz, G. D. Victora, Clonal replacement sustains long-lived germinal centers primed by respiratory viruses. *Cell*. **186**, 131-146.e13 (2023).

21. T. Hägglöf, M. Cipolla, M. Loewe, S. T. Chen, L. Mesin, H. Hartweger, M. A. ElTanbouly, A. Cho, A. Gazumyan, V. Ramos, L. Stamatatos, T. Y. Oliveira, M. C. Nussenzweig, C. Viant, Continuous germinal center invasion contributes to the diversity of the immune response. *Cell*. **186**, 147-161.e15 (2023).

22. L. Mesin, A. Schiepers, J. Ersching, A. Barbulescu, C. B. Cavazzoni, A. Angelini, T. Okada, T. Kurosaki, G. D. Victora, Restricted Clonality and Limited Germinal Center Reentry Characterize Memory B Cell Reactivation by Boosting. *Cell*. **180**, 92-106.e11 (2020).

23. A. Cappione, J. H. Anolik, A. Pugh-Bernard, J. Barnard, P. Dutcher, G. Silverman, I. Sanz, Germinal center exclusion of autoreactive B cells is defective in human systemic lupus erythematosus. *J. Clin. Invest.* **115**, 3205–3216 (2005).

24. J. M. J. Tas, J.-H. Koo, Y.-C. Lin, Z. Xie, J. M. Steichen, A. M. Jackson, B. M. Hauser, X. Wang, C. A. Cottrell, J. L. Torres, J. E. Warner, K. H. Kirsch, S. R. Weldon, B. Groschel, B. Nogal, G. Ozorowski, S. Bangaru, N. Phelps, Y. Adachi, S. Eskandarzadeh, M. Kubitz, D. R. Burton, D. Lingwood, A. G. Schmidt, U. Nair, A. B. Ward, W. R. Schief, F. D. Batista, Antibodies from primary humoral responses modulate the recruitment of naive B cells during secondary responses. *Immunity*. **55**, 1856-1871.e6 (2022).

25. C.-H. Yeh, T. Nojima, M. Kuraoka, G. Kelsoe, Germinal center entry not selection of B cells is controlled by peptide-MHCII complex density. *Nat. Commun.* **9**, 928 (2018).

26. C. Soni, O. A. Perez, W. N. Voss, J. N. Pucella, L. Serpas, J. Mehl, K. L. Ching, J. Goike, G. Georgiou, G. C. Ippolito, V. Sisirak, B. Reizis, Plasmacytoid Dendritic Cells and Type I Interferon Promote Extrafollicular B Cell Responses to Extracellular Self-DNA. *Immunity*. **52**, 1022-1038.e7 (2020).

27. G. J. Brown, P. F. Cañete, H. Wang, A. Medhavy, J. Bones, J. A. Roco, Y. He, Y. Qin, J. Cappello, J. I. Ellyard, K. Bassett, Q. Shen, G. Burgio, Y. Zhang, C. Turnbull, X. Meng, P. Wu, E. Cho, L. A. Miosge, T. D. Andrews, M. A. Field, D. Tvorogov, A. F. Lopez, J. J. Babon, C. A. López, Á. Gónzalez-Murillo, D. C. Garulo, V. Pascual, T. Levy, E. J. Mallack, D. G. Calame, T. Lotze, J. R. Lupski, H. Ding, T. R. Ullah, G. D. Walters, M. E. Koina, M. C. Cook, N. Shen, C. de Lucas Collantes, B. Corry, M. P. Gantier, V. Athanasopoulos, C. G. Vinuesa, TLR7 gain-of-function genetic variation causes human lupus. *Nature*. **605**, 349–356 (2022).

28. J. S. Poovassery, G. A. Bishop, Type I IFN Receptor and the B Cell Antigen Receptor Regulate TLR7 Responses via Distinct Molecular Mechanisms. *J. Immunol.* **189**, 1757–1764 (2012).

29. E. C. Baechler, F. M. Batliwalla, G. Karypis, P. M. Gaffney, W. A. Ortmann, K. J. Espe, K. B. Shark, W. J. Grande, K. M. Hughes, V. Kapur, P. K. Gregersen, T. W. Behrens, Interferon-inducible gene expression signature in peripheral blood cells of patients with severe lupus. *Proc. Natl. Acad. Sci.* **100**, 2610–2615 (2003).

30. S. W. Jackson, H. M. Jacobs, T. Arkatkar, E. M. Dam, N. E. Scharping, N. S. Kolhatkar, B. Hou, J. H. Buckner, D. J. Rawlings, B cell IFN-γ receptor signaling promotes autoimmune germinal centers via cell-intrinsic induction of BCL-6. *J. Exp. Med.* **213**, 733–750 (2016).

31. P. P. Domeier, S. B. Chodisetti, S. L. Schell, Y. I. Kawasawa, M. J. Fasnacht, C. Soni, Z. S. M. Rahman, B-Cell-Intrinsic Type 1 Interferon Signaling Is Crucial for Loss of Tolerance and the Development of Autoreactive B Cells. *Cell Rep.* **24**, 406–418 (2018).

32. A. Das, B. A. Heesters, A. Bialas, J. O'Flynn, I. R. Rifkin, J. Ochando, N. Mittereder, G. Carlesso, R. Herbst, M. C. Carroll, Follicular Dendritic Cell Activation by TLR Ligands Promotes Autoreactive B Cell Responses. *Immunity*. **46**, 106–119 (2017).

33. L. F. Voss, A. J. Howarth, T. R. Wittenborn, S. Hummelgaard, K. Juul-Madsen, K. S. Kastberg, M. K. Pedersen, L. Jensen, A. D. Papanastasiou, T. Vorup-Jensen, K. Weyer, S. E. Degn, "Extrafollicular responses are sufficient to drive initiation of autoimmunity and early disease hallmarks of lupus" (preprint, Immunology, 2022), , doi:10.1101/2022.03.04.482991.

34. G. McDonald, C. O. Medina, M. Pilichowska, J. F. Kearney, R. Shinkura, E. Selsing, H. H. Wortis, T. Honjo, T. Imanishi-Kari, Accelerated Systemic Autoimmunity in the Absence of Somatic Hypermutation in 564Igi: A Mouse Model of Systemic Lupus with Knocked-In Heavy and Light Chain Genes. *Front. Immunol.* **8**, 1094 (2017).

35. Q. Zhang, P. Bastard, Z. Liu, J. Le Pen, M. Moncada-Velez, J. Chen, M. Ogishi, I. K. D. Sabli, S. Hodeib, C. Korol, J. Rosain, K. Bilguvar, J. Ye, A. Bolze, B. Bigio, R. Yang, A. A. Arias, Q. Zhou, Y. Zhang, F. Onodi, S. Korniotis, L. Karpf, Q. Philippot, M. Chbihi, L.

Bonnet-Madin, K. Dorgham, N. Smith, W. M. Schneider, B. S. Razooky, H.-H. Hoffmann,
E. Michailidis, L. Moens, J. E. Han, L. Lorenzo, L. Bizien, P. Meade, A.-L. Neehus, A. C.
Ugurbil, A. Corneau, G. Kerner, P. Zhang, F. Rapaport, Y. Seeleuthner, J. Manry, C.
Masson, Y. Schmitt, A. Schlüter, T. Le Voyer, T. Khan, J. Li, J. Fellay, L. Roussel, M.
Shahrooei, M. F. Alosaimi, D. Mansouri, H. Al-Saud, F. Al-Mulla, F. Almourfi, S. Z. Al-
Muhsen, F. Alsohime, S. Al Turki, R. Hasanato, D. Van De Beek, A. Biondi, L. R. Bettini,
M. D'Angio', P. Bonfanti, L. Imberti, A. Sottini, S. Paghera, E. Quiros-Roldan, C. Rossi,
A. J. Oler, M. F. Tompkins, C. Alba, I. Vandernoot, J.-C. Goffard, G. Smits, I. Migeotte, F.
Haerynck, P. Soler-Palacin, A. Martin-Nalda, R. Colobran, P.-E. Morange, S. Keles, F.
Çölkesen, T. Ozcelik, K. K. Yasar, S. Senoglu, Ş. N. Karabela, C. Rodríguez-Gallego, G.
Novelli, S. Hraiech, Y. Tandjaoui-Lambiotte, X. Duval, C. Laouénan, COVID-STORM
Clinicians, COVID Clinicians, Imagine COVID Group, French COVID Cohort Study
Group, CoV-Contact Cohort, Amsterdam UMC Covid-19 Biobank, COVID Human
Genetic Effort, NIAID-USUHS/TAGC COVID Immunity Group, A. L. Snow, C. L.
Dalgard, J. D. Milner, D. C. Vinh, T. H. Mogensen, N. Marr, A. N. Spaan, B. Boisson, S.
Boisson-Dupuis, J. Bustamante, A. Puel, M. J. Ciancanelli, I. Meyts, T. Maniatis, V.
Soumelis, A. Amara, M. Nussenzweig, A. García-Sastre, F. Krammer, A. Pujol, D. Duffy,
R. P. Lifton, S.-Y. Zhang, G. Gorochov, V. Béziat, E. Jouanguy, V. Sancho-Shimizu, C. M.
Rice, L. Abel, L. D. Notarangelo, A. Cobat, H. C. Su, J.-L. Casanova, G. Foti, G. Bellani,
G. Citerio, E. Contro, A. Pesci, M. G. Valsecchi, M. Cazzaniga, J. Abad, S. Aguilera-
Albesa, O. M. Akcan, I. A. Darazam, J. C. Aldave, M. A. Ramos, S. A. Nadji, G. Alkan, J.
Allardet-Servent, L. M. Allende, L. Alsina, M.-A. Alyanakian, B. Amador-Borrero, Z.
Amoura, A. Antolí, S. Arslan, S. Assant, T. Auguet, A. Azot, F. Bajolle, A. Baldolli, M.

Ballester, H. B. Feldman, B. Barrou, A. Beurton, A. Bilbao, G. Blanchard-Rohner, I.

Blanco, A. Blandinières, D. Blazquez-Gamero, M. Bloomfield, M. Bolivar-Prados, R.

Borie, C. Bosteels, A. A. Bousfiha, C. Bouvattier, O. Boyarchuk, M. R. P. Bueno, J.

Bustamante, J. J. Cáceres Agra, S. Calimli, R. Capra, M. Carrabba, C. Casasnovas, M.

Caseris, M. Castelle, F. Castelli, M. C. De Vera, M. V. Castro, E. Catherinot, M.

Chalumeau, B. Charbit, M. P. Cheng, P. Clavé, B. Clotet, A. Codina, F. Colkesen, F.

Çölkesen, R. Colobran, C. Comarmond, D. Dalmau, D. R. Darley, N. Dauby, S. Dauger, L.

De Pontual, A. Dehban, G. Delplancq, A. Demoule, J.-L. Diehl, S. Dobbelaere, S. Durand,

W. Eldars, M. Elgamal, M. H. Elnagdy, M. Emiroglu, E. H. Erdeniz, S. E. Aytekin, R.

Euvrard, R. Evcen, G. Fabio, L. Faivre, A. Falck, M. Fartoukh, M. Faure, M. F. Arquero, C.

Flores, B. Francois, V. Fumadó, F. Fusco, B. G. Solis, P. Gaussem, J. Gil-Herrera, L.

Gilardin, M. G. Alarcon, M. Girona-Alarcón, J.-C. Goffard, F. Gok, R. González-

Montelongo, A. Guerder, Y. Gul, S. N. Guner, M. Gut, J. Hadjadj, F. Haerynck, R.

Halwani, L. Hammarström, N. Hatipoglu, E. Hernandez-Brito, C. Heijmans, M. S.

Holanda-Peña, J. P. Horcajada, L. Hoste, E. Hoste, S. Hraiech, L. Humbert, A. D. Iglesias,

A. Íñigo-Campos, M. Jamme, M. J. Arranz, I. Jordan, P. Jorens, F. Kanat, H. Kapakli, I.

Kara, A. Karbuz, K. K. Yasar, S. Keles, Y. K. Demirkol, A. Klocperk, Z. J. Król, P.

Kuentz, Y. W. M. Kwan, J.-C. Lagier, B. N. Lambrecht, Y.-L. Lau, F. Le Bourgeois, Y.-S.

Leo, R. L. Lopez, D. Leung, M. Levin, M. Levy, R. Lévy, Z. Li, A. Linglart, B. Loeys, J.

M. Lorenzo-Salazar, C. Louapre, C. Lubetzki, C.-E. Luyt, D. C. Lye, D. Mansouri, M.

Marjani, J. M. Pereira, A. Martin, D. M. Pueyo, J. Martinez-Picado, I. Marzana, A.

Mathian, L. R. B. Matos, G. V. Matthews, J. Mayaux, J.-L. Mège, I. Melki, J.-F. Meritet, O.

Metin, I. Meyts, M. Mezidi, I. Migeotte, M. Millereux, T. Mirault, C. Mircher, M.

Mirsaeidi, A. M. Melián, A. M. Martinez, P. Morange, C. Mordacq, G. Morelle, S. Mouly,

A. Muñoz-Barrera, L. Naesens, C. Nafati, J. F. Neves, L. F. P. Ng, Y. N. Medina, E. N.

Cuadros, J. G. Ocejo-Vinyals, Z. Orbak, M. Oualha, T. Özçelik, Q. Pan-Hammarström, C.

Parizot, T. Pascreau, E. Paz-Artal, S. Pellegrini, R. P. De Diego, A. Philippe, Q. Philippot,

L. Planas-Serra, D. Ploin, J. Poissy, G. Poncelet, M. Pouletty, P. Quentric, D. Raoult, A.-S.

Rebillat, I. Reisli, P. Ricart, J.-C. Richard, N. Rivet, J. G. Rivière, G. R. Blanch, C.

Rodrigo, C. Rodriguez-Gallego, A. Rodríguez-Palmero, C. S. Romero, A. Rothenbuhler, F.

Rozenberg, M. Y. Ruiz Del Prado, J. S. Riera, O. Sanchez, S. Sánchez-Ramón, A. Schluter,

M. Schmidt, C. E. Schweitzer, F. Scolari, A. Sediva, L. M. Seijo, D. Sene, S. Senoglu, M.

R. J. Seppänen, A. S. Ilovich, M. Shahrooei, H. Slabbynck, D. M. Smadja, A. Sobh, X. S.

Moreno, J. Solé-Violán, C. Soler, P. Soler-Palacín, Y. Stepanovskiy, A. Stoclin, F.

Taccone, Y. Tandjaoui-Lambiotte, J.-L. Taupin, S. J. Tavernier, B. Terrier, C. Thumerelle,

G. Tomasoni, J. Toubiana, J. T. Alvarez, S. Trouillet-Assant, J. Troya, A. Tucci, M. V.

Ursini, Y. Uzunhan, P. Vabres, J. Valencia-Ramos, E. Van Braeckel, S. Van De Velde, A.

M. Van Den Rym, J. Van Praet, I. Vandernoot, H. Vatansev, V. Vélez-Santamaria, S. Viel,

C. Vilain, M. E. Vilaire, A. Vincent, G. Voiriot, F. Vuotto, A. Yosunkaya, B. E. Young, F.

Yucel, F. Zannad, M. Zatz, A. Belot, C. Bole-Feysot, S. Lyonnet, C. Masson, P. Nitschke,

A. Pouliet, Y. Schmitt, F. Tores, M. Zarhrate, L. Abel, C. Andrejak, F. Angoulvant, D.

Bachelet, R. Basmaci, S. Behillil, M. Beluze, D. Benkerrou, K. Bhavsar, F. Bompart, L.

Bouadma, M. Bouscambert, M. Caralp, M. Cervantes-Gonzalez, A. Chair, A. Coelho, C.

Couffignal, S. Couffin-Cadiergues, E. D'Ortenzio, C. Da Silveira, M.-P. Debray, D.

Deplanque, D. Descamps, M. Desvallées, A. Diallo, A. Diouf, C. Dorival, F. Dubos, X.

Duval, P. Eloy, V. V. Enouf, H. Esperou, M. Esposito-Farese, M. Etienne, N. Ettalhaoui, N.

Gault, A. Gaymard, J. Ghosn, T. Gigante, I. Gorenne, J. Guedj, A. Hoctin, I. Hoffmann, S. Jaafoura, O. Kafif, F. Kaguelidou, S. Kali, A. Khalil, C. Khan, C. Laouénan, S. Laribi, M. Le, Q. Le Hingrat, S. Le Mestre, H. Le Nagard, F.-X. Lescure, Y. Lévy, C. Levy-Marchal, B. Lina, G. Lingas, J. C. Lucet, D. Malvy, M. Mambert, F. Mentré, N. Mercier, A. Meziane, H. Mouquet, J. Mullaert, N. Neant, M. Noret, J. Pages, A. Papadopoulos, C. Paul, N. Peiffer-Smadja, V. Petrov-Sanchez, G. Peytavin, O. Picone, O. Puéchal, M. Rosa-Calatrava, B. Rossignol, P. Rossignol, C. Roy, M. Schneider, C. Semaille, N. S. Mohammed, L. Tagherset, C. Tardivon, M.-C. Tellier, F. Téoulé, O. Terrier, J.-F. Timsit, T. Trioux, C. Tual, S. Tubiana, S. Van Der Werf, N. Vanel, A. Veislinger, B. Visseaux, A. Wiedemann, Y. Yazdanpanah, L. Alavoine, K. K. A. Amat, S. Behillil, J. Bielicki, P. Bruijning, C. Burdet, E. Caumes, C. Charpentier, B. Coignard, Y. Costa, S. Couffin-Cadiergues, F. Damond, A. Dechanet, C. Delmas, D. Descamps, X. Duval, J.-L. Ecobichon, V. Enouf, H. Espérou, W. Frezouls, N. Houhou, E. Ilic-Habensus, O. Kafif, J. Kikoine, Q. Le Hingrat, D. Lebeaux, A. Leclercq, J. Lehacaut, S. Letrou, B. Lina, J.-C. Lucet, D. Malvy, P. Manchon, M. Mandic, M. Meghadecha, J. Motiejunaite, M. Nouroudine, V. Piquard, A. Postolache, C. Quintin, J. Rexach, L. Roufai, Z. Terzian, M. Thy, S. Tubiana, S. Van Der Werf, V. Vignali, B. Visseaux, Y. Yazdanpanah, M. Van Agtmael, A. G. Algera, F. Van Baarle, D. Bax, M. Beudel, H. J. Bogaard, M. Bomers, L. Bos, M. Botta, J. De Brabander, G. De Bree, M. C. Brouwer, S. De Bruin, M. Bugiani, E. Bulle, O. Chouchane, A. Cloherty, P. Elbers, L. Fleuren, S. Geerlings, B. Geerts, T. Geijtenbeek, A. Girbes, B. Goorhuis, M. P. Grobusch, F. Hafkamp, L. Hagens, J. Hamann, V. Harris, R. Hemke, S. M. Hermans, L. Heunks, M. W. Hollmann, J. Horn, J. W. Hovius, M. D. De Jong, R. Koning, N. Van Mourik, J. Nellen, F. Paulus, E. Peters, T. Van Der Poll, B.

Preckel, J. M. Prins, J. Raasveld, T. Reijnders, M. Schinkel, M. J. Schultz, A. Schuurman, K. Sigaloff, M. Smit, C. S. Stijnis, W. Stilma, C. Teunissen, P. Thoral, A. Tsonas, M. Van Der Valk, D. Veelo, A. P. J. Vlaar, H. De Vries, M. Van Vugt, W. J. Wiersinga, D. Wouters, A. H. (Koos) Zwinderman, D. Van De Beek, L. Abel, A. Aiuti, S. Al Muhsen, F. Al-Mulla, M. S. Anderson, A. A. Arias, H. B. Feldman, D. Bogunovic, A. Bolze, A. Bondarenko, A. A. Bousfiha, P. Brodin, Y. Bryceson, C. D. Bustamante, M. Butte, G. Casari, S. Chakravorty, J. Christodoulou, E. Cirulli, A. Condino-Neto, M. A. Cooper, C. L. Dalgard, A. David, J. L. DeRisi, M. Desai, B. A. Drolet, S. Espinosa, J. Fellay, C. Flores, J. L. Franco, P. K. Gregersen, F. Haerynck, D. Hagin, R. Halwani, J. Heath, S. E. Henrickson, E. Hsieh, K. Imai, Y. Itan, T. Karamitros, K. Kisand, C.-L. Ku, Y.-L. Lau, Y. Ling, C. L. Lucas, T. Maniatis, D. Mansouri, L. Marodi, I. Meyts, J. Milner, K. Mironska, T. Mogensen, T. Morio, L. F. P. Ng, L. D. Notarangelo, A. Novelli, G. Novelli, C. O'Farrelly, S. Okada, T. Ozcelik, R. P. De Diego, A. M. Planas, C. Prando, A. Pujol, L. Quintana-Murci, L. Renia, A. Renieri, C. Rodríguez-Gallego, V. Sancho-Shimizu, V. Sankaran, K. S. Barrett, M. Shahrooei, A. Snow, P. Soler-Palacín, A. N. Spaan, S. Tangye, S. Turvey, F. Uddin, M. J. Uddin, D. Van De Beek, S. E. Vazquez, D. C. Vinh, H. Von Bernuth, N. Washington, P. Zawadzki, H. C. Su, J.-L. Casanova, H. Jing, W. Tung, C. R. Luthers, B. M. Bauman, S. Shafer, L. Zheng, Z. Zhang, S. Kubo, S. D. Chauvin, K. Meguro, E. Shaw, M. Lenardo, J. Lack, E. Karlins, D. M. Hupalo, J. Rosenberger, G. Sukumar, M. D. Wilkerson, X. Zhang, Inborn errors of type I IFN immunity in patients with life-threatening COVID-19. *Science*. **370**, eabd4570 (2020).

36. S. Mantovani, S. Daga, C. Fallerini, M. Baldassarri, E. Benetti, N. Picchiotti, F. Fava, A. Gallì, S. Zibellini, M. Bruttini, M. Palmieri, S. Croci, S. Amitrano, D. Alaverdian, K.

Capitani, S. Furini, F. Mari, I. Meloni, GEN-COVID Multicenter Study, F. Montagnani, M. Tumbarello, I. Rancan, M. Fabbiani, B. Rossetti, L. Bergantini, M. D'Alessandro, P. Cameli, D. Bennett, F. Anedda, S. Marcantonio, S. Scolletta, F. Franchi, M. A. Mazzei, S. Guerrini, E. Conticini, L. Cantarini, B. Frediani, D. Tacconi, C. S. Raffaelli, M. Feri, A. Donati, R. Scala, L. Guidelli, G. Spargi, M. Corridi, C. Nencioni, L. Croci, G. P. Caldarelli, D. Romani, P. Piacentini, M. Bandini, E. Desanctis, S. Cappelli, A. Canaccini, A. Verzuri, V. Anemoli, M. Pisani, A. Ognibene, A. Pancrazzi, M. Lorubbio, M. Vaghi, A. D'Arminio Monforte, F. G. Miraglia, R. Bruno, M. Vecchia, M. Girardis, S. Venturelli, S. Busani, A. Cossarizza, A. Antinori, A. Vergori, A. Emiliozzi, S. Rusconi, M. Siano, A. Gabrieli, A. Riva, D. Francisci, E. Schiaroli, F. Paciosi, A. Tommasi, P. G. Scotton, F. Andretta, S. Panese, S. Baratti, R. Scaggiante, F. Gatti, S. G. Parisi, F. Castelli, E. Quiros-Roldan, M. D. Antoni, I. Zanella, M. Della Monica, C. Piscopo, M. Capasso, R. Russo, I. Andolfo, A. Iolascon, G. Fiorentino, M. Carella, M. Castori, F. Aucella, P. Raggi, R. Perna, M. Bassetti, A. Di Biagio, M. Sanguinetti, L. Masucci, A. Guarnaccia, S. Valente, O. De Vivo, G. Doddato, M. Lista, G. Beligni, F. Valentino, K. Zguro, R. Tita, A. Giliberti, M. A. Mencarelli, C. L. Rizzo, A. M. Pinto, F. Ariani, L. Di Sarno, E. Bargagli, M. Mandalà, A. Giorli, L. Salerni, P. Zucchi, P. Parravicini, E. Menatti, T. Trotta, F. Giannattasio, G. Coiro, F. Lena, G. Lacerenza, C. Mussini, E. Martinelli, L. Tavecchia, M. A. Belli, L. Crotti, G. Parati, M. Sanarico, F. Raimondi, F. Biscarini, A. Stella, T. Bachetti, M. T. La Rovere, S. Ludovisi, M. Bussotti, S. Dei, S. Ravaglia, R. Artuso, E. Andreucci, G. Gori, A. Pagliazzi, E. Fiorentini, A. Perrella, F. Bianchi, P. Bergomi, E. Catena, R. Colombo, S. Luchi, G. Morelli, P. Petrocelli, S. Iacopini, S. Modica, S. Baroni, F. V. Segala, M. Falcone, G. Tiseo, C. Barbieri, T. Matucci, D. Grassi, C. Ferri, F. Marinangeli, F. Brancati, A. Vincenti, V.

Borgo, S. Lombardi, M. Lenzi, M. A. Di Pietro, F. Vichi, B. Romanin, L. Attala, C. Costa, A. Gabbuti, R. Menè, M. Colaneri, P. Casprini, G. Merla, G. M. Squeo, M. Maffezzoni, E. Frullanti, M. U. Mondelli, A. Renieri, Rare variants in Toll-like receptor 7 results in functional impairment and downregulation of cytokine-mediated signaling in COVID-19 patients. *Genes Immun.* **23**, 51–56 (2022).

37. G. D. Victora, T. A. Schwickert, D. R. Fooksman, A. O. Kamphorst, M. Meyer-Hermann, M. L. Dustin, M. C. Nussenzweig, Germinal Center Dynamics Revealed by Multiphoton Microscopy with a Photoactivatable Fluorescent Reporter. *Cell*. **143**, 592–605 (2010).

38. A. Le Bon, C. Thompson, E. Kamphuis, V. Durand, C. Rossmann, U. Kalinke, D. F. Tough, Cutting Edge: Enhancement of Antibody Responses Through Direct Stimulation of B and T Cells by Type I IFN. *J. Immunol.* **176**, 2074–2078 (2006).

39. M. R. Lamprecht, D. M. Sabatini, A. E. Carpenter, CellProfiler™: free, versatile software for automated biological image analysis. *BioTechniques*. **42**, 71–75 (2007).

Acknowledgments:

We thank E.M. Carroll for administrative and technical support and all the members of the M.C.C. lab for help with experiments and feedback on the manuscript. Imaging was performed at the Program in Cellular and Molecular Medicine (PCMM) microscopy core, with technical support from Harry Leung. Flow cytometry and cell sorting were performed at the PCMM flow cytometry core, with technical support from Jodene K. Moore.

Funding:

National Institutes of Health grant R01 AI130307 (MCC)

National Institutes of Health grant R01 AR074105 (MCC)

European Union's Horizon 2020 Research and Innovation Program, Marie Sklodowska-Curie

grant 796988 (TB)

Author contributions:

Conceptualization: MCC, TB

Methodology: MCC, TB, KO, CC, CEP

Investigation: TB, KO, SR

Formal analysis: TB, KO

Visualization: TB

Funding acquisition: MCC TB

Project administration: MCC

Supervision: MCC

Writing – original draft: TB

Writing – review & editing: MCC, TB, KO, CC

Competing interests:

The authors declare that they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials.

Figures:

Figure 1. TLR7-dependent wild-type B cell dominance in spontaneous GCs in 564Igi:WT parabionts.

A. Schematic depiction of the parabiosis model (WT = wild-type).

B. Representative FACS dot plot of the proportion of WT or 564Igi B cells in the circulation of the WT and 564Igi parabionts, at 6 weeks parabiosis.

C. Representative FACS dot plot of the proportion of WT (CD45.2) or WT (CD45.1) cells in the circulation of the WT and WT control parabiosis, at 8-9 weeks.

D. Frequency of wild-type (WT) B cells in the circulation of 564Igi parabiosis partner (grey circles) or WT parabiosis partner (control parabiosis, black circles) over time. Week 1 (564Igi n=11 vs WT n=6), Week 2 (n=5 vs n=6), Week 3 (n=10 vs n6), Week 4 (n=8 vs n=4), Week 5 (n=10 vs n=6), Week 6 (n=10 v n=6), Week 7 (n=9 vs n=2), Week 8-9 (n=10 vs n=6).

E. Frequency of 564Igi (blue circles) or WT (control parabiosis, yellow circles) B cells in the circulation of WT parabiosis partner mouse over time. As in Figure 1B, but week 2 (564Igi n=6 vs WT n=6), Week 6 (n=9 vs n=6).

F. Frequency of 564Igi B cells in WT parabiosis partner (left column), and in the 564Igi partner, the proportion of WT B cells in the total splenic B cell population or within the germinal center (GC) B cell population within the spleen 8 weeks following surgery.

G. Parabiosis pairing of Toll-Like receptor 7 knock-out (TLR KO) mouse with 564Igi mouse for 8 weeks, showing the proportion of TLR7 KO B cells within circulation (n=5), splenic B cell pool (n=6) and splenic GC B cell population (n=6) of the 564Igi parabiosis partner.

H. Parabiosis pairing of Toll-Like receptor 7 knock-out (TLR KO) mouse with 564Igi mouse for 8 weeks, showing the proportion of TLR7 KO B cells within the mesenteric lymph node

1 (mesLN) B cell pool (n=6) and mesLN GC B cell population (n=6) of the 564Igi parabiosis

2 partner.

3 Significance is indicated as ***P<0.001, **P<0.01, *P<0.05, and P>0.05 (not significant (NS))

4 by Mann-Whitney test for Fig 1D, 1E and by Wilcoxon paired test for Fig 1F, 1G, 1H.

Figure 2. Temporary parabiosis highlights WT B cell persistence in the spontaneous

germinal center

A. Schematic representation of parabiosis surgery followed by surgical separation at 2 weeks of

parabiosis and subsequent follow-up.

B. The proportion of 564Igi (blue circles) or WT (Control parabiosis, yellow circles) B cells in

the circulation of WT parabiosis partner mouse over time. Week 1 (564Igi n=8 vs WT n=5),

Week 2 (n=19 vs n=5), Week 3 (n=12 vs n=4), Week 4 (n=11 vs n=4), Week 5 (n=11 vs

n=4), Week 6 (n=9 v n=4), Week 7 (n=7 vs n=4), Week 8-9 (n=8 vs n=3).

C. The proportion of wild-type (WT) B cells in the circulation of 564Igi (grey circles) or WT

(control parabiosis, black circles) parabiosis partner mouse over time. Number of mice as in

Figure 2B.

D. The proportion of 564Igi B cells in the WT parabiosis partner (left graph, n=9), and in the 564Igi parabiosis partner (right graph, n=10), the proportion of WT B cells in the circulation, splenic B cell population or splenic germinal center B cell population, after 2 weeks of parabiosis and subsequent 6 weeks of separation.

Significance is indicated as ***P<0.001, **P<0.01, *P<0.05, and P>0.05 (not significant (NS) by Mann-Whitney test for Fig 2B, 2C and Wilcoxon paired test for Fig 2D (Blood vs Spleen, Spleen vs Spleen GC).

Figure 3. Temporary parabiosis highlights WT B cell persistence and clonal expansion in autoreactive 564Igi GCs.

A. Schematic depiction of parabiosis and separation surgery of AID-Cre[ert2] Confetti mouse with 564Igi mouse to assess clonal evolution of WT B cells in 564Igi partner at day +4, +15, and +31 post-separation surgery.

B. Representative multi-photon images of the splenic GC of the 564Igi parabiosis mouse partner for quantification of color dominance. The scale bars indicate 100 μm.

C. Frequency of most frequent color (1st), second-most frequent color (2nd), and their sum observed in individual splenic GC at Day 3, 15, and 31 post-tamoxifen induction of Confetti

recombination in 564Igi parabiont. Day 3 (2 mice, 10 GC), day 15 (3 mice, 30 GC) and day 30 (4 mice, 18 GC).

D. The divergence index for the GC represented in (C).

E. The number of fate-mapped WT B cells for the GC represented in (C).

Significance is indicated as ***P<0.001, **P<0.01, *P<0.05, and P>0.05 (not significant (NS)) by Mann-Whitney test for Fig 3C,3D (compared to day 3) and ANOVA with posthoc test (Dunn) for Fig 3E.

Figure 4. Wild-type B cells rapidly enter the pre-existing autoreactive germinal center.

A. Schematic depiction of WT B cell adoptive transfer model and follow-up time points.

B. Representative FACs dot plot of WT B cells in the splenic B cell compartment and within
    the splenic germinal center B cell compartment, at day 7. Day 1 (n=1), day 2 (n=1), day 3
    (n=7), day 4 (n=7), day 5 (n=7), day 6 (n=7), day 7 (n=11), day 10 (n=6), day 14 (n=9), day
    21 (n=7), day 56 (n=5).

C. The proportion of WT B cells within the splenic B cell pool (grey line, open circle) and
    splenic germinal center B cell pool (black line, closed circle) over time.

D. Bar charts indicating the frequency (left) and absolute number (right) of GCs populated by
    WT and 564Igi, 564Igi only, or WT only B cells (left). Each symbol indicates a mouse.

E. Pie charts demonstrating the frequency of GCs populated by both WT and 564Igi, 564Igi only or WT only B cells as a frequency of white pulp areas (WPAs) in individual mice. No GCs with only B6.2 B cells were identified.

F. Representative confocal immunofluorescence microscopy image showing GC composition in a splenic section; GCs were identified by GL7 (green) staining and confirmed by the overlapping CD21bright staining of FDC networks in a consecutive slice. Anti-CD45.1 (blue) and anti-CD45.2 (red) staining was used to identify 564Igi- and WT-derived cells, respectively. Arrows indicate GCs with different patterns of WT residency. GCs contained within boxes with roman numerals are magnified in the smaller panels to the right.

G. Bar chart showing the area of GCs with WT and 564Igi and 564Igi only residents. Each symbol indicates a GC.

Wilcoxon paired test: Fig 4B. Fig 4C: One-way ANOVA with Bonferroni's multiple comparison test. Fig 4F: Mann-Whitney test. Values are mean+/- SD. Significance is indicated as ***$P<0.001$, **$P<0.01$, *$P<0.05$, and $P>0.05$ (not significant (NS)).

A

B

C

Figure 5. Molecular requirements for wild-type B cell entry and participation in spontaneous germinal centers.

A. Schematic representation of B1-8 B cell adoptive transfer.

B. Representative FACSplot and gating for identifying NP-specific B cells (left panel). The proportion of Lambda-positive (NP-specific) B cells within the splenic B cell compartment, splenic GC B cells in the 564Igi mouse, 7 days post-transfer (n=6) (right panel).

C. The proportion of germinal B cells derived from WT, MHC2 KO, TLR7 KO, SLE1-Yaa, and IFNAR KO transferred B cells within the splenic GC B cell compartment, 7 days post-transfer. WT transfer n=10, MHC2 KO transfer n=8, TLR7 KO transfer n=11, SLE1-Yaa n=7, IFNAR KO transfer n=8.

1   Significance is indicated as ***P<0.001, **P<0.01, *P<0.05, and P>0.05 (not significant (NS)).

2   The statistical test used is the Mann-Whitney test for Fig 5B, C (compared to WT).