Detecting Regime Shifts: Neurocomputational Substrates for Over- and Underreactions to Change

Mu-Chen Wang[1*]

George Wu[2]

Shih-Wei Wu[1,3*]

1. Institute of Neuroscience, National Yang Ming Chiao Tung University, Taipei, Taiwan

2. Booth School of Business, University of Chicago, Chicago, USA

3. Brain Research Center, National Yang Ming Chiao Tung University, Taipei, Taiwan

Corresponding Authors

Mu-Chen Wang <raccoon65.y@nycu.edu.tw> or Shih-Wei Wu <swwu@nycu.edu.tw >

ABSTRACT

The world constantly changes, with the underlying state of the world shifting from

one regime to another. The ability to detect a regime shift, such as the onset of a

pandemic or the end of a recession, significantly impacts individual decisions as well

as governmental policies. However, determining whether a regime has changed is

usually not obvious, as signals are noisy and reflective of the volatility of the

environment. We designed an fMRI paradigm that examines a stylized regime-shift

detection task. Human participants showed systematic over- and underreaction:

Overreaction was most commonly seen when signals were noisy but when

environments were stable and change is possible but unlikely. By contrast,

underreaction was observed when signals were precise but when environments were

unstable and hence change was more likely. These behavioral signatures are

consistent with the *system-neglect* computational hypothesis, which posits that

sensitivity or lack thereof to system parameters (noise and volatility) is central to

these behavioral biases. Guided by this computational framework, we found that

individual subjects' sensitivity to system parameters were represented by two distinct

brain networks. Whereas a frontoparietal network selectively represented individuals'

sensitivity to signal noise but not environment volatility, the ventromedial prefrontal

cortex (vmPFC) showed the opposite pattern. Further, these two networks were

involved in different aspects of regime-shift computations: while vmPFC correlated

with subjects' beliefs about change, the frontoparietal network represented the

strength of evidence in favor of regime shifts. Together, these results suggest that

regime-shift detection recruits belief-updating and evidence-evaluation networks and

that under- and overreactions arise from how sensitive these networks are to the

system parameters.

SIGNIFICANCE STATEMENT

Judging whether the world has changed, from the onset of a market boom to the end of a pandemic, is ubiquitous. The ability to detect regime shifts not only impacts individual decisions but also governmental policies. However, these judgments are hard to make because the signals we receive are noisy and reflective of the volatility of the environment. We find that people overreact to changes when they receive noisy signals in stable environments, but underreact when facing precise signals in unstable environments. Under- and overreactions can be read out by distinct brain networks according to their sensitivity in responding to different environmental parameters that impact regime changes. This suggests that parameter selectivity at the network level guides regime-shift detection.

1 INTRODUCTION

Judging whether the world has changed is ubiquitous, from public health officials

grappling with whether a pandemic surge has peaked, central banks figuring out

whether inflation is easing, investors discerning whether the electric car market is

getting traction, or romantic partners divining whether a relationship has soured. In all

of these examples, individuals must update their beliefs that the world has changed

based on a noisy signal, such as a drop in positive pandemic cases or a romantic

partner's suddenly mysterious behavior. In some cases, epidemiological or statistical

models provide guidance. However, in many, if not most cases, the determination of

whether a regime shift has occurred is made intuitively (Sanders & Manrodt, 2003).

We investigate intuitive judgments of regime-shift detection using a simple

empirical paradigm (Massey & Wu, 2005; Seifert et al., 2023). Although this

paradigm abstracts away some complications of real-world change detection, it

maintains the most central features of the problem: normatively, regime-shift

judgments reflect the *signals* from the environment as well as knowledge about the

system that produces the signals. The most recent time series of inflation rates,

pandemic cases, and sales of electric cars are all examples of signals. When pandemic

cases continue to decline in recent weeks, one might infer a shift from pandemic to

non-pandemic regime, only to learn a few weeks later that pandemic has resurged.

Indeed, signals such as the latest pandemic cases are seldom precise indications of the

true state of the world. Put differently, signals are, by and large, noisy. The noisier the

signals are, the less *diagnostic* they are of the underlying regime.

In addition, signals are affected by how likely the regime shifts from one to

another (transition probability). These two fundamental features or *system*

parameters—the diagnosticity of the signals and transition probability—can be

conceptualized as two independent aspects of the system that generates the signals.

Previous works on regime-shift detection has found that people tend to overreact to

change when they receive noisy signals (low signal diagnosticity) but nonetheless are

in a stable environment (small transition probability). By contrast, precise signals

(high signal diagnosticity) in an unstable environment (large transition probability)

typically results in underreaction (Benjamin, 2019; Brown & Steyvers, 2009; Massey

& Wu, 2005).

Massey and Wu (2005) proposed that over- and underreactions reflect *system neglect*—the tendency to respond primarily to signals and secondarily to the system parameters that produces the signals. The system-neglect hypothesis was derived from theoretical accounts of the determinants of confidence by Griffin and Tversky (1992). To explain system neglect, consider someone who is making judgments on whether a stock market had shifted from the bear to the bull market regime and has been given information about recent stock returns (signals), how frequent regime shifts happen (transition probability), and how similar the two regimes are (signal diagnosticity). If her judgments are solely based on the signals and not affected by transition probability and signal diagnosticity, she shows a *complete* neglect of the system parameters. Broadly, system neglect describes a lack of sensitivity—compared with normative Bayesian updating—to the system parameters. In the case of regime-shift detection, this leads to insufficient belief revision (i.e., underreaction) in diagnostic and unstable environments, where Bayesian updating requires a larger change in beliefs, and excessive belief change in noisy and stable environments (i.e., overreaction), where Bayesian updating calls for less pronounced belief revision. Empirical patterns akin to system neglect is not only observed in regime-shift detection, but also in other domains such as confidence judgments (Griffin & Tversky, 1992; Kraemer & Weber, 2004), demand forecasting (Kremer et al., 2011), and pricing decisions (Seifert et al., 2023). Under- and overreactions have been an active research topic in financial economics, often measured as reactions to stock market changes or firm news (Baker & Wurgler, 2007; Barberis et al., 1998; Daniel et al., 1998; De Bondt & Thaler, 1985; Nelson et al., 2001).

At the neurobiological level, change detection has been investigated in the context of reinforcement learning in dynamic environments where changes in the state of the world, such as reward distributions, take place during the experiments (Soltani & Izquierdo, 2019). Different behavioral paradigms, most notably reversal learning, and computational models were developed to investigate its neurocomputational substrates (Behrens et al., 2007; Izquierdo et al., 2017; McGuire et al., 2014; Muller et al., 2019; Nassar et al., 2010; Payzan-LeNestour & Bossaerts, 2011; Payzan-LeNestour et al., 2013). Key findings on the neural implementations for such learning include identifying brain areas and networks that track volatility in the environment (rate of change) (Behrens et al., 2007), the uncertainty or entropy of the current state of the environment (Muller et al., 2019), participants' beliefs about change (Kao et

al., 2020; McGuire et al., 2014; Payzan-LeNestour & Bossaerts, 2011), and their uncertainty about whether a change had occurred (Kao et al., 2020; McGuire et al., 2014). Evidence from several of the aforementioned studies (Behrens et al., 2007; Kao et al., 2020; McGuire et al., 2014) suggests that the dorsomedial frontal cortex (DMFC) is critical to learning in dynamic environments, as information about volatility, subjective beliefs and uncertainty about change converge in this brain region.

But how do biases in change detection arise in the brain? Although reinforcement learning studies provide valuable insights into change detection in the learning process, it remains unclear how biases in change detection—under- and overreactions to change—arise at the neural algorithmic and implementation levels. For example, it is unclear how a certain brain area, such as DMFC, that had been shown to represent environmental volatility, would contribute to under- and overreactions to change. In order to systematically characterize under- and overreactions, it would be critical to (1) adopt a well-established behavioral paradigm that robustly elicits these behavioral phenomena and (2) have computational frameworks suitable for developing neural hypotheses regarding under- and overreactions. To address these issues, in this study, we adopted the regime-shift detection task from Massey and Wu (2005) and their system-neglect computational framework. At the behavioral level, the regime-shift task is a well-established paradigm that robustly elicits under- and overreactions to change. At the algorithmic and implementation levels, the system-neglect framework provides a straightforward neurocomputational hypothesis regarding under- and overreactions. It predicts that for brain areas involved in regime-shift detection, under- and overreactions arise from their sensitivity or lack thereof in response to the system parameters.

We replicated previous behavioral findings on under- and overreactions (Massey & Wu, 2005). Using blood-oxygen-level-dependent (BOLD) functional magnetic resonance imaging (fMRI), we reported three key findings. First, we identified two distinct brain networks involved in regime-shift detection, with the ventromedial prefrontal cortex (vmPFC) and ventral striatum in representing subjects' reported beliefs about change and a frontoparietal network in evaluating the strength of change evidence. Second, we found that these two networks selectively respond to different system parameters: while the frontoparietal network represents individual subjects' sensitivity to signal diagnosticity but not transition probability, the vmPFC shows the

opposite pattern. Third, the neural sensitivity profiles were signal-dependent: frontoparietal network only represented individuals' sensitivity to signal diagnosticity when signals consistent with change appeared. By contrast, vmPFC represented individuals' sensitivity to transition probability regardless of whether subjects received signals consistent or inconsistent with change. Such signal-dependent representations led us to further examine and subsequently verify that they are indeed key properties of our system-neglect computational model. Together, these results suggest that regime-shift detection is implemented jointly by a belief-updating network (vmPFC-striatum) and evidence evaluation network (frontoparietal network) and that their sensitivity in response to different environmental parameters contribute to under- and overreactions to change. More broadly, we showed that neural data can reveal important properties of computational models that are overlooked in theoretical treatments and behavioral analyses.

RESULTS

In our *regime-shift detection* task (Fig. 1A), in each trial, subjects saw a series of sequentially presented sensory signals (red or blue balls). They were told that the signals came from one of two regimes, the red regime or the blue regime (Fig. 1B). Regimes were symmetric, for example, with a red regime consisting of 60 red balls and 40 blue balls and the corresponding blue regime consisting of 60 blue balls and 40 red balls. Each trial started with the red regime but could shift to the blue regime before each of the 10 periods in a trial. After seeing a new signal in each period, subjects provided a probability estimate that the current regime was the blue regime, i.e., a posterior probability of a regime shift. They were also instructed that once the regime has shifted from the red to the blue during a trial, the regime would remain in the blue regime until the end of the trial, i.e., the blue regime was a trapping or absorbing state. Our experimental paradigm hence follows Massey and Wu (2005). Note that, during a trial, subjects did not receive feedback—after making probability estimates in each period—on whether the regime had shifted and the monetary bonus earned as a result of accuracy in probability estimates (see *Methods* for details). Hence, subjects had no access to information about accuracy and rewards as she or he was making probability estimates.

We manipulated two system parameters, transition probability and signal diagnosticity (Fig. 1C). Transition probability, q, with possible values are 0.01, 0.05, and 0.1, specified the probability that the regime would shift from the red to the blue regime in any period. Signal diagnosticity, d, with possible values of 1.5, 3, and 9, captured the degree to which the two regimes differed. For example, an environment with high signal diagnosticity (e.g., $d = 9$) indicated that there were 9 times more red balls than blue balls in the red regime (a $90:10$ Red to Blue ratio) and 9 times more blue balls than the red balls in the blue regime (a $90:10$ Blue to Red ratio). Therefore, the weight that a signal (blue or red ball) carried was captured by the signal diagnosticity: in a low diagnostic environment, a blue signal most likely reflects no change in regime ($d = 1.5$, example on the left of Fig. 1D). By contrast, in a highly diagnostic environment, a blue signal very likely reveals a shift in regime ($d = 9$, example on the right of Fig. 1D). At the beginning of each trial, subjects were informed about the transition probability and signal diagnosticity in that trial. In the example trial sequence (Fig. 1A), the transition probability (indicated by "switch probability" in Fig. 1A) is 0.1 while the signal diagnosticity (indicated by "color

ratio" in Fig. 1A) is 1.5, with the red regime consisting of 60 red balls and 40 blue

balls and the blue regime consisting of 40 red balls and 60 blue balls.

To establish the neural representations for regime-shift estimation, we performed

three fMRI experiments ($n = 30$ subjects for each experiment, 90 subjects in total).

Experiment 1 was the main experiment, while Experiments 2 to 3 were control

experiments that ruled out two important confounds (Fig. 1E). The control

experiments were designed to clarify whether any effect of subjects' probability

estimates of a regime shift, P_t, in brain activity can be uniquely attributed to change

detection. Here we considered two major confounds that can contribute to the effect

of P_t. First, since subjects in Experiment 1 made judgments about the probability that

the current regime is the blue regime (which corresponded to probability of regime

change), the effect of P_t did not particularly have to do with change detection. To

address this issue, in Experiment 2 subjects made exactly the same judgments as in

Experiment 1 except that the environments were stationary (no transition from one

regime to another was possible), as in Edwards (1968) classic "bookbag-and-poker

chip" studies. Subjects in both experiments had to estimate the probability that the

current regime is the blue regime, but this estimation corresponded to the estimates of

regime change only in Experiment 1. Therefore, activity that correlated with

probability estimates in Experiment 1 but not in Experiment 2 can be uniquely

attributed to representing regime-shift judgments. Second, the effect of P_t can be due

to motor preparation and/or execution, as subjects in Experiment 1 entered two-digit

numbers with button presses to indicate their probability estimates. To address this

issue, in Experiment 3 subjects performed a task where they were presented with two-

digit numbers and were instructed to enter the numbers with button presses. By

comparing the fMRI results of these experiments, we were therefore able to establish

the neural representations that can be uniquely attributed to the probability estimates

of regime-shift.

Figure 1. The regime-shift detection task. **A.** Trial sequence. In each trial, the subjects saw a

sequence of red and/or blue signals and were told that these signals were drawn from one of

the two regimes, a Red regime and a Blue regime. Both regimes were described as urns

containing red and blue balls. The Red regime contained more red balls, while the Blue

regime contained more blue balls. Each trial always started at the Red regime but could shift

to the Blue regime in any of the 10 periods according to some transition probability (q). At

the beginning of a trial, information about transition probability (shown as "switch"

probability in the illustration) and signal diagnosticity (shown as "color ratio") were revealed

to the subjects. In this example, the transition probability is 0.1 and signal diagnosticity is 1.5.

See main text for more detailed descriptions. **B.** Manipulation of the system parameters, i.e.,

transition probability (q) and signal diagnosticity (d). We independently manipulated the q (3

levels) and d (3 levels), resulting in a 3×3 factorial design. **C.** An example of a particular

combination of the system parameters from the 3×3 design. Here the system that produces

the signals has a $q = 0.01$ transition probability and $d = 1.5$ signal diagnosticity. Signals

were sequentially presented to subjects. After each new signal appeared (a period), subjects

provided a probability estimate (P_t) of a regime shift. **D.** Two example trials sequences. The

example on the left shows the sequence of 10 periods of blue and red signals where $d = 1.5$

and $q = 0.01$. In this example, the regime was never shifted. The example on the right shows

the sequence of periods where $d = 9$ and $q = .1$. In this example, the regime was shifted

from the Red to the Blue regime in Period 3 such that the signals shown starting at this period

were drawn from the Blue regime. **E.** We performed three fMRI experiments (30 subjects in

each experiment) to investigate the neural basis of regime-shift judgments. Experiment 1 was

the main experiment looking at regime shift—which corresponds to $P(Change)$ in the Venn

diagram—while Experiments 2 and 3 were the control experiments that ruled out additional

confounds. In both Experiments 1 and 2, the subjects had to estimate the probability that

signals came from the blue regime. But unlike Experiment 1, in Experiment 2, which

corresponds to $P(Blue)$, no regime shift was possible. In Experiment 3, the subjects were

simply asked to enter a number with a button-press setup identical to Experiments 1 and 2.

Therefore, Experiment 3 (Motor) allowed to rule out motor confounds.

Behavioral evidence for over- and underreactions to change

Our analyses used subjects' probability estimates of a regime shift, P_t, for each

period, $t = 1, \dots, 10$. We found that subjects were in general responsive to the system

parameters, with higher P_t when the transition probability was larger. We also found

that subjects tended to give more extreme P_t under high signal diagnosticity than

low diagnosticity (Fig. S1 in Supplementary Information, *SI*). In addition, we used a

measure of belief revision, $\Delta P_t = P_t - P_{t-1}$. In Fig. 2A we show examples of P_t

and ΔP_t from a subject. On the left, the subject was in a stable environment (small

transition probability, $q = 0.01$) and faced two regimes that were very similar to each other (low signal diagnosticity, $d = 1.5$). The red and blue signals (10 periods) were what the subject encountered during a trial. On the right, the subject was in an unstable environment ($q = 0.1$) and faced two regimes that were very different ($d = 9$).

To examine over- and underreactions to change, we compared subjects' belief revision, $\Delta P_t = P_t - P_{t-1}, t = 2, \dots, 10$, with belief revision predicted by the Bayesian model, $\Delta P_t^B = P_t^B - P_{t-1}^B$ (see Fig. 2A for illustrations). ΔP_t and ΔP_t^B respectively capture how much subjects and a normative Bayesian change probability estimates in response to a new signal. When $\Delta P_t > \Delta P_t^B$, it indicates larger belief revision than the normative Bayesian, i.e., an overreaction. By contrast, $\Delta P_t < \Delta P_t^B$ indicates smaller belief revision, i.e., an underreaction. We therefore use $IO = \Delta P_t - \Delta P_t^B$ as an Index of Overreaction (IO). We found that subjects tended to overreact to change ($IO > 0$) when they received noisy signals (i.e., low signal diagnosticity, $d = 1.5$) and when the environment was stable (small transition probability, $q = 0.01$). By contrast, underreaction ($IO < 0$) was most commonly observed when they were in unstable environments (large transition probability, $q = 0.1$) and with clear signals (i.e., high signal diagnosticity, $d = 9$) (Fig. 2B). These patterns of over- and underreactions were consistent with findings in (Massey & Wu, 2005) and the system-neglect hypothesis, which posits a tendency to respond primarily to the signals and secondarily to the system that generates the signals (Massey & Wu, 2005; Seifert et al., 2023). According to the system-neglect hypothesis, responding secondarily to the system is synonymous with a lack of sensitivity to the system parameters, which leads to underreactions in unstable environments with precise signals, and overreactions in stable environments with noisy signals.

Figure 2. Behavioral results for Experiment 1. **A.** Illustrations of Over- and underreactions. Left column: stable environment ($q = 0.01$) with noisy signals ($d = 1.5$) and the 10 periods of red and blue signals a subject encountered. Right column: unstable environment ($q = 0.1$) with precise signals ($d = 9$). Top row: we plot a subject's actual probability estimates (P_t, solid line) and the normative Bayesian posterior probability (P_t^B, dashed line). Bottom row: belief revision shown by the subject ($\Delta P_t = P_t, -P_{t-1}$, solid line) and the Bayesian belief revision (ΔP_t^B, dashed line). The orange bars represent $\Delta P_t - \Delta P_t^B$, which we define as the Index of Overreaction (IO; vertical axis in orange on the right). **B.** Over- and underreactions to change. The mean IO (across all 30 subjects) is plotted as a function of

transition probability and signal diagnosticity. Subjects overreacted to change if $IO > 0$ and underreacted if $IO < 0$. Error bars represent ± 1 standard error of the mean. **C.** Parameter estimates of the system-neglect model. Left graph: Weighting parameter (α) for transition probability. Right graph: Weighting parameter (β) for signal diagnosticity. Dashed lines indicate parameter values equal to 1, which is required for Bayesian updating. **D-F.** Sensitivity to transition probability and signal diagnosticity are independent. **D.** Correlation between α and β estimates at different levels of transition probability (q_1 to q_3) and signal diagnosticity (d_1 to d_3). All pairwise Pearson correlation coefficients (indicated by the values on the table that were also color coded) were not significantly different from 0 ($p > .05$). **E.** Pearson correlation coefficients of α estimates between different levels of transition probability. All pairwise correlations were significantly different from 0 ($p < .05$). **F.** Pearson correlation coefficients of β estimates between different levels of signal diagnosticity. All pairwise correlations were significantly different from 0 ($p < .05$).

Following Massey and Wu (2005), we quantitatively model these belief revisions using the system-neglect model (see *System-neglect model* in *Methods*). The model is a parameterized version of the normative Bayesian model that allows for distortion of the system parameters via weighting parameters for transition probability (α) and signal diagnosticity (β). In short, α reflects distortion of transition probability, with $\alpha \times q$ in the system-neglect model capturing a decision maker's *effective* transition probability (q). For example, if $\alpha = 4$ when $q = 0.01$, the decision maker effectively treats a 0.01 transition probability as if it were 0.04. By contrast, β captures the extent to which the decision maker overweighs or underweighs signal diagnosticity (d^β) when faced with a signal. For example, if $\beta = 2$ when $d = 1.5$, subjects would treat a blue signal by updating the odds ratio for change by 1.5^2, or 2.25 rather than 1.5.

In the system-neglect model, we estimated the weighting parameters separately for each level of transition probability and signal diagnosticity, i.e., $\alpha_i \times q_i$ and $d_j^{\beta_j}$, where $\alpha_1, \alpha_2,$ and α_3 correspond respectively to transition probabilities of 0.01, 0.05, and 0.1, and $\beta_1, \beta_2,$ and β_3 correspond respectively to signal diagnosticity of 1.5, 3, and 9. In contrast to the Bayesian model which implies $\alpha_i = \beta_j = 1$ for all i, j, the system-neglect model requires that that $\alpha_i > \alpha_{i+1}$ and $\beta_j >$

β_{j+1} because it would effectively capture a lack of sensitivity to the system

parameters.

We fit the model to P_t for each subject separately and found parameter estimates

consistent with system neglect (Fig. 2C). The mean estimates of α were 3.69, 1.04

and 0.65 respectively for $q = 0.01, 0.05$ and 0.10. The parameters indicated that, on

average, when $q = 0.01$, subjects treated as if it were 0.0369. By contrast, when $q =$

0.10, the subjects treated it as if it were 0.065. Thus, a factor of 10 in actual transition

probability (0.01 vs. 0.1) was reduced to a factor of less than 2 (0.0369 vs. 0.065) in

effective transition probability. For signal diagnosticity, the mean parameter estimates

of β were 1.69, 0.77 and 0.57 for $d = 1.5,\ 3$, and 9, respectively. Thus, subjects

updated their beliefs $1.5^{1.69} = 1.98$ when $d_1 = 1.5$ and $9^{0.57} = 3.50$ when $d_3 =$

9. Normatively, the change in odds ratio between the two conditions should have been

$d_3/d_1 = 6$ but, consistent with system neglect, was considerably smaller,

$3.50/1.98 = 1.76$. Together, large parameter estimates ($\alpha > 1, \beta > 1$) at low signal

diagnosticity (noisy signals) and low transition probability (stable environments)

capture overreactions to changes, while small parameter estimates ($\alpha < 1, \beta < 1$) at

large signal diagnosticity (precise signals) and large transition probability (unstable

environments) reflect underreactions to change. These results replicate the findings by

Massey and Wu (2005), with the pattern of over- and underreactions as predicted by

the system neglect hypothesis. Critically, the degree of system neglect can be

captured by the negative trend of the parameter estimate as a function of the system

parameter levels (Fig. 2C): the steeper the slope, the larger the system neglect. We

found a similar pattern on β in Experiment 2 (one of the control experiments) where

environments were stationary (no transition probability) and signal diagnosticity was

manipulated (Fig. S2 in *SI*) (Benjamin, 2019; Tversky et al., 1990).

We performed a parameter recovery analysis to examine whether the fitting

procedure gave reasonable parameter estimates (Wilson & Collins, 2019). First, we

simulated each subject's probability estimation data based on the system-neglect

model by using that subject's parameter estimates. Second, we fitted the system-

neglect model to the simulated data. Third, we computed the correlation across

subjects between the estimated parameters and the parameter values we used to

simulate data. Fourth, we repeated the above steps by adding independent white noise

to the simulated data. Across different levels of noise, we found good parameter

recovery (Pearson's r for transition probability $r \geq 0.9533$ across different noise

levels, Pearson's r for signal diagnosticity $r \geq 0.9515$ across different noise

levels) (Figs. S3 and S4 in *SI*). In addition, we found that the empirical results (Fig.

2C) can be reproduced by the system-neglect model (Figs. S5 and S6 in *SI*). That is,

we used each subject's parameter estimates to compute the period-wise probability

estimates according to the system-neglect model and used these probability estimates

to compute and plot index of overreaction (*IO*). The patterns of *IO* based on the

system-neglect model (Fig. S6 in *SI*) were very similar to those based on subjects'

actual data (Fig. 2B).

We next examined whether the way subjects respond to different system

parameters is similar. It is possible, for example, that subjects who showed stronger

(or weaker) distortion of transition probability (captured by α parameter) also

showed stronger (or weaker) distortion of signal diagnosticity (captured by β

parameter). There was no significant correlation between α and β parameters (Fig.

2D). However, we did find within-parameter correlation: subjects who had a higher

α_i for a transition probability level i also tended to have a higher $\alpha_{i'}$ for a second

transition probability level i' (Fig. 2E), with the same pattern also holding for signal

diagnosticity (Fig. 2F). Together, these results suggested that the way an individual

decision maker responds to information about the probability of change in the

environment (transition probability) has little to do with how she or he responds to

information about the similarity between different regimes (signal diagnosticity). But

individuals are consistent in responding to a particular system parameter (transition

probability or signal diagnosticity) across different levels of the parameter.

fMRI results

We focus our fMRI analyses on addressing three questions. First, what are the brain

regions that correlated with subjects' probability estimates of change and belief

revision? Second, what are the neural representations for the computational variables

contributing to these probability estimates? Third, how might neural responses in the

identified brain areas be associated with under- and overreactions to change?

Ventromedial prefrontal cortex and ventral striatum represent probability

estimates and belief revision on regime shift

Our first analysis is aimed at identifying brain regions that represented our subjects'

regime-shift estimation. To address this question, we used two behavioral measures,

namely the period-by-period probability estimates of regime shift, P_t, and the change

in P_t between successive periods, ΔP_t. P_t can be regarded as the subjects' posterior

probability estimates of regime shift, whereas ΔP_t captures the change in belief

(belief revision) about regime shift in the presence of a new signal (see Fig. 3A for an

example on P_t and ΔP_t).

Figure 3. Neural representations for the updating of beliefs about regime shift. **A.** An

example. Updating is captured by the difference in probability estimates between two

adjacent periods (ΔP_t). The Blue bars reflect the period probability estimates (P_t), while

yellow bars depict ΔP_t. **B.** Whole-brain results on the main experiment (Experiment 1)

showing significant brain regions that correlate with the updating of beliefs about change

(ΔP_t). The clusters in orange represent activity correlated with ΔP_t. The blue clusters

represent activity correlated with the regime-shift probability estimates (P_t). The magenta

clusters represent the overlap between the P_t and ΔP_t clusters. **C-D.** Comparison between

experiments. To rule out visual and motor confounds on the P_t results described in **B**, we

compared the P_t contrast between the main experiment (Experiment 1) and two control

experiments (Experiments 2 and 3). **C.** Whole-brain results on the effect of P_t between Experiments 1 and 2 (Experiment 1 – Experiment 2 on the negative P_t contrast). **D.** Whole-brain results on the effect of P_t between Experiments 1 and 3 (Experiment 1 – Experiment 3 on the negative P_t contrast).

For P_t, we found that the ventromedial prefrontal cortex (vmPFC) and ventral striatum correlated with this behavioral measure of subjects' belief about change. In addition, many other brain regions, including the motor cortex, central opercular cortex, insula, occipital cortex, and the cerebellum also significantly correlated with P_t. (Fig. 3B; see GLM-1 in *Methods,* and Tables S1 to S3 in *SI*, respectively, for significant clusters of activation using Gaussian random field theory, permutation test on threshold-free-cluster-enhancement statistic, and permutations test on cluster-extent statistic).

For ΔP_t, we also found that the vmPFC and ventral striatum were associated with regime shift belief revision (Fig. 3; see GLM-1 in *Methods* and Tables S1 to S3 in *SI* for significant clusters of activation). While many brain regions correlated with regime-shift probability estimates (P_t), only the vmPFC and ventral striatum also correlated with belief revisions, ΔP_t. We examined the robustness of P_t representations in these two regions in several follow-up analyses. First, we implemented a GLM (GLM-2 in *Methods*) that, in addition to P_t, included various task-related variables contributing to P_t as regressors (Fig. S7 in *SI*). Specifically, to account for the fact that the probability of regime change increased over time, we included the *intertemporal prior* as a regressor in GLM-2. The intertemporal prior is the natural logarithm of the odds in favor of regime shift in the *t*-th

period, $\ln\left(\frac{1-(1-q)^t}{(1-q)^t}\right)$, where q is transition probability and $t = 1, ... ,10$ is the period

(see Eq. 1 in *Methods*). It describes normatively how the prior probability of change increased over time regardless of the signals (blue and red balls) the subjects saw during a trial. Including it along with P_t would clarify whether any effect of P_t can otherwise be attributed to the intertemporal prior. Second, we implemented a GLM that replaced P_t with the log odds of $P_t, \ln\left(P_t/(1 - P_t)\right)$ (Fig. S8 in *SI*). Third, we implemented a GLM that examined P_t separately on periods when change-consistent (blue balls) and change-inconsistent (red balls) signals appeared (Fig. S9 in *SI*). Each

of these analyses showed the same pattern of correlations between P_t and activation

in vmPFC and ventral striatum, further establishing the robustness of the P_t findings.

Brain regions shown to correlate with regime-shift probability estimates, P_t,

could be driven by motor response because larger estimates predominantly involved

right-hand finger presses (see *Methods* for details). To rule out motor confounds, we

conducted two control experiments (Experiments 2 and 3) and performed two

analyses. First, we examined the neural correlates of probability estimates (P_t in

GLM-1) in the control experiments (Experiments 2 and 3). Second, we compared the

effect of P_t (GLM-1) between the main experiment (Experiment 1) and the control

experiments. In the first analysis, we found that in both control experiments, vmPFC

and ventral striatum did not significantly correlate with probability estimates P_t at

the whole-brain level (in Experiment 2, no significant clusters of activation at the

whole-brain level; see Table S4 for Experiment 3 in *SI*). In the second analysis, we

found that for both vmPFC and ventral striatum, the regression coefficient of P_t was

significantly different between Experiment 1 and Experiment 2 (Fig. 3C) and between

Experiment 1 and Experiment 3 (Fig. 3D; also see Tables S5 and S6 in *SI*). In a

separate, independent (leave-one-subject-out, LOSO) ROI analysis on vmPFC and

ventral striatum, we also found the same results (Fig. S10 in *SI*). Finally, we note that

in GLM-1, we implemented an "action-handedness" regressor to directly address the

motor-confound issue, that higher probability estimates preferentially involved right-

handed responses for entering higher digits. The action-handedness regressor was

parametric, coding -1 if both finger presses involved the left hand (e.g., a subject

pressed "23" as her probability estimate when seeing a signal), 0 if using one left

finger and one right finger (e.g., "75"), and 1 if both finger presses involved the right

hand (e.g., "90"). Taken together, these results ruled out motor confounds and

suggested that vmPFC and ventral striatum represent subjects' probability estimates

of change (regime shifts) and belief revision.

A frontoparietal network represents key variables for estimating regime shifts

Our second analysis is aimed at identifying brain regions that represented key

variables contributing to regime-shift estimation. Guided by our theoretical

framework and computational models, we focused on two variables, the interaction

between signals and signal diagnosticity and intertemporal prior probability of

change, and set up a new GLM (GLM-2 in *Methods*) to examine these effects.

Our theoretical framework makes two fundamental predictions. First, a signal

should be weighted differently depending on signal diagnosticity, i.e., a blue ball is

stronger evidence for change in a highly diagnostic environment (e.g., $d = 9$) than a

system in which the red and blue regimes are very similar (e.g., $d = 1.5$). To capture

the interaction between signals and signal diagnosticity, we code a blue signal as 1

and a red signal as -1 and multiply the signal code ($s =$1 or -1) by the natural

logarithm of signal diagnosticity, $\ln(d)$ (two examples are shown in Fig. 4A). We

term this interaction, $s \times \ln(d)$, the strength of evidence in favor of change or

strength of change evidence for short. The Bayesian model, as described in *Methods*,

critically depends on d^s, computing posterior odds by multiplying prior odds by the

likelihood ratio. Thus, the log posterior odds were calculated from both the prior odds

and $s \times \ln(d)$. At the whole-brain level, we found that a frontoparietal network

including the dorsal medial prefrontal cortex (dmPFC), lateral prefrontal cortex

(bilateral inferior frontal gyrus, IFG), and the posterior parietal cortex (bilateral

intraparietal sulcus, IPS) represented $s \times \ln(d)$ (Fig. 4A). These brain regions

overlap with what is commonly referred to as the frontoparietal control network

(Buckner et al., 2013; Seeley et al., 2007; Yeo et al., 2011). Among them, dmPFC sits

in the vicinity of dorsomedial frontal cortex (DMFC) shown to represent change

probability and uncertainty about change in reinforcement learning (McGuire et al.,

2014).

The second prediction our theoretical framework offers concerns the prior

probability of a regime shift over time. Specifically, the Bayesian model predicts that

the prior probability should increase over time (see two examples in Fig. 4B), with the

intertemporal prior, in log odds terms, defined as the natural logarithm of the odds in

favor of regime shift in the *t*-th period, $\ln\left(\frac{1-(1-q)^t}{(1-q)^t}\right)$, where q is transition

probability and $t = 1, \ldots, 10$ is the period (see Eq. 1 in *Methods*). With independent

(leave-one-subject-out, LOSO) ROI analysis, we examined whether brain regions in

the frontoparietal network (shown to represent strength of change evidence)

correlated with intertemporal prior and found that all brain regions, with the exception

of dmPFC, in the frontoparietal network correlated with the intertemporal prior (Fig.

4B; dmPFC: $t(29) = -1.69, p = 0.10$; left IFG: $t(29) = 2.20, p = 0.04$; right

IFG: $t(29) = -2.64, p = 0.01;$ left IPS: $t(29) = -2.35, p = 0.03;$ right IPS: $t(29) = -2.07, p = 0.05$). By contrast, brain regions that represented the intertemporal prior, which we found to be in the right fusiform cortex in the occipitotemporal regions, did not correlate with the strength of change evidence (Fig. S11 in *SI*).

Finally, we emphasize that these effects—the strength of change evidence and intertemporal prior—cannot be otherwise attributed to probability estimates (P_t) or belief revision (ΔP_t) because both P_t and ΔP_t were included in GLM-2 where these effects were examined. Taken together, these results suggest that the frontoparietal network is critically involved in representing the two key variables for estimating regime shifts, strength of change evidence and intertemporal prior. See Supplementary Tables S7 to S9, respectively, for information about significant clusters of activation using Gaussian random field theory, permutation test on threshold-free-cluster-enhancement (TFCE) statistic, and permutation test on cluster-extent statistic.

Figure 4. A frontoparietal network represented key variables for regime-shift estimation. **A.** Variable 1: strength of evidence in favor of/against regime shifts (strength of change evidence), as measured by the interaction between signal diagnosticity and signal. Left: two

examples of the interaction between signal diagnosticity (d) and sensory signal (s), where a blue signal is coded as 1 and a red signal is coded as -1. The x-axis represents the time periods, from the first to the last period, in a trial. The y-axis represents the interaction, $\ln(d) \times s$. Right: whole-brain results showing brain regions in a frontoparietal network that significantly correlated with $\ln(d) \times s$. **B.** Variable 2: intertemporal prior probability of change. Two examples of intertemporal prior are shown on the left graphs. To examine the effect of the intertemporal prior, we performed independent region-of-interest analysis (leave-one-subject-out, LOSO) on the brain regions identified to represent strength of evidence (5A). Due to the LOSO procedure, individual subjects' ROIs (a cluster of contiguous voxels) would be slightly different from one another. To visualize such differences, we used the red color to indicate voxels shared by all individual subjects' ROIs, and orange to indicate voxels by at least one subject's ROI. The ROI analysis examined the regression coefficients (mean PE) of intertemporal prior. The * symbol indicates $p < .05$, ** indicates $p < .01$. dmPFC: dorsomedial prefrontal cortex; lIPS: left intraparietal sulcus; rIPS: right intraparietal sulcus; lIFG: left inferior frontal gyrus; rIFG: right inferior frontal gyrus.

Under- and overreactions are associated with selectivity and sensitivity of neural responses to system parameters

The system-neglect hypothesis posits that under- and overreactions arise from a lack of sensitivity to the system parameters. We can measure individual subjects' sensitivity to system parameters using behavioral data (subjects' probability estimates). Meanwhile, we can also measure sensitivity using neural data. In the following analysis, we examined whether there is a match between the behavioral and neural measures of sensitivity to the system parameters. This would allow us to examine, through the system-neglect framework, whether sensitivity in neural responses to the system parameters are associated with under- and overreactions to change.

We focused on the vmPFC-striatum network and frontoparietal network, as they were shown to be involved in regime-shift detection (Figs. 3 and 4). We examined whether these brain networks show selective preference for a particular system parameter, which we refer to as *parameter selectivity*. We also asked whether parameter selectivity is signal-dependent, i.e., different for signals consistent with change (blue signals) or inconsistent with change (red signals).

We started by defining a behavioral measure of sensitivity to the system

parameter. To visualize this measure, we consider two extreme decision makers, a

Bayesian and someone who reacts to signals identically across all systems, which we

term *complete neglect*. In Fig. 5A (left graph), we use signal diagnosticity (d) to

illustrate the pattern of these two decision makers. The vertical axis is $\beta \ln(d)$ and

the horizontal axis is the signal-diagnosticity level (d), where β is the weighting

parameter on signal diagnosticity in the system-neglect model. A Bayesian (open

circles) does not overweight or underweight d, and thus $\beta = 1$. We can then define

the *Bayesian slope* by regressing $\beta_i \ln(d_i)$ against $\ln(d_i)$. In this formulation, the

Bayesian slope is 1 and it reflects the sensitivity of a Bayesian decision maker to

signal diagnosticity. On the other hand, a complete-neglect decision maker is

unresponsive to signal diagnosticity, i.e., $\beta_1 \ln(d_1) = \beta_2 \ln(d_2) = \beta_3 \ln(d_3)$. Hence,

the *complete-neglect slope* should be 0. These two slopes, the Bayesian slope and the

complete-neglect slope, provide the boundaries for system neglect. For each subject,

we computed $\beta_i \ln(d_i)$ at each d_i level, where β_i is the estimate for diagnosticity

d_i fitted to the system-neglect model (see β_i in Fig. 2B). We then estimated each

subject's *behavioral slope* (to distinguish it from the *neural slope* reported later) and

use it as a behavioral measure of sensitivity to signal diagnosticity.

C

D

Signal diagnosticity (*d*)

Transition probability (*q*)

Red · Blue · Red · Blue

vmPFC (x=2)

ventral striatum (y=6, z=0)

Behavioral slope vs. Neural slope | Red / Neural slope | Blue

vmPFC:
- Signal diagnosticity Red: *r*=0.28, *p*=0.133
- Signal diagnosticity Blue: *r*=0.26, *p*=0.171
- Transition probability Red: *r*=-0.38, *p*=0.043*
- Transition probability Blue: *r*=-0.37, *p*=0.047*

ventral striatum:
- Signal diagnosticity Red: *r*=0.33, *p*=0.077
- Signal diagnosticity Blue: *r*=0.27, *p*=0.150
- Transition probability Red: *r*=-0.26, *p*=0.175
- Transition probability Blue: *r*=0.03, *p*=0.870

Figure 5. Estimating and comparing neural measures of sensitivity to system parameters with behavioral measures of sensitivity. **A.** Behavioral measures of sensitivity to system parameters. For each system parameter, we plot the subjectively weighted system parameter against the system parameter level (top row: signal diagnosticity; bottom row: transition probability). For each subject and each system parameter, we estimated the slope (how the subjectively weighted system parameter changes as a function of the system parameter level) and used it as a behavioral measure of sensitivity to the system parameter (behavioral slope). We also show a Bayesian (no system neglect) decision maker's slope (dark green) and the slope of a decision maker who completely neglects the system parameter (in light green; the slope would be 0). A subject with stronger neglect would have a behavioral slope closer to complete neglect. **B.** Comparison of behavioral and neural measures of sensitivity to the system parameters. To estimate neural sensitivity, for each subject and each system parameter, we regressed neural activity of a ROI against the parameter level and used the slope estimate as a neural measure of sensitivity to that system parameter (neural slope). We also estimated the neural slope separately for blue-signal periods (when the subject saw a blue signal) and red-signal periods. We computed the Pearson correlation coefficient (*r*) between the behavioral slope and the neural slope and used it to statistically test whether there is a match between the behavioral and neural slopes. **C.** The frontoparietal network selectively represented individuals' sensitivity to signal diagnosticity (left two columns), but not transition probability (right two columns). Further, neural sensitivity to signal diagnosticity (neural slope) correlated with behavioral sensitivity (behavioral slope) only when a signal in favor of potential change (blue) appeared: all the regions except the right IPS showed

statistically significant match between the behavioral and neural slopes. By contrast, sensitivity to transition probability was not represented in the frontoparietal network. **D.** The vmPFC selectively represented individuals' sensitivity to transition probability ($r = -0.38, p = 0.043$ for red signals; $r = -0.37, p = 0.047$ for blue signals), but not signal diagnosticity ($r = 0.28, p = 0.13$ for red signals; $r = 0.26, p = 0.17$ for blue signals). The ventral striatum did not show selectivity to either transition probability or signal diagnosticity. Error bars represent ± 1 standard error of the mean.

For each subject, we estimated two behavioral slopes, one for $d,$ the signal diagnosticity (top row in Fig. 5A), and the other for $q,$ the transition probability (bottom row in Fig. 5A). The right graphs in Fig. 5A shows the behavioral slope for each of the 30 subjects (top: signal diagnosticity; bottom: transition probability). For signal diagnosticity, 28 out of 30 subjects' behavioral slope were within the boundaries. For transition probability, 27 out of 30 subjects' behavioral slope were within the boundaries (between complete neglect and Bayesian). One subject's (subject 6) behavioral slope for q (transition probability) was approximately 2 and clearly outside the boundaries. This subject's data were excluded for further analysis on q (the right two columns in Fig. 5CD).

We found that, across subjects, system neglect was unique from either Bayesian or complete neglect. Subjects' sensitivity to transition probability, as captured by the behavioral slope in Fig. 5A, deviated significantly from the Bayesian slope (comparing subjects' slope with 1, $t(29) = -10.8, p < .01$, two-tailed) and from complete neglect slope (comparing subjects' slope with 0, $t(29) = 4.8, p < .01$, two-tailed). For signal dignosticity, subjects' sensitivity to signal diagnosticity was also significantly different from both Bayesian ($t(29) = -12.5, p < .01$, two-tailed) and complete neglect ($t(29) = 6.1, p < .01$, two-tailed).

However, subjects were closer to complete neglect than to the Bayesian. We tested this by examining whether the behavioral slope, γ, was significantly greater or smaller than 0.5, the midpoint between complete neglect (slope of 0) and Bayesian (slope of 1). $\gamma - .5 > 0$ indicates that subjects' behavior was in closer alignment with Bayesian. By contrast, $\gamma - .5 < 0$ implies behavior closer to complete neglect. We found that, for both transition probability and signal diagnosticity, the behavioral slope was closer to complete neglect than to Bayesian (transition probability: $t(29) = -2.97, p < .01$; signal diagnosticity: $t(29) = -3.23, p < .01$, two-

tailed). Together, these results suggested that, while subjects did respond to the

system parameters in regime-shift estimation in the correct direction predicted by the

Bayesian model, their sensitivity to the system parameters was closer to complete

neglect than to normative Bayesian.

For the neural data, we defined a neural measure of sensitivity to the system

parameters by estimating how neural responses change as a function of those

parameters. Using the signal diagnosticity parameter as an example, for each subject

and each ROI separately, we regressed average brain activity at each diagnosticity

level against $\ln(d)$. The slope estimate, termed the *neural slope*, from the linear

regression gave us a neural measure of sensitivity to signal diagnosticity. To

investigate whether the neural sensitivity was signal-dependent, i.e., neural sensitivity

in response to the change-consistent signals (blue signals) was different from the

change-inconsistent signals (red signals), we separately estimated the neural slope in

response to blue and red signals.

After obtaining both the behavioral and neural measures of sensitivity to the

system parameters, we then computed the Pearson correlation coefficient between

them. We found that the vmPFC-striatum network and frontoparietal network showed

clear dissociations in how they corresponded with the system parameters. First, the

frontoparietal network represented individual subjects' sensitivity to signal

diagnosticity (left two columns in Fig. 5C), but not transition probability (right two

columns in Fig. 5C). Notably, patterns of parameter selectivity were remarkably

consistent across brain regions in the frontoparietal network: when change-consistent

signals (blue signals) appeared, the neural measure of sensitivity from all brain

regions in the frontoparietal network except the right IPS significantly correlated with

the behavioral measure of sensitivity (second column from the left in Fig. 5B;

dmPFC: $r = 0.48, p = 0.007$; lIFG: $r = 0.5, p = 0.009$; rIFG: $r = 0.4, p = 0.027$;

lIPS: $r = 0.58, p = 0.001$; rIPS: $r = 0.32, p = 0.082$). By contrast, when change-

inconsistent signals (red signals) appeared, all regions within the frontoparietal

network did not significantly correlate with the behavioral measure of sensitivity (first

column from the left in Fig 5B; dmPFC: $r = 0.32, p = 0.083$; lIFG: $r = 0.04, p = 0.848$; rIFG: $r = 0.19, p = 0.312$; lIPS: $r = 0.05, p = 0.787$; rIPS: $r = -0.02, p = 0.914$). We further tested, for each brain region, whether the difference in

correlation was significant using both parametric and nonparametric tests (see

Parametric and nonparametric tests for difference in correlation coefficients in

Methods). The results were identical. In the parametric test, we used the Fisher z transformation to transform the correlation coefficients to the z statistic. Since these correlation coefficients were not independent, we compared them using the test developed in Meng et al. (1992) (see *Methods*). We found that among the five ROIs in the frontoparietal network, two of them, namely the left IFG and left IPS, the difference in correlation was significant (one-tailed z test; left IFG: $z = 1.8908, p = 0.0293$; left IPS: $z = 2.2584, p = 0.0049$). For the remaining three ROIs, the difference in correlation was not significant (dmPFC: $z = 0.9522, p = 0.1705$; right IFG: $z = 0.9860, p = 0.1621$; right IPS: $z = 1.4833, p = 0.0690$). We chose one-tailed test because we already know the correlation under change-consistent signals was significantly greater than 0. In the nonparametric test, we performed nonparametric bootstrapping to test for the difference in correlation. We referred to the correlation between neural and behavioral sensitivity at change-consistent (blue) signals as r_{blue}, and that at change-inconsistent (red) signals as r_{red}. Consistent with the parametric tests, we also found that the difference in correlation was significant in left IFG and left IPS (left IFG: $r_{blue} - r_{red} = 0.46, p = 0.0496$; left IPS: $r_{blue} - r_{red} = 0.5306, p = 0.0041$), but was not significant in dmPFC, right IFG, and right IPS (dmPFC: $r_{blue} - r_{red} = 0.1634, p = 0.1919$; right IFG: $r_{blue} - r_{red} = 0.2123, p = 0.1681$; right IPS: $r_{blue} - r_{red} = 0.3434, p = 0.0631$). In summary, we found that neural sensitivity to signal diagnosticity measured at change-consistent signals significantly correlated with individual subjects' behavioral sensitivity to signal diagnosticity. By contrast, neural sensitivity to signal diagnosticity measured at change-inconsistent signals did not significantly correlate with behavioral sensitivity. The difference in correlation, however, was statistically significant in some (left IPS and left IFG) but not all brain regions within the frontoparietal network.

Second, in contrast to the frontoparietal network, vmPFC in the vmPFC-striatum network showed the opposite pattern of parameter selectivity: vmPFC selectively represented individual subjects' sensitivity to transition probability (right two columns in Fig. 5D), but not to signal diagnosticity (left two columns in Fig. 5D). Selectivity in vmPFC was not signal-dependent: regardless of change-consistent (blue) or change-inconsistent (red) signals, neural sensitivity to transition probability in vmPFC represented individual subjects' sensitivity to transition probability ($r = -0.38, p = 0.043$ for change-inconsistent signals; $r = -0.37, p = 0.047$ for change-consistent signals). By contrast, the ventral striatum did not show selectivity

to either the transition probability or signal diagnosticity (transition probability: $r = $

$-0.26, p = 0.175$ for change-inconsistent signals; $r = 0.03, p = 0.870$ for change-

consistent signals; signal diagnosticity: $r = 0.33, p = 0.077$ for change-inconsistent

signals; $r = 0.27, p = 0.150$ for change-consistent signals). In summary, these

results suggest that vmPFC selectively represented individuals' sensitivity to

transition probability, whereas the frontoparietal network selectively represented

individuals' sensitivity to signal diagnosticity.

Incorporating signal dependency into system-neglect model led to better models

for regime-shift detection

The neural findings on signal dependency (Fig. 5) point to the possibility that

participants might respond to the system parameters differently when facing change-

consistent and change-inconsistent signals. This led us to ask whether building signal

dependency into the system-neglect model would be a better model choice for

subjects' behavioral data (probability estimates of regime shift) than the original

system-neglect model. To examine this question, we built and fit three new versions

of the system-neglect (SN) model (see Table S12 in *SI* for model-fitting summary)

and compared them with the original model (SN-original) (see Table S13 in *SI* for

summary of statistical tests for model comparison). In the signal-dependent β

system-neglect model (SN-SigDep-β model), we estimated the β parameters

separately at change-consistent and change-inconsistent signals. As a result, in this

model there were 6 β parameters—three for change-consistent signals to model each

of the three levels of signal diagnosticity and three for change-inconsistent signals—

and three α parameters that modeled each of the three levels of transition probability

without distinguishing between change-consistent and change-inconsistent signals. In

the signal-dependent α system-neglect model (SN-SigDep-α model), we estimated

the α parameters separately at change-consistent and change-inconsistent signals. As

a result, in this model there were 6 α parameters (three for change-consistent signals

and three for change-inconsistent signals) and three β parameters. In the signal-

dependent α and β system-neglect model (SN-SigDep-$\alpha\beta$ model), we estimated

both α and β parameters separately at change-consistent and change-inconsistent

signals (12 total parameters). Compared with SN-original, we found that SN-SigDep-

β, SN-SigDep-α, and SN-SigDep- $\alpha\beta$ qualitatively described subjects' behavioral

data (belief revision, ΔP_t) better (Fig. 6B-E). Further, we found that estimating α

separately at change-consistent and change-inconsistent signals (SN-SigDep-α, Fig. 6D) model improved model fits than estimating β separately (SN-SigDep-β, Fig. 6C), suggesting that subjects responded to transition probability differently when facing change-consistent and change-inconsistent signals more than to signal diagnosticity. Model comparison using Akaike Information Criterion (AIC) revealed that SN-SigDep-$\alpha\beta$ is the best model, followed by SN-SigDep-α, SN-SigDep-β, and SN-original (Fig. 6F). Together, these results suggest that participants showed system-neglect to both transition probability and signal diagnosticity and that they responded to these system parameters differently when facing change-consistent and change-inconsistent signals. In summary, signal dependency in response to system parameters is a new behavioral finding not reported in the original Massey and Wu (2005) study and is largely inspired by the neural sensitivity findings in the current study.

Figure 6. Model comparison. **A-E.** Modeling results from five competing models. For each

model, we plot subjects' belief revision (ΔP_t) and the model-estimated ΔP_t. Light-colored

dots and dashed lines respectively represent the model-estimated ΔP_t at the individual and

group levels. Dark-colored dots and solid lines indicate individual subjects' ΔP_t and group-

averaged behavioral data respectively. Blue indicates data and model estimates at change-consistent signals; Red indicates data and model estimates at change-inconsistent signals. **A.** Bayesian model. **B.** Original system-neglect model (SN-original). **C.** Signal-dependent β system-neglect model (SN-SigDep-β). **D.** Signal-dependent α system-neglect model (SN-SigDep-α). **E.** Signal-dependent α and β system-neglect model (SN-SigDep-$\alpha\beta$). **F.** Model comparison based on the Akaike Information Criterion (AIC). Lower AIC values indicate better model fits. The bars indicate group mean AIC (averaged across all subjects), while the black dots indicate individual subjects' AIC values. Error bars represent ± 1 standard error of the mean. The * symbol indicates $p < .05$, ** indicates $p < .01$ (paired t-test; see Table S13 in *SI* for summary of statistical tests).

DISCUSSION

In this study, we investigated how humans detect changes in the environments and the neural mechanisms that contribute to how we might under- and overreact in our judgments. Combining a novel behavioral paradigm with computational modeling and fMRI, we discovered that sensitivity to environmental parameters that directly impact change detection is a key mechanism for under- and overreactions. This mechanism is implemented by distinct brain networks in the frontal and parietal cortices and in accordance with the computational roles they played in change detection. By introducing the framework in system neglect and providing evidence for its neural implementations, this study offered both theoretical and empirical insights into how systematic judgment biases arise in dynamic environments.

Regime shifts—the transition from one state of the world to another—are present in many daily situations, from the stock market (a change from the bull to the bear market) to the state of a pandemic. Detecting regime shifts can be challenging for at least two reasons. First, the signals we receive from the environments are often noisy. A signal in favor of potential change, for example a drop in pandemic cases, can either inform a true shift in regime or simply reflect noisy fluctuations. Second, the signals we receive reflect the volatility of the environment: while some environments are more prone to changes, others are not. To capture these two key features in regime-shift detection, we designed an fMRI task based on Massey and Wu (2005) where subjects made probability judgments about regime shifts and where we manipulated the signal diagnosticity and transition probability. Signal diagnosticity captures the level of noise inherent in the signals, while transition probability reflects the volatility of the environment. Replicating Massey and Wu (2005), we found that overreactions to regime shifts take place when participants received noisy signals (low signal diagnosticity) but when the environments were stable (low transition probability). By contrast, when the signals are more precise but the environments were unstable, participants tended to underreact to changes. These results suggest system neglect—people respond primarily to signals and secondarily to the system that generates the signals (Massey & Wu, 2005).

At the neurobiological level, we found that regime-shift detection is jointly implemented by two networks, the vmPFC-striatum network and a frontoparietal network. The vmPFC-striatum network represented subjects' probability estimates of

change and the revision of probability estimates in the presence of new signals (belief revision). By contrast, the frontoparietal network represented strength of change evidence and intertemporal prior probability of change—two key variables contributing to probability estimation. Guided by the system-neglect framework, we found that under- and overreactions to change are closely associated with the sensitivity of these networks in response to the system parameters—transition probability and signal diagnosticity—that impact regime changes. In particular, the vmPFC represented individual subjects' sensitivity to transition probability, whereas the frontoparietal network represented sensitivity to signal diagnosticity. Together, these findings suggest that selectivity and sensitivity of neural responses to system parameters are key mechanisms that give rise to under- and overreactions.

Our work is closely related to the reversal-learning paradigm—the standard paradigm in neuroscience and psychology to study change detection (Fellows & Farah, 2003; Izquierdo et al., 2017; O'Doherty et al., 2001; Schoenbaum et al., 2000; Walton et al., 2010). In a typical reversal-learning task, human or animal subjects choose between two options that differ in the reward magnitude or probability of receiving a reward. Through reward feedback the participants gradually learn the reward contingencies associated with the options and have to update knowledge about reward contingencies when contingencies are switched in order to maximize rewards. While a switch in reward contingencies can be regarded as a kind of regime shift, there are three major differences between the reversal-learning paradigm and our regime-shift task. The first difference is about learning. In the reversal-learning paradigm, the subjects must learn both the reward contingencies and the switches through experience. By contrast, in the regime-shift task, the subjects were explicitly informed about the makeup of different regimes and the transition probability. Therefore, participants do not need to learn about the different regimes and the transition probability through experience. The second difference is the kind of behavioral data collected. In our task, we asked the subjects to estimate the probability of change, whereas in the reversal-learning task, the subjects indicate their choice preferences. The third difference is on reward contingency. In the reversal-learning task, a change is specifically about change in reward contingencies, which is not the case in our task.

We believe that these major differences in task design led to three key insights into change detection from our study. The first insight is on over- and underreactions

to change. At the behavioral level, we were able to identify situations that led to over-

and underreactions. At the theoretical level, we were able to provide a systematic

account for these over- and underreactions in the system-neglect hypothesis. Finally,

at the neurobiological level, we were able to quantify the degree to which individual

subjects neglected the system parameters and use these behavioral measures to

unravel the neural mechanisms that give rise to over- and underreactions to change.

The second insight is on the brain networks associated with change detection. In

particular, we were able to clarify whether the neural systems involved in change

detection in the reversal-learning tasks are contingent on whether rewards are

involved. Since the reversal-learning tasks are about learning the reward

contingencies and the change in reward contingencies, it would be challenging to

infer whether the neural implementations of change detection are dissociable from

reward processing. Indeed, brain regions shown to be involved in the reversal-

learning tasks, the OFC, mPFC, striatum, and amygdala, were also found to be highly

involved in reward-related learning and value-based decision making. In the current

study, unlike in reversal-learning paradigms, regimes were not defined by rewards

(e.g., high reward-probability regime vs. low reward-probability regime in reversal

learning paradigm). Therefore, estimating the probability of regime shifts in our task

did not require considerations for change in reward contingencies. Our findings that

vmPFC and ventral striatum represent probability estimates of change and belief

revision therefore suggest that these brain regions might be part of a common

pathway for change detection in general where changes in the state of the world do

not have to be about changes in reward contingencies.

The third insight has to do with the impact of learning on change detection.

Under the reversal-learning paradigm, it has been challenging to infer whether there

exists a unique change-detection mechanism that is dissociable from reinforcement

learning mechanisms. The way to make such inference is through theory, such as

implementing a prior for state changes (Bartolo & Averbeck, 2020; Costa et al.,

2015). Unlike the reversal-learning task, participants in our task did not have to learn

about the different regimes through experience. Without the confound of

reinforcement learning, our results help clarify the roles of change detection on choice

behavior by suggesting that independent of learning, there exists a specialized

change-detection mechanism in the brain that impacts decision making. This

mechanism involves the participation of the vmPFC-striatum network and the

frontoparietal network, which partially overlap with the brain regions involved in reversal learning. However, it remains to be seen how learning interacts with change detection. Future investigations can address this question by combining the key features of both the reversal-learning paradigm and regime-shift paradigm.

Outside of the reversal-learning paradigm, previous fMRI studies that investigated learning and belief updating in dynamic environments where change takes place regularly had identified brain regions that represent perceived likelihood of change inferred from participants' choice behavior. Payzan-LeNestour et al. (2013) identified that the posterior cingulate cortex, postcentral gyrus, middle temporal gyrus, hippocampus, and insula correlated with subjects' perceived likelihood of change in a multi-arm bandit task. McGuire et al. (2014) found that subjective change-point probability was represented in a large posterior cluster including occipital, inferior temporal, and posterior parietal cortex. In addition, activity in dorsomedial frontal cortex, posterior cingulate cortex, superior frontal cortex, and anterior insula also positively correlated with change probability. Interestingly, both McGuire et al. (2014) and our results found that the ventral striatum negatively correlated with probability estimates of change. This result suggested that ventral striatum represents probability estimates of change irrespective of whether the task was based on a learning-based paradigm (McGuire et al., 2014) or non-learning paradigm where information about task-related variables was explicitly revealed to the participants. Further, both McGuire et al. and our results found the involvement of the dorsomedial prefrontal cortex (dmPFC; or dorsomedial frontal cortex in McGuire et al.) in change detection. Our results further suggest that dmPFC is specialized in weighing the strength of change evidence and represents individual subjects' sensitivity to signal diagnosticity, both of which played important roles in contributing to the over- and underreactions to change.

How might our results relate to value-based decision making? In previous studies, vmPFC had been implicated to dynamically track financial risks that carry potential monetary gains or losses. To understand dynamic computations of risk, Schonberg et al. (2012) used a Balloon Analog Risk Task (BART) where subjects decide whether to inflate a simulated balloon through successive pumps for the potential to win larger gains or incur larger losses (if balloon explodes), or to cash out before the balloon explodes. They found that vmPFC activity decreased as subjects pumped and expanded the balloon, suggesting its involvement in estimating the risk

of potential losses. Since the explosion of balloon can be regarded as a change in the state of the balloon, as the balloon expands the possibility of such a change in state (regime shift) also increases. In this view, the vmPFC result from Schonberg et al. (2012) was consistent with our finding in that vmPFC negatively correlated with probability estimates of regime shift. Together, these results add to the existing literature by suggesting that vmPFC is involved in estimating and updating the state of the world in dynamic environments where changes take place regularly.

Related to OFC function in decision making and reinforcement learning, Wilson et al. (2014) proposed that OFC is involved in inferring the current state of the environment. For example, medial OFC had been shown to represent probability distribution on possible states of the environment (Chan et al., 2016), the current task state (Schuck et al., 2016) and uncertainty or entropy associated with the state of the environment (Muller et al., 2019). In the context of regime-shift detection, regimes can be regarded as states of the environment and therefore a change in regime indicates a change in the state of the environment. Muller et al. (2019) found that in dynamic environments where changes in the state of the environment happen regularly, medial OFC represented the level of uncertainty in the current state of the environment. Our finding that vmPFC represented individual participants' probability estimates of regime shifts suggest that vmPFC and/or OFC are involved in inferring the current state of the environment through estimating whether the state has changed. Our finding that vmPFC represented individual participants' sensitivity to transition probability further suggest that vmPFC and/or OFC contribute to individual participants' biases in state inference (over- and underreactions to change) in how these brain areas respond to the volatility of the environment.

Our results are also closely related to the literature on the neural mechanisms for evidence accumulation in decision making (Gold & Shadlen, 2007; Mante et al., 2013; Philiastides et al., 2010; Roitman & Shadlen, 2002; Yates et al., 2017). In our task, evaluating the signals (red or blue balls) and in particular, the strength of change evidence associated with the signals is central to performing the task. Normatively, such evaluation should depend on the signal diagnosticity. In a highly diagnostic environment, seeing a red ball should signal a strong possibility of being in the red regime, while seeing a blue ball should signal otherwise. By contrast, in a low diagnostic environment, a red (resp. blue) ball is not strongly indicative of a red (resp.

blue) regime. Hence, the evaluation of signals should reflect the interaction between the signals and the diagnosticity of the signals.

We found that this key computation was implemented in a frontoparietal network commonly referred to as the frontoparietal control network (Buckner et al., 2013; Dosenbach et al., 2007; Seeley et al., 2007; Vincent et al., 2008; Yeo et al., 2011). This network was proposed to support adaptive control functions, including initiating control and providing flexibility to adjust the level of control through feedback (Dosenbach et al., 2007). The IPS and dlPFC, part of this network, have also been found to play a major role in the top-down control of attention (Corbetta & Shulman, 2002; Woldorff et al., 2004). In perceptual decision making, the IPS and dlPFC were also shown to represent the accumulation of sensory evidence that leads to the formation of a perceptual decision (Heekeren et al., 2004; Heekeren et al., 2006). Our findings—that activity in this network does not reflect just the sensory signals (red or blue balls) but how these signals should be interpreted through the lens of their diagnosticity—highlights the involvement of the frontoparietal control network in computing the strength of evidence through combining information about signals and knowledge about the precision of those signals.

For the frontoparietal network, we identified its involvement in our task through finding that its activity correlated with strength of change evidence (Fig. 4) and individual subjects' sensitivity to signal diagnosticity (Fig. 5). Conceptually, these two findings reflect how individuals interpret the signals (signals consistent or inconsistent with change) in light of signal diagnosticity. This is because (1) strength of change evidence is defined as signals ($+1$ for signal consistent with change, and -1 for signal inconsistent with change) multiplied by signal diagnosticity and (2) sensitivity to signal diagnosticity reflects how individuals subjectively evaluate signal diagnosticity. At the theoretical level, these two findings can be interpreted through our computational framework in that both the strength of change evidence and sensitivity to signal diagnosticity contribute to estimating the likelihood of change (Equations 1 and 2 in *Methods*).

Our result on the intraparietal sulcus (IPS) being part of the brain network that represents diagnosticity-weighted sensory signals are consistent with previous studies showing that IPS is involved in accumulating sensory evidence over time (Gold & Shadlen, 2007). There are three interesting aspects of our data that add to the current

literature on evidence accumulation. First, IPS representations for sensory evidence need not be in the space of actions. Unlike previous studies showing that IPS represents sensory evidence for potential motor actions, we found that IPS represents the strength of evidence in favor of or against regime shifts. This result points to a more general role of the IPS in estimating the strength of sensory evidence. In fact, our result suggests that it depends on the task goal, which in the current study is to estimate whether a change has taken place. Second, although evidence accumulation is important and necessary for a wide array of cognitive functions, it is not a central requirement for the regime-shift task. Bayesian updating—the framework in which our system-neglect model was built upon—only requires the computation of strength of change evidence associated with the signal shown in the latest period. By showing that IPS representing this quantity, this suggests that IPS is involved in evaluating the latest piece of evidence necessary for belief updating.

In the current study, the central opercular cortex—in addition to the vmPFC—is another brain region that represented the probability estimates of change. Like the vmPFC, activity in this region negatively correlated with the probability estimates of change. This finding is associated with previous findings on change detection using the oddball paradigm. Using the oddball paradigm, it was found that the central opercular cortex is involved in the detection of change, showing stronger activation in blocks containing only the standard stimulus than blocks containing both the standard and deviant stimulus (Hedge et al., 2015) and correlating with ERP P3 signals at the trial-level that reflected differences between standard and deviant stimuli (Warbrick et al., 2009). There are two implications here. First, our findings suggest that the central opercular cortex is not only involved in the detection of change—as revealed by the oddball tasks—but also is involved in the estimation of change where there is uncertainty regarding whether the state of the world had changed. Second, the central opercular cortex may be part of a common pathway for the detection of change across very different tasks such as the oddball paradigm and our regime-shift detection task.

In the current study, our psychometric-neurometric analysis focused on comparing behavioral sensitivity with neural sensitivity to the system parameters (transition probability and signal diagnosticity). We measured sensitivity by estimating the slope of behavioral data (behavioral slope) and neural data (neural slope) in response to the system parameters. Previous studies had adopted a similar approach (Ting et al., 2015a; Vilares et al., 2012; Yang & Wu, 2020). For example,

Vilares et al. (2012) found that sensitivity to prior information (uncertainty in prior distribution) in the orbitofrontal cortex (OFC) and putamen correlated with behavioral measure of sensitivity to the prior. In the current study, transition probability acts as prior in the system-neglect framework (Eq. 2 in *Methods*) and we found that ventromedial prefrontal cortex represents subjects' sensitivity to transition probability. Together, these results suggest that OFC (with vmPFC being part of OFC, see Wallis, 2011) is involved in the subjective evaluation of prior information in both static (Vilares et al., 2012) and dynamic environments (current study). In addition, distinct from vmPFC in representing sensitivity to transition probability or prior, we found through the behavioral-neural slope comparison that the frontoparietal network represents how sensitive individual decision makers are to the diagnosticity of signals in revealing the true state (regime) of the environment. Interestingly, such sensitivity to signal diagnosticity was only present in the frontoparietal network when participants encountered change-consistent signals. However, while most brain areas within this network responded in this fashion, only the left IPS and left IFG showed a significant difference in coding individual participants' sensitivity to signal diagnosticity between change-consistent and change-inconsistent signals. Unlike the left IPS and left IFG, we observed in dmPFC a marginally significant correlation with behavioral sensitivity at change-inconsistent signals as well. Together, these results indicate that while different brain areas in the frontoparietal network responded similarly to change-consistent signals, there was a greater degree of heterogeneity in responding to change-inconsistent signals.

    In summary, our results suggest that an important mechanism for under- and overreactions to change has to do with neural sensitivity to system parameters that impact regime shifts. Importantly, different system parameters appear to recruit distinct brain networks according to their unique computational specializations. Given that under- and overreactions underly a wide array of human judgments, our findings indicate that network-level computational specificity and parameter selectivity are two key building blocks that give rise to human judgment biases.

MATERIALS AND METHODS

The data and analysis code are available at https://osf.io/xh7dy/.

We performed three fMRI experiments (90 subjects in total, 30 subjects for each experiment). Experiment 1 was the main experiment where we investigated the neurocomputational substrates for regime shifts. Experiments 2 and 3 were control experiments. Experiment 2 was designed to rule out brain activity that correlated with probability estimates but was not specifically about regime shifts. Experiment 3 attempted to rule out brain activity that correlated with entering numbers through button presses. In the main text, we only presented the results from Experiment 1. The procedure and results of Experiments 2 and 3 were presented in Supplementary Materials.

Subjects. All subjects gave informed written consent to participate in the study. All subjects were right-handed. The study procedures were approved by the National Yang Ming Chiao Tung University Institutional Review Board. Ninety subjects participated in this study:

- Experiment 1: $n = 30$ subjects; 15 males; mean age: 22.9 years; age range: 20 to 29 yrs.
- Experiment 2: $n = 30$ subjects; 15 males; mean age: 23.3 years; age range, 20 to 30 years.
- Experiment 3: $n = 30$ subjects; 15 males; mean age: 23.7 years; age range: 20 to 34 years.

Subjects were paid 300 New Taiwan dollar (TWD, 1 USD = 30 TWD) for participating the behavioral session and 500 TWD for the fMRI session. Subjects received additional monetary bonus based on his or her performance on probability estimation in Experiments 1 and 2 (Experiment 1: an average of 209 and 212 TWD for the behavioral and fMRI session respectively; Experiment 2: an average of 223 and 206 TWD for the behavioral and fMRI session respectively). In Experiment 3, subjects received the bonus based on their performance for entering the correct number (an average of 243 TWD for the fMRI session).

Procedure

Overview. Experiment 1 consisted of two sessions—a behavioral session followed by an fMRI session—that took place on two consecutive days. Subjects performed the same task in both sessions. The goals of having the behavioral session were to familiarize subjects with the task and to have enough trials—along with the fMRI session—to reliably estimate the parameters of the system-neglect model. Details of Experiments 2 and 3 can be found in the Supplement.

Regime-shift detection task. In this task, the environments the subjects are described as *regimes*. There were two possible regimes and at any point in time the regime can shift from one to another. Subjects judged whether the regime had shifted based on three sources of information: transition probability, signal diagnosticity, and signals. Prior to each trial, the subjects were given information about the transition probability and signal diagnosticity of the upcoming trial. The transition probability described how likely it is that the regime can shift from one to another. The signal diagnosticity described how different the two regimes are. The two regimes—red and blue—were represented by a red urn and a blue urn respectively, with each urn consisting of red and blue balls. The red urn had more red balls and the blue urn had more blue balls. The two urns always shared the same ratio of the dominant ball color to the dominated ball color. That is, the ratio of the red balls to the blue balls in the red urn was same as the ratio of the blue balls to the red balls in the blue urn. Signal diagnosticity was quantitatively defined by this ratio. For example, when the ratio was 9:1, signal diagnosticity (d) was 9. Under this definition, larger signal diagnosticity indicated that the two regimes were more different from each other.

After information about the transition probability and signal diagnosticity were revealed, 10 sensory signals—red or ball balls shown on the computer screen—were sequentially presented to the subjects. Each signal was sampled with replacement from one of the two urns. We also referred to the sequential presentation of these signals as *periods*. By design, each trial always started at the red regime, but the regime can shift from the red to the blue regime at any point in time during a trial, including prior to the sampling of the first signal. In addition, also by design, the regime can shift only once in a trial (i.e., the blue regime was an absorbing or trapping state). That is, once the regime changed from the red to the blue regime, signals would be sampled from the blue urn until the end of the trial. Subjects provided an

estimate of the probability that current regime was the blue regime at each period when a new signal was presented. Since the regime can only shift from the red to the blue, this probability estimate was equivalent to the probability estimate of regime shift.

Stimuli. For each subject and each trial separately, we generated the stimuli, i.e., the sequence of red and blue balls according to the transition probability and signal diagnosticity of the trial. Before the start of each period, we first determined whether the regime would shift from the Red regime (the starting regime) to the Blue regime by sampling from the transition probability. There were two possible outcomes: 1 indicates a change in regime, whereas 0 indicates no change. If the outcome were 1, we would sample from the Blue regime for that period and for all the remaining period(s). If the outcome were 0, we would sample from the Red regime for that period and repeat the same process described above for the next period.

Manipulations of transition probability and signal diagnosticity. We implemented a 3×3 factorial design where there were three possible transition probabilities, denoted as q where $q = [0.01, 0.5, 0.1]$ and three levels of signal diagnosticity, denoted as d where $d = [1.5, 3, 9]$. The transition probability—the probability that the regime shifts from the red to the blue regime—was used to simulate the stability of the environment. Larger transition probabilities indicate less stable environments. The signal diagnosticity was the ratio of dominant balls to the dominated balls in the urns. For example, when $d = 9$, it indicates that the red balls are 9 times more than the blue balls in the red urn and that the blue balls are 9 times more than the red balls in the blue urn. In this case, the two regimes were highly different and hence the signals shown to the subjects should be highly diagnostic of the regime the subjects were currently in.

Session 1: behavioral session

A pre-test block followed by the main task (8 blocks of trials) were implemented in the behavioral session. We implemented the pre-test block to make sure the subjects understood the experimental instructions and to train them to become familiar with

the task, especially the time limit in entering probability estimates (see Trial sequence below for details on the pre-test block). A 3 (transition probability) by 3 (signal diagnosticity) factorial design was implemented, resulting in a total of 9 experimental conditions. Each block consisted of 9 trials (one for each condition, randomized in order) and each trial consisted of 10 periods or equivalently, 10 sequentially presented signals. After the pre-test block, subjects performed the main task that consisted of 8 blocks of trials. The behavioral session took approximately 70 minutes to complete.

Trial sequence. At the beginning of each trial, the subjects were shown information about the transition probability and signal diagnosticity (3s) (Fig. 1A). This was followed by the sequential presentation of 10 signals. At each period, a signal—a red or blue dot sampled from the current regime was shown. Subjects' task was to estimate the probability that the current regime was the blue regime within 4s. The subjects indicated the probability estimate through two button presses. During the experiment, the subjects placed his or her 10 fingers on a 10-button keypad. Each finger was associated with a single button that corresponded to a unique integer value from 0 to 9, with the left little finger for 1, left ring finger for 2, left middle finger for 3, left index finger for 4, left thumb for 5, right thumb for 6, right index finger for 7, right middle finger for 8, right ring finger for 9, and right little finger for 0. To enter the probability estimate, the subjects had to first press the number corresponding to the tens and second the number corresponding to the ones. For example, if the probability estimate was 95%, the subjects first had to press the "9" button and second the "5" button. Once the subjects entered the probability estimate, she or he was not allowed to change it. After providing the probability estimate, they were given a brief feedback (0.5s) on the number they just entered. If they failed to indicate probability estimate within the time limit (4s), a "too slow" message would be shown. At the end of each trial, the subjects received feedback (2s) on the amount of monetary bonus earned in the trial (2s) and information about whether the regime shifted during the trial. If the regime was shifted, the signal that was drawn right after the shift took place would be highlighted in white. This was followed by a variable inter-trial interval (ITI, 1s to 5s in steps of 1s drawn from a discrete uniform distribution).

We implemented a pre-test block at the beginning of the behavioral so as to train the subjects to enter the probability estimate within the time limit (4s). In the pre-test block, we started with a lenient time limit and gradually decreased it. The time limit for the first three trials was 20s, 10s from trial number 4 to 6, and 4s from trial

number 7 to 9. Recall that in each trial, the subjects encountered 10 sequentially presented signals (red and blue balls) and had to provide probability estimate at each period. The subjects therefore entered 30 probability estimates under each of these three time limits—from 20s, 10s, to 4s. After the pre-test block, all the subjects were able to enter the probability estimates within 4s.

Session 2: fMRI session

The fMRI session consisted of 3 blocks (9 trials in each block, with each trial consisting of 10 sequentially presented signals). The task was identical to the behavioral session (except with varying inter-stimulus interval for the purpose of fMRI analysis) and took approximately 60 minutes to complete. The subjects indicated the probability estimate through two button presses. During the experiment, the subjects placed his or her 10 fingers on a 10-button keypad. Each finger was associated with a single button that corresponded to a unique integer value from 0 to 9 (starting from the left pinkie for 1, left ring finger for 2, left middle finger for 3, and etc., to right ring finger for 9, and finally right pinkie for 0). The trial sequence was identical to the behavioral session with a few exceptions. First, each new signal was presented for 4s regardless of when subjects make a response. Second, we added a variable inter-stimulus interval (ISI, 1s to 5s in steps of 1 drawn from a uniform distribution) between two successive signals. We also made the range of ITI to be slightly larger (1s to 7s drawn from a discrete uniform distribution between in steps of 1s) than the behavioral session. The design of variable ISIs and ITIs was to allow better dissociations between events, i.e., between different sensory signals presented during a trial, and between trials for fMRI analysis.

Monetary bonus

To incentivize subjects to perform well in this experiment, they received monetary bonus based on his or her probability estimates. The bonus rule was designed so that the subjects who gave more accurate estimates would earn more bonus. The bonus structure used a quadratic payoff:

$$Bonus(t) = \$30 \times (0.1 - 0.2 \times (P_t - B_t)^2),$$

where P_t is the probability estimate that the current regime was blue at the t-th

period in a trial and B_t is the regime at t ($B_t = 1$ for the blue regime and $B_t = 0$ for

the red regime). For each probability estimate, the bonus therefore ranged from

winning \$3 to losing \$3 TWD. For example, if the subject gave a 99% probability

estimate that the current regime was blue and the current regime was indeed the blue

regime, she would receive a bonus close to 3 TWD. By contrast, if the subjects gave a

1% probability estimate that the current regime was blue but the current regime was

the red regime, she would receive a penalty close to 3 TWD. With 10 probability

estimates given in each trial, the subjects can therefore receive a bonus up to 30 TWD

or a penalty up to 30 TWD in a trial. The subjects did not receive feedback on bonus

after each probability estimate. Instead, at the end of each trial, the subjects received

information about the total amount won or lost in that trial. The final total bonus was

realized by randomly selecting 10 trials at the end of the experiment.

Computational models for regime shift

We examined two computational models for regime shift: the Bayesian model and the

system-neglect model (Massey & Wu, 2005). The Bayesian model was parameter-

free and was used to compare with subjects' probability estimates of regime shift. The

system-neglect model is a quasi-Bayesian model or parameterized version of the

Bayesian model that was fit to the subjects' probability-estimate data. The parameter

estimates of the system-neglect model were further used in the fMRI analysis so as to

identify neural representations for over- and underreactions to change.

Bayesian model

Here we describe the Bayesian posterior odds of shifting to the blue regime given the

period history H_t (Edwards, 1968; Massey & Wu, 2005):

$$\frac{P_t^B}{1-P_t^B} = \frac{\Pr(B_t|H_t)}{\Pr(R|H_t)} = \frac{1-(1-q)^t}{(1-q)^t}\sum_{j=1}^{t}\frac{q(1-q)^{j-1}}{1-(1-q)^t}d^{t+1-j-2\sum_{k=j}^{t}r_k}, \qquad (1)$$

$P_t^B = \Pr(B_t|H_t)$ is the posterior probability that the regime has shifted to the blue

regime at the t-th period, H_t denotes the sequence of history from r_1 to r_t. Here

r_t donates the t-th period, where $r_t = 1$ when the signal at t is red, and $r_t = 0$

when the signal at t is blue. The transition probability and signal diagnosticity are

denoted by q and d respectively.

The posterior odds are the product of the prior odds and the likelihood ratio. The

prior odds, $\frac{1-(1-q)^t}{(1-q)^t}$, indicates that given the transition probability q, the probability

that the regime has shifted at time t, $1-(1-q)^t$, relative to the probability of no

change in regime, $(1-q)^t$. The likelihood ratio, $\sum_{j=1}^{t} \frac{q(1-q)^{j-1}}{1-(1-q)^t} d^{t+1-j-2\sum_{k=j}^{t} r_k}$,

indicates the probability of observing the history of signals H_t given that the regime

has shifted relative to the probability given that the regime has not shifted. This

requires considering all the possibilities on the timing of shift, i.e., the likelihood ratio

that the regime was shifted at $t=j$, $d^{t+1-j-2\sum_{k=j}^{t} r_k}$, weighted by its odds $\frac{q(1-q)^{j-1}}{1-(1-q)^t}$.

Since these possibilities are disjoint events, the likelihood ratio that the regime has

shifted is simply the weighted sum of all the likelihood ratios associated with these

disjoint possibilities.

Index of overreaction to change

To quantify under- and overreactions to change, we derived an Index of Overreaction

(IO). In short, this index reflects the degree to which the subjects overreacted or

underreacted to change such that an index value greater than 0 indicates overreaction

and an index value smaller than 0 suggests underreaction. To compute IO, we

compared the change in probability estimates between two adjacent periods

($\Delta P_t = P_t - P_{t-1}$) with the normative change in probability estimates according to

the Bayesian model $(\Delta P_t^B = P_t^B - P_{t-1}^B)$. Here we use P_t to denote the subject's

probability estimate at the t-th period and P_t^B to denote the Bayesian probability

estimate. We computed IO ($IO = \Delta P_t - \Delta P_t^B$) separately for each subject and for

each condition or combination of transition probability and signal diagnosticity.

Overreaction is defined when the actual change in probability estimates is greater than

the normative change in probability estimates ($\Delta P_t > \Delta P_t^B$). By contrast,

underreaction is defined when the actual change is smaller than the normative change

in probability estimate ($\Delta P_t < \Delta P_t^B$). When $\Delta P_t^e = \Delta P_t^B$, there neither overreaction

nor underreaction to change.

System-neglect model

Following Massey and Wu (2005), we fit the system-neglect model—a quasi-
Bayesian model—to the subjects' probability estimates. In this model, we aimed to
capture system-neglect—that people respond primarily to the signals and secondarily
to the system generating the signals. Responding secondarily to the system indicates
that people with system-neglect would be less sensitive to the changes in the system
parameters compared with the normative Bayesian. This is captured by adding a
weighting parameter to each system parameter level. Hence, two weighting
parameters, α and β, were added to Eq. (1) to transition probability and signal
diagnosticity respectively such that

12
$$\frac{P_t}{1-P_t} = \frac{\Pr(B_t|H_t)}{\Pr(R|H_t)} = \frac{1-(1-\alpha q)^t}{(1-\alpha q)^t} \sum_{j=1}^{t} \frac{q(1-q)^{j-1}}{1-(1-q)^t} d^{\beta\left(t+1-j-2\sum_{k=j}^{t} r_k\right)} \tag{2}$$

13 where P_t is the probability estimate that regime has changed to the blue regime at
14 period t. We separately estimated α for each level of transition probability and β
15 for each level of signal diagnosticity. This was implemented by setting dummy
16 variables for each level of transition probability and signal diagnosticity in Eq. (2)

17
$$\alpha = \alpha_1 Q_1 + \alpha_2 Q_2 + \alpha_3 Q_3 \text{ and } \beta = \beta_1 D_1 + \beta_2 D_2 + \beta_3 D_3.$$

Where Q_i is the dummy variable for transition probability q_i and D_j is the dummy
variable for diagnosticity d_j. α_i and β_j therefore respectively reflect the sensitivity
to different levels of transition probability and signal diagnosticity. For each subject
separately, we performed nonlinear regression (using the *fitnlm* function in
MATLAB) to fit the model to the subject's probability estimates and estimated the
parameters of interest.

Parameter recovery analysis

To examine whether the fitting procedure gave reasonable parameter estimates of the
system-neglect model, we performed a parameter recovery analysis (Wilson &
Collins, 2019). The analysis proceeded in the following steps. First, we simulated
each subject's probability estimation data based on the system-neglect model by using
the parameter estimates obtained for that subject after fitting the system-neglect
model to the subject' probability estimation data. Second, we fit the system-neglect

model to the simulated data. Third, as a measure of parameter recovery, we computed

the correlation across subjects between the estimated parameters and the parameter

values we used to simulate data, where larger correlation indicates better recovery.

Fourth, we repeated the above steps by adding independent white noise to the

simulated data, from a Gaussian distribution with mean 0 and variance σ^2. We

implemented 5 levels of noise with $\sigma_{noise} = \{0.01, 0.05, 0.1, 0.2, 0.3\}$ and examined

the impact of noise on parameter recovery. These noise levels covered the range of

empirical noise levels estimated from the subjects. To estimate each subject's noise

level, we incorporated a noise parameter into the system-neglect model. We assumed

that probability estimates are noisy and modeled them with a Gaussian distribution

where the noise parameter (σ_{noise}) is the standard deviation. At each period, a

probability estimate of regime shift was computed according to the system-neglect

model where Θ is the set of parameters including parameters in the system-neglect

model and the noise parameter. The likelihood function, $L(\Theta)$, is the probability of

observing the subject's probability estimate at period t, p_t, given Θ, $L(\Theta) =$

$P(p_t | \Theta)$. Since we modeled the noisy probability estimates with a Gaussian

distribution, we can therefore express $L(\Theta)$ as $L(\Theta) \sim N(p_t; p_t^{SN}, \sigma_{noise})$ where p_t^{SN}

is the probability estimate predicted by the system-neglect (SN) model at period t. As

a reminder, we referred to a 'period' as the time when a new signal appeared during a

trial (for a given transition probability and signal diagnosticity). To find the maximum

likelihood estimates of the parameters, Θ_{MLE}, we summed over all periods the

negative natural logarithm of likelihood and used MATLAB's fmincon function to

find Θ_{MLE}. Across subjects, we found that the mean noise estimate was 0.1735 and

ranged from 0.1118 to 0.2704 (Fig. S3 in *SI*).

Impact of noise homoscedasticity on parameter estimation

The fitnlm function we used in MATLAB to estimate the parameters of the system-

neglect models assumes homoscedasticity—that noise is constant when fitting the

model. However, it is possible that this assumption was violated, that subjects' actual

noise was heteroscedastic, for example, having larger noise when probability

estimates are around 0.5 and smaller noise at the two extremes (0 and 1). To examine

this possibility, for each subject, we divided probability into 5 intervals ([0.0–0.2),

[0.2–0.4), [0.4–0.6), [0.6–0.8), and [0.8–1.0]) and computed the residual standard

deviation separately for each bin. Here the residual was the difference between

subjects' probability estimates (p_t) and the probability estimates derived from the

system-neglect model based on each subject's parameter estimates (p_t^{SN}),

$residual = p_t - p_t^{SN}$. We found that homoscedasticity was indeed violated—the

standard deviation of residuals was smallest when probability was 0.1 and increased

as a function of probability from 0.1 to 0.5. When probability was larger than 0.5, the

residual standard deviation was similar (Fig. S4A in *SI*). To see how this would affect

parameter estimation, we performed parameter recovery analysis assuming

heteroscedasticity. That is, we simulated subjects' probability estimates using the

empirically estimated, probability-dependent residual standard deviation as the

standard deviation of the Gaussian noise and using the simulated data to estimate

parameters in the system-neglect model. We found that we were able to recover the

parameters well (Fig. S4B in *SI*) and the result was similar to the parameter recovery

assuming homoscedastic noise (Fig. S3 in *SI*). This suggested that heteroscedastic

noise did not impact the accuracy of parameter estimation when homoscedasticity was

assumed.

fMRI data acquisition. All the imaging parameters, including the EPI sequence and

MPRAGE sequence, remained the same throughout the three experiments. For

Experiments 1 and 2, the subjects completed the task in a 3T Siemens MRI scanner

(MAGNETOM Trio) equipped with a 32-channel head array coil. Experiment 3 was

collected at a later date after the same scanner went through a major upgrade (from

Trio to the Prisma system) where a 64-channel head array coil was used. Each subject

completed three functional runs. Before each run, a localizer scan was implemented

for slice positioning. For each run, T2*-weighted functional images were collected

using an EPI sequence (TR=2000ms, TE=30ms, 33 oblique slices acquired in

ascending interleaved order, 3.4×3.4×3.4 mm isotropic voxel, 64×64 matrix in 220

mm field of view, flip angle 90°). To reduce signal loss in the ventromedial prefrontal

cortex and orbitofrontal cortex, the sagittal axis was tilted clockwise up to 30°. Each

run consisted of 9 trials and a total of 374 images. After the functional scans, T1-

weighted structural images were collected (MPRAGE sequence with TR=2530ms,

TE=3.03ms, flip angle=7°, 192 sagittal slices, 1×1×1 mm isotropic voxel, 224×256

matrix in a 256-mm field of view). For each subject, a field map image was also

acquired for the purpose of estimating and partially compensating for geometric

distortion of the EPI image so as to improve registration performance with the T1-weighted images.

fMRI preprocessing. The imaging data were preprocessed with FMRIB's software Library (FSL version 6.0). First, for motion correction, MCFLIRT was used to remove the effect of head motion during each run. Second, FUGUE (FMRIB's Utility for Geometric Unwarping of EPIs) was used to estimate and partially compensate for geometric distortion of the EPI images using field map images collected for the subject. Third, spatial smoothing was applied with a Gaussian kernel with FWHM=6mm. Fourth, a high-pass temporal filtering was applied using Gaussian-weighted least square straight-line fitting with $\sigma = 50s$. Fifth, registration was performed in a two-step procedure, with the field map used to improve the performance of registration. First, EPI images were registered to the high-resolution brain T1-weighted structural image (non-brain structures were removed via FSL's BET (Brain Extraction Tool). Second, the transformation matrix (12-parameter affine transformation) from the T1-weighted image to the Montreal Neurological Institute (MNI) template brain was estimated using FLIRT (FMRIB's Linear Image Registration Tool), followed by nonlinear registration using FNIRT (FMRIB's Non-linear Image Registration Tool) with a 10mm warp resolution. This two-step procedure allowed for transforming the EPI images to the standard MNI template brain.

General Linear Models of BOLD signals

All GLM analyses were carried out in the following steps (Beckmann et al., 2003). First, BOLD time series were pre-whitened with local autocorrelation correction. A first-level FEAT analysis was carried out for each run of each subject. Second, a second-level (subject-level) fixed-effect (FE) analysis was carried out for each subject that combined the first-level FEAT results from different runs using the summary statistics approach. Finally, a third-level (group-level) mixed-effect (ME) analysis using FSL's FLAME module (FMRIB's Local Analysis of Mixed Effects) was carried out across subjects by taking the FE results from the previous level and treating subjects as a random effect (Woolrich et al., 2004). All reported whole-brain results were corrected for multiple comparisons. We first identified clusters of

activation by defining a cluster-forming threshold of the z statistic ($z > 3.1$ or

equivalently $p < .001$) (Eklund et al., 2016; Woo et al., 2014). Then, a family-wise

error corrected p-value of each cluster based on its size was estimated using Gaussian

random field theory (Worsley et al., 1992). In addition, we performed nonparametric

permutation test using the randomize function in FSL (threshold-free cluster

enhancement or TFCE option; (Smith & Nichols, 2009) on all the contrasts reported.

GLM-1. This model was used for Fig 3. The model served two purposes. First, we

used it to examine neural representations for probability estimates (P_t) and belief

revision (ΔP_t) (Fig. 3B). Second, we used this model to compare results between the

main experiment (Experiment 1) and the control experiments (Experiments 2 and 3)

(Fig. 3CD). We implemented the following regressors. At the time of each signal

presentation, we implemented the following regressors: (R1) an indicator regressor

with length equal to the subject's RT, (R2) R1 multiplied by the subject's probability

estimate, P_t, that the signal came from the Blue regime, (R3) R1 multiplied by

difference in the subject's probability estimate between two successive periods,

ΔP_t, which captured the updating of belief about change, (R4) R1 multiplied by the

degree of certainty in probability estimate $|P_t - 0.5|$, (R5) R1 multiplied by the

period number (from 1 to 10). Both positive and negative contrasts of R2 (P_t) and

R3 (ΔP_t) were set up to identify activity that either positively or negatively correlate

with these regressors. At the end of each trial (after subjects saw all 10 signals), we

provided the monetary bonus the subject earned in that trial. We implemented an

indicator regressor (R6) and a parametric regressor for the subject's winning (R7). We

implemented an indicator regressor (length equal to 4s) for the no-response periods

(R8), which corresponded to the period(s) where the subject did not indicate the

probability estimate within the time limit (4s). Finally, to directly address the motor

confound issue, we implemented an action-handedness regressor, (R9), which was R1

multiplied by action-handedness. This regressor served to address the motor

confounds of probability estimates P_t, as higher probability estimates preferentially

involved right-handed responses for entering higher digits and lower estimates

involved left-handed responses. Therefore, at the time of each signal presentation, the

action-handedness regressor coded -1 if both finger presses to enter P_t involved left

hand, 0 if using one left finger and one right finger, and 1 if both finger presses

involved right hand. For Experiment 1, we also performed a GLM analysis that was

identical to GLM-1 except that it did not include the action-handedness regressor and the results on both P_t and ΔP_t were largely identical (Supplementary Table S10). Note that for both Experiment 1 (the main regime-shift experiment) and Experiment 2 (control experiment), P_t indicates the probability estimate that the signal came from the Blue regime. However, for Experiment 3 (control experiment) subjects were instructed to press a two-digit number shown on the screen at each period. Hence, the P_t regressor for Experiment 3 was effectively the instructed number, and the ΔP_t regressor was effectively the difference in instructed number between adjacent periods.

GLM-2. This model was implemented to examine the effects of component variables critically contributing to regime-shift probability estimation. In particular, we used this model to identify the effects on the strength of change evidence (based on R11 below) and the intertemporal prior (based on R9 below) (Fig. 4 in main text). The model was identical to GLM-1 from (R1) to (R8) with the addition of the following regressors: (R9) R1 multiplied by the intertemporal prior, defined as $\ln\left(\frac{1-(1-q)^t}{(1-q)^t}\right)$, where q is the transition probability and t is the period, (R10) R1 multiplied by the natural logarithm of signal diagnosticity, $\ln(d)$, (R11) R1 multiplied by the current signal (1 for blue ball, -1 for red ball), and (R12) the interaction $R9 \times R10$ (interaction between signal diagnosticity and signal). We note that since these component variables (R9 to R12) contributed to regime-shift probability estimation, they were correlated with P_t and ΔP_t. Having both with P_t and ΔP_t in the same model as these component variables would therefore introduce collinearity and reduce the reliability of regression coefficients. While we were aware of this issue, we also recognized that having these correlated regressors in the same model have the advantage of statistically stating that neural activity attributed to a particular variable cannot be otherwise attributed to another variable that correlated with it. We also performed another GLM analysis that was identical to GLM-2 except that it did not include P_t and ΔP_t in the model and the results on strength of change evidence and intertemporal prior shown in Fig. 4 were largely identical (Supplementary Table S11).

GLM-3. This model was the basis of results shown in Fig. 5. We set up two sets of regressors, one set for when the blue signal (signal for potential change) appeared and the other for when the red signal (signal for no change) appeared. For each set, at the time of signal presentation, we included 9 indicator regressors, one for every combination of 3 transition probability levels and 3 signal diagnosticity levels. Based on these 9 regressors, we set up a linear-increasing contrast separately for signal diagnosticity and transition probability. At the subject level (second level), the parameter estimate of these contrasts reflects individual subjects' sensitivity to signal diagnosticity and transition probability were extracted from a given ROI and was used to correlate with individual subjects' behavioral sensitivity to signal diagnosticity and transition probability. The ROIs used were vmPFC, ventral striatum, dmPFC, bilateral IFG, and bilateral IPS. These ROIs were identified by results from GLM 1 and 2 and were constructed in a statistically independent manner using leave-one-subject-out method. See Independent region-of-interest (ROI) analysis in the section below. Same as in the previous two GLMs, at the end of each trial when feedback on the current-trial monetary bonus was revealed, we implemented an indicator regressor and a parametric regressor for the monetary bonus. Finally, we implemented an indicator regressor (length equal to 4s) for no-response periods. These corresponded to the period(s) where the subject did not indicate the probability estimate within the time limit (4s).

Independent regions-of-interest (ROIs) analysis. We performed two kinds of independent ROI analysis—leave-one-subject-out (LOSO) method and functional/structural masks based on previous meta-analysis paper (Bartra et al. (2013) or existing structural atlases. Specifically, we used the functional mask in vmPFC from Bartra et al. (2013) and structural mask for the ventral striatum based on the Harvard-Oxford cortical and subcortical atlases in FSL. For LOSO, based on results from the whole-brain analysis, we created independent and unbiased ROIs using the leave-one-subject-out (LOSO) method (Litt et al., 2011; Ting et al., 2015b). The LOSO method was used to analyze probability estimates and representations for transition probability and signal diagnosticity described above. We performed the analysis for each subject separately in the following steps. First, we identified the significant cluster in a brain region (e.g., dmPFC) that correlated with a contrast of interest (e.g., probability estimates) using all other subjects' data. We referred to this

as the LOSO ROI. Second, we extracted the mean beta value (regression coefficient)

within the LOSO ROI from the subject and used it for further statistical analysis. Note

that the LOSO ROI tended to be different spatially between subjects. To give an idea

of these differences and the spatial distribution of these ROIs, in Figs. 4 and 5 where

the LOSO analysis was performed, we showed both the voxels that were part of the

LOSO ROI of all the subjects in one color and the voxels that were part of at least one

LOSO ROI in another color. Finally, we note that since the LOSO procedure

involves, for each subject separately, performing the leave-one-subject-out inference

at the group level and identifying the significant clusters of activation. Therefore, it is

possible that some subject(s), after the LOSO inference, did not have significant

cluster in some brain region. For the analysis shown in Fig. 5C, for the left IFG ROI,

we had 3 subjects that did not have a significant cluster in this brain region. All other

ROIs had data from all subjects.

Parametric and nonparametric tests for difference in correlation coefficients. We

implemented both parametric and nonparametric tests to examine whether the

difference in Pearson correlation coefficients was significant. We denote the

correlation coefficient between neural and behavioral sensitivity at change-consistent

(blue) signals as r_{blue}, and that at change-inconsistent (red) signals as r_{red}. In the

parametric test, we adopted the approach of Meng et al. (1992) to statistically

compare the two correlation coefficients. This approach specifically tests differences

between dependent correlation coefficients according to the following equation

24
$$z = (z_{r1} - z_{r2})\sqrt{\frac{N-3}{2(1-r_x)h}}$$

26 where N is the number of subjects, z_{ri} is the Fisher z-transformed value of r_i,

27 ($r_1 = r_{blue}$ and $r_2 = r_{red}$), and r_x is the correlation between the neural sensitivity at

28 change-consistent signals and change-inconsistent signals. The computation of h is

29 based on the following equations

31
$$h = \frac{1 - f\overline{r^2}}{1 - \overline{r^2}} = 1 + \frac{\overline{r^2}}{1 - \overline{r^2}}(1 - f)$$

$$f = \frac{1 - r_x}{2(1 - \overline{r^2})}, which\ must\ be \leq 1$$

where $\overline{r^2}$ is the mean of the r_i^2, $(r_1^2 + r_2^2)/2$, and f should be set to 1 if > 1. In

the nonparametric test, we performed nonparametric bootstrapping to test for the

difference in correlation (Efron & Tibshirani, 1994). That is, we resampled with

replacement the dataset (subject-wise) and used the resampled dataset to compute the

difference in correlation. We then repeated the above for 100,000 times so as to

estimate the distribution of the difference in correlation coefficients, tested for

significance and estimated p-value based on this distribution.

ACKNOWLEDGMENTS

This work was supported by the National Science and Technology Council (NSTC) in Taiwan (Grants 108-2410-H-010-012-MY3, 110-2410-H-A49A-504 -MY3 to S.-W.W.) and by the Brain Research Center, National Yang Ming Chiao Tung University from The Featured Areas Research Center Program within the framework of the Higher Education Sprout Project by the Ministry of Education (MOE) in Taiwan. We acknowledge magnetic resonance imaging support from National Yang Ming Chiao Tung University, Taiwan, which is in part supported by the Ministry of Education plan for the top University.

REFERENCES

Baker, M., & Wurgler, J. (2007). Investor sentiment in the stock market. *Journal of economic perspectives*, *21*(2), 129-151.

Barberis, N., Shleifer, A., & Vishny, R. (1998). A model of investor sentiment. *Journal of financial economics*, *49*(3), 307-343.

Bartolo, R., & Averbeck, B. B. (2020). Prefrontal Cortex Predicts State Switches during Reversal Learning. *Neuron*, *106*(6), 1044-1054 e1044. https://doi.org/10.1016/j.neuron.2020.03.024

Bartra, O., McGuire, J. T., & Kable, J. W. (2013). The valuation system: a coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *Neuroimage*, *76*, 412-427.

Beckmann, C. F., Jenkinson, M., & Smith, S. M. (2003). General multilevel linear modeling for group analysis in FMRI. *Neuroimage*, *20*(2), 1052-1063. http://www.sciencedirect.com/science/article/pii/S105381190300435X

Behrens, T. E., Woolrich, M. W., Walton, M. E., & Rushworth, M. F. (2007). Learning the value of information in an uncertain world. *Nat Neurosci*, *10*(9), 1214-1221. https://doi.org/10.1038/nn1954

Benjamin, D. J. (2019). Errors in probabilistic reasoning and judgment biases. *Handbook of Behavioral Economics: Applications and Foundations 1*, *2*, 69-186.

Brown, S. D., & Steyvers, M. (2009). Detecting and predicting changes. *Cogn Psychol*, *58*(1), 49-67. https://doi.org/10.1016/j.cogpsych.2008.09.002

Buckner, R. L., Krienen, F. M., & Yeo, B. T. (2013). Opportunities and limitations of intrinsic functional connectivity MRI. *Nat Neurosci*, *16*(7), 832-837. https://doi.org/10.1038/nn.3423

Chan, S. C., Niv, Y., & Norman, K. A. (2016). A probability distribution over latent causes, in the orbitofrontal cortex. *Journal of Neuroscience*, *36*(30), 7817-7828.

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci*, *3*(3), 201-215. https://doi.org/10.1038/nrn755

Costa, V. D., Tran, V. L., Turchi, J., & Averbeck, B. B. (2015). Reversal learning and dopamine: a bayesian perspective. *J Neurosci*, *35*(6), 2407-2416. https://doi.org/10.1523/JNEUROSCI.1989-14.2015

Daniel, K., Hirshleifer, D., & Subrahmanyam, A. (1998). Investor psychology and security market under-and overreactions. *the Journal of Finance*, *53*(6), 1839-1885.

De Bondt, W. F., & Thaler, R. (1985). Does the stock market overreact? *the Journal of Finance*, *40*(3), 793-805.

Dosenbach, N. U., Fair, D. A., Miezin, F. M., Cohen, A. L., Wenger, K. K., Dosenbach, R. A., Fox, M. D., Snyder, A. Z., Vincent, J. L., Raichle, M. E., Schlaggar, B. L., & Petersen, S. E. (2007). Distinct brain networks for adaptive and stable task control in humans. *Proc Natl Acad Sci U S A*, *104*(26), 11073-11078. https://doi.org/10.1073/pnas.0704320104

Edwards, W. (1968). Conservatism in human information processing. *Formal representation of human judgment*.

Efron, B., & Tibshirani, R. J. (1994). *An introduction to the bootstrap*. Chapman and Hall/CRC.

Eklund, A., Nichols, T. E., & Knutsson, H. (2016). Cluster failure: Why fMRI inferences for spatial extent have inflated false-positive rates. *Proceedings of the National Academy of Sciences*, *113*(28), 7900-7905.

Fellows, L. K., & Farah, M. J. (2003). Ventromedial frontal cortex mediates affective shifting in humans: evidence from a reversal learning paradigm. *Brain*, *126*(8), 1830-1837.

Gold, J. I., & Shadlen, M. N. (2007). The neural basis of decision making. *Annu. Rev. Neurosci.*, *30*, 535-574.

Griffin, D., & Tversky, A. (1992). The weighing of evidence and the determinants of confidence. *Cognitive psychology*, *24*(3), 411-435.

Hedge, C., Stothart, G., Todd Jones, J., Rojas Frias, P., Magee, K. L., & Brooks, J. C. (2015). A frontal attention mechanism in the visual mismatch negativity. *Behav Brain Res*, *293*, 173-181. https://doi.org/10.1016/j.bbr.2015.07.022

Heekeren, H. R., Marrett, S., Bandettini, P. A., & Ungerleider, L. G. (2004). A general mechanism for perceptual decision-making in the human brain. *nature*, *431*(7010), 859-862. https://doi.org/10.1038/nature02966

Heekeren, H. R., Marrett, S., Ruff, D. A., Bandettini, P. A., & Ungerleider, L. G. (2006). Involvement of human left dorsolateral prefrontal cortex in perceptual decision making is independent of response modality. *Proc Natl Acad Sci U S A*, *103*(26), 10023-10028. https://doi.org/10.1073/pnas.0603949103

Izquierdo, A., Brigman, J. L., Radke, A. K., Rudebeck, P. H., & Holmes, A. (2017). The neural basis of reversal learning: An updated perspective. *Neuroscience*, *345*, 12-26. https://doi.org/10.1016/j.neuroscience.2016.03.021

Kao, C.-H., Khambhati, A. N., Bassett, D. S., Nassar, M. R., McGuire, J. T., Gold, J. I., & Kable, J. W. (2020). Functional brain network reconfiguration during learning in a dynamic environment. *Nature communications*, *11*(1), 1682.

Kraemer, C., & Weber, M. (2004). How do people take into account weight, strength and quality of segregated vs. aggregated data? Experimental evidence. *Journal of Risk and Uncertainty*, *29*, 113-142.

Kremer, M., Moritz, B., & Siemsen, E. (2011). Demand Forecasting Behavior: System Neglect and Change Detection. *Management Science*, *57*(10), 1827-1843. https://doi.org/10.1287/mnsc.1110.1382

Litt, A., Plassmann, H., Shiv, B., & Rangel, A. (2011). Dissociating Valuation and Saliency Signals during Decision-Making. *Cerebral Cortex*, *21*(1), 95-102. http://cercor.oxfordjournals.org/content/21/1/95.abstract

Mante, V., Sussillo, D., Shenoy, K. V., & Newsome, W. T. (2013). Context-dependent computation by recurrent dynamics in prefrontal cortex. *nature*, *503*(7474), 78-84.

Massey, C., & Wu, G. (2005). Detecting regime shifts: The causes of under-and overreaction. *Management Science*, *51*(6), 932-947.

McGuire, J. T., Nassar, M. R., Gold, J. I., & Kable, J. W. (2014). Functionally dissociable influences on learning rate in a dynamic environment. *Neuron*, *84*(4), 870-881.

Meng, X.-L., Rosenthal, R., & Rubin, D. B. (1992). Comparing correlated correlation coefficients. *Psychological bulletin*, *111*(1), 172.

Muller, T. H., Mars, R. B., Behrens, T. E., & O'Reilly, J. X. (2019). Control of entropy in neural models of environmental state. *elife*, *8*, e39404.

Nassar, M. R., Wilson, R. C., Heasly, B., & Gold, J. I. (2010). An approximately Bayesian delta-rule model explains the dynamics of belief updating in a changing environment. *Journal of Neuroscience*, *30*(37), 12366-12378.

Nelson, M. W., Bloomfield, R., Hales, J. W., & Libby, R. (2001). The effect of information strength and weight on behavior in financial markets. *Organizational Behavior and Human Decision Processes*, *86*(2), 168-196.

O'Doherty, J., Kringelbach, M. L., Rolls, E. T., Hornak, J., & Andrews, C. (2001). Abstract reward and punishment representations in the human orbitofrontal cortex. *Nature neuroscience*, *4*(1), 95-102.

Payzan-LeNestour, E., & Bossaerts, P. (2011). Risk, unexpected uncertainty, and estimation uncertainty: Bayesian learning in unstable settings. *PLoS computational biology*, *7*(1), e1001048.

Payzan-LeNestour, E., Dunne, S., Bossaerts, P., & O'Doherty, J. P. (2013). The neural representation of unexpected uncertainty during value-based decision making. *Neuron*, *79*(1), 191-201.

Philiastides, M. G., Biele, G., & Heekeren, H. R. (2010). A mechanistic account of value computation in the human brain. *Proceedings of the National Academy of Sciences*, *107*(20), 9430-9435.

Roitman, J. D., & Shadlen, M. N. (2002). Response of neurons in the lateral intraparietal area during a combined visual discrimination reaction time task. *Journal of Neuroscience*, *22*(21), 9475-9489.

Sanders, N. R., & Manrodt, K. B. (2003). Forecasting Software in Practice: Use, Satisfaction, and Performance. *Interfaces*, *33*(5), 90-93. https://doi.org/10.1287/inte.33.5.90.19251

Schoenbaum, G., Chiba, A. A., & Gallagher, M. (2000). Changes in functional connectivity in orbitofrontal cortex and basolateral amygdala during learning and reversal training. *Journal of Neuroscience*, *20*(13), 5179-5189.

Schonberg, T., Fox, C. R., Mumford, J. A., Congdon, E., Trepel, C., & Poldrack, R. A. (2012). Decreasing ventromedial prefrontal cortex activity during sequential risk-taking: an fMRI investigation of the balloon analog risk task. *Frontiers in neuroscience*, *6*, 25510.

Schuck, N. W., Cai, M. B., Wilson, R. C., & Niv, Y. (2016). Human orbitofrontal cortex represents a cognitive map of state space. *Neuron*, *91*(6), 1402-1412.

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., Reiss, A. L., & Greicius, M. D. (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci*, *27*(9), 2349-2356. https://doi.org/10.1523/JNEUROSCI.5587-06.2007

Seifert, M., Ulu, C., & Guha, S. (2023). Decision Making Under Impending Regime Shifts. *Management Science*. https://doi.org/10.1287/mnsc.2022.4661

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage*, *44*(1), 83-98.

Soltani, A., & Izquierdo, A. (2019). Adaptive learning under expected and unexpected uncertainty. *Nat Rev Neurosci*, *20*(10), 635-644. https://doi.org/10.1038/s41583-019-0180-y

Ting, C.-C., Yu, C.-C., Maloney, L. T., & Wu, S.-W. (2015a). Neural mechanisms for integrating prior knowledge and likelihood in value-based probabilistic inference. *Journal of Neuroscience*, *35*(4), 1792-1805.

Ting, C.-C., Yu, C.-C., Maloney, L. T., & Wu, S.-W. (2015b). Neural Mechanisms for Integrating Prior Knowledge and Likelihood in Value-Based Probabilistic Inference. *The Journal of Neuroscience*, *35*(4), 1792-1805.

Tversky, A., Griffin, D., Heath, C., & Slovic, P. (1990). *Decision under conflict: Resolution and confidence in judgment and choice*. Department of Psychology, Stanford University.

Vilares, I., Howard, J. D., Fernandes, H. L., Gottfried, J. A., & Kording, K. P. (2012). Differential representations of prior and likelihood uncertainty in the human brain. *Current Biology*, *22*(18), 1641-1648.

Vincent, J. L., Kahn, I., Snyder, A. Z., Raichle, M. E., & Buckner, R. L. (2008). Evidence for a frontoparietal control system revealed by intrinsic functional connectivity. *J Neurophysiol*, *100*(6), 3328-3342. https://doi.org/10.1152/jn.90355.2008

Walton, M. E., Behrens, T. E., Buckley, M. J., Rudebeck, P. H., & Rushworth, M. F. (2010). Separable learning systems in the macaque brain and the role of orbitofrontal cortex in contingent learning. *Neuron*, *65*(6), 927-939.

Warbrick, T., Mobascher, A., Brinkmeyer, J., Musso, F., Richter, N., Stoecker, T., Fink, G. R., Shah, N. J., & Winterer, G. (2009). Single-trial P3 amplitude and latency informed event-related fMRI models yield different BOLD response patterns to a target detection task. *Neuroimage*, *47*(4), 1532-1544. https://doi.org/10.1016/j.neuroimage.2009.05.082

Wilson, R. C., & Collins, A. G. (2019). Ten simple rules for the computational modeling of behavioral data. *elife*, *8*, e49547.

Wilson, R. C., Takahashi, Y. K., Schoenbaum, G., & Niv, Y. (2014). Orbitofrontal cortex as a cognitive map of task space. *Neuron*, *81*(2), 267-279.

Woldorff, M. G., Hazlett, C. J., Fichtenholtz, H. M., Weissman, D. H., Dale, A. M., & Song, A. W. (2004). Functional parcellation of attentional control regions of the brain. *J Cogn Neurosci*, *16*(1), 149-165. https://doi.org/10.1162/089892904322755638

Woo, C.-W., Krishnan, A., & Wager, T. D. (2014). Cluster-extent based thresholding in fMRI analyses: pitfalls and recommendations. *Neuroimage*, *91*, 412-419.

Woolrich, M. W., Behrens, T. E. J., Beckmann, C. F., Jenkinson, M., & Smith, S. M. (2004). Multilevel linear modelling for FMRI group analysis using Bayesian inference. *Neuroimage*, *21*(4), 1732-1747. http://www.sciencedirect.com/science/article/pii/S1053811903007894

Worsley, K. J., Evans, A. C., Marrett, S., & Neelin, P. (1992). A three-dimensional statistical analysis for CBF activation studies in human brain. *Journal of Cerebral Blood Flow & Metabolism*, *12*(6), 900-918.

Yang, Y.-Y., & Wu, S.-W. (2020). Base rate neglect and neural computations for subjective weight in decision under uncertainty. *Proceedings of the National Academy of Sciences*, *117*(29), 16908-16919.

Yates, J. L., Park, I. M., Katz, L. N., Pillow, J. W., & Huk, A. C. (2017). Functional dissection of signal and noise in MT and LIP during decision-making. *Nat Neurosci*, *20*(9), 1285-1292. https://doi.org/10.1038/nn.4611

Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zollei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*, *106*(3), 1125-1165. https://doi.org/10.1152/jn.00338.2011

Supplementary Information

7   Detecting regime shifts: neurocomputational substrates for over- and
8   underreactions to change

SUPPLEMENTARY METHODS

Experiment 2

The procedure of Experiment 2 was identical to Experiment 1, except that there was no regime shift in this experiment. In other words, we only manipulated the signal diagnosticity in this experiment. The regime that the sensory signals were drawn from (red or blue urn) was randomly determined such that in half of the trials the regime was red and the other half the regime was blue. The order of the regimes across trials was pseudo randomized for each subject separately. Identical to Experiment 1, the subjects in Experiment 2 were instructed to estimate the probability that regime was the blue urn at the presentation of each new signal. Therefore, in both Experiments 1 and 2, the subjects estimated the probability that the current regime was the blue regime. However, it was only in Experiment 1 that the probability estimates conveyed information about the subjects' belief about change (whether the regime had shifted).

Experiment 3

We referred to Experiment 3 as the motor-equivalent task of Experiments 1 and 2. The primary goal of this experiment was to rule out the motor confounds for probability estimates in Experiment 1. We were aware that brain regions whose activity correlated with probability estimates in Experiment 1 can simply reflect entering of numbers through button presses and thus have nothing to do with estimating the probability of change. Therefore, to establish the evidence for the neural representations of probability estimates, we needed to rule out the motor confounds. The motor-equivalent task shared many of the physical and motor aspects of the tasks in Experiments 1 and 2. First, the task structure was identical. In the motor-equivalent task, each trial contained ten rounds. In each round, a hollow dot with no color information was shown to indicate the number of this round. A random number from 0 to 99 was presented at the center of screen. Subjects had to enter the indicated number with two successive button presses within 4 seconds. After this button-press stage there was feedback on the number they just entered, with white given for correct inputs and yellow given for incorrect entries. The reward was $3 for each correct answer and -$3 for incorrect answer. The accumulated reward outcome for each trial was revealed at the end of the trial.

Computational models for Experiment 2

3 Below we describe the computational models (Bayesian model and system-neglect

4 model) for Experiment 2 where subjects performed identical task as in Experiment 1

5 with one exception that no regime shift was possible (transition probability $q = 0$).

6 We fit the system-neglect model and the parameter estimates can be found in

7 Supplementary Fig. 2.

Bayesian model

9 The Bayesian posterior odds of the blue regime at the t-th period given the signal

10 history H_t were calculated by:

11
$$\frac{P_t^B}{1-P_t^B} = \frac{\Pr(B_t|H_t)}{\Pr(R|H_t)} = \frac{\Pr(B)}{\Pr(R)} \times \frac{\Pr(H_t|B)}{\Pr(H_t|R)}.$$

12 where $\frac{\Pr(B)}{\Pr(R)}$ is the prior odds and $\frac{\Pr(H_t|B)}{\Pr(H_t|R)}$ is the likelihood ratio. Given that the

13 probability of being in the blue regime and being in the red regime were equal

14 $\Pr(B) = \Pr(R) = 0.5$, Eq. (3) can be rewritten as

15
$$\frac{P_t^B}{1-P_t^B} = \frac{\Pr(B_t|H_t)}{\Pr(R|H_t)} = \frac{0.5}{0.5} \times \frac{\Pr(H_t|B)}{\Pr(H_t|R)} = \frac{\Pr(H_t|B)}{\Pr(H_t|R)}.$$

16 The likelihood ratio is computed according to the following equation

17
$$\frac{\Pr(H_t|B)}{\Pr(H_t|R)} = d^{t-2\sum_{k=1}^{t} r_k}.$$

18 where d represents signal diagnosticity. The exponent of d captures the difference

19 between the number of the blue balls and red balls from the first to the t-th period. For

20 example, suppose that at the 5th period there are 4 red balls and 1 blue ball. Then

21 $d^{t-2\sum_{k=1}^{t} r_k}$ would be $d^{5-2\times 3} = d^{-3}$ where the exponent is the number of blue balls

22 minus the number of red balls.

System-neglect model

Similar to Experiment 1, we developed and fit the system-neglect model. The system-

neglect model consisted of a weighting parameter β for signal diagnosticity

$$\frac{P_t^B}{1-P_t^B} = \frac{\Pr(B_t|H_t)}{\Pr(R|H_t)} = d^{\beta(t-2\sum_{k=1}^{t} r_k)}.$$

where we separately estimated β for each level of signal diagnosticity

$$\beta = \beta_1 D_1 + \beta_2 D_2 + \beta_3 D_3$$

Here D_n is the dummy variable for diagnosticity for d_n. Note that $\beta = 1$

corresponds to Bayesian estimates.

    SUPPLEMENTARY **R**ESULTS

Figure S1. Probability estimates from all subjects are plotted as histograms separately for each condition—a combination of transition probability and signal diagnosticity. The blue bars represent the actual probability estimates, while the orange bars correspond to the probability estimates predicted by the Bayesian model.

3   **Figure S2.** Experiment 2: estimates of the weighting parameter for signal diagnosticity (β)

4   in the system-neglect model. Dashed lines indicate parameter value equal to 1.

Figure S3. Parameter recovery. We simulated probability estimates according to the system-neglect model. We used each subject's parameter estimates as our choice of parameter values used in the simulation. Using simulated data, we estimated the parameters (α and β) in the system-neglect model. To examine parameter recovery, we plotted the parameter values we used to simulate the data against the parameter estimates we obtained based on simulated data and computed their Pearson correlation. Further, we added different levels of Gaussian white noise with standard deviation $\sigma = [0.01, 0.05, 0.1, 0.2, 0.3]$ to the simulated data to examine parameter recovery and show the results respectively in Fig. A, B, and C. For each noise level, we show the parameter estimates in the left two graphs. In the right two graphs, we plot the parameter estimates based on simulated data against the parameter values used to simulate the data. **A.** Noise $\sigma = 0.01$. **B.** Noise $\sigma = 0.05$. **C.** Noise $\sigma = 0.1$. **D.** Noise $\sigma = 0.2$. **E.** Noise $\sigma = 0.3$. **F.** Empirically estimated noise (σ) of each subject. Each bar represents a subject's estimated noise level.

A

B

Figure S4. Impact of noise homoscedasticity on parameter estimation. **A.** Empirically estimated residual standard deviation. Mean residual standard deviation (across subjects, black data points) in the five probability intervals, [0.0–0.2), [0.2–0.4), [0.4–0.6), [0.6–0.8), and [0.8–1.0], were 0.1015, 0.1296, 0.1987, 0.1929, and 0.2061, respectively. Error bars represent ± 1 standard error of the mean. **B.** Parameter recovery results assuming heteroscedastic noise. We performed parameter recovery using the empirically estimated, probability-dependent residual variance shown in A (the mean residual standard deviation estimates). Conventions are the same as in Fig. S3.

Figure S5. Probability estimates from the actual and simulated data. **A.** Histogram of
subjects' actual probability estimation data collapsed across all conditions (left graph) and
simulated probability estimation data under three different noise levels (Noise $\sigma =$
$0.01, 0.05, 0.1$). Descriptions of how we performed simulations can be seen in Supplementary
Fig. S3. **B.** Subjects data are plotted as histograms separately for each condition.

Figure S6. B. System-neglect model can well-describe subjects' over- and underreactions to

change. We fit the system-neglect model to each individual subject' probability estimates and

used the resulting parameter estimates to compute each subject's probability estimates under

the system-neglect model (P_t^{SN}). We then used P_t^{SN} to compute Index of Overreaction

(IO). Here, IO was computed by subtracting belief revision predicted by the Bayesian model

($\Delta P_t^B = P_t^B - P_{t-1}^B$) from belief revision estimated by system-neglect model ($\Delta P_t^{SN} =$

$P_t^{SN} - P_{t-1}^{SN}$). Formally, $IO = \Delta P_t^{SN} - \Delta P_t^B$. The mean IO (across all subjects; indicated

by the bars) is plotted as a function of transition probability and signal diagnosticity. Solid

symbols represent data points from individual subjects. Error bars represent ± 1 standard

error of the mean. The patterns of over- and underreactions here resembled those based on

actual data (Fig. 2B), suggesting that the system-neglect model can describe subjects' over-

and underreactions well.

1 **Robustness checks on the probability estimates of regime shifts (Figs. S7 to S9)**

x=-2   y=-18   z=-8

Figure S7. Whole-brain results on activity that significantly correlated with probability

estimates of regime shift (P_t) in GLM-2. In GLM-2, we implemented variables important for

estimating regime shifts as parametric regressors in addition to P_t. In particular, the

intertemporal prior that captured the prior probability of change increasing as a function of

time was included in the GLM-2. Still, we observed P_t correlates in the vmPFC and ventral

striatum after controlling for these variables. Color for significant activations: blue indicates

negative correlation with P_t; yellow indicates positive correlation with P_t. Familywise

error-corrected at $p < .05$ using Gaussian random field theory with a cluster-forming

threshold $z > 3.1$.

x=-2   y=-18   z=-8

Figure S8. Whole-brain results on activity that significantly correlated with the subjects' log

odds estimates of regime shift, $\ln(P_t/(1 - P_t))$, where P_t represents subjects' probability

estimates. In this analysis, we replaced the parametric regressor of P_t with the log odds of

regime shifts in GLM-1. Color for significant activations: blue indicates negative correlation

with $\ln(P_t/(1 - P_t))$. Familywise error-corrected at $p < .05$ using Gaussian random field

theory with a cluster-forming threshold $z > 3.1$.

Figure S9. Probability estimates of regime shifts (P_t) separately analyzed for change-consistent (blue) and change-inconsistent (red) signals. Whole-brain results on activity that significantly correlated with probability estimates of regime shift for change-consistent and change-inconsistent signals. We observed that vmPFC correlated with P_t for both types of signals. Blue color showing significant activations indicates negative correlation with P_t. Familywise error-corrected at $p < .05$ using Gaussian random field theory with a cluster-forming threshold $z > 3.1$.

Figure S10. Independent region-of-interest (ROI) analysis in vmPFC and ventral striatum on

probability estimates across the three experiments (Experiment 1 was the main experiment on

regime-shift detection, Experiments 2 and 3 were control experiments). The vmPFC ROI we

used was based on Bartrat et al. (2013). The ventral striatum mask was based on nucleus

accumbens mask from Harvard-Oxford Cortical Structural Atlas. For each subject and each

ROI, we extracted the mean parameter estimates (PE) of the probability estimates contrast in

GLM-1. **A.** vmPFC ROI. Experiment 1: One-sample t test, $t(29) = -3.82$, $p < 0.01$;

Experiment 2: One-sample t test, $t(29) = 0.36$, $p = 0.71$; Experiment 3: One-sample t test,

$t(29) = -1.11$, $p = 0.28$; Experiments 1 $-$ Experiment 2: two-sample t test, $t(58) =$

$-3.67, p < 0.01$; Experiments 1 $-$ Experiment 3: two-sample t test, $t(58) = -3.12$, $p <$

0.01;. **B.** Ventral striatum ROI. Experiment 1: $(29) = -3.06$, $p < 0.01$; Experiment 2:

$t(29) = 0.44$, $p = 0.67$; Experiment 3: $t(29) = -0.93$, $p = 0.36$; Experiments 1 $-$

Experiment 2: $t(58) = -2.55$, $p = 0.01$; Experiments 1 $-$ Experiment 3: $t(58) = -1.95$,

$p = 0.06$. The * symbol indicates $p<0.05$ (two-tailed), and ** symbol indicates $p<0.01$ (two-

tailed).

Figure S11. Key variables for regime-shift computations. **A.** Whole-brain results on the intertemporal prior of regime shift. **B.** Using the intertemporal prior ROI (left graph: magenta indicates voxels shared by the LOSO ROI of all subjects; blue indicates voxels of LOSO ROI of at least one subject) to examine the regression coefficients of the strength of evidence in favor of change, $\ln(d) \times$ signal. The mean parameter estimates (mean PE), i.e., regression coefficient, was not significantly different from 0 (one-sample t test, $t(29) = 0.54, p = 0.59$, two-tailed).

1   **Table S1.** Experiment 1: Probability estimates (P_t) and belief revision (ΔP_t) contrasts

2   based on GLM-1. Cluster-level inference using Gaussian random field theory (familywise

3   error corrected at $p < .05$ with a cluster-forming threshold $z > 3.1$).

Probability estimates P_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	**z-max**	**z-max(x,y,z)**
Central Opercular Cortex	R	26990	6.13	(62,-6,6)
Cingulate Gyrus, anterior division	-	989	4.51	(0,18,36)
Belief revision ΔP_t (positive correlation)				
Frontal Orbital Cortex	L	3105	4.94	(-24,18,-10)
Cingulate Gyrus, anterior division	L	2589	4.34	(-4,26,30)
Frontal Orbital Cortex	R	1796	4.57	(26,20,-12)
Right Cerebral White Matter	R	1557	4.67	(12,-14,-4)
Lateral Occipital Cortex, superior division	L	1366	4.34	(-28,-82,34)
Postcentral Gyrus	R	1262	4.71	(60,-18,36)
Postcentral Gyrus	L	1088	4.48	(-58,-22,36)
Postcentral Gyrus	L	587	4.25	(-34,-26,68)

1 **Table S2.** Experiment 1: Probability estimates (P_t) and belief revision (ΔP_t) contrasts

2 based on GLM-1. Permutation tests based on threshold-free-cluster-enhancement (TFCE)

3 statistic.

Probability estimates P_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	p_{max}	**1-p$_{max}$(x,y,z)**
Occipital Pole	R	62107	0	(18,-98,22)
Frontal Pole	L	676	0.036	(-32,40,34)
Belief revision ΔP_t (positive correlation)				
Frontal Orbital Cortex	L	75881	0.002	(-22,14,-16)
Frontal Pole	R	424	0.045	(34,36,32)
Supramarginal Gyrus, posterior division	L	53	0.048	(-68,-46,10)

1 **Table S3.** Experiment 1: Probability estimates (P_t) and belief revision (ΔP_t) contrasts
2 based on GLM-1. Permutation tests based on cluster extent.
3

Probability estimates P_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	p_{max}	**1-p$_{max}$(x,y,z)**
Lateral Occipital Cortex, inferior division	R	5326	0	(56,-66,-12)
Temporal Fusiform Cortex, anterior division	L	1989	0.001	(-38,-8,-32)
Planum Polare	R	926	0.003	(52,6,-4)
Middle Temporal Gyrus, anterior division	L	777	0.004	(-52,-4,-18)
Postcentral Gyrus	R	570	0.006	(44,-18,36)
Right Cerebral White Matter	R	394	0.009	(36,0,-30)
Cingulate Gyrus, anterior division	R	156	0.027	(2,30,16)
Middle Temporal Gyrus, posterior division	L	89	0.044	(-62,-36,0)
Precuneus Cortex	-	80	0.048	(0,-56,56)
Belief revision ΔP_t (positive correlation)				
Cingulate Gyrus, anterior division	L	3082	0.002	(-4,44,4)
Frontal Orbital Cortex	L	2083	0.002	(-24,14,-26)
Lingual Gyrus	R	1274	0.004	(2,-78,0)
Frontal Orbital Cortex	R	708	0.008	(14,12,-18)
Planum Polare	R	582	0.01	(44,-2,-20)
Brain-Stem	L	566	0.01	(-4,-36,-28)
Postcentral Gyrus	L	390	0.014	(-58,-20,24)
Supramarginal Gyrus, anterior division	R	366	0.014	(70,-20,24)
Lateral Occipital Cortex, superior division	L	233	0.024	(-32,-78,30)

Precentral Gyrus	L	207	0.028	(-28,-8,54)
Precentral Gyrus	L	188	0.03	(-54,6,24)
Precentral Gyrus	R	168	0.035	(56,8,24)

Table S4. Experiment 3: Instructed number (IN_t) and difference in instructed number

(ΔIN_t) contrasts based on GLM-1. For Experiment 3, IN_t represents the two-digit number

subjects were instructed to press at each period, and ΔIN_t represents the difference in

number between successive periods. IN_t is the control for P_t in Experiment 1, and ΔIN_t is

the control for ΔP_t in Experiment 1. Cluster-level inference using Gaussian random field

theory (familywise error corrected at $p < .05$ with a cluster-forming threshold $z > 3.1$).

Instructed number IN_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	**z-max**	**z-max(x,y,z)**
Precentral Gyrus	R	1320	4.66	(38,-22,62)
Postcentral Gyrus	L	521	4.08	(-40,-36,62)
Brain-Stem	L	419	4.22	(-4,-28,-10)
Cerebellar Left V	L	401	4.8	(-16,-50,-20)
Middle Frontal Gyrus	L	316	4.22	(-40,36,32)
Precentral Gyrus	R	294	3.95	(56,2,38)
Right Thalamus	R	248	4.31	(8,-18,0)
Superior Frontal Gyrus	L	223	3.87	(-20,-8,70)
Difference in instructed number ΔIN_t (positive correlation)				
Precentral Gyrus	L	1489	5.09	(-40,-20,54)

1   **Table S5.** Experiment 1 $-$ Experiment 2 based on the probability estimates (P_t) contrast in

2   GLM-1. Cluster-level inference using Gaussian random field theory (familywise error

3   corrected at $p < .05$ with a cluster-forming threshold $z > 3.1$).

Experiment 1 $>$ Experiment 2 on negative probability estimates contrast				
Cluster	**Hemisphere**	**Cluster size**	**z-max**	**z-max(x,y,z)**
Paracingulate Gyrus	L	7726	4.93	(-8,38,-8)
Lateral Occipital Cortex, superior division	L	3457	4.95	(-16,-86,46)
Postcentral Gyrus	R	328	3.82	(48,-20,64)

1 **Table S6.** Experiment 1 − Experiment 3: based on the probability estimates (P_t) contrast

2 in GLM-1. Cluster-level inference using Gaussian random field theory (familywise error

3 corrected at p<0.05 with a cluster-forming threshold z>3.1).

Experiment 1 > Experiment 2 on negative probability estimates contrast				
Cluster	**Hemisphere**	**Cluster size**	**z-max**	**z-max(x,y,z)**
Superior Temporal Gyrus, anterior division	L	4099	5.04	(-60,2,-2)
Occipital Pole	R	2656	5.21	(16,-98,28)
Central Opercular Cortex	R	1575	5.41	(62,-6,8)
Experiment 1 > Experiment 2 on positive probability estimates contrast				
Supramarginal Gyrus, anterior division	L	328	4	(-52,-38,44)

1 **Table S7.** Experiment 1: GLM-2. Cluster-level inference using Gaussian random field theory

2 (familywise error corrected at $p < .05$ with a cluster-forming threshold $z > 3.1$).

Probability estimates P_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	**z-max**	$z - \max (x, y, z)$
Lingual Gyrus	L	11504	6.2	(-14,-52,-12)
Central Opercular Cortex	R	9872	6.19	(60,-8,6)
Probability estimates (positive correlation)				
Postcentral Gyrus	L	1680	5.45	(-46,-26,56)
Belief revision ΔP_t (positive correlation)				
Left Cerebral White Matter	L	717	4.33	(-20,18,-12)
Cingulate Gyrus, anterior division	-	566	4.39	(0,32,-8)
Frontal Orbital Cortex	R	475	4.17	(20,8,-16)
Postcentral Gyrus	L	333	4.02	(-30,-28,64)
Intertemporal prior (negative correlation)				
Occipital Fusiform Gyrus	R	1833	4.61	(36,-74,-16)
Lateral Occipital Corte	R	223	4.04	(36,-66,-56)
Lateral Occipital Cortex	L	204	4.25	(-40,-82,-8)
$\ln(d)\times$ **signal (positive correlation)**				
Middle Frontal Gyrus	R	988	4.17	(36,10,30)
Superior Frontal Gyrus	R	821	4.56	(6,34,48)
Superior Parietal Lobule	L	620	4.26	(-40,-52,56)
Supramarginal Gyrus	R	604	4.29	(46,-40,48)
Middle Frontal Gyrus	L	277	4.21	(-46,34,30)

1 **Table S8** Experiment 1: GLM-2. Permutation tests based on the threshold-free-cluster-

2 enhancement (TFCE) statistic.

Probability estimates P_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	p_{max}	$1 - p_{max}(x, y, z)$
Postcentral Gyrus	R	31490	0	(38,-24,50)
Frontal Pole	R	165	0.038	(16,36,-20)
Probability estimates P_t (positive correlation)				
Postcentral Gyrus	L	439	0.012	(-42,-28,54)
Belief revision ΔP_t (positive correlation)				
Right Caudate	R	2857	0.027	(12,14,-4)
Cingulate Gyrus, anterior division	R	1873	0.027	(4,26,24)
Postcentral Gyrus	L	707	0.034	(-42,-22,54)
Insular Cortex	L	427	0.041	(-42,6,-4)
Frontal Medial Cortex	L	320	0.04	(-4,40,-14)
Frontal Pole	L	40	0.048	(-2,60,28)
Insular Cortex	R	20	0.048	(40,6,-14)
Intertemporal prior (negative correlation)				
Lateral Occipital Cortex, inferior division	R	2117	0.006	(38,-74,14)
Frontal Pole	R	185	0.038	(36,42,36)
Frontal Pole	R	114	0.043	(40,44,-6)
Middle Frontal Gyrus	R	14	0.048	(34,28,50)
Right Thalamus	R	13	0.046	(22,-28,-2)
Frontal Pole	R	1	0.05	(32,48,-8)
Frontal Pole	R	1	0.05	(30,52,-6)

1   **Table S9.** Experiment 1: GLM-2. Permutation tests based on cluster extent.

Probability estimates P_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	p_{max}	$1 - p_{max}(x, y, z)$
Postcentral Gyrus	R	1770	0.001	(42,-20,38)
Lateral Occipital Cortex, inferior division	R	1085	0.003	(60,-68,-6)
Lateral Occipital Cortex, inferior division	L	674	0.006	(-42,-84,10)
Temporal Fusiform Cortex, posterior division	L	597	0.007	(-36,-18,-32)
Heschl's Gyrus	R	527	0.008	(54,-12,2)
Temporal Fusiform Cortex, anterior division	R	441	0.01	(34,-2,-30)
Cerebellar Left V	L	235	0.022	(-16,-50,-24)
Middle Temporal Gyrus, anterior division	L	193	0.027	(-52,-4-18)
Frontal Medial Cortex	L	134	0.039	(-10,34,-18)
Lingual Gyrus	L	134	0.039	(-6,-48,-2)
Probability estimates P_t (positive correlation)				
Supramarginal Gyrus, anterior division	L	1012	0.002	(-44,-30,40)
Belief revision ΔP_t (positive correlation)				
Cingulate Gyrus, anterior division	-	450	0.016	(0,34,14)
Frontal Orbital Cortex	R	351	0.021	(16,12,-16)
Frontal Orbital Cortex	L	300	0.024	(-22,10,-18)
Postcentral Gyrus	L	215	0.035	(-38,-24,46)
Intertemporal prior (negative correlation)				
Temporal Occipital Fusiform Cortex	R	1687	0.002	(34,-60,24)

Lateral Occipital Cortex, superior division	R	220	0.042	(38,-70,32)
ln(d)× signal (positive correlation)				
Superior Parietal Lobule	L	384	0.022	(-38,-52,40)

1 **Table S10.** Experiment 1: GLM-1 without the action-handedness regressor. Cluster-level

2 inference using Gaussian random field theory (familywise error corrected at $p < .05$ with a

3 cluster-forming threshold $z > 3.1$).

Probability estimates P_t (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	**z-max**	**z-max(x,y,z)**
Temporal Occipital Fusiform Cortex	L	27876	6.63	(-24,-54,-22)
Supplementary Motor Cortex	R	1096	4.72	(12,0,46)
Probability estimates P_t (positive correlation)				
Postcentral Gyrus	L	705	4.73	(-46,-28,54)
Belief revision ΔP_t (positive correlation)				
Frontal Orbital Cortex	L	8939	5.02	(-24,20,-10)
Frontal Orbital Cortex	R	2244	4.7	(20,6,-14)
Lateral Occipital Cortex, superior division	L	1786	4.45	(-28,-82,34)
Postcentral Gyrus	L	1752	4.61	(-58,-24,38)
Postcentral Gyrus	R	1053	4.61	(60,-18,36)
Precentral Gyrus	L	361	3.92	(-52,6,32)

1 **Table S11.** Experiment 1: GLM-2 without P_t and ΔP_t regressors. Cluster-level inference

2 using Gaussian random field theory (familywise error corrected at $p < .05$ with a cluster-

3 forming threshold $z > 3.1$).

Intertemporal prior (negative correlation)				
Cluster	**Hemisphere**	**Cluster size**	**z-max**	$z - \mathbf{max}\ (x, y, z)$
Temporal Occipital Fusiform Cortex	R	2101	5.3	(44,-54,-20)
Lateral Occipital Cortex, inferior division	L	976	4.49	(-40,-82,-8)
Paracingulate Gyrus	R	316	4.12	(14,34,30)
Right Thalamus	R	240	4.32	(22,-28,-4)
Precuneus Cortex	R	230	4.13	(6,-82,48)
Frontal Pole	R	186	3.73	(42,44,34)
$\ln(d)\times$ **signal (positive correlation)**				
Supramarginal Gyrus, posterior division	L	945	4.58	(-50,-42,56)
Supramarginal Gyrus, posterior division	R	923	4.44	(46,-40,48)
Middle Frontal Gyrus	R	785	4.19	(46,12,50)
Superior Frontal Gyrus	R	521	4.49	(6,32,50)
Middle Frontal Gyrus	L	226	4.1	(-46,34,30)

Table S12. Model-fitting summary.

Model	Likelihood	Number of Parameters	AIC
SN-original	356.27 $\pm$ 44.7	6	-700.54 ± 89.49
SN-SigDep-β	383.08 $\pm$ 44.69	9	-748.15 ± 89.37
SN-SigDep-α	398.61 $\pm$ 45.78	9	-779.21 ± 91.57
SN-SigDep-$\alpha\beta$	410.03 $\pm$ 46.37	12	-796.07 ± 92.74

1   **Table S13.** Model comparison using paired t-test with Bonferroni correction.

Model	t test $(df = 29)$	p-value	Bonferroni-corrected p-value
SN-SigDep-β − SN-original	5.15	0.0000	0.0001
SN-SigDep-α − SN-original	6.93	0.0000	0.0000
SN-SigDep-$\alpha\beta$ − SN-original	8.56	0.0000	0.0000
SN-SigDep-α − SN-SigDep-β	4.01	0.0004	0.0023
SN-SigDep-$\alpha\beta$ − SN-SigDep-β	6.55	0.0000	0.0000
SN-SigDep-$\alpha\beta$ − SN-SigDep-α	5.03	0.0000	0.0001

Task Instruction (in Chinese) for Experiment 1

All subjects were instructed in Chinese with this document. We also included the English translation version after the Chinese version.

實驗介紹

簡介：

想像我們手邊有兩個箱子：紅箱子與藍箱子。紅箱子中有比較多的紅球；藍箱子中有比較多的藍球。我們會從箱子中抽球給您看，您會知道球的顏色，但不知道球是從紅箱子或藍箱子抽出的。您要做的作業為<mark>猜測該球是從藍箱子抽出的機率為何</mark>。

Color ratio= [60   Color ratio= [40

抽球規則：

1. 一題會抽出10顆球。一次抽出一顆球。
2. 抽球並不會都從同一個箱子。一開始設定的箱子為紅箱子，但從抽第一顆球開始，電腦便有<mark>一定的機率轉換</mark>成由藍箱子抽球。
3. 箱子最多只會轉換一次。一旦轉換到藍箱子， 就不會再轉換回紅箱子。
4. 球取後會被放回原箱子。故在同一題中兩箱子個紅藍球比例會是固定的。且兩箱子的顏色比例是相反的。

實驗流程

3 首先，您會看到該題「箱子間轉換的機率」與「箱子中紅籃球的比例」

4 您會看到電腦從箱子中抽出一顆球，看到球後即可估計<mark>球是從藍箱子抽出的機</mark>

5 <mark>率</mark>。每題共有 10 次觀察。

7
8

9 在實驗中，每題都會提供給您以下的資訊以幫助您做機率的估計，這些資訊在

10 您做估計時都會提供給您：

11 1. 紅箱子中紅藍球的比例。(您可以依據這個比例知道藍箱子的比例)

12 2. 轉換機率：在每次觀察中，由紅箱子抽球轉換到由藍箱子抽球的機率。

13
14

什麼是轉換機率？

假設轉換機率為 25％，那表示每次觀察電腦都有 25％的機率由紅箱子轉換到

由藍箱子抽球。這表示第一次觀察您所看到的球有 25％的機率來自藍箱子，75

％的機率來自紅箱子。一旦轉換成由藍箱子抽球，那麼直到這一題十次觀察結

束，都會由藍箱子抽球。若第一次觀察到的球是由紅箱抽出，那麼下一次觀察

的球是由藍箱子抽出的機率也會是 25％。每次觀察的轉換機率為獨立。因此，

轉換機率其實就是在每次觀察中，由紅箱子抽球轉換到由藍箱子抽球的機率。

當然，電腦也有可能 10 次都由紅箱抽球，或 10 次都由藍箱抽球。

範例

酬賞: 一題（十次嘗試）最多可獲得 30 元，實驗結束時會抽十題實現。

計算方式 30× ($0.10 - ($0.20 × Error^2)) 猜測越近可以得到越多錢

Period 1:

猜測球是從藍箱子抽出的機率為 10%，實際上是由紅箱子抽出。

30*($0.10 - ($0.20 × 0.1^2) = 2.94

Period 7:

猜測球是從藍箱子抽出的機率為 10%，實際上是由藍箱子抽出。

30*($0.10 - ($0.20 × 0.9^2) = -1.86

Task Instruction for Experiment 1

Brief Introduction:

Imagine two boxes, red and blue, placed in front of you. In the red box, there are more

red balls. In the blue box, there are more blue balls, as shown below. We will draw

balls from one of the boxes and show them to you. You will see the color of the

drawn balls, but you will not know whether they come from the red or blue box. Your

task is to <mark>estimate the probability that the drawn ball comes from the blue box</mark>.

Color ratio= [60   Color ratio= [40

Ball drawing rules：

1. On each trial, we will draw a total of 10 balls. We will draw one ball at a time.

2. We will not always draw from the same box. In the beginning, we will always start

from the red box. But before we draw the first ball, there is a certain probability that

the box we draw from would switch to the blue box.

3. A shift would happen at most one time during a trial. Once a shift to the blue box

happens, it is no longer possible to shift again from the blue to the red box.

4. Once a ball is drawn and shown to you, it would be returned back to the box.

Therefore, during a trial the ratio of blue and red balls in both boxes would be fixed.

And the color ratio is opposite between the two boxes.

Experiment Procedure

At the beginning of a trial, you will first see information about the transition probability and the color ratio of this trial. Then, you will see a ball drawn from a box (an observation). When you see the ball, ==estimate the probability that it comes from the blue box==. There are a total of 10 observations in a trial.

During the experiment, in each trial we will give you the following information to help you estimate probability:

1. The ratio of the red and blue balls in the red box (you can use this ratio to know the proportion of red and blue balls in the blue box).

2. Transition probability: this is the probability that, at each observation, we shift from drawing the ball from the red box to the blue box.

What is transition probability?

Suppose that transition probability is 25%. This indicates that at each observation

there is a 25% chance that the box switches from the red box to the blue box. It means

that at the first observation, there is a 25% chance that the drawn ball comes from the

blue box, and a 75% chance that the ball comes from the red box. During a trial, once

the box transitions from the red to the blue box, all the remaining balls will be drawn

from the blue box. If the drawn ball at the first observation comes from the red box,

then at the second observation the probability that the ball comes from the blue box

would also be 25%. At each observation the transition probability is independent. In

other words, transition probability is, at each observation, the probability that the box

switches from the red to the blue box. Of course, it is possible that all the 10

observations come from the red box, or all 10 observations comes from the blue box.

Example

Reward: you can earn at most 30 dollars in a trial (10 observations). At the end of the

experiment, we will randomly select 10 trials to realize your payoffs.

Reward calculation formula: $30 \times$ (\$0.10 - (\$0.20 $\times$ Error^2)). This means that when

your estimate is closer to the true box where the ball is drawn you would receive more

money.

Observation 1:

You guess that the probability that the ball comes from the blue box is 10%.

However, it was in fact coming from the red box. As a result, you win

$30*$(\$0.10 - (\$0.20 $\times 0.1^2$)) = 2.94 at this observation.

Observation 7:

You guess that the probability that the ball comes from the blue box is 10%.

However, it was in fact coming from the red box. As a result, you lose

$30*$(\$0.10 - (\$0.20 $\times 0.9^2$)) = -1.86 at this observation.

1 Demo of the trial sequence

2 Here we show the sequence of two example trials.

4
5

Period 1
Estimation feedback
0.5s

ISI fixation
1-5s

Period 2
Estimation
4s

Period 2
Estimation feedback
0.5s

ISI fixation
1-5s

Period 3
Estimation
4s

Period 3
Estimation feedback
0.5s

1
2

ISI fixation
1-5s

Period 4
Estimation
4s

Period 4
Estimation feedback
0.5s

ISI fixation
1-5s

Period 5
Estimation
4s

Period 5
Estimation feedback
0.5s

ISI fixation
1-5s

2
3

Period 6
Estimation
4s

Period 6
Estimation feedback
0.5s

ISI fixation
1-5s

Period 7
Estimation
4s

Period 7
Estimation feedback
0.5s

ISI fixation
1-5s

Period 8
Estimation
4s

Period 8
Estimation feedback
0.5s

ISI fixation
1-5s

Period 9
Estimation
4s

Period 9
Estimation feedback
0.5s

ISI fixation
1-5s

Period 10
Estimation
4s

Period 10
Estimation feedback
0.5s

ISI fixation
1-5s

win $27

Reward feedback
2s

ITI
1-5s

Trial information
3s

Period 1
Estimation
4s

Period 1
Estimation feedback
0.5s

3
4

ISI fixation
1-5s

Period 2
Estimation
4s

Period 2
Estimation feedback
0.5s

ISI fixation
1-5s

Period 3
Estimation
4s

Period 3
Estimation feedback
0.5s

ISI fixation
1-5s

Period 4
Estimation
4s

Period 4
Estimation feedback
0.5s

ISI fixation
1-5s

Period 5
Estimation
4s

Period 5
Estimation feedback
0.5s

ISI fixation
1-5s

Period 6
Estimation
4s

Period 6
Estimation feedback
0.5s

ISI fixation
1-5s

Period 7
Estimation
4s

Period 7
Estimation feedback
0.5s

ISI fixation
1-5s

Period 8
Estimation
4s

Period 8
Estimation feedback
0.5s

ISI fixation
1-5s

Period 9
Estimation
4s

Period 9
Estimation feedback
0.5s

ISI fixation
1-5s

3
4

Period 10
Estimation
4s

Period 10
Estimation feedback
0.5s

ISI fixation
1-5s

Reward feedback
2s

ITI
1-5s

2
3
4

A

$$\Delta P_t = P_t - P_{t-1}$$

Legend: P_t (blue), ΔP_t (orange)

Left plot — y-axis: Probability estimates; x-axis: Period (t)

Right plot — y-axis: ΔP_t; x-axis: Period (t)

B

vmPFC

x=-4   y=32   z=-6   R   L

ventral striatum

y=10   z=-6   R   L

P_t: -3.6 to -6.2

ΔP_t: 3.6 to 5

P_t and ΔP_t

C

Expt. 1 - Expt. 2

x=-4   y=10   z=-6

-5.4 to -3.1

D

Expt. 1 - Expt. 3

x=-4   y=10   z=-6

-5.4 to -3.1

A

ln(*d*)×signal

d = 1.5
q = 0.01

d = 9
q = 0.1

▲shift

Period (*t*)

R   L

4.6

3.1

x=4   y=28   z=46

B

Intertemporal prior

q = 0.01

q = 0.1

Period (*t*)

dmPFC

x=4

rIFG   lIFG

y=28

rIPS   lIPS

z=48

mean PE

*   *   *   *

dmPFC   lIFG   rIFG   lIPS   rIPS