# Foveal RGCs develop abnormal calcium dynamics weeks after photoreceptor ablation

# Abstract

### Objective or purpose
Physiological changes in retinal ganglion cells (RGCs) have been reported in rodent models of photoreceptor (PR) loss but this has not been investigated in primates. By expressing both a calcium indicator (GCaMP6s) and an optogenetic actuator (ChrimsonR) in foveal RGCs of a macaque, we reactivate RGCs *in vivo* and assess activity in the weeks and years following PR loss.

### Design
We used an in vivo calcium imaging approach to record optogenetically evoked activity in deafferented RGCs in primate fovea. Cellular scale recordings were made longitudinally over a 10 week period following photoreceptor ablation and compared to RGC responses from RGCs that had lost photoreceptor input more than two years prior.

### Participants
Three eyes received photoreceptor ablation, OD of a male *Macacus rhesus* (M1), OS of a female *Macacus fascicularis* (M2) and OD of a male *Macacus fascicularis* (M3). Two animals were used for *in vivo* recording, one for histological assessment.

### Methods
Cones were ablated with an ultrafast laser delivered through an adaptive optics scanning light ophthalmoscope (AOSLO). A 0.5 s, 25Hz stimulus was delivered to optogenetically stimulate the deafferented RGCs, and the resulting GCaMP fluorescence rise and decay were recorded from RGCs with an AOSLO. These measurements were repeated in the 10 weeks immediately after photoreceptor ablation and at 2 years.

### Main Outcome measures
The rise time, decay constant and response magnitude of the deafferented RGCs was derived from the GCaMP fluorescence following optogenetic stimulation. These measurements were made at the cellular scale from 221(M1) and 218 (M2) RGCs *in vivo.*

### Results
The mean time to peak calcium was stable in deafferented RGCs over the 10 weeks post ablation (paired t-test, $p<0.001$), while the mean decay constant of the calcium response decreased by 1.5 fold 1.6±0.5 s to 0.6±0.3 s within 10 weeks in subject 1 and 2.1 fold (2.5±0.5 s to 1.2±0.2 s SD) within 8 weeks in subject 2.

### Conclusions
The presence of optogenetic responses 2 years after PR loss and the stable rise time is promising for vision restoration therapies. However, the 2-fold reduction in the decay constant of the calcium response indicates that abnormal dynamics develop in the inner retina weeks after PR loss.

# Introduction

In diseases such as age-related macular degeneration and retinitis pigmentosa, photoreceptor cell death leads to irreversible vision loss. A range of vision restoration therapies are currently under development that aim to restore light sensitivity to the retina including optogenetic and, optoelectronic prostheses, and regenerative stem cell transplantation. However, all current vision restoration approaches that act at the retinal level require reactivation of the remaining retinal architecture to restore interpretable signals to the brain. These approaches therefore require not only structural, but high quality functional preservation of the inner-retinal cells following extended periods of vision loss.

Previous studies have demonstrated RGC survival following long-term photoreceptor loss both in mice[1,2] and primates[3–5]. While RGCs may survive for many years in cases of advanced retinal degeneration, there is evidence of cytopathologies and ultrastructural 'whorls' suggestive of organelle stress[6]. Physiological studies of RGCs in murine models of retinal degeneration report increased spontaneous activity, including higher oscillatory firing frequency and an increased sustained firing rate[7–10]. In primates, structural rewiring of the midget pathway has been reported following acute photoreceptor loss[11], but whether physiological changes occur in primate RGCs is unknown. If altered physiology does develop in primate RGCs in the absence of photoreceptor input, this may compromise the efficacy of vision restoration therapies. In this study, we investigated the functional preservation of RGCs after photoreceptor ablation with the aim of understanding how deafferented RGCs in the fovea respond to restored activity as a function of time after photoreceptor loss.

In humans, high acuity vision is mediated by the specialized anatomy and physiology of the fovea. Previous attempts to restore vision in patients with retinal prostheses have been most successful when the fovea is stimulated[12], and optimizing therapies at this location may provide a pathway toward improved visual outcomes[13,14]. Due to the similarity in foveal anatomy, physiology, and downstream processing, the non-human primate model is the gold standard animal model for preclinical development of vision restoration therapies. While naturally occurring retinal degeneration models have recently been identified in NHPs[15,16], they are not widely available at present. Here we adopt a photoreceptor ablation approach using a femtosecond laser exposure delivered through an adaptive optics scanning laser ophthalmoscope (AOSLO). This results in acute photoreceptor loss that is laterally and axially confined[17].

In the absence of photoreceptors, investigating the functional status of deafferented RGCs requires an alternative stimulation approach. While electrical stimulation of RGCs in retinal explants is possible, longitudinal monitoring of responses in these preparations can't be performed without sacrificing animals at each time point. In this study we take an *in vivo* approach, optically stimulating and recording from deafferented RGCs for weeks, months, and years after photoreceptor ablation by combining adaptive optics (AOSLO) with the optogenetic actuator ChrimsonR and the calcium indicator GCaMP6s. Calcium imaging with AOSLO

enables a longitudinal readout of the intracellular calcium response to stimulation at the cellular scale *in vivo*[18]. In this study, we presented a brief optogenetic stimulus focused onto deafferented RGCs, and recorded optogenetic mediated calcium rise and decay in the weeks and years following photoreceptor loss.

## Methods

**Animal Care**

The primates in this study were housed in an AAALAC-approved facility, where the animals were provided with free access to water and nutritious chow together with various treats and vegetables. Primates were provided with daily enrichment including 2-4 pieces of manipulata such as mirrors and puzzle feeders rotating between animals, daily movies and music, and rotating access to larger free roaming space with swings and elevated perches. A team of 4 full-time veterinarians, 5 veterinary technicians, and an animal behavior specialist monitored animal health, body condition and checked for any signs of discomfort daily. Subjects recieves ambient diffused illumination during daytime (equivalent to 620 nm 0.05mW at pupil plane), significantly lower than Chrimson activation threshold[19], thus deafferented RGCs were only stimulated during the optogenetic trials. This study was performed in strict accordance with the Association for Research in Vision and Ophthalmology (ARVO) statement for use of animals, and the recommendations in the Guide for Care and Use of Laboratory Animals of the National Institutes of Health. All procedures were approved by the University of Rochester Committee in animal resources (PHS Assurance No. 1160743209A1).

**Co-expression of GCaMP6s and td-Tomato labeled ChrimsonR**

AAV2-CAG-tdTomato-ChrimsonR and AAV2-CAG-GCaMP6s were synthesized by the University of Pennsylvania vector core and were intravitreally injected into the left eye of one female *Macaca fascicularis* (M1) 65 weeks before recording, the right eye of one male 11 (M2) weeks before recording, and the right eye of a male *Macaca fascicularis* (M3) 164 weeks before euthanasia. Before vector injection, the ocular tissues were sterilized with 50% diluted Betadine and a lid speculum was placed. The viral vector was injected into the middle of the vitreous from a location ~3 mm behind the limbus using a 30-gauge needle with a tuberculin syringe. Neutralizing antibodies to AAV2 were measured in serum prior to purchase of monkeys and are expressed as the maximum serum dilution at which an antibody response is seen. For both NHPs no antibody response was seen at the minimal dilution, 1:5. Both animals received daily subcutaneous cyclosporine A injections to suppress the immune system 1-4 weeks prior to and after injection of the viral vectors. A starting dose of 6 mg/kg was titrated into a therapeutic range of 150–200 ng/mL by monitoring blood trough levels. Once the desired blood trough level was achieved, that dose was maintained for several months after injection of the viral vector. Signs of intraocular inflammation were monitored by slit lamp examination, fundus photography, and observation of the pupillary response.

**Photoreceptor Ablation by ultrafast laser exposure**

A patch of photoreceptors (1.2° × 0.9° subject M1, 0.9° × 0.8° subject M2 1st ablation, 2.7° × 2.5° subject M2 2nd ablation) was ablated in the central fovea by 106 ms exposure to an ultrafast pulsed, 730 nm laser focused onto the photoreceptor layer (PRL) using the AOSLO (average power of 4.48 Wcm$^{-2}$ and a repetition rate of 80 MHz). Studies by Dhakal *et al.* show that this approach creates damage to the PRL that is both axially and laterally confined. Furthermore as the photoreceptors in the fovea are laterally displaced from the RGCs they drive, the deafferented RGCs under study here are never directly exposed to the high intensity laser during PRL ablation. Femtosecond laser ablation has previously been used as a model of vision loss in retinal photoreceptor transplantation studies[20] and to demonstrate optogenetic restoration of RGC activity [5,18].

**Fundus imaging/OCT/cSLO Imaging**

Imaging was performed with a conventional scanning light ophthalmoscope (Heidelberg specialist), OCT, and a fundus camera prior and after PRL ablation weekly. A fundus camera (Topcon TRC 50ex) with customized fluorescence filters for GCaMP imaging (excitation 466/40 nm, emission 520/28) and tdTomato imaging (excitation 549/25 nm and emission 586/20 nm) were used to assess the fluorescence of GCaMP6s and tdTomato-ChrimsonR independently. The axial impact of laser exposure was monitored with OCT.

**Adaptive optics calcium imaging**

Foveal RGCs were stimulated and recorded through an AOSLO system described in Gray *et al.* To summarize, a Shack-Hartmann wavefront sensor and a deformable mirror at subject pupil plane detect and compensate ocular aberrations in a closed-loop correction system using 847 nm laser diode (Thorlabs) in an SLO. A 796 nm superluminescent-diode laser source (Superlum) was focused on the photoreceptor mosaic layer, and reflectance images collected through the 25.6 Hz adaptive optics SLO with a 2-airy disk, confocal pinhole by a photomultiplier tube (Hamamatsu). GCaMP6s was excited by focusing a 488 nm laser source (Qioptics) at the RGC layer over an an area of 398x398 um$^2$ and emitted photons detected in 517/20 nm band using an 8-airy disk pinhole to maximize signal collection.

**Visual stimulation**

Both optogenetic and photoreceptor stimulation was delivered through the 25 Hz AOSLO system. Both types of trials began with a period of adaptation to the imaging light (67.5 sec for the optogenetic trials and 60 sec for the photoreceptor trials). To drive the optogenetic actuator ChrimsonR, 640 nm laser 0.5 sec single flash stimulus was focused onto the RGC layer, (mean luminance 21.37 mw/cm$^2$, 205x205 um$^2$). The calcium responses are recorded for 22.5 s following stimulation to observe the calcium rise and decay. To examine photoreceptor-mediated responses, a 561 nm, 0.025 Hz square wave uniform flicker stimulus with a mean luminance of 0.25 mw/cm$^2$ was delivered to central photoreceptors for 80 seconds while GCaMP recordings were made from the laterally displaced foveal RGCs they drive.

**Experimental Design**

To ensure sufficient GCaMP and ChrimsonR expressions in intact RGCs before ablation, the photoreceptor and optogenetic stimulation test were performed twice in the 1 week before PRL ablation. In the 1-2 weeks following the PRL ablation, a photoreceptor stimulation test was delivered twice to ensure complete loss of photoreceptor inputs in recorded RGCs. Optogenetic stimulation test was also performed and recorded both in the recent deafferented area and a long-term deafferented area at the same eccentricity on the opposite side of the RGC ring 5-6 times over the 10 week period following PRL ablation in subject M1 and M2. Control experiments included only presenting 488 nm stimulation to confirm the response was optogenetically driven, and a ChrimsonR stimulus only trial to ensure no bleed through of the 640 nm stimulus into the detection channel.

**Data analysis**

To correct for eye motion, *in vivo* fluorescence videos containing the calcium imaging data were co-registered to simultaneously acquired, higher signal to noise IR reflectance images of the photoreceptor mosaic. For each imaging location, a reference frame was used to perform a strip-based cross-correlation[21–23] to remove translation and rotation. Motion corrected reflectance and fluorescence videos were summed over time to generate higher signal to noise reflectance images of the photoreceptor ablation site and fluorescence images of intact and deafferented RGCs for manual segmentation. To ensure the same group of cells were tracked longitudinally, fluorescence GCaMP images of RGCs over weeks are registered to week 1 after ablation before segmentation.

To extract GCaMP recordings for each RGC, the segmentation mask was applied to the motion-corrected fluorescence videos and the initial 67.5s adaptation period was removed. The mean signal within each cell mask was computed for a single frame (subject M1 n=96, subject M2 n=109).

In optogenetic stimulation trials, each trace was separated into a rising phase and a decay phase based on the frame containing the maximum fluorescence intensity. Rise time was calculated as the time between the stimulus onset and the frame containing the maximum fluorescence intensity. To determine the decay constant for single traces the data was first normalized by subtracting the average baseline fluorescence in the 100 frames prior to stimulus onset and dividing by the maximum fluorescence intensity. The resulting traces were fitted with an exponential function using a least-square approach. The mean responses presented in figures 2 and 3 were created by taking the mean of the normalized single cell traces from all segmented RGCs for each location.

The response magnitude was quantified in the following ways:

**Absolute response magnitude** To report the response magnitude excluding changes in signal baseline, $F_{LT}$-$F0_{LT}$ & $F_{ST}$-$F0_{ST}$ indexes are the difference between the maximum fluorescence signal (F) in a single cell and the pre-stimulus baseline (F0).

**Normalized response magnitude** Then the normalized response magnitude is calculated by dividing single RGC $F_{ST}$-$F0_{ST}$ indexes by the mean of $F_{LT}$-$F0_{LT}$ recorded in the same imaging session to exclude changes in eye quality or imaging condition over weeks, and the sequence of experiments between sessions is not varied over the sessions.

**Normalized pre-stimulus baseline** Normalized F0 value is acquired by dividing single short-term deafferented RGC calcium floor level by the mean of long-term deafferented RGC calcium floor level recorded within the same imaging session, the sequence of experiments between sessions is not varied over the sessions.

**The sensitivity index** ((F-F0) / std(F0)) is measured by dividing individual short-term deafferented RGC's F-F0 index by the standard deviation F0 value, F-F0 & F0 are calculated with the same method described above.

For photoreceptor stimulation trials, the 60-s adaptation period is removed from the registered video, and the mean signal within each cell mask is computed for each frame of the recording period. The resulting fluorescence traces were multiplied by a Hann windowing function to remove ringing artifacts and Fourier transformed into the frequency domain as described previously[18].

## Light Safety
The total light exposure at each retinal location was calculated for each recording session, with all light sources taken into account (488 nm at 0.71 mw cm$^{-2}$, 796 nm at 1.30 mw cm$^{-2}$, and 843 nm at 0.19 mw cm$^{-2}$), optogenetic stimulation (640 nm at 21.37 mw cm$^{-2}$), and photoreceptor stimulation (561 nm at 0.25 mw cm$^{-2}$). Overall light exposure at a single retinal location did not exceed the American National Standard Institute (American National Standard for the Safe Use of Lasers ANSI Z136.1–2007) maximum permissible exposure of the human retina scaled by the ratio of numerical aperture difference between human and macaque (Photoreceptor ablation session excluded). The power of each light source was measured before the imaging session at the pupil plane, and accumulation of laser exposure between sessions is not considered since all sessions were separated by at least six days.

## Histology
Subject 3 was euthanized with an intravenous injection of Euthasol and perfused with 1 L of heparinized saline for 23 min and 1 L of 4% paraformaldehyde for 7 min. After enucleation, the anterior segment was removed and the posterior segment was immersion-fixed in 4% paraformaldehyde for 90 min. After fixation, the posterior segment was washed for 5 min in PBS and cryoprotected in 10%, 20%, and 30% sucrose. Retinal pieces were embedded in Cryogel embedding compound (Leica) and frozen in isopentane cooled in liquid nitrogen. 20-μm thick cryosections were collected and dried for 60 min at 35°C on a slide warmer before storage at −80°C. To assess the cytoplasmic to nuclear ratio of GCaMP6s expression in RGCs, nuclei were counterstained with Hoescht 33342. Samples were imaged on a laser scanning confocal microscope (Zeiss LSM 880) with the 405- and 561-laser lines using a Zeiss Plan-Apochromat 60×/1.4 N.A. oil immersion objective. Tiled image stacks were stitched in Image J software. To

measure the cytoplasmic to nuclear ratio for GCaMP6s, line profiles were drawn to transect the center of each RGC in the image stack. Intensity profile plots of GCaMP6s and Hoescht fluorescence were exported to Igor Pro 9 (Wavemetrics) for further analysis. Intensity measurements were background subtracted using areas of the image without GCaMP6s fluorescence.

# Results

## Localized photoreceptor ablation in primate fovea deafferents RGCs allowing characterization of optogenetic mediated calcium dynamics

Intravitreal co-injection of the viral vectors AAV2-CAG-GCaMP6s and AAV2-CAG-tdTomato:ChrimsonR resulted in a ring of fluorescent RGCs expressing the calcium indicator (Fig 1A and Supplementary Fig 1A, 1B) and the optogenetic actuator tdTomato-ChrimsonR in all 3 animals (Fig 1B and Supplementary Fig 1C, 1D) as observed in previous studies[5,18].

AOSLO NIR reflectance imaging of the cone photoreceptor mosaic *in vivo* confirms photoreceptor damage following laser exposure (Fig 1C). Optical coherence tomography (OCT) shows the absence of the foveal photoreceptor layer (PRL) post exposure (Supp.1X). Figure 1.D shows GCaMP fluorescence from foveal RGCs one week after PRL ablation (9 weeks after the initial injection of AAV2-CAG-GCaMP6s and AAV2-CAG-ChrimsonR-tdTomato). As previously reported[5,18] fluorescence fundus imaging shows a reduction in fluorescence from the RGC ring that is spatially consistent with the ablation of photoreceptors in the temporal hemifield of the central fovea. The loss of photoreceptor response is confirmed by presenting a 0.025 Hz square wave flicker stimulus to the central fovea. Figure 1E shows 0.025Hz GCaMP responses from RGCs prior to laser exposure and the absence of these signals from the same RGCs post exposure as anticipated.

To assess RGC calcium dynamics in these deafferented RGCs we presented a 0.5s, 640 nm, 0.9 mW stimulus focused at the ganglion cell layer through AOSLO to optogenetically stimulate RGCs. The optogenetic mediated calcium response leads to an increase of GCaMP fluorescence and then gradual decay. Figure 1F shows the optogenetic mediated calcium response recorded from deafferented RGCs 1 week after PRL ablation.

**Fig. 1: Optogenetic mediated calcium response triggered in deafferented primate foveal RGCs.** (A) Fundus fluorescence image of td-Tomato ChrimsonR expression 7 weeks after injection of the viral vector and 1 week prior to photoreceptor ablation. (B) AOSLO NIR reflectance image of photoreceptor ablation site 1-week following ablation. (C) Fundus fluorescence image of GCaMP6s expression 7 weeks after injection of viral vector and 1 week prior to photoreceptor ablation. (D) Fundus fluorescence image of GCaMP nine weeks after injection of viral vector and 1-week post photoreceptor ablation. The dark region of the RGC ring has photoreceptor inputs ablated and the activity dependent fluorescence is correspondingly reduced. (E) Clear 0.025Hz GCaMP response is observed from RGCs on the right side of the RGC ring before photoreceptor ablation, while no GCaMP response is observed from the same group of RGCs following photoreceptor ablation, supporting loss of photoreceptor inputs. (F) Optogenetic mediated GCaMP response to a 0.5 s 640 nm stimulation 1 week following laser ablation. Stimulus duration is shown in the red bar, AOSLO deafferented RGC image with low fluorescence expression in top right. All scale bars are 100 μm, and all data were acquired from subject M1.

## Calcium responses to optogenetic stimulation decay more rapidly in RGCs within weeks of PRL ablation

The same stimulation and recording paradigm was used to interrogate the calcium rise and decay in a patch of RGCs deafferented for 1 week (nasal: short-term deafferented), and also in an eccentricity matched control patch of RGCs deafferented for 2 yrs (temporal: long-term deafferented). By recording from these two regions during each session we were able to use

long-term deafferented RGC responses as a control group to assess the stability of the imaging platform.

The NIR reflectance AOLSO image of the photoreceptor mosaic and OCT data show the two regions of photoreceptor ablation (Figure 2 A-B) in subject M2. The decreased activity dependent RGC GCaMP fluorescence in Figure 2D reflects the loss of photoreceptor input, consistent with the previous study[5]. Deafferentation was confirmed at the 1 week timepoint post ablation on the nasal side of the RGC ring, with no response to a 0.45 Hz photoreceptor stimulation detected (Figure 2D). Figure 2G-H shows the mean decay constant of the calcium response decreased 2.1 fold (from 2.5±0.5 s to 1.2±0.2 s SD) in the 8 weeks post PR ablation ($p$<0.001, paired $t$-test, n=109), with 87% of this decrease occurring within the first 5 weeks. The decay constant did not decrease further from week 8 to 10, and the calcium rise and decay speed stabilized from 101-122 weeks following photoreceptor loss (Figure 2E-F).

**Fig. 2: Foveal RGCs show faster calcium decay in the five weeks following PRL ablation in subject M2.** (A) OCT scan through the foveal region shows two localized regions of photoreceptor loss. (B) AOSLO NIR reflectance image of two photoreceptor ablation sites 122 weeks following the first ablation (left) and 10 weeks following the second ablation (right). (C) Fundus fluorescence image of GCaMP6s fluorescence 1 week following the second PRL ablation. (C) Fundus fluorescence image of GCaMP6s expression 7 weeks after injection of viral vector and 1 week prior to photoreceptor ablation. Two dark regions of the RGC

ring have photoreceptor inputs ablated on the nasal and the temporal side of the RGC ring, reflecting the laser PRL ablation performed 2 years and 1 week before the study. The nasal side corresponds to the short-term deafferented RGCs, and the temporal side corresponds to the long-term deafferented RGCs. (D) A 0.45 Hz GCaMP response is observed in RGCs before photoreceptor ablation, while no GCaMP response is observed from the same group of RGCs following photoreceptor ablation. (E) Mean GCaMP response to optogenetic stimulation from 109 long-term deafferented RGCs from weeks 1-10. (F) GCaMP fluorescence decay constants from 109 individual long-term deafferented RGCs from weeks 1-10. (G) Mean GCaMP response to optogenetic stimulation from 109 short-term deafferented RGCs from weeks 1-10. (H) GCaMP fluorescence decay constants from 109 individual short-term deafferented RGCs from weeks 1-10.

## Stable calcium decay in RGCs with intact photoreceptors and in RGCs prior to PRL ablation

The experimental paradigm was repeated in a second subject (M1) this time comparing the optogenetic response in deafferented RGCs to the optogenetic response in RGCs with intact photoreceptors. Photoreceptor-mediated calcium responses to a 0.1 Hz flickering stimulus incident on foveal photoreceptors were recorded 1 week prior to, and 5 weeks post PRL ablation in a single field of view. (Figure 3A-B). Laser ablation to subject M1 PRL layers shown in removes photoreceptor inputs in only half of the RGC recording area at lower eccentricity, shown in PRL reflectance image in Figure 3C. After ablation, periodic RGC responses to photoreceptor stimulation were absent in the deafferented zone (Figure 3D-E). Similar to animal M2, the mean decay constant of the optogenetically-evoked calcium responses in the deafferented RGCs decreased 1.5 fold (from 1.6±0.5 s to 0.6±0.3 s SD) in the 10 weeks post PR ablation ($p<0.001$, paired $t$-test, Figure 3F). In comparison, RGCs that had intact photoreceptors that were recorded in the same field of view showed relatively stable decay kinetics over the 10 week period . Use of a spatially-limited optogenetic stimulus in the fovea ensured that the calcium responses from these cells were not photoreceptor mediated (Figure 3G, Appendix 1). Optogenetic stimulation of RGCs with intact photoreceptors in this region and in the deafferented region 1 week prior to photoreceptor ablation produced a stable decay constant record (2.2±0.5 s), confirming that these results are not explained by a temporary increase in decay constant relating to e.g inflammation post ablation.

**Fig. 3: Calcium response to PR stimulation and optogenetic decay constant in RGCs with photoreceptor loss and intact photoreceptor in a second primate.** (A) AOSLO GCaMP fluorescence image shows two study regions of RGCs with normal GCaMP expression level at higher eccentricity (blue) & RGCs with reduced GCaMP expression level at lower eccentricity (red) 1 week before photoreceptor ablation. (B) RGC photoreceptor mediated GCaMP fluorescence responses to 0.1 Hz on-off stimulation in deafferented (blue) and disturbed (red) areas 1 week before photoreceptor ablation. (C) AOSLO GCaMP fluorescence image shows two study regions of RGCs with normal GCaMP expression level at higher eccentricity (blue) & RGCs with reduced GCaMP expression level at lower eccentricity (red) 5 weeks post photoreceptor ablation. (D) AOSLO NIR reflectance image of photoreceptor ablation sites 5 weeks following the photoreceptor ablation. (E) RGC photoreceptor mediated GCaMP fluorescence responses to 0.1 Hz on-off stimulation in deafferented (blue) and disturbed (red) areas 5 weeks post photoreceptor ablation. (F) Normalized optogenetic mediated GCaMP fluorescence intensity was recorded from 129 short-term deafferented RGCs from 1 week before PRL ablation to 10 weeks post-ablation in subject M1.

(G) Optogenetic mediated GCaMP fluorescence decay constant from 129 short-term deafferented RGCs from 1 week before PRL ablation to 10 weeks post-ablation. (H) Optogenetic mediated GCaMP fluorescence decay constant from 92 intact RGCs from 1 week before PRL ablation to 10 weeks post-ablation. All scale bars are 100 μm, and all data are acquired from subject M1.

## The optogenetic mediated calcium rise time is unaffected by deafferentation

The mean time to peak calcium response reflects the speed of the optogenetic mediated calcium influx. In animal M1, the mean rise time did not change significantly prior to and post PRL ablation (Week -1, 1.47±0.33s SD) & (Week 5, 1.44±0.14s SD). In animal M2, the mean rise time also did not differ significantly between the long-term (1.43±0.17s SD) and the short-term (1.45±0.30s SD) deafferented RGCs during the 10-week period (unpaired $t$-test, $p$=0.05, n=654) (Figure 4A-B). These data suggest that photoreceptor ablation does not alter the rise time of calcium responses in downstream RGCs.

**Fig. 4: Optogenetic mediated calcium rise time in deafferented and intact RGCs.** (A) Optogenetic mediated calcium rise time in response to a 0.5 second 25Hz pulsed 640 nm stimulation in non deafferented RGCs (n=182) and deafferented RGCs (n=257) in subject M1 . Standard deviations of the RGC rise time are labeled with gray bars, and the mean of RGC rise times are shown in black dashed lines. Each data point shows a single RGC rise time recorded within an imaging session. (B) Optogenetic mediated calcium rise time in response to a 0.5 second pulsed 640 nm stimulation in RGCs deafferented for 2

years (long-term, n=107), and subject 2 RGCs deafferented for 1-10 weeks (short-term, n=106) in animal M2. Each data point shows the mean RGC rise time collected from a single cell over the six imaging sessions.

## An optogenetic mediated calcium response can be evoked in deafferented RGCs 2.8 years following viral transduction and 2.3 years after photoreceptor ablation

In subject M2, robust optogenetic mediated responses were recorded 2.8 years post-injection of the viral vector carrying the ChrimsonR transgene. The RGCs in this area were deafferented for over 2.3 years at the time of recording. The decay constant in long-term deafferented RGCs was 1.6 ± 0.2s S.D.. The calcium rise time in these long-term deafferented RGCs (1.43 ± 0.10s) is not significantly different from RGCs with intact photoreceptors, indicating the speed of the optogenetic mediated calcium response is not impacted by long-term photoreceptor loss.

**Fig. 5: Mean optogenetic mediated calcium response recorded from 109 RGCs deafferented over 2.3 years.** (A) Optogenetic mediated GCaMP response to a 0.5 second 640 nm stimulation 2.3 years following photoreceptor ablation and 2.8 years following initial vector injection, the stimulation duration is shown by the red bar.

## The decay constant is independent of response amplitude, stimulus power, and chronic expression of GCaMP6s

To assess the impact of potentially varying optical parameters, response magnitude and protein expression on the calcium decay constant, we calculated the sensitivity index of responses and the normalized response magnitude to control for these parameters. In addition, we performed control experiments using a range of optogenetic stimulus powers and conducted a histological investigation of intact and deafferented RGCs in subject M3 with chronic GCaMP6s expression.

The sensitivity index scales the absolute response magnitude by the standard deviation of the baseline fluorescence, providing a discriminability metric for the response amplitude independent of fluorescence intensity. The division controls for  parameters that may potentially

vary between sessions such as the clarity of the anterior optics, system alignment, AO performance, pinhole position and the level of GCaMP expression. Figure 6A reveals the sensitivity index of responses in deafferented RGCs is stable during the 10 week recording period, suggesting changes in the magnitude of the response do not account for the reported reduction in the decay constant.

Figure 6B shows the decay constant from deafferented RGCs under optogenetic stimulation powers ranging from 0.8 - 1 mW. The decay constant in deafferented RGCs shows no linear trend with stimulus power, suggesting the changes we observed in the speed of calcium decay are not linked to the power of stimulus or the level of optogenetic actuator expression changing over time. Additional analysis of how the absolute response magnitude varies is presented in Figure S2.

**Fig. 6: Control experiments in insensitivity index, optogenetic stimulus power, GCaMP6s Nucleus expression level.** (A) Optogenetic sensitivity index

for short-term deafferented RGCs in the 10 weeks following photoreceptor ablation. (B) Optogenetic mediated calcium response decay constant in deafferented RGCs with three stimulation powers. Data collected from 109 short-term deafferented RGCs and 109 long-term deafferented RGCs in subject M2. (C,D), Confocal image (maximal z-projection) showing GCaMP6s expression (green) in a non-deafferented (normal, C) and deafferented (D) region of the same retina. The foveal slope region is indicated. ROIs are shown enlarged on the right with and without a nuclear counterstain (Hoescht 33342). E, Violin/bar plots showing the nuclear:cytosolic ratio of GCaMP6s for RGCs in normal (n=152 cells) and deafferented (n=140 cells) retinal regions from subject 3. Horizontal line shows median, whiskers show min/max. Dots show data from individual cells. Data are pooled from two different sections of the same retina. Statistical comparison was with an unpaired, two-tailed t-test.

Previous studies in mouse cortical neurons have shown that long-term expression of GCaMP3 can lead to nuclear accumulation and prolongation of the decay of calcium signals, suggesting altered physiology [24]. However, recent longitudinal studies with GCaMP7s in rodents showed no abnormal light-evoked activity in RGCs following long-term expression of GCaMP7s[25]. To investigate whether changes in GCaMP6s localization could account for the observed changes in calcium dynamics, we quantified the nuclear to cytosolic (N/C) ratio of GCaMP6s in deafferented and intact RGCs in subject M3 (Figure 6E-F). In prior studies, a higher GCaMP6s N/C ratio was linked to slower decay of calcium signals. We found a significantly higher N/C ratio in deafferented compared to intact RGCs (0.44 ± 0.16 vs 0.31 ± 0.14 S.D.) . These findings suggest that the faster decay time constant of calcium signals in deafferented RGCs cannot be explained by nuclear accumulation associated with GCaMP toxicity.

## Discussion

By co-expressing the calcium indicator GCaMP6s and optogenetic actuator ChrimsonR in vivo, we have demonstrated the development of abnormal calcium dynamics in adult primate foveal RGCs just weeks after photoreceptor loss. The timescale over which these changes were observed is similar to that associated with the development of retinal hyperactivity in rodent models[7–10]. A 1.5-2-fold increase in the speed of calcium clearance was observed within 5 weeks in subjects M1 and M2 respectively. The decay constant in short-term deafferented RGCs recorded from subject M2 also approached the long-term deafferented RGC responses after week 8. Data from both subjects suggest that physiological changes occurred within weeks of photoreceptor loss. While there was an initial difference in calcium decay constant between the two subjects, it is important to note that both deafferented RGC groups approach a similar decay constant of 1.3 s and 1 s by 10 weeks after photoreceptor loss. The initial difference between the two subjects may be related to the substantial age difference between these animals (M1 7 years old, M1 X years old). Data from post-mortem human retina has suggested that exposure to age-related stressors may generate differences in physiology[26].

The stable decay constant recording in intact RGCs in subject M1 indicates the abnormal calcium clearance observed in both subjects was not an immediate RGC response to PRL ablation or optogenetic actuator stimulation. We also demonstrated that with adaptive optics with active closed-loop tracking, the optogenetic stimulation of displaced RGCs will not trigger responses from nearby cells with photoreceptor crosstalk, confirming the initial longer decay constant prior to PRL ablationConsistent with the reduction of GCaMP6s and tdTomato-ChimsonR fluorescence intensity in deafferented RGCs in the weeks of PRL ablation reported by McGregor *et al.*, the normalized response magnitude shows a significant reduction within 1 week following photoreceptor ablation. The optogenetic response sensitivity index, various optogenetic stimulus power, and histological assessment of chroicical GCaMP6s experssion suggest the response magnitude and GCaMP expression level do not directly account for gradual changes in the decay constant.

While this is the first report of abnormal calcium dynamics in primate retina, studies in a murine model of glaucoma have also revealed abnormal calcium dynamics developing in RGCs following axon crush, although interestingly the decay constant increases after injury[27]. This is different from the more rapid decay observed in this study, but there are a number of important model and species differences to note. In the glaucoma model, lengthening of the decay constant appears to precede rapid cell death[27], but in the photoreceptor ablation model the inner retina remains structurally intact despite many years of photoreceptor loss. It is possible that the shorter decay constant we observe is the consequence of a protective adaptation to higher levels of intracellular calcium generated by hyperactivity. Studies in other parts of the primate visual system have observed abnormal calcium homeostasis following ablation[28]. It is also possible that the altered calcium clearance develops following photoreceptor loss to maintain calcium homeostasis. Additionally. primate foveal RGCs are largely midget ganglion cells, which are absent in mice. This study represents the first observations of this phenomenon and more work is required to understand the mechanism giving rise to this effect, and whether RGC types are differentially affected. To fully investigate the impact of physiological changes on restored vision, psychophysical tests are necessary.

Although physiological changes are observed in RGC calcium dynamics following photoreceptor loss, it is important to note that the optogenetic mediated calcium rise time was stable in intact and deafferented RGCs in both subjects and across all time points. Furthermore, robust optogenetic mediated responses are triggered in RGCs even 2.3 years following photoreceptor loss, and 2.8 years after viral transduction of the optogenetic actuator. This is a promising result with current clinical trials having reported optogenetic mediated light sensitivity remaining 1.6 years after administration of the therapy in patients[29].

# Acknowledgements

We thank Bill Merigan and David Williams for accommodating access to the 1P and 2P AOSLOs and sharing animal subjects. We thank Jacqueline Gayet for technical assistance with histology. We thank the vector core at the Perelman School of Medicine, University of

Pennsylvania and the Genetically-Encoded Neuronal Indicator and Effector (GENIE) Project and the Janelia Research Campus of the Howard Hughes Medical Institute, specifically Vivek Jayaraman, PhD, Douglas S. Kim, PhD, Loren L. Looger, PhD, and Karel Svoboda, PhD. This work is supported by National Institutes of Health through grants through grants U24 EY033275 (to J.E.M.), and NIH EY024265 (to T.P.) and P30 Core Grants NIH EY001319 (University of Rochester) and NIH EY003176 (UC Berkeley). Additional funding was provided by the Steven E. Feldon Scholarship to J.E.M. and through an unrestricted grant to the Flaum Eye Institute from Research to Prevent Blindness (https://www.rpbusa.org/rpb/grants-and-research/grants/rpb-unrestricted-grants/rpb-unrestricted-grants/). Confocal imaging was conducted in the CRL Molecular Imaging Center at UC Berkeley (RRID:SCR_017852), supported by the Helen Wills Neuroscience Institute.

# Reference:

1. Margolis DJ, Newkirk G, Euler T, Detwiler PB. Functional Stability of Retinal Ganglion Cells after Degeneration-Induced Changes in Synaptic Input. *J Neurosci*. 2008;28(25):6526. doi:10.1523/JNEUROSCI.1533-08.2008
2. Mazzoni F, Novelli E, Strettoi E. Retinal Ganglion Cells Survive and Maintain Normal Dendritic Morphology in a Mouse Model of Inherited Photoreceptor Degeneration. *J Neurosci*. 2008;28(52):14282. doi:10.1523/JNEUROSCI.4968-08.2008
3. Santos A, Humayun MS, de Juan E Jr, et al. Preservation of the Inner Retina in Retinitis Pigmentosa: A Morphometric Analysis. *Arch Ophthalmol*. 1997;115(4):511-515. doi:10.1001/archopht.1997.01100150513011
4. Jones BW, Pfeiffer RL, Ferrell WD, Watt CB, Marmor M, Marc RE. Retinal remodeling in human retinitis pigmentosa. *Spec Issue Cell Remodel Retina Opt Nerve Inj Dis*. 2016;150:149-165. doi:10.1016/j.exer.2016.03.018
5. McGregor JE, Kunala K, Xu Z, et al. Optogenetic therapy restores retinal activity in primate for at least a year following photoreceptor ablation. *Mol Ther*. 2022;30(3):1315-1328. doi:10.1016/j.ymthe.2021.09.014
6. Pfeiffer RL, Marc RE, Jones BW. Persistent remodeling and neurodegeneration in late-stage retinal degeneration. *Prog Retin Eye Res*. 2020;74:100771. doi:10.1016/j.preteyeres.2019.07.004
7. Stasheff SF. Emergence of Sustained Spontaneous Hyperactivity and Temporary Preservation of off Responses in Ganglion Cells of the Retinal Degeneration (rd1) Mouse. *J Neurophysiol*. 2008;99(3):1408-1421. doi:10.1152/jn.00144.2007
8. Kolomiets B, Dubus E, Simonutti M, Rosolen S, Sahel JA, Picaud S. Late histological and functional changes in the P23H rat retina after photoreceptor loss. *Neurobiol Dis*. 2010;38(1):47-58. doi:10.1016/j.nbd.2009.12.025
9. Telias M, Denlinger B, Helft Z, Thornton C, Beckwith-Cohen B, Kramer RH. Retinoic Acid Induces Hyperactivity, and Blocking Its Receptor Unmasks Light Responses and Augments Vision in Retinal Degeneration. *Neuron*. 2019;102(3):574-586.e5. doi:10.1016/j.neuron.2019.02.015
10. Denlinger B, Helft Z, Telias M, Lorach H, Palanker D, Kramer RH. Local photoreceptor degeneration causes local pathophysiological remodeling of retinal neurons. *JCI Insight*. 2020;5(2):132114. doi:10.1172/jci.insight.132114
11. Akiba R, Boniec S, Uyama H, et al. Rewiring of the adult midget pathway in a primate model of acute photoreceptor loss. *Invest Ophthalmol Vis Sci*. 2022;63(7):3132-3132.
12. Stingl K, Bartz-Schmidt KU, Besch D, et al. Subretinal Visual Implant Alpha IMS – Clinical

trial interim report. *Sight Restor Prosthet Optogenetics Gene Ther*. 2015;111:149-160. doi:10.1016/j.visres.2015.03.001

13. Damle S, Lo YH, Freeman WR. High Visual Acuity Retinal Prosthesis: Understanding Limitations and Advancements Toward Functional Prosthetic Vision. *RETINA*. 2017;37(8). https://journals.lww.com/retinajournal/Fulltext/2017/08000/High_Visual_Acuity_Retinal_Prosthesis_.1.aspx

14. McGregor JE, Williams DR, Merigan WH. Functional Assessment of Vision Restoration. In: Bowes Rickman C, Grimm C, Anderson RE, Ash JD, LaVail MM, Hollyfield JG, eds. *Retinal Degenerative Diseases*. Springer International Publishing; 2019:145-149.

15. Moshiri A, Chen R, Kim S, et al. A nonhuman primate model of inherited retinal disease. *J Clin Invest*. 2019;129(2):863-874. doi:10.1172/JCI123980

16. Peterson SM, McGill TJ, Puthussery T, et al. Bardet-Biedl Syndrome in rhesus macaques: A nonhuman primate model of retinitis pigmentosa. *Exp Eye Res*. 2019;189:107825. doi:10.1016/j.exer.2019.107825

17. Dhakal KR, Walters S, McGregor JE, et al. Localized Photoreceptor Ablation Using Femtosecond Pulses Focused With Adaptive Optics. *Transl Vis Sci Technol*. 2020;9(7):16-16. doi:10.1167/tvst.9.7.16

18. McGregor JE, Godat T, Dhakal KR, et al. Optogenetic restoration of retinal ganglion cell activity in the living primate. *Nat Commun*. 2020;11(1):1703. doi:10.1038/s41467-020-15317-6

19. Klapoetke NC, Murata Y, Kim SS, et al. Independent optical excitation of distinct neural populations. *Nat Methods*. 2014;11(3):338-346. doi:10.1038/nmeth.2836

20. Aboualizadeh E, Phillips MJ, McGregor JE, et al. Imaging Transplanted Photoreceptors in Living Nonhuman Primates with Single-Cell Resolution. *Stem Cell Rep*. 2020;15(2):482-497. doi:10.1016/j.stemcr.2020.06.019

21. Yang Q, Zhang J, Nozato K, et al. Closed-loop optical stabilization and digital image registration in adaptive optics scanning light ophthalmoscopy. *Biomed Opt Express*. 2014;5(9):3174-3191. doi:10.1364/BOE.5.003174

22. McGregor JE, Yin L, Yang Q, et al. Functional architecture of the foveola revealed in the living primate. *PLOS ONE*. 2018;13(11):e0207102. doi:10.1371/journal.pone.0207102

23. Hu X, Yang Q. Real-time correction of image rotation with adaptive optics scanning light ophthalmoscopy. *J Opt Soc Am A*. 2022;39(9):1663-1672. doi:10.1364/JOSAA.465889

24. Tian L, Hires SA, Mao T, et al. Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators. *Nat Methods*. 2009;6(12):875-881. doi:10.1038/nmeth.1398

25. Li L, Feng X, Fang F, et al. Longitudinal in vivo Ca2+ imaging reveals dynamic activity changes of diseased retinal ganglion cells at the single-cell level. *Proc Natl Acad Sci*. 2022;119(48):e2206829119. doi:10.1073/pnas.2206829119

26. Abbas F, Becker S, Jones BW, et al. Revival of light signalling in the postmortem mouse and human retina. *Nature*. 2022;606(7913):351-357. doi:10.1038/s41586-022-04709-x

27. Shiga Y, Rangel Olguin AG, Alarcon-Martinez L, et al. Light-evoked RGC calcium dynamics are altered in glaucoma: live imaging evidence of abnormal calcium clearance. *Invest Ophthalmol Vis Sci*. 2022;63(7):1850-1850.

28. Gutierrez C, Cusick CG. Effects of chronic monocular enucleation on calcium binding proteins calbindin-D28k and parvalbumin in the lateral geniculate nucleus of adult rhesus monkeys. *Brain Res*. 1994;651(1-2):300-310. doi:10.1016/0006-8993(94)90710-2

29. Sahel JA, Boulanger-Scemama E, Pagot C, et al. Partial recovery of visual function in a blind patient after optogenetic therapy. *Nat Med*. 2021;27(7):1223-1229. doi:10.1038/s41591-021-01351-4