Title

Vehicle-mounted cameras reveal negative impact of the Fukushima Daiichi

nuclear power plant accident on large bird abundance via paddy field

abandonment.

Author details

Nao Kumada[1], Keita Fukasawa[1], Akira Yoshioka[2], Naoe Tsuda[3], Hirofumi Ouchi[2]

[1] Biodiversity Division, National Institute for Environmental Studies, Tsukuba, Ibaraki,
Japan

[2] Fukushima Regional Collaborative Research Center, National Institute for
Environmental Studies, Miharu, Fukushima, Japan

[3] Center for Climate Change Adaptation, National Institute for Environmental Studies,
Tsukuba, Ibaraki, Japan

Corresponding author

Nao Kumada

Biodiversity Division, National Institute for Environmental Studies

16-1 Onogawa, Tsukuba City, Ibaraki Prefecture, Japan

Email: kumada.nao@nies.go.jp

Abstract

Farmland bird populations are declining due to farmland abandonment and agricultural

intensification. Effective conservation strategies require appropriate monitoring approaches,

including efficient and scalable survey methods. In this study, we developed a large-scale

monitoring method for herons and egrets (Ardeidae) using vehicle-mounted video cameras and

distance sampling models that incorporate location uncertainty. The survey was conducted in and

around the evacuation zone of the Fukushima Daiichi nuclear power plant accident. A total of

7,031 km of roadside video footage was recorded, covering 24.41 km² of farmland. Ardeidae were

observed only outside the evacuation zone and were entirely absent within it. Predicted abundance

differed greatly between areas inside (0.0279 ± 0.0307/km²) and outside (4.57 ± 5.36/km²) the

evacuation zone. Incorporating location uncertainty into the distance sampling model had little

effect on the estimates (4.57 ± 5.36 vs. 4.51 ± 5.29/km² with and without integrating location

uncertainty, respectively). This suggests that our video-based roadside survey method is robust to

location uncertainty in structured landscapes such as Japanese rice paddies. The accuracy may be

attributed to the study system, where levees and roads divide paddy fields, limiting potential error

in observer-target distances Our method can be applied to other open habitats, such as natural

grasslands and wetlands, especially in areas lacking artificial markers, by incorporating

measurement uncertainty into the model. This combination of roadside surveys with vehicle-

mounted cameras and distance sampling provides a practical and transferable approach for

monitoring large-bodied species in open landscapes, enhancing both the efficiency and spatial

coverage of biodiversity assessments.

Keywords

Abandoned paddy fields, Vehicle-mounted video camera, Agricultural birds, Distance sampling,

Roadside survey, UAV

Highlights

・Broad-scale farmland abandonment following the Fukushima Daiichi nuclear power plant

accident led to sharp declines in large farmland bird abundance.

・Efficient large-scale roadside surveys were conducted using vehicle-mounted video cameras.

・Ardeidae abundance was estimated using distance sampling models that incorporated location

uncertainty.

・The survey method is robust and applicable to open habitats such as farmlands, grasslands, and

wetlands.

1 Introduction

Population sizes of many farmland birds are declining due to agricultural intensification and the abandonment of cultivation (Fujioka, Don Lee, Kurechi, & Yoshida, 2010; Fuller et al., 1995; Siriwardena et al., 1998; Stanton, Morrissey, & Clark, 2018). To implement appropriate conservation strategies for farmland birds, information on their abundances and related environmental factors is essential. Various methods have been developed for monitoring farmland birds, including point count, transect count, territory mapping, and capture-recapture methods (Birrer et al., 2007; Chiron, Filippi-Codaccioni, Jiguet, & Devictor, 2010; Gillings & Fuller, 1998). These methods are often labor-intensive and costly to conduct over a large area. Cost-effective methods for monitoring are needed to better understand population patterns and environmental drivers of farmland birds at a large spatial scale and hence to develop effective conservation measures.

In Japan, the Fukushima Daiichi nuclear power plant accident in 2011 led to uninhabited areas due to an evacuation order for the contaminated area, and all agricultural activities within those areas were stopped (Yoshioka, Mishima, & Fukasawa, 2015). Although observable effects of ionizing radiation at the population level on terrestrial organisms were unlikely (UNSCARE, 2017), land abandonment in the evacuation zone can have indirect effects. These environmental changes are thought to have a significant impact on the organisms that use farmland. Therefore,

extensive monitoring in the farmland throughout the region is needed to evaluate the effect of

farmland abandonment in the evacuation zone. However, with respect to radiation protection, it

is advisable for researchers not to stay in the evacuation zone for a long time, necessitating an

efficient monitoring scheme. Main farmland in the region is paddy fields, which are used by a lot

of herons and egrets (Ardeidae). Therefore, they are suitable indicator species of the paddy field

environment. They are the main predators in paddy fields and are particularly vulnerable to the

abandonment of paddy fields (Amano & Katayama, 2009; Lane & Fujioka, 1998). Their large

size and conspicuous whitish body color make Ardeidae foraging in paddy fields easily detectable

from roads.

Vehicle-mounted cameras offer a novel and efficient approach for wildlife monitoring along

roads. These systems enable continuous, non-intrusive image collection over large areas while

minimizing the need for prolonged field presence. The roadside survey approach, involving

species observations from moving vehicles, has the capacity to collect data for species abundance

indices across broad areas at low cost (Abella, Spencer, Hoines, & Nazarchyk, 2009; Catry,

Moreira, Deus, Silva, & Águas, 2015; Milton & Dean, 1998). Recently, vehicle-mounted camera

footage—including from platforms such as Google Street View—has been used to monitor birds

and other organisms in various habitats (Deus, Silva, Catry, Rocha, & Moreira, 2016; Olea &

Mateo-Tomás, 2013; Rousselet et al., 2013). However, applications for estimating animal density

have been limited to areas with clearly defined and uninterrupted backgrounds (e.g., birds nesting

on cliffs) , and challenges remain in open habitats such as farmlands, where detection ranges are

uncertain. Ignoring this uncertainty can lead to biased estimates of abundance and environmental

effects (Buckland, Marsden, & Green, 2008).

Distance sampling can resolve the issue of bias in abundance estimation by video-based

roadside survey. Distance sampling estimates animal density or abundance from the distance

between the observer and the detected individual, accounting for the decrease in detection

probability due to distance (Buckland et al., 2001; Fox et al., 2017; Hara, Yamaji, Yokota,

Thaitakoo, & Sampei, 2018; Howe, Buckland, Després-Einspenner, & Kühl, 2017; Kvasnes,

Pedersen, & Nilsen, 2018; Stober, Prieto-Gonzalez, Smith, Marques, & Thomas, 2017). Distance

sampling can be used to estimate trends in farmland bird populations in roadside surveys.

However, there might be uncertainty in the distance between the observer and the detected

individual measured from video images. For robust abundance estimation, an independent field

test of location uncertainty in distance sampling is required (Hefley, Boyle, & Mohankumar,

2020).

Comparison of the observer–target distances measured by roadside survey and unmanned

aerial vehicles (UAVs) is expected to be a feasible independent field test. UAVs have become an

effective tool for monitoring animal populations and individual locations (Iwamoto, Nogami,

Ichinose, & Takeda, 2022; Witczuk, Pagacz, Zmarz, & Cypel, 2018). Although their application

to large spatial areas remains costly, this method is highly accurate because UAVs can measure

animal positions from almost directly above. If the same individuals are observed by both roadside

survey and UAV, location uncertainty in roadside survey can be quantified. Integrating location

uncertainty obtained by these small-scale measurements into a stochastic distance sampling

model will allow us to moderate cost–accuracy trade-offs and estimate abundance at a large spatial

scale.

The purpose of this study was to estimate the population density of Ardeidae in rice fields in

and around the evacuation zone of the Fukushima Daiichi nuclear power plant accident by

roadside survey. We surveyed the locations of Ardeidae by using a vehicle-mounted camera and

then estimated population densities by using a distance sampling model incorporating location

errors. The magnitude of location errors by roadside survey was determined by a field test

comparing roadside survey with UAV-mounted cameras. To evaluate the importance of

accounting for location uncertainty, population densities were compared between analyses with

and without location errors. We demonstrated the impact of paddy field abandonment due to the

evacuation order by comparing the population density with a counter-factual scenario without an

evacuation order.

2   Materials and methods

2.1   Roadside survey of Ardeidae around the evacuation zone

The roadside survey with vehicle-mounted cameras of abandoned and cultivated farmlands

was conducted in eastern Fukushima, which includes the evacuation zone of the Fukushima

Daiichi nuclear power plant accident (Fig. 1). The evacuation zone was designated after the

Fukushima Daiichi nuclear power plant accident in March 2011 and is divided into three

categories according to radiation levels: Difficult-to-Return Area, Restricted-Residential Area,

and Preparing for the Lifting of the Evacuation Order Area (Cabinet Office, Government of Japan,

2013). In all three categories, cropland cultivation did not resume until 2016, except in some test

fields. The study area covered farmlands within 18 municipalities located inside and outside the

evacuation zone.

From early May to early July between 2014 and 2016, roadside video recording surveys were

conducted on farmlands by vehicle-mounted cameras. Video cameras with GPS (VIRB Elite or

VIRB X; Garmin Ltd.) were mounted on the side windows of a vehicle, which was driven on

roads adjacent to farmlands for video recording (Fig. 2). Videos recorded by the two camera

models (VIRB Elite or VIRB X) simultaneously at the same location showed no obvious

differences. Although the video cameras were mounted on both sides of a car in many cases, some

surveys were carried out using a video camera mounted on one side only. The GPS trajectories of

the car were also recorded simultaneously. Videos were shot at 1920 × 1080 p/30 fps for all models

and saved in mp4 file format. The speed of the car during the survey varied depending on the

speed limit (60 km/h or less) and road congestion; however, the frame rate was sufficiently high

to detect Ardeidae under all speeds in the study.

To identify farmland within the viewshed from the vehicle-mounted cameras, the farmland

polygons in the direction of the camera and not hidden by obstacles such as forest and houses

were selected as follows (Fig. 3). Farmland polygons in this area were created from aerial

photographs taken in 2006–2015 (see Appendix 1 for detailed methods). The farmland polygons

included upland fields, pastures, and abandoned farmlands, and more than 70% of the farmland

area was paddy fields in Fukushima prefecture before the nuclear power plant accident (in 2010)

(Ministry of Agriculture, Forestry and Fisheries, 2020). First, we extracted the farmland polygons

in the direction of the camera and intersecting within a straight-line distance of 10 m from the

trajectories of the car; these were defined as the viewshed. Ten meters was set as an approximate

threshold for adjacency (i.e., no obstacles) between the road and farmland. Second, we

sequentially included farmlands that were within 10 meters from the viewshed obtained in the

previous step. Farmlands were not included in the viewshed when the distance from the movement

trajectory was more than 500 m because this was too far to detect Ardeidae. We visually verified

that the selection criteria could effectively extract farmlands within the viewshed of the camera.

For subsequent analysis, only paddy field polygons were used, which were selected from the

identified polygons based on existing information (Ministry of Agriculture, Forestry and Fisheries,

2021) and aerial photographs.

We played the videos back and searched for Ardeidae in paddy fields. Then, we determined

the locations on the map following the same method used to test location errors described above.

Because it was difficult to identify species from the video images correctly, except for grey heron,

we merged all species of Ardeidae for further analyses. The following Ardeidae may be found in

the study area: grey heron *Ardea cinerea*, great egret *A. cinerea*, intermediate egret *A. intermedia*,

little egret *Egretta garzetta*, and cattle egret *Bubulcus ibis*. All of these Ardeidae are colonial and

breed in the summer. In addition, flooding status (i.e., flooded or not) was recorded for the paddy

fields where Ardeidae were found.

2.2   Field test of location errors of Ardeidae using UAVs

Location errors of the vehicle-mounted camera were validated by comparison with the

accurate values determined by UAV images (UAV-truth). The survey was conducted on paddy

fields in two regions, the district of Inashiki in Ibaraki Prefecture and the city of Minamisoma in

Fukushima Prefecture. These paddy fields are about 5 square kilometers and are used for foraging

by Ardeidae. The survey was conducted in Inashiki in July 2019 and May 2020 and in

Minamisoma in June 2020. During the survey period, Ardeidae in the paddy fields were recorded

simultaneously by a vehicle-mounted video camera (VIRB X; Garmin Ltd.) and a video camera

in a UAV (Phantom 4; DJI). The GPS trajectories of vehicle-mounted video cameras were

recorded by VIRB X simultaneously.

Locations of Ardeidae were determined from the movies captured by the vehicle-mounted

cameras. If the researcher observed Ardeidae in the video, the movement trajectory information

recorded at the time the video was taken was used to determine the location of the camera. Then,

using information on the camera location with orthographical aerial imagery layers as background

maps (GSI Tiles) and views surrounding Ardeidae (paddy levees, houses, telegraph poles, etc.) in

the video image, the locations of Ardeidae were manually determined and recorded using QGIS

ver.3.4.3.

To obtain correct locations of Ardeidae, georeferenced images obtained by the UAV were

used. Georeferencing was performed using Georeferencer (Transformation type: Projective,

Resampling method: Linear) in QGIS. The measurement errors of locations obtained by vehicle-

mounted cameras were calculated as the Euclidean distances between the locations of the

Ardeidae determined by the vehicle-mounted cameras and the UAV-truth values.

2.3   Statistical analysis

We employed a fully model-based approach to estimate the abundance of Ardeidae and the

effect of land abandonment based on a distance sampling considering individual location

uncertainty. As our observations consisted of the camera-based distance sampling results with

location uncertainty and the independent field test of location errors based on correct locations

determined by UAV, we integrated the likelihoods and estimated all parameters simultaneously.

A common approach for model-based distance sampling without location uncertainty is to

treat the occurrence of individuals in relation to the distance from an observer as a realization of

an inhomogeneous Poisson process (IPP) (Johnson, Laake, & Ver Hoef, 2010). The likelihood

of distance sampling without location uncertainty is expressed by the following equation

(Johnson, Laake, & Ver Hoef, 2010).

216
$$L_{IPP}\left(\alpha, \beta, \gamma | \mathbf{Y}\right) = \prod_{i=1}^{n} \lambda\left(\mathbf{y}_i | \alpha, \beta\right) \delta(\mathbf{y}_i | \gamma) \exp\left(-\int_{\mathcal{S}} \lambda\left(\mathbf{z} | \alpha, \beta\right) \delta(\mathbf{z} | \gamma) d\mathbf{z}\right)$$

(eqn. 1)

where $\mathbf{Y} = (\mathbf{y}_1, \mathbf{y}_2, ..., \mathbf{y}_n)$ are true locations of observed individuals, (α, β, γ) are model parameters,

and $\mathcal{S}$ is the spatial domain(s) that is potentially observable. $\lambda(\mathbf{z} | \alpha, \beta)$ is an intensity function of the

population density at location $\mathbf{z}$ and is commonly specified as follows.

$\lambda(\mathbf{z} | \alpha, \beta) = \exp(\alpha + \beta x(\mathbf{z}))$

where α is the intercept, $\beta = (\beta_1, ..., \beta_p)'$ is a vector of regression coefficients, and $x(\mathbf{z})$ is a vector of

covariates on population density at location $\mathbf{z}$. $\delta(\mathbf{z} | \gamma)$ is the detection probability, which is expressed

as a thinning function as follows.

$\delta(z) = \exp(-\gamma D(\mathbf{z}))$

where $D(\mathbf{z})$ is the Euclidian distance from an observer to location $\mathbf{z}$.

Let us consider the case when observed location $\mathbf{Y}' = (\mathbf{y}'_1, \mathbf{y}'_2, ..., \mathbf{y}'_n)$ is assumed to vary around the

true location $\mathbf{Y}$ due to location error. The likelihood of IPP incorporating location uncertainty can be

obtained by integrating out $\mathbf{y}_i$ over the surveyed region (Hefley et al., 2020).

232
$$L(\alpha, \beta, \gamma, \boldsymbol{\sigma}|\mathbf{Y}') = \prod_{i=1}^{n} \int_{\mathcal{S}_i} p_{\mathrm{err}}(\mathbf{y}'_i|\mathbf{y}_i, \boldsymbol{\sigma})\lambda(\mathbf{y}_i)\delta(\mathbf{y}_i)\mathrm{d}\mathbf{y}_i \exp\left(-\int_{\mathcal{S}} \lambda(\mathbf{z})\delta(\mathbf{z})dz\right)$$

(eqn. 2, parameters α, $\boldsymbol{\beta}$, γ were not included on the right side for simplicity)

where $p_{\mathrm{err}}(\mathbf{y}'|\mathbf{y}, \boldsymbol{\sigma})$ is a probability distribution of location error, $\boldsymbol{\sigma}$ is a vector of location error

parameters, and $\mathcal{S}_i$ is subset of $\mathcal{S}$ in which $\mathbf{y}_i$ exist. For simplicity, we assume that the location error

is isotropic and normally distributed.

$p_{\mathrm{err}}(\mathbf{y}'|\mathbf{y}, \boldsymbol{\sigma}) = N_2(\mathbf{y}'|\mathbf{y}, f(\mathbf{y}', \mathbf{y}, \boldsymbol{\sigma})\mathbf{I})$ (eqn. 3)

where $f(\mathbf{y}', \mathbf{y}, \boldsymbol{\sigma})$ is a function of the standard deviation depending on the true location. In general,

the greater the distance from the observer, the lower the accuracy of the observation position. The

relationship was assumed to be described by a power function of Euclidean distances between $\mathbf{y}'$ and

$\mathbf{y}$, $D(\mathbf{y}', \mathbf{y})$, with two parameters $\boldsymbol{\sigma} = (\sigma_0, \sigma_1)$ as follows.

$f(\mathbf{y}', \mathbf{y}, \boldsymbol{\sigma}) = \sigma_0 D(\mathbf{y}', \mathbf{y})^{\sigma_1}$ (eqn. 4)

The likelihood function required integration over spatial domains, which is difficult to solve

analytically. A practical approach is discretizing spatial domains into grid cells and replacing

integration   with   piecewise   summation   as   follows.

237
$$\int_{\mathcal{S}_i} \mathrm{p_{err}}\left(\mathbf{y}_i^{'} \, | \mathbf{y}_i\right)\lambda(\mathbf{y}_i)\delta(\mathbf{y}_i)\mathrm{d}\mathbf{y}_i \approx \sum_{q \subset \mathcal{S}_i} A_q \mathrm{p_{err}}\left(\mathbf{y}_i^{'} \, | \mathbf{y}_q\right)\lambda(\mathbf{y}_q)\delta(\mathbf{y}_q)$$

238
$$\int_{\mathcal{S}} \lambda(\mathbf{z})\delta(\mathbf{z})d\mathbf{z} \approx \sum_{q \subset \mathcal{S}} A_q\lambda(\mathbf{z}_q)\delta(\mathbf{z}_q)$$

where $\mathbf{z}_q$ and A_q are the centroid and area of grid cell q, respectively.

The location error parameter $\boldsymbol{\sigma}$ in the likelihood function described above is not identifiable

without an independent survey of location error. In this study, the location error and UAV-truth were

included simultaneously in the distance sampling model for the estimation of all parameters. When

the true location is available, the error distribution (eqn. 3 and 4) can be written equivalently by the

Rayleigh distribution (i.e., Weibull distribution with shape parameter 2) for the Euclidean distance

between the observed and true location.

248
$$\mathrm{p_{err}}(D|\boldsymbol{\sigma}) = D/(\sigma_0^{\,2}D^{2\sigma_1})\exp\left(-D/(2\sigma_0^{\,2}D^{2\sigma_1})\right)$$

246   Given the vector of distances obtained by the field test $D(\mathbf{y'}, \mathbf{y}) = (D(y'_1, y_1), ..., D(y'_m, y_m))$, the joint

247   likelihood is given as follows.

249     $\mathrm{L}\left(\alpha, \beta, \gamma, \boldsymbol{\sigma}|\mathbf{Y}^{'}, \mathbf{D}(\mathbf{y'}, \mathbf{y})\right)$

250
$$= \prod_{i=1}^{n} \sum_{q \subset \mathcal{S}_i} A_q \mathrm{p_{err}}\left(\mathbf{y}_i^{'} \, | \mathbf{y}_q\right)\lambda(\mathbf{y}_q)\delta(\mathbf{y}_q) \exp\left(-\sum_{q \subset \mathcal{S}} A_q\lambda(\mathbf{z}_q)\delta(\mathbf{z}_q)\right)\prod_{j=1}^{m} \mathrm{p_{err}}(D_j)$$

Both the maximum likelihood method and Bayesian method can be used for parameter estimation.

When the Bayesian method is applied, prior information and constraints on parameters can be

incorporated as a prior distribution.

We estimated the effects of the evacuation order (evacuation zone or not) (Cabinet Office,

Government of Japan, 2013), the area of water bodies (rivers and lakes) (Ogawa et al., 2013)

around the farmlands, and the year of the study on the abundance of Ardeidae. We defined

farmland polygons within the viewshed of the camera as a multi-part domain S and discretized it

into grid cells with a resolution of 10 m. Then, these variables and the Euclidean distances from

the camera were assigned to each cell. Even when location error is taken into account, the error

is reasonably assumed to be within a paddy field polygon because paddy levees act as sharp

borders dividing paddy field polygons. Therefore, we assumed that the subdomain S_i containing

the true location of the observed individual i was restricted to a paddy field polygon where the

individual was observed.

The IPP model is often limited by overfitting, and it is common to apply regression shrinkage

techniques. In this study, we applied the Bayesian default prior (Gelman, Jakulin, Pittau, & Su,

2007) to stabilize the coefficients and to prevent overfitting. We applied a weakly informative

prior, Student's *t*-test (df = 5, mean = 0, scale = 2.5), to regression coefficients $\boldsymbol{\beta}$ to avoid

overfitting. For other parameters, improper flat priors p(θ) = 1 were used. The maximum *a*

posteriori (MAP) estimate of each parameter was calculated using a quasi-Newton algorithm.

To estimate credible intervals of parameters, we obtained posterior samples by importance

sampling (Gelman et al., 2013) with a proposal distribution that approximates the posterior. For

the proposal distribution, we used a multivariate t-distribution with four degrees of freedom

whose mean vector and scale matrix are the MAP estimate and the inverse of the Hessian matrix

at the MAP, respectively. We drew 10,000 samples from the proposal distribution and obtained

3,000 posterior samples.

We calculated the abundance of Ardeidae by applying MAP estimates of each parameter to

grid cells in paddy field polygons throughout the study area, with and without accounting for

measurement error, and compared the two abundance results. The predicted values were

aggregated for each 1-km square grid. To evaluate the effect of paddy field abandonment in the

evacuation zone on the population size of Ardeidae, we predicted the population size in the study

area under a counter-factual scenario in which no evacuation order had been given. All analyses

in this study were implemented using R (R Core Team, 2021) (Appendix 2).

3   Results

3.1   Field test of location error of Ardeidae

In total, 78 Ardeidae (24 in July 2019, 31 in May 2020 in Inashiki and 23 in June 2020 in

Minamisoma) were located with GIS by the vehicle-mounted camera and the camera on the UAV.

The Euclidean distance between the locations of the Ardeidae determined by the vehicle-mounted

camera and the true locations determined by the UAV was defined as the measurement error (Fig.

4). The location errors were increased with the distance between the observer and the Ardeidae.

3.2   Roadside survey with vehicle-mounted cameras and abundance estimates of Ardeidae

The total distances traveled while recording with cameras were 3,675 km in the evacuation

zone and 3,356 km outside the evacuation zone. The areas of farmland polygons (including

abandoned fields in response to the evacuation order), selected as the viewshed, were 10.07 km^2

in and 14.34 km^2 around the evacuation zone.

A total of 56 individual Ardeidae were observed around the evacuation zone. No Ardeidae were

observed in the evacuation zone (Fig. 5). All Ardeidae were found in flooded paddy fields. The

frequency distribution of observed distances to Ardeidae was more concentrated near zero than

the distribution of distances to roads for all grid cells of paddy field polygons. Although all

polygons had a median distance of 133.3 m from roads, the polygons containing Ardeidae

discovery sites had a median distance of 47.1 m and a maximum of 104.4 m (Fig. 6).

The MAP estimates and summary statistics obtained by importance sampling are shown in

Table 1. The water body area had a positive association with the abundance of Ardeidae.

Abundance was lower in the evacuation zone than outside the zone. Although the signs (i.e.,

positive or negative) of parameter estimates were similar between the models with and without

the location error, the parameter values differed slightly. The distance from the camera had a

negative association with the probability of detection (Fig. 4). The probability decreased to half

at 35.4 m (95% credible interval: 29.6–42.8 m) from the camera. The scaling factor and power

value when measurement errors are expressed as a power function of the distance from the camera

were 0.644 and 0.614. The measurement errors tended to increase with increasing distance from

the camera on the car but exhibited a concave downward trend.

The predicted abundance of Ardeidae were substantially larger outside the evacuation zone

than inside the zone (Table 2, Fig. 7). This trend did not differ between the models with and

without location error. The predicted total population of Ardeidae in the study area was 7,820

individuals. Assuming that the evacuation zone was never designated, the predicted population

would have increased by 2,647 individuals to 10,467 individuals (Fig. 8).

4 Discussion

We developed an efficient method for monitoring large birds in open areas by combining roadside

surveys using vehicle-mounted cameras and distance sampling. By incorporating measurement

error into the distance sampling framework, we addressed the challenge posed by the lack of a

clearly defined observation area a common limitation of roadside surveys. The spatially explicit

approach enabled us to estimate the impact of large-scale land abandonment on Ardeidae

populations. Our survey revealed a large gap in Ardeidae distribution in the evacuation zone and

allowed us to quantify the magnitude of this impact as a loss of abundance (Fig. 8, Table 2). The

estimated number of foraging individuals around the evacuation zone was consistent with a

previous field study that reported approximately 30 individuals per km^2 (Lane & Fujioka, 1998) ,

supporting the reliability of our abundance estimates.

The direct effects of radiation exposure from the Fukushima Daiichi nuclear power plant

accident on wildlife populations and ecosystems are not large enough to be observable at the

population level (UNSCEARE 2017). However, the evacuation orders in response to the nuclear

accident have resulted in short-term changes in land use and human activities (Tamaoki, 2016;

Yoshioka et al., 2015), and these changes may have important impacts on wildlife populations

and ecosystems (Fujimoto, Mitsunaga, & Takeuchi, 2015; Lyons, Okuda, Hamilton, Hinton, &

Beasley, 2020; Matsushima, Ihara, Inaba, & Horiguchi, 2021; Yoshioka et al., 2015). The strong

negative impact of the evacuation order on Ardeidae populations can be attributed to the decrease

in food availability due to the widespread abandonment of paddy fields. Paddy field abandonment

leads to the growth of dense vegetation, which is not a common foraging environment for

Ardeidae (Katayama, Osawa, Amano, & Kusumoto, 2015; Maeda, 2005). In addition, Ardeidae

feed on fish and frogs in flooded paddy fields (Tojo, 1996), which can be maintained by

cultivation (Maeda, 2001). Our results clearly demonstrated that paddy field abandonment in

evacuation zones affects the distribution of birds over a wide area. These findings are consistent

with the observation that the long-term abandonment of farmland reduces the biodiversity of

farmlands (Koshida & Katayama, 2018). Since 2016, the evacuation zone has been reduced and

agricultural cultivation has begun to resume. In the future, the effect of resumption should be

tested by continued monitoring following the protocols developed in this study.

While the roadside survey approach benefits from the ability to efficiently survey a wide

area, the use of vehicle-mounted cameras will further improve efficiency. In this study, 237 hours

of video were recorded to search for Ardeidae. In a field survey, experts would be required for the

full time; however, the roadside survey method with vehicle-mounted cameras does not require

expertise in bird identification in the field. In addition, the work of sorting paddy fields without

Ardeidae, accounting for the majority of each video, does not require expertise; instead, experts

only need to check images with suspected Ardeidae. Crowdsourcing this sorting process can

promote the participation of birders in remote areas who are interested in the survey. In the future,

automated counting may be possible (Norouzzadeh et al., 2018).

An accurate distance between the observer and target is required for distance sampling to

estimate density; accordingly, when the distance accuracy could not be guaranteed, it is necessary

to account for measurement error in distance sampling (Hefley et al., 2020). The incorporation of

measurement error in the location of Ardeidae did not significantly affect density estimates. This

result showed that the distance accuracy of roadside survey with vehicle-mounted cameras in our

study was sufficient for density estimation by distance sampling. The robust performance of the

roadside survey can be explained by the study system. The Ardeidae were found in paddy fields,

and the location error could not exceed the size of one rice field separated by paths or levees.

However, use of a UAV to estimate location errors for incorporation into a distance sampling

model may be an effective approach in situations where measurement errors are likely such as

natural grasslands and wetlands, where artificial markers are absent. Roadside survey by vehicle-

mounted cameras may be a feasible strategy for various environments by using this method.

The detection probability depends on the distance from the observer as well as various

covariates, in which case multiple-covariate distance sampling is appropriate (Marques &

Buckland, 2003; Thomas et al., 2010). In this study, we did not consider the evacuation zone as a

covariate, despite the potential effect of this parameter on the detection probability, because no

Ardeidae were observed in the evacuation zone. However, we believe the absence of Ardeidae in

the evacuation zone reflects differences in true population densities resulting from the

abandonment of rice field because dense vegetation is generally not suitable for foraging by these

species (Katayama, Osawa, Amano, & Kusumoto, 2015; Maeda, 2005). These species were not

found in the evacuation zone, even near roads in which detectability is less affected by the

vegetation.

The escape behavior of birds during surveys might biases population estimates (Hutto &

Young, 2003; Roberts & Schnell, 2006). At paddy fields bordering roads, Ardeidae may fly away

382 to avoid the approach of vehicles carrying cameras. Ardeidae at a distance from the camera can

383 therefore be overlooked, which can lead to biases in the detection function and population

384 estimates. However, the videos obtained in this study did not reveal any Ardeidae in flight, which

385 could be a response to an approaching vehicle. In fact, waterbirds are not substantially impacted

386 by approaching vehicles (McLeod, Guay, Taysom, Robinson, & Weston, 2013). Similarly, in this

387 study, the effect of approaching vehicles was considered negligible for Ardeidae. If this method

388 is applied to broader taxa, the use of fisheye lenses or 360-degree cameras may be a viable option

389 to capture individuals before they are affected by an observer.

390  In our survey, we estimated the number of Ardeidae using paddy fields; however, these

391 estimates cannot be interpreted as total counts for the study area. The survey was conducted during

392 the breeding season, and one member of each nesting pair always stays in the nest. In addition,

393 because Ardeidae fly over a wide area in search of good foraging sites, it is thought that they

394 frequently move in and out of the evacuation zone and travel to and from the study area. It is

395 necessary to consider whether the number of Ardeidae obtained in this study is an indicator of the

396 quality of the paddy field as a foraging site. However, we believe that the density of Ardeidae

397 foraging in paddy fields is a viable indicator of biodiversity in agricultural land because it is a

398 common parameter for assessing the quality of paddy fields as foraging sites (Amano & Katayama,

399 2009; Katayama et al., 2019; Lane & Fujioka, 1998).

Another limitation of our study is that we were unable to identify the Ardeidae at the species

level. We considered the total abundance of Ardeidae. The five species in the study area have

overlapping ecological niches; they forage in rice fields and often share colonies (Mashiko &

Toquenaga, 2014). In addition, the number of Ardeidae is related to the abandonment of rice field

and taxon diversity and can therefore be used as an indicator of species richness on paddy fields

(Katayama, Mashiko, Koshida, & Yamaura, 2021; Katayama et al., 2019; Koshida & Katayama,

2018). In the future, technological improvements such as increased camera resolution may make

it possible to identify species and evaluate responses.

5 Conclusion

The combination of roadside surveys using vehicle-mounted cameras and distance sampling

provided a reliable and efficient method to estimate the abundance and distribution of large birds

in open agricultural landscapes. This approach allowed us to quantify the strong negative impact

of paddy field abandonment on Ardeidae populations in the evacuation zone following the

Fukushima Daiichi nuclear power plant accident.

Thanks to the structured landscape of paddy fields, our system was robust to location

measurement error; however, by incorporating measurement error into the model, it can also be

applied to other open habitats such as wetlands and grasslands. The approach is scalable, cost-

efficient, and compatible with emerging technologies such as UAVs, street-level imagery

platforms (e.g., Mapillary https://www.mapillary.com), and machine learning-based species

recognition (Norouzzadeh et al., 2021) and distance estimation (Johanns, Haucke, & Steinhage,

2022). It holds promise for wide-scale biodiversity monitoring and conservation planning in

agricultural systems and beyond.

424 DATA ACCESSIBILITY

425 Data will be uploaded to the Dryad Digital Repository after acceptance.

428 Acknowledgements

429 We are grateful to Fukushima Robot Test Field for helping field work. Yoshio Mishima

430 contributed to create the farmland polygon data but declined to join as a co-author. This study was

431 partially supported by JSPS KAKENHI Grant number 21H03656.

432 Authors' contributions

433 Nao Kumada, Keita Fukasawa and Akira Yoshioka conceived the ideas and designed the

434 methodology; All the authors collected the data; Nao Kumada and Keita Fukasawa analysed the

435 data; Nao Kumada, Keita Fukasawa and Akira Yoshioka led the writing of the manuscript. All the

436 authors reviewed and contributed critically to the draft and gave final approval for publication.

Conflict of interest

439   The authors declare no conflict of interest.

References

Abella, S. R., Spencer, J. E., Hoines, J., & Nazarchyk, C. (2009). Assessing an exotic plant surveying program in the Mojave Desert, Clark County, Nevada, USA. *Environmental Monitoring and Assessment*, *151*(1–4), 221–230. doi: 10.1007/s10661-008-0263-0

Amano, T., & Katayama, N. (2009). Hierarchical movement decisions in predators: Effects of foraging experience at more than one spatial and temporal scale. *Ecology*, *90*(12), 3536–3545. doi: 10.1890/08-1910.1

Birrer, S., Spiess, M., Herzog, F., Jenny, M., Kohli, L., & Lugrin, B. (2007). The Swiss agri-environment scheme promotes farmland birds: but only moderately. *Journal of Ornithology*, *148*(2), 295–303. doi: 10.1007/s10336-007-0237-y

Buckland, S.T., Anderson, D. R., Burnham, K. P., Laake, J. L., Borchers, D. L., & Thomas, L. (2001). *Introduction to distance sampling: estimating abundance of biological populations*. Oxford (United Kingdom) Oxford Univ. Press.

Buckland, S. T., Marsden, S. J., & Green, R. E. (2008). Estimating bird abundance: Making methods work. *Bird Conservation International*, *18*, S91–S108. doi: 10.1017/S0959270908000294

Cabinet Office, Government of Japan. About revision of the evacuation zone (in Japanese). (2013). http://www.meti.go.jp/earthquake/nuclear/pdf/131009/131009_02a.pdf. Accessed 17 January 2023.

Catry, F. X., Moreira, F., Deus, E., Silva, J. S., & Águas, A. (2015). Assessing the extent and the environmental drivers of Eucalyptus globulus wildling establishment in Portugal: results from a countrywide survey. *Biological Invasions*, *17*(11), 3163–3181. doi: 10.1007/s10530-015-0943-y

Chiron, F., Filippi-Codaccioni, O., Jiguet, F., & Devictor, V. (2010). Effects of non-cropped landscape diversity on spatial dynamics of farmland birds in intensive farming systems.

Biological Conservation, *143*(11), 2609–2616. doi:

https://doi.org/10.1016/j.biocon.2010.07.003

Deus, E., Silva, J. S., Catry, F. X., Rocha, M., & Moreira, F. (2016). Google Street View as an

alternative method to car surveys in large-scale vegetation assessments. *Environmental*

Monitoring and Assessment, *188*(10), 560. doi: 10.1007/s10661-016-5555-1

Fox, C. H., Huettmann, F. H., Harvey, G. K. A., Morgan, K. H., Robinson, J., Williams, R., &

Paquet, P. C. (2017). Predictions from machine learning ensembles: Marine bird

distribution and density on Canada's Pacific coast. *Marine Ecology Progress Series*, *566*,

199–216. doi: 10.3354/meps12030

Fujimoto, R., Mitsunaga, T., & Takeuchi, M. (2015). Influence of evacuation orders on the

occurrence of the wild boar Sus scrofa in farmlands in the northern evacuation zone of the

accident at Tokyo Electric Power Company's Fukushima Daiichi Nuclear Power Station

(in Japanese with English abstract). *Mammalian Science*, *55*(2), 145–154.

Fujioka, M., Don Lee, S., Kurechi, M., & Yoshida, H. (2010). Bird use of rice fields in Korea

and Japan. *Waterbirds*, *33*(SPEC.ISSUE.1), 8–29. doi: 10.1675/063.033.s102

Fuller, R. J., Gregory, R. D., Gibbons, D. W., Marchant, J. H., Wilson, J. D., Baillie, S. R., &

Carter, N. (1995). Population Declines and Range Contractions among Lowland Farmland

Birds in Britain. *Conservation Biology*, *9*(6), 1425–1441. doi: 10.1046/j.1523-

1739.1995.09061425.x

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013).

Bayesian Data Analysis (third edition). New York: Chapman and Hall/CRC

Gelman, A., Jakulin, A., Pittau, M. G., & Su, Y.-S. (2007). A weakly informative default prior

distribution for logistic and other regression models. *The Annals of Applied Statistics*, *2*.

doi: 10.2139/ssrn.1010421

Gillings, S., & Fuller, R. J. (1998). Changes in bird populations on sample lowland English

farms in relation to loss of hedgerows and other non-crop habitats. *Oecologia*, *116*(1),

120–127. doi: 10.1007/s004420050570

Hara, Y., Yamaji, K., Yokota, S., Thaitakoo, D., & Sampei, Y. (2018). Dynamic wetland

mosaic environments and Asian openbill habitat creation in peri-urban Bangkok. *Urban*

Ecosystems, *21*(2), 305–322. doi: 10.1007/s11252-017-0718-6

Hefley, T. J., Boyle, W. A., & Mohankumar, N. M. (2020). *Accounting for location uncertainty*

in distance sampling data. (2020). Retrieved from http://arxiv.org/abs/2005.14316

Howe, E. J., Buckland, S. T., Després-Einspenner, M. L., & Kühl, H. S. (2017). Distance

sampling with camera traps. *Methods in Ecology and Evolution*, *8*(11), 1558–1565. doi:

10.1111/2041-210X.12790

Hutto, R. L., & Young, J. S. (2003). On the Design of Monitoring Programs and the Use of

Population Indices: A Reply to Ellingson and Lukacs. *Wildlife Society Bulletin*, 31(3),

903–910. Retrieved from http://www.jstor.org/stable/3784617

Iwamoto, M., Nogami, S., Ichinose, T., & Takeda, K. (2022). Unmanned aerial vehicles as a

useful tool for investigating animal movements. *Methods in Ecology and Evolution*, *13*(5),

969–975. doi: 10.1111/2041-210X.13829

Johanns, P., Haucke, T., & Steinhage, V. (2022). Automated distance estimation for wildlife

camera trapping. *Ecological Informatics*, *70*(February), 101734. doi:

10.1016/j.ecoinf.2022.101734

Johnson, D. S., Laake, J. L., & Ver Hoef, J. M. (2010). A model-based approach for making

ecological inference from distance sampling data. *Biometrics*, *66*(1), 310–318. doi:

10.1111/j.1541-0420.2009.01265.x

Katayama, N., Mashiko, M., Koshida, C., & Yamaura, Y. (2021). Agriculture, Ecosystems and

Environment Effects of rice field abandonment rates on bird communities in mixed

farmland – woodland landscapes in Japan. *Agriculture, Ecosystems and Environment*,

319(June), 107539. doi: 10.1016/j.agee.2021.107539

Katayama, N., Osada, Y., Mashiko, M., Baba, Y. G., Tanaka, K., Kusumoto, Y., … Natuhara,

Y. (2019). Organic farming and associated management practices benefit multiple wildlife

taxa: A large-scale field study in rice paddy landscapes. *Journal of Applied Ecology*,

56(8), 1970–1981. doi: 10.1111/1365-2664.13446

Katayama, N., Osawa, T., Amano, T., & Kusumoto, Y. (2015). Are both agricultural

intensification and farmland abandonment threats to biodiversity? A test with bird

communities in paddy-dominated landscapes. *Agriculture, Ecosystems and Environment*,

214, 21–30. doi: 10.1016/j.agee.2015.08.014

Koshida, C., & Katayama, N. (2018). Meta-analysis of the effects of rice field abandonment on

biodiversity in Japan. *Conservation Biology*, *32*(6), 1392–1402. doi: 10.1111/cobi.13156

Kvasnes, M. A. J., Pedersen, H. C., & Nilsen, E. B. (2018). Quantifying suitable late summer

brood habitats for willow ptarmigan in Norway. *BMC Ecology*, *18*(1), 1–13. doi:

10.1186/s12898-018-0196-6

Lane, S. J., & Fujioka, M. (1998). The impact of changes in irrigation practices on the

distribution of foraging egrets and herons (Ardeidae) in the rice fields of central Japan.

Biological Conservation, *83*(2), 221–230. doi: 10.1016/S0006-3207(97)00054-2

Lyons, P. C., Okuda, K., Hamilton, M. T., Hinton, T. G., & Beasley, J. C. (2020). Rewilding of

Fukushima's human evacuation zone. *Frontiers in Ecology and the Environment*, *18*(3),

127–134. doi: 10.1002/fee.2149

Maeda, T. (2001). Patterns of bird abundance and habitat use in rice fields of the Kanto Plain,

central Japan. *Ecological Research*, *16*(3), 569–585. doi: 10.1046/j.1440-

1703.2001.00418.x

Maeda, T. (2005). Bird use of rice field strips of varying width in the Kanto Plain of central

Japan. *Agriculture, Ecosystems and Environment*, *105*(1–2), 347–351. doi:

10.1016/j.agee.2004.02.007

Marques, F. F. C., & Buckland, S. T. (2003). Incorporating Covariates into Standard Line

Transect Analyses. *Biometrics*, *59*(4), 924–935. doi: 10.1111/j.0006-341X.2003.00107.x

Mashiko, M., & Toquenaga, Y. (2014). Grey Heron (Ardea cinerea) Expansion Promotes the

Persistence of Mixed-species Heron Colonies. *WATERBIRDS*, *37*(4), 362–370. doi:

10.1675/063.037.0403

Matsushima, N., Ihara, S., Inaba, O., & Horiguchi, T. (2021). Assessing the impact of large-

scale farmland abandonment on the habitat distributions of frog species after the

Fukushima nuclear accident. *Oecologia*, *196*(4), 1219–1232. doi: 10.1007/s00442-021-

04991-y

McLeod, E. M., Guay, P. J., Taysom, A. J., Robinson, R. W., & Weston, M. A. (2013). Buses,

cars, bicycles and walkers: The influence of the type of human transport on the flight

responses of waterbirds. *PLoS ONE*, *8*(12). doi: 10.1371/journal.pone.0082008

Milton, S. J., & Dean, W. R. J. (1998). Alien plant assemblages near roads in arid and semi-arid

South Africa. *Diversity and Distributions*, *4*(4), 175–187. doi: 10.1046/j.1472-

4642.1998.00024.x

Ministry of Agriculture, Forestry and Fisheries, Government of Japan. Provision of agricultural

percel data (in Japanese). (2021). https://www.maff.go.jp/j/tokei/porigon/. Accessed 15

October 2024.

Norouzzadeh, M. S., Morris, D., Beery, S., Joshi, N., Jojic, N., & Clune, J. (2021). A deep

active learning system for species identification and counting in camera trap images.

Methods in Ecology and Evolution, *12*(1), 150–161. doi: 10.1111/2041-210X.13504

Norouzzadeh, M. S., Nguyen, A., Kosmala, M., Swanson, A., Palmer, M. S., Packer, C., &

Clune, J. (2018). Automatically identifying, counting, and describing wild animals in

camera-trap images with deep learning. *Proceedings of the National Academy of Sciences*

of the United States of America, *115*(25), E5716–E5725. doi: 10.1073/pnas.1719367115

Ogawa, M., Takenaka, A., Kadoya, T., Ishihama, F., Yamano, H., & Akasaka, M. (2013). A

comprehensive new land-use classification map for Japan for biodiversity assessment and

species distribution modeling (in Japanese with English abstract). *Japanese Journal of*

Conservation Ecology, *18*(1), 69–76.

Olea, P. P., & Mateo-Tomás, P. (2013). Assessing Species Habitat Using Google Street View:

A Case Study of Cliff-Nesting Vultures. *PLoS ONE, 8*(1), 1–8. doi:

10.1371/journal.pone.0054582

R Core Team (2021). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Roberts, J. P., & Schnell, G. D. (2006). Comparison of survey methods for wintering grassland
birds. *JOURNAL OF FIELD ORNITHOLOGY*, *77*(1), 46–60. doi: 10.1111/j.1557-
9263.2006.00024.x

Rousselet, J., Imbert, C.-E., Dekri, A., Garcia, J., Goussard, F., Vincent, B., … Rossi, J.-P.
(2013). Assessing Species Distribution Using Google Street View: A Pilot Study with the
Pine Processionary Moth. *PLoS ONE*, *8*(10), e74918. doi: 10.1371/journal.pone.0074918

Siriwardena, G. M., Baillie, S. R., Buckland, S. T., Fewster, R. M., Marchant, J. H., & Wilson,
J. D. (1998). Trends in the abundance of farmland birds: A quantitative comparison of
smoothed Common Birds Census indices. *Journal of Applied Ecology*, *35*(1), 24–43. doi:
10.1046/j.1365-2664.1998.00275.x

Stanton, R. L., Morrissey, C. A., & Clark, R. G. (2018). Analysis of trends and agricultural
drivers of farmland bird declines in North America: A review. *Agriculture, Ecosystems*
and Environment, *254*(December 2017), 244–254. doi: 10.1016/j.agee.2017.11.028

Stober, J. M., Prieto-Gonzalez, R., Smith, L. L., Marques, T. A., & Thomas, L. (2017).
Techniques for estimating the size of low-density gopher tortoise populations. *Journal of*
Fish and Wildlife Management, *8*(2), 377–386. doi: 10.3996/012017-JFWM-005

Tamaoki, M. (2016). Studies on radiation effects from the Fukushima nuclear accident on wild
organisms and ecosystems. *Global Environmental Research*, *20*, 73–82.

Thomas, L., Buckland, S. T., Rexstad, E. A., Laake, J. L., Strindberg, S., Hedley, S. L., …
Burnham, K. P. (2010). Distance software: design and analysis of distance sampling
surveys for estimating population size. *Journal of Applied Ecology*, *47*(1), 5–14. doi:
10.1111/j.1365-2664.2009.01737.x

Tojo, H. (1996). Habitat Selection, Foraging Behaviour and Prey of Five Heron Species in
Japan. *Japanese Journal of Ornithology*, *45*(3), 141–158. doi: 10.3838/jjo.45.141

UNSCEARE. (2017) FUKUSHIMA 2017 WHITE PAPER.
https://www.unscear.org/unscear/en/publications/whitepapers.html Accessed 17 January
2023.

Witczuk, J., Pagacz, S., Zmarz, A., & Cypel, M. (2018). Exploring the feasibility of unmanned
aerial vehicles and thermal imaging for ungulate surveys in forests - preliminary results.
International Journal of Remote Sensing, *39*(15–16), 5504–5521. doi:
10.1080/01431161.2017.1390621

Yoshioka, A., Mishima, Y., & Fukasawa, K. (2015). Pollinators and Other Flying Insects inside
and outside the Fukushima Evacuation Zone. *PLOS ONE*, *10*(11), e0140957. doi:
10.1371/journal.pone.0140957

Figures and tables presented alongside each individual caption

Figure 1 Map of survey routes (black lines) in Eastern Fukushima prefecture. Dark grey area

shows the evacuation zone of the Fukushima Daiichi nuclear power plant accident. Light grey

areas show farmland polygons.

Figure 2 Photo showing a video camera with GPS mounted on a side window of a vehicle (a) and

photo, captured from a video recorded by a vehicle-mounted camera, showing Ardeidae

(in a red circle) in a paddy field (b).

Figure 3 Illustration of the selection of farmlands within the viewshed. A survey route (bold line)

and farmland polygons (surrounding lines) are shown. The polygons that meet the criteria (see

text) are selected as viewsheds (shaded areas).

Figure 4 Relationship between the distance from cameras to Ardeidae and location errors. Black

line and grey area show the maximum likelihood estimates and 95% credible interval for the

regression function.

Figure 5 Location of Ardeidae observed by roadside survey.

640 Figure 6 Histograms of the distance from the video camera to all grid cells of farmland polygons
641 (light grey bars) and grid cells containing Ardeidae (dark grey bars).
642

Figure 7 Prediction map of number of Ardeidae per square kilometer. Only grids that contain

farmland area are included. The evacuation zone is circled by solid black lines.

Figure 8 Map of the difference between the two predicted values for the numbers of Ardeidae per

square kilometer. The actual value and the value obtained when assuming a counterfactual

condition that the evacuation zone was not established were compared.

Table 1 MAP estimates and summary statistics via importance sampling with and without the

location error.

Parameter	With the location error					Without the location error				
	Map	Mean	S.D.	CI2.5	CI97.5	Map	Mean	S.D.	CI2.5	CI97.5
Intercept	6.55	7.11	1.48	4.79	10.4	6.54	7.12	1.48	4.76	10.4
Distance	-0.0190	-0.0191	0.00180	-0.0228	-0.0156	-0.0188	-0.0189	0.00183	-0.0227	-0.0155
Water	0.567	0.561	0.0820	0.391	0.717	0.567	0.561	0.0823	0.392	0.716
Evac zone	-5.28	-5.86	1.46	-9.02	-3.59	-5.28	-5.89	1.45	-9.14	-3.60
Year(2015)	0.190	0.203	0.263	-0.292	0.760	0.192	0.218	0.265	-0.294	0.749
Year(2016)	-0.876	-0.863	0.270	-1.37	-0.316	-0.875	-0.853	0.267	-1.36	-0.325

659　Table 2 Predicted values for the number of Ardeidae per square kilometer using the model with

660　and without location error.

Area category	With the location error		Without the location error	
	Mean	S.D.	Mean	S.D.
Outside the evacuation zone	4.57	5.36	4.51	5.29
Inside the evacuation zone	0.0279	0.0307	0.0276	0.0303