■ Title

Distance sampling with vehicle-mounted cameras for the estimation of farmland bird abundance over a wide area

■ Author details

Nao Kumada[1], Keita Fukasawa[1], Akira Yoshioka[2], Naoe Tsuda[3], Hirofumi Ouchi[2]

[1] Biodiversity Division, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

[2] Fukushima Regional Collaborative Research Center, National Institute for Environmental Studies, Miharu, Fukushima, Japan

[3] Center for Climate Change Adaptation, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

■ Abstract

1. Agricultural birds are declining due to farmland abandonment and agricultural intensification, and monitoring approaches, including efficient survey methods, are crucial for the development of effective conservation strategies. Roadside surveys (RS), especially those using vehicle-mounted video cameras, can be used to efficiently search for targets; however, they are limited by the unknown detection range. Distance sampling (DS), in which animal density or abundance is estimated from the distance between the observer and the detected individual, can provide a complimentary approach to address this limitation of RS.

2. We developed a DS model robust to location uncertainty based on video-based RS. We

integrated location errors determined by an independent field test using RS and unmanned

aerial vehicles (UAVs) into a stochastic DS model. We estimated the abundance of herons

and egrets in farmlands in and around the evacuation zone of the Fukushima Daiichi nuclear

power plant accident by combining RS and DS.

3. The video-based RS covered 7,031 km of roads and 24.41 km^2 of farmlands. All herons and

egrets were observed around the evacuation zone and none were detected in the zone. The

predicted abundance of herons and egrets differed substantially between areas inside

(0.0158 ± 0.0174/km^2) and outside of the zone (4.62 ± 5.41/km^2).

4. Integrating location uncertainty into the DS model did not affect predicted heron and egret

population densities (4.62 ± 5.41 vs. 4.66 ± 5.45/km^2, with or without integrating location

uncertainty, respectively). Accordingly, our survey method combining RS and DS is robust

to location uncertainty. The study system (i.e., herons and egrets in farmlands) and the

inability of location error to exceed the size of one farmland separated by levees may

contribute to the accuracy of RS in this study.

5. *Synthesis and Applications:* Combining RS with vehicle-mounted cameras and DS

considering location uncertainty is widely applicable to open land species and can improve

the efficiency of monitoring. When large location errors are expected by conventional

approaches, DS models incorporating location uncertainty determined by RS and UAVs

could be effective for abundance estimation over wide areas.

■ Keywords

43   Roadside survey, Distance sampling, Action camera, Agricultural birds, Abandoned farmlands,

44   UAV, Fukushima Daiichi nuclear power plant accident

■ Introduction

Population sizes of many farmland birds are declining due to agricultural intensification and the

abandonment of cultivation (Fujioka, Don Lee, Kurechi, & Yoshida, 2010; Fuller et al., 1995;

Siriwardena et al., 1998; Stanton, Morrissey, & Clark, 2018). To implement appropriate

conservation strategies for farmland birds, information on their abundances and related

environmental factors is essential. Various methods have been developed for monitoring

farmland birds, including point count, transect count, territory mapping, and capture-recapture

methods (Birrer et al., 2007; Chiron, Filippi-Codaccioni, Jiguet, & Devictor, 2010; Gillings &

Fuller, 1998). Survey methods for birds are often labor-intensive and costly to conduct over a

large area. Cost-effective methods for monitoring are needed to better understand population

patterns and environmental drivers of farmland birds at a large spatial scale and hence to

develop effective conservation measures.

The roadside survey (RS) approach, involving species searches from vehicles on the road,

has the capacity to obtain data for species abundance indices over a wide area at a low cost

(Catry, Moreira, Deus, Silva, & Águas, 2015; Milton & Dean, 1998). Recently, RS using

vehicle-mounted cameras, including Google Street View, has been used to monitor birds in

various habitats (Olea & Mateo-Tomás, 2013; Rousselet et al., 2013). However, RS with digital

images has been limited to areas where there are clearly no gaps in the search area (e.g., birds

nesting on a cliff, Olea & Mateo-Tomás, 2013). A major difficulty in applying vehicle-mounted

cameras to farmland bird monitoring as well as other open-land habitats is that the detection

range is unknown. Ignoring the effect of detection range on the number of detected individuals

can lead to biases in estimates of abundance and the impact of environmental drivers (Buckland,

Marsden, & Green, 2008).

Distance sampling (DS) can resolve the issue of bias in abundance estimation by video-

based RS. DS estimates animal density or abundance from the distance between the observer

and the detected individual, accounting for the decrease in detection probability due to distance

(Buckland et al., 2001; Fox et al., 2017; Hara, Yamaji, Yokota, Thaitakoo, & Sampei, 2018;

Howe, Buckland, Després-Einspenner, & Kühl, 2017; Kvasnes, Pedersen, & Nilsen, 2018). DS

can be used to estimate trends in farmland bird populations in RS. However, there might be

uncertainty in the distance between the observer and the detected individual measured from

video images. For robust abundance estimation, an independent field test of location uncertainty

in DS is required (Hefley, Boyle, & Mohankumar, 2020).

Comparison of the observer–target distances measured by RS and unmanned aerial

vehicles (UAVs) is expected to be a feasible independent field test. UAVs have become an

effective tool for monitoring animal populations and individual locations (Iwamoto, Nogami,

Ichinose, & Takeda, 2022; Witczuk, Pagacz, Zmarz, & Cypel, 2018). Although their application

to large spatial areas remains costly, this method is highly accurate because UAVs can measure

animal positions from almost directly above. If the same individuals are observed by both RS

and UAV, location uncertainty in RS can be quantified. Integrating location uncertainty obtained

by these small-scale measurements into a stochastic DS model will allow us to moderate cost–

accuracy trade-offs and estimate abundance at a large spatial scale.

In Japan, the Fukushima Daiichi nuclear power plant accident in 2011 led to uninhabited

areas and the cessation of agricultural activities due to an evacuation order for the contaminated

area (Yoshioka, Mishima, & Fukasawa, 2015). Although observable effects of ionizing radiation

at the population level on terrestrial organisms were unlikely (UNSCARE, 2017), land

abandonment in the evacuation zone can have indirect effects. These environmental changes are

thought to have a significant impact on the organisms that use farmlands. Therefore, extensive

monitoring in the farmlands throughout the region is needed to evaluate the effect of farmland

abandonment in the evacuation zone. However, with respect to radiation protection, it is

advisable for researchers not to stay in the evacuation zone for a long time, necessitating an

efficient monitoring scheme. Farmland birds such as herons and egrets are suitable indicator

species of the farmland environment. They are the main predators in farmlands and are

susceptible to the abandonment of rice fields (Amano & Katayama, 2009; Lane & Fujioka,

1998). Their large size and conspicuous whitish body color make herons and egrets foraging in

farmlands easily detectable from roads.

The purpose of this study was to estimate the population density of herons and egrets in

farmlands in and around the evacuation zone of the Fukushima Daiichi nuclear power plant

accident by RS. In particular, we surveyed the locations of herons and egrets by using a vehicle-

mounted camera and then estimated population densities by using a DS model incorporating

location errors. The magnitude of location errors for herons and egrets by RS was determined by

a field test comparing RS with UAV-mounted cameras. To evaluate the importance of

accounting for location uncertainty, population densities were compared between analyses with

and without location errors. We demonstrated the impact of farmland abandonment due to the

evacuation order by comparing the population density with a counter-factual scenario without

an evacuation order.

■ Materials and methods

Field test of location errors of herons and egrets using UAVs

Location errors of the vehicle-mounted camera were validated by comparison with the

accurate values determined by UAV images (UAV-truth). The survey was conducted on

farmlands in two regions, the district of Inashiki in Ibaraki Prefecture and the city of

Minamisoma in Fukushima Prefecture. These farmlands are about 5 square kilometers and are

used for foraging by herons and egrets. The survey was conducted in Inashiki in July 2019 and

May 2020 and in Minamisoma in June 2020. During the survey period, herons and egrets in the

farmlands were recorded simultaneously by a vehicle-mounted video camera (VIRB X; Garmin

Ltd.) and a video camera in a UAV (Phantom 4; DJI). The GPS trajectories of vehicle-mounted

video cameras were recorded by VIRB X simultaneously.

Locations of herons and egrets were determined from the movies captured by the vehicle-

mounted cameras. If the researcher observed herons and egrets in the video, the movement

trajectory information recorded at the time the video was taken was used to determine the

location of the camera. Then, using information on the camera location with orthographical

aerial imagery layers as background maps (GSI Tiles) and views surrounding herons and egrets

(paddy levees, houses, telegraph poles, etc.) in the video image, the locations of herons and

egrets were manually determined and recorded using QGIS ver.3.4.3.

To obtain correct locations of herons and egrets, georeferenced images obtained by the

UAV were used. Georeferencing was performed using Georeferencer (Transformation type:

Projective, Resampling method: Linear) in QGIS. The measurement errors of locations obtained

by vehicle-mounted cameras were calculated as the Euclidean distances between the locations

of the herons and egrets determined by the vehicle-mounted cameras and the UAV-truth values.

Roadside survey of herons and egrets around the evacuation zone

The RS with vehicle-mounted cameras of abandoned and cultivated farmlands was

conducted in eastern Fukushima, which includes the evacuation zone of the Fukushima Daiichi

nuclear power plant accident (Fig. 1). The evacuation zone was designated after the Fukushima

Daiichi nuclear power plant accident in March 2011 and is divided into three categories

according to radiation levels: Difficult-to-Return Area, Restricted-Residential Area, and

Preparing for the Lifting of the Evacuation Order Area (Cabinet Office, Government of Japan,

2013). In all three categories, cropland cultivation did not resume until 2016, except in some

test fields. The study area covered farmlands within 18 municipalities located inside and outside

the evacuation zone. Farmland polygons in this area were created from aerial photographs taken

in 2006–2015 (see Appendix 1 for detailed methods). The farmland polygons included upland

fields, pastures, and abandoned farmlands, and more than 70% of the farmland area was rice

fields in Fukushima prefecture before the nuclear power plant accident (in 2010) (Ministry of

Agriculture, Forestry and Fisheries, 2020).

From early May to early July between 2014 and 2016, roadside video recording surveys

were conducted on farmlands by vehicle-mounted cameras. Video cameras with GPS (VIRB

Elite or VIRB X; Garmin Ltd.) were mounted on the side windows of a vehicle, which was

driven on roads adjacent to farmlands for video recording (Fig. 2). Videos recorded by the two

camera models (VIRB Elite or VIRB X) simultaneously at the same location showed no obvious

differences. Although the video cameras were mounted on both sides of a car in many cases,

some surveys were carried out using a video camera mounted on one side only. The GPS

trajectories of the car were also recorded simultaneously. Videos were shot at 1920×1080 p/30

fps for all models and saved in mp4 file format. The speed of the car during the survey varied

depending on the speed limit (60 km/h or less) and road congestion; however, the frame rate

was sufficiently high to detect herons and egrets under all speeds in the study.

To identify farmlands within the viewshed from the vehicle-mounted cameras, farmland

polygons in the direction of the camera and not hidden by obstacles such as forest and houses

were selected as follows (Fig. 3). First, we extracted the farmland polygons in the direction of

the camera and intersecting within a straight-line distance of 10 m from the trajectories of the

car; these were defined as the viewshed. Ten meters was set as an approximate threshold for

adjacency (i.e., no obstacles) between the road and farmland. Second, we sequentially included

farmlands that were within 10 meters from the viewshed obtained in the previous step.

Farmlands were not included in the viewshed when the distance from the movement trajectory

was more than 500 m because this was too far to detect herons and egrets. We visually verified

that the selection criteria could effectively extract farmlands within the viewshed of the camera.

We played the videos back and searched for herons and egrets in farmlands. Then, we

determined the locations on the map following the same method used to test location errors

described above. Because it was difficult to identify species from the video images correctly,

except for grey heron, we merged all species of herons and egrets for further analyses. The

following herons and egrets may be found in the study area: grey heron *Ardea cinerea*, great

egret *A. cinerea*, intermediate egret *A. intermedia*, little egret *Egretta garzetta*, and cattle egret

Bubulcus ibis. All of these herons and egrets are colonial and breed in the summer. In addition,

flooding status (i.e., flooded or not) was recorded for the rice fields where herons and egrets

were found.

Statistical analysis

We employed a fully model-based approach to estimate the abundance of herons and egrets

and the effect of land abandonment based on a DS considering individual location uncertainty.

As our observations consisted of the camera-based DS results with location uncertainty and the

independent field test of location errors based on correct locations determined by UAV, we

integrated the likelihoods and estimated all parameters simultaneously.

A common approach for model-based DS without location uncertainty is to treat the

occurrence of individuals in relation to the distance from an observer as a realization of an

inhomogeneous Poisson process (IPP) (Johnson, Laake, & Ver Hoef, 2010). The likelihood of

DS without location uncertainty is expressed by the following equation (Johnson, Laake, & Ver

Hoef, 2010).

$$L_{IPP}(\alpha,\ \beta,\gamma|\mathbf{Y}) = \prod_{i=1}^{n} \lambda(\mathbf{y}_i|\alpha,\ \beta\ )\delta(\mathbf{y}_i|\gamma)\exp\left(-\int_{\mathcal{S}} \lambda(\mathbf{z}|\alpha,\ \beta\ )\delta(\mathbf{z}|\gamma)d\mathbf{z}\right)$$

(eqn. 1)

where $\mathbf{Y} = (\mathbf{y}_1, \mathbf{y}_2, ..., \mathbf{y}_n)$ are true locations of observed individuals, (α, β, γ) are model parameters,

and $\mathcal{S}$ is the spatial domain(s) that is potentially observable. $\lambda(\mathbf{z}|\alpha,\ \beta\ )$ is an intensity function of the

population density at location $\mathbf{z}$ and is commonly specified as follows.

$\lambda(\mathbf{z}|\alpha, \beta) = \exp(\alpha + \beta\mathrm{x}(\mathbf{z}))$

where α is the intercept, $\beta = (\beta_1,...,\beta_p)'$ is a vector of regression coefficients, and x($\mathbf{z}$) is a vector of

covariates on population density at location $\mathbf{z}$. $\delta(\mathbf{z}|\gamma)$ is the detection probability, which is expressed

as a thinning function as follows.

$\delta(\mathbf{z}) = \exp(-\gamma D(\mathbf{z}))$

where $D(\mathbf{z})$ is the Euclidian distance from an observer to location $\mathbf{z}$.

Let us consider the case when observed location $\mathbf{Y}' = (\mathbf{y}'_1, \mathbf{y}'_2, ..., \mathbf{y}'_n)$ is assumed to vary

around the true location $\mathbf{Y}$ due to location error. The likelihood of IPP incorporating location

uncertainty can be obtained by integrating out $\mathbf{y}_i$ over the surveyed region (Hefley et al., 2020).

$$L(\alpha, \beta, \gamma, \boldsymbol{\sigma}|\mathbf{Y}') = \prod_{i=1}^{n} \int_{\mathcal{S}_i} p_{\mathrm{err}}(\mathbf{y}'_i|\mathbf{y}_i, \boldsymbol{\sigma})\lambda(\mathbf{y}_i)\delta(\mathbf{y}_i)d\mathbf{y}_i \exp\left(-\int_{\mathcal{S}} \lambda(\mathbf{z})\delta(\mathbf{z})d\mathbf{z}\right)$$

(eqn. 2, parameters α, $\boldsymbol{\beta}$, γ were not included on the right side for simplicity)

where $p_{\mathrm{err}}(\mathbf{y}'|\mathbf{y}, \boldsymbol{\sigma})$ is a probability distribution of location error, $\boldsymbol{\sigma}$ is a vector of location error

parameters, and $\mathcal{S}_i$ is subset of $\mathcal{S}$ in which $\mathbf{y}_i$ exist. For simplicity, we assume that the location error

is isotropic and normally distributed.

$p_{\mathrm{err}}(\mathbf{y}'|\mathbf{y}, \boldsymbol{\sigma}) = N_2(\mathbf{y}'|\mathbf{y}, f(\mathbf{y}', \mathbf{y}, \boldsymbol{\sigma})\mathbf{I})$ (eqn. 3)

where $f(\mathbf{y}', \mathbf{y}, \boldsymbol{\sigma})$ is a function of the standard deviation depending on the true location. In general,

the greater the distance from the observer, the lower the accuracy of the observation position. The

relationship was assumed to be described by a power function of Euclidean distances between $\mathbf{y}'$

and $\mathbf{y}$, $D(\mathbf{y}', \mathbf{y})$, with two parameters $\boldsymbol{\sigma} = (\sigma_0, \sigma_1)$ as follows.

$f(\mathbf{y}, \mathbf{y}, \boldsymbol{\sigma}) = \sigma_0 D(\mathbf{y}, \mathbf{y})^{\sigma_1}$ (eqn. 4)

The likelihood function required integration over spatial domains, which is difficult to solve

analytically. A practical approach is discretizing spatial domains into grid cells and replacing

integration   with   piecewise   summation   as   follows.

$$\int_{\mathcal{S}_i} \mathrm{p_{err}}\left(\mathbf{y}_i^{'} \mid \mathbf{y}_i\right) \lambda(\mathbf{y}_i) \delta(\mathbf{y}_i) \mathrm{d}\mathbf{y}_i \approx \sum_{q \subset \mathcal{S}_i} A_q \mathrm{p_{err}}\left(\mathbf{y}_i^{'} \mid \mathbf{y}_q\right) \lambda(\mathbf{y}_q) \delta(\mathbf{y}_q)$$

$$\int_{\mathcal{S}} \lambda(\mathbf{z}) \delta(\mathbf{z}) d\mathbf{z} \approx \sum_{q \subset \mathcal{S}} A_q \lambda(\mathbf{z}_q) \delta(\mathbf{z}_q)$$

where $\mathbf{z}_q$ and A_q are the centroid and area of grid cell q, respectively.

The location error parameter $\boldsymbol{\sigma}$ in the likelihood function described above is not identifiable

without an independent survey of location error. In this study, the location error and UAV-truth were

included simultaneously in the DS model for the estimation of all parameters. When the true location

is available, the error distribution (eqn. 3 and 4) can be written equivalently by the Rayleigh distribution

(i.e., Weibull distribution with shape parameter 2) for the Euclidean distance between the observed and

true location.

$$\mathrm{p_{err}}(D \mid \boldsymbol{\sigma}) = D/\left(\sigma_0^2 D^{2\sigma_1}\right) \exp\left(-D/\left(2\sigma_0^2 D^{2\sigma_1}\right)\right)$$

226   Given the vector of distances obtained by the field test $D(\mathbf{y}\square, \mathbf{y}) = (D(y'_1, y_1), ..., D(y'_m, y_m))$, the joint

227   likelihood is given as follows.

$$L\left(\alpha, \beta, \gamma, \boldsymbol{\sigma} \mid \mathbf{Y}^{'}, \mathbf{D}(\mathbf{y}, \mathbf{y})\right)$$

$$= \prod_{i=1}^{n} \sum_{q \subset \mathcal{S}_i} A_q \mathrm{p_{err}}\left(\mathbf{y}_i^{'} \mid \mathbf{y}_q\right) \lambda(\mathbf{y}_q) \delta(\mathbf{y}_q) \exp\left(-\sum_{q \subset \mathcal{S}} A_q \lambda(\mathbf{z}_q) \delta(\mathbf{z}_q)\right) \prod_{j=1}^{m} \mathrm{p_{err}}(D_j)$$

Both the maximum likelihood method and Bayesian method can be used for parameter

estimation. When the Bayesian method is applied, prior information and constraints on

parameters can be incorporated as a prior distribution.

We estimated the effects of the evacuation order (evacuation zone or not) (Cabinet Office,

Government of Japan, 2013), the area of water bodies (rivers and lakes) (Ogawa et al., 2013)

around the farmlands, and the year of the study on the abundance of herons and egrets. We

defined farmland polygons within the viewshed of the camera as a multi-part domain S and

discretized it into grid cells with a resolution of 10 m. Then, these variables and the Euclidean

distances from the camera were assigned to each cell. Even when location error is taken into

account, the error is reasonably assumed to be within a farmland polygon because paddy levees

act as sharp borders dividing farmland polygons. Therefore, we assumed that the subdomain S_i

containing the true location of the observed individual i was restricted to a farmland polygon

where the individual was observed.

In this study, we applied the Bayesian default prior (Gelman, Jakulin, Pittau, & Su, 2007)

to stabilize the coefficients and to prevent overfitting. The IPP model is often limited by

overfitting, and it is common to apply regression shrinkage techniques. We applied a weakly

informative prior, Student's t-test (df = 5, mean = 0, scale = 2.5), to regression coefficients $\boldsymbol{\beta}$ to

avoid overfitting. For other parameters, improper flat priors p(θ) = 1 were used. The maximum

a posteriori (MAP) estimate of each parameter was calculated using a quasi-Newton algorithm.

To estimate credible intervals of parameters, we obtained posterior samples by importance

sampling (Gelman et al., 2013) with a proposal distribution that approximates the posterior. For

the proposal distribution, we used a multivariate t-distribution with four degrees of freedom

whose mean vector and scale matrix are the MAP estimate and the inverse of the Hessian matrix

at the MAP, respectively. We drew 10,000 samples from the proposal distribution and obtained

3,000 posterior samples.

We calculated the abundance of herons and egrets by applying MAP estimates of each

parameter to grid cells in farmland polygons throughout the study area, with and without

accounting for measurement error, and compared the two abundance results. The predicted

values were aggregated for each 1-km square grid. To evaluate the effect of farmland

abandonment in the evacuation zone on the population size of herons and egrets, we predicted

the population size in the study area under a counter-factual scenario in which no evacuation

order had been given. All analyses in this study were implemented using R (R Core Team, 2021)

(Appendix 2).

■   **Results**

Field test of location error of herons and egrets

In total, 78 herons and egrets (24 in July 2019, 31 in May 2020 in Inashiki and 23 in June

2020 in Minamisoma) were located with GIS by the vehicle-mounted camera and the camera on

the UAV. The Euclidean distance between the locations of the herons and egrets determined by

the vehicle-mounted camera and the true locations determined by the UAV was defined as the

measurement error (Fig. 4).

RS with vehicle-mounted cameras and abundance estimates of herons

and egrets

The total distances traveled while recording with cameras were 3,675 km in the evacuation

zone and 3,356 km outside the evacuation zone. The areas of farmland polygons (including

abandoned fields in response to the evacuation order), selected as the viewshed, were 10.07 km^2

in and 14.34 km^2 around the evacuation zone.

A total of 56 individual herons and egrets were observed around the evacuation zone. No

herons or egrets were observed in the evacuation zone (Fig. 5). All herons and egrets were

found in flooded rice fields. The frequency distribution of observed distances to herons and

egrets was more concentrated near zero than the distribution of distances to roads for all grid

cells of farmland polygons. Although all polygons had a median distance of 133.3 m from roads,

the polygons containing heron and egret discovery sites had a median distance of 47.1 m and a

maximum of 104.4 m (Fig. 6).

The MAP estimates and summary statistics obtained by importance sampling are shown in

Table 1. The water body area had a positive association with the abundance of herons and egrets.

Abundance was lower in the evacuation zone than outside the zone. Although the signs (i.e.,

positive or negative) of parameter estimates were similar between the models with and without

the location error, the parameter values differed slightly. The distance from the camera had a

negative association with the probability of detection (Fig. 4). The probability decreased to half

at 35.4 m (95% credible interval: 29.6–42.8 m) from the camera. The scaling factor and power

value when measurement errors are expressed as a power function of the distance from the

camera were 0.644 and 0.614. The measurement errors tended to increase with increasing

distance from the camera on the car but exhibited a concave downward trend.

The predicted abundance of herons and egrets were substantially larger outside the

evacuation zone than inside the zone (Table 2, Fig. 7). This trend did not differ between the

models with and without location error. The predicted total population of herons and egrets in

the study area was 7,820 individuals. Assuming that the evacuation zone was never designated,

the predicted population would have increased by 2,647 individuals to 10,467 individuals (Fig.

8).

■   Discussion

We developed an efficient method for monitoring farmland birds using RS with vehicle-

mounted cameras and DS. By performing DS considering location error, we resolved the issue

of the lack of a clear observation range of RS. The spatially explicit approach enabled us to

estimate the impact of large-scale land abandonment on heron and egret populations. The

estimated number of foraging individuals around the evacuation zone did not deviate

significantly from the value obtained by a researcher's count in a previous study of

approximately 30 individuals/km^2 (Lane & Fujioka, 1998) and is considered a reasonable

estimate.

While the RS approach benefits from the ability to efficiently survey a wide area, the use

of vehicle-mounted cameras will further improve efficiency. For example, in this study, 237

hours of video was recorded to search for herons and egrets. In a field survey, experts would be

required for the full time period; however, the RS method with vehicle-mounted cameras does

not require expertise in bird identification in the field. In addition, the work of sorting farmlands

without herons and egrets, accounting for the majority of each video, does not require expertise;

instead, experts only need to check images with suspected herons and egrets. Crowdsourcing

this sorting process can promote the participation of birders in remote areas who are interested

in the survey. In the future, automated counting may be possible (Norouzzadeh et al., 2018).

An accurate distance between the observer and target is required for DS to estimate

density; accordingly, when the distance accuracy could not be guaranteed, it is necessary to

account for measurement error in DS (Hefley et al., 2020). The incorporation of measurement

error in the location of egrets and herons did not significantly change density estimates. This

result showed that the distance accuracy of RS with vehicle-mounted cameras in our study was

sufficient for density estimation by DS. The study system can explain the good performance of

RS. The herons and egrets were found in farmlands, and the location error could not be larger

than the size of one farmland separated by paths or levees. However, use of a UAV to estimate

location errors for incorporation into a DS model may be an effective approach in situations

where measurement errors are likely. RS by vehicle-mounted cameras may be a feasible strategy

for various situations by using this method.

The direct effects of radiation exposure from the Fukushima Daiichi nuclear power plant

accident on wildlife populations and ecosystems are not large enough to be observable at the

population level (UNSCEARE 2017). However, the evacuation orders in response to the nuclear

accident have resulted in short-term changes in land use and human activities (Tamaoki, 2016;

Yoshioka et al., 2015), and these changes may have important impacts on wildlife populations

and ecosystems (Fujimoto, Mitsunaga, & Takeuchi, 2015; Lyons, Okuda, Hamilton, Hinton, &

Beasley, 2020; Matsushima, Ihara, Inaba, & Horiguchi, 2021; Yoshioka et al., 2015). The strong

negative impact of the evacuation order on heron and egret populations can be attributed to the

decrease in food availability due to the widespread abandonment of farmlands. Farmland

abandonment leads to the growth of dense vegetation, which is not a common foraging

environment for herons and egrets (Katayama, Osawa, Amano, & Kusumoto, 2015; Maeda,

2005). In addition, herons and egrets feed on fish and frogs in flooded rice fields (Tojo, 1996),

which can be maintained by cultivation (Maeda, 2001). Our results clearly demonstrated that

cropland abandonment in evacuation zones affects the distribution of birds over a wide area.

These findings are consistent with the observation that the long-term abandonment of farmland

reduces the biodiversity of farmlands (Koshida & Katayama, 2018). Since 2016, the evacuation

zone has been reduced and agricultural cultivation has begun to resume. In the future, the effect

of resumption should be tested by continued monitoring following the protocols developed in

this study.

The detection probability depends on the distance from the observer as well as various

covariates, in which case multiple-covariate DS is appropriate (Marques & Buckland, 2003;

Thomas et al., 2010). In this study, we did not consider the evacuation zone as a covariate,

despite the potential effect of this parameter on the detection probability, because no herons or

egrets were observed in the evacuation zone. However, we believe the absence of herons and

egrets in the evacuation zone reflects differences in true population densities resulting from the

abandonment of farmland because dense vegetation is generally not suitable for foraging by

these species (Katayama, Osawa, Amano, & Kusumoto, 2015; Maeda, 2005). These species

were not found in the evacuation zone, even near roads in which detectability is less affected by

the vegetation.

The escape behavior of birds during surveys might result in biases in population estimates

(Hutto & Young, 2003; Roberts & Schnell, 2006). At farmlands bordering roads, herons and

egrets may fly away to avoid the approach of vehicles carrying cameras. Herons and egrets at a

distance from the camera can therefore be overlooked, which can lead to biases in the detection

function and population estimates. However, the videos obtained in this study did not reveal any

herons and egrets in flight, which could be a response to an approaching vehicle. In fact,

waterbirds are not substantially impacted by approaching vehicles (McLeod, Guay, Taysom,

Robinson, & Weston, 2013). Similarly, in this study, the effect of approaching vehicles was

considered negligible for herons and egrets. If this method is applied to broader taxa, the use of

fisheye lenses or 360-degree cameras may be a viable option to capture individuals before they

are affected by an observer.

In our survey, we estimated the number of herons and egrets using farmlands; however,

these estimates cannot be interpreted as total counts for the study area. The survey was

conducted during the breeding season, and one member of each nesting pair always stays in the

nest. In addition, because herons fly over a wide area in search of good foraging sites, it is

thought that they frequently move in and out of the evacuation zone and travel to and from the

study area. It is necessary to consider whether the number of herons obtained in this study is an

indicator of the quality of the farmland as a foraging site. However, we believe that the density

of herons and egrets foraging in farmlands is a viable indicator of biodiversity in agricultural

land because it is a common parameter for assessing the quality of farmlands as foraging sites

(Amano & Katayama, 2009; Katayama et al., 2019; Lane & Fujioka, 1998).

Another limitation of our study is that we were unable to identify the herons and egrets at

the species level. We considered the total abundance of herons and egrets. The five species in

the study area have overlapping ecological niches; they forage in farmlands and often share

colonies (Mashiko & Toquenaga, 2014). In addition, the number of herons and egrets is related

to the abandonment of farmland and taxon diversity and can therefore be used as an indicator of

species richness on farmlands (Katayama, Mashiko, Koshida, & Yamaura, 2021; Katayama et

al., 2019; Koshida & Katayama, 2018). In the future, technological improvements such as

increased camera resolution may make it possible to identify species and evaluate responses.

The combination of RS using vehicle-mounted cameras and DS made it possible to determine

the abundance of large farmland birds and their limiting factors in and around the evacuation

zone. This method could be applied to the monitoring of large birds and mammals in open areas,

including tidal flats, grasslands, and agricultural fields; some of these species are endangered

and others cause agricultural conflicts. Furthermore, when combined with an infrared camera,

this method might be used to estimate the population of nocturnal mammals. In recent years,

open street-level imagery, such as Mapillary (https://www.mapillary.com), has become widely

available, with the potential to minimize field survey costs. With advances in image recognition

technology for species detection, identification (Norouzzadeh et al., 2021), and distance

estimation (Johanns, Haucke, & Steinhage, 2022), our approach may enable global monitoring.

■ Authors' contributions

399 N.K., K.F. and A.Y. conceived the ideas and designed the methodology; All the authors

400 collected the data; N.K. and K.F. analysed the data; N.K., K.F. and A.Y. led the writing of the

401 manuscript. All the authors reviewed and contributed critically to the draft and gave final

402 approval for publication.

■ Acknowledgements

405 We are grateful to Fukushima Robot Test Field for helping field work. Yoshio Mishima

406 contributed to create the farmland polygon data but declined to join as a co-author. This

407 study was partially supported by JSPS KAKENHI Grant number 21H03656.

■ References

410 Amano, T., & Katayama, N. (2009). Hierarchical movement decisions in predators: Effects of

411 foraging experience at more than one spatial and temporal scale. *Ecology*, *90*(12), 3536–

412 3545. doi: 10.1890/08-1910.1

413 Birrer, S., Spiess, M., Herzog, F., Jenny, M., Kohli, L., & Lugrin, B. (2007). The Swiss agri-

414 environment scheme promotes farmland birds: but only moderately. *Journal of*

415 *Ornithology*, *148*(2), 295–303. doi: 10.1007/s10336-007-0237-y

416 Buckland, S.T., Anderson, D. R., Burnham, K. P., Laake, J. L., Borchers, D. L., & Thomas, L.

417 (2001). *Introduction to distance sampling: estimating abundance of biological populations.*

418 Oxford (United Kingdom) Oxford Univ. Press.

419 Buckland, S. T., Marsden, S. J., & Green, R. E. (2008). Estimating bird abundance: Making

420 methods work. *Bird Conservation International*, *18*, S91–S108. doi:

421 10.1017/S0959270908000294

422 Cabinet Office, Government of Japan. About revision of the evacuation zone (in Japanese).

423 (2013). http://www.meti.go.jp/earthquake/nuclear/pdf/131009/131009_02a.pdf. Accessed

424 17 January 2023.

425 Catry, F. X., Moreira, F., Deus, E., Silva, J. S., & Águas, A. (2015). Assessing the extent and

426 the environmental drivers of Eucalyptus globulus wildling establishment in Portugal:

427 results from a countrywide survey. *Biological Invasions*, *17*(11), 3163–3181. doi:

428 10.1007/s10530-015-0943-y

429 Chiron, F., Filippi-Codaccioni, O., Jiguet, F., & Devictor, V. (2010). Effects of non-cropped

430 landscape diversity on spatial dynamics of farmland birds in intensive farming systems.

431 *Biological Conservation*, *143*(11), 2609–2616. doi:

432 https://doi.org/10.1016/j.biocon.2010.07.003

433 Fox, C. H., Huettmann, F. H., Harvey, G. K. A., Morgan, K. H., Robinson, J., Williams, R., &

434 Paquet, P. C. (2017). Predictions from machine learning ensembles: Marine bird

435 distribution and density on Canada's Pacific coast. *Marine Ecology Progress Series*, *566*,

436 199–216. doi: 10.3354/meps12030

437 Fujimoto, R., Mitsunaga, T., & Takeuchi, M. (2015). Influence of evacuation orders on the

438 occurrence of the wild boar Sus scrofa in farmlands in the northern evacuation zone of the

439 accident at Tokyo Electric Power Company's Fukushima Daiichi Nuclear Power Station

440 (in Japanese with English abstract). *Mammalian Science*, *55*(2), 145–154.

441 Fujioka, M., Don Lee, S., Kurechi, M., & Yoshida, H. (2010). Bird use of rice fields in Korea

442 and Japan. *Waterbirds*, *33*(SPEC.ISSUE.1), 8–29. doi: 10.1675/063.033.s102

443 Fuller, R. J., Gregory, R. D., Gibbons, D. W., Marchant, J. H., Wilson, J. D., Baillie, S. R., &

444 Carter, N. (1995). Population Declines and Range Contractions among Lowland Farmland

445 Birds in Britain. *Conservation Biology*, *9*(6), 1425–1441. doi: 10.1046/j.1523-

446 1739.1995.09061425.x

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013).

Bayesian Data Analysis (third edition). New York: Chapman and Hall/CRC

Gelman, A., Jakulin, A., Pittau, M. G., & Su, Y.-S. (2007). A weakly informative default prior

distribution for logistic and other regression models. *The Annals of Applied Statistics*, *2*.

doi: 10.2139/ssrn.1010421

Gillings, S., & Fuller, R. J. (1998). Changes in bird populations on sample lowland English

farms in relation to loss of hedgerows and other non-crop habitats. *Oecologia*, *116*(1),

120–127. doi: 10.1007/s004420050570

Hara, Y., Yamaji, K., Yokota, S., Thaitakoo, D., & Sampei, Y. (2018). Dynamic wetland

mosaic environments and Asian openbill habitat creation in peri-urban Bangkok. *Urban

Ecosystems*, *21*(2), 305–322. doi: 10.1007/s11252-017-0718-6

Hefley, T. J., Boyle, W. A., & Mohankumar, N. M. (2020). *Accounting for location uncertainty

in distance sampling data*. (2020). Retrieved from http://arxiv.org/abs/2005.14316

Howe, E. J., Buckland, S. T., Després-Einspenner, M. L., & Kühl, H. S. (2017). Distance

sampling with camera traps. *Methods in Ecology and Evolution*, *8*(11), 1558–1565. doi:

10.1111/2041-210X.12790

Hutto, R. L., & Young, J. S. (2003). On the Design of Monitoring Programs and the Use of

Population Indices: A Reply to Ellingson and Lukacs. *Wildlife Society Bulletin*, 31(3),

903–910. Retrieved from http://www.jstor.org/stable/3784617

Iwamoto, M., Nogami, S., Ichinose, T., & Takeda, K. (2022). Unmanned aerial vehicles as a

useful tool for investigating animal movements. *Methods in Ecology and Evolution*, *13*(5),

969–975. doi: 10.1111/2041-210X.13829

Johanns, P., Haucke, T., & Steinhage, V. (2022). Automated distance estimation for wildlife

camera trapping. *Ecological Informatics*, *70*(February), 101734. doi:

10.1016/j.ecoinf.2022.101734

Johnson, D. S., Laake, J. L., & Ver Hoef, J. M. (2010). A model-based approach for making

ecological inference from distance sampling data. *Biometrics*, *66*(1), 310–318. doi:

10.1111/j.1541-0420.2009.01265.x

Katayama, N., Mashiko, M., Koshida, C., & Yamaura, Y. (2021). Agriculture, Ecosystems and

Environment Effects of rice field abandonment rates on bird communities in mixed

farmland – woodland landscapes in Japan. *Agriculture, Ecosystems and Environment*,

319(June), 107539. doi: 10.1016/j.agee.2021.107539

Katayama, N., Osada, Y., Mashiko, M., Baba, Y. G., Tanaka, K., Kusumoto, Y., … Natuhara, Y.

(2019). Organic farming and associated management practices benefit multiple wildlife

taxa: A large-scale field study in rice paddy landscapes. *Journal of Applied Ecology*, *56*(8),

1970–1981. doi: 10.1111/1365-2664.13446

Katayama, N., Osawa, T., Amano, T., & Kusumoto, Y. (2015). Are both agricultural

intensification and farmland abandonment threats to biodiversity? A test with bird

communities in paddy-dominated landscapes. *Agriculture, Ecosystems and Environment*,

214, 21–30. doi: 10.1016/j.agee.2015.08.014

Koshida, C., & Katayama, N. (2018). Meta-analysis of the effects of rice field abandonment on

biodiversity in Japan. *Conservation Biology*, *32*(6), 1392–1402. doi: 10.1111/cobi.13156

Kvasnes, M. A. J., Pedersen, H. C., & Nilsen, E. B. (2018). Quantifying suitable late summer

brood habitats for willow ptarmigan in Norway. *BMC Ecology*, *18*(1), 1–13. doi:

10.1186/s12898-018-0196-6

Lane, S. J., & Fujioka, M. (1998). The impact of changes in irrigation practices on the

distribution of foraging egrets and herons (Ardeidae) in the rice fields of central Japan.

Biological Conservation, *83*(2), 221–230. doi: 10.1016/S0006-3207(97)00054-2

Lyons, P. C., Okuda, K., Hamilton, M. T., Hinton, T. G., & Beasley, J. C. (2020). Rewilding of

Fukushima's human evacuation zone. *Frontiers in Ecology and the Environment*, *18*(3),

127–134. doi: 10.1002/fee.2149

Maeda, T. (2001). Patterns of bird abundance and habitat use in rice fields of the Kanto Plain,

central Japan. *Ecological Research*, *16*(3), 569–585. doi: 10.1046/j.1440-

1703.2001.00418.x

Maeda, T. (2005). Bird use of rice field strips of varying width in the Kanto Plain of central

Japan. *Agriculture, Ecosystems and Environment*, *105*(1–2), 347–351. doi:

10.1016/j.agee.2004.02.007

Marques, F. F. C., & Buckland, S. T. (2003). Incorporating Covariates into Standard Line

Transect Analyses. *Biometrics*, *59*(4), 924–935. doi: 10.1111/j.0006-341X.2003.00107.x

Mashiko, M., & Toquenaga, Y. (2014). Grey Heron (Ardea cinerea) Expansion Promotes the

Persistence of Mixed-species Heron Colonies. *WATERBIRDS*, *37*(4), 362–370. doi:

10.1675/063.037.0403

Matsushima, N., Ihara, S., Inaba, O., & Horiguchi, T. (2021). Assessing the impact of large-

scale farmland abandonment on the habitat distributions of frog species after the

Fukushima nuclear accident. *Oecologia*, *196*(4), 1219–1232. doi: 10.1007/s00442-021-

04991-y

McLeod, E. M., Guay, P. J., Taysom, A. J., Robinson, R. W., & Weston, M. A. (2013). Buses,

cars, bicycles and walkers: The influence of the type of human transport on the flight

responses of waterbirds. *PLoS ONE*, *8*(12). doi: 10.1371/journal.pone.0082008

Milton, S. J., & Dean, W. R. J. (1998). Alien plant assemblages near roads in arid and semi-arid

South Africa. *Diversity and Distributions*, *4*(4), 175–187. doi: 10.1046/j.1472-

4642.1998.00024.x

Norouzzadeh, M. S., Morris, D., Beery, S., Joshi, N., Jojic, N., & Clune, J. (2021). A deep
active learning system for species identification and counting in camera trap images.
Methods in Ecology and Evolution, *12*(1), 150–161. doi: 10.1111/2041-210X.13504

Norouzzadeh, M. S., Nguyen, A., Kosmala, M., Swanson, A., Palmer, M. S., Packer, C., &
Clune, J. (2018). Automatically identifying, counting, and describing wild animals in
camera-trap images with deep learning. *Proceedings of the National Academy of Sciences*
of the United States of America, *115*(25), E5716–E5725. doi: 10.1073/pnas.1719367115

Ogawa, M., Takenaka, A., Kadoya, T., Ishihama, F., Yamano, H., & Akasaka, M. (2013). A
comprehensive new land-use classification map for Japan for biodiversity assessment and
species distribution modeling (in Japanese with English abstract). *Japanese Journal of*
Conservation Ecology, *18*(1), 69–76.

Olea, P. P., & Mateo-Tomás, P. (2013). Assessing Species Habitat Using Google Street View:
A Case Study of Cliff-Nesting Vultures. *PLoS ONE*, *8*(1), 1–8. doi:
10.1371/journal.pone.0054582

R Core Team (2021). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Roberts, J. P., & Schnell, G. D. (2006). Comparison of survey methods for wintering grassland
birds. *JOURNAL OF FIELD ORNITHOLOGY*, *77*(1), 46–60. doi: 10.1111/j.1557-
9263.2006.00024.x

Rousselet, J., Imbert, C.-E., Dekri, A., Garcia, J., Goussard, F., Vincent, B., … Rossi, J.-P.
(2013). Assessing Species Distribution Using Google Street View: A Pilot Study with the
Pine Processionary Moth. *PLoS ONE*, *8*(10), e74918. doi: 10.1371/journal.pone.0074918

Siriwardena, G. M., Baillie, S. R., Buckland, S. T., Fewster, R. M., Marchant, J. H., & Wilson, J.
D. (1998). Trends in the abundance of farmland birds: A quantitative comparison of
smoothed Common Birds Census indices. *Journal of Applied Ecology*, *35*(1), 24–43. doi:
10.1046/j.1365-2664.1998.00275.x

Stanton, R. L., Morrissey, C. A., & Clark, R. G. (2018). Analysis of trends and agricultural
drivers of farmland bird declines in North America: A review. *Agriculture, Ecosystems*
and Environment, *254*(December 2017), 244–254. doi: 10.1016/j.agee.2017.11.028

Tamaoki, M. (2016). Studies on radiation effects from the Fukushima nuclear accident on wild
organisms and ecosystems. *Global Environmental Research*, *20*, 73–82.

Thomas, L., Buckland, S. T., Rexstad, E. A., Laake, J. L., Strindberg, S., Hedley, S. L., …
Burnham, K. P. (2010). Distance software: design and analysis of distance sampling
surveys for estimating population size. *Journal of Applied Ecology*, *47*(1), 5–14. doi:
10.1111/j.1365-2664.2009.01737.x

Tojo, H. (1996). Habitat Selection, Foraging Behaviour and Prey of Five Heron Species in

Japan. *Japanese Journal of Ornithology*, *45*(3), 141–158. doi: 10.3838/jjo.45.141

UNSCEARE. (2017) FUKUSHIMA 2017 WHITE PAPER.

https://www.unscear.org/unscear/en/publications/whitepapers.html Accessed 17 January

2023.

Witczuk, J., Pagacz, S., Zmarz, A., & Cypel, M. (2018). Exploring the feasibility of unmanned

aerial vehicles and thermal imaging for ungulate surveys in forests - preliminary results.

International Journal of Remote Sensing, *39*(15–16), 5504–5521. doi:

10.1080/01431161.2017.1390621

Yoshioka, A., Mishima, Y., & Fukasawa, K. (2015). Pollinators and Other Flying Insects inside

and outside the Fukushima Evacuation Zone. *PLOS ONE*, *10*(11), e0140957. doi:

10.1371/journal.pone.0140957

 Figures and tables

572   Figure 1 Map of survey routes (black lines) in Eastern Fukushima prefecture. Dark grey area

573   shows the evacuation zone of the Fukushima Daiichi nuclear power plant accident. Light grey

574   areas show farmland polygons.

575
576 Figure 2 Photo showing a video camera with GPS mounted on a side window of a vehicle (a)
577 and photo, captured from a video recorded by a vehicle-mounted camera, showing a heron in a
578 rice field (b).
579

580
581 Figure 3 Illustration of the selection of farmlands within the viewshed. A survey route (bold
582 line) and farmland polygons (surrounding lines) are shown. The polygons that meet the criteria
583 (see text) are selected as viewsheds (shaded areas).
584

586　Figure 4 Relationship between the distance from cameras to herons and egrets and location

587　errors. Black line and grey area show the maximum likelihood estimates and 95% credible

588　interval for the regression function.

590   Figure 5 Location of herons and egrets observed by roadside survey.

Figure 6 Histograms of the distance from the video camera to all grid cells of farmland

polygons (light grey bars) and grid cells containing herons and egrets (dark grey bars).

Figure 7 Prediction map of number of herons and egrets per square kilometer. Only grids that

contain farmland area are included. The evacuation zone is circled by solid black lines.

Figure 8 Map of the difference between the two predicted values for the numbers of herons and
egrets per square kilometer. The actual value and the value obtained when assuming a
counterfactual condition that the evacuation zone was not established were compared.

Table 1 MAP estimates and summary statistics via importance sampling with and without the
location error.

Parameter	With the location error					Without the location error				
	Map	Mean	S.D.	CI2.5	CI97.5	Map	Mean	S.D.	CI2.5	CI97.5
Intercept	7.15	7.71	1.43	5.41	10.8	7.16	7.78	1.53	5.39	11.3
Distance	−0.0196	−0.0197	0.00180	−0.0234	−0.0162	−0.0197	−0.0199	0.00175	−0.0233	−0.0166
Water	0.545	0.541	0.0696	0.399	0.674	0.546	0.540	0.0701	0.395	0.671
Evac zone	−5.86	−6.43	1.40	−9.50	−4.16	−5.86	−6.50	1.52	−9.96	−4.18
Year(2015)	−0.139	−0.132	0.234	−0.568	0.335	−0.140	−0.135	0.230	−0.580	0.338
Year(2016)	−1.08	−1.07	0.230	−1.52	−0.624	−1.08	−1.07	0.234	−1.50	−0.578

609
610
611

612 Table 2 Predicted values for the number of herons and egrets per square kilometer using the
613 model with and without location error.

Area category	With the location error		Without the location error	
	Mean	S.D.	Mean	S.D.
Outside the evacuation zone	4.62	5.41	4.66	5.45
Inside the evacuation zone	0.0158	0.0173	0.0159	0.0174

614
615