Detection and characterization of novel luchacoviruses, genus Alphacoronavirus, in meso-carnivores in the northeastern United States

Ximena A. Olarte-Castillo[1,2], Laura Plimpton[3], Holly McQueary[2,4], Yining Sun[2,4], Y. Tina Yu[2,4], Sarah Cover[2,4], Amy N. Richardson[2,4], Jennifer K. Grenier[5], Kevin J. Cummings[4], Elizabeth Bunting[4], Maria Diuk-Wasser[3], David Needle[6], Krysten Schuler[7], Michael J. Stanhope[4], Gary Whittaker[1,4], Laura B. Goodman[2,4*]

1. Department of Microbiology & Immunology, College of Veterinary Medicine, Cornell University, Ithaca, NY, 14853, USA
2. James A. Baker Institute for Animal Health, Cornell University College of Veterinary Medicine, Ithaca, NY, 14853, USA
3. Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, NY, USA
4. Department of Public and Ecosystem Health, College of Veterinary Medicine, Cornell University, Ithaca, NY, 14853, USA.
5. Transcriptional Regulation and Expression Facility, Biotechnology Resource Center, Institute of Biotechnology, Cornell University
6. New Hampshire Veterinary Diagnostic Laboratory, College of Life Sciences and Agriculture, University of New Hampshire
7. Cornell Wildlife Health Lab, Animal Health Diagnostic Center, Cornell College of Veterinary Medicine, 240 Farrier Road, Ithaca, NY 14853

*Corresponding author: laura.goodman@cornell.edu

Abstract

Small to mid-sized carnivores, known as meso-carnivores, comprise a group of highly diverse mammals, many of which can easily adapt to anthropogenically disturbed environments. Due to commonly inhabiting urban landscapes, high population densities, and wide home ranges, wild meso-carnivores may get exposed to or spread pathogens to other species. To date, several coronaviruses (family *Coronaviridae*) have been detected and characterized in domesticated and farmed meso-carnivores, but knowledge of the role of meso-carnivores in the epidemiology of coronaviruses (CoVs) remains limited. To increase our understanding of CoVs circulating in wild meso-carnivores, we screened 300 samples, including feces and tissues, from 9 species of free-ranging wild meso-carnivores in the northeastern United States collected between 2016 and 2022. Using a pan-CoV nested PCR approach, we detected CoVs in fecal samples of animals in three species: fisher (*Pekania pennanti*), bobcat (*Lynx rufus*) and red fox (*Vulpes vulpes*). Next generation sequencing revealed that all the detected viruses belonged to the *Luchacovirus* subgenus within the *Alphacoronavirus* genus, previously reported only in rodents and lagomorphs (i.e., rabbits and pikas). Genetic comparison of the 3'-end of the genome (~12,000 bp) revealed that although the viruses detected group with, and have a genetic organization similar to other luchacoviruses, they are genetically distinct to luchacoviruses from rodents and lagomorphs. Genetic characterization of the spike (S) protein revealed that these luchacoviruses do not have an identifiable S1/S2 cleavage site but do have highly variable structural loops within S1 that comprise protease cleavage motifs or very similar cleavage motifs to those identified in certain pathogenic CoVs. This study highlights the importance of characterizing the S protein from CoVs in various wild animal species to target specific regions for further experimental analysis and epidemiologic monitoring.

Importance

Spillback of SARS-CoV-2 into free-ranging animal species has been well documented, but the repertoire of other coronaviruses that could serve as sources for recombination in many hosts is unknown. The cryptic circulation of possible animal-associated variant SARS-CoV-2 lineages in urban sewage surveillance, and canine alphacoronavirus infections in humans, highlight a critical need to survey animal populations in the US. In general, no large-scale pathogen discovery work has been conducted in North America; studies on other continents have largely focused on bats. While bats are a well-known reservoir for coronavirus emergence, the potential risks for other mammals are very poorly understood. Assessing the potential of US wildlife species as pathogen reservoirs is vastly under-explored and is essential to prepare for future emerging threats. Our focus here was on α and β CoV discovery from the Northeast USA as a potential "hotspot" area with high density of humans and urban wildlife. This study identifies novel alphacoronaviruses circulating in this area and explores their potential zoonotic risk based on Spike gene sequence.

Introduction

The *Coronaviridae* family encompasses diverse viruses with single-stranded, positive sense RNA genomes. Coronaviruses (CoVs) are currently classified into four genera (*Alpha-*, *Beta-*, *Gamma*, and *Deltacoronavirus*, (1)), that group viruses known to infect birds (mostly gamma- and deltacoronaviruses) and mammals (mostly alpha- and betacoronaviruses), including domesticated and wild animal species, and humans (2). The potential host range of CoVs is coordinated by the spike (S) protein, which is involved in receptor binding via its S1 domain (3). The S protein also mediates membrane fusion with target cells through its S2 domain (4). In general, cleavage of S can occur during virion biosynthesis at a cleavage site between the S1 and S2 domains (S1/S2

cleavage site) which activates the S protein exposing the receptor binding region within the S1

domain, promoting receptor binding (5). After receptor binding, S is also cleaved in the S2 domain

close to the fusion peptide (S2' cleavage site), promoting membrane fusion (6). Coronavirus entry

is often regulated through the S cleavage sites. In the *Betacoronavirus* genera, MERS-CoV (from

the Merbecovirus clade) can be cleaved by furin at both cleavage sites (S1/S2 and S2', (5)), while

SARS-CoV-2 is the only known member of the Sarbecovirus clade that has a furin cleavage site

in S1/S2 (7) , which is a key determinant of its emergence (8). In the *Alphacoronavirus* genera,

CoVs like feline and canine CoV type 2 (FCoV-2 and CCoV-2, respectively), transmissible

gastroenteritis virus and human CoV 229E do not have an identifiable S1/S2 cleavage site (9),

while swine acute diarrhea syndrome CoV (SADS-CoV), an emergent and highly pathogenic CoV

that possibly originated from bat CoV HKU2 does have an S1/S2 cleavage site (10). The S protein

of SADS-CoV also has an additional cleavage site in the S1 domain whose cleavage together with

that on the S1/S2 site is essential for cell-cell fusion (11). Overall, multiple cleavage sites may

contribute to membrane fusion and virus entry, processes that are crucial determinants for an

efficient infection and the disease outcome.

The advancement in sequencing technologies has contributed to the surge of viral

discovery which together with comparative genetic analyses can help increase our understanding

of how CoV exploit their genetic variation for host adaptation and pathogenesis (3). As the S

protein of several CoVs has been crystalized, it is possible to identify and map regions within this

protein that interact with the host for receptor binding (12) and cleavage-mediated activation or

fusion (13). Comparative genetic analyses of the S protein between CoVs from different hosts can

help guide experimental assays to evaluate the importance of these regions and assess the impact

of certain mutations on CoV host range and/or pathogenicity. For example, furin, tryspin-like

proteases and cathepsins, among other proteases, can cleave the S protein after recognizing certain

amino acid sequence motifs (14). Mutations and/or insertions/deletions in these motifs can impact

the virus-host interactions. For example, the S1/S2 cleavage site of FCoV type 1 (FCoV-1) and

SARS-CoV-2 is located in exposed loops easily accessible to proteases (15-17). Mutations in these

loops that disrupt the furin cleavage motif results in highly pathogenic variants for FCoV-1 (18)

and variants with decreased pathogenicity for SARS-CoV-2 (19). Likewise, mutations in the S2'

cleavage site can inhibit viral fusion which in MERS-CoV results in reduced viral infectivity (20).

Therefore, identifying and mapping in the tertiary structure of S potential cleavage sites in CoVs

detected in different species can shed light on the processes CoVs may use to acquire cleavage

sites (21) and its relation to host range, pathogenicity and evolution of CoVs (22).

Luchacovirus is a subgenus of *Alphacoronavirus* (also known as rodent or murine

Alphacoronavirus), that includes viruses infecting rodents (order Rodentia) and rabbits (order

Lagomorpha). Although currently classified as alphacoronaviruses, the S gene of luchacoviruses

is more similar to and groups with betacoronaviruses and not alphacoronaviruses, likely as a result

of an early recombination event (23). To date, luchacoviruses have only been detected in 12 species

of rodents (including rats, mice, and voles) and 2 species of lagomorphs (including rabbits and

plateau pika, Table 1). Even though luchacoviruses are highly diverse and have been reported

from various rabbit and rodent species in different countries in Asia, Europe and the U.S. (24),

they form a monophyletic group in the phylogenies involving the S and other genes and share

similar genetic features in their genomes, suggesting a long association with rodents and rabbits

(23). In general, the S gene of these viruses is highly variable and the sequence of one of these

viruses (JC34, accession number KX964649) obtained from a Chevrier's field mouse (*Apodemus*

chevrieri) in China in 2011 (25) has a furin cleavage motif (S-R-R-A-R | A, where S is Serine, R

is Arginine, A is Alanine and | is where the cleavage occurs) in its S1 domain. Interestingly this

motif has a very similar amino acid sequence to the S1/S2 cleavage site of SARS-CoV-2 (P/H-R-

R-A-R | S, where P is proline, H is Histidine (22)), which is an atypical furin recognition motif

(the canonical furin cleavage motif is X-R-R-X-K/R-R | S, where K is lysine and X is any residue).

Although experimentally the luchacovirus JC34 cleavage motif is not cleaved by human furin (26),

continued surveillance of these viruses focusing on targeted regions is essential to understand if

there is variation in regions with motifs of interest and their relevance in the origin and emergence

of new CoVs.

The meso-carnivores are a group of small to mid-size carnivores of economic and ecologic

relevance that include domesticated and farm animals (dogs, cats, and American mink, *Canis lupus*

familiaris, Felis catus, Mustela vison, respectively) as well as various wild species. In the

northeastern United States (U.S.), there are 14 species of meso-carnivores (27), including

ubiquitous species like raccoons (*Procyon lotor*), red foxes (*Vulpes vulpes*), and striped skunks

(*Mephitis mephitis*). Due to their diverse behavior and ecology, meso-carnivores can thrive in

various habitats including urban and rural areas (28). In urban areas they may get exposed to

pathogens from or spread infectious diseases to domesticated animals. For example, free-ranging

red foxes and coyotes *(Canis latrans)* in Pennsylvania are exposed to canine distemper virus

(CDV) and canine parvovirus (CPV), two viruses that can cause deadly infections in a wide range

of carnivores (29). In rural areas they may act as reservoirs for pathogens that may spill over into

other species of wild carnivores. Indeed, skunks and raccoons in the eastern U.S. can maintain the

sylvatic cycle of rabies independently of domesticated animals (30). Despite their potential

importance in the spread of infectious diseases, our knowledge of the viral diversity hosted by

free-ranging wild meso-carnivore species in the U.S. remains limited (31).

Table 1. Species of rodents and lagomorphs from which uchacoviruses have been reported.

Order	Species	Common name
Rodentia	*Rattus novergicus*	Brown rat
	Rattus losea	Lesser field rat
	Rattus nitidus	Himalayan field rat
	Berylmys bowersi	Bower's white-toothed rat
	Mus musculus	House mouse
	Apodemus chevrieri	Chevrier's field mouse
	Apodemus agrarius	Striped field mouse
	Microtus arvalis	Common vole
	Eothenomys fidelis	Oriental vole
	Eothenomys melanogaster	Pere David's vole
	Myodes rufocanus	Grey red-backed vole
	Microtus agrestis	Field or short-tailed vole
Lagomorpha	*Oryctolagus cuniculus*	European rabbit
	Ochotona curzoniae	Plateau pika

Alpha- and betacoronaviruses can infect diverse species of meso-carnivores causing a

range of diseases. Alphacoronaviruses like mink CoV (32) and ferret enteric CoV (33) cause

enteric diseases in mink and ferret, respectively, while ferret systemic CoV (33) and FCoV (34)

can cause highly fatal diseases in ferrets and domesticated cats, respectively. Betacoronaviruses

like canine respiratory CoV are widely spread throughout the U.S. causing respiratory signs in

domesticated dogs (35). Like their domesticated counterpart, wild meso-carnivores also host

different CoVs, including the divergent novel alphacoronaviruses identified in free-ranging Asian

leopard cats (*Prionailurus bengalensis*) and Chinese ferret badgers (*Melogale moschata*) in

southern China (36). Meso-carnivores also play a role in the epidemiology of CoVs relevant to

humans including SARS-CoV-2 which can infect minks and ferrets (37), species that can also

infect humans (38). Likewise, recent studies have found CCoV-2, a common *Alphacoronavirus*

of domesticated dogs and wild canids, in humans (39). This zoonotic event was related to extensive

recombination with other alphacoronaviruses and the possible loss of function of the N-terminal

region of the S1 domain (40). Meso-carnivores may also be susceptible to CoVs from domesticated

and farmed meso-carnivores, due to their close genetic relatedness. For example, the sequence of

the receptor for CCoV-2 and FCoV-2 (Aminopeptidase N) is highly similar (>92.7%) between

members of the Felidae and Canidae families (41). Overall, both urban and rural wild meso-

carnivore carry a high diversity of viruses (42) that need to be further explored to understand the

complex epidemiology of CoVs in wild populations, especially those highly adapted to

anthropogenically disturbed areas.

The aim of this study was to perform a longitudinal surveillance of CoV in free-ranging

wild meso-carnivore in the northwestern region of the U.S. Using next generation sequencing,

comparative genetic and phylogenetic analyses, we provide the first molecular characterization,

focusing on the S gene, of luchacoviruses detected in three species of meso-carnivore in the U.S.

and show that the luchacoviruses detected in the U.S. to date, are highly diverse when compared

to those reported elsewhere.

Methods

<u>Sample collection</u>

Between 2016 and 2022, tissues and feces from 9 species of meso-carnivore were opportunistically

collected from carcasses turned in for routine necropsy to either the New York State or New

Hampshire veterinary diagnostic laboratories. We collected samples from bobcat, fisher, American

mink, coyote, red fox, racoon, striped skunk, river otter (*Lontra canadensis*), and gray fox

(*Urocyon cinereoargenteus*). In 2021 we also collected samples from 4 species of rodents

including chipmunk (subfamily Xerinae), Eastern cottontail (*Sylvilagus floridanus*), Eastern gray

squirrel (*Sciurus carolinensis*), and North American porcupine (*Erethizon dorsatum*).

Between August and October 2021, we collected and screened 30 rectal swabs from 17

raccoons and 7 striped skunks living in an urban green-space in New York City, NY. Animals

were captured in Tomahawk Live Traps baited with cat food. Traps were baited and set at dusk

and checked at dawn. Captured animals were anesthetized using ketamine-dexmedetomidine

administered by a veterinarian in the field. Each sedated animal was given a routine health exam

and rectal swabs were collected from each individual using sterile polyester tipped swabs by gently

inserting the swab into the anus and rotating the rod on its axis for 10-15 seconds. Swabs were

then individually stored in 1.5mL screw top microcentrifuge tubes with 500uL of RNAlater

(Thermo Fisher Scientific).. Samples were transported on ice, incubated at 4°C overnight and

transferred to -80°C for long term storage prior to RNA extraction. After sample collection,

animals were returned to their cage, monitored during recovery, and released at the point of

capture. Capture and sampling of NYC meso-carnivore was performed under New York State

Department of Environmental Conservation (NYSDEC) permit #2704 and were approved under

Columbia University's IACUC protocols #AC-AABA1469 and #AC-AABA1472

Pancoronavirus screening

Total RNA was obtained from both tissues and feces using the MagMax *mir*Vana Total RNA

isolation kit (Thermo Fisher Scientific) using the KingFisher Apex Purification System (Thermo

Fisher Scientific). Prior to RNA extraction between 20 to 40 mg of sample was disrupted using

the Bead Ruptor Elite (OMNI International) and 800 ul of lysis buffer. For the disruption we used

the MagMAX Microbiome Bead Tubes for feces and 500mg of ~2.5mm Zirconia/Silica beads

(Biospec Products) for tissues. The obtained RNA was reverse transcribed using the SuperScript

VILO Master Mix (Thermo Fisher Scientific). CoV RNA was detected using a pancoronavirus

nested PCR with primers targeting the conserved RdRp gene of CoVs as previously published

(Watanabe et al., 2010) and using the Platinum II Taq Hot-Start DNA Polymerase (Thermo Fisher

Scientific). After PCR, samples were run in a 1% agarose gel stained with GelRed (Biotium) and

visualized in a Gel Doc XR System (Biorad). Positive samples were Sanger sequenced at the

Biotechnology Resource Center (BRC) at Cornell University.

CoV-positive samples were further sequenced by RNA-seq with a sequencing depth of 100

million reads at the Transcriptional Regulation and Expression Facility at Cornell

(RRID:SCR_022532). RNA-seq libraries were prepared using the NEBNext Ultra II Directional

RNA Library Prep Kit for Illumina (New England Biolabs, NEB) using 200-600ng total RNA as

input. Subtraction of both bacterial and mammalian rRNA was done using the NEBNext rRNA

Depletion Kits (NEB), and sequenced on a Novaseq 6000 instrument (Novogene). RNA Reads

generated were uploaded to CZ ID (www.czid.org) to find matches to nucleotide and protein

sequences available in NCBI. To close gaps from the sequences obtained after RNAseq, we

designed primers based on the known sequences and performed PCR on the previously obtained

cDNA using Platinum II Taq Hot-Start DNA Polymerase (Thermo Fisher Scientific). Obtained

products were sequenced in a MinION Mk1C (Oxford Nanopore Technologies, ONT) using a

Flow Cell R10.4 (ONT) and the Native Barcoding Kit 24 V14 (ONT).

<u>Genetic and phylogenetic analyses</u>

In this study we obtained the partial sequences of the RdRp gene (401nt) for the three

positive samples (W291/Bobcat/2021, W145/fisher/2022 and W317/red fox/2022), the partial

sequence of the S gene (2,561nt), that includes the complete S1 domain and a partial sequence of

the S2 domain, of one of our positive samples (W291/Bobcat/2021) and the complete 3'-end of

the genome (12,214nt) of one of our positive samples (W317/red fox/2022). The obtained

sequences were deposited in GenBank (accession numbers OQ756331-35). The three partial

sequences of the RdRp (401nt) and the two complete sequences of the S1 domain of the S gene

(1674nt) obtained in this study were each aligned with homologous sequences from 60 and 36

other CoVs from the four families, respectively. These alignments were obtained using the Clustal

Omega algorithm (43) and each was used to compute the best-fitting nucleotide substitution model

and to construct a maximum likelihood (ML) tree using MEGA 11 (44).

We detected open reading frames (ORFs) in the complete sequence of the 3'-end of the

genome (12,214nt) obtained from one of our samples and 15 other luchacoviruses sequences

available using Geneious Prime 2023.0 (Dotmatics). We aligned these 16 sequences using the

MUSCLE algorithm (45) after which we deleted the regions where gaps were identified and then

generated a similarity plot using SimPlot 5.1 (46). We also used this alignment to find possible

recombination break points using the Recombination Detection Program (RDP) 5 (47) using the

RDP method (48). Additionally, we aligned the partial S gene sequence obtained from one of our

samples (2,561nt) with the 16 luchacovirus sequences mentioned previously using the MUSCLE

algorithm (45). This alignment was used to generate a similarity plot using SimPlot 5.1 (46). These

17 partial S sequences were translated into their respective amino acid sequences and aligned with

the homologous amino acid sequences of two alphacoronaviruses, SADS-CoV (MT294722), bat

CoVs HKU2 (NC_009988), and three betacoronaviruses including SARS-CoV-2 (QQN72495), a

rodent CoV (MT820632) and a rabbit CoV (JN874560) using the Clustal Omega algorithm (43).

This alignment was used to compare regions in S in which known cleavage sites occur including

the S1/S2 and S2' regions of SARS-CoV-2 (49), and regions in the S1 domain in which

Luchacovirus JC34 (22) and SADS-CoV (11) have predicted and characterized furin cleavage

sites, respectively. We detected potential cathepsin cleavage sites by looking for specific amino

acid motifs previously identified for human cathepsins L, V, K, S, F, B (14). We also used ProP

(50) to predict furin cleavage sites (ProP values >0.5). We mapped the regions of interest on the

tertiary structure of the S protein of SADS-CoV which is the most closely related

Alphacoronavirus to the *Luchacovirus* clade that has an available crystalized tertiary structure

(51). The colors of the protein and the mapping of the regions of interest were done using Chimera

X (52).

Results

From 177 individual meso-carnivores we collected 300 samples: 145 fecal samples and 155 tissues

samples (Table 2) including liver (43%), lung (44%), brain (2.8%), kidney (6%), small intestine

(2.8%), and heart (1.4%) from 9 species of meso-carnivore (Table 3). Using a nested PCR

pancoronavirus screening method we found 3 samples positive for CoV RNA, all from feces

(Table 2) from three different meso-carnivore species (Table 3) including samples from a bobcat

collected in October 2021 (W291/bobcat/2021) and a fisher (W145/fisher/2022) and red fox

(W317/red fox/2022) collected in January and February 2022, respectively (Table 4). The bobcat

and red fox samples were collected in New Hampshire and the fisher was collected in New York.

Screening of lung from the bobcat (W291/bobcat/2021) and red fox (W317/red fox/2022), as well

as lung liver, kidney and heart from the fisher (W145/fisher/2022) revealed that none of these

tissues were positive for CoV RNA. Of the 30 fecal samples from raccoons and skunks sampled

in 2021 in New York City, none was positive for any CoV. For rodents, we collected 36 samples

in 2021, including feces (50%) and lung (50%). We did not find CoV in any of these samples

(Supplementary table 1).

Table 2. Results of the CoV screening of meso-carnivore according to sample type.

Sample type	Total	Positive	Negative
Tissues	155	0	151
Feces	145	3	142
Total	300	3	293

273 **Table 3**. Results of the CoV screening of meso-carnivore according to species.

Species	Total	Positive	Negative
Fisher	124	1	123
Bobcat	40	1	39
Raccoon	39	0	39
Coyote	24	0	24
Otter	21	0	21
Red fox	21	1	18
Skunk	18	0	18
Gray fox	8	0	5
Mink	5	0	4
Total	300	3	291

276 **Table 4**. Results of the CoV screening of meso-carnivore according to year of collection.

Year of collection	Total	Positive	Negative
2016	1	0	1
2017	2	0	2
2018	47	0	47
2019	71	0	71
2020	18	0	18
2021	123	1	122
2022	38	2	36
Total	300	3	297

278 **Supplementary Table 1**. Rabbit and rodent screening results according to sample type and
279 species.

Species	Collection year	Sample type	Number of samples	Number of positive samples
Chipmunk	2021	Feces	6	0
Chipmunk	2021	Lung	6	0
Eastern cottontail	2021	Feces	4	0
Eastern cottontail	2021	Lung	4	0
Gray squirrel	2021	feces	6	0
Gray squirrel	2021	Lung	6	0
Porcupine	2021	Feces	2	0
Porcupine	2021	Lung	2	0
			36	0

280
281
For the three positive samples obtained from the meso-carnivores, we initially sequenced a partial

region of the RdRp gene (401bp). The percentage sequence similarity of this region between these

three sequences was 81.4%. A phylogeny including these three sequences plus 60 others from

CoVs in the four genera revealed that all sequences obtained in this study are within the

Luchacovirus clade (Figure 1). Both W145/fisher/2022 and W317/red fox/2022 clustered in a

group together with viruses from rabbit and vole reported in Europe, while W291/Bobcat/2021 did

not group with any specific group of luchaconviruses. The Luchacovirus cluster grouped together

with SADS-CoV and bat CoV HKU2 (Figure 1, emphasized with blue arrows).

290
Figure 1. Maximum likelihood phylogenetic tree of a partial region of the RdRp gene of 63
coronaviruses (CoVs) from the four genera, including the three obtained in this study
(highlighted in fuchsia). The Luchacovirus clade is within the *Alphacoronavirus* group, each
shown within labelled brackets. The CoVs most closely related to the Luchacovirus group
(SADS-CoV and bat CoV HKU2) are emphasized with blue arrows. For all viruses the virus
name, host species, year and country of collection, and Genebank accession numbers are shown
at the tip of the tree. Numbers at the branches indicate bootstrap percentage values from 1000
replicates. Branches with support <50 were collapsed. Nucleotide substitution model used:
HKY+I+G.

300
301

Additional sequences were obtained for W317/red fox/2022 and W291/bobcat/2021 but not for

W145/fisher/2022 due to low quality and quantity of RNA in this sample. For W317/red fox/2022

we obtained a partial genome sequence (12,214nt) that includes the complete S gene and for

W291/Bobcat/2021we obtained a partial sequence of S gene (2,561nt) containing the complete

sequence of the S1 domain and a partial sequence of the S2 domain. A phylogeny constructed

using the nucleotide sequences of the S1 domain of the two variants obtained in this study and 36

other CoVs from the four families revealed that our sequences also grouped within the

Luchacovirus cluster and that this group was not closely related to the *Alphacoronavirus* group

(Figure 2). In general, there were three main groups (bootstrap support of 100%) within the

Luchacovirus cluster: Group 1 includes viruses from voles and rat from China, Group 2 includes

viruses from rat from China and the UK and Group 3 includes viruses from mice and rat from

China (Figure 2). There were 4 luchacoviruses, including the two reported in this study, that did

not cluster within these three groups. W317/red fox/2022 was placed in a group with one of the

other divergent luchacoviruses obtained from a pika (P83/plateau pika/China), outside Group 3

while W291/Bobcat/2021clustered outside groups 1 and 2. The fourth divergent luchacovirus,

UMN2020/mouse/2018/USA, collected from a mouse in the U.S., did not group with any other

luchacovirus (Figure 2). Each of the three defined groups have an overall higher percentage

sequence similarity in the S1 domain (>92%) than when compared to the other groups (group 1 vs

2 50%, group 1 vs 3 56.2% and group 2 vs 3 51.8%) or to the 4 luchacoviruses that were not within

these groups, including the two viruses obtained in this report, W317/red fox/2022 and

W291/Bobcat/2021 (<64%), as well as P83/plateau pika/China (<69%) and the most divergent

one, UMN2020/mouse/2018/USA (<52%). Like in the phylogeny of the partial RdRp gene, the

Luchacovirus cluster grouped together with SADS-CoV and bat CoV HKU2 (Figure 2,

emphasized with blue arrows).

Figure 2. Maximum likelihood tree of the complete nucleotide sequence of the S1 domain of the
S gene of 38 coronaviruses from the four genera including the two obtained in this study
(highlighted in fuchsia). In this phylogeny the Luchacovirus clade is not within the
Alphacoronavirus group, each shown within labelled brackets. Within the Luchacovirus group
the three well-supported groups (bootstrap of 100%) are shown in different colors (Group 1 in
blue, Group 2 in orange, Group 3 in green). Four luchacoviruses, included the ones identified in
this study do not cluster within these three groups. For all viruses the virus name, host species,
year, and place of collection, and Genebank accession number are shown at the tip of the tree.
Numbers at the branches indicate bootstrap percentage values from 1000 replicates. Branches
with bootstrap support of <50 were collapsed. Nucleotide substitution model used: GTR+I+G.
337

The amino acid sequences of the partial S protein (853 amino acids), containing the complete S1

domain and a partial region of the S2 domain of W317/red fox/2022 and W291/bobcat/2021 were

52.1% similar. We compared the amino acid sequences of the four regions in S known in other

CoVs to have protease cleavage sites between the two viruses detected in this study, 15 other

luchacoviruses, 2 alphacoronaviruses and 3 betacoronaviruses. The genetic comparison of these

four regions revealed that revealed that two of these regions are located in the C-terminal domain

(CTD) of S1 (Figure 3). Region 1 includes the furin cleavage site of SADS-CoV (R-Y-V-R | I,

Figure 3A), whose sequence differs in only one residue (residue 458 in Figure 3A) with the

respective sequence from luchacoviruses from groups 1 and 2, the divergent variants

UMN2020/mouse/2018/USA and P83/plateau pika/China (**S**-Y-V- R | I), and bat CoV HKU2 (**K**-

Y-V-R | I). For these variants, a single nucleotide change is sufficient to revert the residue change

detected between these viruses (Figure 3B). Compared to group 3 luchacoviruses and the two

viruses obtained in this study, it differs in two residues (458 and 457, Figure 3A). In the tertiary

structure of SADS-CoV region 1 is located on a beta sheet within the CTD (Figure 3C). Region 2

is highly variable and includes a weakly predicted furin cleavage motif detected in Group 3

luchacoviruses (R-R-A-R | A, ProP score 0.28) and motif F-R | S in W317/red fox/2022/USA

consistent with a cathepsin L cleavage site (Figure 3A). In this region divergent luchacoviruses

UMN2020/mouse/2018/USA and W291/Bobcat/2021 have highly different sequences (Figure

3A). Region 3 includes the S1/S2 cleavage site, which is observed in betacoronaviruses like SARS-

CoV-2 and two rodent betacoronaviruses (ProP scores >0.5), but not in luchacoviruses (Figure

3A). Region 4 includes the S2'cleavage site in which furin cleavage motifs were not predicted for

any Luchacovirus. Regions 2, 3 and 4 are located in exposed loops in the tertiary structure of

SADS-CoV (Figure 3C).

Figure 3. A. Graphical representation of the S protein showing the S1 and S2 domains (top) and multiple sequence alignments (bottom) of four regions of interest within S (shown in violet circles) in which cleavage motifs are found (underlined in blue are the predicted cleavage sites and in violet are experimentally proven furin cleavage sites). The S1 domain is divided into the N-terminal domain (NTD, in green) and the C-terminal domain (CTD, in yellow). In the amino acid alignments polar residues are in orange, acidic residues are in green, basic residues are in blue and

nonpolar residues are in grey. In the left side of the alignment the virus name/host/year of
collection/place of collection is shown. Above the alignments are the amino acid location of each
region based on the S protein sequence of rodent Luchacovirus JC34 (KX964649). From left to
right there are four regions of interest: (1) Alphacoronavirus SADS-CoV has a furin cleavage site
94 amino acids upstream of its S1/S2 cleavage site that differs in only one residue when compared
to four luchacoviruses and bat CoV HKU2 (shown with yellow triangles), (2) luchacoviruses have
a region 37 amino acids upstream of the end of domain S1 that is highly variable and in which
there are two different cleavage motifs, (3) luchacoviruses do not have an S1/S2 cleavage region
in comparison to SADS-CoV and selected betacoronaviruses (SARS-CoV2 and two rodent
coronaviruses) and (4) only SARS-CoV2 has an S2' cleavage motif. B. Amino acid sequences and
the respective codons (on top) of region 1, in which SADS-CoV has a cleavage site and bat CoV
HKU2 and the luchacoviruses shown differ by a single residue (458) to that of SADS-CoV
(R458S, R458K, respectively), which abrogates the cleavage site. Inspection of the codon
sequences shows that only a single nucleotide change (highlighted in yellow) is necessary to revert
this residue change. C. Tertiary structure of one of the monomers of the S protein of SADS-CoV
showing the four regions in which we found cleavage sites in luchacoviruses or related alpha and
betacoronaviruses. For this figure we used the same colors as in A.

The 3'-end of the genome of luchacoviruses has a general organization that includes four structural

proteins S, envelope (E), matrix (M), and nucleocapsid (N) and three to five non-structural ORFs,

one or two between ORF1b and the S protein (ORF2 and 2b), ORF3 between S and E, ORF6 M

and N, ORF8 within N and ORF9 after N (Figure 4). In this region the three luchacovirus groups

differed in the number of ORF2, group 1 has ORF2 and 2b, group 2 only has ORF2 and group 3

does not have any ORF2 (Figure 4). The obtained partial genome of W317/red fox/2022 had a

main similar organization to group 2 luchacoviruses with only one ORF2 (Figure 4), but differed

in that it does not have an ORF8 or its internal transcription signal and ORF9 was shorter than

those from the other luchacoviruses, due to a 69-nucleotide deletion (Figure 4). W317/red fox/2022

shared low overall genetic similarity at the 3'-end of the genome when compared to other

luchacoviruses (<86%, Figure 5A). In comparison, the luchacoviruses within groups 1, 2, 3 shared

an overall high nucleotide similarity (pairwise percentage similarity >94.2%). Divergent

luchacoviruses UMN2020/mouse/2018/USA and P83/plateau pika/China had the same genomic

organization as luchacoviruses from groups 3 and 2, respectively. However, genetic comparison

revealed an overall low similarity throughout the 3'-end of the genome between these

luchacoviruses and those in groups 3 and 2 (UMN2020/mouse/2018/USA vs group 3 <82% and

P83/plateau pika/China vs group 2 <88%, Supplementary Figure 1). We did not find any

recombination breakpoints in this region of the genome between the 16 luchacoviruses included

in the analysis (see Methods). The partial S gene of W291/Bobcat/2021was overall genetically

different from other luchacoviruses, including W317/red fox/2022 (<80%, Figure 5B).

409
Figure 4. Graphical representation of the 3'-end of the genome of Luchacovirus, that includes
the structural (S, E, M, N) and non-structural (ORFs 2, 2b, 6, 8, 9) genes. Sixteen sequences,
including the one obtained in this study (on top in fuchsia) are shown. Above each genome
representation is the name of each gene, below is the size of the corresponding proteins (number
of amino acids). If some proteins varied in size between the viruses, all size values are shown
from shortest to longest. On the left side of the figure the name of the virus/host species/year of
collection/country of collection of all sequences included are shown. There are three genomic
organizations that match the three groups (groups 1, 2 and 3 in blue, orange, and green,
respectively) identified in Figure 2. The three genomic organizations differ in the existence of
ORFs between ORF1b and S (ORF2 and 2b). The sequence obtained in this study, W317/red
fox/2022 (top), has a similar organization to group 2 luchacoviruses as it only has one ORF2 but
it differs from other luchacoviruses because it lacks ORF8 and ORF9 is shorter. Divergent
luchacoviruses P83/plateau pika/China and UMN2020/mouse/2018/USA have the same genomic
organization of viruses from groups 2 and 3, respectively, but do not belong to these groups and
are overall genetically different (Supplementary Figure 1).

426
Figure 5. Similarity plot of the nucleotide sequence of (A) the 3'-end of the genome of
W317/red fox/2022 (top, used as reference) and (B) a partial region of the S gene of W291/bob
cat/2021 (top, used as reference) compared to 15 other luchacoviruses including those in group 1
(in blue), 2 (in orange), and 3 (in green), and divergent luchacoviruses
UMN2020/mouse/2018/USA (in yellow) and P83/plateau pika/China (in violet). The location of
each gene or gene domain is shown above each the graph. The graphs were constructed using a
window of 200bp, a step of 20nt and the kimura 2 parameter distance model.

Supplementary Figure 1. Similarity plots comparing the nucleotide sequence of the 3'-end of
the genome of UMN2020/mouse/2018/USA (in violet) with group 3 luchacoviruses (in green,
top) and P83/plateau pika/China (red) with group 2 luchacoviruses (in orange, bottom). The
graphs were constructed using a window of 200bp, a step 20nt and the kimura 2 parameter
distance model. and Kimura-2-parameter distance model with a sliding window of 200
nucleotides and a step size of 20 bases.
440

441

442
443
444
445 **Discussion**
446
Coronaviruses are important pathogens that have a wide host range and can result in subclinical

infection or cause a variety of diseases in both domesticated and wild mammals. Within the

Alphacoronavirus genus, the *Luchacovirus* sub-genus includes CoVs reported to date only in

rodents and rabbits. In this study we report for the first time luchacoviruses in three species of

meso-carnivore including bobcat (W291/bobcat/2021), fisher (W145/fisher/2022), and red fox

(W317/red fox/2022, Figure 1). The phylogeny of the S1 domain of the S protein and the general

genomic organization of the rodent and rabbit luchacoviruses reported to date showed that there

are three main groups (Groups 1- 3, Figures 2 and 4), two of which were already observed using

fewer sequences (53). Phylogenetic and genetic comparison of the S1 domain of two of the viruses

obtained in this study (W291/bobcat/2021 and W317/red fox/2022) revealed that this region is

genetically different from the luchacoviruses reported to date in rodents and rabbits (<75%, Figure

5) and that these viruses did not group within the three groups that comprise most of the

luchacoviruses reported to date (Figure 2). Although in general CoVs have highly variable S1

regions, further genetic comparison of the 3'-end of the genome revealed that luchacoviruses

within the three established groups are highly similar (>94.2% similarity) but W317/red fox/2022

is overall divergent (<86% similarity, Figure 5). Likewise, the genetic organization of the 3'-end

of the genome of W317/red fox/2022 had certain differences with the other luchacoviruses

including a shorter ORF9 and the absence of ORF8 (Figure 4). These results support the idea that,

with the sequences available to date, the two viruses reported in this study (W291/bob cat/2021

and W317/red fox/2022) are genetically different from other luchacoviruses reported in rodents

and rabbits. As in the phylogeny of the partial RdRp gene W317/red fox/2022 and

W145/fisher/2022 grouped with viruses of vole and rabbit from which only partial sequences exists

(Figure 1), additional sequences of these viruses or from these hosts will help clarify if meso-

carnivores host luchacoviruses divergent from those hosted by rodents and rabbits or if there is

cross-species transmission.

Some CoVs have wide host ranges, possibly due to the use of protein receptors like

angiotensin converting enzyme 2 (ACE-2), dipeptidyl peptidase 4 (DPP4) or aminopeptidase N

(APN), that are highly conserved between diverse species (41). As the S1 domain of the S protein

of CoVs interacts with the receptor, signatures of viral adaptation to the receptor of different

species have been identified in this region which has help increase our understanding of the

molecular mechanisms of the adaptation and emergence of CoVs in various species (12). Much

less is known about the relation between the cleavage of S and the emergence of pathogenic CoV

variants, however the presence of a unique cleavage site in the S1/S2 of SARS-CoV2, that to date

has not been found in closely related viruses (54), has highlighted the importance of studying the

mechanisms CoVs may use to acquire functional cleavage (22). For example, genetic comparison

of the S1/S2 region between SARS-CoV-2 and closely related CoVs from European bats revealed

that a single nucleotide substitution is sufficient to produce a functional furin cleavage site

identical to that of SARS-CoV-2 in two bat CoV variants, an essential molecular determinant of

their zoonotic potential (21). The Luchacovirus clade was closely related to SADS-CoV and bat

CoV HKU2 (Figure 1 and 2), which are also characterized for having a recombinant S gene that

is more similar to betacoronaviruses than alphacoronaviruses (55). To date the receptor of

luchacoviruses, SADS-CoV or bat CoV HKU2 remains unknown (56); in this study we directed

our genetic analysis of the S protein on finding regions with putative cleavage sites and the genetic

comparison of these regions with other CoVs like SADS-CoV (11) and SARS-CoV-2 in which the

cleavage of S has been experimentally studied.

SADS-CoV is a highly pathogenic CoV that infects pigs and has a unique cleavage site (R-

Y-V-R | I) in the S1 domain (region 1 in Figure 3) that is essential for cell fusion and may have an

impact in the pathogenicity of this virus and its observed broad cell range (57). Comparative

genetic analyses have shown that bat CoV HKU2 is closely related to SADS-CoV, suggesting that

bat CoVs may also be involved in the emergence of pathogenic CoVs of veterinary importance

(51). It has been previously shown that a substitution of a single reside in this cleavage site (residue

458 in Figure 3) is enough to abrogate the cleavage of S (11). In this study we show that several

luchacoviruses,   including   those   in   groups   1   and   2   and   divergent   variants

UMN2020/mouse/2018/USA and P83/plateau pika/China, have a cleavage motif in this region that

differ only in this essential residue to that of SADS-CoV (R458S, Figure 3A) and that a single

nucleotide substitution is sufficient to revert this residue (Figure 3B). Likewise, bat CoV HKU2

also differs only in this residue when compared to SADS-CoV (R458K, Figures 3A and B).

However, we show that unlike SADS-CoV and bat CoV HKU2, luchacoviruses do not have an

S1/S2 cleavage site (Figure 3A). As cleavage in both S1 and S1/S2 sites is required for cleavage

and cell fusion of SADS-CoV (11), more than a single non-synonymous change would be

necessary for luchacoviruses to acquire a cleavage mechanism like the one of pathogenic SADS-

CoV. In contrast to luchacoviruses, most rodent betacoronaviruses have a predicted S1/S2

cleavage site (78% of available S sequences, (22)), more than 13 times of what was predicted for

bat betacoronaviruses (6% of available sequences, (22)). *In vitro* experiments have shown that

SADS-CoV can successfully infect and replicate in several rodent cell lines (56); therefore,

surveillance of rodents, meso-carnivore, pigs, and bats targeting luchacoviruses, rodent

betacoronaviruses and SADS-CoV and characterizing their S genes is essential to detect if there is

any recombination or variation in these key regions and its relationship with pathogenicity and

host range.

A previous study found that Luchacovirus JC34 had a predicted furin cleavage site (R-R-

A-R | A) in the S1 domain similar to that of the S1/S2 cleavage site SARS-CoV2 (22). In this

study we identified two additional luchacoviruses (all from group 3, Figure 2) that also have this

cleavage site (Figure 3). Genetic comparison of this region among luchacoviruses revealed that it

is a highly variable loop (region 2 in Figure 3) in which we also identified motifs for cathepsin L

or V cleavage (F-R | S) in divergent luchacovirus P83/plateau pika/China (Figure 3A). As cleavage

motifs need to be accessible to proteases, we mapped this region in the tertiary structure of SADS-

CoV and show that it is located in an exposed loop, like region 3 (S1/S2 cleavage site) and region

4 (S2' cleavage site, Figure 3C). Therefore, identifying cleavage sites in divergent CoVs together

with their mapping on crystalized tertiary structures can help find regions that may be critical in

the context of the virus-host interactions, and which can be targeted for future laboratory testing.

Aside from the two viruses reported in this study, we also found two additional luchacoviruses

(UMN2020/mouse/2018/USA (58) and P83/plateau pika/China (59)) that are highly divergent

(Figure 2, supplementary figure 1), thus showing that luchacoviruses are highly diverse and

surveillance targeting both conserved and variable regions like the S1 may be essential to

understand CoV diversity and evolution. With the results shown in this study we increased the

dataset of sequences from luchacoviruses extending the known host range of these viruses and

showing possible regions of interest in the S protein for future surveillance.

We did not find recombination breakpoints in the genomic region analyzed between the

luchacovirus sequences available to date. However possible recombination breakpoints have been

detected in the ORF1ab between viruses of groups 1, 2 and 3 (53). This may indicate that these

viruses are infecting different hosts at the same time. In this study we are reporting divergent

luchacoviruses in three species of widely spread meso-carnivore, whose diet include rabbits and

rodents, and thus it would be interesting to assess if the predator-prey interactions play a role in

the spread of CoVs in wild populations. However, we did not find luchacoviruses in the rodent

and rabbit samples screened in this study (Supplementary Table 1) possibly due to our smaller

sample size (36 samples) compared to that of the meso-carnivore (300 samples) in which we found

luchacoviruses in 1% (3 positive samples). In previous studies from China (25) and Europe (60) it

has been shown that luchacoviruses are less prevalent than rodent and rabbit betacoronaviruses

(0.8% to 1.1% of positive samples were luchacoviruses compared to 6.1 to 11.8% for rodent

betacoronaviruses). However, we found that aside from the two variants reported in this study from

meso-carnivores in the U.S, we also found that UMN2020/mouse/2018/USA which was collected

from a mouse bought at a pet store in the USA in 2018 (58) is also highly variable and did not

group with other luchacoviruses collected in mice from China (Figure 1 and 2) despite having the

same genetic organization (Figure 4). Therefore, to date luchacoviruses recovered from rodents

and meso-carnivore in the U.S. are highly variable when compared to those from Asia and Europe

(Figures 2 and 5). It is necessary to continue the surveillance of wildlife in the U.S. to increase our

knowledge of the viral diversity hosted in native species. Since all the recovered viral sequences

herein reported were obtained from fecal samples and not from tissue samples, we show it is

possible to perform a successful surveillance of alphacoronaviruses in meso-carnivore in a non-

invasive manner by screening fecal samples.

Acknowledgements

This work was funded by the Cornell Public and Ecosystem Health Impact Award. Capacity for

coronavirus discovery screening was supported (FOA PAR-18-604) by the US Food and Drug

Administration's Veterinary Laboratory Investigation and Response Network under grant

1U18FD006993-01.

The authors thank Melisa A. Fadden for assistance with sample storage and retrieval; Christian

Lange for assistance with screening protocols; Guillaume Reboul and Joe Flint for technical

support. RNA-seq was conducted by scientists in the Transcriptional Regulation and Expression

Facility at Cornell University (RRID: SCR_022532).

Data Availability

Sequences obtained in this study are deposited in GenBank (accession numbers OQ756331-35).

571
572   **References**
573

574   1.   Siddell SG, Walker PJ, Lefkowitz EJ, Mushegian AR, Adams MJ, Dutilh BE,
575   Gorbalenya AE, Harrach B, Harrison RL, Junglen S, Knowles NJ, Kropinski AM,
576   Krupovic M, Kuhn JH, Nibert M, Rubino L, Sabanadzovic S, Sanfaçon H, Simmonds P,
577   Varsani A, Zerbini FM, Davison AJ. 2019. Additional changes to taxonomy ratified in a
578   special vote by the International Committee on Taxonomy of Viruses (October 2018).
579   Arch Virol 164:943-946.
580   2.   Cui J, Li F, Shi Z-L. 2019. Origin and evolution of pathogenic coronaviruses. Nature
581   Reviews Microbiology 17:181-192.
582   3.   Li F. 2015. Receptor recognition mechanisms of coronaviruses: a decade of structural
583   studies. J Virol 89:1954-64.
584   4.   Belouzard S, Millet JK, Licitra BN, Whittaker GR. 2012. Mechanisms of coronavirus cell
585   entry mediated by the viral spike protein. Viruses 4:1011-33.
586   5.   Millet JK, Whittaker GR. 2015. Host cell proteases: Critical determinants of coronavirus
587   tropism and pathogenesis. Virus Res 202:120-34.
588   6.   Belouzard S, Chu VC, Whittaker GR. 2009. Activation of the SARS coronavirus spike
589   protein via sequential proteolytic cleavage at two distinct sites. Proc Natl Acad Sci U S A
590   106:5871-6.
591   7.   Wu Y, Zhao S. 2020. Furin cleavage sites naturally occur in coronaviruses. Stem Cell
592   Res 50:102115.
593   8.   Chan YA, Zhan SH. 2022. The Emergence of the Spike Furin Cleavage Site in SARS-
594   CoV-2. Mol Biol Evol 39.
595   9.   Jaimes JA, Whittaker GR. 2018. Feline coronavirus: Insights into viral pathogenesis
596   based on the spike protein structure and function. Virology 517:108-121.
597   10.   Gong L, Li J, Zhou Q, Xu Z, Chen L, Zhang Y, Xue C, Wen Z, Cao Y. 2017. A New
598   Bat-HKU2-like Coronavirus in Swine, China, 2017. Emerg Infect Dis 23:1607-9.
599   11.   Kim J, Yoon J, Park JE. 2022. Furin cleavage is required for swine acute diarrhea
600   syndrome coronavirus spike protein-mediated cell - cell fusion. Emerg Microbes Infect
601   11:2176-2183.
602   12.   Li F. 2016. Structure, Function, and Evolution of Coronavirus Spike Proteins. Annual
603   Review of Virology 3:237-261.

13. Jaimes JA, Millet JK, Stout AE, Andre NM, Whittaker GR. 2020. A Tale of Two
Viruses: The Distinct Spike Glycoproteins of Feline Coronaviruses. Viruses 12.
14. Choe Y, Leonetti F, Greenbaum DC, Lecaille F, Bogyo M, Brömme D, Ellman JA, Craik
CS. 2006. Substrate profiling of cysteine proteases using a combinatorial peptide library
identifies functionally unique specificities. J Biol Chem 281:12824-32.
15. Yang TJ, Chang YC, Ko TP, Draczkowski P, Chien YC, Chang YC, Wu KP, Khoo KH,
Chang HW, Hsu SD. 2020. Cryo-EM analysis of a feline coronavirus spike protein
reveals a unique structure and camouflaging glycans. Proc Natl Acad Sci U S A
117:1438-1446.
16. Tang T, Jaimes JA, Bidon MK, Straus MR, Daniel S, Whittaker GR. 2021. Proteolytic
Activation of SARS-CoV-2 Spike at the S1/S2 Boundary: Potential Role of Proteases
beyond Furin. ACS Infect Dis 7:264-272.
17. Hatmal MM, Alshaer W, Al-Hatamleh MAI, Hatmal M, Smadi O, Taha MO, Oweida AJ,
Boer JC, Mohamud R, Plebanski M. 2020. Comprehensive Structural and Molecular
Comparison of Spike Proteins of SARS-CoV-2, SARS-CoV and MERS-CoV, and Their
Interactions with ACE2. Cells 9.
18. Licitra BN, Millet JK, Regan AD, Hamilton BS, Rinaldi VD, Duhamel GE, Whittaker
GR. 2013. Mutation in spike protein cleavage site and pathogenesis of feline coronavirus.
Emerg Infect Dis 19:1066-73.
19. Johnson BA, Xie X, Bailey AL, Kalveram B, Lokugamage KG, Muruato A, Zou J,
Zhang X, Juelich T, Smith JK, Zhang L, Bopp N, Schindewolf C, Vu M, Vanderheiden
A, Winkler ES, Swetnam D, Plante JA, Aguilar P, Plante KS, Popov V, Lee B, Weaver
SC, Suthar MS, Routh AL, Ren P, Ku Z, An Z, Debbink K, Diamond MS, Shi PY,
Freiberg AN, Menachery VD. 2021. Loss of furin cleavage site attenuates SARS-CoV-2
pathogenesis. Nature 591:293-299.
20. Millet JK, Whittaker GR. 2014. Host cell entry of Middle East respiratory syndrome
coronavirus after two-step, furin-mediated activation of the spike protein. Proc Natl Acad
Sci U S A 111:15214-9.
21. Sander AL, Moreira-Soto A, Yordanov S, Toplak I, Balboni A, Ameneiros RS, Corman
V, Drosten C, Drexler JF. 2022. Genomic determinants of Furin cleavage in diverse
European SARS-related bat coronaviruses. Commun Biol 5:491.
22. Stout AE, Millet JK, Stanhope MJ, Whittaker GR. 2021. Furin cleavage sites in the spike
proteins of bat and rodent coronaviruses: Implications for virus evolution and zoonotic
transfer from rodent species. One Health 13:100282.
23. Tsoleridis T, Chappell JG, Onianwa O, Marston DA, Fooks AR, Monchatre-Leroy E,
Umhang G, Müller MA, Drexler JF, Drosten C, Tarlinton RE, McClure CP, Holmes EC,
Ball JK. 2019. Shared Common Ancestry of Rodent Alphacoronaviruses Sampled
Globally. Viruses 11.
24. Wu Z, Lu L, Du J, Yang L, Ren X, Liu B, Jiang J, Yang J, Dong J, Sun L, Zhu Y, Li Y,
Zheng D, Zhang C, Su H, Zheng Y, Zhou H, Zhu G, Li H, Chmura A, Yang F, Daszak P,
Wang J, Liu Q, Jin Q. 2018. Comparative analysis of rodent and small mammal viromes
to better understand the wildlife origin of emerging infectious diseases. Microbiome
6:178.
25. Ge XY, Yang WH, Zhou JH, Li B, Zhang W, Shi ZL, Zhang YZ. 2017. Detection of
alpha- and betacoronaviruses in rodents from Yunnan, China. Virol J 14:98.

26. Choi A, Singleton DT, Stout AE, Millet JK, Whittaker GR. 2021. In vitro and
computational analysis of the putative furin cleavage site (RRARS) in the divergent spike
protein of the rodent coronavirus AcCoV-JC34 (sub-genus luchacovirus). bioRxiv
doi:10.1101/2021.12.16.473025:2021.12.16.473025.
27. Ray JC. 2000. Meso-carnivore of Northeastern North America: Status
and Conservation Issues. WCS Working Papers No 15. Wildlife Conservation Society, Bronx,
New York.
28. Prange S, Gehrt SD, Wiggers EP. 2003. Demographic Factors Contributing to High
Raccoon Densities in Urban Landscapes. The Journal of Wildlife Management 67:324-
333.
29. Kimpston CN, Hatke AL, Castelli B, Otto N, Tiffin HS, Machtinger ET, Brown JD, Van
Why KR, Marconi RT. 2022. High Prevalence of Antibodies against Canine Parvovirus
and Canine Distemper Virus among Coyotes and Foxes from Pennsylvania: Implications
for the Intersection of Companion Animals and Wildlife. Microbiol Spectr 10:e0253221.
30. Sabeta CT, Mansfield KL, McElhinney LM, Fooks AR, Nel LH. 2007. Molecular
epidemiology of rabies in bat-eared foxes (Otocyon megalotis) in South Africa. Virus
Res 129:1-10.
31. Roemer GW, Gompper ME, Van Valkenburgh B. 2009. The Ecological Role of the
Mammalian Mesocarnivore. BioScience 59:165-173.
32. Vlasova AN, Halpin R, Wang S, Ghedin E, Spiro DJ, Saif LJ. 2011. Molecular
characterization of a new species in the genus Alphacoronavirus associated with mink
epizootic catarrhal gastroenteritis. J Gen Virol 92:1369-1379.
33. Murray J, Kiupel M, Maes RK. 2010. Ferret coronavirus-associated diseases. Vet Clin
North Am Exot Anim Pract 13:543-60.
34. Tekes G, Thiel HJ. 2016. Feline Coronaviruses: Pathogenesis of Feline Infectious
Peritonitis. Adv Virus Res 96:193-218.
35. Erles K, Brownlie J. 2008. Canine respiratory coronavirus: an emerging pathogen in the
canine infectious respiratory disease complex. Vet Clin North Am Small Anim Pract
38:815-25, viii.
36. Dong BQ, Liu W, Fan XH, Vijaykrishna D, Tang XC, Gao F, Li LF, Li GJ, Zhang JX,
Yang LQ, Poon LL, Zhang SY, Peiris JS, Smith GJ, Chen H, Guan Y. 2007. Detection of
a novel and highly divergent coronavirus from asian leopard cats and Chinese ferret
badgers in Southern China. J Virol 81:6920-6.
37. Kim YI, Kim SG, Kim SM, Kim EH, Park SJ, Yu KM, Chang JH, Kim EJ, Lee S, Casel
MAB, Um J, Song MS, Jeong HW, Lai VD, Kim Y, Chin BS, Park JS, Chung KH, Foo
SS, Poo H, Mo IP, Lee OJ, Webby RJ, Jung JU, Choi YK. 2020. Infection and Rapid
Transmission of SARS-CoV-2 in Ferrets. Cell Host Microbe 27:704-709.e2.
38. Oude Munnink BB, Sikkema RS, Nieuwenhuijse DF, Molenaar RJ, Munger E,
Molenkamp R, van der Spek A, Tolsma P, Rietveld A, Brouwer M, Bouwmeester-
Vincken N, Harders F, Hakze-van der Honing R, Wegdam-Blans MCA, Bouwstra RJ,
GeurtsvanKessel C, van der Eijk AA, Velkers FC, Smit LAM, Stegeman A, van der Poel
WHM, Koopmans MPG. 2021. Transmission of SARS-CoV-2 on mink farms between
humans and mink and back to humans. Science 371:172-177.
39. Vlasova AN, Diaz A, Damtie D, Xiu L, Toh T-H, Lee JS-Y, Saif LJ, Gray GC. 2021.
Novel Canine Coronavirus Isolated from a Hospitalized Patient With Pneumonia in East
Malaysia. Clinical Infectious Diseases 74:446-454.

40. Zehr JD, Pond SLK, Martin DP, Ceres K, Whittaker GR, Millet JK, Goodman LB,
Stanhope MJ. 2022. Recent Zoonotic Spillover and Tropism Shift of a Canine
Coronavirus Is Associated with Relaxed Selection and Putative Loss of Function in NTD
Subdomain of Spike Protein. Viruses 14:853.
41. Olarte-Castillo XA, Dos Remedios JF, Heeger F, Hofer H, Karl S, Greenwood AD, East
ML. 2021. The virus-host interface: Molecular interactions of Alphacoronavirus-1
variants from wild and domestic hosts with mammalian aminopeptidase N. Mol Ecol
30:2607-2625.
42. Campbell SJ, Ashley W, Gil-Fernandez M, Newsome TM, Di Giallonardo F, Ortiz-Baez
AS, Mahar JE, Towerton AL, Gillings M, Holmes EC, Carthey AJR, Geoghegan JL.
2020. Red fox viromes in urban and rural landscapes. Virus Evol 6:veaa065.
43. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of
high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol
7:539.
44. Tamura K, Stecher G, Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics
Analysis Version 11. Mol Biol Evol 38:3022-3027.
45. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32:1792-7.
46. Lole KS, Bollinger RC, Paranjape RS, Gadkari D, Kulkarni SS, Novak NG, Ingersoll R,
Sheppard HW, Ray SC. 1999. Full-length human immunodeficiency virus type 1
genomes from subtype C-infected seroconverters in India, with evidence of intersubtype
recombination. J Virol 73:152-60.
47. Martin DP, Varsani A, Roumagnac P, Botha G, Maslamoney S, Schwab T, Kelz Z,
Kumar V, Murrell B. 2021. RDP5: a computer program for analyzing recombination in,
and removing signals of recombination from, nucleotide sequence datasets. Virus Evol
7:veaa087.
48. Martin D, Rybicki E. 2000. RDP: detection of recombination amongst aligned sequences.
Bioinformatics 16:562-3.
49. Peacock TP, Goldhill DH, Zhou J, Baillon L, Frise R, Swann OC, Kugathasan R, Penn R,
Brown JC, Sanchez-David RY, Braga L, Williamson MK, Hassard JA, Staller E, Hanley
B, Osborn M, Giacca M, Davidson AD, Matthews DA, Barclay WS. 2021. The furin
cleavage site in the SARS-CoV-2 spike protein is required for transmission in ferrets.
Nature Microbiology 6:899-909.
50. Duckert P, Brunak S, Blom N. 2004. Prediction of proprotein convertase cleavage sites.
Protein Eng Des Sel 17:107-12.
51. Yu J, Qiao S, Guo R, Wang X. 2020. Cryo-EM structures of HKU2 and SADS-CoV
spike glycoproteins provide insights into coronavirus evolution. Nature Communications
11:3070.
52. Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE.
2018. UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein
Sci 27:14-25.
53. Wang W, Lin XD, Zhang HL, Wang MR, Guan XQ, Holmes EC, Zhang YZ. 2020.
Extensive genetic diversity and host range of rodent-borne coronaviruses. Virus Evol
6:veaa078.

54. Coutard B, Valle C, de Lamballerie X, Canard B, Seidah NG, Decroly E. 2020. The spike
glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent
in CoV of the same clade. Antiviral Res 176:104742.
55. Fu X, Fang B, Liu Y, Cai M, Jun J, Ma J, Bu D, Wang L, Zhou P, Wang H, Zhang G.
2018. Newly emerged porcine enteric alphacoronavirus in southern China: Identification,
origin and evolutionary history analysis. Infect Genet Evol 62:179-187.
56. Edwards CE, Yount BL, Graham RL, Leist SR, Hou YJ, Dinnon KH, 3rd, Sims AC,
Swanstrom J, Gully K, Scobey TD, Cooley MR, Currie CG, Randell SH, Baric RS. 2020.
Swine acute diarrhea syndrome coronavirus replication in primary human cells reveals
potential susceptibility to infection. Proc Natl Acad Sci U S A 117:26915-26925.
57. Guo Y-Y, Wang P-H, Pan Y-Q, Shi R-Z, Li Y-Q, Guo F, Xing L. 2021. The
Characteristics of Spike Glycoprotein Gene of Swine Acute Diarrhea Syndrome
Coronavirus Strain CH/FJWT/2018 Isolated in China. Frontiers in Veterinary Science 8.
58. Fay EJ, Balla KM, Roach SN, Shepherd FK, Putri DS, Wiggen TD, Goldstein SA,
Pierson MJ, Ferris MT, Thefaine CE, Tucker A, Salnikov M, Cortez V, Compton SR,
Kotenko SV, Hunter RC, Masopust D, Elde NC, Langlois RA. 2022. Natural rodent
model of viral transmission reveals biological features of virus population dynamics. J
Exp Med 219.
59. Zhu W, Yang J, Lu S, Jin D, Wu S, Pu J, Luo X-l, Liu L, Li Z, Xu J. 2021. Discovery and
Evolution of a Divergent Coronavirus in the Plateau Pika From China That Extends the
Host Range of Alphacoronaviruses. Frontiers in Microbiology 12.
60. Monchatre-Leroy E, Boué F, Boucher JM, Renault C, Moutou F, Ar Gouilh M, Umhang
G. 2017. Identification of Alpha and Beta Coronavirus in Wildlife Species in France:
Bats, Rodents, Rabbits, and Hedgehogs. Viruses 9.
764