BrainAGE as a measure of maturation during early adolescence

Lucy B. Whitmore[1], Sara J. Weston[1], Kathryn L. Mills[1,2]

Affiliations:

[1]University of Oregon

[2]University of Oslo

Author Note

No competing interests exist.

Correspondence concerning this article should be addressed to Lucy B. Whitmore, Department

of Psychology, 1227 University of Oregon, Eugene, OR 97403. E-mail: lwhitmor@uoregon.edu

Abstract

The Brain-Age Gap Estimation (BrainAGE) is an important new tool that purports to evaluate brain maturity when used in adolescent populations. However, it is unclear whether BrainAGE tracks with other maturational metrics in adolescence. In the current study, we related BrainAGE to metrics of pubertal and cognitive development using both a previously validated model and a novel model trained specifically on an early adolescent population. The previously validated model was used to predict BrainAGE in two age bands, 9-11 and 11-13 years old, while the novel model was used with 9-11 year olds only. Across both models and age bands, an older BrainAGE was related to more advanced pubertal development. The relationship between BrainAGE and cognition was less clear, with conflicting relationships across the two models. The results of the current study provide initial evidence that BrainAGE tracks with some metrics of maturation, including pubertal development. However, the conflicting results between BrainAGE and cognition lead us to question the utility of these models for non-biological processes.

Introduction

Over the last 25 years, we have learned a considerable amount about the prolonged development of the human brain, due in part to advances in neuroimaging technologies and methods (Bethlehem et al., 2022; Mills et al., 2014; Raznahan et al., 2011; Wierenga et al., 2014). Brain development progresses differently across individuals (Mills et al., 2021), with different patterns and trajectories theorized to relate to outcomes such as psychopathology (Shaw et al., 2010). While longitudinal data is necessary to track individual trajectories, most neuroimaging studies that examine the brain across the lifespan are cross-sectional, and there has been an increasing desire to find ways to measure individual differences in brain maturation using methods that are compatible with cross-sectional data. The Brain-Age Gap Estimation (BrainAGE) has been proposed as one approach to assess the maturity of an individual's brain, as this measure reflects the difference between an individual's chronological age and their age as predicted by machine learning algorithms trained on neuroimaging data, often structural MRI (Brown et al., 2012; Franke et al. 2010). With this approach, an individual receives an estimated brain age that can differ from their chronological age.

BrainAGE as a measure of aging and pathology

BrainAGE has most frequently been used in older adult populations, as it was originally proposed as a tool to research and assess risk for aging-related diseases (Franke et al., 2010). For older adults, differences between an individual's estimated brain age and chronological age have been thought to reflect potential deviations from a normative aging trajectory related to various health concerns. This has been supported by previous work that has demonstrated having a BrainAGE that is older than one's chronological age is related to metrics and outcomes of cognitive aging and decline, such as Alzheimer's, in older adults (Franke & Gaser, 2012; 2019;

Biondo et al., 2022). Additionally, across adulthood, having an older BrainAGE than one's chronological age has also been associated with psychopathologies including depression and schizophrenia (Han et al., 2021; 2022; Schnack et al., 2016).

BrainAGE in adolescence

Although BrainAGE is more commonly used in investigations of older adult populations, researchers are starting to use BrainAGE as a tool to advance current theories in developmental cognitive neuroscience. As BrainAGE was not originally developed as a measure of individual differences in adolescent brain maturation, the implications for interpreting BrainAGE are less clear in this population than in older populations where an older-appearing brain is more straightforwardly related to aging-related diseases. Conventionally, differences between BrainAGE and chronological age have been interpreted as reflecting accelerated or decelerated brain maturation in children and adolescents, though this has not been validated with longitudinal data (Franke et al., 2012; Lewis et al., 2018).

BrainAGE prediction based on structural MRI has been shown to accurately predict age in adolescents as measured by mean absolute errors in the one- to two-year range (Franke et al., 2012). Since the inception of using BrainAGE in adolescent populations, there have been efforts to link an adolescent's BrainAGE to psychopathologies, which often emerge during this time. Recently, BrainAGE has shown promise as a potential biomarker of mental health issues in adolescent populations. Notably, having a BrainAGE that is older than one's chronological age has been linked to major depression, functional impairment, and risk of psychosis (Drobonin et al., 2022; Chung et al., 2018; Copley et al., 2021). This is consistent with the idea that deviations from typical developmental trajectories play a role in the development of psychological disorders, as hypothesized by Shaw et al. (2010). Similarly, the Stress Acceleration Hypothesis

(Callaghan & Tottenham, 2016) proposes that early life stress accelerates neural development, potentially resulting in short-term benefits but long-term vulnerability to psychopathology.

However, in recent years BrainAGE has also been theorized to reflect more than just accelerated or decelerated brain maturation (Vidal-Pineiro et al., 2021, Ball et al., 2021, Kelly et al., 2022). In adults, BrainAGE has been found to relate to birth weight and polygenic brain age scores (Videl-Pineiro et al., 2021). Additionally, adult BrainAGE varies over the course of the menstrual cycle (Franke et al., 2015). In children and adolescents, BrainAGE has been related to genetic factors (Brouwer et al., 2021). Despite this, the success of BrainAGE prediction in adolescent populations suggests its potential use as a clinically-relevant biomarker; however, given a lack of clarity around interpretations of adolescent BrainAGE, additional validation is needed before BrainAGE can be recommended for clinical and applied settings. In particular, it remains unclear as to which aspects of maturation are captured by this measure and which are not. A challenge to validation is the lack of conceptual clarity around the concept of "brain maturity" and what it means for a brain to be mature (Somerville, 2016).

The validation of any measure interpreted as reflecting a maturational process is an important step, and one that has been accomplished for developmental metrics such as the Tanner stages, also known as the Tanner Scale (Tanner, 1962). The Tanner stages, developed in the 1960s, describe and quantify physical changes associated with puberty. The stages were validated by observing adolescents and correlating physical changes with the corresponding stage, making it a widely accepted measure of physical maturation in clinical and research settings. In the case of adolescent BrainAGE, we would expect a similar relationship to be present, where BrainAGE is related to other maturational metrics that develop over adolescence.

In the current study, we explore the relationship of BrainAGE to two domains of changes that are occurring in this time: pubertal and cognitive development.

Brain maturation and puberty

Pubertal development is often measured in stages, such as Tanner stages or using the Pubertal Development Scale (Tanner, 1962; Petersen et al., 1988). Tanner stages measure physical development, based on changes in pubic hair, genitalia, and breast growth (for females). Based on these metrics, Tanner Stage 1 represents pre-puberty, Stages 2-4 represent mid/intermediate puberty, and Stage 5 represents reproductive maturity. The Pubertal Development Scale examines similar, though not entirely overlapping, measures as Tanner stages and includes questions about characteristics such as body hair, growth in height, and other secondary sex characteristics (Petersen et al., 1988). Pubertal hormones such as testosterone or estradiol are also commonly examined in relation to pubertal development, but no current scale exists to assess pubertal development progress based on levels of certain hormones.

Independent of age, pubertal stage has been related to changes in brain structures, both cortical and subcortical. While cross-sectional relationships between subcortical volumes and pubertal development have found mixed results, longitudinal work has more consistently found a positive relationship between pubertal stage and amygdala and hippocampus volumes and a negative relationship between pubertal stage and nucleus accumbens, caudate, putamen, and globus pallidus volumes (Goddings et al., 2014; Herting and Sowell, 2017). Pubertal stage has also been found to be a better predictor of development than age in a number of subcortical regions, including the caudate, pallidum, and hippocampus (Wierenga et al., 2018).

A previous literature review found that both cross-sectional and longitudinal studies report widespread patterns of reductions in cortical gray matter in relation to pubertal development, both stage and timing (Vijayakumar et al., 2018). Notably, significant negative associations have been found between the pubertal hormone testosterone and cortical thinning in regions including the left posterior cingulate, precuneus, dlPFC and ACC in males and right somatosensory cortex in females (Nguyen, 2012). Additionally, change in Tanner stage has been related to rates of change in cortical surface area, where a main effect of stage was found for decreases in left precuneus surface area (Herting et al., 2015). More recent work has shown relationships between pubertal stage and increased cortical thinning in frontal and parietal cortices, as well as some temporal regions, independent of age (Vijayakumar et al., 2021).

In terms of pubertal development, recent work exists on the relationship between puberty and adolescent BrainAGE. Using convolutional neural nets, Holm et al. (2023) found a positive relationship between BrainAGE and parent-report PDS scores. Additionally, Dehestani et al. (2023) found a positive relationship between BrainAGE and "puberty age", a new measure proposed to assess pubertal timing based on observed physical development (PDS) and pubertal hormones (testosterone and DHEA).

Brain maturation and cognition

In addition to pubertal changes, adolescents undergo a number of cognitive changes, such as improved working memory performance and inhibitory control (Cromer et al. 2015; Davidson et al. 2006). Performance on the NIH Toolbox Cognition Battery, a commonly used measure of cognition, has been shown to improve cross-sectionally and longitudinally over early-to-mid adolescence (Anokhin et al., 2022). Additionally, working memory improvements in adolescence

have been found to be associated with cortical volume reductions in bilateral prefrontal and posterior parietal regions and in regions around the central sulci (Tamnes et al, 2015).

BrainAGE and cognition in adolescence

Past research on cognition and BrainAGE in adolescence has found mixed results, with some work indicating that a positive BrainAGE is related to faster processing speed (Erus et al., 2015) and others finding that a negative BrainAGE was associated with better cognitive performance (Lewis et al., 2018). Additionally, some studies have found very small or non-significant effects (Ball et al., 2017; 2021; Khudrakham et al., 2015, Kelly et al., 2022).

These studies exhibit a multitude of sources of variation, spanning from model features employed to train BrainAGE models, to participant age ranges, as well as the types of cognitive measures employed. In the studies described above, model features employed encompassed cortical thickness exclusively, T1 white/gray contrast, cortical thickness/volume/area, and subcortical volume, as well as the combination of gray matter, white matter, and ventricular measures. Parcellation methods varied significantly as well. Cognitive measurements employed in the studies were also highly varied, including IQ, measures of speed and accuracy on the Penn Computerized Neurocognitive Battery, and the NIH Toolbox Cognition Battery (Gur et al., 2012; Gershon et al., 2013). Although all of the studies incorporated adolescent participants, the age ranges varied significantly and the distribution of cognitive data across the age range was not always clearly indicated. Age ranges in certain studies were as broad as 8-22, 3-21, and 4.5-18.5 years, while Kelly et al. (2022) was restricted to cognition scores from participants who were 13 years old.

Current Study

In the current study, we examine the relationship between BrainAGE and metrics of pubertal and cognitive maturation within two narrow age bands in early adolescence, 9-11 years and 10-13 years. For BrainAGE models trained on structural MRI data, such as the models in this paper, we expect that BrainAGE would track with maturational metrics that are reflected in changes to brain structure and change during these age ranges, including pubertal and cognitive development.

The current study used two different BrainAGE models in order to provide additional confidence in our results by comparing models trained on both wide and narrow age ranges in order to balance specificity and replication. This strategy enabled us to balance providing the models with extensive data from our age range of interest, while not overly restricting our training data and possible predictions. In Study 1, we used an existing, previously validated BrainAGE model trained on a wide age range to predict BrainAGE in a sample of early adolescents (Drobonin et al. 2022). Owing to the large sample size of the ABCD study, we were able to address our questions using a split sample approach as recommended by Saragosa-Harris et al. (2022). We opted for this approach to determine if relationships would replicate across two samples, increasing our confidence in any reported findings.

In Study 2, we trained and tested a new BrainAGE model trained specifically on our age range of interest (9-11 years old). The development of a BrainAGE model trained specifically on data from early adolescents is novel, and its usage reflects the unique structural changes occurring during this age range. The goal of the novel model is to capture the specific changes happening in this range, as opposed to the dynamic changes occurring over all of adolescence. While BrainAGE models exist that include adolescents, none sample specifically for early

adolescents. The novel model is available for public use and can be found at

https://github.com/LucyWhitmore/BrainAGE-Maturation.

      Using predictions from the novel model, BrainAGE estimates were examined in

association with youth-report pubertal development, parent-report pubertal development, and

cognition scores.

      As the goal of the study was to examine relationships between BrainAGE and other

forms of maturation and determine whether these relationships replicate across models and age

bands, the methods of Study 1 and Study 2 were kept as similar as possible. In particular, model

training code, analysis procedures, and pubertal/cognitive measures were identical across Study

1 and Study 2. Differences in samples and procedures are described where present.

Study 1

Participants

Model Training

      The model used in Study 1 was a previously validated BrainAGE model, created for use

in Drobonin et al. (2022). The model was trained on a composite of six samples, with

participants ranging from 9-19 years old. Inclusion criteria included being within the age range

of 9-19 years old and passing automated MRI quality control procedures. Additionally, only

participants without psychiatric disorders and with an IQ over 75 were included in model

training and testing.

Analyses

      Adolescents included in the analyses were participants in the Adolescent Brain Cognitive

Development (ABCD) Study. The ABCD Study is a large longitudinal study that has recruited

participants from 21 sites across the United States. Informed consent was obtained from all

participants and their parents. For further details on recruitment procedures in the ABCD Study, see Garavan et al. (2018).

While the ABCD Annual Release 4.0 included both baseline data and data from the one- and two-year follow-ups, only baseline and the two-year follow-up included imaging data. Baseline imaging data were available for 11,878 participants between the ages of 9-11 years old. After excluding participants with missing or low-quality structural MRI data, 11,402 participants remained (M = 9.92 years old, SD = 0.63 years). Follow-up imaging data were available for 7,827 participants between the ages of 10 and 13. After excluding participants with missing or low-quality structural MRI data, 7,696 participants remained (M = 11.94 years old, SD = 0.65 years). Additional participant demographic details are available in *Table 1*.

Table 1. Participant demographics.

		Baseline (Model 1)	*Follow-Up*	*Baseline (Model 2)*
Sex	*Female*	48% (*N*=5471)	46.3% (*N* = 3566)	48.4% (*N* = 2759)
	Male	52% (*N* = 5931)	53.7% (*N*= 4129)	51.6% (*N* = 2943)
Race/ Ethnicity	*Non-Hispanic White*	53.3% (*N*=5962)	55.1% (*N*=4242)	51.6% (*N*=2946)
	Hispanic/Latino	20.5% (*N*=2332)	19.3% (*N*=1487)	21.5% (*N*=1223)
	Non-Hispanic Black	15.4% (*N*=1754)	13.1% (*N*=1010)	14.3% (*N*=815)
	Asian	2.1% (*N*=244)	1.9% (*N* =145)	2.1% (*N* =122)
	Native American/ Alaska Native	0.5% (*N* =56)	0.6% (*N* =44)	0.5% (*N* =30)
	Multiracial	12% (*N* =1,370)	12.1% (*N* =931)	12.1% (*N* =688)

	Additional Race	0.4% (*N* =40)	0.4% (*N* =31)	0.3% (*N* =18)
Household Income	*<$5000*	3.4 % (*N* =392)	2.5% (*N* =196)	3.3% (*N* =188)
	$5,000 - $11,999	3.5% (*N* =398)	2.7% (*N* =211)	3.5% (*N* =199)
	$12,000 - $15,999	2.3% (*N* =265)	1.8% (*N* =140)	2.3% (*N* =130)
	$16,000 - $24,999	4.4% (*N* =497)	3.2% (*N* =246)	4.2% (*N* =238)
	$25,000 - $34,999	5.6% (*N* =633)	5.2% (*N* =402)	5.7% (*N* =324)
	$35,000 - $49,999	7.8% (*N* =889)	6.8% (*N* =524)	8.3% (*N* =471)
	$50,000 - $74,999	12.5% (*N* =1,429)	12.7% (*N* =980)	13.1% (*N* =747)
	$75,000 - $99,999	13.3% (*N* =1,520)	13.3% (*N* =1,026)	12.8% (*N* =725)
	$100,000 - $199,999	28.1 % (*N* =3,201)	31.4% (*N* =2,413)	27.6% (*N* =1,575)
	≥ $200,000	10.6% (*N* =1,208)	12.49% (*N* =961)	10.6% (*N* =603)
	Missing	8.5% (*N* =970)	7.7% (*N* =596)	8.8% (*N* =501)
Model Training & Testing	*# Participants (Training)*	1,621	1,621	5,701
	Age Range (Training)	9-19 y, m = 13.5 ± 3.04	9-19 y, m = 13.5 ± 3.04	8.9–11.08 y, m = 9.92 ± 63
	# Participants (Testing)	11,402	7,695	5,701
	Age Range (Testing)	8.9–11.08 y, m = 9.92 ±.63	10.58-13.83 y, m = 11.94 ± .65	8.9–11.08 y, m = 9.92 ± .63

Materials & Methods

MRI Acquisition

For details on the acquisition of MRI images used to train the model used in Study 1, see

Drobonin et al. (2022). For participants included in the analyses, detailed imaging procedures

can be found in Casey et al. (2018) and Hagler et al. (2019). Harmonized protocols were used

across 21 sites to collect imaging data using one of three 3T scanner platforms (Phillips,

Siemens, or GE) and a 32-channel head coil; 3-dimensional T1-weighted scans were collected

with 1-mm voxel resolution.

MRI Processing and Model Features

Model features included cortical gray matter volume and surface area measurements, as

well as bilateral global and subcortical volume measurements. Cortical features were obtained

using the Freesurfer default Desikan-Killiany atlas, while bilateral global and subcortical volume

measures were obtained using the Freesurfer output.

The Drobonin et al. (2022) model was originally trained on 189 structural features, but

only 175 of these features were available in the ABCD dataset. Therefore, for the current study,

the model was used to predict BrainAGE based on the 175 available features (Supplementary

Table 1).

BrainAGE Prediction

For the baseline sample, the Drobinin et al. (2022) model was used to predict BrainAGE

for 11,402 adolescents, who were then split into two samples of 5,701 participants each for

further analyses. The two samples did not differ in age ($t(11395.91) = 0.26$, $p = .796$, $d < 0.01$) or

sex ($\chi^2(1) = 1.42$, $p = .233$).

For the two-year follow-up sample, the model was used to predict BrainAGE for 7,696

adolescents, who were then split into two samples of 3,848 participants for further analyses. The

two samples did not differ in age ($t(7692.42) = -0.09$, $p = .930$, $d < 0.01$) or sex (χ^2 (1) = 0.02, p =.885).

Bias Correction

Like other machine learning models, BrainAGE models can be susceptible to prediction bias towards the group mean (Le et al., 2018). In the case of BrainAGE models, this bias often takes the form of younger participants being more likely to be predicted as slightly older than their chronological age, while older participants are predicted to be slightly younger. To adjust for this age bias, we followed the bias correction procedures described in Smith et al. (2019). Briefly, the bias correction procedures require fitting a linear model to the validation set and extracting the intercept and slope. To obtain a corrected estimate, the intercept is subtracted from predicted age, and then value is then divided by the slope. The slope and intercept are generalizable to new data, allowing these coefficients to be used to correct test set bias (Peng et al., 2021). Additionally, age was included as a covariate in analyses to further correct for age-related bias.

Behavioral & Survey Measures

Youth & Parent Report Pubertal Development

Both participants and their parents completed the Pubertal Development Scale (PDS; Petersen et al., 1988; Barch et al., 2018). The PDS is a five-item scale that measures pubertal stage. Of the five items, three are asked of every participant (related to growth in height, skin changes, and body hair growth), and two are dependent on the participants' assigned sex at birth. The female version asks about the onset of menstruation and breast growth, while the male version asks about facial hair and vocal changes.

For each participant, mean scores were calculated for each of their self-report and parent-report

scores (α = .53-.81, see Supplementary Table 2 for reliability estimates of the measures).

Participants with incomplete data were excluded from the corresponding analyses.

Cognition

Cognition was measured using the NIH Toolbox Cognition Battery (Bleck et al., 2013;

Gershon et al., 2013b; Hodes et al., 2013). The Toolbox uses seven cognitive tasks, which

measure domains such as cognitive control, working memory, set shifting, and reading ability.

Table 2. NIH Toolbox Cognition Tasks

NIH Toolbox Flanker®	Cognitive Control/Attention
NIH Toolbox List Sorting Working Memory Test®	Working Memory; Categorization; Information Processing
NIH Toolbox Dimensional Change Card Sort®	Flexible thinking; concept formation; set shifting
NIH Toolbox Oral Reading Recognition Test®	Reading Ability; Language; Academic Achievement
NIH Toolbox Pattern Comparison Processing Speed®	Processing Speed; Information Processing
NIH Toolbox Picture Sequence Memory Test®	Visuospatial sequencing & memory
NIH Toolbox Picture Vocabulary Test®	Language; Verbal intellect

NIH Toolbox cognition tasks and their respective cognitive domains.

Summary scores were provided in the ABCD 4.0 data release for each participant based

on their performance on the seven tasks, listed in Table 1 and described in detail in Luciana et al.

(2018). Summary scores are presented as the participants' performance compared to those in the

NIH Toolbox normative sample, which includes nationally representative data from participants

who are between 3-85 years old. Complete cognition summary scores were only available at

baseline. Distributions of all maturational metrics are available in Table 3.

Table 3. Descriptive statistics

Study	Sample	BrainAGE - Corrected		Youth-Report Puberty		Parent-Report Puberty		Cognition	
		Mean ± SD	Range	Mean ± SD	Range	Mean ± SD	Range	Mean ± SD	Range
Study 1, Baseline	Sample 1	0.67 ± 1.75	-4.45 - 8.0	1.71 ± 0.46	1.0 - 4.0	1.60 ± 0.47	1.0 - 4.0	86.38 ± 9.06	53 - 117
	Sample 2	0.7 ± 1.73	-4.2 - 8.48	1.72 ± 0.48	1.0 - 4.0	1.60 ± 0.48	1.0 - 4.0	86.31 ± 9.15	44 - 115
Study 1, Follow-Up	Sample 1	0.18 ± 1.98	-5.02 - 7.7	2.07 ± 0.65	1.0 - 4.0	2.12 ± 0.71	1.0 - 4.0	--	--
	Sample 2	0.20 ± 1.99	-5.64 - 7.74	2.08 ± 0.64	1.0 - 4.0	2.12 ± 0.69	1.0 - 4.0	--	--
Study 2	--	0.0 ± 1.09	-4.36 - 4.26	1.72 ± 0.48	1.0 - 4.0	1.60 ± 0.48	1.0 - 4.0	86.31 ± 9.15	44 - 115

Statistical Analyses

Statistical analyses were performed in RStudio (R version 4.1.1, RStudio version 2021.09.0). Associations with maturational metrics were analyzed with linear models, using the *lm()* function. Separate models were created for the relationship between BrainAGE and each maturational metric, where BrainAGE was regressed on each metric of puberty and cognition. Age was included as a covariate to reduce age-related bias in BrainAGE estimates. We report unstandardized effects and standard errors.

Results

BrainAGE Model Performance

In the original validation of the model, the model performed with a corrected mean absolute error (MAE) of 1.98 (Drobonin et al., 2022). This error is in line with prior adolescent BrainAGE models, which have performed with MAEs frequently in the 1-2 year range (Franke

& Gaser, 2019). The existing model performed with a MAE of 2.32 before correction for age

bias, and a corrected MAE of 1.45 on the ABCD data (baseline) and an uncorrected MAE of 1.3

(corrected MAE 1.58) at follow-up. Model performance is shown in *Figure 1*. A slight age bias

was observed, as is common in these types of models, and was discussed previously. BrainAGE

(predicted age subtracted from chronological age) had a correlation of -0.1 with chronological

age at baseline, and -0.01 at follow-up.

Figure 1. Age versus model predictions for the previously validated model. A) Scan/chronological age versus model-predicted age for the baseline sample. The line represents an exact relationship between chronological age and predicted age. B) Scan/chronological age versus the age-corrected brain-age gap, for the baseline sample. The regression line shows a slight age-related bias, with younger participants predicted to be slightly older than their chronological age. C) Similar to a), scan/chronological age versus predicted age for the follow-up sample. D) Similar to b) scan/chronological age versus the age-corrected brain-age gap for the follow-up sample. Again, the regression line shows a slight amount of age-bias, but less than at baseline.

BrainAGE and Maturational Metrics

In both samples from the baseline wave (9-11 years old), a higher/more positive BrainAGE was related to more advanced youth- and parent-report pubertal development (Table 4). For youth-report pubertal development, (b_{youth} = 0.28, *se* = 0.07; 0.32, se = 0.07, *p* < .001 in both samples) *(Figure 2a)*. For parent-report pubertal development, (b_{parent} = 0.27, *se* = 0.05; 0.38, *se* = 0.05, *p* < .001 in both samples) *(Figure 2b)*. In both baseline sub-samples, a higher/more positive BrainAGE was also related to lower cognition scores (b_{cog} = -0.01, *se* = 0.003; -0.01, se = 0.003, *p* < .001 in both samples) *(Figure 2c)*.

Table 4. Summary of results.

| | | Youth-Report Puberty | | | Parent-Report Puberty | | | Cognition | | |
		beta	SE	p	beta	SE	p	beta	SE	p
Study 1, Baseline	Sample 1	0.28	0.07	<.001	0.27	0.05	<.001	-0.01	0.003	<.001
	Sample 2	0.32	0.07	<.001	0.38	0.05	<.001	-0.01	0.003	<.001
Study 1, Follow-Up	Sample 1	0.50	0.05	<.001	0.57	0.05	<.001	--	--	--
	Sample 2	0.35	0.05	<.001	0.49	0.05	<.001	--	--	--

Study 2	--	0.07	0.04	<.001	0.12	0.03	<.001	0.001	0.002	<.001

Figure 2. Results - Previously validated model, baseline sample. a) Distributions of brain-age gaps, grouped by pubertal stage (youth-report). Means for each group are represented with vertical lines. b) Distributions of brain-age gaps, grouped by pubertal stage (parent-report). Means for each group are represented with vertical lines. c) Distributions of brain-age gaps, grouped by quartile of cognition scores, compared to others in the sample. Only the highest and lowest quartiles are shown. Means for each group are represented with vertical lines.

In both samples of the two-year follow-up data, a higher/more positive BrainAGE was related to more advanced youth- and parent-report pubertal development. For youth-report pubertal development, (b_{youth}= 0.50, *se* = 0.05; 0.35, *se* = 0.05, *p* < .001) *(Figure 3a)*. For parent-report pubertal development, (b_{parent}=0.57, *se* = 0.05; 0.49, *se* = 0.05, *p* < .001 in both samples)

(Figure 3b). As discussed above, composite cognition scores were not available at the two-year

follow-up.

Figure 3. Results - Previously validated model, follow-up sample. a) Distributions of brain-age gaps, grouped by pubertal stage (youth-report). Means for each group are represented with vertical lines. b) Distributions of brain-age gaps, grouped by pubertal stage (parent-report). Means for each group are represented with vertical lines.

Study 2

Participants

Samples and Models

<u>Model Training and Analyses</u>

       Adolescents included in the model training, testing, and analyses were participants in the

Adolescent Brain Cognitive Development (ABCD) Study, described in Study 1. Data included in

Study 2 were from the baseline sample (9-11 years old) and are identical to the baseline sample

described in Study 1.

<u>MRI Acquisition</u>

       Detailed imaging procedures can be found in Casey et al. (2018) and Hagler et al. (2019).

Harmonized protocols were used across 21 sites to collect imaging data using one of three 3T

scanner platforms (Phllips, Siemens, or GE) and a 32-channel head coil. 3-dimensional T1-weighted scans were collected with 1-mm voxel resolution.

Materials & Methods

MRI Processing and Model Features

The model was trained on and used to predict BrainAGE from the same 175 structural features used in the prediction for Study 1. A complete list of the features used in the training and prediction of Study 2 is available in *Supplementary Table 1*.

BrainAGE Prediction

For Study 2, available data were split into training, validation, and test sets. Half the sample (5,701 participants) were reserved for the test set. Of the remaining 5,701 participants, eighty percent ($N = 4,559$) were used for model training, and twenty percent ($N = 1,142$) were reserved for model validation. Model training followed the same procedures as the Study 1 model, originally described in Drobonin et al. (2022). Ten-fold cross-validation was repeated 10 times, stratified by scan age. Model training was performed using tidymodels (version 0.2.0; Kuhn & Wickham, 2020), with the XGBoost ML algorithm (version 1.5.2.1; Chen & Guestrin, 2016). Scan age was predicted using the cortical and subcortical features in Supplementary Table 1. The best model was the one with the smallest MAE (mean absolute error), which was expressed in years. Following model validation, the brain age gap was calculated in the hold-out sample by subtracting the actual scan age/chronological age from the predicted age. Descriptive statistics for BrainAGE predictions and maturational metrics are available in Table 3.

Results

BrainAGE Model Performance

Model 2

The best baseline model performed with a MAE of 0.14 in cross-validation. In our hold-out validation sample, the model performed with a MAE of 0.49. Finally, in the sample reserved for analyses, the model performed with a MAE of 0.49 (uncorrected for age bias) and 0.86 (corrected). Model performance is shown in *Figure 4*. Model performance was improved over existing models, including the model used in Study 1. The model showed a slight age bias, again due to the effect of regression to the mean. Corrected BrainAGE had a correlation of -0.20 with chronological age (Figure 4b).

Figure 4. Age versus model predictions for the novel model. a) Scan/chronological age versus model-predicted age for the baseline sample. The line represents an exact relationship between chronological age and predicted age. b) Scan/chronological age versus the age-corrected brain-age gap estimate (BrainAGE), for the baseline sample. The regression line shows a slight age-related bias, with younger participants predicted to be slightly older than their chronological age.

Neuroanatomical Contributions

The top contributors to the model included the brainstem volume, left frontal pole volume, and the posterior, mid-posterior, and anterior corpus callosum volume. *Figure 5* shows the cortical and subcortical features with the greatest importance to the model prediction.

A

B

Cortical Volume

C

Cortical Area

D

Figure 5. Neuroanatomical contributors to the novel model. A) Top 20 features ranked by importance to the model, across domains. B) Variable importance of cortical volume features, visualized on 2-dimensional brain. C) Variable importance of cortical area features, visualized on 2-dimensional brain. D) Variable importance of subcortical features, visualized on 2-dimensional brain. Left: Coronal view. Right: Sagittal view.

To contextualize our model features, we compared our findings to those of past adolescent BrainAGE models, including those of Drobonin et al. (2022) and Kelly et al. (2022). While BrainAGE model features cannot always be directly compared due to different brain parcellations and reporting methods, there were a number of high-contributing features in the novel model that are shared with those found in existing work. Shared with Drobonin et al. (2022), we found that the brainstem, posterior corpus callosum, rostral middle frontal volume, lateral ventricle, amygdala, and ventral diencephalon were high contributors to the model. Additionally, we compared our results with those of the model from Kelly et al. (2022) and found that both models shared volumes of frontal regions, the brainstem, and ventricles as high contributors.

BrainAGE and Maturational Metrics

A higher BrainAGE was related to more advanced youth- and parent-report pubertal development (Table 4). For youth-report pubertal development, (b_{youth} = 0.07, se = 0.04, $p <$

.001) *(Figure 6a)*. For parent-report pubertal development, (b_{parent} = 0.12, *se* = 0.03, *p* <.001)

(Figure 6b). In addition, a higher/more positive BrainAGE was related to higher cognition scores

(b_{cog} = 0.001, *se* = 0.002, *p* < .001) *(Figure 6c)*.

Figure 6. Results - Novel model. A) Distributions of brain-age gaps, grouped by pubertal stage (youth-report). Means for each group are represented with vertical lines. B) Distributions of brain-age gaps, grouped by pubertal stage (parent-report). Means for each group are represented with vertical lines. C) Distributions of brain-age gaps, grouped by quartile of cognition scores, compared to others in the sample. Only the highest and lowest quartiles are shown. Means for each group are represented with vertical lines.

Discussion

The current study used two BrainAGE models—one previously validated on a wide age

range across adolescence and one trained on a specific, narrow age range in early adolescence—

to quantify the relationship between BrainAGE and metrics of pubertal and cognitive

development. Using the previously validated model, we established that BrainAGE is positively

related to both youth- and parent-report pubertal development, and this relationship holds across

two age ranges (9-11, 10-13 years) and two subsamples within each of those age ranges (Table

3). This association was replicated using our novel model created from an independent sample of

9-11 year olds.

While the relationship between BrainAGE and NIH Toolbox Cognition scores was

consistently negatively related across subsamples of estimates derived from the validated model,

these results were in conflict with the positive relationship between BrainAGE and cognition

scores found in the estimates derived from the novel model. NIH Toolbox Cognition summary

scores were not calculated for the two-year follow-up data within ABCD Release 4.0, so it was

not possible to examine whether these relationships replicate across waves.

The results of the current study provide initial evidence that BrainAGE tracks with some

metrics of maturation, including pubertal development. However, the conflicting results between

BrainAGE and cognition lead us to question the utility of these models for non-biological

processes. The improvement in MAE between the age groups may be due to the fact that the

younger age group was on the lower end/extreme of the age range used to train the model, which

might have had fewer participants in the training set than in the older group, so the model would

be more accurate for the older age group. Potential sources of the improved MAE (mean absolute

error) could be the large amount of data, the narrow age range used in both training and testing (a

novel feature of the study), and the use of independent samples of data derived from the same

study (ABCD) for training, testing, and prediction in Study 2. In line with our findings, Ball et al.

(2017) suggested that image-based models predict most accurately during periods when the rate

of anatomical change is greatest. The current study provides a novel BrainAGE model trained on

a large sample of early adolescents, which can be used with developmental samples such as

ABCD. The novel model is available for use and can be found at

https://github.com/LucyWhitmore/BrainAGE-Maturation.

The current study used composite cognition scores which collapse across multiple

domains. To accurately assess the relationship between BrainAGE and cognition, we may need

to look at individual domains that show dramatic change in early adolescence, such as executive

function (Tervo-Clemmens et al., 2022, preprint). While the NIH Toolbox Cognition Battery is

generally sensitive to change over our age range of interest, specific domains may be more or

less strongly related to BrainAGE (Bauer & Zalazo, 2013).

While the NIH Toolbox is intended to be used over a lifespan age range, measures

targeted specifically at constructs that change in early adolescence may provide a clearer view of

how BrainAGE relates to cognition in this age range. Furthermore, the lack of complete

composite cognition scores at the two-year follow-up limits our ability to examine the

relationship between BrainAGE and cognition in multiple age bands. In future work, we may be

able to see whether the inconsistent relationship persists in older adolescents.

Additionally, a robust relationship was observed between BrainAGE and puberty across

subsamples, waves, models, and reporting methods. These results provide some evidence that

BrainAGE tracks with known metrics of pubertal maturation. The relationship between

BrainAGE and pubertal development strengthened between waves, possibly due to increased

representation of later pubertal stages in the older age range. In future work, these analyses can

be extended to track the relationship between BrainAGE and puberty within different age bands

in adolescence and illustrate when BrainAGE and puberty are most closely tied. Additionally,

youth- and parent-report pubertal development are known to vary in their reliability across age,

with parent-report being more reliable at younger ages, and youth-report being more reliable at

older ages (Terry et al., 2016). In the current study, we have shown that both reporting methods are consistently linked to BrainAGE. Future work may be able to illustrate when, or if, one method of reporting is no longer associated with BrainAGE.

To make matters more complicated, there are reciprocal connections between many of the constructs that have been related to BrainAGE. Here, we showed that BrainAGE is related to pubertal development, and past work has shown BrainAGE is related to psychopathology as well as adversity and, in adults, early life events such as low birth weight (Drobonin et al., 2022; Vidal-Pineiro et al., 2021). However, earlier pubertal timing has also been related to increased risk for psychopathology, and both psychopathology and early pubertal timing are related to environmental factors (Barendse et al., 2022; Graber et al., 1995; 2013; Mclaughlin et al., 2012). Therefore, it remains difficult to untangle these relationships and mechanisms.

Additionally, more longitudinal BrainAGE work is needed to fully characterize the relationship between BrainAGE and maturational metrics. The current study provides evidence that BrainAGE is positively correlated with pubertal development in early adolescence. However, it's still unknown how these relationships change over time, or if they remain consistent. Additionally, many of the features used to train BrainAGE models have huge variability and individual differences in adolescence, and brain maturity is more related to trajectories of brain development than overall values, which is not accounted for in most BrainAGE models (Mills et al., 2021). In current models, a participant may be considered younger or older because of the starting size of their brain, while they are in fact still following a normative developmental trajectory.

In summary, BrainAGE is a powerful tool that has promise as a clinical biomarker and as a tool to inform basic research. However, as BrainAGE gains traction within the field of

developmental cognitive neuroscience, we must ensure that we are careful about our interpretations of its predictions and associations and that we are confident in what we are actually measuring when we talk about brain maturity. As the results of brain maturity studies have implications for legal and other policy domains, it is vital that researchers provide contextualization for BrainAGE and be clear about what we can and cannot infer about different forms of maturity. BrainAGE is a powerful and promising tool, but we need to ensure that we have a strong theoretical background before widely incorporating it into studies of adolescent development and maturation.

Conclusions

The present study demonstrates that BrainAGE is positively correlated with pubertal development in a sample of early adolescents, and that this result is replicable across subsamples, models, and multiple age bands. However, the relationship between BrainAGE and cognition in early adolescence remains unclear. Future work should examine these relationships in later adolescence and utilize cognition measures that are optimized to examine changes across adolescence.

References

Anokhin, A. P., Luciana, M., Banich, M., Barch, D., Bjork, J. M., Gonzalez, M. R., Gonzalez, R., Haist, F., Jacobus, J., Lisdahl, K., McGlade, E., McCandliss, B., Nagel, B., Nixon, S. J., Tapert, S., Kennedy, J. T., & Thompson, W. (2022). Age-related changes and longitudinal stability of individual differences in ABCD Neurocognition measures. *Developmental Cognitive Neuroscience*, *54*, 101078. https://doi.org/10.1016/j.dcn.2022.101078

Aubert-Broche, B., Fonov, V. S., García-Lorenzo, D., Mouiha, A., Guizard, N., Coupé, P., Eskildsen, S. F., & Collins, D. L. (2013). A new method for structural volume analysis of longitudinal brain MRI data and its application in studying the growth trajectories of anatomical brain structures in childhood. *NeuroImage*, *82*, 393–402. https://doi.org/10.1016/j.neuroimage.2013.05.065

Ball, G., Adamson, C., Beare, R., & Seal, M. L. (2017). Modelling neuroanatomical variation during childhood and adolescence with neighbourhood-preserving embedding. *Scientific Reports*, *7*(1), Article 1. https://doi.org/10.1038/s41598-017-18253-6

Ball, G., Kelly, C. E., Beare, R., & Seal, M. L. (2021). Individual variation underlying brain age estimates in typical development. *NeuroImage*, *235*, 118036. https://doi.org/10.1016/j.neuroimage.2021.118036

Barch, D. M., Albaugh, M. D., Avenevoli, S., Chang, L., Clark, D. B., Glantz, M. D., Hudziak, J. J., Jernigan, T. L., Tapert, S. F., Yurgelun-Todd, D., Alia-Klein, N., Potter, A. S., Paulus, M. P., Prouty, D., Zucker, R. A., & Sher, K. J. (2018). Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale

and description. *Developmental Cognitive Neuroscience*, *32*, 55–66.

https://doi.org/10.1016/j.dcn.2017.10.010

Barendse, M. E. A., Byrne, M. L., Flournoy, J. C., McNeilly, E. A., Guazzelli Williamson, V.,

Barrett, A.-M. Y., Chavez, S. J., Shirtcliff, E. A., Allen, N. B., & Pfeifer, J. H. (2022).

Multimethod assessment of pubertal timing and associations with internalizing

psychopathology in early adolescent girls. *Journal of Psychopathology and Clinical*

Science, *131*(1), 14–25. https://doi.org/10.1037/abn0000721

Bauer, P. J., & Zelazo, P. D. (2013). Ix. Nih Toolbox Cognition Battery (cb): Summary,

Conclusions, and Implications for Cognitive Development. *Monographs of the Society*

for Research in Child Development, *78*(4), 133–146. https://doi.org/10.1111/mono.12039

Bethlehem, R. a. I., Seidlitz, J., White, S. R., Vogel, J. W., Anderson, K. M., Adamson, C.,

Adler, S., Alexopoulos, G. S., Anagnostou, E., Areces-Gonzalez, A., Astle, D. E.,

Auyeung, B., Ayub, M., Bae, J., Ball, G., Baron-Cohen, S., Beare, R., Bedford, S. A.,

Benegal, V., … Alexander-Bloch, A. F. (2022). Brain charts for the human lifespan.

Nature, *604*(7906), Article 7906. https://doi.org/10.1038/s41586-022-04554-y

Biondo, F., Jewell, A., Pritchard, M., Aarsland, D., Steves, C. J., Mueller, C., & Cole, J. H.

(2022). Brain-age is associated with progression to dementia in memory clinic patients.

NeuroImage: Clinical, *36*, 103175. https://doi.org/10.1016/j.nicl.2022.103175

Blakemore, S., Burnett, S., & Dahl, R. E. (2010). The role of puberty in the developing

adolescent brain. *Human Brain Mapping*, *31*(6), 926–933.

https://doi.org/10.1002/hbm.21052

Bleck, T. P., Nowinski, C. J., Gershon, R., & Koroshetz, W. J. (2013). What is the NIH Toolbox, and what will it mean to neurology? *Neurology*, *80*(10), 874–875. https://doi.org/10.1212/WNL.0b013e3182872ea0

Brouwer, R. M., Schutte, J., Janssen, R., Boomsma, D. I., Hulshoff Pol, H. E., & Schnack, H. G. (2021a). The Speed of Development of Adolescent Brain Age Depends on Sex and Is Genetically Determined. *Cerebral Cortex*, *31*(2), 1296–1306. https://doi.org/10.1093/cercor/bhaa296

Brouwer, R. M., Schutte, J., Janssen, R., Boomsma, D. I., Hulshoff Pol, H. E., & Schnack, H. G. (2021b). The Speed of Development of Adolescent Brain Age Depends on Sex and Is Genetically Determined. *Cerebral Cortex*, *31*(2), 1296–1306. https://doi.org/10.1093/cercor/bhaa296

Brown, T. T., Kuperman, J. M., Chung, Y., Erhart, M., McCabe, C., Hagler, D. J., Venkatraman, V. K., Akshoomoff, N., Amaral, D. G., Bloss, C. S., Casey, B. J., Chang, L., Ernst, T. M., Frazier, J. A., Gruen, J. R., Kaufmann, W. E., Kenet, T., Kennedy, D. N., Murray, S. S., … Dale, A. M. (2012). Neuroanatomical Assessment of Biological Maturity. *Current Biology*, *22*(18), 1693–1698. https://doi.org/10.1016/j.cub.2012.07.002

Bunge, S. A., & Whitaker, K. J. (2012). Brain Imaging: Your Brain Scan Doesn't Lie About Your Age. *Current Biology*, *22*(18), R800–R801. https://doi.org/10.1016/j.cub.2012.07.032

Callaghan, B. L., & Tottenham, N. (2016). The Stress Acceleration Hypothesis: Effects of early-life adversity on emotion circuits and behavior. *Current Opinion in Behavioral Sciences*, *7*, 76–81. https://doi.org/10.1016/j.cobeha.2015.11.018

Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M., Soules, M. E., Teslovich, T., Dellarco, D. V., Garavan, H., Orr, C. A., Wager, T. D., Banich, M. T., Speer, N. K., Sutherland, M. T., Riedel, M. C., Dick, A. S., Bjork, J. M., Thomas, K. M., … Dale, A. M. (2018). The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites. *Developmental Cognitive Neuroscience*, *32*, 43–54. https://doi.org/10.1016/j.dcn.2018.03.001

Chen, T., & Guestrin, C. (2016). XGBoost: A Scalable Tree Boosting System. *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, 785–794. https://doi.org/10.1145/2939672.2939785

Cheng, T. W., Magis-Weinberg, L., Guazzelli Williamson, V., Ladouceur, C. D., Whittle, S. L., Herting, M. M., Uban, K. A., Byrne, M. L., Barendse, M. E. A., Shirtcliff, E. A., & Pfeifer, J. H. (2021). A Researcher's Guide to the Measurement and Modeling of Puberty in the ABCD Study® at Baseline. *Frontiers in Endocrinology*, *12*. https://www.frontiersin.org/articles/10.3389/fendo.2021.608575

Chung, Y., Addington, J., Bearden, C. E., Cadenhead, K., Cornblatt, B., Mathalon, D. H., McGlashan, T., Perkins, D., Seidman, L. J., Tsuang, M., Walker, E., Woods, S. W., McEwen, S., van Erp, T. G. M., Cannon, T. D., & for the North American Prodrome Longitudinal Study (NAPLS) Consortium and the Pediatric Imaging, N., and Genetics (PING) Study Consortium. (2018). Use of Machine Learning to Determine Deviance in Neuroanatomical Maturity Associated With Future Psychosis in Youths at Clinically High Risk. *JAMA Psychiatry*, *75*(9), 960–968. https://doi.org/10.1001/jamapsychiatry.2018.1543

Cromer, J. A., Schembri, A. J., Harel, B. T., & Maruff, P. (2015). The nature and rate of

cognitive maturation from late childhood to adulthood. *Frontiers in Psychology*, *6*.

https://www.frontiersin.org/articles/10.3389/fpsyg.2015.00704

Cropley, V. L., Tian, Y., Fernando, K., Mansour L., S., Pantelis, C., Cocchi, L., & Zalesky, A.

(2021). Brain-Predicted Age Associates With Psychopathology Dimensions in Youths.

Biological Psychiatry: Cognitive Neuroscience and Neuroimaging, *6*(4), 410–419.

https://doi.org/10.1016/j.bpsc.2020.07.014

Davidson, M. C., Amso, D., Anderson, L. C., & Diamond, A. (2006). Development of cognitive

control and executive functions from 4 to 13 years: Evidence from manipulations of

memory, inhibition, and task switching. *Neuropsychologia*, *44*(11), 2037–2078.

https://doi.org/10.1016/j.neuropsychologia.2006.02.006

*Developmental brain changes during puberty and associations with mental health problems |

Elsevier Enhanced Reader*. (n.d.). https://doi.org/10.1016/j.dcn.2023.101227

Dorn, L. D., & Biro, F. M. (2011). Puberty and Its Measurement: A Decade in Review. *Journal

of Research on Adolescence*, *21*(1), 180–195. https://doi.org/10.1111/j.1532-

7795.2010.00722.x

Dorn, L. D., Dahl, R. E., Woodward, H. R., & Biro, F. (2006). Defining the Boundaries of Early

Adolescence: A User's Guide to Assessing Pubertal Status and Pubertal Timing in

Research With Adolescents. *Applied Developmental Science*, *10*(1), 30–56.

https://doi.org/10.1207/s1532480xads1001_3

Dosenbach, N. U. F., Nardos, B., Cohen, A. L., Fair, D. A., Power, J. D., Church, J. A., Nelson,

S. M., Wig, G. S., Vogel, A. C., Lessov-Schlaggar, C. N., Barnes, K. A., Dubis, J. W.,

Feczko, E., Coalson, R. S., Pruett, J. R., Barch, D. M., Petersen, S. E., & Schlaggar, B. L.

(2010). Prediction of Individual Brain Maturity Using fMRI. *Science (New York, N.Y.)*, *329*(5997), 1358–1361. https://doi.org/10.1126/science.1194144

Drobinin, V., Gestel, H. V., Helmick, C. A., Schmidt, M. H., Bowen, C. V., & Uher, R. (2022). The Developmental Brain Age Is Associated With Adversity, Depression, and Functional Outcomes Among Adolescents. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, *7*(4), 406–414. https://doi.org/10.1016/j.bpsc.2021.09.004

Erus, G., Battapady, H., Satterthwaite, T. D., Hakonarson, H., Gur, R. E., Davatzikos, C., & Gur, R. C. (2015). Imaging Patterns of Brain Development and their Relationship to Cognition. *Cerebral Cortex*, *25*(6), 1676–1684. https://doi.org/10.1093/cercor/bht425

Franke, K., & Gaser, C. (2012). Longitudinal changes in individual BrainAGE in healthy aging, mild cognitive impairment, and Alzheimer's disease. *GeroPsych: The Journal of Gerontopsychology and Geriatric Psychiatry*, *25*, 235–245. https://doi.org/10.1024/1662-9647/a000074

Franke, K., & Gaser, C. (2019). Ten Years of BrainAGE as a Neuroimaging Biomarker of Brain Aging: What Insights Have We Gained? *Frontiers in Neurology*, *10*. https://www.frontiersin.org/articles/10.3389/fneur.2019.00789

Franke, K., Hagemann, G., Schleussner, E., & Gaser, C. (2015). Changes of individual BrainAGE during the course of the menstrual cycle. *NeuroImage*, *115*, 1–6. https://doi.org/10.1016/j.neuroimage.2015.04.036

Franke, K., Luders, E., May, A., Wilke, M., & Gaser, C. (2012). Brain maturation: Predicting individual BrainAGE in children and adolescents using structural MRI. *NeuroImage*, *63*(3), 1305–1312. https://doi.org/10.1016/j.neuroimage.2012.08.001

Franke, K., Ziegler, G., Klöppel, S., & Gaser, C. (2010). Estimating the age of healthy subjects from T1-weighted MRI scans using kernel methods: Exploring the influence of various parameters. *NeuroImage*, *50*(3), 883–892. https://doi.org/10.1016/j.neuroimage.2010.01.005

Fuhrmann, D., Madsen, K. S., Johansen, L. B., Baaré, W. F. C., & Kievit, R. A. (2022). The midpoint of cortical thinning between late childhood and early adulthood differs between individuals and brain regions: Evidence from longitudinal modelling in a 12-wave neuroimaging sample. *NeuroImage*, *261*, 119507. https://doi.org/10.1016/j.neuroimage.2022.119507

Garavan, H., Bartsch, H., Conway, K., Decastro, A., Goldstein, R. Z., Heeringa, S., Jernigan, T., Potter, A., Thompson, W., & Zahs, D. (2018). Recruiting the ABCD sample: Design considerations and procedures. *Developmental Cognitive Neuroscience*, *32*, 16–22. https://doi.org/10.1016/j.dcn.2018.04.004

Gershon, R. C., Wagster, M. V., Hendrie, H. C., Fox, N. A., Cook, K. F., & Nowinski, C. J. (2013). NIH Toolbox for Assessment of Neurological and Behavioral Function. *Neurology*, *80*(11 Supplement 3), S2–S6. https://doi.org/10.1212/WNL.0b013e3182872e5f

Goddings, A.-L., Mills, K. L., Clasen, L. S., Giedd, J. N., Viner, R. M., & Blakemore, S.-J. (2014). The influence of puberty on subcortical brain development. *NeuroImage*, *88*, 242–251. https://doi.org/10.1016/j.neuroimage.2013.09.073

Graber, J. A. (2013). Pubertal timing and the development of psychopathology in adolescence and beyond. *Hormones and Behavior*, *64*(2), 262–269. https://doi.org/10.1016/j.yhbeh.2013.04.003

Graber, J. A., Brooks-Gunn, J., & Warren, M. P. (1995). The Antecedents of Menarcheal Age:

Heredity, Family Environment, and Stressful Life Events. *Child Development*, *66*(2),

346–359. https://doi.org/10.2307/1131582

Gur, R. C., Richard, J., Calkins, M. E., Chiavacci, R., Hansen, J. A., Bilker, W. B., Loughead, J.,

Connolly, J. J., Qiu, H., Mentch, F. D., Abou-Sleiman, P. M., Hakonarson, H., & Gur, R.

E. (2012). Age group and sex differences in performance on a computerized

neurocognitive battery in children age 8–21. *Neuropsychology*, *26*(2), 251–265.

https://doi.org/10.1037/a0026712

Hagler, D. J., Hatton, S. N., Cornejo, M. D., Makowski, C., Fair, D. A., Dick, A. S., Sutherland,

M. T., Casey, B., Barch, D. M., Harms, M. P., Watts, R., Bjork, J. M., Garavan, H. P.,

Hilmer, L., Pung, C. J., Sicat, C. S., Kuperman, J., Bartsch, H., Xue, F., … Dale, A. M.

(2019). Image processing and analysis methods for the Adolescent Brain Cognitive

Development Study. *NeuroImage*, *202*, 116091.

https://doi.org/10.1016/j.neuroimage.2019.116091

Han, L. K. M., Dinga, R., Hahn, T., Ching, C. R. K., Eyler, L. T., Aftanas, L., Aghajani, M.,

Aleman, A., Baune, B. T., Berger, K., Brak, I., Filho, G. B., Carballedo, A., Connolly, C.

G., Couvy-Duchesne, B., Cullen, K. R., Dannlowski, U., Davey, C. G., Dima, D., …

Schmaal, L. (2021). Brain aging in major depressive disorder: Results from the ENIGMA

major depressive disorder working group. *Molecular Psychiatry*, *26*(9), Article 9.

https://doi.org/10.1038/s41380-020-0754-0

Han, L. K. M., Dinga, R., Leenings, R., Hahn, T., Cole, J. H., Aftanas, L. I., Amod, A. R.,

Besteher, B., Colle, R., Corruble, E., Couvy-Duchesne, B., Danilenko, K. V., Fuentes-

Claramonte, P., Gonul, A. S., Gotlib, I. H., Goya-Maldonado, R., Groenewold, N. A.,

Hamilton, P., Ichikawa, N., … Schmaal, L. (2022). A large-scale ENIGMA multisite

replication study of brain age in depression. *Neuroimage: Reports*, *2*(4), 100149.

https://doi.org/10.1016/j.ynirp.2022.100149

Havelock, J. C., Auchus, R. J., & Rainey, W. E. (2004). The Rise in Adrenal Androgen

Biosynthesis: Adrenarche. *Seminars in Reproductive Medicine*, *22*(04), 337–347.

https://doi.org/10.1055/s-2004-861550

Herting, M. M., Gautam, P., Spielberg, J. M., Dahl, R. E., & Sowell, E. R. (2015). A

Longitudinal Study: Changes in Cortical Thickness and Surface Area during Pubertal

Maturation. *PLOS ONE*, *10*(3), e0119774. https://doi.org/10.1371/journal.pone.0119774

Herting, M. M., & Sowell, E. R. (2017). Puberty and structural brain development in humans.

Frontiers in Neuroendocrinology, *44*, 122–137.

https://doi.org/10.1016/j.yfrne.2016.12.003

Hodes, R. J., Insel, T. R., Landis, S. C., & Research, O. behalf of the N. B. for N. (2013). The

NIH Toolbox: Setting a standard for biomedical research. *Neurology*, *80*(11 Supplement

3), S1–S1. https://doi.org/10.1212/WNL.0b013e3182872e90

Holm, M. C., Leonardsen, E. H., Beck, D., Dahl, A., Kjelkenes, R., de Lange, A.-M. G., &

Westlye, L. T. (2022). *Linking brain maturation and puberty during early adolescence

using longitudinal brain age prediction in the ABCD cohort* [Preprint]. Psychiatry and

Clinical Psychology. https://doi.org/10.1101/2022.05.16.22275146

Holm, M. C., Leonardsen, E. H., Beck, D., Dahl, A., Kjelkenes, R., de Lange, A.-M. G., &

Westlye, L. T. (2023). Linking brain maturation and puberty during early adolescence

using longitudinal brain age prediction in the ABCD cohort. *Developmental Cognitive

Neuroscience*, *60*, 101220. https://doi.org/10.1016/j.dcn.2023.101220

Holm, M. C., Leonardsen, E. H., Beck, D., Dahl, A., Kjelkenes, R., Lange, A.-M. G. de, & Westlye, L. T. (2022). *Linking brain maturation and puberty during early adolescence using longitudinal brain age prediction in the ABCD cohort* (p. 2022.05.16.22275146). medRxiv. https://doi.org/10.1101/2022.05.16.22275146

Kelly, C., Ball, G., Matthews, L. G., Cheong, J. L., Doyle, L. W., Inder, T. E., Thompson, D. K., & Anderson, P. J. (2022). Investigating brain structural maturation in children and adolescents born very preterm using the brain age framework. *NeuroImage*, *247*, 118828. https://doi.org/10.1016/j.neuroimage.2021.118828

Khundrakpam, B. S., Tohka, J., & Evans, A. C. (2015). Prediction of brain maturity based on cortical thickness at different spatial resolutions. *NeuroImage*, *111*, 350–359. https://doi.org/10.1016/j.neuroimage.2015.02.046

Kuhn, M., & Wickham, H. (2020). *Tidymodels: A collection of packages for modeling and machine learning using tidyverse principles.* https://www.tidymodels.org

Le, T. T., Kuplicki, R. T., McKinney, B. A., Yeh, H.-W., Thompson, W. K., Paulus, M. P., Tulsa 1000 Investigators, Aupperle, R. L., Bodurka, J., Cha, Y.-H., Feinstein, J. S., Khalsa, S. S., Savitz, J., Simmons, W. K., & Victor, T. A. (2018). A Nonlinear Simulation Framework Supports Adjusting for Age When Analyzing BrainAGE. *Frontiers in Aging Neuroscience*, *10*. https://www.frontiersin.org/articles/10.3389/fnagi.2018.00317

Lewis, J. D., Evans, A. C., & Tohka, J. (2018). T1 white/gray contrast as a predictor of chronological age, and an index of cognitive performance. *NeuroImage*, *173*, 341–350. https://doi.org/10.1016/j.neuroimage.2018.02.050

Luciana, M., Bjork, J. M., Nagel, B. J., Barch, D. M., Gonzalez, R., Nixon, S. J., & Banich, M. T. (2018). Adolescent neurocognitive development and impacts of substance use:

Overview of the adolescent brain cognitive development (ABCD) baseline

neurocognition battery. *Developmental Cognitive Neuroscience*, *32*, 67–79.

https://doi.org/10.1016/j.dcn.2018.02.006

McLaughlin, K. A., Greif Green, J., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler,

R. C. (2012). Childhood Adversities and First Onset of Psychiatric Disorders in a

National Sample of US Adolescents. *Archives of General Psychiatry*, *69*(11), 1151–1160.

https://doi.org/10.1001/archgenpsychiatry.2011.2277

Mills, K. L., Goddings, A.-L., Herting, M. M., Meuwese, R., Blakemore, S.-J., Crone, E. A.,

Dahl, R. E., Güroğlu, B., Raznahan, A., Sowell, E. R., & Tamnes, C. K. (2016).

Structural brain development between childhood and adulthood: Convergence across four

longitudinal samples. *NeuroImage*, *141*, 273–281.

https://doi.org/10.1016/j.neuroimage.2016.07.044

Mills, K. L., Siegmund, K. D., Tamnes, C. K., Ferschmann, L., Wierenga, L. M., Bos, M. G. N.,

Luna, B., Li, C., & Herting, M. M. (2021). Inter-individual variability in structural brain

development from late childhood to young adulthood. *NeuroImage*, *242*, 118450.

https://doi.org/10.1016/j.neuroimage.2021.118450

Nguyen, T.-V., McCracken, J., Ducharme, S., Botteron, K. N., Mahabir, M., Johnson, W., Israel,

M., Evans, A. C., Karama, S., & for the Brain Development Cooperative Group. (2013).

Testosterone-Related Cortical Maturation Across Childhood and Adolescence. *Cerebral

Cortex*, *23*(6), 1424–1432. https://doi.org/10.1093/cercor/bhs125

Peng, H., Gong, W., Beckmann, C. F., Vedaldi, A., & Smith, S. M. (2021). Accurate brain age

prediction with lightweight deep neural networks. *Medical Image Analysis*, *68*, 101871.

https://doi.org/10.1016/j.media.2020.101871

Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of

pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*,

17(2), 117–133. https://doi.org/10.1007/BF01537962

Pfeifer, J. H., & Allen, N. B. (2021). Puberty Initiates Cascading Relationships Between

Neurodevelopmental, Social, and Internalizing Processes Across Adolescence. *Biological*

Psychiatry, *89*(2), 99–108. https://doi.org/10.1016/j.biopsych.2020.09.002

Potter, A. S., Dube, S. L., Barrios, L. C., Bookheimer, S., Espinoza, A., Feldstein Ewing, S. W.,

Freedman, E. G., Hoffman, E. A., Ivanova, M., Jefferys, H., McGlade, E. C., Tapert, S.

F., & Johns, M. M. (2022). Measurement of gender and sexuality in the Adolescent Brain

Cognitive Development (ABCD) study. *Developmental Cognitive Neuroscience*, *53*,

101057. https://doi.org/10.1016/j.dcn.2022.101057

Raznahan, A., Shaw, P., Lalonde, F., Stockman, M., Wallace, G. L., Greenstein, D., Clasen, L.,

Gogtay, N., & Giedd, J. N. (2011). How Does Your Cortex Grow? *Journal of*

Neuroscience, *31*(19), 7174–7177. https://doi.org/10.1523/JNEUROSCI.0054-11.2011

Saragosa-Harris, N. M., Chaku, N., MacSweeney, N., Guazzelli Williamson, V., Scheuplein, M.,

Feola, B., Cardenas-Iniguez, C., Demir-Lira, E., McNeilly, E. A., Huffman, L. G.,

Whitmore, L., Michalska, K. J., Damme, K. S., Rakesh, D., & Mills, K. L. (2022). A

practical guide for researchers and reviewers using the ABCD Study and other large

longitudinal datasets. *Developmental Cognitive Neuroscience*, *55*, 101115.

https://doi.org/10.1016/j.dcn.2022.101115

Schnack, H. G., van Haren, N. E. M., Nieuwenhuis, M., Hulshoff Pol, H. E., Cahn, W., & Kahn,

R. S. (2016). Accelerated Brain Aging in Schizophrenia: A Longitudinal Pattern

Recognition Study. *The American Journal of Psychiatry*, *173*(6), 607–616.

https://doi.org/10.1176/appi.ajp.2015.15070922

Segalowitz, S. J., & Rose-Krasnor, L. (1992). The construct of brain maturation in theories of

child development. *Brain and Cognition*, *20*(1), 1–7. https://doi.org/10.1016/0278-

2626(92)90058-T

Shaw, P., Gogtay, N., & Rapoport, J. (2010). Childhood psychiatric disorders as anomalies in

neurodevelopmental trajectories. *Human Brain Mapping*, *31*(6), 917–925.

https://doi.org/10.1002/hbm.21028

Sisk, C. L., & Foster, D. L. (2004). The neural basis of puberty and adolescence. *Nature

Neuroscience*, *7*(10), Article 10. https://doi.org/10.1038/nn1326

Smith, S. M., Vidaurre, D., Alfaro-Almagro, F., Nichols, T. E., & Miller, K. L. (2019).

Estimation of brain age delta from brain imaging. *NeuroImage*, *200*, 528–539.

https://doi.org/10.1016/j.neuroimage.2019.06.017

Somerville, L. H. (2016). Searching for Signatures of Brain Maturity: What Are We Searching

For? *Neuron*, *92*(6), 1164–1167. https://doi.org/10.1016/j.neuron.2016.10.059

Steinbeis, N., & Crone, E. A. (2016). The link between cognitive control and decision-making

across child and adolescent development. *Current Opinion in Behavioral Sciences*, *10*,

28–32. https://doi.org/10.1016/j.cobeha.2016.04.009

Tamnes, C. K., Herting, M. M., Goddings, A.-L., Meuwese, R., Blakemore, S.-J., Dahl, R. E.,

Güroğlu, B., Raznahan, A., Sowell, E. R., Crone, E. A., & Mills, K. L. (2017).

Development of the Cerebral Cortex across Adolescence: A Multisample Study of Inter-

Related Longitudinal Changes in Cortical Volume, Surface Area, and Thickness. *The*

Journal of Neuroscience, *37*(12), 3402–3412. https://doi.org/10.1523/JNEUROSCI.3302-16.2017

Tamnes, C. K., Walhovd, K. B., Grydeland, H., Holland, D., Østby, Y., Dale, A. M., & Fjell, A. M. (2013). Longitudinal Working Memory Development Is Related to Structural Maturation of Frontal and Parietal Cortices. *Journal of Cognitive Neuroscience*, *25*(10), 1611–1623. https://doi.org/10.1162/jocn_a_00434

Tanner, J. M. (1962). *Growth at adolescence, 2nd ed*. Springfield, Ill.

Terry, M. B., Goldberg, M., Schechter, S., Houghton, L. C., White, M. L., O'Toole, K., Chung, W. K., Daly, M. B., Keegan, T. H. M., Andrulis, I. L., Bradbury, A. R., Schwartz, L., Knight, J. A., John, E. M., & Buys, S. S. (2016). Comparison of Clinical, Maternal, and Self Pubertal Assessments: Implications for Health Studies. *Pediatrics*, *138*(1), e20154571. https://doi.org/10.1542/peds.2015-4571

Tervo-Clemmens, B., Calabro, F. J., Parr, A. C., Fedor, J., Foran, W., & Luna, B. (2022). *A Canonical Trajectory of Executive Function Maturation During the Transition from Adolescence to Adulthood*. PsyArXiv. https://doi.org/10.31234/osf.io/73yfv

Thompson, W. K., Barch, D. M., Bjork, J. M., Gonzalez, R., Nagel, B. J., Nixon, S. J., & Luciana, M. (2018). The structure of cognition in 9 and 10 year-old children and associations with problem behaviors: Findings from the ABCD study's baseline neurocognitive battery. *Developmental Cognitive Neuroscience*, *36*, 100606. https://doi.org/10.1016/j.dcn.2018.12.004

Veldhuis, J. D. (1996). Neuroendocrine mechanisms mediating awakening of the human gonadotropic axis in puberty. *Pediatric Nephrology (Berlin, Germany)*, *10*(3), 304–317. https://doi.org/10.1007/BF00866767

Vidal-Pineiro, D., Wang, Y., Krogsrud, S. K., Amlien, I. K., Baaré, W. F., Bartres-Faz, D.,

  Bertram, L., Brandmaier, A. M., Drevon, C. A., Düzel, S., Ebmeier, K., Henson, R. N.,

  Junqué, C., Kievit, R. A., Kühn, S., Leonardsen, E., Lindenberger, U., Madsen, K. S.,

  Magnussen, F., … Fjell, A. (2021). Individual variations in 'brain age' relate to early-life

  factors more than to longitudinal brain change. *ELife*, *10*, e69995.

  https://doi.org/10.7554/eLife.69995

Vijayakumar, N., Op de Macks, Z., Shirtcliff, E. A., & Pfeifer, J. H. (2018). Puberty and the

  human brain: Insights into adolescent development. *Neuroscience & Biobehavioral*

  Reviews, *92*, 417–436. https://doi.org/10.1016/j.neubiorev.2018.06.004

Vijayakumar, N., Youssef, G. J., Allen, N. B., Anderson, V., Efron, D., Hazell, P., Mundy, L.,

  Nicholson, J. M., Patton, G., Seal, M. L., Simmons, J. G., Whittle, S., & Silk, T. (2021).

  A longitudinal analysis of puberty-related cortical development. *NeuroImage*, *228*,

  117684. https://doi.org/10.1016/j.neuroimage.2020.117684

Wierenga, L., Langen, M., Ambrosino, S., van Dijk, S., Oranje, B., & Durston, S. (2014).

  Typical development of basal ganglia, hippocampus, amygdala and cerebellum from age

  7 to 24. *NeuroImage*, *96*, 67–72. https://doi.org/10.1016/j.neuroimage.2014.03.072

Wierenga, L. M., Bos, M. G. N., Schreuders, E., Vd Kamp, F., Peper, J. S., Tamnes, C. K., &

  Crone, E. A. (2018). Unraveling age, puberty and testosterone effects on subcortical brain

  development across adolescence. *Psychoneuroendocrinology*, *91*, 105–114.

  https://doi.org/10.1016/j.psyneuen.2018.02.034

Contributions

Contributed to conception and design: LBW, KLM
Contributed to acquisition of data: ABCD
Contributed to analysis and interpretation of data: LBW, KLM, SJW
Drafted and/or revised the article: LBW, KLM, SJW
Approved the submitted version for publication: LBW, KLM, SJW

Acknowledgements

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children aged 9-10 and follow them over 10 years into early adulthood. The ABCD Study is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123, U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators. The ABCD data repository grows and changes over time. The ABCD data used in this report came from https://dx.doi.org/10.15154/1528926.

We thank Dr. Vlad Drobonin for making his scripts and validated BrainAGE model available for use.

Funding

This study was funded by University of Oregon College of Arts and Sciences. Author KLM was supported by the Research Council of Norway (RCN) grant number 288083.

Competing Interests
No competing interests exist.

Data Availability Statement
Analysis scripts can be found at: https://github.com/LucyWhitmore/BrainAGE-Maturation
The publicly available existing model, as well as training and testing scripts, can be found at: https://github.com/GitDro/DevelopmentalBrainAge. Data are part of the ABCD 4.0 Data Release, and are available with an NIH Data Use Certification. The 4.0 release can be found at: http://dx.doi.org/10.15154/1523041

Supplementary Materials

Supplementary Table 1. Brain features

	Volume	Area	Model 1 Training (Not Included in Prediction)
Cortical	caudal anterior cingulate caudal middle frontal cuneus entorhinal fusiform inferior parietal inferior temporal isthmus cingulate lateral occipital lateral orbitofrontal lingual medial orbitofrontal middle temporal parahippocampal paracentral pars opercularis pars orbitalis pars triangularis pericalcarine postcentral posterior cingulate precentral precuneus rostral anterior cingulate rostral middle frontal superior frontal superior parietal superior temporal supramarginal transverse temporal left insula left bankssts frontal pole temporal pole Total GM	caudal anterior cingulate caudal middle frontal cuneus entorhinal fusiform inferior parietal inferior temporal isthmus cingulate lateral occipital lateral orbitofrontal lingual medial orbitofrontal middle temporal parahippocampal paracentral pars opercularis pars orbitalis pars triangularis pericalcarine postcentral posterior cingulate precentral precuneus rostral anterior cingulate rostral middle frontal superior frontal superior parietal superior temporal supramarginal transverse temporal left insula left bankssts frontal pole temporal pole	BrainSeg_Vol BrainSeg_Vol_No_Vent BrainSeg_Vol_No_Vent_Surf Total GM SupraTentorial_Vol_No_Vent SupraTentorial_No_Vent_Voxel_Count Mask BrainSegVol_eTIV_Ratio MaskVol_eTIV_Ratio Vessel Choroid Plexus Optic Chiasm
Subcortical	Thalamus Caudate		

	Putamen Pallidum Lateral Ventricle Inferior Lateral Ventricle Hippocampus Amygdala Accumbens Left Ventral DC 3rd Ventricle 4th Ventricle BrainStem Posterior CC MidPosterior Central CC MidAnterior CC Anterior CC Total GM		
Other	Cerebellum WM Cerebellum Intercranial CSF SupraTentorial		

Supplementary Table 2. Reliability Measures

	Youth-Report PDS		Parent-Report PDS	
	Female	Male	Female	Male
Baseline	.60	.53	.70	.64
Follow-Up	.76	.73	.81	.79

Measure reliability, Cronbach's a as calculated in psych::omega()