Albumin-binding RNAi Conjugate for Carrier Free Treatment of Arthritis

Juan M. Colazo[1], Ella N. Hoogenboezem[1], Veeraj Shah[1], Fang Yu[1], Justin H. Lo[1], Alexander G. Sorets[1], Nora Francini[1], Hongsik Cho[2], Danielle L. Michell[3], Kasey C. Vickers[3], Katherine N. Gibson-Corley[4], Karen A. Hasty[2], Leslie Crofford[5], Rebecca S. Cook[1,6], and Craig L. Duvall[1*]

[1]Department of Biomedical Engineering, Vanderbilt University, Nashville, TN, USA.
[2]Department of Orthopaedic Surgery and Biomedical Engineering, UTHSC, Memphis VA Medical Center, Memphis, TN, USA.
[3]Department of Medicine, Division of Cardiovascular Medicine, Vanderbilt University Medical Center, Nashville, TN, USA.
[4]Department of Pathology, Microbiology, and Immunology, Division of Comparative Medicine, Vanderbilt University, Nashville, TN, USA.
[5]Department of Medicine, Division of Rheumatology, Vanderbilt University, Nashville, TN, USA.
[6]Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, TN, USA.

*Corresponding Author:
Craig L. Duvall, Ph.D.
Cornelius Vanderbilt Professor of Biomedical Engineering
Vanderbilt University
email: craig.duvall@vanderbilt.edu
mailing address: 5824 Stevenson Center, Nashville, TN 37232

Keywords:
Albumin, siRNA, siRNA conjugates, lipid conjugates, matrix metalloproteinase (MMP), MMP13, siRNA, RNAi, osteoarthritis, multi-joint osteoarthritis, K/BxN, rheumatoid arthritis, extra-hepatic delivery, pharmacokinetics, pharmacodynamics, DMOAD, DMARD

Editorial Summary: Lipophilic siRNA conjugates optimized for albumin binding and "hitchhiking" can be leveraged to achieve preferential delivery to and gene silencing activity within arthritic joints. Chemical stabilization of the lipophilic siRNA enables intravenous siRNA delivery without lipid or polymer encapsulation. Using siRNA sequences targeting MMP13, a key driver of arthritis-related inflammation, albumin hitchhiking siRNA diminished MMP13, inflammation, and manifestations of osteoarthritis and rheumatoid arthritis at molecular, histological, and clinical levels, consistently outperforming clinical standards of care and small molecule MMP antagonists.

Running Title: RNAi conjugate for arthritis

Abstract

Osteoarthritis (**OA**) and rheumatoid arthritis (**RA**) are joint diseases that are associated with pain and lost quality of life. No disease modifying OA drugs are currently available. RA treatments are better established but are not always effective and can cause immune suppression. Here, an MMP13-selective siRNA conjugate was developed that, when delivered intravenously, docks onto endogenous albumin and promotes preferential accumulation in articular cartilage and synovia of OA and RA joints. MMP13 expression was diminished upon intravenous delivery of MMP13 siRNA conjugates, consequently decreasing multiple histological and molecular markers of disease severity, while also reducing clinical manifestations such as swelling (RA) and joint pressure sensitivity (RA and OA). Importantly, MMP13 silencing provided more comprehensive OA treatment efficacy than standard of care (steroids) or experimental MMP inhibitors. These data demonstrate the utility of albumin 'hitchhiking' for drug delivery to arthritic joints, and establish the therapeutic utility of systemically delivered anti-MMP13 siRNA conjugates in OA and RA.

Introduction

Osteoarthritis (**OA**) is a degenerative joint disease that can be idiopathic or secondary to injury, in the case of post-traumatic osteoarthritis (**PTOA**). OA is characterized by cartilage loss, synovial inflammation, and formation of osteophytes, all of which contribute to pain and lost joint function[1]. Patients often develop broader multi-joint osteoarthritis (**MJOA**), which can have debilitating impact on quality of life[2]. The autoimmune disease rheumatoid arthritis (**RA**) has independent etiology and is characterized by severe synovial and systemic inflammation, bone erosion, and cartilage damage[3,4]. Current OA treatment options are palliative and include non-steroidal anti-inflammatory drugs (**NSAIDS**) or corticosteroids, approaches that do not counteract the root causes of disease nor prevent OA progression; corticosteroids may in fact worsen cartilage thinning[5-7]. Current treatments for RA include synthetic disease-modifying antirheumatic drugs (**DMARD**s) and biologic DMARDs such as tumor necrosis factor (**TNF**) inhibitors, interleukin (**IL**) inhibitors, and B-cell inhibitors. These treatments are efficacious, although they can suppress the immune system, and their benefits often wane over the long-term[8].

There is extensive overlap between OA and RA in terms of the downstream molecular effectors that drive joint degeneration[9]. Both are associated with inflammation and upregulation of extracellular matrix (**ECM**) degrading matrix metalloproteinases (MMPs)[10]. MMPs erode cartilage, and the degradation byproducts have inflammatory signaling properties that further induce expression of inflammatory cytokines and MMPs, perpetuating a chronic, catabolic cycle that leads to deep cartilage erosion, chondrocyte loss, and subchondral bone exposure[11-13].

Therapeutic MMP inhibition is a logical approach for OA and RA treatment. Development of MMP-selective antagonists is critical, as broad spectrum MMP inhibitors cause musculoskeletal toxicities in humans, likely due to universal disruption of normal tissue homeostasis[14-16]. Among MMPs, MMP13 is perhaps the most efficient at cleaving the key ECM components, collagen II[17] and fibronectin[18], producing inflammatory fragments. Further, knockout mouse studies pinpoint MMP13 as a key proteolytic driver of cartilage loss in arthritis models[19]. Lastly, MMP13 expression is relatively confined to joint tissues and diseased or injured tissues characterized by fibrosis or inflammation[20]. This narrow expression pattern allowing for a potentially wider therapeutic window with minimal toxicities. However, this hypothesis has yet to be tested fully, as clinical development of selective MMP inhibitors has been challenged by the extensive structural similarities between the catalytic sites of the various MMPs[21].

RNA interference (RNAi) provides an ideal technique for gene-selective inhibition, as specificity is conferred by the genetic sequence. The translatability of siRNA therapeutics is also now well-established based on the FDA approval of lipid nanoparticle and GalNAc-siRNA conjugate drugs, both of which have been successfully applied in humans for gene silencing in the liver[22,23]. Selective MMP13 gene knockdown using siRNA sequences would overcome the hurdle of structural similarity between MMPs, thus providing an opportunity to target MMP13 in the context of OA and RA. This notion is supported by recent PTOA model studies in which intra-articular delivery of MMP13 silencing siRNA nanoparticles (siNPs) protected against knee joint load-induced cartilage loss[24,25].

To improve upon these relatively-complex technologies for sustained MMP13 siRNA delivery via intra-articular injection of siNPs, we sought an approach in which carrier-free siRNA conjugates delivered intravenously might preferentially accumulate within arthritic joints. A systemically delivered approach avoids the potential joint damage that can occur from repeated intra-articular injections and better enables treatment of MJOA and RA. Relative to siNPs, a molecularly-defined conjugate is also simpler to manufacture and does not come with the added risk of carrier toxicities that often narrow the therapeutic window of cationic lipid or polymer siNP formulations.

Development of carrier-free siRNA conjugates for rheumatic diseases is an unchartered but high-impact endeavor. Toward this goal, experimental siRNA-lipid conjugates that commandeer circulating plasma albumin for delivery to arthritic joints were engineered and tested in models of OA and RA. Albumin has unique promise as a therapeutic siRNA carrier, given its long circulation half-life (20 days)[26] and natural

tendency to accumulate in inflamed tissues, including arthritic joints[27-30]. Although covalent conjugation of siRNA to albumin has been explored to some extent[31,32], lipid-siRNA conjugates that enable reversible, in situ association with native circulating albumin has been proof-of-concept tested only in cancer studies. We show here that systemic delivery of an albumin-hitchhiking siRNA-lipid conjugate targeting MMP13 achieves robust delivery, MMP13 silencing, and therapeutic efficacy in joints afflicted by OA and RA.

Results and discussion

Circulating albumin robustly accumulates in PTOA joints.

To determine the extent to which circulating albumin accumulates in arthritic joints, we intravenously (i.v.) delivered Cy5-labeled mouse serum albumin (MSA-Cy5) into mice with induced PTOA. In this model, the left knee undergoes repeated loading to induce damage and inflammation, while the right knee remains unchallenged (**Figure 1A**). At 24 hrs after delivery, abundant MSA-Cy5 was observed in the synovium and cartilage of PTOA knees but not in contralateral control knees (**Figure 1B, Supplementary Figure 1**). In contrast, Cy5-conjugated poly(ethylene glycol) (PEG-Cy5) did not accumulate preferentially in PTOA knees (**Figure 1C**). Further, i.v. delivery of the albumin binding dye Evans Blue resulted in preferential accumulation in synovia and cartilage of PTOA knees, but not contralateral healthy knees (**Figure 1D,E, Supplementary Figures 2,3**). Interestingly, mRNAs of genes encoding the albumin delivery / transport factors caveolin-1 and secreted protein acidic and rich in cysteine (SPARC) were elevated in PTOA joint tissues (**Supplementary Figure 3**), consistent with previous reports of SPARC upregulation in arthritic joints[33-36]. Interestingly, interstitial SPARC expression in tumors correlates with efficacy of albumin-based therapeutics in head and neck cancers[37]. These findings support the idea that hitchhiking on endogenous circulating albumin might be used to skew the biodistribution of siRNA conjugates to arthritic joints following i.v. delivery.

Enhanced stability, albumin binding, and PTOA joint accumulation of optimized siRNA-lipid conjugates.

RNA therapeutics i.v. administered carrier-free would be highly vulnerable to degradation by endo- and exonucleases. To stabilize the siRNA sequence, we engineered blunt-ended 19-mer RNA strands with alternating 2′-OMe and 2′-F ribose modifications in a "zipper" pattern and replaced phosphodiesters with PS linkages for the last two bases of the 5□- and 3□-termini of the sense and anti-sense strands (**Figure 1F,G**). Stability of the zipper-modified siRNA was assessed in synovial fluid from a treatment-naive OA patient, revealing complete siRNA stability through at least 24 hrs at 37ºC, whereas a more lightly modified dicer-substrate siRNA was >50% degraded within 1 hr **(Figure 1H)**. Given the stability of the zipper-modified siRNA, all further siRNA modifications were done using this modification pattern.

The zipper siRNAs were end-modified with bivalent C18 lipids intended to promote binding of the siRNA into the natural fatty acid (**FA**) binding pockets of albumin (**Figure 1I**)[38]. A library of bivalent lipid structures were synthesized as recently reported[39]. Briefly, a splitter phosphoramidite (<) was added at the 5□ end of the sense strand, followed by addition of 0 (EG_0), 1 (EG_6), 3 (EG_{18}), or 5 (EG_{30}) hexa-ethylene glycol (EG_6) phosphoramidite spacers to each branch. Both branches were terminally appended with a C_{18} lipid, thus generating siRNA<$(EG_xL)_2$, where X is the number of EG repeats linking the splitter to each terminal C18.

Association of the different siRNA conjugates and the parent zipper siRNA structures with synovial fluid albumin was assessed using fast protein liquid chromatography (**FPLC**). A proportion of each of the siRNA<$(EG_XL)_2$ conjugates was found in the albumin-containing synovial fluid fractions (**Figure 1J**), whereas unconjugated siRNA was not. Notably, >80% of the total siRNA<$(EG_{18}L)_2$ detected by FPLC was found in the albumin-containing fraction, higher than any other siRNA-lipid conjugate tested. This observation was consistent with the previously-measured high affinity binding of siRNA<$(EG_{18}L)_2$ to albumin (K_D = 30 nM)[39].

Cy5-labeled siRNA<(EG$_X$L)$_2$ constructs were next delivered i.v. (1 mg/kg) to mice with PTOA induced in the left knee (**Figure 2A**). While each accumulated to a greater extent in PTOA knees over healthy contralateral knees, siRNA<(EG$_{18}$L)$_2$ achieved the highest accumulation PTOA joints (**Figure 2B, Supplementary Figure 4**). In contrast to the >4-fold enrichment of siRNA<(EG$_{18}$L)$_2$ in PTOA knees over healthy knees, parental siRNA and cholesterol-conjugated siRNA (**siRNA-Chol**) each showed only a 1.5- and 2-fold enrichment to PTOA vs healthy knees (**Figure 2C**). Histological cryosections of PTOA knee synovium and cartilage illustrated holistic joint tissue penetration and cellular uptake of siRNA<(EG$_{18}$L)$_2$, while siRNA-Chol was observed with greatly reduced intensity in these PTOA joint compartments (**Figure 2D, Supplementary Figure 5**). Organ biodistribution studies performed by ex vivo Cy5 imaging showed that liver was the main off-target delivery site for both siRNA<(EG$_{18}$L) and siRNA-Chol, while free siRNA accumulated primarily in kidneys (**Supplementary Figure 6**), a major site of siRNA clearance.

Interactions between albumin and siRNA<(EG$_{18}$L)$_2$ were further assessed by FPLC of synovial fluid from OA, RA, or healthy joints. Like what was observed using healthy patient-derived synovial fluid, Cy5-siRNA<(EG$_{18}$L)$_2$ eluted primarily in the albumin-containing fractions of OA patient and RA patient-derived synovial fluid samples, but with substantially greater intensity in arthritic samples (**Figure 2E, F**). In contrast, the parent Cy5-siRNA remained predominantly unbound. Western blotting confirmed the increased albumin content in OA- and RA-derived synovial fractions B12 and C1, the same fractions that siRNA<(EG$_{18}$L)$_2$ was predominantly associated with (**Supplementary Figure 7**). Complementary native gel electrophoresis analyses of synovial fluid fractions B12-C1 revealed colocalization of Cy5-siRNA<(EG$_{18}$L)$_2$ with a Coomassie-stained protein of 66 kDa, the known molecular weight of albumin. Taken together, these findings allowed us to assign siRNA<(EG$_{18}$L)$_2$ as our lead siRNA-lipid conjugate for albumin hitchhiking to arthritic joints, motivating a focus on this construct for subsequent therapeutic studies. We also found that siRNA<(EG$_{18}$L)$_2$ interacted with albumin across multiple species, including human, mouse, and guinea pig, supporting its utility for all of the OA and RA models used herein**.

siRNA<(EG$_{18}$L)$_2$ accumulation in PTOA joints following subcutaneous and intra-articular delivery.

Subcutaneous injection of siRNA<(EG$_{18}$L)$_2$ was tested as an alternative to the i.v. delivery route, given the utility of subcutaneous injections for patient self-administration of biologic drugs. Subcutaneous delivery of Cy5-siRNA<(EG$_{18}$L)$_2$ (2mg/kg) enabled preferential siRNA accumulation in PTOA knees over contralateral unchallenged knees, with similar organ biodistribution profile as i.v. delivery (**Extended Data Fig 1A-C**). However, subcutaneous delivery of siRNA<(EG$_{18}$L)$_2$ at 2 mg/kg did not achieve the absolute level of siRNA accumulation in PTOA knees seen with i.v. delivery at 1 mg/kg (**Extended Data Fig 1D**). Notably, a large amount of the Cy5-siRNA<(EG$_{18}$L)$_2$ dose was retained at the injection site (data not shown), perhaps due to its rapid interaction with subcutaneous fat or cells at the injection site[40]. Intra-articular (**i.a.**) delivery, a common clinical route for steroid delivery, was also tested (0.25 mg/kg), revealing greater retention in the PTOA over healthy knee and greater retention than Cy5-siRNA at 48 hrs post-injection, achieving penetration into cartilage and synovia (**Extended Data Fig 1E-G**). While i.a.-delivered Cy5-siRNA<(EG$_{18}$L)$_2$ was retained primarily in the knee, Cy5-siRNA redistributed to kidneys, reflecting its relatively rapid removal from the joint by synovial drainage via vasculature and lymphatics, underscoring the importance of albumin-binding moieties for retention of siRNA<(EG$_{18}$L)$_2$ within the joint space.

Potent MMP13 knockdown in PTOA-afflicted joints by siRNA<(EG$_{18}$L)$_2$

siRNA sequences against mouse and guinea pig MMP13 (**siMMP13**) were screened in mouse ATDC5 chondrogenic cells and primary guinea pig chondrocytes, respectively, for MMP13 knockdown potency (**Supplementary Figure 8**). Lead candidate sequences were then synthesized with zipper modifications, enhancing siMMP13 stability in serum (**Supplementary Figure 9**). Transfection of zipper-modified siRNAs achieved potent MMP13 knockdown activity in TNFα-stimulated ATDC5 cells (**Supplementary Figure 9**). Zipper-pattern siMMP13 was further modified with diacyl lipids as described above to enable albumin binding, thus generating siMMP13<(EG$_{18}$L)$_2$. Importantly, siMMP13<(EG$_{18}$L)$_2$, but not free siMMP13, provided robust carrier-free (no lipofection reagent), *Mmp13* knockdown and efficient cellular uptake in ATDC5 cells (**Supplementary Figure 9,10**). Cell uptake of siMMP13<(EG$_{18}$L)$_2$ was partially diminished in

the presence of excess albumin but increased upon treatment with pro-inflammatory TNFα stimulation, even in the presence of excess albumin.

MMP13silencing by siMMP13<(EG$_{18}$L)$_2$ in PTOA-affected mouse knees was first assessed 5 days after i.v. treatment, finding >50% *Mmp13* knockdown with a 5 mg/kg dose, and >60% with a 10 mg/kg dose (**Extended Data Fig 2A**). Neither free siMMP13 nor siMMP13-Chol (each at 10mg/kg, i.v.) significantly affected *Mmp13* levels in PTOA knees. A single intra-articular (i.a.) injection of siMMP13<(EG$_{18}$L)$_2$ at 1 mg/kg also decreased *Mmp13* in PTOA knees (**Extended Data Fig 2B**). While subcutaneous delivery of siMMP13<(EG$_{18}$L)$_2$ at a relatively high dose (50 mg/kg) diminished *Mmp13* in PTOA knees (**Extended Data Fig 2C**), a lower dose (20 mg/kg) reduced *Mmp13* in PTOA knees only when co-delivered with mouse albumin (1:5 molar ratio). These results show promise for use of siMMP13<(EG$_{18}$L)$_2$ with multiple delivery routes. However, subcutaneous delivery requires more development to promote injection site absorption.

Immunohistochemical (**IHC**) analysis of PTOA knees revealed abundant MMP13 in articular cartilage, synovium, and meniscus compared to what was seen in healthy knees. PTOA-induced MMP13 was markedly reduced in each PTOA knee compartment at 5 days after treatment with siMMP13<(EG$_{18}$L)$_2$ delivered at either 5 or 10 mg/kg. However, neither siMMP13-Chol nor siMMP13 impacted MMP13 levels in PTOA knees (**Extended Data Fig 2D**). MMP13 protein was undetectable in liver or kidneys by IHC (**Extended Data Fig 2E**). RNA expression analyses similarly did not detect *Mmp13* in liver, although low levels were found in kidney, albeit 6000-fold lower than that detected in PTOA knees (**Extended Data Fig 2E**). Regardless, kidney *Mmp13* gene expression was unaffected in siMMP13<(EG$_{18}$L)$_2$-treated mice (**Extended Data Fig 2F**), supporting the idea that siMMP13<(EG$_{18}$L)$_2$ might have a wide therapeutic window, with minimal risk of molecularly on-target side effects in two of the primary organs commonly associated with siRNA clearance, liver and kidney.

Sustained stability and bioavailability of siMMP13<(EG$_{18}$L)$_2$ enables potent and durable MMP13 knockdown in PTOA joints following systemic administration.

Mice were subjected to mechanical left knee loading (3 times / week) for 37 days. They were i.v. treated with a single dose of Cy5-siRNA<(EG$_{18}$L)$_2$ at 10mg/kg after the first week of loading, and longitudinal measurements of Cy5 retention in PTOA knees were performed at time points throughout the next 30 days (**Extended Data Fig 3A**). At day 30, remarkable Cy5 signal was retained in cartilage/meniscus and synovial tissues of PTOA knees treated systemically with Cy5-siRNA<(EG$_{18}$L)$_2$ (**Extended Data Fig 3B)**. Again, preferential siRNA<(EG$_{18}$L)$_2$ accumulation in PTOA knees over contralateral control knees was observed (**Extended Data Fig 3C,D)**. Importantly, a single siMMP13<(EG$_{18}$L)$_2$ treatment maintained knockdown of MMP13 transcript and protein in PTOA joints through day 30 (**Extended Data Fig 3E,F**). Intra-articular dosing of siRNA<(EG$_{18}$L)$_2$ (1mg/kg) was similarly assessed, again revealing preferential retention of Cy5-siRNA<(EG$_{18}$L)$_2$ in PTOA knees over contralateral healthy knees through 30 days post-treatment (**Extended Data Fig 4**).

In contrast to the sustained siRNA<(EG$_{18}$L)$_2$ retention in PTOA joints, minimal Cy5 fluorescence was seen in other organs at 30 days post-treatment (**Supplementary Figure 11**). In a side-by-side comparison, an i.v. dose of 10 mg/kg and intra-articular dose of 1 mg/kg siRNA<(EG$_{18}$L)$_2$ had remarkably similar area under the curve (**AUC**) profiles, suggesting that an approximately 10-fold higher dose should be used for the i.v. relative to the i.a. delivery route for siRNA<(EG$_{18}$L)$_2$ arthritic knee joint treatments. These trends were confirmed by a peptide nucleic acid (**PNA**) hybridization assay that enables absolute siRNA quantification in tissue. This measurement showed preferential accumulation of siRNA<(EG$_{18}$L)$_2$, but not free siRNA, in PTOA knees over contralateral healthy knees, and similar siRNA levels (measured as ng siRNA / mg tissue) when delivered at 10 mg/kg i.v. and at 1 mg/kg i.a (**Supplementary Figure 11D**).

siMMP13<(EG$_{18}$L)$_2$ diminishes molecular, histological, and clinical manifestations of PTOA.

Therapeutic efficacy of siMMP13<(EG$_{18}$L)$_2$ was tested in mouse PTOA model after 5 weeks of mechanical knee loading (3 times / week). Mice were treated with siMMP13<(EG$_{18}$L)$_2$ or siControl<(EG$_{18}$L)$_2$ (10 mg/kg) on day 7 and again on day 21 (**Figure 3A**). The pressure-pain threshold in PTOA knees was measured by algometer, revealing a 45% reduction in threshold pressure tolerance in siControl<(EG$_{18}$L)-treated mice, while siMMP13<(EG$_{18}$L)$_2$ lost only 25% (**Figure 3B**). The siMMP13<(EG$_{18}$L)$_2$ cohort showed relief of joint sensitivity to a level similar to mice receiving intra-articular delivery of the FDA-approved sustained release corticosteroid Zilretta (8 mg/kg) or 3X week intraperitoneal treatment with the experimental MMP13-selective small molecule inhibitor CL-82198[19] (10 mg/kg) (**Figure 3A,B**). However, treatment with CL-82198 on the same schedule as siMMP13<(EG$_{18}$L)$_2$ (once every two weeks) produced a less favorable pain tolerance in PTAO knees, as did treatment with the pan-MMP inhibitor Marimastat (10 mg/kg, i.p., 3 times weekly).

Pan-MMP activity in PTOA joints was assessed *in vivo* on treatment day 30 using MMPSense, an MMP substrate that fluoresces upon proteolytic cleavage. Intravital imaging revealed significantly elevated MMP-driven fluorescence in PTOA knees of mice treated with siControl<(EG$_{18}$L)$_2$ as compared to healthy knees (**Figure 3C**). Decreased pan-MMP activity was seen in PTOA knees of mice treated with siMMP13<(EG$_{18}$L)$_2$, Zilretta, and Marimastat, although the MMP13-selective small molecule inhibitor CL-82198 did not significantly diminish pan-MMP activity in PTOA knees. A fluorescently labeled monoclonal antibody against aberrantly exposed CII collagen (mAbCII) was used to assess relative cartilage damage[41-43], finding significantly elevated mAbCII localization to PTOA knees (**Figure 3C**), thus confirming PTOA-induced cartilage damage in this model. However, mAbCII intensity in PTOA knees was diminished significantly in siMMP13<(EG$_{18}$L)$_2$-treated samples, approximating the mAbCII signal in healthy knees. Similar trends were observed for serum-based measurements of collagen C2C degradation fragments, a biomarker of cartilage degradation[44]. PTOA-induced mAbCII signal and serum C2C levels were also diminished by CL-82198 when dosed with higher frequency (3 times per week), underscoring the causative role ofMMP13 in PTOA-induced collagen fragmentation and cartilage damage and confirming that MMP13 targeting by siMMP13<(EG$_{18}$L)$_2$ reduces structural OA manifestations. The long-acting steroid Zilretta had little impact on mAbCII binding in PTOA knees or on serum C2C levels, consistent with observations that steroids temporarily control arthritis pain but do not target underlying joint destruction and may even accelerate cartilage thinning[45].

Histological analyses of PTOA joints collected on treatment day 30 revealed profound disruption of cartilage architecture, proteoglycan loss, synovial thickening, osteophyte formation, and meniscal mineralization (**Figure 3D**). In PTOA knees treated with siMMP13<(EG$_{18}$L)$_2$, cartilage morphology, proteoglycan content, synovial structure, and meniscal anatomy were each relatively maintained, as quantitated using the Osteoarthritis Research Society International (OARSI) scoring criteria[46,47] for OA progression (**Figure 3E, Supplementary Table 1**). Degenerative joint disease scoring (**Supplementary Table 1**) confirmed significantly less PTOA joint structural changes in mice treated with siMMP13<(EG$_{18}$L)$_2$ versus those treated with non-targeting siControl<(EG$_{18}$L)$_2$. Unlike what was seen with siMMP13<(EG$_{18}$L)$_2$, modest reductions in OA progression and joint degeneration were achieved in PTOA joints treated with CL-82198, although these trends did not reach statistical significance. OA progression and joint degeneration scores were not impacted by treatment with Zilretta or Marimistat.

Using micro-computed tomography (**microCT**) to quantify osteophytes and ectopic mineralization in PTOA mouse knees, the significant protection conferred by siMMP13<(EG$_{18}$L)$_2$ against meniscal mineralization and osteophyte formation was confirmed (**Extended Data Fig 5A-C**). Importantly, pathologically elevated MMP13 expression in PTOA cartilage, meniscus, and synovial compartments was reduced by treatment with siMMP13<(EG$_{18}$L)$_2$, correlating with decreased presence of degradation fragments (**Extended Data Fig 5D-G**). These data show that MMP13 silencing in arthritic joints provides both molecular and clinically tangible benefits, reducing inflammation, clinical pain, articular cartilage loss, and arthritis-associated secondary manifestations.

MMP13 knockdown in PTOA knees reprograms pro-inflammatory gene expression patterns.

To understand the gene expression patterns that underlie the therapeutic outcomes associated with MMP13 silencing, we first measured the impact of PTOA on a small panel of pro-inflammatory genes with previously described roles in OA, including *Ngf*, *Il1b*, *Tnf*, and *Cdkn2a*. These genes were each elevated in mouse PTOA knees compared to healthy knees (**Extended Data Fig 6A**). We found that PTOA-induced levels of *Ngf*, *Il1b*, *Tnf*, and *Cdkn2a* were significantly dampened as early as 5 days following i.v. delivery of siMMP13<(EG$_{18}$L)$_2$ at 10 mg/kg, gene expression changes that were sustained through day 30. These findings motivated a broader and unbiased analysis PTOA-induced gene expression changes in knee tissue, and the impact of MMP13 knockdown therein, using the nanoString nCounter Inflammation 254 gene panel. Unsupervised analysis of gene expression illustrated extensive PTOA-induced gene expression changes, including genes associated with the p38 mitogen associated protein kinase (**MAPK**) stress signaling pathway (**Extended Data Fig 6B, Supplementary File S1**). The serine-threonine kinase p38 MAPK phosphorylates a network of substrates that respond to cell stress[48]. Relevant to OA, p38 MAPK signaling through its substrate MAPK-associated protein kinase 2 (**MK2**) regulates chondrocyte differentiation, cellular senescence, MMP upregulation and pro-inflammatory cytokine production[49]. MK2 signaling within human arthritic chondrocytes drives expression of prostaglandin E2, MMP3, and MMP13, leading to arthritic pain, inflammation, and cartilage degradation[50]. Interestingly, we found that the PTOA-induced p38 MAPK gene cluster was the gene enrichment set most potently suppressed in siMMP13<(EG$_{18}$L)$_2$ treated samples (**Extended Data Fig 6C**).

Treatment with siMMP13<(EG$_{18}$L)$_2$ also significantly suppressed IL-1, chemokine, and TNFα / NFκB pathway-associated gene clusters. Notably, these clusters include several genes associated with OA pathogenesis, including those associated with chondrocyte proliferation (*Mef2d*), synovial fibroblast prolfieration (*Myc*, *Jun*, *Tgfb2*, *Pdgfa*)[51,52], angiogenesis (*Pdgfba*, *Flt1*, *Hif1a*), ECM degradation (*Mmp3*, *Masp1*), inflammatory signaling through Nuclear Factor-κB, (*Nfκb1*, *Rela*, *Relb*), cytokines (*Il1b*, *Stat1*), chemokines (*Cxcl1*, *Cxcl10*, *Ccl3*, *Cxcr4*), and toll-like receptors (*TLR1,-2, -4, -5, -6, and -8*).

Given that many inflammatory and stress response factors induce MMP13 expression and that proteolytic matrix degradation fragments further potentiate inflammation and expression of stress response factors,[53] these data suggest that siMMP13<(EG$_{18}$L)$_2$ uncouples a feed-forward inflammation-degradation cycle that drives arthritic joint pathology. These data support this hypothesis, given that selective MMP13 inhibition in arthritic joints suppresses expression of both pro-inflammatory factors and transcription factors that promote MMP13 expression.

K/BxN Serum Transfer Arthritis (STA) Multi-Joint Rheumatoid/Inflammatory Arthritis Mouse Model

A role for MMP13 has been reported in rheumatoid arthritis (**RA**)[4,54-56], including studies in which daily dosing with an experimental MMP13 small molecule inhibitor decreased RA pathogenesis in 2 models[54]. We used the transgenic K/BxN serum transfer RA model to test therapeutic efficacy of siMMP13<(EG$_{18}$L)$_2$. In this model, KRN, a T-cell receptor that binds to an endogenous glucose-6-phosphate isomerase (**GPI**) peptide presented by the IAg7 major histocompatibility complex class II (**MHC-II**) allele, drives production of anti-GPI autoantibodies and GPI-directed auto-immunity that localizes to articular cartilage/joint tissues. K/BxN mouse serum transiently confers RA to wild-type mouse serum recipients. MMP13 expression is increased in inflamed joints of K/BxN serum recipients, and MMP13-deficient mice are partially protected from K/BxN serum-induced RA[56]. Carrier-free siRNA delivery in RA has not yet been explored to our knowledge, but siMMP13<(EG$_{18}$L)$_2$ systemic administration for delivery to multiple-afflicted joints is potentially advantageous. Interestingly, albumin-based delivery approaches have been shown to facilitate therapeutic outcomes in RA preclinical-studies[29,30,57-65]. Importantly, i.v. delivery of albumin-binding Evans Blue to mice resulted in concentrated Evans Blue accumulation in arthritic forepaws and hindpaws of K/BxN serum recipients as compared to unchallenged mice (**Extended Data Fig 7A-7B**). Likewise, i.v. Cy5-labeled MSA delivered i.v. accumulated in arthritic forepaws and hindpaws of K/BxN serum recipients, but not in unchallenged wild-type mice (**Extended Data Fig 7C**). Expression of Caveolin-1 and SPARC was elevated in hindpaws of K/BxN recipient mice (**Extended Data Fig 7D**). These data confirm increased albumin retention in RA-affected joints, supporting therapeutic testing of albumin-binding si<(EG$_{18}$L)$_2$ in RA models.

Mice with induced RA showed accumulation of i.v. injected Cy5-siRNA$<$(EG$_{18}$L)$_2$ within multiple inflamed joints (forepaw, wrist, hindpaw, ankle, knee) (**Figure 4A; Supplementary Figure 12**) at substantially higher levels than what was seen in healthy mice (**Extended Data Fig 7E-7G**) and exceeding that achieved by treatment with Cy5-siRNA or Cy5-siRNA-Chol (**Figure 4A**). Fluorescence microscopy confirmed increased siRNA$<$(EG$_{18}$L)$_2$ in fore- and hindpaws of K/BxN recipients, with specific accumulation in soft tissue and cartilage (**Figure 4B**). Longevity of siRNA retention was assessed in limbs of K/BXN serum recipients following a single dose Cy5 -siRNA$<$(EG$_{18}$L)$_2$. Intravital fluorescence was detected in limbs throughout the 30 days that mice were monitored (**Supplementary Figure 12**), supporting the notion of albumin-mediated delivery and retention of siRNA$<$(EG$_{18}$L)$_2$ in multiple afflicted joints in RA models.

siMMP13$<$(EG$_{18}$L)$_2$ therapy reduces manifestations and symptoms of RA.

Strong *Mmp13* induction was seen in forepaws, knees, and hindpaws of K/BxN serum recipients as compared to healthy mice (**Figure 4C**). At treatment day 8 following a single 10 mg/kg dose of siMMP13$<$(EG$_{18}$L)$_2$, mRNA and protein was decreased in forepaws, knees, and hindpaws **(Figure 4C and Extended Data Fig 8A)**. Like with PTOA, MMP13 silencing in this RA model reduced expression of pro-inflammatory genes encoding cyclooxygenase 2 (COX2), IL-6, TNF-α, and IL-1B **(Supplementary Figure 13)**. Clinical parameters of RA were also reduced upon MMP13 knockdown, including ankle swelling, pressure pain threshold, arthritis clinical score, disease severity index (**Figure 4D-4F**), and the number of afflicted joints per mouse (**Supplementary Figure 14**). In parallel studies, K/BxN serum recipients were treated with a single dose of Cl-82198 (10 mg/kg, i.p.) or methylprednisolone (10 mg/kg, i.p.), neither of which improved ankle swelling (**Figure 4D-F**). Although methylprednisolone, but not CL-82198, improved clinical score and pressure tolerance, the effect size was not as pronounced as what was seen with MMP13$<$(EG$_{18}$L)$_2$. Similarly, the disease severity score was somewhat improved upon treatment with CL-82198, but to a lesser extent than what was seen with siMMP13$<$(EG$_{18}$L)$_2$. Further, MMP13 silencing resulted in decreased mAbCII binding, indicative of cartilage preservation (**Extended Data Fig 8B**) and decreased total MMP activity (**Extended Data Fig 8C**).

Histologic examination of toluidine blue-stained cartilage harvested on treatment day 8 in K/BxN recipient mice revealed substantial cartilage destruction in hindpaws, knees, and forepaws, a phenotype that was diminished by siMMP13$<$(EG$_{18}$L)$_2$ treatment (**Figure 5A-5B** and **Extended Data Fig 9A-9B**; lower magnification images in **Supplementary Figures 15-18**). Inflammation scoring illustrated similar trends **(Figure 5C and Extended Data Fig 9C-D)**. MicroCT studies of bone structure in untreated K/BxN serum recipients revealed erosive bone disease characterized by increased bone surface to volume ratios, decreased bone mineral density (BMD), and decreased bone volume / total volume (**BV/TV**) ratio (**Figure 5D-5G**; and **Supplementary Figure 16**). Remarkably, treatment with siMMP13$<$(EG$_{18}$L)$_2$ protected K/BxN serum recipients from inflammation-induced changes in bone surface to volume, BMD, and BV/TV measurements. Further, bone erosions characteristic of inflammatory arthritis were evident in forepaws, knees, and hindpaws of untreated K/BxN serum, but were minimal or absent in mice treated with siMMP13$<$(EG$_{18}$L)$_2$ (**Extended Data Fig 9E**). In contrast, Cl-82198 and methylprednisolone did not diminish features of erosive bone disease detected by microCT. Serum C2C degradation fragments were significantly elevated in untreated, CL-82198 treated, and methylprednisolone-treated K/BxN serum recipients (**Extended Data Fig 10A**), while treatment with siMMP13$<$(EG$_{18}$L)$_2$ significantly reduced serum C2C fragments. These findings were confirmed using IHC for C1,2C in K/BxN serum recipients hindpaw/ankle and knee joint tissues **(Extended Data Fig 10B)**.

Hindpaw ankle articular cartilage and synovial RNA was harvested from K/BxN serum transfer recipients on treatment day 8 and assessed using the nanoString nCounter mouse inflammation expression panel. These studies identified several gene clusters that were elevated in untreated K/BxN serum transfer recipients as compared to healthy mice (**Supplementary File S1**), many of which were downregulated in samples collected from K/BxN serum transfer recipients treated with siMMP13$<$(EG$_{18}$L)$_2$ (**Extended Data Fig 10C**). Treated RA animals had gene expression patterns that mirroredthe changes seen in PTOA

knees upon MMP13 knockdown (*e.g.,* IL-1 signaling, NF-kB signaling, and chemokine pathways were suppressed by treatment, **Extended Data Fig 10D**). Additionally, the pro-inflammatory COX-2 encoding gene *Ptgs2*, the chemokine genes *Cxcl5, Ccl7*, *Ccl19*, and the RA-associated macrophage genes *Tnfsf14* and *Tr2* were significantly reduced in K/BxN hindpaws treated with siMMP13<(EG$_{18}$L)$_2$. Thus, the molecular markers of inflammation that characterize RA are dampened by MMP13 knockdown achieved by i.v. delivery of siMMP13<(EG$_{18}$L)$_2$.

Importantly, serum levels of toxicity markers, including blood urea nitrogen (**BUN**), alanine transaminase (**ALT**), aspartate amino transferase (**AST**), creatine phosphokinase (**CPK**), lactate dehydrogenase (**LDH**), and glucose, were detected within the normal ranges, lacking significant differences between healthy, PTOA, or K/BxN serum recipient mice in any of the treatment groups (**Supplementary Figure 19)**. Liver, kidney, lung, heart, and spleen of mice within all groups also showed a normal gross and histological structure (**Supplementary Figure 20**).

MMP13 silencing in a Dunkin-Hartley guinea pig ACL transection (ACLT) large animal model

ACL transection (**ACLT**) was done on the left knee of Dunkin Hartley guinea pigs (**DHGP**s) to establish proof of concept in PTOA in a larger species (**Figure 6A**). Like humans, DHGPs develop OA with aging, with the medial knee compartment showing loss of proteoglycans, fibrillation, cloning of chondrocytes, osteophyte formation, and subchondral bone sclerosis[66]. ACLT is used to mimic PTOA and accelerate the spontaneous OA phenotype[67]. MMP13 is strongly expressed and highly localized with OA cartilage lesions in DHGPs[66]. Cy5-siRNA<(EG$_{18}$L)$_2$ was delivered i.v. on post-surgical day 13 (1 mg/kg, i.v.), and knees were imaged *ex vivo* to measure Cy5 fluorescence on day 14, revealing significantly increased Cy5-siRNA<(EG$_{18}$L)$_2$ accumulation in ACLT knees over unaltered knees (**Figure 6B**), while free Cy5-siRNA did not exhibit preferential accumulation in ACLT damaged knees. Ex vivo Cy5 imaging of organs showed abundant free siRNA in kidneys, while the albumin binding siRNA<(EG$_{18}$L)$_2$ was found at highest levels in ACLT-damaged knees and liver, but not in healthy knees nor in kidneys (**Figure 6C**). Fluorescence microcopy confirmed delivery of Cy5-siRNA<(EG$_{18}$L)$_2$ but not free siRNA, to synovia of ACLT-damaged knees, with strong penetration into cartilage (**Figure 6D**). Intravenous siMMP13<(EG$_{18}$L)$_2$ delivered on post-surgical day 7 at 10 mg/kg decreased *Mmp13* expression in ACLT-damaged knees by >80% (**Figure 6E**), and MMP13 protein expression by >70% on day 14 (**Figure 6F-G)**.

Conclusion

A unique albumin-binding RNAi conjugate (siRNA<(EG$_{18}$L)$_2$) was engineered for MMP13-selective gene targeting in the context of arthritis. This optimized siRNA-lipid conjugate was capable of preferential accumulation and long-term retention within arthritic joints, robustly decreasing arthritis-induced MMP13 and interrupting the forward-feeding cycle of inflammatory joint damage to limit arthritis progression. Overall, this albumin-binding siRNA-L$_2$ system shows promise as a systemic and/or local anti-MMP13-specific therapy. This siRNA delivery platform is readily amenable to modular engineering for targeting other disease-driving genes beyond the scope of arthritis.

Supplementary Files

File S1. Excel file of gene names and fold expression changes in knee RNA samples harvested from healthy mice, untreated PTOA mice, and PTOA mice treated with siMMP13<(EG$_{18}$L)$_2$ as well as hindpaw RNA samples harvested from healthy mice, untreated K/BxN serum recipient mice, and K/BxN serum recipients treated with siMMP13<(EG$_{18}$L)$_2$.

Author Contributions

J.M.C. and C.L.D. conceptualized the study and experiments. J.M.C., E.N.H., J.H.L, A.G.S, and N.F, synthesized and characterized the siRNA conjugates and assisted with PNA hybridization. J.M.C. performed *in vitro* siRNA studies. J.M.C., V.S., F.Y. performed *in vivo* pharmacokinetic studies. J.M.C.

performed *in vivo* confocal microscopy imaging and analyses. D.L.M and K.C.V performed and supervised SEC synovial fluid analysis. J.M.C., V.S., F.Y. performed *in vivo* therapeutic studies including MMPsense, qRT-PCR, microCT, and histological tissue preparation. K.N.G-C performed pathologist-blinded OARSI, DJD, and K/BxN STA scoring of histological sections. L.C. provided K/BxN serum and experimental insight for inflammatory arthritis studies. H.C. and K.A.H. provided mAbCII680 reagent and experimental insight for *in vivo* cartilage damage studies. R.C. and C.L.D. provided study mentorship and oversight. J.M.C., R.C, C.L.D. co-wrote the manuscript with input from all authors.

Acknowledgements and Funding

The authors acknowledge the assistance of the Vanderbilt Translational Pathology Shared Resource (TPSR), which is supported by NCI/NIH Cancer Center Support Grant 2P30 CA068485-14 and the Shared Instrumentation Grant S10 OD023475 for the Leica Bond Rx. Micro CT analyses was supported in part by the NIH (S10RR027631-01). The histological technical assistance of Cindy Lowe and Karen Thompson and the microCT technical assistance by Sasidhar Uppuganti is acknowledged. We are grateful for funding from the DOD (DOD CDMRP OR130302), NIH (NIH R01 CA224241, NIH R01 EB019409, NIH R01 AR078666, NIH R01 CA260958, NIH R21 AR078636, NIH T32GM007347), Canadian Natural Sciences and Engineering Research Council (NSERC), the Rheumatology Research Foundation (RRF), and the VA Merit Awards BX004151 and BX004472. The content in this report is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

Data Availability Statement

The main data supporting the results in this study are available within the paper and its Supplementary Information. NanoString datasets are available at the Gene Expression Omnibus under accession identifier GSE233714. The remaining raw and analysed datasets from the study are too large to be publicly shared, but they are available for research purposes from the corresponding author on reasonable request.

Conflict of Interest

The authors declare no conflict of interest.

References

Loeser, R. F. Osteoarthritis year in review 2013: biology. *Osteoarthritis and Cartilage* **21**, 1436-1442, doi:http://dx.doi.org/10.1016/j.joca.2013.05.020 (2013).

Gullo, T. R. *et al.* Defining multiple joint osteoarthritis, its frequency and impact in a community-based cohort. *Seminars in Arthritis and Rheumatism* **48**, 950-957, doi:10.1016/j.semarthrit.2018.10.001 (2019).

McInnes, I. B. & Schett, G. The pathogenesis of rheumatoid arthritis. *The New England journal of medicine* **365**, 2205-2219, doi:10.1056/NEJMra1004965 (2011).

Shiozawa, S., Tsumiyama, K., Yoshida, K. & Hashiramoto, A. Pathogenesis of joint destruction in rheumatoid arthritis. *Arch Immunol Ther Exp (Warsz)* **59**, 89-95, doi:10.1007/s00005-011-0116-3 (2011).

Bradley, J. D., Brandt, K. D., Katz, B. P., Kalasinski, L. A. & Ryan, S. I. Comparison of an antiinflammatory dose of ibuprofen, an analgesic dose of ibuprofen, and acetaminophen in the treatment of patients with osteoarthritis of the knee. *The New England journal of medicine* **325**, 87-91, doi:10.1056/NEJM199107113250203 (1991).

Hochberg, M. C. *et al.* American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res (Hoboken)* **64**, 465-474 (2012).

McAlindon, T. E. *et al.* OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage* **22**, 363-388, doi:10.1016/j.joca.2014.01.003 (2014).

Benjamin, O., Goyal, A. & Lappin, S. L. in *StatPearls*   (2022).
Bolduc, J. A., Collins, J. A. & Loeser, R. F. Reactive oxygen species, aging and articular cartilage homeostasis. *Free Radic Biol Med* **132**, 73-82, doi:10.1016/j.freeradbiomed.2018.08.038 (2019).
Young, I. C. *et al.* A novel compressive stress-based osteoarthritis-like chondrocyte system. *Exp Biol Med (Maywood)* **242**, 1062-1071, doi:10.1177/1535370217699534 (2017).
Jasin, H. E., Noyori, K., Takagi, T. & Taurog, J. D. Characteristics of anti-type II collagen antibody binding to articular cartilage. *Arthritis and rheumatism* **36**, 651-659 (1993).
Hollander, A. P. *et al.* Damage to type II collagen in aging and osteoarthritis starts at the articular surface, originates around chondrocytes, and extends into the cartilage with progressive degeneration. *J Clin Invest* **96**, 2859-2869, doi:10.1172/JCI118357 (1995).
Loeser, R. F. Integrins and chondrocyte-matrix interactions in articular cartilage. *Matrix biology : journal of the International Society for Matrix Biology* **39**, 11-16, doi:10.1016/j.matbio.2014.08.007 (2014).
Krzeski, P. *et al.* Development of musculoskeletal toxicity without clear benefit after administration of PG-116800, a matrix metalloproteinase inhibitor, to patients with knee osteoarthritis: a randomized, 12-month, double-blind, placebo-controlled study. *Arthritis Res Ther* **9**, R109, doi:10.1186/ar2315 (2007).
Molina, J. R. *et al.* A phase I and pharmacokinetic study of the selective, non-peptidic inhibitor of matrix metalloproteinase BAY 12-9566 in combination with etoposide and carboplatin. *Anticancer Drugs* **16**, 997-1002 (2005).
Clutterbuck, A. L., Asplin, K. E., Harris, P., Allaway, D. & Mobasheri, A. Targeting matrix metalloproteinases in inflammatory conditions. *Current drug targets* **10**, 1245-1254 (2009).
Howes, J. M. *et al.* The recognition of collagen and triple-helical toolkit peptides by MMP-13: sequence specificity for binding and cleavage. *J Biol Chem* **289**, 24091-24101, doi:10.1074/jbc.M114.583443 (2014).
Zhang, X., Chen, C. T., Bhargava, M. & Torzilli, P. A. A Comparative Study of Fibronectin Cleavage by MMP-1, -3, -13, and -14. *Cartilage* **3**, 267-277, doi:10.1177/1947603511435273 (2012).
Wang, M. *et al.* MMP13 is a critical target gene during the progression of osteoarthritis. *Arthritis Res Ther* **15**, R5, doi:10.1186/ar4133 (2013).
Takaishi, H., Kimura, T., Dalal, S., Okada, Y. & D'Armiento, J. Joint diseases and matrix metalloproteinases: a role for MMP-13. *Curr Pharm Biotechnol* **9**, 47-54, doi:10.2174/138920108783497659 (2008).
Liu, J. & Khalil, R. A. Matrix Metalloproteinase Inhibitors as Investigational and Therapeutic Tools in Unrestrained Tissue Remodeling and Pathological Disorders. *Progress in molecular biology and translational science* **148**, 355-420, doi:10.1016/bs.pmbts.2017.04.003 (2017).
Kristen, A. V. *et al.* Patisiran, an RNAi therapeutic for the treatment of hereditary transthyretin-mediated amyloidosis. *Neurodegener Dis Manag* **9**, 5-23, doi:10.2217/nmt-2018-0033 (2019).
Adams, D. *et al.* Efficacy and safety of vutrisiran for patients with hereditary transthyretin-mediated amyloidosis with polyneuropathy: a randomized clinical trial. *Amyloid* **30**, 1-9, doi:10.1080/13506129.2022.2091985 (2023).
Bedingfield, S. K. *et al.* Amelioration of post-traumatic osteoarthritis via nanoparticle depots delivering small interfering RNA to damaged cartilage. *Nature Biomedical Engineering* **5**, 1069-1083, doi:10.1038/s41551-021-00780-3 (2021).
Bedingfield, S. K. *et al.* Top-Down Fabricated microPlates for Prolonged, Intra-articular Matrix Metalloproteinase 13 siRNA Nanocarrier Delivery to Reduce Post-traumatic Osteoarthritis. *ACS Nano* **15**, 14475-14491, doi:10.1021/acsnano.1c04005 (2021).
Sleep, D., Cameron, J. & Evans, L. Albumin as a versatile platform for drug half-life extension. *Biochimica et Biophysica Acta* (2013).

Hsueh, M. F., Khabut, A., Kjellstrom, S., Onnerfjord, P. & Kraus, V. B. CARTILAGE PERMEABILITY ASSESSMENT BASED ON PROTEOMIC ANALYSIS OF PLASMA PROTEIN PENETRATION. *Osteoarthritis and Cartilage* **23**, A132-A133, doi:10.1016/j.joca.2015.02.863 (2015).

Mannik, M. & Person, R. E. DEEP PENETRATION OF ANTIBODIES INTO THE ARTICULAR-CARTILAGE OF PATIENTS WITH RHEUMATOID-ARTHRITIS. *Rheumatology International* **14**, 95-102, doi:10.1007/bf00300809 (1994).

Wunder, A. *et al.* Albumin-based drug delivery as novel therapeutic approach for rheumatoid arthritis. *J Immunol* **170**, 4793-4801, doi:10.4049/jimmunol.170.9.4793 (2003).

Liu, M. *et al.* Selective delivery of interleukine-1 receptor antagonist to inflamed joint by albumin fusion. *BMC Biotechnol* **12**, 68, doi:10.1186/1472-6750-12-68 (2012).

Lau, S. *et al.* Enhanced extravasation, stability and in vivo cardiac gene silencing via in situ siRNA-albumin conjugation. *Mol Pharm* **9**, 71-80, doi:10.1021/mp2002522 (2012).

Bienk, K. *et al.* An albumin-mediated cholesterol design-based strategy for tuning siRNA pharmacokinetics and gene silencing. *J Control Release* **232**, 143-151, doi:10.1016/j.jconrel.2016.04.013 (2016).

Sanchez, C. *et al.* Chondrocyte secretome: a source of novel insights and exploratory biomarkers of osteoarthritis. *Osteoarthritis Cartilage* **25**, 1199-1209, doi:10.1016/j.joca.2017.02.797 (2017).

Sward, P., Frobell, R., Englund, M., Roos, H. & Struglics, A. Cartilage and bone markers and inflammatory cytokines are increased in synovial fluid in the acute phase of knee injury (hemarthrosis)--a cross-sectional analysis. *Osteoarthritis Cartilage* **20**, 1302-1308, doi:10.1016/j.joca.2012.07.021 (2012).

Nakamura, S. *et al.* Enhancement of SPARC (osteonectin) synthesis in arthritic cartilage. Increased levels in synovial fluids from patients with rheumatoid arthritis and regulation by growth factors and cytokines in chondrocyte cultures. *Arthritis and rheumatism* **39**, 539-551, doi:10.1002/art.1780390402 (1996).

Nanba, Y. *et al.* Expression of osteonectin in articular cartilage of osteoarthritic knees. *Acta Med Okayama* **51**, 239-243, doi:10.18926/AMO/30790 (1997).

Desai, N., Trieu, V., Damascelli, B. & Soon-Shiong, P. SPARC Expression Correlates with Tumor Response to Albumin-Bound Paclitaxel in Head and Neck Cancer Patients. *Transl Oncol* **2**, 59-64, doi:10.1593/tlo.09109 (2009).

Hoogenboezem, E. N. & Duvall, C. L. Harnessing albumin as a carrier for cancer therapies. *Adv Drug Deliv Rev* **130**, 73-89, doi:10.1016/j.addr.2018.07.011 (2018).

Hoogenboezem, E. *et al.* Structural Optimization of siRNA Conjugates for Albumin Binding Achieves Effective MCL1-Targeted Cancer Therapy. *bioRxiv*, 2023.2002.2014.528574, doi:10.1101/2023.02.14.528574 (2023).

Brown, K. M. *et al.* Expanding RNAi therapeutics to extrahepatic tissues with lipophilic conjugates. *Nature biotechnology* **40**, 1500-1508, doi:10.1038/s41587-022-01334-x (2022).

Eichaker, L. R., Cho, H., Duvall, C. L., Werfel, T. A. & Hasty, K. A. Future nanomedicine for the diagnosis and treatment of osteoarthritis. *Nanomedicine : nanotechnology, biology, and medicine* **9**, 2203-2215, doi:10.2217/nnm.14.138 (2014).

O'Grady, K. P. *et al.* Drug-Free ROS Sponge Polymeric Microspheres Reduce Tissue Damage from Ischemic and Mechanical Injury. *ACS Biomaterials Science & Engineering*, doi:10.1021/acsbiomaterials.6b00804 (2017).

Cho, H., Pinkhassik, E., David, V., Stuart, J. M. & Hasty, K. A. Detection of early cartilage damage using targeted nanosomes in a post-traumatic osteoarthritis mouse model. *Nanomedicine : nanotechnology, biology, and medicine* **11**, 939-946, doi:10.1016/j.nano.2015.01.011 (2015).

Ameye, L. G. *et al.* The chemical biomarkers C2C, Coll2-1, and Coll2-1NO2 provide complementary information on type II collagen catabolism in healthy and osteoarthritic mice. *Arthritis and rheumatism* **56**, 3336-3346, doi:10.1002/art.22875 (2007).

McAlindon, T. E. *et al.* Effect of Intra-articular Triamcinolone vs Saline on Knee Cartilage Volume and Pain in Patients With Knee Osteoarthritis: A Randomized Clinical Trial. *JAMA* **317**, 1967-1975, doi:10.1001/jama.2017.5283 (2017).

Glasson, S. S., Chambers, M. G., Van Den Berg, W. B. & Little, C. B. The OARSI histopathology initiative - recommendations for histological assessments of osteoarthritis in the mouse. *Osteoarthritis Cartilage* **18 Suppl 3**, S17-23, doi:10.1016/j.joca.2010.05.025 (2010).

Bolon, B. *et al.* Rodent preclinical models for developing novel antiarthritic molecules: comparative biology and preferred methods for evaluating efficacy. *J Biomed Biotechnol* **2011**, 569068, doi:10.1155/2011/569068 (2011).

Coulthard, L. R., White, D. E., Jones, D. L., McDermott, M. F. & Burchill, S. A. p38(MAPK): stress responses from molecular mechanisms to therapeutics. *Trends Mol Med* **15**, 369-379, doi:10.1016/j.molmed.2009.06.005 (2009).

Li, Z. *et al.* p38MAPK Signaling Pathway in Osteoarthritis: Pathological and Therapeutic Aspects. *J Inflamm Res* **15**, 723-734, doi:10.2147/JIR.S348491 (2022).

Jones, S. W. *et al.* Mitogen-activated protein kinase-activated protein kinase 2 (MK2) modulates key biological pathways associated with OA disease pathology. *Osteoarthritis Cartilage* **17**, 124-131, doi:10.1016/j.joca.2008.05.001 (2009).

Almasry, S. M., Soliman, H. M., El-Tarhouny, S. A., Algaidi, S. A. & Ragab, E. M. Platelet rich plasma enhances the immunohistochemical expression of platelet derived growth factor and vascular endothelial growth factor in the synovium of the meniscectomized rat models of osteoarthritis. *Ann Anat* **197**, 38-49, doi:10.1016/j.aanat.2014.10.006 (2015).

Solomon, L. A., Bérubé, N. G. & Beier, F. Transcriptional regulators of chondrocyte hypertrophy. *Birth Defects Res C Embryo Today* **84**, 123-130, doi:10.1002/bdrc.20124 (2008).

Adair-Kirk, T. L. & Senior, R. M. Fragments of extracellular matrix as mediators of inflammation. *Int J Biochem Cell Biol* **40**, 1101-1110, doi:10.1016/j.biocel.2007.12.005 (2008).

Jungel, A. *et al.* Effect of the oral application of a highly selective MMP-13 inhibitor in three different animal models of rheumatoid arthritis. *Ann Rheum Dis* **69**, 898-902, doi:10.1136/ard.2008.106021 (2010).

Kazantseva, M. G., Hung, N. A., Highton, J. & Hessian, P. A. MMP expression in rheumatoid inflammation: the rs11568818 polymorphism is associated with MMP-7 expression at an extra-articular site. *Genes Immun* **14**, 162-169, doi:10.1038/gene.2012.65 (2013).

Singh, A. *et al.* Collagenase-3 (MMP-13) deficiency protects C57BL/6 mice from antibody-induced arthritis. *Arthritis Res Ther* **15**, R222, doi:10.1186/ar4423 (2013).

Granirer, L. W. A study of the albumin and globulin content in postpartum plasma and its use in rheumatoid arthritis. *Science (New York, N.Y.)* **111**, 204, doi:10.1126/science.111.2878.204 (1950).

Elewaut, D. Right on target: coupling methotrexate to albumin to treat rheumatoid arthritis. *Rheumatology (Oxford)* **43**, 1067-1068, doi:10.1093/rheumatology/keh242 (2004).

Fiehn, C. *et al.* Albumin-coupled methotrexate (MTX-HSA) is a new anti-arthritic drug which acts synergistically to MTX. *Rheumatology (Oxford)* **43**, 1097-1105, doi:10.1093/rheumatology/keh254 (2004).

Fiehn, C., Kratz, F., Sass, G., Muller-Ladner, U. & Neumann, E. Targeted drug delivery by in vivo coupling to endogenous albumin: an albumin-binding prodrug of methotrexate (MTX) is better than MTX in the treatment of murine collagen-induced arthritis. *Ann Rheum Dis* **67**, 1188-1191, doi:10.1136/ard.2007.086843 (2008).

Byeon, H. J. *et al.* Human serum albumin-TRAIL conjugate for the treatment of rheumatoid arthritis. *Bioconjugate chemistry* **25**, 2212-2221, doi:10.1021/bc500427g (2014).

Bilthariya, U., Jain, N., Rajoriya, V. & Jain, A. K. Folate-conjugated albumin nanoparticles for rheumatoid arthritis-targeted delivery of etoricoxib. *Drug Dev Ind Pharm* **41**, 95-104, doi:10.3109/03639045.2013.850705 (2015).

Al-Rahim, A. M. *et al.* Folate-methotrexate loaded bovine serum albumin nanoparticles preparation: an in vitro drug targeting cytokines overwhelming expressed immune cells from rheumatoid arthritis patients. *Anim Biotechnol*, 1-17, doi:10.1080/10495398.2021.1951282 (2021).

Lyu, J. *et al.* Treatment of Rheumatoid Arthritis by Serum Albumin Nanoparticles Coated with Mannose to Target Neutrophils. *ACS Appl Mater Interfaces* **13**, 266-276, doi:10.1021/acsami.0c19468 (2021).

Liu, L. *et al.* Secreted Protein Acidic and Rich in Cysteine Mediated Biomimetic Delivery of Methotrexate by Albumin-Based Nanomedicines for Rheumatoid Arthritis Therapy. *ACS Nano* **13**, 5036-5048, doi:10.1021/acsnano.9b01710 (2019).

Huebner, J. L., Otterness, I. G., Freund, E. M., Caterson, B. & Kraus, V. B. Collagenase 1 and collagenase 3 expression in a guinea pig model of osteoarthritis. *Arthritis and rheumatism* **41**, 877-890, doi:10.1002/1529-0131(199805)41:5<877::aid-art16>3.0.co;2-# (1998).

Thomas, N. P. *et al.* Synovial inflammation plays a greater role in post-traumatic osteoarthritis compared to idiopathic osteoarthritis in the Hartley guinea pig knee. *BMC Musculoskelet Disord* **18**, 556-556, doi:10.1186/s12891-017-1913-6 (2017).

a Knee loading PTOA model

Mouse left knee loading

Mechanical loading 9N, 250 cycles

i.v. delivery of Cy5-Albumin, Cy5-PEG or Evans Blue

1 3 5 7 9 11 13 days

Tissue analysis

b Intravital Cy5-albumin imaging

Healthy knee   OA knee

Color Scale
Min = 3.12e8
Max = 5.01e8

Radiance (p/sec/cm3/sr) X 10⁸

Average Radiance (x10⁸)

Healthy Knee   OA Knee

***

c Intravital Cy5 imaging

Healthy knee   OA knee

Cy5-Albumin

Cy5-PEG

Radiance Ratio [OA Knee / Healthy Knee]

Cy5-Albumin   Cy5-PEG

*

d Evans Blue intravital imaging

Healthy knee   OA knee

Color Scale
Min = 2.89e9
Max = 5.00e9

Radiance (p/sec/cm3/sr) X 10⁹

Radiant efficiency (x10⁹)

Healthy knee   OA knee

**

e Evans Blue extraction

Healthy knee   OA knee

knee

extract

Evans Blue (ng) / mg tissue

Healthy knee   OA knee

**

f
Ribose ● and its modifications

● 2'-O-Me RNA   ● 2'-F RNA

Base   Base

OMe   F

Backbone modifications

⌒ Phosphodiester   ⌒ Phosphorothioate

O=P-O⁻   S=P-O⁻

Terminus modifications

Hydrophilic EG Spacer

Branching Molecule

Lipid Tail

g
Dicer substrate siRNA

5'   3'
3'   5'

Zipper pattern siRNA

5'   3'
3'   5' Phos

● RNA   ● DNA

h siRNA stability in OA-derived synovial fluid

Dicer substrate siRNA   Zipper pattern siRNA

hours 0 1 2 4 6 8 24   0 1 2 4 6 8 24

i si<(EG$_X$L)$_2$

si<(EG$_0$L)$_2$ 5' P

si<(EG$_6$L)$_2$ 5' P

si<(EG$_{18}$L)$_2$ 5' P

si<(EG$_{30}$L)$_2$ 5' P

j

Cy5 Fluorescence

EG$_0$
EG6
EG$_{18}$
EG$_{30}$
siRNA

lipoprotein bound   albumin bound   unbound

A01 A03 A05 A07 A09 A11 B01 B03 B05 B07 B09 B11 C01 C03 C05 C07 C09 C11 D01

siRNA   EG$_6$
EG$_0$ EG$_{18}$
        EG$_{30}$

% albumin bound

Figure 1. Albumin-binding lipophilic siRNA constructs for delivery to albumin-enriched PTOA joints.

A-D) Left knee mechanical loading was used to model PTOA in mice, followed by i.v. delivery of MSA-Cy5 (N = 5), PEG-Cy5 (N = 3), and Evans Blue (N = 4). Cy5 and Evans Blue were measured by fluorescence imaging 24 hrs after delivery. Representative images are shown. **A)** Procedural time is shown. **B)** MSA-Cy5 fluorescence in paired PTOA and healthy knees was measured. **C)** Ratio of MSA-Cy5 and PEG-Cy5 fluorescence in PTOA / healthy knees for each mouse treated with was measured. **D)** Evans Blue fluorescence in paired contralateral PTOA and healthy knees was measured.

E) Contralateral PTOA and healthy knees were excised 24 hours after Evans Blue delivery. Evans Blue was extracted and measured by densitometry (N = 3). Representative etissues and resulting extract is shown.

F) Chemical modifications of siRNA ribose, backbone, and terminus.

G) Schematic of synthetic dicer substrate siRNA and alternating 2'F-, 2'OMe-modified "Zipper" siRNA.

H) Representative gel electrophoresis of dicer substrate and zipper pattern siRNA sequences after incubation in synovial fluid collected from treatment-naïve OA patient.

I) Schematic representation of the si<$(EG_XL)_2$ series of zipper-modified siRNAs with divalent lipid end-modifications with variations in EG content.

J) Size exclusion chromatograph (left panel) of healthy human synovial fluid incubated with si<$(EG_XL)_2$ variants. The percentage of total si<$(EG_XL)_2$ detected in albumin fractions is shown (right panel). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

a Mouse Left Knee

Mechanical Loading, 3x / Week: 9N, 250 Cycles

Tissue analysis

1 3 5 7 days

Intravenous delivery: Cy5-si<(EG$_X$L)$_2$ series, Cy5-siRNA-Chol or Cy5-siRNA

b Ex vivo Cy5 imaging

si<(EG$_0$L)$_2$ si<(EG$_6$L)$_2$ si<(EG$_{18}$L)$_2$ si<(EG$_{30}$L)$_2$

Healthy OA   Healthy OA   Healthy OA   Healthy OA
knee knee   knee knee   knee knee   knee knee

Radiant Efficiency [Radiance/uW/cm^2] × 10^9

Color Scale: Min = 2.50e8   Max = 4.00e9

Radiant Efficiency (×10^9)

EG0 EG6 EG18 EG30

Healthy Knee   OA Knee

c Intravital Cy5 imaging

siRNA   siRNA-Chol   siRNA<(EG$_{18}$L)$_2$

Healthy OA   Healthy OA   Healthy OA
knee knee   knee knee   knee knee

Radiance (p/sec/cm3/sr) × 10^{10}

Color Scale: Min = 7.57e8   Max = 1.56e9

Average Radiance

siRNA
siRNA-Chol
siRNA<(EG$_{18}$L)$_2$

Healthy knee + - + - + -
OA knee   - + - + - +

d Cy5 / DAPI imaging in mouse knee

Cy5-siRNA   Cy5-siRNA-Chol   Cy5-siRNA<(EG$_{18}$L)$_2$

OA+ knee   OA+ knee   OA+ knee   Healthy knee

Synovium

Cartilage

siRNA   siRNA-Chol   siRNA<(EG$_{18}$L)$_2$

Synovium   Cartilage

Mean Fluor. Intensity   Mean Fluor. Intensity

e siRNA binding in patient synovial fluid

Patient synovial fluid

Cy5-siRNA:
Healthy
OA
RA

Cy5-siRNA<(EG$_{18}$L)$_2$:
Healthy
OA
RA

albumin bound

unbound

lipoprotein bound

Cy5 Fluorescence

Fractions: A01 A03 A05 A07 A09 A11 B01 B03 B05 B07 B09 B11 C01 C03 C05 C07 C09 C11 D01

f Albumin-bound siRNA in patient synovial fluid

AUC (Cy5 / fractions B11-C1)

Patient synovial fluid

Healthy
OA   Cy5-siRNA
RA

Healthy
OA   Cy5-siRNA<(EG$_{18}$L)$_2$
RA

Figure 2. Accumulation of the albumin-binding siRNA-lipid conjugate siRNA<(EG$_{18}$L)$_2$ in mouse PTOA knee joints following i.v. delivery.

A) Unilateral, left knee mouse loading protocol/timeline used to characterize si<(EG$_X$L)$_2$ series pharmacokinetics.

B) Ex-vivo IVIS images of knee joints and quantification of IVIS signal.

C) Representative intravital images after 24 hours of Cy5 fluorescence are shown (left panel). Quantification of average Cy5 fluorescence after 24 hours (right panel).

D) Cryosections of knees from treated mice were imaged for Cy5 and DAPI fluorescence in cartilage and synovial tissue.

E-F) Size exclusion chromatography was used to measure Cy5 fluorescence in fractions of Cy5-labelled siRNAs incubated with human synovial fluid taken from a healthy, osteoarthritis, and rheumatoid arthritis patient. A representative chromatograph is shown (**E**). Cy5 fluorescence within the albumin-containing fractions was quantitated as an area under the curve (AUC) analysis (**F**). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 3. Systemic delivery of siMMP13<(EG$_{18}$L)$_2$ decreases PTOA-induced hyperalgesia and joint degeneration. Bilateral knee loading [3X/wk, 5 wks] was used as a long-term PTOA mouse model for comparing therapeutic impact of siMMP13<(EG$_{18}$L)$_2$, Zilretta, CL-82198 and Marimastat.
A) Schematic timeline for knee loading and treatment.
B) Mechanical hyperalgesia was measured via algometer through day 35 (left panel). AUC of average Smalgo readings / time was assessed. N = 8.
C) Total MMP activity in mouse knees was measured using MMPsense 750 Fast (N = 8; left panel). mAbCII binding in knee joints was measured by fluorescence imaging (N = 8; middle panel). C2C fragments in mouse serum was measured by ELISA (N = 3; right panel).
D) Knee joints were stained with Toluidine Blue (upper panels, femoral condyles shown) and H&E (lower panels, synovium lining and meniscus shown). Asterisks indicate the following: healthy articular surface (white).; loss of articular surface (red); healthy synovial lining / meniscus (yellow); synovial thickening / meniscal calcification (black).
E) Joint cartilage damage was quantitated with the OARSI osteoarthritis cartilage histopathology assessment system (upper panel) and Degenerative Joint Disease (DJD) score (lower panel). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 4. Systemic delivery enables multi-joint accumulation of siMMP13<(EG$_{18}$L)$_2$, MMP13 knockdown and diminished arthritis progression in a mouse RA model.

A-B) Four days after K/BxN serum transfer, mice were treated with Cy5-siRNA<(EG$_{18}$L)$_2$, Cy5-siRNA-Chol, and Cy5-siRNA (1 mg/kg). Joints of forepaw, hindpaw, and knee were collected 24 hrs later, and assessed by IVIS imaging (**A**) or fluorescence microscopy (**B**). Representative images are shown. N = 6.

C) Relative *Mmp13* mRNA in forepaws, knees, and hindpaws of healthy, untreated wild-type mice, or in K/BxN serum recipients treated with si-Control<(EG$_{18}$L)$_2$ or siMMP13 <(EG$_{18}$L)$_2$. was measured by RT-qPCR. N = 6.

D-F) K/BxN serum recipients treated were treated with si-MMP13<(EG$_{18}$L)$_2$, methylprednisolone, or CL-82198. Schematic timeline for K/BxN serum transfer and treatment is shown (**D**). Images of hindpaws were collected on treatment day 10 (**E**) Representative images are shown. Ankle width change, clinical score, algometer ankle joint hyperalgesia, and severity index were measured for 10 days after treatment (**F**). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 5. Systemic siMMP13<(EG$_{18}$L)$_2$ treatment protects against cartilage / bone destruction and synovial inflammation in a mouse RA model. K/BxN serum recipient mice were treated with siMMP13<(EG$_{18}$L)$_2$, methylprednisolone, or CL-82198.

A-C) Histological analysis of toluidine blue-stained ankle cartilage sections (**A**, upper panels) and H&E-stained hindpaw sections (**A**, lower panels) were used for cartilage destruction scoring (**B**) and inflammation scoring (**C**), respectively. Asterisks indicate the following: yellow: articular surface; red: articular surface damage. Black: healthy synovium. White: synovial inflammation.

D-G) MicroCT-based **3D** renderings of hindpaws scans (D, representative images) were used to measure bone surface:volume ratio (**E**), bone mineral density (**F**) and bone fraction (**G**). White arrows in D show bone erosions *P < 0.05, **P < 0.01, ***P < 0.001,

Figure 6. MMP13 silencing upon systemic siMMP13<(EG$_{18}$L)$_2$ delivery in guinea pig ACL transection model of arthritis.

A) Timeline for ACL transection and siRNA delivery in Dunkin-Hartley guinea pigs is shown.

B-D) Cy5 fluorescence was measured ex vivo in healthy and ACLT knees (**B**) and organs (**C**) of guinea pigs by IVIS, and by fluorescence microscopy of sagittal ACLT knee cartilage / synovial cryosections **(D)** at 24 hrs after i.v. delivery of Cy5-siRNA or Cy5-siRNA<(EG$_{18}$L)$_2$ at 1 mg/kg. Representative images are shown. In D, Blue = DAPI, Red = Cy5 siRNA.

E) Guinea Pig *Mmp13* mRNA expression (ACLT/Healthy Knee) as determined by qRT-PCR.

F-G) MMP13 immunofluorescence in cartilage / synovial cryosections of ACLT knee joints that were untreated (left) or treated with siMMP13<(EG$_{18}$L)$_2$ (10 mg/kg, i.v.**)**. White asterisk: synovium; red asterisk; femoral condyle cartilage. Representative images are shown (**F**). MMP13 immunofluorescence intensity was quantitated using morphometric software (**G**). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Figure 1. Exploring different routes of delivery in mechanically-loaded PTOA model.

A-C) Subcutaneous (SubQ) delivery of Cy5-siRNA<(EG$_{18}$L)$_2$ (2 mg/kg) was studied in a TOA mouse model. Timeline for mechanical knee loading and treatment is shown (A). Intravital (B) and ex vivo (C) Cy5 fluorescence imaging was used to measure Cy5-siRNA<(EG$_{18}$L)$_2$ biodistribution to paired healthy and loaded knees and organs 24 hrs after delivery.N = 3.

D) Quantification of SubQ vs. intravenous (i.v.) delivery route organ biodistribution 24 hours after injection.

E-G) Intra-articular delivery of Cy5-siRNA<(EG$_{18}$L)$_2$ (1 mg/kg) was studied in a PTOA mouse model. Timeline for mechanical knee loading and treatment is shown (**E-F**). Ex vivo Cy5 fluorescence imaging was used to measure Cy5-siRNA<(EG$_{18}$L)$_2$ biodistribution to paired healthy and loaded knees and organs 48 hrs after delivery. N = 3. (**G**). Cryohistology of the loaded knee joints 48 hours after intra-articular injection with a focus on cartilage and synovial tissues. N = 3. *P < 0.05, **P < 0.01.

Extended Data Figure 2. MMP13 knockdown in PTOA joints by systemic delivery of siMMP13<(EG$_{18}$L)$_2$.
A) *Mmp13* mRNA was measured by RT-qPCR in healthy and PTOA knees of mice 5 days after i.v. delivery of siMMP13<(EG$_{18}$L)$_2$, siMMP13-Chol, or free siMMP (5 and 10 mg/kg, where indicated).
B) *Mmp13* mRNA was measured by RT-qPCR in PTOA mouse knees of mice 5 days after i.v., i.a., or subcutaneous siMMP13<(EG$_{18}$L)$_2$ delivery at the doses indicated.
C) *Mmp13* mRNA was measured by RT-qPCR in PTOA mouse knees of mice 5 days subcutaneous siMMP13<(EG$_{18}$L)$_2$ delivery at the doses indicated.
D-E) MMP13 IHC in PTOA knees (D), liver and kidneys (E), 5 days after i.v. delivery (10mg/kg) siMMP13<(EG$_{18}$L), siMMP13-Chol, free siMMP13, or no treatment.
F) Relative *Mmp13* mRNA expression in liver and kidneys of mice (relative to knee *Mmp13* levels).
G) Relative *Mmp13* mRNA in kidneys, 5 days post-injection with 10 mg/kg siMMP13<(EG$_{18}$L)$_2$, free siMMP13, or PBS.
*P < 0.05, **P < 0.01, ***P < 0.001

Extended Data Figure 3. siMMP13<(EG$_{18}$L)$_2$ demonstrates long-term retention and MMP13 silencing in PTOA knee joints.

A) Timeline for mouse left knee mechanical loading (3 times / week) and single treatment (day 7) with Cy5-siRNA<(EG$_{18}$L)$_2$ or siMMP13<(EG$_{18}$L)$_2$.

B-D) Cy5-siRNA was measured on day 30 post-treatment by confocal microscopy of knee joint cryosections (**B**, representative images), and at time points from 0-30 days post-treatment by longitudinal intravital Cy5 imaging (**C**). Semiquantitative analysis of the AUC (Cy5 fluorescence / days) was calculated (**D**) (n = 4).

E-F). Knee joint *Mmp13* was measured at the indicated time points by RT-qPCR (**E**) and IHC (**F**) in healthy or PTOA knee tissues from mice treated with or without siMMP13<(EG$_{18}$L)$_2$ collected at the indicated time points. N = 5. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig 4. Healthy and PTOA knee joint delivery and retention after intra-articular delivery over a 30-day time course.

A) Representative IVIS images of healthy and PTOA mouse knee joints over 30 days after a single 1 mg/kg intra-articular (IA) injection of Cy5-siRNA<(EG$_{18}$L)$_2$ or Cy5-siRNA.

B) Semiquantitative analysis of time-course fluorescent radiant efficiency within healthy and OA mouse knee joints over 30 days (N = 3) and AUC based on the fluorescence intensity profiles.

C) Semiquantitative analysis of time-course fraction of retention within healthy and OA mouse knee joints over 30 days and AUC based on the fraction of retention profiles.

D) Representative confocal microscopy of knee joint cryosections at the 30-day endpoint showing IA Cy5-siRNA<(EG$_{18}$L)2, and IA Cy5-siRNA in loaded, PTOA, knees with a specific focus on synovial and cartilage/meniscus tissues. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig 5. Systemic albumin-hitchhiking siMMP13<(EG$_{18}$L)$_2$ treatment provides whole knee joint protection by reducing ssteophyte formation and pathologic mineralization meanwhile by MMP13 protein and collagen degradation fragments.

A) MicroCT-based 3-dimensional (3D) renderings of meniscal/ectopic mineralization and osteophyte growth in healthy and PTOA mice treated with siMMP13<(EG$_{18}$L)$_2$ or siControl<(EG$_{18}$L)$_2$.

B) Measurements of femoral and tibial osteophyte size at largest outgrowth from normal cortical bone structure.

C) Ectopic mineralization [hydroxyapatite (HA)] in menisci and in the form of osteophytes was measured.

D-F) MMP13 IHC analyses of cartilage (D), meniscus (E), and synovium (F) and showing reduced MMP13 protein levels when treated with siMMP13<(EG$_{18}$L)$_2$.

G) C1,2C collagen 2 degradation fragment IHC. Samples here are from the PTOA therapeutic study in Main Figure 3. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig 6. Systemic siMMP13<(EG₁₈L)₂ delivery alters inflammatory gene expression profiles in mechanically-loaded PTOA joints.

A) qRT-PCR of IL-1B, TNFalpha, NGF, and P16INK4a (Cdkn2a) at days 5, 10, 20, and 30 after a single i.v. injection of 10 mg/kg siMMP13<(EG₁₈L)₂.

B) Unsupervised sorting of treatment groups as quantified by nanoString at day 10 after treatment with 10 mg/kg siMMP13<(EG₁₈L)₂. Gene expression is shown as high- (green) or low-expression (red) sorted vertically by differences between treatment groups.

C) Gene cluster expression changes at day 10 post-treatment. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

a Evans Blue extraction

b Evans Blue accumulation

c Cy5-labeled albumin accumulation

d Albumin transport-associated genes

e Biodistribution of Cy5-labeled siRNA<(EG₁₈L)₂ in K/BxN STA model

1. Right forepaw
2. Right knee
3. Right hindpaw
4. Heart
5. Lung
6. Spleen
7. Liver
8. Kidneys
9. Left forepaw
10. Left knee
11. Left hindpaw

f siRNA<(EG₁₈L)₂ accumulation

g Hindpaw — Knee

Extended Data Fig 7. Characterizing albumin-based drug delivery in the K/BxN STA model.

A-C) Evans Blue or Cy5-MSA was delivered i.v. to wild type (healthy) and K/BxN serum recipient mice 4 days after serum transfer. Joints of forepaws and hindpaws were used for Evans Blue extraction (**A**, representative images), measuring Evans Blue absorbance in extracts, or for visualization of Evans Blue by fluorescence microscopy in joint cryosections (**B**). MSA-Cy5 delivery to joints was assessed by intravital Cy5 fluorescence imaging (**C**).

D) qRT-PCR was used to measure relative gene expression of SPARC and Caveolin-1 in healthy and K/BxN arthritis hind paws.

E-G) Cy5-siRNA<(EG₁₈L)₂ (1 mg/kg, i.v.) was delivered to healthy and K/BxN serum recipient mice. Cy5 fluorescence in organs (**E**) and limb joints (**F**) were measured ex vivo at 24 hrs, and in knee and hindpaw cryosections by fluorescence microscopy (**G**)

*P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig 8. Systemic siMMP13<(EG$_{18}$L)$_2$ delivery silences MMP13 and protects cartilage in the multi-joint K/BxN STA model. K/BxN STA mice were treated with or without siMMP13<(EG$_{18}$L)$_2$ (10 mg/kg, i.v.), then assessed at day 10.

A) MMP13 IHC analysis of cartilage and synovium in multiple joints was used to assess MMP13 expression.

B) Fluorescent mAbCII was delivered to mice, then detected by ex vivo fluorescence imaging (IVIS) to measure relative cartilage damage in healthy and K/BxN STA mice (representative images shown).

C) MMPsense 750 Fast was delivered to mice, then detected by ex vivo fluorescence imaging (IVIS) used to measure pan-MMP activity.

*P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig 9. Systemic delivery of siMMP13<(EG₁₈L)₂ protects cartilage and bone while reducing inflammation in multiple joints of the K/BxN STA model.

A-B) Toluidine Blue stained histological knee and forepaw sections (**A**) were scored using the Cartilage Destruction Score (**B**). Hindpaw data is shown in Figure 5.

C-D) H&E-stained histological knee and forepaw sections (**C**) were scored using the Inflammation Score (**D**). Hindpaw data is shown in Figure 5.

E) Histological bone destruction score quantification of hindpaw, knee, and forepaw joint tissues. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig 10. Systemic siMMP13<(EG₁₈L)₂ delivery reduces cartilage degradation fragments and diminishes the inflammatory gene expression profile in the K/BxN STA model.

A) Relative serum C2C collagen 2 degradation fragment was measured on day 10 by ELISA.

B) C1,2C collagen 2 degradation fragment was assessed by IHC on day 10.

C-D) RNA harvested from K/BxN recipient mouse joints was assessed by nanoString nCounter analysis for the Mouse Inflammatory 294 Gene Expression panel. Clustering of gene expression changes [high- (green) or low- (red)] sorted vertically by differences between treatment groups (**C**). Treatment group-associated changes in gene enrichment clusters ion treatment day 10 (**D**). *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

a

b

c

Supplementary Figure 1. Albumin accumulates in cartilage and synovial tissues of mechanically loaded OA knees to a greater extent than Poly(ethylene glycol).

A) Representative sagittal cryosections (20X) of knee joints used for assessing MSA-Cy5 (N=5) and PEG-Cy5 (N=3). Fluorescent signal within cartilage and synovial compartments of PTOA mouse knee joints.

B) Fluorescent signal within cartilage and synovial compartments of PTOA and healthy mouse knee joints treated with MSA-Cy5.

C) Representative sagittal cryosections (20X) of knee joints assessing MSA-Cy5 and synovial blood vessels.

Representative Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

	Right knee (Healthy)	Left knee (OA)

Denuded hindlimb

1. **Harvest hindlimbs, 24 hrs after Evans Blue delivery**

Knee

2. **Knee dissection**

Extracted Evans Blue

3. **Extract Evans Blue from tissue, 55ºC in formamide**

$y = 0.0916x + 0.0084$
$R^2 = 0.9903$

4. **Measure absorbance of extract against an Evans Blue standard curve.**

Absorbance Signal Intensity vs *ng Evans Blue*

Supplementary Figure 2. Representative samples and processing protocol with standard curve and equation for Evans Blue ng/mg tissue quantification.

Supplementary Figure 3. Characterization of albumin delivery/transport in the mechanical loading PTOA model.

A) Unilateral (left knee loaded, right knee unloaded) mechanical loading protocol/timeline used and injection protocol for Evans Blue dye (N=4).

B) Intravital (IVIS) quantification of Evans Blue dye.

C) Representative cryohistology images (20X) of sagittal-sectioned knee joints 24 hours after intravenous Evans Blue injection (Blue = DAPI, Red = Evans Blue). On right is quantification of Evans Blue cryohistologyimages with a specific focus on cartilage and synovial tissue (N=3-6).

D) qRT-PCR of cartilage/synovial tissue mix for SPARC, FcRn, Caveolin-1 (N=3-4).

Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 4. Biodistribution of systemically delivered lipophilic Cy5-si<(EG$_X$L)$_2$ series in mechanically loaded animals.

A) Representative ex-vivo IVIS images for systemic Cy5-conjugated siRNA molecules.

B) Organ biodistribution quantification (N=3).

Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 5. Albumin-hitchhiking siRNA<$(EG_{18}L)_2$ reaches cartilage and synovial tissues within loaded, arthritic knee joints.

A) Confocal microscopy of knee joint tissues of mice injected with Cy5-conjugated siRNA molecules – siRNA, siChol, and siRNA<$(EG_{18}L)_2$.

B) Confocal microscopy of siRNA<$(EG_{18}L)_2$ (red) in loaded knees stained for blood vessels in the synovium/extra-articular tissues (green).

C) Microscope image quantification of Cy5-siRNA molecules in synovial and cartilage tissues within loaded, arthritic knee joints (N=3-6).

TD – Transmission Detector. Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 6. Albumin-hitchhiking siRNA<(EG$_{18}$L)$_2$ demonstrates preferential delivery to loaded, arthritic knee joints.

A) Unilateral mechanical knee loading protocol/timeline used and injection schedule for systemic delivery of Cy5-conjugated siRNA molecules.

B) Representative ex-vivo IVIS images.

C) Organ biodistribution representative images and quantification (N=5-7).

D) Representative IVIS images of cartilage and synovial tissues dissected out of arthritic and healthy knee joints and quantification of IVIS signal (N=3).

Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 7. siRNA<(EG$_{18}$L)$_2$, but not free siRNA, binds to albumin in human synovial fluid, with increased binding in specimens from OA- and RA-affected joints.

A and B are based on the SEC graph in main figure 1.

A) Western blot for human albumin in B12 and C1 synovial fluid fractions shows increased albumin in OA- and RA-derived specimens.

B) Fractions B12 and C7 were fractionated by native SDS-PAGE, assessing both Cy5 fluorescence (left panel) and total proteins (coomassie blue). Fraction C7 was used as a negative control, non-albumin containing fraction where unbound siRNA elutes. Red arrow indicates molecular weight of human albumin (66 kDa).

C) Gel electrophoresis mobility shift assay demonstrating tendency of siRNA<(EG$_{18}$L)$_2$ to bind/interact with albumin in a species-independent manner.

Mouse Sequence Name	Antisense Strand	Sense Strand
siMMP13.1 (Lead Sequence Candidate)	5'UUUCUCAUGAUGUCUAAGGUU3'	5'CCUUAGACAUCAUGAGAAAUU3'
siMMP13.1 (with 19-mer Zipper Chemisty) PHO: Phosphate, *: Phosphorothioate, f: 2'fluoro, Ome: 2'O-methyl	5'(PHO)(OMeU)*(fU)*(OMeU)(fC)(OMeU)(fC) (OMeA)(fU)(OMeG)(fA)(OMeU)(fG) (OMeU)(fC)(OMeU)(fA)(OMeA)*(fG)*(OMeG)3' MW = 6303 g/mole	5' (fC)*(OMeC)*(fU)(OMeU)(fA)(OMeG) (fA)(OMeC)(fA)(OMeU)(fC)(OMeA) (fU)(OMeG)(fA)(OMeG)(fA)*(OMeA)*(fA) 3' MW = 6263.1 g/mole
siMMP13.2	5'UCACUGUAGACUUCUUCAGUU3'	5'CUGAAGAAGUCUACAGUGAUU3'
siMMP13.3	5'UUUCUCGGAGUCUAUCAACUU3'	5'GUUGAUAGACUCCGAGAAAUU3'
siMMP13.4	5'AAACUCAAGAUCUUCCUCAUU3'	5'UGAGGAAGAUCUUGAGUUUUU3'
siMMP13.5	5'UUUCUCGGAGUCUAUCAACUU3'	5'GUUGAUAGACUCCGAGAAAUU3'
siMMP13.6	5'AAACUCAAGAUCUUCCUCAUU3'	5'UGAGGAAGAUCUUGAGUUUUU3'
siMMP13.7	5'UAAAGAUCAUGGUUUCUCCUU3'	5'GGAGAAACCAUGAUCUUUAUU3'
siControl (Targeting firefly luciferase)	5'UUCAUUAUCAGUGCAAUUG3'	5'CAAUUGCACUGAUAAUGAA3'
siLUC used as a siNEG sequence → Targets Firefly Luciferase (Not in the mouse genome) *For Fluorescent Cy5 Version, switch 5'PHO for 5'Cy5	5'(PHO)(OMeU)*(fU)*(OMeC)(fA)(OMeU)(fU)(OMeA)(fU)(OMeC)(fA)(OMeG)(fU)(OMeG)(fC)(OMeA)(fA)(OMeU)*(fU)*(OMeG)3' MW = 6287 g/mole	5'(fC)*(OMeA)*(fA)(OMeU)(fU)(OMeG)(fC)(OMeA)(fC)(OMeU)(fG)(OMeA)(fU)(OMeA)(fA)(OMeU)(fG)*(OMeA)*(fA)3' MW = 6264.1 g/mole

Guinea Pig Sequence Name	Antisense Strand	Sense Strand
siMMP13.1	5'UAUCUGUCCAGUAUUCAUCUU3'	5'GAUGAAUACUGGACAGAUAUU3'
siMMP13.2	5'UUAAUGUCAUCAUAUCUCCUU3'	5'GGAGAUAUGAUGACAUUAAUU3'
siMMP13.3 (Lead Sequence Candidate)	5'AUGCCCGCAAGAUUUACUGUU3'	5'CAGUAAAUCUUGCGGGCAUUU3'
siMMP13.3 (19-mer Zipper Chemistry used for Therapeutic Studies) PHO = Phosphate, * = Phosphorothioate, f = 2'fluoro, OMe = 2'O-methyl	5'(PHO)(OMeA)*(fU)*(OMeG)(fC)(OMeC)(fC)(OMeG)(fC)(OMeA)(fA)(OMeG)(fA)(OMeU)(fU)(OMeU)(fA)(OMeC)*(fU)*(OMeG)3' MW = 6206.9 g/mole 6344.2 g/mole	5'(fC)*(OMeA)*(fG)(OMeU)(fA)(OMeA)(fA)(OMeA)(fU)(OMeC)(fU)(OMeU)(fG)(OMeC)(fG)(OMeG)(fG)*(OMeC)*(fA)3' MW = 6344.2 g/mole
siMMP13.4	5'UUGCCUGUCACCUCUAAGCUU3'	5'GCUUAGAGGUGACAGGCAAUU3'

Supplementary Figure 8. Mouse and guinea pig siRNA sequences and modifications used herein.

Supplementary Figure 9. MMP13 siRNA sequence screening across species, carrier free activity, and stability.

A-C) *Mmp13* mRNA was measured in TNFα-stimulated murine chondrogenic ATDC5 cells by RT-qPCR following transfection of candidate murine MMP13 siRNA sequences (**A**); Lipofectamine-mediated delivery of zipper-modified relative to unmodified siMMP13 sequences (**B**); Carrier-free silencing activity of siMMP13<$(EG_{18}L)_2$ (**C**) (N=3-4).

D-E) *Mmp13* mRNA was measured in TNFα-stimulated primary guinea pig chondrocytes by RT-qPCR following lipofectamine-mediated delivery of commercially available guinea pig MMP13 siRNA sequences (**D**) or zipper-modified guinea pig siMMP13 (**E**).

F) Serum stability (60% FBS, fetal bovine serum) at 37°C of 'zipper' modified or Dicer substrate mouse siMMP13.

Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 10. Carrier-Free Mediated Cell Uptake in Chondrogenic Murine ATDC5 Cells.

A) Flow cytometry histograms of cellular uptake 2 hours after carrier-free treatment of Cy5-conjugated siRNA or siRNA$<$(EG$_{18}$L)$_2$ at a concentration of 100 nM in serum-free, serum-free + supplemented mouse serum albumin (10x, 1 μM), or serum-containing media. Data are quantified as fold change of siRNA$<$(EG$_{18}$L)$_2$/siRNA cell uptake in differing media conditions (N=3).

B) Quantification of cellular uptake by flow cytometry in ATDC5 cells treated with or without murine TNFα (20 ng/mL) (N=3).

C) Representative confocal microscopy images of ATDC5 cellular uptake (Cy5 fluorescence) after 4 hours of treatment with or without murine TNFα (20 ng/mL).

Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 11. Pharmacokinetic and biodistribution analyses of intra-venous vs. intra-articular routes.

A) Representative IVIS organ biodistribution images of all 3 groups at the 30-day endpoint.

B) AUC analysis for i.v. siRNA<$(EG_{18}L)_2$ (10 mg/kg), intra-articular siRNA (1 mg/kg), and Intra-articular siRNA<$(EG_{18}L)_2$ (1 mg/kg) throughout the entire 30-day intravital IVIS retention study (N=3-4).

C) Quantification of organ biodistribution from images in A (N=3-4).

D) PNA hybridization assay showing ng/mg antisense strand of the siRNA per mg of knee joint tissue at 24 hrs (i.v.) or 48 hrs (i.a.) (N=3).

Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 12. Biodistribution and pharmacokinetics of K/BxN STA model.
A-B) Cy5-siRNA<(EG$_{18}$L)$_2$, Cy5-siRNA-Chol, or Cy5-siRNA was delivered i.v. (1 mg/kg) to healthy and K/BxN STA mice, and assessed 24 hrs later. Intravital Cy5 measurements in healthy and K/BxN STA mice in each treatment group (A). Organ Cy5 biodistribution was measured ex vivo (B) (N=3 mice).
C) Longevity of Cy5-siRNA<(EG$_{18}$L)$_2$ retention in PTOA knees following single dose of Cy5-siRNA<(EG$_{18}$L)$_2$ (10mg/kg, i.v.) was assessed by longitudinal intravital imaging, and endpoint imaging of knees and organs done ex vivo on day 30 (N=3 mice).
Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Hindpaw

IL-6

IL-1b

TNF

COX-2

- Healthy joints, untreated
- K/BxN arthritic joints, untreated
- K/BxN arthritic joints, siMMP13<$(EG_{18}L)_2$

Knee

IL-6

IL-1b

TNF

COX-2

- Healthy joints, untreated
- K/BxN arthritic joints, untreated
- K/BxN arthritic joints, siMMP13<$(EG_{18}L)_2$

Supplementary Figure 13. qRT-PCR of genes of importance in rheumatoid arthritis in the K/BxN therapeutic study at the day 10 endpoint (N=5-7). Statistics markers: $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$.

Supplementary Figure 14. Further characterization of K/BxN therapeutic study.
A-B) Clinical score of forepaws (A) and hindpaws (B) (N=7 mice).
C) Percentage of mice exhibiting affliction of all paw joints following K/BxN serum transfer (N=7 mice).
D) Representative images showing inflammation / swelling in limbs of K/BxN serum recipients on treatment day 8 (10 days after serum transfer). Insets show higher power image of forepaws (upper inset) and hind paws (lower inset). Images were used for clinical scoring, with a maximal clinical score of 3 per paw, total score of 12 per mouse.

Supplementary Figure 15. Toluidine Blue and H&E of knee joint tissue for the K/BxN therapeutic study.

Supplementary Figure 16. Toluidine Blue of forepaw / wrist cartilage of K/BxN serum recipients. Panels on left show 8X magnification. Black rectangles in left panels are the areas shown at higher magnification (20X) in right panels. Yellow rectangles in right panels indicate the areas of higher magnification shown in extended data figures.

Supplementary Figure 17. H&E of forepaw/wrist tissue for the K/BxN therapeutic study.

Supplementary Figure 18. Bone loss and ankle joint assessments in K/BxN therapeutic study.
Top: 3D reconstructions of the calcaneus bone (used for microCT parameter quantification). Middle: lower magnification Tolluidine Blue of hindpaw/ankle cartilage. Bottom: lower magnification H&E of hindpaw/ankle tissue.

Supplementary Figure 19. Systemic toxicity analyses in K/BxN and PTOA model therapeutic studies.
A) Toxicology analyses in serum in the PTOA therapeutic study at the endpoint (N=4 mice).
B) Toxicology analyses in serum in the K/BxN therapeutic study at the endpoint (N=7 mice).
Statistics markers: *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Supplementary Figure 20. H&E-stained sections of organs (liver, kidney, lung, heart, and spleen) from mice enrolled in the different treatment groups in the PTOA and K/BxN therapeutic studies. Organs were harvested at the end of the study for analysis, and a healthy (no disease, no treatment) mouse is included as a control.

UV Elution Spectra

LC-MS Spectra

LC-MS Spectra (Zoom)

Supplementary Figure 21. LC-MS characterization of control sense siRNA sequence.

UV Elution Spectra

LC-MS Spectra

LC-MS Spectra (Zoom)

Supplementary Figure 22. LC-MS characterization of control sense – cholesterol modifier siRNA sequence.

UV Elution Spectra

LC-MS Spectra

Luc Sense (siNEG) - <(EG$_{18}$L)$_2$ Modifier

Theoretical MW = 6264.1 g/mole + 3027 g/mole = 9291.1 g/mole

Z=3 = 3096.6
(3096.6x3)+3 = 9292.8 g/mole

LC-MS Spectra (Zoom)

Supplementary Figure 23 LC-MS characterization of control sense – <(EG$_{18}$L)$_2$ modifier siRNA sequence.

Supplementary Figure 24. LC-MS characterization of control antisense – siRNA sequence.

Supplementary Figure 25. LC-MS characterization of control antisense – Cy5 modifier siRNA sequence.

UV Elution Spectra

LC-MS Spectra

MMP13 Antisense

Theoretical MW = 6303 g/mole

Z=3 = 2099.9
(2099.9x3)+3 = 6302.7 g/mole

LC-MS Spectra (Zoom)

Supplementary Figure 26. LC-MS characterization of mouse MMP13 antisense – siRNA sequence.

UV Elution Spectra

LC-MS Spectra

MMP13 Sense - Regular siRNA

Theoretical MW = 6263.1 g/mole

Z=3 = 2086.3
(2086.3x3)+3 = 6261.9 g/mole

LC-MS Spectra (Zoom)

Supplementary Figure 27. LC-MS characterization of mouse MMP13 sense – siRNA sequence.

MMP13 Sense - <(EG$_{18}$L)$_2$ Modifier

Theoretical MW = 6263.1 g/mole + 3027 g/mole = 9290.1 g/mole

Z=3 = 3096.6

(3096.3x3)+3 = 9291.9 g/mole

Supplementary Figure 28. LC-MS characterization of mouse MMP13 sense – <(EG$_{18}$L)$_2$ modifier siRNA sequence

Supplementary Figure 29. LC-MS characterization of guinea pig MMP13 sense siRNA sequence

UV Elution Spectra

RT: 0.00 - 34.98 SM: 5G

NL: 2.18E7
TIC MS
220221_GP_MMP13_S
_L2_10-
6_220221201435

NL: 2.18E7
TIC F: FTMS - p ESI
Full ms
[350.00-4000.00] MS
220221_GP_MMP13_S
_L2_10-
6_220221201435

NL: 2.41E-1
External channel 1
Analog
220221_GP_MMP13_S
_L2_10-
6_220221201435

LC-MS Spectra

220221_GP_MMP13_S_L2_10-6_220221201435#503-520 RT: 24.42-24.87 AV: 18 NL: 5.99E4
F: FTMS - p ESI Full ms [350.00-4000.00]

LC-MS Spectra (Zoom)

Supplementary Figure 30. LC-MS characterization of guinea pig MMP13 sense – <(EG$_{18}$L)$_2$ modifier siRNA sequence

Supplementary Figure 31. LC-MS Characterization of Guinea Pig MMP13 Antisense – siRNA Sequence.