A TOPBP1 Allele Causing Male Infertility Uncouples XY Silencing Dynamics From Sex Body Formation

Carolline F. R. Ascencao[1*], Jennie R. Sims[1*], Alexis Dziubek[1], William Comstock[1], Elizabeth A. Fogarty[1], Jumana Badar[1], Raimundo Freire[2,3,4], Andrew Grimson[1], Robert S. Weiss[5], Paula E. Cohen[5,#] and Marcus Smolka[1,#]

1. Department of Molecular Biology and Genetics, Cornell University, United States;
2. Fundación Canaria del Instituto de Investigación Sanitaria de Canarias (FIISC), Unidad de Investigación, Hospital Universitario de Canarias, La Laguna, Santa Cruz de Tenerife, Spain;
3. Instituto de Tecnologías Biomédicas, Universidad de La Laguna, 38200 La Laguna, Spain;
4. Universidad Fernando Pessoa Canarias, Las Palmas de Gran Canaria, Spain;
5. Department of Biomedical Sciences, Cornell University, United States.

* Contributed equally

\# Co-corresponding authors

ABSTRACT

Meiotic sex chromosome inactivation (MSCI) is a critical feature of meiotic prophase I progression in males. While the ATR kinase and its activator TOPBP1 are key drivers of MSCI within the specialized sex body (SB) domain of the nucleus, how they promote silencing remains unclear given their multifaceted meiotic functions that also include DNA repair, chromosome synapsis and SB formation. Here we report a novel mutant mouse harboring mutations in the TOPBP1-BRCT5 domain. *Topbp1$^{B5/B5}$* males are infertile, with impaired MSCI despite displaying grossly normal events of early prophase I, including synapsis and SB formation. Specific ATR-dependent events are disrupted including phosphorylation and localization of the RNA:DNA helicase Senataxin. *Topbp1$^{B5/B5}$* spermatocytes initiate, but cannot maintain ongoing, MSCI. These findings reveal a non-canonical role for the ATR-TOPBP1 signaling axis in MSCI dynamics at advanced stages in pachynema and establish the first mouse mutant that separates ATR signaling and MSCI from SB formation.

Introduction

During Prophase I, the SPO11 nuclease and its cofactors induce programmed DNA double-strand breaks (DSBs) that are then recognized by the DNA damage response (DDR) machinery to promote recombination between homologous chromosomes[1–4]. Proper chromosome synapsis achieved through the formation of the proteinaceous synaptonemal complex, together with homologous recombination (HR)-mediated DNA repair[5] are critical for the formation of crossovers that ensure the correct segregation of chromosomes and the formation of healthy and genetically diverse haploid gametes[6]. Chromosomes that fail to synapse as prophase I progresses trigger a process referred to as meiotic silencing of unsynapsed chromatin (MSUC)[7–10] to silence genes at unsynapsed regions. In the heterogametic sex (male mammals), the X and Y chromosomes pose a unique challenge for meiotic progression since they bear homology only at the pseudoautosomal region (PAR). The non-homologous arms of the sex chromosomes remain unsynapsed and must therefore undergo a sex-chromosome specific manifestation of MSUC, termed meiotic sex-chromosome inactivation (MSCI)[8,11–13]. MSCI is critical for normal prophase I progression through two complementary mechanisms, the silencing of toxic Y-linked genes, such as *Zfy1* and *Zfy2*, that enforce the pachytene checkpoint[13,14] and through the sequestration of the DDR machinery at the X and Y chromosomes, away from the autosomes, during early pachynema[10].

The apical serine-threonine kinase Ataxia Telangiectasia mutated and Rad-3 related (ATR), is a master regulator of DNA repair, checkpoints and silencing during prophase I in spermatocytes. In response to DSBs and asynapsis, ATR activation promotes a range of downstream effects, including recombinational DNA repair, crossing over, chromosome synapsis, cell cycle arrest and potentially apoptosis[5,15–18]. During leptonema and zygonema, shortly after DSB formation, ATR and its cofactor ATRIP are recruited to RPA-coated regions of single strand DNA (ssDNA) that accumulate upon 5'-3' resection of both ends of DSBs[19]. ATR activation requires recruitment of TOPBP1 (topoisomerase 2 binding protein 1), a multi-BRCT (BRCA C-terminus motif) domain protein that stimulates ATR kinase activity through its ATR-activation domain (AAD)[19–21]. In addition to activating ATR, TOPBP1 also serves as a scaffold for a range of DDR factors, interacting with, and often recruiting them via its multiple BRCT domains[22–29]. TOPBP1 is

composed of nine BRCT domains, which are protein-interacting modules that typically recognize

phosphorylated motifs[22,30,31]. Through the recognition of phosphoproteins, TOPBP1 is able to

assemble multisubunit complexes to promote discrete pathways[5,22–29,32–34]. TOPBP1 interacts with

the C-terminal tail of RAD9, a component of the 9-1-1 PCNA-like clamp that is loaded at 5'

recessed junctions adjacent to DSBs[35]. The 9-1-1-TOPBP1 complex is essential for canonical ATR

signaling during prophase I[32–34]. Interaction of a protein with TOPBP1 may facilitate its

phosphorylation by ATR[19], suggesting a potential role for TOPBP1 in the control of ATR substrate

selection[19]. How the 9-1-1-TOPBP1-ATR signaling axis and the different TOPBP1 interactions

coordinate DNA repair, checkpoints and silencing during meiosis remains incompletely

understood.

As cells progress from zygonema into pachynema, ATR and TOPBP1 localize to the unsynapsed

axes of the X and Y chromosomes[36–38], leading to phosphorylation of the histone variant H2AX

on serine 139 (γH2AX), a major hallmark of MSCI[16,18]. During establishment of MSCI, a phase

separated structure termed the sex body is formed[39], allowing the confinement of ATR signaling,

DDR factors and silencing to the X and Y[10,11] as part of a checkpoint that induces cell death if

DDR proteins aberrantly accumulate and remain on autosomes[10]. Recruitment of ATR and

TOPBP1 to unsynapsed regions of the XY requires a distinct set of factors compared to their mode

of recruitment to autosomal DSB sites mentioned above, and involves factors such as BRCA1 and

HORMAD[40,41]. Activated ATR phosphorylates H2AX at the unsynapsed cores of the X and Y

chromosomes, a signaling that is propagated to chromatin loops of the X and Y, via a feed-forward

process mediated by recruitment of the MDC1 adaptor, which further recruits and mobilizes

additional TOPBP1-ATR complexes, therefore spreading ATR signaling to promote the broad

chromosome-wide silencing required for MSCI[42]. It has also been proposed that the initiation of

MSCI sequesters DDR proteins from autosomes to the X and Y chromosomes to prevent excessive

DDR signaling at autosomes from activating cellular checkpoints that can stop meiotic

progression[10].

Despite mounting evidence pointing to the importance of ATR and TOPBP1 for MSCI, the precise

mechanisms by which they promote sex body formation and XY silencing remain unknown.

Moreover, it remains unclear how these two processes are spatiotemporally coordinated and how

ATR and TOPBP1 mediate such coordination. Since both proteins are essential for organismal

viability[43–49], conditional knockouts[16,18,34] or hypomorphic[15] models have been used to explore

their roles during prophase I in spermatocytes. However, given the strong pleiotropic effects in

these models, especially in DSB repair, synapsis, MSCI and sex body formation, it is difficult to

dissect the distinct molecular mechanisms involved, and to untangle direct versus indirect effects.

Here we present a separation-of-function mouse mutant that deconvolutes TOPBP1-dependent

ATR signaling in male meiosis. We generated mice bearing multiple mutations in BRCT-domain

5 (*Topbp1^B5/B5* mice) that are viable and grossly indistinguishable from wild-type littermates; yet,

the males are sterile, having reduced testes size, reduced seminiferous tubule cellularity, and a

complete loss of sperm. Strikingly, while *Topbp1^B5/B5* spermatocytes fail to progress into diplotene,

they display largely normal chromosome synapsis, sex body formation, recruitment of DDR

proteins to the X and Y, and DNA repair during prophase I, in sharp contrast to previous models

of TOPBP1 or ATR impairment[18,34]. Single-cell RNA sequencing data showed that while MSCI

is initiated in *Topbp1^B5/B5*, the dynamics of silencing progression and reinforcement is defective,

which is accompanied by a defect in the localization of the RNA:DNA helicase Senataxin to

chromatin loops of the XY chromosomes. We propose that the *Topbp1^B5/B5* is a separation of

function mutant that allows the untangling of XY silencing from sex body formation and DDR

recruitment to the XY, representing a unique model to study the establishment, maintenance,

reinforcement and progression of MSCI.

Results

A TOPBP1 mutant separating its role in fertility from organismal viability.

Topbp1 knockout mice exhibit strong defects early in embryonic development, reaching blastocyst stage but not progressing beyond E.8, with embryos likely dying at the preimplantation stage[48]. In the context of meiosis, conditional knockout of *Topbp1* in spermatocytes leads to pleiotropic effects, including defects in chromosome synapsis, impaired recruitment of DDR factors to XY chromosomes, defects in condensation of the XY chromosomes, abnormal formation of the sex body, lack of γH2AX spreading to chromatin loops, all of which contribute to a strong MSCI defect as indicated by the complete absence of downstream markers of MSCI, such as USP7, H3K9me3, poly-ubiquitination and sumoylation[5,34]. The availability of a separation of function mutant for *Topbp1* is therefore needed to dissect its distinct roles in development, organismal maintenance, and multiple meiotic processes such as DNA repair, silencing and sex body formation. To generate a separation of function mouse model, we inserted eight charge reversal point mutations in BRCT domain 5 of TOPBP1 (hereafter referred to as *Topbp1$^{B5/B5}$*) CRISPR/Cas9 (Figure 1A). After validating the point mutations through Sanger sequencing, (Supplemental Figure 1), we found that *Topbp1$^{B5/B5}$* mice were viable, with no difference in body mass when compared to *WT* littermates (Figures 1B and 1C), and no sensitivity to Ionizing Radiation (IR) (Figures 1D). Strikingly, *Topbp1$^{B5/B5}$* mice displayed male-specific infertility (Figure 1E), with a three-fold reduction in testis size (Figures 1F and 1G) and complete lack of spermatozoa (Figure 1H). H&E-stained histological testis sections revealed mainly spermatogonia and spermatocytes within the seminiferous epithelium (Figure 1I) together with an increased number of TUNEL-positive spermatocytes in seminiferous tubules (Figure 1J and 1K). Cytological evaluation of surface-spread spermatocytes from *Topbp1$^{B5/B5}$* revealed the presence of meiotic prophase I stages from leptonema to pachynema but a total absence of diplonema-staged spermatocytes (Figure 1L). Moreover, the staining of the synaptonemal complex proteins SYCP1 and SYCP3 revealed normal pachytene entry and no gross defects in chromosome synapsis, distinct from previous models of ATR and TOPBP1 conditional depletion[18,34]. Furthermore, unlike reported DDR CKO (conditional knockouts) models such as *Rad1* and *Brca1*[10,29,36], *Topbp1$^{B5/B5}$* pachytene spermatocytes reach mid-pachynema, as demonstrated by the accumulation of signal for H1t[50] (Figure 1L).

***Topbp1^B5/B5* MEFs display no detectable DDR defects.**

To assess the possibility of a somatic phenotype we derived mouse embryonic fibroblasts (MEFs)

from *Topbp1^B5/B5* and wild-type littermates at embryonic day E13.5. Consistent with the observed

organismal viability and the lack of IR sensitivity, *Topbp1^B5/B5* MEFs showed no sensitivity to

hydroxyurea (replication stress) or phleomycin (DSBs) in a long-term cell survival assay (Figure

2A-D and Supplemental Figure 2). Genotoxic stress activates the apical kinases ATR, ATM and

DNA-PKcs[51–53] to trigger a signaling cascade that promotes DNA repair and cell cycle arrest via

activation of the downstream checkpoint kinases CHK1 and CHK2[54–56]. Similar to wild-type

MEFs, *Topbp1^B5/B5*MEFs were able to activate DDR signaling responses when challenged with

hydroxyurea and phleomycin as demonstrated by the induction of established damage-induced

phosphorylation of CHK1, CHK2, RPA and KAP1 (Figure 2E). In addition, *Topbp1^B5/B5* MEFs

were able to recruit MDC1 to γH2AX-marked DSB foci when subjected to IR (Figure 2F) and

showed no increased number of micronuclei, a known marker of defective DNA damage

responses[57], when compared to *Topbp1^+/+* MEFs (Supplemental Figure 3A-B).

The BRCT-5 domain of TOPBP1 is known to interact with the DDR factors 53BP1[22,24,28], MDC1[58]

and BLM[27] through phospho-protein binding modules. To investigate which, if any, of these

interactions are disrupted upon mutating the eight BRCT5 residues, we ectopically expressed Flag-

TOPBP1-WT or Flag-TOPBP1-B5 in HEK293T cells. We found that binding of Flag-TOPBP1-

B5 to BLM and 53BP1 was impaired, as expected (Supplemental Figure 4A-B). Moreover, we

noticed a 2-fold reduction in protein levels of Flag-TOPBP1-B5 compared to Flag-TOPBP1-WT,

which could be explained by the loss of interaction with BLM (that was proposed to lead to protein

stabilization[59,60]), or by protein misfolding caused by the 8 eight K to E/D mutations. In addition,

the reduction in protein levels was detected on MEFs (Supplemental Figure 5). In either case, the

results presented here show that the TOPBP1-B5 mutant offers a unique model to separate roles

of TOPBP1 in male meiosis from the canonical DDR functions of TOPBP1 in somatic cells.

***Topbp1^B5/B5* spermatocytes display normal markers of canonical ATR signaling, chromosome**

synapsis, DNA repair, sex body formation, and DDR protein localization at the X and Y

TOPBP1 and ATR play multiple roles in spermatocytes during prophase I. Mice conditionally

depleted for TOPBP1[34] or ATR[16,18] display severe defects in chromosome synapsis, DNA DSB

repair, sex body formation and MSCI, as well as impaired recruitment of DDR factors to the XY[5].

Strikingly, analysis of *Topbp1*[B5/B5] spermatocytes via meiotic spreads revealed normal repair of

DNA DSBs, with γH2AX staining grossly unchanged at pachynema, being confined only to the

XY chromosomes and being excluded from the autosomes (Figure 3A-B). RAD51 localized only

to the X and Y chromosomes during mid-pachytene (Figure 3C-D), indicating normal DSB repair

at the autosomes. We were unable to detect chromosome synapsis abnormalities in *Topbp1*[B5/B5]

spermatocytes, as *Topbp1*[B5/B5] mutant spermatocytes transit from zygotene into pachytene with

normal patterns of HORMAD1 and HORMAD2 localization (Supplemental Figure 6A-B) and

with SYCP1 overlapping with SYCP3 on chromosome cores from all autosomal chromosomes

during pachynema (Figures 1L and 3E). Since ATR orthologs regulate crossing over in budding

yeast [1] and in *Drosophila melanogaster* [61], we also investigated the localization of factors involved

in regulating DNA crossovers, including MLH1 and MLH3. In *Topbp1*[B5/B5] spermatocytes, while

the number of MLH1 foci were significantly increased, the number of MLH3 foci did not differ

significantly (Supplemental Figure 7A and 7B). Due to the lack of diplotene cells, and any other

stage after pachynema, we were not able to test whether *Topbp1*[B5/B5] display defects in crossing

over.

The sex body appeared normal in its shape and was normally formed in *Topbp1*[B5/B5] spermatocytes

(Figures 3A and Supplemental Figures 8 and 9A), exhibiting only subtle defects/delays in the

spreading of the γH2AX signal on the PAR and pericentromeric regions (Supplemental Figure 9B-

G). Although the defect was subtle in its severity, it accounted for 56% of all γH2AX-stained mid-

pachytene cells. Importantly, TOPBP1 was normally localized to X and Y chromosomes during

prophase I (Supplemental Figure 10 and Figure 3F-H). Similarly, localization of the TOPBP1

interactors ATR, BRCA1, MDC1, and 53BP1 in *Topbp1*[B5/B5] spermatocytes spreads was also

indistinguishable from that observed in *Topbp1*[+/+] spermatocytes (Figures 4A-F and Supplemental

Figures 11). Markers of ATR signaling were also mostly normal, as measured by its canonical

targets pMDC1 T4 (Supplemental Figure 6C-D), pCHK1 S345 (Supplemental Figure 6E-F), and

pHORMAD2 (Supplemental Figure 6G-H). Notably, we did observe that phosphorylation of

CHK1 on S317 was significantly decreased in *Topbp1*[B5/B5] when compared to *Topbp1*[+/+]

spermatocytes (Figure 4G-H). However, since *Chk1* CKO spermatocytes complete prophase I and

differentiate into spermatozoa, with only minor defects such a delay in the removal of γH2AX

from autosomes[62], the observed defect in Chk1 S317 phosphorylation is unlikely to be the cause

of the infertility observed in *Topbp1* B5 mutants. Overall, as summarized in Figure 4I, *Topbp1*$^{B5/B5}$

spermatocytes appear to progress normally through early stages of prophase I up until the end of

pachynema as demonstrated by largely normal localization of markers for DNA repair,

chromosome synapsis and ATR signaling. These findings are surprising because the lack of sex

body formation, synapsis defects or unrepaired DSBs, which are the expected causes of the drastic

loss of diplotene cells and the lack of spermatozoa, were not observed.

Defective phosphorylation and XY localization of the RNA:DNA helicase SETX in

***Topbp1*$^{B5/B5}$ spermatocytes**

With the exception of CHK1 phosphorylation at serine 317, the analysis of other canonical markers

of ATR signaling on meiotic prophase I spreads did not reveal obvious defects in their distribution

or intensity at the XY body (Figure 4G-H and Supplemental Figure 6C-H). Since altered CHK1

regulation is unlikely to be the cause of the drastic defect in meiotic progression observed in

Topbp1$^{B5/B5}$ males, we investigated whether other branches of ATR signaling were altered in

Topbp1$^{B5/B5}$ spermatocytes by performing unbiased quantitative phosphoproteomics based on

TMT (Tandem Mass Tag[63]) labeling of whole testes. Following a similar approach previously used

by our group[64], we analyzed the phosphoproteome of *Topbp1*$^{B5/B5}$ and *Topbp1*$^{+/+}$ testes, and then

compared the results to our previously reported dataset comparing the phosphoproteome of testes

from mice treated with vehicle or the ATR inhibitor AZ20[64] (Figure 5A). The resulting plot

revealed that ATR-dependent signaling is not drastically impaired in *Topbp1*$^{B5/B5}$ testes, as

opposed to the marked impairment of ATR signaling previously observed in testes of *Rad1* CKO

mice[64]. This finding is in agreement with the results from meiotic spreads of *Topbp1*$^{B5/B5}$ showing

no detectable defects in canonical markers of ATR signaling described above. Nonetheless, our

phosphoproteomic analysis did reveal phosphorylation sites mildly disrupted in *Topbp1*$^{B5/B5}$ testes

compared to *Topbp1*$^{+/+}$ testes. In particular, we noticed that several phosphorylation sites in the

RNA:DNA helicase, Senataxin (SETX)[65], including a phosphorylation in the preferred motif for

ATR phosphorylation (S/T-Q), were down-regulated in *Topbp1*$^{B5/B5}$ mice (Figure 5B).

Interestingly, SETX was previously associated with meiotic ATR functions[66,67] and found to be

indispensable for MSCI[66,67]. Moreover, we recently reported that mice treated with the ATR
inhibitor AZ20 or lacking *Rad1* display reduced phosphorylation of SETX at S/T-Q site and SETX
mislocalization during pachynema[64]. Based on our findings, we propose that TOPBP1 regulates
SETX distribution and/or function during meiosis, and that defects in meiotic progression and
fertility observed in *Topbp1[B5/B5]* mice might be associated with SETX dysfunction. Consistent with
this possibility, we found that pachytene spermatocytes from *Topbp1[B5/B5]* display significantly
decreased spreading of SETX to XY chromatin loops while still displayed SETX at the unsynapsed
axes of the X and Y chromosomes (Figures 5C-F). Overall, these findings reveal that *Topbp1[B5/B5]*
spermatocytes display largely normal progression through mid-pachynema, as demonstrated by
normal distribution of a range of markers of meiotic progression, including canonical markers of
ATR signaling. However, in-depth phosphoproteomic and imaging analyses identify specific
defects in the regulation of SETX, a target of ATR signaling during prophase I[29,64] and a key factor
required for MSCI[66,67].

***Topbp1[B5/B5]* spermatocytes initiate MSCI but fail to promote full XY silencing**
During mid-pachynema, spermatocytes that fail to properly silence the X and Y chromosomes
arrest and trigger apoptosis-induced cell death[9–11,68]. In mice, the MSCI process initiates in
leptonema[69], and during early pachynema key events occur at the XY chromosomes, such as
exclusion of RNA polymerase 2 (RNA Pol2), recruitment of DDR proteins and chromatin
remodelers and establishment of heterochromatin marks[10,70] to maintain the active silencing of the
X and Y chromosomes[17]. This process leads to the formation of a membrane-less phase separated
structure termed the sex body[39,71]. Spermatocytes from *Topbp1[B5/B5]* males formed a sex body of
grossly normal appearance (Supplemental Figure 8), with undisrupted patterns of the sex body
markers CHD4, SUMO, and USP7 (Supplemental Figure 12). *Topbp1[B5/B5]* pachytene
spermatocytes also display normal patterns of a range of chromatin marks, including H3K9ac,
H3K9me3, H3K27ac, H3K36me3, H3K4me3, H4K16ac, H4ac, H2AK116ub, H3K4me1, as well
as proper exclusion of RNA Pol2 from the sex body (Supplemental Figures 13-17). Collectively
the evidence presented herein shows that *Topbp1[B5/B5]* mutants are able to form grossly normal XY
bodies, with proper localization of over 28 markers, the severe loss of diplotene cells and the
reduction of SETX at the chromatin loops of the X and Y lead us to speculate that *Topbp1[B5/B5]*
spermatocytes may still have a defective MSCI. To investigate potential MSCI defects, we

employed single cell RNA sequencing (sc-RNA-seq) in germ cells (Figure 6A) using a similar approach recently used to follow spermatogenesis progression and to evaluate MSCI in mammals[69,72,73]. Using the 10X platform we performed sc-RNA-seq on a germ cell-enriched population of cells extracted from adult *Topbp1*[+/+] and *Topbp1*[B5/B5] testes (Figure 6A). The data were analyzed as previously described[69,73] using signature genes as markers of different stages in spermatogenesis such as, *Sal4* and *Dmrt1* for spermatogonia, *Dazl* for early spermatocytes, *Id4*, *Sycp3* and *Shcbp1l* for late spermatocytes, *Acrv1* for round spermatids, and *Oaz3* and *Prm2* for elongated spermatids (Supplemental Figures 18 and 19). *Col1a2*, *Acta2*, *Vcam1*, *Lnsl3*, *laptm5*, *Hbb-bt*, *Ptgds* and *Wt1* were used as markers of somatic cells (Supplemental Figure 20). Analysis of the testicular transcriptome of *Topbp1*[+/+] males revealed 47 sub-clusters of cells covering spermatogonia, spermatocytes, spermatids, and somatic cells (Figure 6B-D and Supplemental Figures 21-22). In sharp contrast, analysis of *Topbp1*[B5/B5] germ cell population revealed only somatic, spermatogonia and the initial populations of spermatocytes (Figure 6D). This result corroborates the H&E-stained testes histological sections (Figure 1I) and meiotic spreads (Figure 1L). Importantly, Pearson correlation values from all RNA reads between cell groups separated by cluster and genotype, demonstrated the similarity of spermatocytes from *Topbp1*[+/+] and *Topbp1*[B5/B5] (Supplemental Figure 23). As shown in Figure 6E, we were able to monitor the dynamics of X chromosome silencing in the early-stage spermatocytes and compare the results between RNA from *Topbp1*[+/+] and *Topbp1*[B5/B5] males. Strikingly, *Topbp1*[B5/B5] early-stage spermatocytes could initiate MSCI and promote robust, albeit incomplete, X chromosome silencing. Both X and Y chromosomes showed increased gene expression levels in the last spermatocyte stage captured in *Topbp1*[B5/B5] males when compared to RNA from wild type males (Figure 6E and Supplemental Figure 24). Moreover, although certain X-genes from *Topbp1*[B5/B5] pachytene cells consistently demonstrated defects in silencing by being expressed in numerous cells, other genes were only expressed in a small number of cells. This highlights the non-uniformity of the MSCI defect in all pachytene cells (Supplemental Figure 24).

MSCI is a dynamic process that involves the sequestration of DDR factors from the autosomes to the X and Y chromosomes as cells enter pachynema[10,17] as well as the inactivation of specific X and Y genes that lead to cell death if expressed at this stage (the so called "killer genes")[13,14]. Similar to previous reports based on mutants or treatments that impaired MSCI[16–18,29,34,74,75],

transcriptomics profiles from *Topbp1*[B5/B5] testicular cells showed an increased number of stage 3

spermatocytes (SP3 - pachytene) expressing the killer genes *Zfy1* and *Zfy2* compared to *Topbp1*[+/+]

(Supplemental Figures 25-26). Other genes typically used to illustrate MSCI defects, such as

Kdm6a, *lamp2*, *Zfx*, *Uba1y* and *Rhox13* were also expressed in a higher number of SP3 cells in

Topbp1[B5/B5] (Supplemental Figures 25-26). In the case of *Scml2*, *Topbp1*[B5/B5] cells not only

displayed an increased number of SP3 (pachytene) cells expressing it but also displayed an

increase in expression levels in these cells compared to *Topbp1*[+/+] testes (Supplemental Figures

25-26). In summary, various Y and X genes that had previously been shown to be expressed in

other MSCI-defective mutants were found de-repressed in *Topbp1*[B5/B5] spermatocytes.

Detailed analysis of the scRNA-seq data for the X-linked genes monitored during the early

spermatocyte stages revealed important differences in the silencing dynamics of these X genes

between *Topbp1*[B5/B5] and *Topbp1*[+/+] spermatocytes. 233 out of the roughly 700 genes present on

the X chromosome had reads detected from pre-leptotene to spermatocyte 3 clusters for both

Topbp1[+/+] and *Topbp1*[B5/B5] cells, and therefore were used during the downstream analysis. As

shown in Figure 7A, we clustered the X-linked genes into 3 distinct categories based on the change

of RNA level between the stages: reduced, unaltered, or increased silencing (Figures 7A). SP1

(spermatocyte 1) was defined as leptotene using the gene marker *Gm960*[76] (Supplemental Figure

27), and SP2 (spermatocyte 2) as zygotene due to its profile of low expression of *Gm960* and high

expression of *Dazl*. SP3 (spermatocyte 3) was defined as pachytene due to its lower expression of

Dazl and increased expression of *Id4* and *Shcbp1l*. When comparing the distribution of genes in

these clusters between *Topbp1*[B5/B5] and *Topbp1*[+/+] spermatocytes (RNA level normalized to pre-

leptotene stage) (Figure 7B), we observed no major differences in the pre-leptotene to

spermatocyte 1 (PL to SP1) and in the spermatocyte 1 to spermatocyte 2 (SP1 to SP2) transitions.

In contrast, when comparing the spermatocyte 2 to spermatocyte 3 (SP2 to SP3) transition between

Topbp1[B5/B5] and *Topbp1*[+/+] spermatocytes we noticed a major difference in the distribution of the

clusters, with 74 genes in *Topbp1*[B5/B5] versus 4 in *Topbp1*[+/+] exhibiting reduced silencing (Figure

7B). Moreover, while 134 genes showed unaltered silencing, and only 24 increased silencing in

Topbp1[B5/B5] spermatocytes during the SP2 to SP3 transition, 43 genes showed unaltered silencing

and 185 increased silencing in *Topbp1*[+/+] (Figure 7B). *Xiap*, *Zc3h12b* and *Cetn2* are examples of

X-linked genes displaying altered silencing behaviors in *Topbp1*[B5/B5] spermatocytes (Figure 7C).

The difference of expression between *Topbp1*$^{+/+}$and *Topbp1*$^{B5/B5}$ was markedly higher in the SP2

to SP3 transition compared to the other transitions (Figure 7D-E) and was used to further split

genes into 3 categories based on the severity of the silencing defect in *Topbp1*$^{B5/B5}$: no defect (13

genes), mild (45 genes) or strong defect (170 genes) (Figure 7F). Notably, the severity of the

silencing defect of a gene had some correlation with its RNA level in the pre-leptotene stage, with

highly expressed genes having a higher tendency to have a more severe silencing defect (Figure

7G and Supplemental Figure 28). Taken together, these data characterize the specific silencing

defect in *Topbp1*$^{B5/B5}$ spermatocytes and point to a specific role for TOPBP1 in ensuring proper

silencing dynamics after an initial wave of MSCI, likely through later waves of silencing

reinforcement. Our data is consistent with the notion that silencing of the X and Y chromosomes

is a dynamic process that needs active and continuous engagement by the ATR-TOPBP1 signaling

axis. Since the majority of the mouse models of male infertility accumulate pleiotropic defects,

with disrupted MSCI and absence of sex body, the *Topbp1*$^{B5/B5}$ mouse reported here provides a

unique model of DDR impairment in which MSCI can be uncoupled from sex body formation

(Figure 8).

Discussion

In male meiosis I, DDR factors such as ATR, TOPBP1, BRCA1, and the 9-1-1 complex play crucial roles in DNA repair, chromosome synapsis, recombination, sex body formation, and silencing[5,15,16,18,29,34,36,40]. Conditional depletion of these factors results in pleiotropic phenotypes from compound effects in multiple processes, with cells ultimately undergoing apoptotic-induced cell death during the pachytene checkpoint. Here, we report a mutant mouse model capable of deconvoluting TOPBP1's roles during meiosis I in males, separating its role in silencing from its roles in DNA repair, synapsis and checkpoint signaling (Figure 8). While $Topbp1^{B5/B5}$ spermatocytes initiate XY silencing with similar dynamics as observed in $Topbp1^{+/+}$, these cells fail to complete silencing at the final steps of MSCI. Furthermore, not all X genes in $Topbp1^{B5/B5}$ spermatocytes were silenced; instead, some genes were only partially silenced while others exhibited increased expression after initial silencing. Interestingly, a sex body is formed that is morphologically indistinguishable from the sex body in wild-type animals. Several heterochromatin markers, as well as multiple canonical markers of sex body formation localize properly in the sex body of $Topbp1^{B5/B5}$ mice. Overall, these findings suggest a non-canonical role for the ATR-TOPBP1 signaling axis in ensuring proper XY silencing dynamics during pachynema. This is the first DDR mutant that separates XY silencing from sex body formation, and that separates TOPBP1's role in spermatogenesis from its roles in organismal viability.

The $B5$ allele reported here, which carries eight lysine to glutamic/aspartic acid substitutions in BRCT domain 5, is the first mutation shown to impair the meiotic silencing function of TOPBP1 in spermatocytes without severely disturbing TOPBP1's role in synapsis and sex body formation. Consistent with this being a separation of function mutant, $Topbp1^{B5/B5}$ males are viable and grossly normal, while completely sterile, whereas $Topbp1$ null, or AAD mutated, mice are embryonic lethal[48,49]. Moreover, depletion of TOPBP1 in mammalian cell lines triggers a robust G2/M arrest followed by cell senescence and loss of viability[48] $Topbp1^{B5/B5}$ MEFs do not display issues with cell proliferation or DDR defects. Based on these observations, and the finding of a silencing defect in $Topbp1^{B5/B5}$ spermatocytes, it is likely that the role of TOPBP1 in silencing documented here could be specifically relevant in the context of male meiosis. It is important to note that other mutants previously reported to have XY silencing defects during meiotic prophase

I, such as *Ago4* null and *Dicer* CKO, do not result in complete loss of sperm production, but a sub-

fertility phenotype[75,77]. Therefore, we speculate that the specific type of silencing defect in

Topbp1[B5/B5] spermatocytes is particularly toxic, similar to other mutations in the DDR pathway

that result in MSCI defects, which would explain the highly penetrant defect in sperm production.

The model that distinct TOPBP1 interactions mediate distinct ATR signaling pathways offers a

potential explanation for why *Topbp1[B5/B5]* have specific defects in silencing without noted effects

in other key processes regulated by ATR, such as synapsis. In addition to binding and activating

ATR through its AAD domain[5,21], TOPBP1 can bind to several proteins through its BRCT

domains[47] and act as a scaffolding protein to bring substrates in close proximity to ATR, thus

facilitating the propagation of specific ATR signaling pathways. Experiments using ectopic

expression of Flag-tagged TOPBP1 in HEK293T cells revealed that the set of mutations in *B5*

disrupt the ability of TOPBP1 to interact with 53BP1 and BLM, as expected from previous

reports[22,24,27,28,60]. However, it is unclear if the same interactions are also disrupted in

spermatocytes, or whether additional TOPBP1 interactions mediated by its BRCT5 specifically

during meiotic prophase I are also disrupted. Apart from disrupting protein interactions, it is also

possible that the observed changes in TOPBP1 protein stability in the *Topbp1 B5* mutant can

contribute to impairing its roles in silencing. Such change in protein stability is consistent with a

previous report showing that the TOPBP1-BLM interaction contributes to protein stabilization[59,60].

Further work will be necessary to determine if the phenotypes observed in *Topbp1[B5/B5]*

spermatocytes are caused specifically by disruption of specific protein interactions, or by a

combination of disrupted interactions and reduced protein abundance. It is worth mentioning that

the *Topbp1[B5/B5]* phenotype is distinct from the *Topbp1* CKO despite the reduction in protein

abundance.

Our finding that SETX localization to chromatin loops of the XY is impacted in *Topbp1[B5/B5]*

pachytene spermatocytes, together with our previous report that SETX undergoes ATR-dependent

phospho-regulation in spermatocytes, suggest that an ATR-TOPBP1-SETX signaling axis is

important for the silencing reinforcement in late MSCI. Genetic data support that impairment of

this specific signaling axis would impact silencing without impacting synapsis. For example,

Topbp1 CKO, *Rad1* CKO, and *Atr* CKO spermatocytes display strong synapsis defect and

defective entry in pachynema[18,34], whereas *Setx* null spermatocytes complete autosomal synapsis,

while still displaying MSCI defects. On the other hand, is it likely that ATR signaling is controlling

a specific aspect of SETX function since *Topbp1*[B5/B5] spermatocytes do not share all defects

observed in *Setx* null spermatocytes, as noted by the localization of γH2AX, sumoylation events,

ubH2A and ATR at chromatin loops, which are defective in *Setx* null spermatocytes but normal in

Topbp1[B5/B5] spermatocytes. Moreover, *Topbp1*[B5/B5] pachytene spermatocytes, but not *Setx* null

spermatocytes, are able to reach the stage of crossover designation with MLH1 positive cells.

Taken together, these observations suggest that *Topbp1*[B5/B5] pachytene spermatocytes progress

further in pachynema when compared to *Setx* null spermatocytes and are consistent with a model

in which a ATR-TOPBP1 control only specific(s) mode of SETX regulation[29,64].

The model involving SETX as a potential factor by which ATR controls silencing late in MSCI

opens exciting directions to explore the interface of ATR-TOPBP1 with RNA processing. Given

the established role for SETX in the resolution of R-loops[78], it is tempting to speculate that

silencing defects in *Topbp1*[B5/B5] mutant may be related to aberrant accumulation of RNA-DNA

hybrids that may affect removal of nascent mRNAs that is necessary for imposing silencing. In

support of this model, R-loops affect chromatin architecture at promoters and interfere with the

recruitment of transcription factors and chromatin remodelers, as observed in regions harboring

CpG islands where R-loops prevent the action of DNA methyltransferases, thus preventing

silencing[79]. Notably, highly transcribed genes, which tend to accumulate more R-loops[80],

displayed increased silencing defects in the *Topbp1*[B5/B5] mutant.

While we have provided strong evidence to suggest a defect in later stages of MSCI as the cause

of the cell death observed in *Topbp1*[B5/B5] spermatocytes, we cannot exclude the potential

contribution of other defects, beyond silencing, to the loss of diplotene cells. The increased number

of MLH1 foci suggested an altered recombination pattern, possibly impairing the ratios of class I

and class II crossovers. The BRCT 5 domain of TOPBP1 interacts with the BLM helicase[27,59],

which has been found to play a role in meiotic recombination in yeast and mice[81,82]. *Blm* CKO

mice display severe defects in prophase I progression in spermatocytes, including, incorrect

pairing and synapsis of homologs, and defective processing of recombination intermediates,

leading to increased chiasmata[82]. These observations raise the possibility that impaired meiotic

progression and cell death in *Topbp1^B5/B5* spermatocytes is a combination of defects in MSCI and

recombination. *Topbp1^B5/B5* spermatocytes do not progress beyond pachytene hence we were not

able to visualize chiasmata and directly infer whether or not *Topbp1^B5/B5* is defective in crossing

over.

Our findings showing that TOPBP1 plays a specific role in silencing reinforcement after the first

wave of MSCI are consistent with the recently proposed notion that the establishment and

maintenance of MSCI is a dynamic process[17,64]. Also consistent with this notion is our recent

finding showing that ATR signaling is itself also highly dynamic and constantly being cycled[64].

For example, using mice treated with the ATR inhibitor (ATRi) AZ20 for four hours, we found

that such a short treatment is already sufficient to cause a complete loss of γH2AX, pMDC1 and

SETX localization from the XY chromosomes[17,64]. Furthermore, germ cells subjected to ATRi for

24 hours showed complete recovery of γH2AX only three hours after release from ATRi

treatment[17]. We propose that TOPBP1 acts on this phospho-cycle to ensure proper silencing

reinforcement and maintenance, potentially by counteracting the engagement of anti-silencing

factors that dynamically enter the sex body and need to be actively antagonized at the XY. Future

work involving the characterization of possible unknown interactors of the BRCT domain 5 of

TOPBP1, as well as functional dissection of ATR targets in MSCI, is essential to understand how

TOPBP1 modulates the silencing machinery and shapes silencing dynamics.

In summary, our study presents a unique model for investigating the role of DDR factors in XY

silencing. By allowing the uncoupling of MSCI progression from sex body formation, the

Topbp1^B5/B5 mutant enables the study of MSCI dynamics during key stages late in pachynema.

Notably, the inability of *Topbp1^B5/B5* to sustain or reinforce silencing opens the possibility of

uncovering new insights into the MSCI-dependent pachytene checkpoint.

STAR*Methods

Key resource table

Reagent type	Designation	Source or Reference	Identifiers	Additional information
Primer	p3xflag-TOPBP1-Flag-N-terminal-F	IDT	PCR primer	attcatcgatagatctgata ATGTCCAGAAAT GACAAAGA
Primer	p3xflag-TOPBP1-Flag-N-terminal-R	IDT	PCR primer	tagagtcgactggtaccgat ttagTGTACTCTAG GTCGTT
Primer	hTopbp1_K1317A _R	IDT	Site-directed mutagenesis	tgccactgaggctaaatac gcctcgtttcgaagtggatg t
Primer	hTopbp1_K1317A _F	IDT	Site-directed mutagenesis	acatccacttcgaaacgag gcgtatttagcctcagtggc a
Primer	hTopbp1_K155A_ F	IDT	Site-directed mutagenesis	caggtttgcagcaactaaat atgctttgctaccaacttctc ctgca
Primer	hTopbp1_K155A_ R	IDT	Site-directed mutagenesis	tgcaggagaagttggtagc aaagcatatttagttgctgca aacctg
Primer	hTopbp1_K250A_ R	IDT	Site-directed mutagenesis	tctcttggcacactcatacg cctgaccttttggttcttgc
Primer	hTopbp1_K250A_ F	IDT	Site-directed mutagenesis	gcaagaaccaaaaggtca ggcgtatgagtgtgccaag aga
Primer	musTopbp1_BRCT 5-Fwd genotyping	IDT	PCR primer	tgcatttccattaaccaacct c
Primer	musTopbp1_BRCT 5 Rev genotyping	IDT	PCR primer	ggtagagttcaaatgtgtgt catg
Antibody	phospho H2A.X (Ser139)	Millipore	Cat# 05-636;RRID:AB_ 309864	IF (meiotic spreads) 1:50000 IF 1:2000

Antibody	SCP3 antibody	Abcam	Cat# ab97672;RRID: AB_10678841	IF (meiotic spreads) 1:1000
Antibody	SYCP3	Lenzi et. al, 2005[83]	Custom	IF (meiotic spreads) 1:10000
Antibody	SCP1	Abcam	ab15090	IF (meiotic spreads) 1:1000
Antibody	Rad51	Millipore	PC130	IF (meiotic spreads) 1:1000
Antibody	ATR	Cell signaling	2790	IF (meiotic spreads) 1:1000
Antibody	TOPBP1	Danielsen et.al, 2008[84]	Custom	IF (meiotic spreads) 1: 500 Western blot 1:1000
Antibody	phospho-Chk1 (ser317)	Cell Signaling	12302	IF (meiotic spreads) 1: 100 Western blot 1:1000
Antibody	H1T	A gift from Dr. Mary Ann Handel[50]	Custom	IF (meiotic spreads) 1: 500
Antibody	HORMAD2	Wojtasz et. al, 2012[85]	Custom	IF (meiotic spreads) 1: 500
Antibody	HORMAD1	Wojtasz et. al, 2012[85]	Custom	IF (meiotic spreads) 1: 500
Antibody	phospho HORMAD2 (S271)	Wojtasz et. al, 2009	Custom	IF (meiotic spreads) 1: 500
Antibody	GAPDH	Thermo Fisher Scientific	AM4300	Western blot 1:5000
Antibody	β-Actin	Cell Signaling	4967	Western blot 1:5000
Antibody	phospho MDC1 (T4)	Abcam	Ab35967	IF (meiotic spreads) 1: 500
Antibody	SETX	Abcam	Ab220827	IF (meiotic spreads) 1: 100
Antibody	SETX	Novus	NBP1-94712SS	Western blot 1:1000

		Biologicals		
Antibody	MLH1	BD Biosciences	550838	IF (meiotic spreads) 1: 200
Antibody	MLH3	A gift from Dr. Paula Cohen[86]	Custom	IF (meiotic spreads) 1: 200
Antibody	CHK1	Santa Cruz	sc-8408	Western blot 1:1000
Antibody	CHK2	Millipore	05-649 (clone7)	Western blot 1:1000
Antibody	phospho CHK1 (S345)	Cell Signaling	2341	IF (meiotic spreads) 1: 200 Western blot 1:1000
Antibody	phospho RPA (S4/S8)	Bethyl	A300-245A,	Western blot 1:1000
Antibody	RPA (made against full length RPA2 expressed and purified in *E. Coli* and injected to rabbit)	N/A	Custom	Western blot 1:1000
Antibody	KAP1	Bethyl	a700-014-T	Western blot 1:1000
Antibody	phospho KAP1 (S824)	Bethyl	A300-767A-T	Western blot 1:1000
Antibody	53BP1	Cell signaling	4937	Western blot 1:1000
Antibody	53BP1	Novus Biologicals	NB100-304	IF (meiotic spreads) 1: 200
Antibody	BLM	Abcam	ab2179	Western blot 1:500
Antibody	Rad9	Bethyl	A300-890A-T	Western blot 1:1000
Antibody	BRCA1	Kakarougkas et. al, 2013[87]	Custom	IF (meiotic spreads) 1: 200
Antibody	BRCA1	Proteintech	22362-1-AP	Western blot 1:1000
Antibody	MDC1	Modzelewski et. al, 2015[88]	Custom	IF (meiotic spreads) 1: 200 IF

				1: 200
Antibody	RNA pol 2	Millipore	05-623	IF (meiotic spreads) 1: 2000
Antibody	phospho RNA pol 2 (Ser2)	Millipore	04-1571	IF (meiotic spreads) 1: 400
Antibody	phospho RNA pol 2 (Ser5)	Millipore	04-1572	IF (meiotic spreads) 1: 400
Antibody	H3K9ac	Abclonal	A7255	IF (meiotic spreads) 1: 200
Antibody	H3K9me3	Active Motif	39766	IF (meiotic spreads) 1: 200
Antibody	H3K4me1	Abclonal	A2355	IF (meiotic spreads) 1: 200
Antibody	CHD4	Abclonal	A10557	IF (meiotic spreads) 1: 200
Antibody	Sumo_2/3	Proteintech	G67154-1-1g	IF (meiotic spreads) 1: 200
Antibody	USP7	Proteintech	66514-1-Ig	IF (meiotic spreads) 1: 200
Antibody	H3K27ac	Active Motif	39134	IF (meiotic spreads) 1: 200
Antibody	H3K36ac	Active Motif	61102	IF (meiotic spreads) 1: 200
Antibody	H3K4me3	Active Motif	39160	IF (meiotic spreads) 1: 200
Antibody	H4K16ac	Abclonal	A5280	IF (meiotic spreads) 1: 200
Antibody	H4c	Millipore	06-598	IF (meiotic spreads) 1: 200
Antibody	H2AK199ub	Abcam	Ab193203	IF (meiotic spreads) 1: 200
Antibody	Goat anti-rabbit IgG (H + L) highly	Thermo Fisher Scientific	A-11034	IF (meiotic spreads) 1: 1000

	cross-adsorbed secondary antibody, Alexa Fluor 488			
Antibody	Goat anti-mouse IgG (H + L) antibody, Alexa Fluor 488 conjugated	Thermo Fisher Scientific	A-11017	IF (meiotic spreads) 1: 1000
Antibody	Goat anti-rabbit IgG (H + L) antibody, Alexa Fluor 594 conjugated	Thermo Fisher Scientific	A-11012	IF (meiotic spreads) 1: 1000
Antibody	Goat anti-mouse IgG (H + L) highly cross-adsorbed secondary antibody, Alexa Fluor Plus 594	Thermo Fisher Scientific	A32742	IF (meiotic spreads) 1: 1000
Antibody	Goat anti-guinea pig IgG (H + L) highly cross-adsorbed secondary antibody, Alexa Fluor 647	Thermo Fisher Scientific	A-21450	IF (meiotic spreads) 1: 1000
Commercial assay or kit	ApopTag Plus Peroxidase In Situ Apoptosis Kit	Millipore	S7101	N/A
Software, algorithm	GraphPad Prism 9	GraphPad Prism	RRID:SCR_002798	N/A
Software	Cellranger v7.0.0	10X Genomics	https://support.10xgenomics.com/single-cell-gene-expression/software/downloads/3.1	N/A

Software	R v4.2.1	R Foundation	RRID:SCR_00 1905	N/A
Software	Seurat v4.3.0	Hao et. al, 2013[89]	RRID:SCR_00 7322	N/A
Software	sva v3.44.0	Leek et. al, 2023[90]	RRID:SCR_01 2836	N/A
Software	SingleR v1.10.0	Aran et. al, 2019[91]	RRID:SCR_02 3120	N/A
Software	Reshape2 v1.4.4	Wickham, 2007[92]	RRID:SCR_02 2679	N/A
Software	biomaRt v2.52.0	Durinck et. al, 2009[93]	RRID:SCR_01 9214	N/A
Software	ggpubr v0.5.0	Kassambara, 2009[94]	RRID:SCR_02 1139	N/A
Software	Tidyverse v1.3.2	Wickham et. al, 2019[95]	RRID:SCR_01 9186	N/A
Software	ggnewscale v0.4.8	Campitelli	https://eliocam p.github.io/ggn ewscale/index.h tml	N/A
Software	Harmony v0.1.1	Korsunsky et.	RRID:SCR_02	N/A

		al, 2019[96]	2206	
Software	Org.Mm.eg.db v3.15.0	Carlson, 2019[97]	RRID:SCR_02 3488	N/A
Software	Corrplot v0.92	Wei et. al, 2017[98]	RRID:SCR_02 3081	N/A
Software	SeuratWrappers v0.3.1	Satija Lab	RRID:SCR_02 2555	N/A
Software	gghalves v0.1.4	Tiedemann	https://github.c om/erocoar/ggh alves	N/A

Resource availability

Lead contact

Further information and requests for resources and data should be directed to and will be fulfilled
by the lead contact, Marcus Smolka (email: mbs266@cornell.edu).

Materials availability

This study generated a unique antibody, RPA.

Mice, genotyping and treatment of mice with IR

All mice used in this work were handled following federal and institutional guidelines under a
protocol approved by the Institutional Animal Care and Use Committee (IACUC) at Cornell
University. CRISPR/Cas9 editing was used to engender the *Topbp1B5* allele and performed by the
Cornell Mouse Modification core facility. To this end, the online tools CRISPR gold and
ChopChop were used to generate high quality (guide score >9) CRISPR guide RNAs targeting the
intronic sequences neighboring the genomic sequence of *Mus musculus topbp1* exon 13. The

CRISPR crRNAs (purchased from IDT) harboring the sequences ccaactcaggtcggccgctcttg and

cctcgattagtcctcaaggcgag (PAM sites are underlined), both targeting the reverse strand, the repair

template (below), and CAS9 RNA were injected on embryos from super ovulated, plugged

C57BL/6J female mice crossed to C57BL/6J stud males. 2-cell stage embryos were then implanted

on pseudo pregnant females and pups were genotyped after one week old.

Repair template:

acagcagggcttctctgtgtaaccctctctccgtagaccagtttggccttgatcgaactcaggtcggccgctcttgcctctagagtcctgggat

taaaggcgtgcactgccaccacccagagtatgtttctctgacattaaccatgctattattttttttaaaatgagctaattgtgtgttcatttgctttattt

ccatgtaaaattttagTGTTCAAGAATTCTTTGTTGACGAAGCCAATGCAGAGGAAGGCATGCT

CGCCAGCACACATCTTATAGTGGAGGAACCGACTGGTTCCGAATACGAAGCTGCAG

AGGAATGGAGTTTGCCGGCAGTTAACATTTCATGGCTCTTAGAAACTGCGGAAATCG

GGGAGGAAGCAGATGAAAACCATTTTCTGGTTGACAACGCACCTAAACAAGgttagaagt

ccttgtttttttttttatgtattttacaacttgatggtttctgaaatagggatgttccagtacttgctttaaaacatttgtatgaccctaacctcagtcagtg

gtgcttacttcagaacccctgagtgaaacacggaaagcagatcaatgaagaagcgcatcagggtcaacggtcgattagtcctcaaggcga

gtgacgagaaggtgacccccga atggctgttagaagcagttttttata (purchased from IDT as a G-block)

Intronic sequence is shown in lower case, exon 13 sequence is shown in capital letters; underline

represents the mutated residues, pink represents the mutated PAM residues, in orange is shown the

guide RNA sequences and in red, the targeted amino acid sequences. Of note, although this repair

template encodes for 11 amino acid changes, only 8 were successfully inserted into the mouse

topbp1 exon 13 locus. For mice genotyping, the following primers were used 5'-

tgcatttccattaaccaacctc-3' and 5'- ggtagagttcaaatgtgtgtcatg-3' (also shown at key resources table).

For irradiation, *Topbp1^{+/+}* and *Topbp1^{B5/B5}* mice were placed in a 137 Cesium-sealed source

irradiator (J.L. Shepherd and Associates) with a rotating turntable and irradiated with 7 Gy IR.

MEFs and cell survival assays

MEFs were prepared from E13.5 mouse embryos as previously described[99]. Briefly, embryos were

dissected and mechanically disrupted using pipette aspiration until homogenous. Cells were

allowed to settle, and the supernatant was transferred into Dulbecco's modified medium

supplemented with 10% fetal bovine serum, 1% penicillin/streptomycin and 1% nonessential

amino acids. Following four days of growth, cells were then immortalized by transduction with a

large-T antigen lentivirus. Subsequently, cells were selected with 10 µg/mL puromycin.

For colony survival assays, 500 cells were seeded per 10 cm dish, allowed to adhere for 24h and

treated with Phleomycin or Hydroxyurea for 24h (drug concentrations are displayed on figure 2A-

D and Supplemental Figure 2). In the following day, cells were released and let to form colonies

for 10-15 days. Cells were then washed once with PBS, fixed in 100% methanol for 1h, stained

with 0.1% crystal violet (MP Biomedicals, 152511) solution overnight, and then washed with

distilled water before imaging and counting.

For accessing DDR and checkpoint responses via western blot, 2×10^6 cells were seeded on a 60

cm dish, allowed to adhere for 24h, and treated with Phleomycin or Hydroxyurea for 3h (drug

concentrations are displayed on Figure 2).

Cell culture

HEK-293T cells were cultured in Dulbecco's modified medium supplemented with 10% fetal calf

serum, 1% penicillin/streptomycin and 1% nonessential amino acids. Immortalized MEFs were

cultured in Dulbecco's modified medium supplemented with 10% fetal calf serum, 1%

penicillin/streptomycin, 1% nonessential amino acids, and 1% glutamine supplementation. All

cells were kept at 37°C and 5% CO_2. All the cell lines were regularly tested for mycoplasma

contamination with the Universal Mycoplasma Detection Kit (ATCC). HEK-293T cells were

transfected using homemade polyethylenimine (Polysciences, Inc.). 36h after transfection, cells

were treated with 1mM HU (Hydroxyurea) and then harvested for immunoprecipitation

experiments.

Plasmids

The full-length TOPBP1 CDS was cloned on a p3xflag vector (Milipore/Sigma E7658) using

Gibson assembly (NEB) following the manufacturer's instructions. The p3xflag-TOPBP1 was

used as a template to generate p3xflag-TOPBP-K155A, p3xflag-TOPBP-K250A, p3xflag-

TOPBP-K704A, p3xflag-TOPBP-K1317A through site-directed mutagenesis using prime STAR

master mix (Takara) and Gibson assembly to generate p3xflag-TOPBP-KE, using the following

G-block (IDT) containing the 8 charge-reversal point mutations at the BRCT 5 domain of

TOPBP1:

AACGAATCCAATGCAGAAGAAGGCATGTTTGCCAGTACTCATCTTATACTGGAAGA

ACGTGGTGGCTCTGAATATGAAGCTGCAAAGAAGTGGAATTTACCTGCCGTTACTAT

AGCTTGGCTGTTGGAGACTGCTAGAACGGGAGAAGA.

All primers used for the cloning are shown in the key resources table.

Immunoblotting

Cells were harvested and lysed in modified RIPA buffer (50 mM Tris–HCl, pH7.5, 150 mM NaCl,

1% Tergitol, 0.25% sodium deoxycholate, 5 mM ethylenediaminetetraacetic acid (EDTA))

supplemented with complete EDTA-free protease inhibitor cocktail (Roche), 1 mM

phenylmethylsulfonyl fluoride (PMSF) and 5 mM NaF. Whole cell lysates, after sonication, were

cleared by 15 min centrifugation at 17,000 xg at 4∘C. Twenty micrograms of protein extract were

mixed with 3X sodium dodecyl sulfate sample buffer and resolved by SDS-PAGE. Gels were

transferred on polyvinylidene difluoride membranes and immunoblotted using standard

procedures. Western blot signal was acquired with a Chemidoc Imaging System (BioRad).

Antibody information is provided in the key resources table.

Immunofluorescence

MEFs were grown on coverslips and then submitted to IR, 5 gy and allowed to recover for 1.5h at

37°C and 5% CO_2. Cells were then fixed using 3.7% formaldehyde in phosphate-buffered saline

(PBS) for 10 min at room temperature (RT). Fixed cells were then washed 3x with PBS,

permeabilized for 5 min with 0.2% Triton X-100/PBS at RT and blocked in 10% bovine serum

albumin/PBS for 20 min at RT. Coverslips were incubated first with primary antibodies for 2h at

RT, followed by three washes with PBS, and then for 1 hour with relative secondary antibodies.

After incubation with secondary antibodies, coverslips were washed three times with PBS and then

mounted on glass microscope slides using DAPI–Vectashield mounting medium (Vector

Laboratories). Slides were imaged on a Leica DMi8 Microscope with a Leica DFC9000 GTC

camera using the LAS X (Leica Application Suite X) software with a 100x objective. For

micronuclei scoring, ∼150–200 cells/replicate were counted. The two-tailed Student's t-test was

used for statistical analysis. Antibody information is provided in the key resources table.

Meiotic Spreads

Meiotic surface spreads were performed from 8- to 12-week-old mice as described by Kolas et
al.,[100] .Briefly, decapsulated testis from mice were incubated on ice in a hypotonic extraction buffer
for 45 min. Tubules were then minced into single- cell suspension in 100 mM sucrose, and cells
were spread on slides coated with 1% PFA with 0.15% TritionX- 100 and incubated in a
humidifying chamber for 4h. For immunostaining, slides were blocked using 10% goat serum and
3% BSA, followed by incubation overnight with primary antibody (listed in the STAR methods)
at 4°C in a humidifying chamber. Secondary antibodies were incubated at 37°C for 2h in the dark,
and slides were then cover-slipped using antifade mounting medium (2.3% DABCO, 20 mM Tris
pH 8.0, 8 µg DAPI in 90% glycerol). Slides were imaged on a Leica DMi8 Microscope with a
Leica DFC9000 GTC camera using the LAS X (Leica Application Suite X) software. For every
condition, a minimum of 50 images from at least two independent mice were acquired. To quantify
fluorescence intensity, the LAS X software quantification tool was used as previously described[64].
Antibody information is provided in the key resources table.
589

3D-Structured Illumination super resolution Microscope (3D-SIM)
Higher resolution images were acquired using an ELYRA 3D-Structured Illumination Super
Resolution Microscopy (3D-SIM) from Carl Zeiss with ZEN Black software (Carl Zeiss AG,
Oberkochen, Germany). Images are shown as maximum intensity projections of z stack images.
To reconstruct high-resolution images, raw images were computationally processed by ZEN
Black. The brightness and contrast of images were adjusted using ImageJ (National Institutes
of Health, USA). Antibody information is provided in the key resources table.
597

Fertility assays
For fertility testing, 8 weeks old *Topbp1$^{B5/B5}$* females and C57BL/6 males or C57BL/6 females and
Topbp1$^{B5/B5}$ males were singly housed, where pregnancies were monitored for a period of one
month. Viable pups were counted on the first day of life. For *Topbp1$^{B5/B5}$* males, breeding cages
remained active for a period of six months and at no time pregnant females nor birth of pups were
detected. No noticeable defects were found on fertility of Topbp*1$^{B5/+}$*, males or females (data not
shown).
605

606 **TUNEL**

607   TUNEL assay was conducted using the Apoptag kit (EMD Millipore) following the

608   manufacturer's instructions. The data were quantified in Image Scope by counting the number of

609   positive cells per tubule for 100 tubules of each genotype, 3 mice each. Statistical differences

610   between *Topbp1*[+/+] and *Topbp1*[B5/B5] were analyzed using Welch's unpaired t- test in GraphPad.

Hematoxylin and eosin staining

Adult testes - from 12 weeks old mice - were dissected and incubated in Bouin's fixative overnight,

washed during 30 min each in 30%, then 50% and then 70% ethanol. The 70% ethanol wash was

repeated 3 times more. Testes were then embedded in paraffin. 5 µm sections were mounted on

slides. After rehydration in Safe Clear Xylene Substitute followed by decreasing amounts of

ethanol, slides were stained with hematoxylin followed by eosin. The slides were then gradually

dehydrated by incubation in increasing concentrations of ethanol before mounting using toluene

mounting medium.

Epididymal sperm counts

Caudal and epididymides from 8- to 12-week-old mice were minced with fine forceps at 37°C in

a Petri dish containing Dulbecco's modified medium supplemented with 10% fetal calf serum, 1%

penicillin/streptomycin, 4% BSA, and 1% nonessential amino acids. Samples were then incubated

at 37°C for 30 min allowing sperm to swim out into the media, then fixed in 10% neutral-buffered

formalin (1:25 dilution). Sperm cells were counted using a hemocytometer and analyzed

statistically using the two-tailed Student's t-test between *Topbp1*[+/+] and *Topbp1*[B5/B5].

Enrichment of testes phosphopeptides and TMT labeling

The enrichment of testes phosphopeptides and TMT labeling were done as described previously[64].

Briefly, whole decapsulated testes were collected from 8- to 12-week-old mice after which tissue

was subject to lysis, protein quantification and normalization, denaturation, alkylation,

precipitation, digestion, and solid-phase extraction (SPE) C18 cartridge clean up as described by

Sims et. al[64]. Lyophilized tryptic peptides were then subject to phosphopeptide enrichment using

a High-Select Fe-NTA Phosphopeptide Enrichment Kit according to the manufacturer's protocol

(Cat# A32992, Thermo Scientific). Phosphopeptide samples were dried in silanized glass shell

vials, resuspended in 50 mM HEPES, and labeled with 100 µg of TMT sixplex Isobaric Label

Reagents (Thermo Scientific) using 3 TMT channels for each condition ($Topbp1^{+/+}$ and

$Topbp1^{B5/B5}$). The TMT-labeling reaction was done at room temperature for 1h and quenched with

5% hydroxylamine for 15 minutes. After quenching, TMT-labeled peptides from all six channels

were pooled, acidified with 0.1% TFA, and desalted using a SPE 1cc C18 cartridge (Sep-Pak C18

cc vac cartridge, 50 mg Sorbent, WAT054955, Waters). Bound TMT-labeled phosphopeptides

were eluted with 80% acetonitrile, 0.1% acetic acid in water before being dried via vacuum

concentrator.

Mass spectrometric analysis of TMT-labeled phosphopeptides

The dried TMT-labeled phosphopeptides were prefractionated using offline HILIC HPLC prior to

being analyzed by mass-spectrometry as described by Sims et. al[64]. The LC-MS/MS was

performed on an UltiMate 3000 RSLC nano chromatographic system coupled to a Q-Exactive HF

mass spectrometer (Thermo Fisher Scientific). The chromatographic separation was achieved via

a 35cm-long 100 µm inner diameter column packed in-house with 3 µm C18 reversed-phase resin

(Reprosil Pur C18AQ 3 µm). The Q-Exactive HF was operated in data-dependent mode with

survey scans acquired in the Orbitrap mass analyzer over the range of 380–1800 m/z with a mass

resolution of 120,000. MS/MS spectra were performed after selecting the top 7 most abundant +2,

+3, or +4 ions and a precursor isolation window of 0.7 m/z. Selected ions were fragmented by

Higher-energy Collisional Dissociation (HCD) with normalized collision energies of 28, with

fragment mass spectra acquired in the Orbitrap mass analyzer with a monitored first mass of 100

m/z, mass resolution of 15,000, AGC target set to 1×10^5, and maximum injection time set to 100

ms. A dynamic exclusion window of 30 seconds was specified.

Phosphoproteomic data analysis

The Trans Proteomic Pipeline (TPP) version 6.0.0 was used for phosphopeptide identification and

quantification. MS data were converted to mzXML using msConvert as packaged with the TPP,

after which spectral data files were searched using the Comet search engine (v2021 rev 1)[101].

Peptide identifications were validated using PeptideProphet, phosphorylation site localization was

performed using PTM Prophet, and TMT channel quantification was performed using Libra.

Results from Libra were exported as tab-delimited files for further processing via R scripts as

previously described[64]. The mass spectrometry phosphoproteomics data have been deposited to

the ProteomeXchange Consortium via the PRIDE [1] partner repository with the dataset identifier PXD042199.

Immunoprecipitation

The immunoprecipitation (IP) experiments were performed as described by Liu et. al[22]. Briefly, cell pellets were lysed in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% tergitol, 0.25% sodium deoxycholate, and 5 mM EDTA, supplemented with EDTA-free protease inhibitor cocktail, 5 mM sodium fluoride, 10 mM β-glycerolphosphate, 1 mM PMSF, and 0.4 mM sodium orthovanadate. The protein extracts were cleared by 10-min centrifugation and then were incubated with anti-FLAG agarose beads (Sigma-Aldrich) for 16h at 4°C. The beads were then washed four times with the same buffer used for IP and then eluted using three resin volumes of the elution buffer (100 mM Tris-HCl, pH 8.0, and 1% SDS, and 1 mM DTT).

Mass spectrometric analysis of immunoprecipitates

HEK-293T cells were grown in stable isotope labeling with amino acids in cell culture (SILAC) as previously described[22] and transfected as described above. Cells were treated with 1 mM HU for 16h before harvesting. Flag-TOPBP1 was immunoprecipitated using anti-FLAG agarose beads. Immunoprecipitates were then prepared for mass spectrometry analysis by reduction, alkylation, precipitation, and digestion by trypsin. The peptides were desalted, dried, and then fractionated by hydrophilic interaction chromatography as previously described[22]. Fractions were dried and analyzed by liquid chromatography–tandem mass spectrometry using a mass spectrometer (Q-Exactive HF Orbitrap; Thermo Fisher Scientific). The capillary column was 35 cm long with a 100-µm inner diameter, packed in-house with 3 µm C18 reversed-phase resin (Reprosil Pur C18AQ 3 µm). Peptides were separated over an 70-minute linear gradient of six to 40% acetonitrile in 0.1% formic acid at a flow rate of 300 nl/min as described previously[102]. Xcalibur 2.2 software (Thermo Fischer Scientific) was used for the data acquisition, and The Q-Exactive HF was operated in data-dependent mode with survey scans acquired in the Orbitrap mass analyzer over the range of 380–1800 m/z with a mass resolution of 120,000. The maximum ion injection time for the survey scan was 100 ms with a 3e6 automatic gain-control target ion. Tandem mass spectrometry spectra were performed by selecting up to the 20 most abundant ions with a charge state of 2, 3, or 4 and with an isolation window of 1.2 m/z. Selected ions were

fragmented by higher energy collisional dissociation with a normalized collision energy of 28, and

the tandem mass spectra was acquired in the Orbitrap mass analyzer with a mass resolution of

17,500 (at m/z 200). The Trans Proteomic Pipeline (TPP) version 6.0.0 was used for peptide

identification and SILAC quantification. MS data were converted to mzXML using msConvert as

packaged with the TPP, after which spectral data files were searched using the Comet search

engine (v2021 rev 1)[101]. The following parameters were used in the database search: semitryptic

requirement, a mass accuracy of 15 ppm for the precursor ions, a differential modification of

8.0142 D for lysine and 10.00827 D for arginine, and a static mass modification of 57.021465 D

for alkylated cysteine residues. Peptide identifications were validated using PeptideProphet and

SILAC quantification was performed using XPRESS as described previously[102]. The mass

spectrometry data have been deposited to the ProteomeXchange Consortium via the PRIDE [1]

partner repository with the dataset identifier PXD042199.

Total germ cells preparation

Mice testis were collected from 8- to 12-week-old mice (n = 5 mice, 10 testis for $Topbp1^{+/+}$, and

n = 20-30 mice, 40-60 testis for $Topbp1^{B5/B5}$), and dissociated using standard protocols for germ

cell extraction[73]. Briefly, decapsulated testes were held in 10 ml of preheated (35°C) DMEM-F12

buffer containing 2 mg of Collagenase 1A and DNAse 7 mg/ml, on a 50 ml conical tube. The

collagenase digestion was performed at a 35°C shaker water-bath, 150 rpm during 5 min. The

collagenase digestion was stopped by the addition of 40 ml of DMEM-F12 and the tubules were

let to decantate for 1 min. The supernatant was removed and added another 40 ml of DMEM-F12

to further wash the residual collagenase and remove somatic and excessive sperm cells. Next, the

tubules were digested using 10 ml of DMEM-F12 buffer containing 5 mg of trypsin (Thermo

Fisher 27250018) on a 50 ml conical tube and the reaction was carried out on a 35°C shaker water-

bath, 150 rpm during 5 min. Digested tubules were strained on a 100 μm strainer containing 3 ml

of FBS (100% Fetal Bovine Serum, heat inactivated, Sigma F4135-500mL). Total germ cells were

centrifuged at 300 xg for 5 min, 4°C and checked for single cells and viability.

Flow cytometry analysis

Total germ cell extracts were stained with Vybrant dye cycle (VDG) (Invitrogen) 100 μM for 30

min at room temperature, kept on dark, and rocking. After staining, cells were sorted as previously

described[103] aiming to enrich for spermatocytes, and using the Sony MA900 fluorescent activated

cell sorter (FACS), tuned to emit at 488 nm and with a 100 μm nozzle. Laser power was set to

collect VDG-emitted fluorescence in FL1. Sorted cells were collected on 1.5 ml tubes containing

0.5 ml of DMEM-F12 buffer + 10% FBS. The FACS was done at the Flow Cytometry Facility -

Cornell University.

Single-Cell RNA sequencing

Flow sorted cells were submitted to the Cornell DNA Sequencing Core Facility and processed on

the 10X Genomics Chromium System targeting 5000–7000 cells per sample as described

previously[73] using the 10X Genomics Chromium Single Cell 3′ RNA-seq v2 kit to generate the

sequencing libraries, which were then tested for quality control on an ABI DNA Fragment

Analyzer and ran on a NextSeq platform with 150 base-pair reads. The sequencing was carried out

to an average depth of 98M reads (range 77M-124M); on average, 91% of reads (range 89%-92%)

and then mapped to the reference genome.

Single-cell RNA sequencing data analysis

Count matrices were generated for each sample using the Cell Ranger counts function and then

imported into Seurat and integrated. Cells were filtered based on gene number, UMI, counts, and

mitochondrial percentage. Cells with less than 500 genes, less than 1000 UMIs, or more than 5%

of mitochondrial reads were excluded from further analysis (Supplemental Figure 29). Raw counts

were normalized using Seurat NormalizeData using default parameters and the top 4000 variable

features were identified using the FindVariableFeatures function using ExpMean for the mean

function and LogVMR for the dispersion function. Principal components were calculated from

variable genes and used with Harmony to correct for batch effect. Harmony embeddings

dimensions 1-20 were used for a Shared Nearest Neighbor graph with k=30, unsupervised

clustering with a resolution of 4, and Uniform Manifold Approximation and Projection (UMAP)

analysis. Cell types were manually identified by use of marker genes, and the SingleR package

was used to confirm cell identity. Chromosome X/autosome 9 ratios were calculated by taking the

mean gene expression of all genes on chromosome 9 or X for a cell and dividing it by the mean

gene expression for all autosomal genes in a cell. Ratios were visualized using ggplot and the

introdataviz package geom_split_violin function. The single cell heatmap was generated using the

DoHeatmap function. The clustered analysis of the X-linked genes shown in Figure 7A, was done

by splitting the detected 233 X-linked genes into 3 distinct categories based on the difference of

RNA level between the spermatocyte stages and normalized by pre-leptotene: reduced silencing

(> 1), unaltered silencing ($\geq$ -1 or $\leq$ 1), or increased silencing ($<$ -1). To further split the X-linked

genes into 3 categories based on the severity of the silencing defect (detected at spermatocyte 3

stage) in $Topbp1^{B5/B5}$: no defect (13 genes), mild (45 genes) or strong defect (170 genes), the

difference in expression between SP3 genes from $Topbp1^{+/+}$ or from $Topbp1^{B5/B5}$ was calculated.

The categories were defined as: Strong > 2.5, Mild $\geq$ 1 or $\leq$ 2.5 and no defect <1. The Single-cell

RNAseq data generated in this study have been deposited in the NCBI GEO database under

accession number GSE232588.

Data and code availability

All data and original codes reported in this work, and any additional information required will be

shared by the lead contact upon request.

Acknowledgments

We thank the members of Cohen lab, Weiss lab, Grimson lab and Smolka lab for comments and

suggestions. We thank Dr. Fenghua Hu and Dr. Tony Bretscher for the use of the microscopes and

Beatriz Almeida for technical assistance. We thank Dr. Lydia Tesfa and the BRC Flow Cytometry

Facility (RRID:SCR_021740) for training and assistance with FACS at the Cornell Institute of

Biotechnology, Peter Schweitzer and the BRC Genomics Facility (RRID:SCR_021727) at the

Cornell Institute of Biotechnology for sequencing experiments, Johana M. de la Cruz and the BRC

Imaging Facility (RRID:SCR_021741) at the Cornell Institute of Biotechnology for training and

assistance with the Elyra Super Resolution Microscope and the 3D-SIM imaging. We thank Marco

Hiler and the BMS and Stem Cell Pathology Unit at the Cornell Veterinary Research Tower for

training and assistance with imaging histological sections using ScanScope. We thank Dr. John

Schimenti, Dr. Robert Munroe, Mr. Christian Abratte, and Cornell's Stem Cell and Transgenic

Core Facility for generating the $Topbp1^{B5/B5}$ mouse. We thank Dr. Paula Cohen for the MLH3

antibody, and Dr. Mary Ann Handel for the H1T antibody.

Author contributions

M.B.S, C.F.R.A and J.R.S. conceived the project. M.B.S and C.F.R.A wrote the paper. C.F.R.A
and J.R.S contributed equally with conceptualization, data curation and formal analysis. C.F.R.A,
J.R.S and E.A.F. performed experiments. A.D and A.G contributed with data curation and formal
analysis. W.C and J.B contributed with data analysis. M.B.S, C.F.R.A, R.W. and P.E.C,
contributed with funding acquisition. R.F. contributed with resources.
798

Declaration of interests

800 No competing interests declared.

Inclusion and diversity

803 We support inclusive, diverse, and equitable conduct of research.

Funding

This work was supported by Eunice Kennedy Shriver National Institute of Child Health and
Human Development (R01-HD095296) to M.B.S and R.W, National Institute of Health – General
Medicine (R35-GM141159) to M.B.S, National Institute of Health (R01HD041012) to P.E.C, the
Eunice Kennedy Shriver National Institute of Child Health and Human Development
(P50HD104454) to A.G., the Spanish Agencia Estatal de Investigación [PID2019-109222RB-
I00/AEI/10.13039/501100011033]; European Union Regional Funds (FEDER) to R. F., and 2022
CVG Scholar Award from the Center of Vertebrates Genomics and the 2022 ASCB International
training Scholarship to C.F.R.A.

814
815

References

1. Subramanian, V.V., and Hochwagen, A. (2014). The meiotic checkpoint network: step-by-step
through meiotic prophase. Cold Spring Harb. Perspect. Biol. *6*, a016675.

2. Handel, M.A., and Schimenti, J.C. (2010). Genetics of mammalian meiosis: Regulation, dynamics
and impact on fertility. Nature Reviews Genetics *11*, 124–136. 10.1038/nrg2723.

3. Keeney, S., Giroux, C.N., and Kleckner, N. (1997). Meiosis-specific DNA double-strand breaks are
catalyzed by Spo11, a member of a widely conserved protein family. Cell *88*. 10.1016/S0092-
8674(00)81876-0.

4. Joshi, N., Brown, M.S., Bishop, D.K., and Börner, G.V. (2015). Gradual Implementation of the
Meiotic Recombination Program via Checkpoint Pathways Controlled by Global DSB Levels. Mol.
Cell *57*. 10.1016/j.molcel.2014.12.027.

5. Pereira, C., Smolka, M.B., Weiss, R.S., and Brieño-Enríquez, M.A. (2020). ATR signaling in
mammalian meiosis: From upstream scaffolds to downstream signaling. Environmental and
Molecular Mutagenesis *61*, 752–766. 10.1002/em.22401.

6. Gray, S., and Cohen, P.E. (2016). Control of Meiotic Crossovers: From Double-Strand Break
Formation to Designation. Annu. Rev. Genet. *50*, 175–210.

7. Burgoyne, P.S., Mahadevaiah, S.K., and Turner, J.M.A. (2009). The consequences of asynapsis for
mammalian meiosis. Nature Reviews Genetics *10*. 10.1038/nrg2505.

8. Turner, J.M.A., Mahadevaiah, S.K., Ellis, P.J.I., Mitchell, M.J., and Burgoyne, P.S. (2006).
Pachytene asynapsis drives meiotic sex chromosome inactivation and leads to substantial
postmeiotic repression in spermatids. Dev. Cell. 10.1016/j.devcel.2006.02.009.

9. Turner, J.M.A. (2015). Meiotic Silencing in Mammals. Annu. Rev. Genet. *49*. 10.1146/annurev-
genet-112414-055145.

10. Abe, H., Alavattam, K.G., Hu, Y.C., Pang, Q., Andreassen, P.R., Hegde, R.S., and Namekawa, S.H.
(2020). The Initiation of Meiotic Sex Chromosome Inactivation Sequesters DNA Damage Signaling
from Autosomes in Mouse Spermatogenesis. Curr. Biol. *30*, 408-420.e5.

11. Turner, J.M.A. (2007). Meiotic sex chromosome inactivation. Development *134*, 1823–1831.

12. Alavattam, K.G., Abe, H., Sakashita, A., and Namekawa, S.H. (2018). Chromosome spread analyses
of meiotic sex chromosome inactivation. In Methods in Molecular Biology (Humana Press Inc.), pp.
113–129.

13. Royo, H., Polikiewicz, G., Mahadevaiah, S.K., Prosser, H., Mitchell, M., Bradley, A., De Rooij,
D.G., Burgoyne, P.S., and Turner, J.M.A. (2010). Evidence that meiotic sex chromosome
inactivation is essential for male fertility. Curr. Biol. *20*, 2117–2123.

14. Vernet, N., Mahadevaiah, S.K., de Rooij, D.G., Burgoyne, P.S., and Ellis, P.J.I. (2016). Zfy genes
are required for efficient meiotic sex chromosome inactivation (MSCI) in spermatocytes. Hum. Mol.
Genet. *25*, 5300–5310.

15. Pacheco, S., Maldonado-Linares, A., Marcet-Ortega, M., Rojas, C., Martínez-Marchal, A., Fuentes-

Lazaro, J., Lange, J., Jasin, M., Keeney, S., Fernández-Capetillo, O., et al. (2018). ATR is required
to complete meiotic recombination in mice. Nat. Commun. *9*. 10.1038/s41467-018-04851-z.

16. Royo, H., Prosser, H., Ruzankina, Y., Mahadevaiah, S.K., Cloutier, J.M., Baumann, M., Fukuda, T.,
Höög, C., Tóth, A., de Rooij, D.G., et al. (2013). ATR acts stage specifically to regulate multiple
aspects of mammalian meiotic silencing. Genes and Development *27*. 10.1101/gad.219477.113.

17. Abe, H., Yeh, Y.-H., Munakata, Y., Ishiguro, K.-I., Andreassen, P.R., and Namekawa, S.H. (2022).
Active DNA damage response signaling initiates and maintains meiotic sex chromosome
inactivation. Nat. Commun. *13*, 7212.

18. Widger, A., Mahadevaiah, S.K., Lange, J., Elinati, E., Zohren, J., Hirota, T., Pacheco, S.,
Maldonado-Linares, A., Stanzione, M., Ojarikre, O., et al. (2018). ATR is a multifunctional
regulator of male mouse meiosis. Nat. Commun. *9*. 10.1038/s41467-018-04850-0.

19. Cimprich, K.A., and Cortez, D. (2008). ATR: An essential regulator of genome integrity. Nature
Reviews Molecular Cell Biology *9*. 10.1038/nrm2450.

20. Liu, Y., and Smolka, M.B. (2016). TOPBP1 takes RADical command in recombinational DNA
repair. J. Exp. Med. *213*. 10.1084/jem.2132oia2.

21. Mordes, D.A., Glick, G.G., Zhao, R., and Cortez, D. (2008). TopBP1 activates ATR through ATRIP
and a PIKK regulatory domain. Genes and Development *22*, 1478–1489.

22. Liu, Y., Cussiol, J.R., Dibitetto, D., Sims, J.R., Twayana, S., Weiss, R.S., Freire, R., Marini, F.,
Pellicioli, A., and Smolka, M.B. (2017). TOPBP1Dpb11 plays a conserved role in homologous
recombination DNA repair through the coordinated recruitment of 53BP1Rad9. J. Cell Biol. *216*,
623–639.

23. Leimbacher, P.A., Jones, S.E., Shorrocks, A.M.K., de Marco Zompit, M., Day, M., Blaauwendraad,
J., Bundschuh, D., Bonham, S., Fischer, R., Fink, D., et al. (2019). MDC1 Interacts with TOPBP1 to
Maintain Chromosomal Stability during Mitosis. Mol. Cell *74*, 571-583.e8.

24. Cescutti, R., Negrini, S., Kohzaki, M., and Halazonetis, T.D. (2010). TopBP1 functions with 53BP1
in the G1 DNA damage checkpoint. EMBO J. *29*, 3723–3732.

25. Leung, C.C.Y., Gong, Z., Chen, J., and Glover, J.N.M. (2011). Molecular basis of BACH1/FANCJ
recognition by TopBP1 in DNA replication checkpoint control. J. Biol. Chem. *286*, 4292–4301.

26. Delacroix, S., Wagner, J.M., Kobayashi, M., Yamamoto, K.I., and Karnitz, L.M. (2007). The Rad9-
Hus1-Rad1 (9-1-1) clamp activates checkpoint signaling via TopBP1. Genes and Development *21*,
1472–1477.

27. Blackford, A.N., Nieminuszczy, J., Schwab, R.A., Galanty, Y., Jackson, S.P., and Niedzwiedz, W.
(2015). TopBP1 Interacts with BLM to Maintain Genome Stability but Is Dispensable for
Preventing BLM Degradation. Mol. Cell *57*, 1133–1141.

28. Bigot, N., Day, M., Baldock, R.A., Watts, F.Z., Oliver, A.W., and Pearl, L.H. (2019).
Phosphorylation-mediated interactions with topbp1 couple 53bp1 and 9-1-1 to control the g1 DNA
damage checkpoint. Elife *8*, 1–28.

29.   Pereira, C., Arroyo-Martinez, G.A., Guo, M.Z., Downey, M.S., Kelly, E.R., Grive, K.J.,
891   Mahadevaiah, S.K., Sims, J.R., Faca, V.M., Tsai, C., et al. (2022). Multiple 9-1-1 complexes
892   promote homolog synapsis, DSB repair, and ATR signaling during mammalian meiosis. Elife *11*.
893   10.7554/ELIFE.68677.

894   30.   Manke, I.A., Lowery, D.M., Nguyen, A., and Yaffe, M.B. (2003). BRCT repeats as phosphopeptide-
895   binding modules involved in protein targeting. Science *302*, 636–639.

896   31.   Rodriguez, M., Yu, X., Chen, J., and Songyang, Z. (2003). Phosphopeptide binding specificities of
897   BRCA1 COOH-terminal (BRCT) domains. J. Biol. Chem. *278*, 52914–52918.

898   32.   Jeon, Y., Park, M.K., Kim, S.M., Bae, J.S., Lee, C.W., and Lee, H. (2019). TopBP1 deficiency
899   impairs the localization of proteins involved in early recombination and results in meiotic
900   chromosome defects during spermatogenesis. Biochem. Biophys. Res. Commun. *508*.
901   10.1016/j.bbrc.2018.12.001.

902   33.   Perera, D., Perez-Hidalgo, L., Moens, P.B., Reini, K., Lakin, N., Syväoja, J.E., San-Segundo, P.A.,
903   and Freire, R. (2004). TopBP1 and ATR Colocalization at Meiotic Chromosomes: Role of
904   TopBP1/Cut5 in the Meiotic Recombination Checkpoint. Mol. Biol. Cell *15*, 1568–1579.

905   34.   ElInati, E., Russell, H.R., Ojarikre, O.A., Sangrithi, M., Hirota, T., De Rooij, D.G., McKinnon, P.J.,
906   and Turner, J.M.A. (2017). DNA damage response protein TOPBP1 regulates X chromosome
907   silencing in the mammalian germ line. Proc. Natl. Acad. Sci. U. S. A. *114*, 12536–12541.

908   35.   Parrilla-Castellar, E.R., Arlander, S.J.H., and Karnitz, L. (2004). Dial 9–1–1 for DNA damage: the
909   Rad9–Hus1–Rad1 (9–1–1) clamp complex. DNA Repair *3*, 1009–1014.

910   36.   Broering, T.J., Alavattam, K.G., Sadreyev, R.I., Ichijima, Y., Kato, Y., Hasegawa, K., Camerini-
911   Otero, R.D., Lee, J.T., Andreassen, P.R., and Namekawa, S.H. (2014). BRCA1 establishes DNA
912   damage signaling and pericentric heterochromatin of the X chromosome in male meiosis. J. Cell
913   Biol. *205*, 663–675.

914   37.   Reini, K., Uitto, L., Perera, D., Moens, P.B., Freire, R., and Syväoja, J.E. (2004). TopBPI localises
915   to centrosomes in mitosis and to chromosome cores in meiosis. Chromosoma *112*. 10.1007/s00412-
916   004-0277-5.

917   38.   Moens, P.B., Tarsounas, M., Morita, T., Habu, T., Rottinghaus, S.T., Freire, R., Jackson, S.P.,
918   Barlow, C., and Wynshaw-Boris, A. (1999). The association of ATR protein with mouse meiotic
919   chromosome cores. Chromosoma *108*. 10.1007/s004120050356.

920   39.   Xu, Y., and Qiao, H. (2021). A Hypothesis: Linking Phase Separation to Meiotic Sex Chromosome
921   Inactivation and Sex-Body Formation. Front Cell Dev Biol *9*, 674203.

922   40.   Turner, J.M.A., Aprelikova, O., Xu, X., Wang, R., Kim, S., Chandramouli, G.V.R., Barrett, J.C.,
923   Burgoyne, P.S., and Deng, C.-X. (2004). BRCA1, histone H2AX phosphorylation, and male meiotic
924   sex chromosome inactivation. Curr. Biol. *14*, 2135–2142.

925   41.   Shin, Y.-H., Choi, Y., Erdin, S.U., Yatsenko, S.A., Kloc, M., Yang, F., Jeremy Wang, P., Meistrich,
926   M.L., and Rajkovic, A. (2010). Hormad1 Mutation Disrupts Synaptonemal Complex Formation,
927   Recombination, and Chromosome Segregation in Mammalian Meiosis. PLoS Genetics *6*, e1001190.
928   10.1371/journal.pgen.1001190.

42.   Ichijima, Y., Ichijima, M., Lou, Z., Nussenzweig, A., Daniel Camerini-Otero, R., Chen, J.,
Andreassen, P.R., and Namekawa, S.H. (2011). MDC1 directs chromosome-wide silencing of the
sex chromosomes in male germ cells. Genes and Development *25*, 959–971.

43.   O'Driscoll, M., Dobyns, W.B., Van Hagen, J.M., and Jeggo, P.A. (2007). Cellular and clinical
impact of haploinsufficiency for genes involved in ATR signaling. Am. J. Hum. Genet. *81*.
10.1086/518696.

44.   Brown, E.J., and Baltimore, D. (2000). ATR disruption leads to chromosomal fragmentation and
early embryonic lethality. Genes and Development *14*. 10.1101/gad.14.4.397.

45.   De Klein, A., Muijtjens, M., Van Os, R., Verhoeven, Y., Smit, B., Carr, A.M., Lehmann, A.R., and
Hoeijmakers, J.H.J. (2000). Targeted disruption of the cell-cycle checkpoint gene ATR leads to
early embryonic lethality in mice. Curr. Biol. *10*. 10.1016/S0960-9822(00)00447-4.

46.   O'Driscoll, M. (2009). Mouse models for ATR deficiency. DNA Repair *8*.
10.1016/j.dnarep.2009.09.001.

47.   Yamane, K., Wu, X., and Chen, J. (2002). A DNA damage-regulated BRCT-containing protein,
TopBP1, is required for cell survival. Mol. Cell. Biol. *22*, 555–566.

48.   Jeon, Y., Ko, E., Lee, K.Y., Ko, M.J., Park, S.Y., Kang, J., Jeon, C.H., Lee, H., and Hwang, D.S.
(2011). TopBP1 deficiency causes an early embryonic lethality and induces cellular senescence in
primary cells. J. Biol. Chem. *286*, 5414–5422.

49.   Zhou, Z.W., Liu, C., Li, T.L., Bruhn, C., Krueger, A., Min, W.K., Wang, Z.Q., and Carr, A.M.
(2013). An Essential Function for the ATR-Activation-Domain (AAD) of TopBP1 in Mouse
Development and Cellular Senescence. PLoS Genet. *9*. 10.1371/journal.pgen.1003702.

50.   Inselman, A., Eaker, S., and Handel, M.A. (2003). Temporal expression of cell cycle-related
proteins during spermatogenesis: establishing a timeline for onset of the meiotic divisions.
Cytogenet. Genome Res. *103*, 277–284.

51.   Falck, J., Coates, J., and Jackson, S.P. (2005). Conserved modes of recruitment of ATM, ATR and
DNA-PKcs to sites of DNA damage. Nature *434*, 605–611.

52.   Blackford, A.N., and Jackson, S.P. (2017). ATM, ATR, and DNA-PK: The Trinity at the Heart of
the DNA Damage Response. Mol. Cell *66*, 801–817.

53.   Maréchal, A., and Zou, L. (2013). DNA damage sensing by the ATM and ATR kinases. Cold Spring
Harb. Perspect. Biol. *5*. 10.1101/cshperspect.a012716.

54.   Hartwell, L.H., and Kastan, M.B. (1994). Cell cycle control and cancer. Science *266*, 1821–1828.

55.   Shiloh, Y. (2003). ATM and related protein kinases: safeguarding genome integrity. Nat. Rev.
Cancer *3*, 155–168.

56.   Lanz, M.C., Dibitetto, D., and Smolka, M.B. (2019). DNA damage kinase signaling: checkpoint and
repair at 30 years. EMBO J. *38*, e101801.

57.   Kwon, M., Leibowitz, M.L., and Lee, J.-H. (2020). Small but mighty: the causes and consequences

of micronucleus rupture. Experimental & Molecular Medicine *52*, 1777–1786. 10.1038/s12276-020-
00529-z.

58. Wang, J., Gong, Z., and Chen, J. (2011). MDC1 collaborates with TopBP1 in DNA replication
checkpoint control. J. Cell Biol. *193*, 267–273.

59. Balbo Pogliano, C., Ceppi, I., Giovannini, S., Petroulaki, V., Palmer, N., Uliana, F., Gatti, M.,
Kasaciunaite, K., Freire, R., Seidel, R., et al. (2022). The CDK1-TOPBP1-PLK1 axis regulates the
Bloom's syndrome helicase BLM to suppress crossover recombination in somatic cells. Sci Adv *8*,
eabk0221.

60. Wang, J., Chen, J., and Gong, Z. (2013). TopBP1 controls BLM protein level to maintain genome
stability. Mol. Cell *52*, 667–678.

61. Carpenter, A.T. (1979). Recombination nodules and synaptonemal complex in recombination-
defective females of Drosophila melanogaster. Chromosoma *75*, 259–292.

62. Abe, H., Alavattam, K.G., Kato, Y., Castrillon, D.H., Pang, Q., Andreassen, P.R., and Namekawa,
S.H. (2018). CHEK1 coordinates DNA damage signaling and meiotic progression in the male
germline of mice. Hum. Mol. Genet. *27*, 1136–1149.

63. Thompson, A., Schäfer, J., Kuhn, K., Kienle, S., Schwarz, J., Schmidt, G., Neumann, T., Johnstone,
R., Mohammed, A.K.A., and Hamon, C. (2003). Tandem mass tags: a novel quantification strategy
for comparative analysis of complex protein mixtures by MS/MS. Anal. Chem. *75*, 1895–1904.

64. Sims, J.R., Faça, V.M., Pereira, C., Ascenção, C., Comstock, W., Badar, J., Arroyo-Martinez, G.A.,
Freire, R., Cohen, P.E., Weiss, R.S., et al. (2022). Phosphoproteomics of ATR signaling in mouse
testes. Elife *11*. 10.7554/eLife.68648.

65. Cohen, S., Puget, N., Lin, Y.-L., Clouaire, T., Aguirrebengoa, M., Rocher, V., Pasero, P., Canitrot,
Y., and Legube, G. (2018). Senataxin resolves RNA:DNA hybrids forming at DNA double-strand
breaks to prevent translocations. Nat. Commun. *9*, 533.

66. Yeo, A.J., Becherel, O.J., Luff, J.E., Graham, M.E., Richard, D., and Lavin, M.F. (2015). Senataxin
controls meiotic silencing through ATR activation and chromatin remodeling. Cell Discovery *1*.
10.1038/celldisc.2015.25.

67. Becherel, O.J., Yeo, A.J., Stellati, A., Heng, E.Y.H., Luff, J., Suraweera, A.M., Woods, R., Fleming,
J., Carrie, D., McKinney, K., et al. (2013). Senataxin Plays an Essential Role with DNA Damage
Response Proteins in Meiotic Recombination and Gene Silencing. PLoS Genet. *9*.
10.1371/journal.pgen.1003435.

68. Ichijima, Y., Sin, H.S., and Namekawa, S.H. (2012). Sex chromosome inactivation in germ cells:
Emerging roles of DNA damage response pathways. Cellular and Molecular Life Sciences *69*.
10.1007/s00018-012-0941-5.

69. Lau, X., Munusamy, P., Ng, M.J., and Sangrithi, M. (2020). Single-Cell RNA Sequencing of the
Cynomolgus Macaque Testis Reveals Conserved Transcriptional Profiles during Mammalian
Spermatogenesis. Dev. Cell *54*, 548-566.e7.

70. Khalil, A.M., Boyar, F.Z., and Driscoll, D.J. (2004). Dynamic histone modifications mark sex

chromosome inactivation and reactivation during mammalian spermatogenesis. Proc. Natl. Acad.
Sci. U. S. A. *101*, 16583–16587.

71. Alavattam, K.G., Maezawa, S., Andreassen, P.R., and Namekawa, S.H. (2022). Meiotic sex
chromosome inactivation and the XY body: a phase separation hypothesis. Cellular and Molecular
Life Sciences *79*. 10.1007/s00018-021-04075-3.

72. Jung, M., Wells, D., Rusch, J., Ahmad, S., Marchini, J., Myers, S.R., and Conrad, D.F. (2019).
Unified single-cell analysis of testis gene regulation and pathology in five mouse strains. eLife *8*.
10.7554/elife.43966.

73. Grive, K.J., Hu, Y., Shu, E., Grimson, A., Elemento, O., Grenier, J.K., and Cohen, P.E. (2019).
Dynamic transcriptome profiles within spermatogonial and spermatocyte populations during
postnatal testis maturation revealed by single-cell sequencing. PLoS Genet. *15*, e1007810.

74. Hirota, T., Blakeley, P., Sangrithi, M.N., Mahadevaiah, S.K., Encheva, V., Snijders, A.P., ElInati,
E., Ojarikre, O.A., de Rooij, D.G., Niakan, K.K., et al. (2018). SETDB1 Links the Meiotic DNA
Damage Response to Sex Chromosome Silencing in Mice. Dev. Cell *47*, 645-659.e6.

75. Modzelewski, A.J., Holmes, R.J., Hilz, S., Grimson, A., and Cohen, P.E. (2012). AGO4 regulates
entry into meiosis and influences silencing of sex chromosomes in the male mouse germline. Dev.
Cell *23*, 251–264.

76. Chen, Y., Zheng, Y., Gao, Y., Lin, Z., Yang, S., Wang, T., Wang, Q., Xie, N., Hua, R., Liu, M., et
al. (2018). Single-cell RNA-seq uncovers dynamic processes and critical regulators in mouse
spermatogenesis. Cell Res. *28*, 879–896.

77. Greenlee, A.R., Shiao, M.-S., Snyder, E., Buaas, F.W., Gu, T., Stearns, T.M., Sharma, M.,
Murchison, E.P., Puente, G.C., and Braun, R.E. (2012). Deregulated sex chromosome gene
expression with male germ cell-specific loss of Dicer1. PLoS One *7*, e46359.

78. Bennett, C.L., and La Spada, A.R. (2018). Senataxin, A Novel Helicase at the Interface of RNA
Transcriptome Regulation and Neurobiology: From Normal Function to Pathological Roles in Motor
Neuron Disease and Cerebellar Degeneration. In RNA Metabolism in Neurodegenerative Diseases,
R. Sattler and C. J. Donnelly, eds. (Springer International Publishing), pp. 265–281.

79. Santos-Pereira, J.M., and Aguilera, A. (2015). R loops: new modulators of genome dynamics and
function. Nat. Rev. Genet. *16*, 583–597.

80. Marnef, A., and Legube, G. (2020). m6A RNA modification as a new player in R-loop regulation.
Nat. Genet. *52*, 27–28.

81. Rockmill, B., Fung, J.C., Branda, S.S., and Roeder, G.S. (2003). The Sgs1 helicase regulates
chromosome synapsis and meiotic crossing over. Curr. Biol. *13*, 1954–1962.

82. Holloway, J.K., Morelli, M.A., Borst, P.L., and Cohen, P.E. (2010). Mammalian BLM helicase is
critical for integrating multiple pathways of meiotic recombination. J. Cell Biol. *188*, 779–789.

83. Lenzi, M.L., Smith, J., Snowden, T., Kim, M., Fishel, R., Poulos, B.K., and Cohen, P.E. (2005).
Extreme heterogeneity in the molecular events leading to the establishment of chiasmata during
meiosis i in human oocytes. Am. J. Hum. Genet. *76*, 112–127.

84.   Danielsen, J.M.R., Larsen, D.H., Freire, R., Falck, J., Bartek, J., Lukas, J., and Others (2009).
1042   HCLK2 is required for activity of the DNA damage response kinase ATR. J. Biol. Chem. *284*,
1043   4140–4147.

1044   85.   Wojtasz, L., Cloutier, J.M., Baumann, M., Daniel, K., Varga, J., Fu, J., Anastassiadis, K., Stewart,
1045   A.F., Reményi, A., Turner, J.M.A., et al. (2012). Meiotic DNA double-strand breaks and
1046   chromosome asynapsis in mice are monitored by distinct HORMAD2-independent and -dependent
1047   mechanisms. Genes Dev. *26*, 958–973.

1048   86.   Holloway, J.K., Sun, X., Yokoo, R., Villeneuve, A.M., and Cohen, P.E. (2014). Mammalian
1049   CNTD1 is critical for meiotic crossover maturation and deselection of excess precrossover sites. J.
1050   Cell Biol. *205*, 633–641.

1051   87.   Kakarougkas, A., Ismail, A., Katsuki, Y., Freire, R., Shibata, A., and Jeggo, P.A. (2013). Co-
1052   operation of BRCA1 and POH1 relieves the barriers posed by 53BP1 and RAP80 to resection.
1053   Nucleic Acids Res. *41*, 10298–10311.

1054   88.   Modzelewski, A.J., Hilz, S., Crate, E.A., Schweidenback, C.T.H., Fogarty, E.A., Grenier, J.K.,
1055   Freire, R., Cohen, P.E., and Grimson, A. (2015). Dgcr8 and Dicer are essential for sex chromosome
1056   integrity during meiosis in males. J. Cell Sci. *128*, 2314–2327.

1057   89.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk,
1058   A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*,
1059   3573-3587.e29.

1060   90.   Leek, J.T., Johnson, W.E., Parker, H.S., Fertig, E.J., and Jaffe, A.E. sva: Surrogate Variable
1061   Analysis R package version 3.10. 0. DOI10 B.

1062   91.   Aran, D., Looney, A.P., Liu, L., Wu, E., Fong, V., Hsu, A., Chak, S., Naikawadi, R.P., Wolters, P.J.,
1063   Abate, A.R., et al. (2019). Reference-based analysis of lung single-cell sequencing reveals a
1064   transitional profibrotic macrophage. Nat. Immunol. *20*, 163–172.

1065   92.   Wickham, H. (2007). Reshaping Data with the reshape Package. J. Stat. Softw. *21*, 1–20.

1066   93.   Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for the
1067   integration of genomic datasets with the R/Bioconductor package biomaRt. Nat. Protoc. *4*, 1184–
1068   1191.

1069   94.   Kassambara, A. (2019). GGPlot2 Essentials: Great Data Visualization in R (Independently
1070   Published).

1071   95.   Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,
1072   Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the tidyverse. J. Open Source Softw. *4*,
1073   1686.

1074   96.   Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner,
1075   M., Loh, P.-R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-cell
1076   data with Harmony. Nat. Methods *16*, 1289–1296.

1077   97.   Carlson, M., Falcon, S., Pages, H., and Li, N. org. Mm. eg. db: Genome wide annotation for Mouse.
1078   R package version.

1079   98.   Wei, T., and Simko, V. (2017). R package "corrplot": Visualization of a Correlation Matrix
1080   (Version 0.84).

1081   99.   Balmus, G., Zhu, M., Mukherjee, S., Lyndaker, A.M., Hume, K.R., Lee, J., Riccio, M.L., Reeves,
1082   A.P., Sutter, N.B., Noden, D.M., et al. (2012). Disease severity in a mouse model of ataxia
1083   telangiectasia is modulated by the DNA damage checkpoint gene Hus1. Hum. Mol. Genet. *21*,
1084   3408–3420.

1085   100.   Kolas, N.K., Svetlanov, A., Lenzi, M.L., Macaluso, F.P., Lipkin, S.M., Liskay, R.M., Greally, J.,
1086   Edelmann, W., and Cohen, P.E. (2005). Localization of MMR proteins on meiotic chromosomes in
1087   mice indicates distinct functions during prophase I. J. Cell Biol. *171*, 447–458.

1088   101.   Eng, J.K., Jahan, T.A., and Hoopmann, M.R. (2013). Comet: An open-source MS/MS sequence
1089   database search tool. PROTEOMICS *13*, 22–24. 10.1002/pmic.201200439.

1090   102.   Bastos de Oliveira, F.M., Kim, D., Cussiol, J.R., Das, J., Jeong, M.C., Doerfler, L., Schmidt, K.H.,
1091   Yu, H., and Smolka, M.B. (2015). Phosphoproteomics reveals distinct modes of Mec1/ATR
1092   signaling during DNA replication. Mol. Cell *57*, 1124–1132.

1093   103.   Rodríguez-Casuriaga, R., Santiñaque, F.F., Folle, G.A., Souza, E., López-Carro, B., and Geisinger,
1094   A. (2014). Rapid preparation of rodent testicular cell suspensions and spermatogenic stages
1095   purification by flow cytometry using a novel blue-laser-excitable vital dye. MethodsX *1*, 239–243.

1096

Figure legends

1098

Figure 1. A new TOPBP1 mutant separating its role in fertility from organismal viability.

A) Schematic showing mutations in the *Topbp1 B5* allele. B) Body mass (*Topbp1$^{+/+}$* Mean = 25.26, SD =

2.38; *Topbp1$^{B5/B5}$* Mean = 26.43, SD = 2.28, n = 9) and C) appearance of *Topbp1$^{B5/B5}$* and *Topbp1$^{+/+}$*

littermate mice. D) Effect of full body IR (7 Gy) on changes in body mass of *Topbp1$^{B5/B5}$* and *Topbp1$^{+/+}$*

littermate mice E) Breeding scheme and resulting litters. F) and G) Comparison of testes size (*Topbp1$^{+/+}$*

Mean = 0.038, SD = 0.006; *Topbp1$^{B5/B5}$* Mean = 0.011, SD = 0.004, n = 9), and H) sperm count, of

Topbp1$^{B5/B5}$ and *Topbp1$^{+/+}$* littermate mice (*Topbp1$^{+/+}$* Mean = 2.1 x 10^7, SD = 6x 10^6; *Topbp1$^{B5/B5}$* Mean =

0.0, SD = 0.0, n = 9). I) H&E-stained histological testes sections displaying loss of cellularity in *Topbp1$^{B5/B5}$*.

Green arrow = spermatogonia, Red arrow = healthy spermatocyte, Blue arrow = dying spermatocyte. J) and

K) TUNEL assay performed on histological testes sections (*Topbp1$^{+/+}$* Mean = 0.36, SD = 0.15; *Topbp1$^{B5/B5}$*

Mean = 4.80, SD = 0.43, n = 3). L) Meiotic spreads stained for SYCP3, SYCP1 and H1t. **** = p< 0.0001,

n = number of mice. p-Values were calculated using Welch's unpaired *t*-test in GraphPad.

1111

Figure 2. *Topbp1$^{B5/B5}$* MEFs display no detectable DDR defects.

A) MEFs obtained from littermate mice of indicated genotypes were subjected to clonogenic survival assay

in the indicated concentrations of hydroxyurea, or C) phleomycin. B) and D) Quantification of clonogenic

survival assays from A and C, respectively, performed in biological and experimental triplicates. D)

Western blot for indicated DDR markers in MEFs obtained from *Topbp1$^{B5/B5}$* and *Topbp1$^{+/+}$* littermate mice.

The data from MEFs were performed using littermate pairs and validated using a second pair of *Topbp1$^{B5/B5}$*

and *Topbp1$^{+/+}$* littermate mice. E) Immunofluorescence of MDC1 and phosphoH2AX_S139-stained nuclei

from *Topbp1$^{B5/B5}$* and *Topbp1$^{+/+}$* MEFs treated with IR (7 Gy).

1120

Figure 3. Markers of DNA repair, synapsis, sex body formation and TOPBP1 localization are mostly

normal in *Topbp1$^{B5/B5}$* spermatocytes.

A) Meiotic spreads showing *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* spermatocytes stained with SYCP3 and γH2AX,

prepared as described in methods. B) Quantification of γH2AX-stained pachytene spreads, upper

graph autosomal chromosomes (each dot represents the average of signal from all autosomes in

each mouse) and bottom, XY body. C) Meiotic spreads showing *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* pachytene

spermatocytes stained with SYCP3 and RAD51. D) Quantification of RAD51 foci/cell of mid-pachytene

meiotic spreads. E) Meiotic spreads showing *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* pachytene spermatocytes stained

with SYCP3 and SYCP1. F) Meiotic spreads showing *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* pachytene spermatocytes

stained with SYCP3 and TOPBP1. Quantification of TOPBP1-stained pachytene spreads G) X and Y

chromosome cores, and H) X and Y chromatin loops. n = number of mice. p-Values were calculated

using Welch's unpaired *t*-test in GraphPad.

1133

Figure 4. *Topbp1^{B5/B5}* spermatocytes display normal localization of ATR, BRCA1, MDC1 and

pMDC1-T4.

A-H) Meiotic spreads stained for indicated proteins and related quantification in *Topbp1^{+/+}* and *Topbp1^{B5/B5}*

pachytene spermatocytes. I) Table summarizing the normal or disrupted ATR and TOPBP1-

dependent events during male fertility accessed in *Topbp1^{B5/B5}*. n = number of mice. p-Values were

calculated using Welch's unpaired *t*-test in GraphPad.

1140

Figure 5. Defective SETX phosphorylation and localization in *Topbp1^{B5/B5}* spermatocytes.

A) Scatter plot of phosphoproteomic datasets corresponding to *Topbp1^{+/+}*/*Topbp1^{B5/B}* (Y axis) and

Topbp1^{+/+}(vehicle)/*Topbp1^{+/+}*(AZ20) (X axis) from whole testes of mice. B) SETX phosphopeptides

identified in the *Topbp1^{+/+}*/*Topbp1^{B5/B}* phosphoproteomic experiment shown in A. Red: reduced in

Topbp1^{B5/B} mutant; Blue: unchanged. C) Meiotic spreads showing pachytene spermatocytes from *Topbp1^{+/+}*

and *Topbp1^{B5/B5}* mice stained with SYCP3 and SETX in regular immunofluorescence, and D) 3D-SIM

analysis of meiotic spread described in C. E) and F) Quantification of the intensity of SETX chromatin

loops and cores of the XY chromosomes, respectively. * p < 0.05, n = number of mice. p-Values were

calculated using Welch's unpaired *t*-test in GraphPad.

1150

Figure 6. Single cell RNAseq reveals that *Topbp1^{B5/B5}* spermatocytes initiate MSCI but fail to promote

full silencing.

A) scRNAseq workflow for isolation and purification of single cells for RNA-seq. B) UMAP analysis of

sub-clusters captured in the scRNAseq. C) UMAP analysis of sub-clusters captured in the scRNAseq of

Topbp1^{+/+}. D) UMAP analysis of sub-clusters captured in the scRNAseq of *Topbp1^{+/+}* (gray) and

Topbp1^{B5/B5} (blue). E) Violin plots displaying the ratio of the average expression of X chromosome genes

by the average expression of chromosome 9 genes at different stages of spermatogenesis for *Topbp1^{+/+}* and

Topbp1^{B5/B5} cells.

1159

1160   **Figure 7. Topbp1 regulates silencing dynamics of X genes at the Spermatocyte 3 stage.**

1161   A) Illustration of the gene markers used to define spermatocyte 1 as leptotene, spermatocyte 2 as zygotene,

1162   and spermatocyte 3 as pachytene; hypothetical examples illustrating the categorization of transitions in

1163   silencing dynamics between the stages of pre-leptotene (PL), spermatocyte 1 (SP1), spermatocyte 2 (SP2),

spermatocyte 3 (SP3). B) Number of genes in each of the categories described in A, during the different stage transitions. C) Examples of genes with altered silencing dynamics in the $Topbp1^{B5/B5}$, red = reduced silencing, blue = unaltered silencing and green = increased silencing D) Scatter plot showing the difference in RNA level between $Topbp1^{+/+}$ and $Topbp1^{B5/B5}$ for each of the indicated stage transitions. E) Scatter plot showing expression level of X-chromosome genes, normalized to pre-leptotene levels, in $Topbp1^{+/+}$ (gray) and $Topbp1^{B5/B5}$ (blue) at SP3. F) Graph plotting expression levels of X-chromosome genes, normalized to pre-leptotene levels, in $Topbp1^{+/+}$ (Y axis) and $Topbp1^{B5/B5}$ (X axis) and split in three categories based on the severity of silencing defect. G) Box plot showing PL expression levels of X-chromosome genes in each of the categories of silencing defect severity shown in F.

Figure 8. A new TOPBP1 mutant separates XY silencing from sex body formation.

Schematic of sub-stages of meiotic prophase I. In Wild-type mice, MSCI initiates following the accumulation of the DDR proteins at the XY chromosomes. During mid-pachytene, the XY body is fully formed, and transcription is restricted to the autosomes. In Atr or $Topbp1$ CKOs, the sex body is not formed, and the DDR proteins are not sequestered to the XY. Asynapsis events and transcription of toxic genes at the sex chromosomes are observed, triggering mid-pachytene arrest. In $Topbp1^{B5/B5}$, MSCI initiates, the sex body is normally formed with normal recruitment of DDR proteins to the X and Y chromosomes, yet cells fail in the reinforcement/maintenance of silencing. Cells progress through mid-pachytene but not into diplotene.

 Supplemental material

 Supplemental Figure legends

 Supplemental Figure 1. Genotyping of *Topbp1$^{B5/B5}$* mice.

 Sanger sequencing confirms the eight charge reversal point mutations on the BRCT 5 domain of TOPBP1

 from homozygous *Topbp1$^{B5/B5}$* mice.

 Supplemental Figure 2. Survival assays of *Topbp1$^{B5/B5}$* MEFs.

 A) MEFs obtained from littermate mice (second independent litter) of indicated genotypes were subjected

 to clonogenic survival assay in the indicated concentrations of hydroxyurea, or B) phleomycin.

 Supplemental Figure 3. Absence of an increased number of micronuclei in *Topbp1$^{B5/B5}$* MEFs.

 Immunofluorescence showing DAPI-stained nuclei and micronuclei quantification of *Topbp1$^{+/+}$* and

 Topbp1$^{B5/B5}$ A) MEFs #1 and B) MEFs #2. (MEFs #2 is a second independent litter).

 Supplemental Figure 4. TOPBP1-B5 shows disruption in binding 53BP1 and BLM.

 A) HEK-293T cells were transfected with plasmids expressing wild-type TOPBP1 or indicated BRCT

 mutants (K155A - BRCT 1, K250A - BRCT 2, K704A - BRCT 5, K1713A - BRCT 7, and B5 - 8 mutations

 in BRCT 5), with a 3xFlag tag. After cell lysis and anti-Flag IP, lysate and elution were analyzed by western

 blot for the indicated proteins. B) IP-mass spectrometry of flagTOPBP1 wild type versus flag-TOPBP1-B5

 was performed in silac-labeled HEK-293T cells. Cell transfections, lysis, protein IP, and elution were

 performed as described in A. From left to right, the TOPBP1 interactors were plotted from highest to lowest

 binding. The size of circles represents the number of peptide hits.

 Supplemental Figure 5. *Topbp1$^{B5/B5}$* MEFs show lower TOPBP1 protein abundance than *Topbp1$^{+/+}$*

 MEFs.

 Western blot probing TOPBP1 in MEFs, *Topbp1$^{+/+}$*, *Topbp1$^{B5/+}$,* and *Topbp1$^{B5/B5}$* in two biological

 replicates.

 Supplemental Figure 6. *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes show normal HORMAD1,

 HORMAD2, pHORMAD1 and pCHK1_S345 localization and intensities.

 A) *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* pachytene spermatocytes stained with HORMAD1 and SYCP3, B)

 HORMAD2 and SYCP3, and C) pMDC1_T4 and SYCP3. D) Quantification of *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$*

1218 mid-pachytene meiotic spreads stained with pMDC1_T4. E) *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* mid-pachytene

1219 spermatocytes stained with pCHK1_S345 and SYCP3. F) Quantification of *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* mid-

1220 pachytene meiotic spreads stained with pCHK1_S345. G) *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* mid-pachytene

1221 spermatocytes stained with pHORMAD1_S271 and SYCP3. H) Quantification of *Topbp1$^{+/+}$* and

1222 *Topbp1$^{B5/B5}$* mid-pachytene meiotic spreads stained with pHORMAD1_S271. n = number of mice.

1223

1224 **Supplemental Figure 7. *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes show an increased number of**

1225 **MLH1 foci/cell but no difference in the number of MHL3 foci/cell.**

1226 A) Representative images and quantification of *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes

1227 stained with MLH1 and SYCP3, and B) MLH3. n = number of mice. *** $p < 0.001$. p-Values were

1228 calculated using Welch's unpaired t-test in GraphPad.

1229

1230 **Supplemental Figure 8. *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes form a normal sex body.**

1231 *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes stained with DAPI and SYCP3.

1232

Supplemental Figure 9. *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes show subtle defects/delays in the

spreading of γH2AX at the XY chromosomes.

A) Quantification and representative images of *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes showing normal

spreading of γH2AX in the XY chromosomes B) showing a defect/delay in the spreading of γH2AX at the

PAR, C) showing a defect/delay in the spreading of γH2AX at the pericentromeric region of the Y

chromosome, D) showing a spreading of γH2AX into an autosomal chromosome fused to the X

chromosome, and E) showing other types of defects/delays in the spreading of γH2AX into the XY

chromosomes. F) Percentage of the previously described defects found in *Topbp1$^{+/+}$* or G) *Topbp1$^{B5/B5}$* mid-

pachytene spermatocytes. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and **** $p < 0.0001$. p-Values were

calculated using Welch's unpaired t-test in GraphPad. n = number of mice.

1243

1244 **Supplemental Figure 10. TOPBP1 is normally localized in all stages of prophase I in *Topbp1$^{B5/B5}$*.**

1245 Meiotic spreads showing *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* spermatocytes stained with SYCP3 and TOPBP1.

1246

1247 **Supplemental Figure 11. 53BP1 localization in *Topbp1$^{B5/B5}$* mid-pachytene spermatocytes.**

1248 Meiotic spreads and quantification of pachytene spermatocytes from *Topbp1$^{+/+}$* and *Topbp1$^{B5/B5}$* mice

1249 stained with SYCP3 and 53BP1. n = number of mice.

1250

Supplemental Figure 12. *Topbp1^{B5/B5}* pachytene spermatocytes do not exhibit any difference in CHD4, Sumo2-3, and USP7 intensities and localization compared to *Topbp1^{+/+}*.

A) Representative images and quantification of *Topbp1^{+/+}* and *Topbp1^{B5/B5}* mid-pachytene spermatocytes stained with CHD4 and SYCP3, B) Sumo2.3 and SYCP3, and C) USP7 and SYCP3. n = number of mice.

Supplemental Figure 13. *Topbp1^{B5/B5}* pachytene spermatocytes show no difference in H3K9ac or K3K9me3 staining compared to *Topbp1^{+/+}*.

A) Representative images and quantification of *Topbp1^{+/+}* and *Topbp1^{B5/B5}* mid-pachytene spermatocytes stained with H3K9ac and SYCP3. B) Representative images and quantification of *Topbp1^{+/+}* and *Topbp1^{B5/B5}* early, mid, or late-pachytene spermatocytes stained with H3K9me3 and SYCP3 *Topbp1^{B5/B5}* n = number of mice.

Supplemental Figure 14. *Topbp1^{B5/B5}* pachytene spermatocytes display no difference in H3K27ac, H3K36me3 or K3K4me3 staining compared to *Topbp1^{+/+}*.

A) Representative images and quantification of *Topbp1^{+/+}* and *Topbp1^{B5/B5}* mid-pachytene spermatocytes stained with H3K27ac and SYCP3, and B) H3K36me3 and SYCP3. C) Representative images and quantification of *Topbp1^{+/+}* and *Topbp1^{B5/B5}* early and mid-pachytene spermatocytes stained with H3K4me3 and SYCP3. n = number of mice.

Supplemental Figure 15. *Topbp1^{B5/B5}* pachytene spermatocytes display no difference in H4K16ac, H4ac or H2AK166ub staining compared to *Topbp1^{+/+}*.

A) Representative images and quantification of *Topbp1^{+/+}* and *Topbp1^{B5/B5}* mid-pachytene spermatocytes stained with H4K16ac and SYCP3, B) H4 and SYCP3, and C) H2AK166ub and SYCP3. n = number of mice.

Supplemental Figure 16. From early pachytene to beginning of late pachytene, *Topbp1^{B5/B5}* spermatocytes exhibit the same intensity and localization pattern of H3K4me1 compared to *Topbp1^{+/+}*.

Representative images of *Topbp1^{+/+}* and *Topbp1^{B5/B5}* early, mid and late pachytene spermatocytes stained with H3K4me1 and SYCP3.

Supplemental figure 17. *Topbp1^{B5/B5}* pachytene spermatocytes exhibit no differences in the intensity or localization patterns of RNApol2, pRNApol2_Ser2 or pRNApol2_ser5 compared to *Topbp1^{+/+}*.

A) Representative images and quantification of $Topbp1^{+/+}$ and $Topbp1^{B5/B5}$ mid-pachytene spermatocytes stained with RNApol2 and SYCP3, B) pRNApol2_Ser2 and SYCP3, and C) pRNApol2_Ser5 and SYCP3. n = number of mice.

Supplemental Figure 18. Genes used as markers to define the stages of the germ cells in the scRNAseq analysis.
UMAP plots showing markers for Spermatogonia (*Sal1* and *Dmrt1*), Spermatocytes (*Dazl*, *Id4*, *Sycp3*, and *Shcbp1l*), and Spermatids (*Acrv1*, *Oaz3* and *Prm2*).

Supplemental Figure 19. Genes used as markers to track spermatogenesis progression and form the 47 sub-cluster for all cells captured in the 10X platform in the scRNAseq analysis.
Bubble plot displaying gene markers and their respective pattern of expression during spermatogenesis.

Supplemental Figure 20. Genes used as markers of somatic cells captured in the scRNAseq analysis.
UMAP plots show markers for the somatic cells captured on the germ cell-enriched population, Endothelial and Hematopoietic (*Col1a2*, *Vcam1*, *Laptm5* and *Insl3*), Smooth muscle (*Acta2*) and Sertoli (*Ptgds* and *Wt1*).

Supplemental Figure 21. Analysis of sub-clusters of all cells captured in the 10X platform.
UMAP plot displaying the 47 sub-clusters of cells captured in the scRNAseq analysis.

Supplemental Figure 22. Analysis of the main 24 sub-clusters captured in the germ cells-enriched scRNAseq analysis.
UMAP plot showing 24 sub-clusters for spermatogonia, spermatocytes, round spermatids, spermatids, Sertoli, smooth muscle and endothelial and hematopoietic cells captured in the 10X platform.

Supplemental Figure 23. Spermatocytes from $Topbp1^{+/+}$ and $Topbp1^{B5/B5}$ show correlation greater than 0.9.
Plot of Pearson correlation values between cell groups separated by cluster and genotype using log-normalized average expression of all genes.

Supplemental Figure 24. $Topbp1^{B5/B5}$ pachytene spermatocytes show increased expression of X-linked genes.

1317   Heatmap displaying the expression levels of X-chromosome genes in spermatocyte 3 stage (pachytene) of

1318   *Topbp1$^{B5/B5}$* on the right and *Topbp1$^{+/+}$* on the left. Each line represents a gene, and each row is a cell.

1319

Supplemental Figure 25. *Topbp1$^{B5/B5}$* pachytene spermatocytes show increased expression of XY

"killer genes" and other X and Y genes typically used to illustrate MSCI defects.

Violin plots showing the level of expression and number of cells expressing the XY chromosome genes in

SP3 (pachytene): *Zfy1*, *Zfy2*, *Kdm6a*, *Lamp2*, *Scml2*, *Zfx*, *Uba1y* and *Rhox13* in *Topbp1$^{B5/B5}$* and in

Topbp1$^{+/+}$. The Wilcoxon test was used to calculate the p-values. **$p < 0.01$, ***$p < 0.001$, and **** $p <$

0.0001.

1326

1327   **Supplemental Figure 26. *Topbp1$^{B5/B5}$* pachytene spermatocytes show increased expression of XY-**

1328   **linked genes.**

1329   Bar plots showing the number of cells expressing the XY chromosome genes in SP3 (pachytene): *Zfy1*,

1330   *Zfy2*, *Kdm6a*, *Lamp2*, *Scml2*, *Zfx*, *Uba1y* and *Rhox13* in *Topbp1$^{B5/B5}$* and in *Topbp1$^{+/+}$*. Fisher's exact test

1331   was used to calculate the p-values. **$p < 0.01$, ***$p < 0.001$, and **** $p < 0.0001$.

1332

1333   **Supplemental Figure 27. Analysis of the gene marker used to define the zygotene stage.**

1334   UMAP analysis of the gene *Gm960* was used as a gene marker to define the zygotene stage.

1335

1336   **Supplemental Figure 28. Expression levels of genes with no defects, mild defects, and strong defects**

1337   **in *Topbp1$^{B5/B5}$* and *Topbp1$^{+/+}$*.**

1338   Boxplot showing the expression levels of the clusters of genes with no defects, mild defects, and strong

1339   defects from pre-leptotene to Spermatocyte 3 (pachytene) in *Topbp1$^{B5/B5}$* and *Topbp1$^{+/+}$*. The Wilcoxon

1340   rank-sum test was used to calculate the p-values. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and **** $p < 0.0001$.

1341

1342   **Supplemental Figure 29. Quality control of the scRNAseq data.**

1343   Violin plots show the quality control for the scRNAseq data, displaying the distribution of gene counts,

1344   UMI counts, and mitochondrial gene percentage per library before filtering data.

1345

Figure 1

A

TOPBP1

BRCT0 | BRCT1 | BRCT2 | BRCT3 | BRCT4 | BRCT5 | BRCT6 | ATR Activation Domain | BRCT7 | BRCT8

VRKANAKKGMLASTHLIVKEPTGSKYEAAKKWSLPAVNISWLLETARIGKR   *Topbp1*^(+/+)
VDEANAEEGMLASTHLIVEEPTGSEYEAAKKWSLPAVNISWLLETARIGEE   *Topbp1*^(B5/B5)

B Body mass (g) — ns — *Topbp1:* +/+ B5/B5

C +/+ B5/B5 — *Topbp1*

D IR 7Gy Δweight (%) vs Days — *Topbp1*^(+/+), *Topbp1*^(B5/B5)

E
♀ *Topbp1*^(+/+) X ♂ *Topbp1*^(B5/B5) Breeding cages = 7 → Litters = 0
♀ *Topbp1*^(B5/B5) X ♂ *Topbp1*^(+/+) Breeding cages = 7 → Litters = 10

F *Topbp1:* +/+ B5/B5

G Testes mass/ Body mass — **** — *Topbp1:* +/+ B5/B5

H Sperm count — **** — *Topbp1:* +/+ B5/B5

I *Topbp1*^(+/+) 20X 40X; *Topbp1*^(B5/B5) 20X 40X

J *Topbp1*^(+/+) 20X; *Topbp1*^(B5/B5) 20X

K Apoptotic cells/Tubule — **** — *Topbp1:* +/+ B5/B5

L *Topbp1*^(+/+) *Topbp1*^(B5/B5) — SYCP3 SYCP1 H1T
Leptonema, Zygonema, Early Pachynema, Mid-Pachynema, Diplonema (Not detected)

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8