Variation in thermal sensitivity of diapause development among individuals and over time

drives life history timing patterns in an insect pest

Short title: Thermal sensitivity of insect diapause

Jantina Toxopeus[1,*], Edwina J. Dowle[2], Lalitya Andaloori, Gregory J. Ragland

Department of Integrative Biology, University of Colorado, Denver, 1151 Arapahoe St, Denver

CO, 80204, United States of America

[1]Present address: Department of Biology, St. Francis Xavier University, 2321 Notre Dame Ave,

Antigonish NS, B2G 2W5 Canada

[2]Present address: Department of Anatomy, University of Otago, Dunedin, New Zealand

*Corresponding author

Author emails: jtoxopeu@stfx.ca (JT), eddy.dowle@otago.ac.nz (EJD),

lalitya.andaloori@cuanschutz.edu (LA), gregory.ragland@ucdenver.edu (GJR)

Abstract

Physiological time is important for understanding the development and seasonal timing

of ectothermic animals, but has largely been applied to developmental processes that occur

during spring and summer such as morphogenesis. There is a substantial knowledge gap in the

relationship between temperature and development during winter, a season that is increasingly

impacted by climate change. Most temperate insects overwinter in diapause, a developmental

process with little obvious morphological change. We used principles from the physiological

time literature to measure and model the thermal sensitivity of diapause development rate in the

apple maggot fly *Rhagoletis pomonella*, a univoltine fly whose diapause duration varies

substantially within and among populations. We showed that diapause duration could be

modelled with simple linear relationships between diapause development rate and temperature.

Low temperatures were necessary for most individuals to complete diapause, similar to many

insects. However, we also found evidence for an ontogenetic shift in the thermal sensitivity of

diapause: diapause development proceeded more quickly at high temperatures later in diapause

and in the absence of quiescence, a phenomenon not previously reported. Increasingly warmer

temperatures during and after winter may impact the phenology of this and other insects with

temperature-sensitive diapause, impacting their viability.

Introduction

Physiological time is an important concept for understanding the development and seasonal timing of ectothermic animals (Taylor 1981; Buckley 2022). The cellular and physiological processes that underlie morphogenesis (development of an animal's body size, shape and structure) are temperature-dependent (Kipyatkov and Lopatina 2010; Shi et al. 2011; Damos and Savopoulou-Soultani 2012). The vast majority of animals on our planet are ectothermic, so the chronological time (i.e., number of hours/days) required to reach a developmental landmark depends on environmental temperature. Most natural environments exhibit substantial temperature variation; chronological time is thus a poor metric of life history timing (van Straalen 1983; Trudgill et al. 2005; Rebaudo and Rabhi 2018). Rather, ectothermic development is usually described in physiological time, conceptualized by the relationship between development rate (governed by physiological processes) and environmental temperature (Taylor 1981; van Straalen 1983). Environmental temperature variability, duration of temperature exposure, and ontogeny can complicate the calculation of physiological time (Kingsolver and Woods 2016; Sinclair et al. 2016; Buckley 2022), but physiological time models have been well-characterized in insects and can facilitate predictions of population dynamics during the growing season – i.e., during spring, summer, and fall (Powell and Bentz 2009; Régnière et al. 2012; Sridhar and Reddy 2013; Scranton and Amarasekare 2017). However, we have comparatively little insight into whether and how the same concept of physiological time applies to developmental processes that occur during winter. The relative dearth of information on the relationship between temperature and development during winter represents a substantial knowledge gap, especially if we want to understand the consequences of climate warming on insect phenology (Marshall et al. 2020; Buckley 2022).

Most temperate insects overwinter in diapause (Fig. 1 – green and yellow), a dormant,

stress-tolerant state that delays growth, morphogenesis, and reproduction until conditions

become permissive in spring or summer (Tauber and Tauber 1976; Hand et al. 2016; Wilsterman

et al. 2021). Many insects spend the bulk of their life in diapause, either the majority of a

calendar year (Fig. 1A) or sometimes multiple years (Tauber et al. 1986; Hanski 1988; Hahn and

Denlinger 2011; Moraiti et al. 2014; Dupuis et al. 2016). If diapause ends during winter, the

insect often enters another form of dormancy called 'quiescence' – a lack of morphogenesis due

to non-permissive environmental conditions (Fig. 1B – dark blue; Jenkins et al. 2001; Hodek

2002; Koštál 2006). Unlike diapause, where dormancy is maintained during transient or early

exposure to permissive environments, quiescent individuals are not recalcitrant and will

immediately resume post-diapause processes such as growth and morphogenesis when

conditions become permissive. Diapause and post-diapause quiescence can synchronize

individuals to have similar life history timing, facilitating mate finding and supporting

reproductive output (Tauber and Tauber 1976; Hand et al. 2016; Wilsterman et al. 2021). Given

the prevalence of diapause in overwintering insects, any attempts to apply physiological time to

overwintering development must consider this important process.

To expand the concept of physiological time to diapause, we must first establish that

diapause is indeed a form of development. In contrast to growth and morphogenesis (Fig. 1 -

gray) during warmer or more permissive periods of the year, there is often no obvious

morphological change (morphogenesis) during diapause (but see Shingleton et al. 2003), leading

some authors to prefer alternatives to the term 'diapause development' (Hodek 1996). However,

diapause is clearly a dynamic developmental process, where 'development' may primarily

consist of progressive or cyclic changes in the molecular cellular milieu (Andrewartha 1952;

Hodek 1996; Koštál 2006), including differential expression of often thousands of genes over

time (Koštál et al. 2017; Dowle et al. 2020; Pruisscher et al. 2022). Indeed, diapause can have

distinct phases (e.g., initiation, maintenance, termination; cf. Koštál 2006), each of which must

complete before post-diapause processes such as morphogenesis can resume.

Figure 1. Example durations of diapause for insects that typically enter diapause in late
summer or early autumn. (A) Some insects remain in diapause under warm (yellow) and cool
(green) conditions for most of the year. **(B)** Many insects complete diapause during winter when
it is cool (green), and enter a post-diapause dormancy called quiescence (blue), resuming post-
diapause development (morphogenesis; gray) immediately when temperatures become
permissive in the spring or early summer. **(C)** Warm summer or autumn conditions can cause
individuals to complete diapause rapidly, referred to as 'weak diapause' or 'shallow diapause' in
some species **(D)** Non-diapause individuals never enter diapause and may complete
morphogenesis prior to winter.

Temperature has some well-described effects on diapause, but few studies have

systematically investigated thermal sensitivity of diapause development. Chilling at low

temperatures appears to be required for completion of the diapause program in many temperate

species (Denlinger 2002), although there are many exceptions to this 'rule' (Hodek 2002; Zhu et

al. 2009; Chen et al. 2014). In addition, warm temperatures can cause some insects to complete

diapause relatively quickly (Fig. 1C) or never enter diapause (Fig. 1D; Denlinger 2002; Koštál

2006; Dambroski and Feder 2007; Toxopeus et al. 2021; Calvert et al. 2022). A common

experimental approach examines the impact of different chilling or winter durations to find the

chilling threshold time required for diapause termination (e.g., Feder et al. 1997; Sgolastra et al.

2010; Higaki and Toyama 2012; Moraiti et al. 2014; Lehmann et al. 2017). However,

experiments that manipulate only chilling duration do not provide much insight into the

underlying thermal sensitivity of the diapause development process, and the potential impact of

warming autumn and winter conditions.

Models of physiological time that measure morphogenesis during warm conditions are

well-described, and can inform our approach to modelling diapause development. Nonlinear,

hump-shaped functions relating environmental temperature to morphogenic development rate

generally fit empirical data quite well (Kipyatkov and Lopatina 2010; Shi et al. 2011; Damos and

Savopoulou-Soultani 2012; Kingsolver and Woods 2016; Sinclair et al. 2016). These models, a

class of thermal performance curves, predict zero development below or above the low and high

temperature limits, respectively. Between these lower and upper thermal limits, as temperatures

increases morphogenic development rate increases to its highest value at the thermal maximum

(c. 20 – 40 °C for many insects; Taylor 1981), and then sharply decreases back to a rate of zero

as temperatures become too warm. The time to complete a developmental process at a particular

temperature is the inverse of the development rate at that temperature (Kipyatkov and Lopatina

2010; Damos and Savopoulou-Soultani 2012), so development duration is shortest at the thermal

maximum. Simpler slope plus intercept (straight line) models may also reasonably approximate

the relationship between morphogenic development rate and temperature between the lower

thermal limit and the thermal maximum (Trudgill et al. 2005; Kipyatkov and Lopatina 2010).

Ontogenetic variation in thermal sensitivity of development is another important source of

variation in development times/rates that is well described and often incorporated into predictive

models (Yurk and Powell 2010; Lopatina et al. 2014; Sinclair et al. 2016; Kutcherov 2020).

Finally, genetic variation within populations may also explain a substantial portion of remaining,

interindividual variation in development time that is often observed even under highly controlled

laboratory conditions (Kingsolver et al. 2004)

We propose that diapause duration can be modelled such that physiological time serves

as the diapause timer (cf. Hodek 2002). To do this, we can use a thermal performance curve of

diapause development rate. Given the well-known chilling relationships described above, these

diapause curves likely have very different shapes compared to functions describing morphogenic

(non-diapause) development with maximum rates at high temperatures (cf. Hilbert et al. 1985).

To our knowledge, there are no existing empirical estimates of thermal performance curves for

diapause development, so we start with simple, hypothetical models that cover the most plausible

forms of the relationship (Fig. 2 insets). Diapause phenology experiments often involve

incubating insects at two different, consecutive temperatures (chilling then warming; e.g., Rull et

al. 2016; Lindestad et al. 2020; Toxopeus et al. 2021), so we present the predicted diapause and

total development times associated with these simple thermal sensitivity experiments (Fig. 2), as

expanded upon below.

152
Figure 2. Hypothesized effect of a single (A) chronological or (B – D) physiological diapause
timer on development duration under different combinations of chilling and warming.
Transition from cold to warm temperature indicated by dashed line. Development duration
includes diapause development under cold (green) and warm (yellow) conditions, and post-
diapause (e.g., morphogenic) development (gray) that can only occur under warm conditions. If
individuals remain chilled after diapause completes (arrows), cold-induced post-diapause
quiescence (dark blue) occurs until animals are returned to warm temperatures. Insets:
relationship between temperature (T) and diapause development rate (r) at a low (chilling) and
high (warming) temperatures based on an example thermal sensitivity (TS) of the diapause timer.
The equations that explain the relationships between development rates, chill duration, and
development time are fully explained in Appendix A. Multiple diapause timers may exist (e.g.,
for different stages of diapause), which would result in more complex relationships between
chilling duration and development duration (not shown).
166
167
Our models make several simplifying assumptions about the environmental sensitivity of

diapause and about how an observer measures diapause duration. We do not make any

assumptions about how diapause is induced, but do assume that temperature alone (and not

photoperiod or other cues) influences the duration of diapause. This assumption applies

reasonably well to many insects that complete diapause without additional environmental cues

regardless of whether diapause was induced by a non-temperature cue (Tauber and Tauber 1976;

Hodek 1996; Denlinger 2022). Directly observing the precise timing of the end of diapause is

often not practical. This is because overt, morphological changes (morphogenesis) associated

with resumption of post-diapause growth and development often occur well after hormonal and

metabolic pulses that delineate the true physio-developmental transition (Hodek 1996, 2002;

Denlinger 2002). Thus, we further assume that a measurement proxy for diapause duration also

includes some amount of post-diapause development (morphogenesis; Fig. 2 - gray), culminating

in a clear phenotypic marker such as a life stage transition (e.g., adult emergence of a pupal

diapausing insect) or a behavior (e.g., emergence from hibernacula). Finally, we assume that

though diapause may progress at relatively cold temperatures, post-diapause development only

proceeds at relatively warm temperatures. If diapause ends during low temperature exposure,

quiescence occurs (Fig. 2 – dark blue), which delays post-diapause development because low

temperatures impede morphogenesis (Jenkins et al. 2001; Hodek 2002; Koštál 2006).

Consider a naïve model in which diapause development progresses at a constant rate over

time, independent of temperature. Under this chronological timer, diapause duration is

approximately the same regardless of thermal environment (Fig. 2A) – some proportion of

diapause development (and post-diapause development) occurs during chilling and the rest of

diapause development occurs during warming. The only treatment that can increase overall

development duration is chilling the insects for longer than their diapause duration (e.g., Fig. 2A;

longest chill duration), which delays resumption of post-diapause development while insects are

in chilling-induced quiescence. More nuanced chronological diapause timers are possible, e.g.,

chronological timers that are "started" by chilling or warming, but we focus here on the simplest

model as a point of comparison to models that incorporate physiological time (Fig. 2B-D).

Realistically, we expect temperature to influence diapause development in some way.

First, consider the classic "chill-dependent" diapause (or chilling threshold) model (cf.

Dambroski and Feder 2007), where diapause can only progress at low temperatures. In this case,

we predict that the physiological diapause timer has a fairly narrow thermal sensitivity, such that

the rate of diapause development (r) is high at low temperatures, but zero at high temperatures

(Fig. 2B inset). Individuals must be chilled for a sufficient time (chilling threshold) to complete

diapause, and those that experience no or short chilling durations never complete 100% of the

diapause program, and therefore cannot proceed to post-diapause development when warmed

(Fig. 2B). Alternatively, the physiological diapause timer might have a broad thermal sensitivity,

with a thermal maximum for diapause development rate at low (Fig. 2C inset) or high (Fig. 2D

inset) temperatures. If low temperatures accelerate the diapause timer, diapause duration and

chilling duration have an inverse relationship: long chilling durations result in short development

times (Fig. 2C). Under this diapause timer, the longer the individual is in a cold environment, the

greater the proportion of diapause completed prior to warming, so less time is needed at warm

temperatures to complete diapause. If high temperatures accelerate the diapause timer, we expect

a positive correlation between chill duration and development time, with the shortest diapause

duration in individuals that are only exposed to warm temperatures due to high rates of diapause

development at these temperatures (Fig. 2D).

Just as thermal sensitivity may vary among individuals and over time during

morphogenic development (Yurk and Powell 2010; Lopatina et al. 2014; Sinclair et al. 2016;

Kutcherov 2020), diapause development performance curves may differ among individuals and

over time. For example, early diapause (e.g., initiation, maintenance) may have a different

thermal sensitivity than late diapause (e.g., termination), which we have not captured in our

simple models (Fig. 2). However, each thermal sensitivity model in Figure 2 can be applied to a

distinct stage of diapause rather than diapause as a whole, if needed. To avoid confusion between

development rate and changes in development rate across diapause development, we refer to the

latter as changes or variation across ontogeny.

In this study we fit these naïve and physiological time models (Fig. 2) to measurements

of diapause duration across simple cold/warm temperature treatments in the univoltine apple

maggot fly *Rhagoletis pomonella* (Diptera; Tephritidae). *Rhagoletis pomonella* diapause as

pupae under the soil for most of the year (Fig. 1A), and this diapause duration is not highly

photosensitive (Prokopy 1968; Feder et al. 1997), making *R. pomonella* a good model to

examine the impact of temperature on diapause development. In addition, intra/inter-population

variation in diapause related to seasonality has been intensively studied in *R. pomonella*

(Dambroski and Feder 2007; Lyons-Sobaski and Berlocher 2009; Powell et al. 2020; Calvert et

al. 2022). Our study demonstrates that diapause duration can be modelled using the principles of

physiological time, with implications for the effect of changing climates on the life history

timing of many insect species. Some elements of the models may be somewhat idiosyncratic to

R. pomonella, but we argue that accounting for ontogenetic and interindividual variation will

likely be similarly critical in other species.

Materials and Methods

Rhagoletis pomonella as a diapause model

     Rhagoletis pomonella has recently evolved to be an agricultural pest, laying its eggs in fruits of apple trees (*Malus domesticus*) that were introduced to North America several hundred years ago. It natively infests hawthorn (*Crataegus* spp.) trees throughout North America, and both apple- and haw-infesting populations mate and oviposit in the late summer/early autumn (Dean and Chapman 1973). Larvae develop within fruit, burrow into the soil to pupate, and then most pupae enter a long diapause and spend up to 10 months in the soil, eclosing as adults the following summer (e.g., Fig. 1A; Dean and Chapman 1973). Differences in life history timing (e.g., between apple- and hawthorn-infesting *R. pomonella*; (Bush 1969; Feder et al. 1993) are driven by differences in diapause duration; the duration of post-diapause morphogenesis leading to adult eclosion in *R. pomonella* varies little among individuals at a given temperature (Powell et al. 2020). The lower temperature limit for this pupal morphogenesis is c. 6°C, and above this the rate of morphogenic development increases approximately linearly with temperature up to 26°C (Reissig et al. 1979). If *R. pomonella* pupae complete diapause at low temperatures ($< 6°C$) they will enter post-diapause quiescence until temperatures become permissive ($> 6°C$) for post-diapause development (morphogenesis).

     There are three well-described, discrete diapause phenotypic classes in *R. pomonella* that vary in diapause intensity, or recalcitrance to develop despite permissive temperatures (Dambroski and Feder 2007; Calvert et al. 2022). In the absence of low temperatures, a small proportion ($<10\,\%$) of *R. pomonella* pupae avert diapause completely (non-diapause phenotype; ND), developing directly into adults without ever entering dormancy (e.g., Fig. 1D; Prokopy

1968; Dambroski and Feder 2007; Calvert et al. 2022). Of the remaining pupae that enter

diapause (as determined by a dramatic suppression of metabolic rate; cf. Ragland et al. 2009),

most exhibit the prolonged chill-dependent diapause (CD) phenotype and remain in diapause for

an extended period of several months (e.g., Fig. 1A; Feder et al. 1997; Dambroski and Feder

2007; Calvert et al. 2022). A small portion (< 20 %) of pupae exhibit the weak diapause

phenotype (WD; cf. Toxopeus et al. 2021), also termed shallow diapause (Dambroski and Feder

2007; Calvert et al. 2022); they enter diapause, but will terminate diapause (increase metabolic

rate) after only a few weeks and develop into adults if held at warm temperatures (e.g., Fig. 1C).

These three diapause intensity phenotypes are relatively discrete, with multimodal distributions

of eclosion and even distinct allele frequencies (Feder et al. 1997; Dambroski and Feder 2007;

Calvert et al. 2022).

Intra- and inter-population variation in diapause development is pervasive and well-

characterized in nature, including variation in diapause intensity (Tauber and Tauber 1972;

Dingle et al. 1990; Masaki 2002; Chen et al. 2013; Geisert and Meinke 2013; Papanastasiou and

Papadopoulos 2014). Thus, we argue that the relative complexity that we observe *R. pomonella*

diapause is important for understanding the fairly general diversity in diapause phenotypes

across temperate insects. In *R. pomonella* we also have the advantage of a large body of literature

on diapause variation that allows us to realistically parameterize our models.

Insect collection and diapause incidence

We collected hawthorn fruits (*Crataegus* spp.) infested with *R. pomonella* in Denver CO

during August 2017 and 2018, brought them to the lab, and collected emerging larvae and pupae

as previously described (Toxopeus et al. 2021). Briefly, we placed fruits in wire mesh baskets

suspended over plastic trays in an environmentally-controlled room (c. 22°C). We collected

newly formed pupae three times per week, and transferred them to a Percival DR-36VL

incubator (Percival Scientific, Perry IA) set to 21°C, 14:10 L:D and 80% R.H. for 10 days.

Diapause *R. pomonella* pupae suppress their metabolic rate during these 10 days, after which

metabolic rate stabilizes (Ragland et al. 2009). We defined 10 d post-pupariation as "time zero"

in our experiments.

To estimate the proportion of *R. pomonella* pupae in each diapause phenotypic class (ND,

WD, CD), we measured CO_2 production in a subset of 2018 pupae at 10 d and 45 d post-

pupariation at 21°C using stop-flow respirometry as previously described (Ragland et al. 2009;

Toxopeus et al. 2021). A small number (6 out of 200) with intermediate metabolic rates were

excluded because they could not be unambiguously assigned to a diapause phenotype. Because

we wanted to focus our analysis on diapause development, we excluded ND pupae from our

subsequent analyses, either based on metabolic rate, or very short (< 40 days; cf. Feder et al.

1997; Toxopeus et al. 2021) times to complete development to adulthood after chilling

treatments (see below).

Temperature treatments and development tracking

<u>Experiment 1: Different durations of chilling</u>

To determine whether diapause was governed by physiological timers, and whether

diapause development was accelerate at low or high temperatures, we compared the relationship

between chilling duration and diapause duration in *R. pomonella* to the predictions in Figure 2.

At time zero (10 d post-pupariation), pupae were separated into groups of c. 100 individuals. One

group was kept at 21°C for the duration of the experiment (no chilling). All other groups were

transferred directly to simulated winter (chilling at 4 °C and c. 85% R.H) for one of several

chilling durations ranging from 2 to 29 weeks. Following chilling, pupae were transferred back

to 21°C. Each experimental group was checked three times per week at 21°C to determine when

adults eclosed from the puparia, the marker that diapause and post-diapause development

(morphogenesis) were complete. For each treatment group, we calculated the mean

developmental duration (including time at 4°C and 21°C), the coefficient of variation (CoV) of

mean developmental duration, and the proportion of pupae that eclosed as adults. Summaries of

development duration only included individuals that eclosed as adults.

Mean diapause duration and the incidence of quiescence in *R. pomonella* was determined

in each temperature treatment group based on two assumptions. 1) At 4°C, pupal morphogenesis

(i.e. post-diapause development) cannot occur (Reissig et al. 1979), but diapause development

can occur, based on the observation that pupae can complete development following prolonged

chilling (Feder et al. 1997; Powell et al. 2020; Toxopeus et al. 2021). 2) At 21°C, time to

complete post-diapause pupal morphogenesis is relatively invariant among individuals, so the

time to complete development is largely driven by the time it takes to complete diapause (Feder

et al. 1997; Powell et al. 2020). If post-chill developmental duration is approximately equal to

the time to complete morphogenesis (c. 30 days), we can infer that the diapause process

completed at 4°C and post-diapause quiescence may have occurred (e.g., longest chill duration

bars in Fig. 2). However, if the post-chill developmental duration is greater than that required to

complete morphogenesis, we can infer that only a portion of the diapause process was completed

at 4°C and the remaining proportion of diapause development occurred at 21°C prior to post-

diapause morphogenesis (e.g., all chill durations except the longest in Fig. 2C, 2D). In this latter

case, cold-induced quiescence does not occur, so diapause duration (e.g., green + yellow bars in

Fig. 2) is the difference between age at eclosion (e.g., total height of bars in Fig. 2) and the

duration of morphogenesis (e.g., gray bars in Fig. 2).

<u>Experiment 2: Chilling for the same duration but at different ages</u>

339     To detect ontogenetic shifts in the thermal sensitivity of diapause development, we

compared developmental duration of *R. pomonella* pupae exposed to the same chilling durations

but at different chronological ages. We included additional temperature treatments in which

groups of c. 100 pupae were kept at 21°C for a warming period (2 to 6 weeks beyond time zero)

prior to chilling (4°C) for 2, 3, 6, or 9 weeks. Pupae were then transferred back to 21°C and

eclosion was tracked as described above. We compared developmental duration in these

treatments to pupae that were chilled at time zero (10 d post-pupariation) for 2, 3, 6, or 9 weeks.

If chilling at different chronological ages resulted in a decrease or increase in total development

time, this would suggest that the thermal sensitivity of the diapause timer can vary with

ontogeny, i.e., it can change over the course of diapause development.

Diapause timer computational simulations

351     Initial inspection of our empirical data suggested the results could not be explained by

one of the simple relationships between diapause development rate and temperature (Fig. 2; see

Results for additional details). Indeed, due to variation in the proportion of individuals that

eclosed (completed diapause and post-diapause development) following each temperature

treatment in Experiment 1 and the presence of WD and CD diapause phenotypes, it was clear

that we needed to account for interindividual and ontogenetic variation in thermal sensitivity of

diapause development. To do this, we developed a simulation model that incorporated these

factors. The full derivation of the model equations and an explanation of model assumptions

(summarized in Table A1) are available in Appendix A, and briefly described below. All

simulations and analyses were conducted in R v4.1.0 (R Core Team 2022), and our code is

available at https://anonymous.4open.science/r/diapause-thermal-sensitivity-0DDC/.

Based on estimated proportions of diapause intensity phenotypes (see Results), we

simulated a population that included 150 WD and 850 CD individuals, each with its own

diapause timer(s) that specified diapause development rate at 4°C and 21°C. We modeled one

diapause timer for each WD individual, and two diapause timers (early and late diapause

development) for each CD individual, accounting for ontogenetic variation in diapause

development in CD individuals. Within each diapause phenotype (WD, CD), diapause

development rates varied among simulated individuals. In addition, we incorporated some

chilling-related mortality for both diapause phenotypes based on empirical observations. Total

development duration (diapause and post-diapause development) was calculated from

development rates (Eq'ns 1.1 – 2.4) for each simulated individual that was 'exposed' to 4°C

(chilling) at time zero for between 0 and 400 days, followed by warming at 21°C, similar to

Experiment 1 above.

Total development time (diapause and post-diapause development) of each WD

individual as a function of time at 4°C was simulated with the following equations that included

a single diapause timer:

$Y = \frac{r-R}{r}X + \frac{1}{r} + \frac{1}{r_g}$ when $0 \leq X < \frac{1}{R}$ (Eq'n 1.1)

$Y = X + \frac{1}{r_g}$ when $\frac{1}{R} \leq X < X_z$ (Eq'n 1.2)

$Y = 0$ when $X \geq X_z$ (Eq'n 1.3)

where Y is the time to complete development since time zero, X is the duration of

chilling (d), R is the rate of diapause development at 4°C (d^{-1}; e.g., green circle in Fig. 2 insets),

r is the rate of diapause development at 21°C (d^{-1}; e.g., yellow triangle in Fig. 2 insets), r_g is the

rate of post-diapause development (morphogenesis) at 21°C (d^{-1}), and X_z is the duration of

chilling (d) that causes mortality in WD individuals. The inverse of each rate is the time to

complete that process (diapause or morphogenesis) at the respective temperature. When $Y = 0$

(e.g., Eq'n 1.3), that indicates that an individual was unable to complete development and adult

eclosion never occurred, so development time is effectively 0 d. We use $Y = 0$ rather than

recording simulated individuals as dead because (as described below for CD individuals) some

individuals fail to complete development because they are alive but temperatures are not

permissive for development (e.g., no chilling and shortest chilling treatment in Fig. 2B).

To model total development time (diapause and post-diapause development) of each CD

individual, we included two diapause timers – one each for early (*e*) and late (*l*) diapause

development – in the following equations:

$Y = 0$ when $X < \frac{1}{R_e}$ (Eq'n 2.1)

$$Y = \frac{r_l - R_l}{r_l}X + \frac{R_l + R_e}{R_e r_l} + \frac{1}{r_g} \text{ when } \frac{1}{R_e} \leq X < \frac{1}{R_e} + \frac{1}{R_l} \quad \text{(Eq'n 2.2)}$$

$$Y = X + \frac{1}{r_g} \text{ when } \frac{1}{R_e} + \frac{1}{R_l} \leq X < X_z \quad \text{(Eq'n 2.3)}$$

$$Y = 0 \text{ when } X \geq X_z \quad \text{(Eq'n 2.4)}$$

where R_e is the rate of early diapause development at 4°C (d^{-1}), R_l is the rate of late

diapause development at 4°C (d^{-1}), r_l is the rate of late diapause development at 21°C (d^{-1}), X_z is

the duration of chilling (d) that causes mortality in CD individuals, and all other symbols are as

described for the WD individuals. Early diapause must complete before late diapause can begin.

We assumed that early diapause development did not proceed at warm temperatures (21°C), i.e.

chilling was required for completion of early diapause (cf. Fig. 2B), and so individuals did not

complete development and eclose ($Y = 0$) if chilling duration was insufficient (less than $1/R_e$) for

individuals to complete early diapause (Eq'n 2.1).

Our simulations used the equations above and accounted for interindividual variation in

diapause development rates among simulated individuals in the following way. The rates of

diapause development for each of the simulated WD individuals (R and r in Eq'ns 1.1 – 1.2) and

CD individuals (r_l, R_e, R_l in Eq'ns 2.1 – 2.3) were randomly sampled from inverse Gaussian

distributions of rates with the parameters described in Table A2. We then calculated the same

summary metrics (mean and CoV development duration, proportion eclosion) for our simulated

population as for our empirical population for each chill duration used in Experiment 1,

including only eclosed ($Y > 0$) individuals from summaries of development duration (as we did

for our empirical population). We compared the fit of our simulation summary metrics to the

empirical summary metrics by calculating the sum of squared residuals (SSR) of mean

development duration and the χ^2 of proportion eclosion between the two datasets. We used

multiple iterations of these simulations to numerically solve for the parameters of our rate

distributions (μ, λ in Table A2) that minimized the SSR and χ^2 values.

While the assumptions in our final simulation model were justified by empirical data (see

Appendix A), we also tested the sensitivity of the model to these assumptions. To do this, we

either modified or removed parameters from Eq'ns 1.1 – 2.4, ran our simulation, calculated the

SSR and χ^2 parameters as described above, and visually compared the fit of the simulation

summary metrics to the empirical summary metrics. The alternative simulations we ran included:

1) only one diapause phenotype (Eq'ns 2.1 – 2.4) instead of distinct WD and CD phenotypes;

2) both WD and CD phenotypes, but no ontogenetic variation in diapause development rates in

CD individuals; 3) mortality due to chilling in only one of or neither WD and CD phenotypes.

All alternative models are summarized in Supplementary Information and a subset of the most

relevant alternative simulations are included in the main text.

Results

The results detailed below broadly support the hypothesis that insect development during

winter-like conditions can be explained by timers reflecting thermally sensitive, physiological

time. The thermal sensitivity of these timers differ between diapause intensity phenotypes,

among individuals, and across ontogeny. While there are multiple potential interpretations of the

empirical data, below we focus on interpretations that were supported by our subsequent

simulation models.

Diapause timers vary among diapause intensity phenotypes

Based on the proportion of individuals that completed development under different

chilling and warming treatments in Experiment 1, our data suggested that weak diapause (WD)

and chill-dependent diapause (CD) phenotypes have different diapause timers. Consistent with a

number of previous observations (Dambroski and Feder 2007; Toxopeus et al. 2021; Calvert et

al. 2022), we confirmed the presence of ND (non-diapause), WD, and CD *R. pomonella* pupae

reared under constant warm conditions using metabolic rate measurements (Fig. 3A). We note

that these diapause phenotypes are relatively distinct rather representing a continuous

distribution of diapause intensities. Under constant warmth, all ND flies completed their eclosion

before WD flies began eclosing, and most CD flies remained in diapause under these conditions

(did not eclose at all; Fig. S1, S2). CD pupae were most abundant (82%), followed by WD

(15%), and ND (3%; Fig. 3B), the latter of which was excluded from our analyses (see Methods

for details).

456
Figure 3. Diapause incidence and the effect of chilling on diapause completion in *Rhagoletis pomonella*. **(A)** Mass-specific metabolic rate after 10 or 45 days of constant warmth post-pupariation was used to determine diapause phenotype: non-diapause (ND), weak diapause (WD), and chill-dependent diapause (CD). Each open circle represents one individual. Filled black circles represent the mean of each group. Error bars represent 95% confidence intervals; small error bars are obscured by symbols. **(B)** The proportion of ND, WD, and CD pupae inferred from the metabolic rate data in **(A)** and duration of total development (Fig. S2) in groups that were not metabolically-phenotyped. **(C)** Each point represents the proportion of c. 100 pupae that completed diapause and post-diapause development (eclosed as adults) following exposure to chilling (4°C) for a specific duration at time zero (10 d post-pupariation). Error bars represent the standard error of proportion. Black line is a cumulative distribution fit with a mean ± standard deviation of 60 ± 15 d chilling.

The two types of diapause timers in this population included a rarer one that did not

require chilling, and a common one that required moderate to prolonged chilling. As previously

documented (Feder et al. 1997; Dambroski and Feder 2007; Toxopeus et al. 2021), some

diapause *R. pomonella* pupae were able to complete diapause and eclose as adults following a

range of temperature treatments, from no chilling (warming only) to prolonged chilling (>200

days) at 4°C (Fig. 3C). Approximately 15% of our population eclosed with little to no chilling (0

– 4 weeks at 4°C; Fig. 3C), which is similar to the proportion of WD pupae in our population

(Fig. 3B). We therefore inferred that the WD pupae had a diapause timer that did not require

chilling to complete the diapause program. However, as chill duration increased, the proportion

eclosion increased in a sigmoidal pattern to a maximum of 91% (Fig. 3C), suggesting that the

remaining CD pupae had a diapause timer that required a threshold cold exposure (chilling) to

successfully complete the diapause program. We confirmed that the majority of uneclosed pupae

following short chill treatments appeared to be alive and undeveloped (Fig. S1), suggesting that

they were unable to complete diapause, rather than dead, or developed but unable to leave the

puparium.

Variation in diapause timers among individuals

The relationship between chill duration and proportion eclosion in Experiment 1 (Fig.

3C) also suggested that there was substantial interindividual variation in diapause timers among

individuals within each diapause phenotype. This is best illustrated here in CD pupae, but our

simulation models (see below) also confirm interindividual variation in WD diapause timers. If

we assumed that 15% of eclosed *R. pomonella* are WD individuals (see Fig. 3B), any proportion

eclosion above 15% indicated completion of development by CD individuals. Thus, the increase

in proportion eclosion following between c. 50 and 100 days of chilling (Fig. 3C) represented an

increase in the number of CD individuals that had experienced sufficient chilling to complete the

diapause program. This chilling threshold was quite variable among individuals in the

population; 60 days of chilling was required for 50% of flies to eclose, and more than 90 days of

chilling was required for maximum (91%) proportion eclosion (Fig. 3C). This sigmoidal pattern

in proportion eclosion can be explained by continuous variation among CD individuals in rate of

diapause development at 4°C (Fig. 3C). Those that eclosed after shorter chill durations were

likely able to complete a larger proportion of the diapause program at 4°C than other individuals

due to higher diapause development rates at 4°C, resulting in less time at 21°C required to

complete the diapause program.

Diapause is governed by physiological timers

When *R. pomonella* were exposed to increasing chill durations in Experiment 1, the mean

total development time increased (Fig. 4), suggesting that the diapause timers were temperature-

sensitive, as expected for physiological timers. However, low temperatures did not universally

accelerate development (decrease mean development time), as the classic chilling model (Fig.

2B) would predict. Indeed, at first glance, the positive relationship between chilling duration and

mean total development time (Fig. 4) is broadly consistent with a diapause timer that is

accelerated by high temperatures and has a broad thermal sensitivity (Fig. 2D). However, as

discussed above, the proportion of eclosed individuals varied with chill time due to the

requirement for chilling in most (likely CD) individuals, which is partially consistent with the

predictions of the classic, chilling only model (Fig. 2B). So, it seems that none of the simple

models individually explain all of the observations, especially given that WD and CD pupae

respond differently to chilling.

Figure 4. The effect of chilling on the time to complete diapause and post-diapause
development in *Rhagoletis pomonella*. Pupae were exposed to chilling (4°C; green) for a
specific duration at time zero (10 d post-pupariation) followed by warming (21°C). Each violin
plot represents the distribution of times to complete development from 10 d post-pupariation to
adult eclosion, with means represented by points, and error bars representing 95% confidence
intervals. Small error bars are obscured by symbols. Bars below the violin plots show the mean
duration of diapause development at 4°C (green – dictated by the chilling time) and 21°C
(yellow - estimated). We assumed a mean value of 30 days for post-diapause development
(morphogenesis; gray) at 21°C; gray bars are thus invariant across treatments.

Diapause timers can have broad thermal sensitivity

Although chilling was required for completion of diapause in most of our population, the

diapause program had a fairly broad thermal sensitivity for both WD and CD pupae. First, we

reasoned that diapause could progress at the two temperatures we used in Experiment 1: 4°C and

21°C, as explained below. Second, we also found diapause could complete after "chilling" at the

relatively warm temperature of 15°C (Fig. S3). We assumed the time to complete post-diapause

development was c. 30 days at 21°C for all individuals because this morphogenic process is

largely invariant in *R. pomonella* (Powell et al. 2020). Under this assumption, we inferred that

the average diapause program completed at 21°C (post-chill) for all chill treatments in

Experiment 1 because the mean time to eclose following chilling (yellow + gray bars; Fig. 4)

was always greater than the mean time to complete post-diapause morphogenesis at 21°C (gray

bars; Fig. 4). Therefore, pupae were able to complete part of the diapause program at 4°C and the

remaining proportion at 21°C (green and yellow bars; Fig. 4). This was likely true for WD pupae

(all or most of the eclosed individuals in the 2 – 4 weeks chilling treatments) and CD pupae (all

or most of the eclosed individuals after prolonged chilling; Fig. 4). Consistent with this finding,

none of our treatments included a clear period of cold-induced post-diapause quiescence (cf.

longest chill treatments in Fig. 2). Indeed, data collected by Feder et al. (1997) suggested that

more than 40 weeks of chilling are required for CD *R. pomonella* from hawthorn fruits to

complete diapause at low temperatures and enter quiescence, which is longer than the maximum

chilling duration (29 weeks) we used in our study.

 Thermal sensitivity of the diapause timer can change during ontogeny

While early chilling seemed to be required for completion of diapause in CD pupae

(Fig. 3C), prolonged cold exposure slowed down diapause development (Fig. 4). This suggested

that CD pupae had at least two diapause timers: one that governed early diapause development

and was accelerated by low temperatures (chilling), and another that governed late diapause

development and was accelerated by high temperatures (warming). Though a similar pattern is

expected when comparing diapause to post-diapause development in most insects (Jenkins et al.

2001; Hodek 2002), we emphasize that this ontogenetic shift in thermal sensitivity of

development rate clearly occurs during diapause in CD *R. pomonella* because our chill durations

were too short to induce quiescence.

Indeed, when we exposed groups of *R. pomonella* pupae to identical cold treatments at

different ages during Experiment 2, cold treatments later in development increased diapause

duration (slowed development) compared to cold treatments initiated at time zero (Fig. 5).

Because the age at chill exposure affected total development time, this suggested that the rate of

diapause development at low temperatures could change during diapause ontogeny. However,

there were multiple factors at play that could influence mean development time in these

treatments (Fig. 5), including differences in the proportion of individuals that eclosed (Fig. S1)

and how many of those individuals were CD vs. WD. Disentangling these effects empirically

was challenging because it is only possible to metabolically phenotype CD and WD accurately

when these individuals are held at warm temperatures for at least 45 days pre-chill (Dambroski

and Feder 2007; Toxopeus et al. 2021; Calvert et al. 2022), which we could not do in all

treatments. To address this issue, we developed simulation models to investigate whether

interindividual and temporal variation in diapause timer thermal sensitivity could explain the

trends we observed in our empirical diapause development data.

Figure 5. The effect of chilling at different ages on the on the time to complete diapause and post-diapause development in *Rhagoletis pomonella*. Pupae were exposed to chilling (4°C) for a specific duration at time zero (10 d post-pupariation), or 2 – 6 weeks after time zero followed by warming (21°C). Each violin plot represents the distribution of times to complete from 10 d post-pupariation to adult eclosion, with means represented by points, and error bars representing 95% confidence intervals. Small error bars are obscured by symbols. Bars below the violin plots show the mean duration of diapause development at 4°C (green – dictated by chilling time) and 21°C (yellow - estimated), and assumed duration of morphogenesis at 21°C (gray).

Accurate models must incorporate variation in diapause timers between diapause phenotypes,

among individuals, and across ontogeny of diapause development

A simulation model incorporating interindividual and ontogenetic variation in diapause

development rates, as well as differences between diapause phenotypes, provided an exceptional

fit to our empirical observations. We simulated a population of *R. pomonella* consisting of two

groups (WD and CD) with different diapause timers (Fig. 6A, B), and additional assumptions

described in Table A1. Our simulated population had approximately the same proportion of WD

(15%) and CD (85%) pupae as our empirical population (Fig. 3B). In the simulated population,

each WD pupa had one diapause timer with a broad thermal sensitivity (Fig. 6A), while each CD

pupa had two diapause timers: one for early diapause development (pre-threshold) progressing

only at low temperatures, and one for late diapause development with a broad thermal sensitivity

(Fig. 6B). Both the thermal sensitivities of WD and late CD development overlapped with the

thermal sensitivity of post-diapause development (Reissig et al. 1979). This allowed completion

of the diapause program at warm temperatures, followed immediately by post-diapause

development.

We modelled mean and interindividual variation in diapause development rate at two

temperatures (4°C and 21°C). Development time had a Gaussian distribution (Fig. S2), so we

modelled interindividual variation in development rates with an inverse Gaussian distribution.

However, the final simulation model fit to empirical data did not differ substantially if we

modelled rates with a Gaussian distribution instead (Fig. S4). Interindividual variation in

development rate was higher at the temperatures that were closer to the thermal optima for both

WD and CD populations, and was high early in diapause development for the CD population

(Fig. 6A, B). The final simulation model fit to empirical data was sensitive to both these

assumptions (Figs. S5, S6). We also assumed that development rates within an individual were

correlated across temperature (and time, in the case of CD pupae). That is, a simulated individual

with a development rate higher than the mean at 4°C would also have a development rate higher

than the mean at 21°C. However, the final simulation model fit to empirical data did not differ

substantially if we violated this assumption (Figs. S5, S6).

Figure 6. Summary of simulated diapause development in *Rhagoletis pomonella*.
Mean and standard deviation of diapause development rates in a simulated population of **(A)** 150
weak diapause (WD) pupae and **(B)** 850 chill-dependent diapause (CD) pupae under cold (green)
and warm (yellow) conditions during early and late diapause development. **(C)** Mean and
standard deviation of time to complete diapause in simulated WD pupae and early and late
diapause in simulated CD pupae at 4°C or 21°C. 0 indicates failure to complete the
developmental process. **(D-F)** Summary metrics of development time and proportion eclosion
phenotypes versus chill time for empirical data (symbols) and fitted simulation models (lines).
The full simulation model (black line, Eq'ns 1.1 – 2.4) is compared to models that only included
a single diapause phenotype (purple dotted line) or did not allow for ontogenetic shifts in thermal
sensitivity (TS) of diapause development rates (pink dashed line). Gray shading indicates the
95% confidence interval of the simulation summary metrics in (D) and (F). Error bars represent
standard error of the (D) mean or (F) proportion in empirical data. Small error bars are obscured
by symbols. CoV, coefficient of variation of mean development time.

In our simulation, only the WD pupae could complete diapause development at low or

high temperatures, while CD pupae required at least some chilling to complete diapause

development. The WD rate of diapause development was approximately twice as high at 21°C

than 4°C (Fig. 6A), resulting in faster completion of diapause (shorter development times) at

21°C (Fig. 6C). We note that WD pupae could have more than one diapause timer (e.g., different

timers in early vs. late development), but this added complexity was not necessary to explain the

general trends in our empirical data. For each simulated CD pupa, the early diapause timer only

allowed diapause to progress at low temperatures, and after some threshold chill duration (68.2 ±

16.4 d; Fig. 6C), a diapause timer with a different thermal sensitivity allowed CD pupae to

complete late diapause development at 4°C or 21°C (Fig. 6B). In late diapause, diapause

development rates were approximately four times higher at the warmer temperature (Fig. 6B),

resulting in faster completion of late diapause at 21°C than 4°C (Fig. 6C). We acknowledge that

a small proportion of CD pupae can likely complete pre-threshold diapause development at 21°C

in actual *R. pomonella* populations (Dambroski and Feder 2007), but that this pre-threshold

development is so slow that individuals did not complete development within the measurement

period of our experiments.

Using the full model described in the methods, our simulated population (Fig. 6A, B)

produced a close fit to the empirical data for mean development time (Fig. 6D), variation in

development time (Fig. 6E), and proportion eclosion (Fig. 6F) following different chilling

durations. If we only used one diapause phenotype (CD), or used two phenotypes (WD and CD)

but did not allow for ontogenetic variation in rates for CD pupae, the simulation produced a

considerably poorer fit to the empirical data (Fig. 6D-F). This reinforced the conclusions from

our empirical dataset that WD and CD are distinct phenotypes, rather than different ends of a

continuum of diapause intensity. Our full simulation model captured the prominent

discontinuity/transition in development time (Fig. 6D) and proportion eclosion (Fig. 6F) at

~60 days chilling caused by a transition in high proportion of WD to high proportion of CD

phenotypes among eclosed individuals. It likewise captured the decline in proportion eclosion at

long chill times (Fig. 6F) driven by overwinter mortality. The simulated variation in

development time (Fig. 6E) was over-estimated for moderate chill treatments (6 – 14 weeks).

Attempts to modify the distributions of diapause development rates (including pre-threshold

diapause development at 21°C) to improve the fit for development time metrics resulted in poor

fit for proportion eclosion or CoV metrics (see Fig. S7-9 for alternative simulations that modify

these parameters). A more complex model (e.g., more than two diapause timers) could

potentially improve the fit, but would not change the core principles of our simulation: diapause

is a temperature-sensitive process that varies among individuals, across diapause phenotypes,

and – in the case of CD pupae – across ontogeny.

Discussion

Our study demonstrates that principles of physiological time that predict developmental

duration during the growing season in ectotherms can also predict diapause developmental

duration under various simulated winter conditions. A few studies have shown that different

chilling temperatures impact diapause duration, with the fastest diapause development (shortest

diapause duration) at temperatures between 0°C and 7°C (Irwin et al. 2001; Sgolastra et al. 2010;

Xiao et al. 2013; Lehmann et al. 2017). Others have suggested that the impact of chilling

changes over time during diapause, with late chilling causing an increase in diapause duration

(Nomura and Ishikawa 2000; Xiao et al. 2013). Here we use data from *R. pomonella* to

parameterize well-fitting models that incorporate similar principles. Our experimental design

exposing each group of *R. pomonella* to two temperatures (chilling and warming) for different

durations and at different ages facilitated detection of at least two distinct phases of diapause

development, each with its own thermal sensitivity of diapause development rate. It likely would

have been difficult to detect this ontogenetic shift in thermal sensitivity of diapause development

in a classic thermal performance curve experiment, where multiple temperatures treatments were

used but each group of insects was exposed to a single temperature (e.g., Irwin et al. 2001;

Sgolastra et al. 2010; Xiao et al. 2013; Lehmann et al. 2017). Although we do not know the exact

shape of the diapause development rate thermal performance curves in *R. pomonella*, our simple

linear models are sufficient to explain most of the impacts of chilling and warming on two

diapause phenotypes in our populations under laboratory conditions.

Our experiments on thermal sensitivity of development in chill-dependent diapause

revealed only two distinct stages of diapause (early and late) rather than the three stages

(initiation, maintenance, termination) conceptualized in a broadly accepted ecophysiological

model of diapause development (Koštál 2006; Koštál et al. 2017). This is most likely because

diapause initiation in *R. pomonella* is relatively short; it takes less than 10 days post-pupariation

for pupae to suppress their metabolic rate (Ragland et al. 2009) and subsequently enter the

maintenance stage. Because we started our experiments 10 days post-pupariation (after

initiation), the 'early diapause' stage we describe in CD pupae likely represents maintenance,

while the 'late diapause' stage likely represents a long and gradual process of diapause

termination, also termed horotely (Hodek 1996, 2002). Additional work is needed to determine

whether this pattern (different stages of diapause development have different thermal

sensitivities) applies broadly to other insects, especially those that experience a broad range of

temperatures during diapause or do not routinely experience quiescence.

Increasing environmental temperatures will likely affect insect diapause development

across multiple seasons, with implications for life history timing. It is already known that warm

pre-winter conditions can negatively impact some insects by causing aversion or early

termination of diapause (Denlinger 2002; Koštál 2006; Toxopeus et al. 2021). In addition, warm

temperatures can cause energy drain that leads to mortality (Sinclair 2015; Roberts et al. 2021;

Nielsen et al. 2022). As autumn temperatures increase and winter warm spells become more

frequent (Williams et al. 2015; Marshall et al. 2020), early diapause development in chill-

dependent *R. pomonella* will likely slow down, resulting in an overall delay in the resumption of

post-diapause morphogenesis in the following summer or fall, and a possible mismatch in adult

eclosion and host fruit availability. Conversely, the predicted global increase in spring

temperatures (Williams et al. 2015; Marshall et al. 2020) may shorten diapause duration in

R. pomonella because this warming is likely to coincide with the later diapause development

process that is accelerated at high temperatures. Thus, any models that attempt to predict insect

phenology based on rates of post-diapause morphogenesis (e.g., Reissig et al. 1979) should take

diapause termination timing into account, especially for species with a temperature-dependent

diapause termination.

The physiological or cellular processes that determine diapause duration still remain

elusive (Hand et al. 2016). There are well-characterized hormonal responses that govern the

transition from diapause to post-diapause development in photoperiodically- or

pharmacologically-terminated diapause (Denlinger 2002; Denlinger et al. 2012). However, if

diapause termination is a gradual, horotelic process, diapause duration is likely governed by

some kind of diapause timer that is temperature-sensitive. For example, in a chill-dependent

diapause (or stage of diapause), the time to complete diapause (or a stage of diapause) may

depend on a particular physiological or cellular process that is active at low temperatures. There

could be a single candidate molecule that must be accumulated or degraded over time, such as a

particular gene product (e.g., *aldose reductase* and *sorbitol dehydrogenase* in *Pyrrhocoris*

apterus; Koštál et al. 2008), protein (e.g., p26 in *Artemia franciscana*; King and MacRae 2012)

or metabolite (e.g., alanine in *Pieris napi*; Lehmann et al. 2018); sorbitol in *Bombyx mori*, (Chino

1958). Conversely, there could be a suite of genes or molecules involved in regulating diapause

duration. For example, multiple studies have shown that gene expression changes over time in

diapause (Robich and Denlinger 2005; Kim et al. 2006; Rinehart et al. 2007; Reynolds and Hand

2009; Emerson et al. 2010; Ragland et al. 2010; Urbanski et al. 2010; Kankare et al. 2016; Green

and Kronforst 2019), although few have studied this process at low temperatures (Salminen et al.

2015; Koštál et al. 2017; Dowle et al. 2020; Pruisscher et al. 2022).

Our simulation and empirical datasets, and the transcriptomic studies cited above are both

highly consistent with temperature-modulated developmental processes dictating the duration of

diapause. Many of the temperature-sensitive genes in late diapause development of *R. pomonella*

are related to cell cycle and developmental progression (Meyers et al. 2016), although there is no

obvious morphological change during diapause. Indeed, genetic variants associated with

diapause duration in *R. pomonella* are enriched for genes also associated with egg-to-adult

developmental duration in *Drosophila melanogaster* (Dowle et al. 2020). In addition, neural

differentiation and remodeling (brain development) might occur during 6 months (42 weeks) of

chilling (Kharva et al. 2022), further suggesting that many developmental regulatory pathways

govern diapause duration just as they govern 'typical' (non-diapause) developmental duration.

We suggest that this may be a general phenomenon. Diapause has evolved rapidly and repeatedly

across the arthropod phylogeny (Ragland and Keep 2017; Denlinger 2022), suggesting that

diapausing clades likely coopt existing developmental machinery from non-diapausing ancestors

to initiate and regulate diapause development. This evolutionary hypothesis would also predict

that models of physiological time would fit developmental data from both diapause and non-

diapause development.

Conclusions

Applying the principles of physiological time to overwintering *R. pomonella* revealed

that diapause is temperature-sensitive, thermal sensitivity of diapause development rate can

change over time, and substantial portions of the diapause program progress more quickly at

warm temperatures. We also showed that the combination of two diapause timers (early and late

diapause) can account for interindividual synchronization of post-diapause development, even in

the absence of quiescence. Understanding the thermal sensitivity of overwintering development

(even if that development occurs in a "dormant" state) is important for predicting how warming

temperatures will impact the phenology of ectothermic animals. Additional work to elucidate the

cellular and physiological basis for diapause timers is still needed, with implications for

understanding how diapause has evolved.

Acknowledgements

The authors would like to thank M. Calvert, M. Davenport, N. Sanaei, E. Rader, and M. Haigh

for assistance with collecting processing *R. pomonella* in the lab. We would also like to thank J.

Kingsolver for comments on an earlier version of this manuscript. This work was funded by

National Science Foundation IOS 1700773 and DEB 1638951 grants to GJR.

Data and Code Accessibility

All analysis code and data referenced in this manuscript is available at

https://anonymous.4open.science/r/diapause-thermal-sensitivity-0DDC/.

Author Contributions Statement

JT, EJD, and GJR conceptualized the experiments. JT, EJD, and LA conducted the experiments.

JT and EJD analyzed the data. JT and GJR drafted the manuscript. All authors contributed to

editing and approving the final manuscript.

Literature Cited

Andrewartha, H. G. 1952. Diapause in relation to the ecology of insects. Biological Reviews 27:50–107.

Blanckenhorn, W. U. 1997. Altitudinal life history variation in the dung flies *Scathophaga stercoraria* and *Sepsis cynipsea*. Oecologia 109:342–352.

Buckley, L. B. 2022. Temperature-sensitive development shapes insect phenological responses to climate change. Current Opinion in Insect Science 52:100897.

Bush, G. L. 1969. Sympatric host race formation and speciation in frugivorous flies of the genus *Rhagoletis* (Diptera, Tephritidae). Evolution 23:237–251.

Calvert, M. B., M. M. Doellman, J. L. Feder, G. R. Hood, P. Meyers, S. P. Egan, T. H. Q. Powell, et al. 2022. Genomically correlated trait combinations and antagonistic selection contributing to counterintuitive genetic patterns of adaptive diapause divergence in *Rhagoletis* flies. Journal of Evolutionary Biology 35:146–163.

Chen, C., Q.-W. Xia, S. Fu, X.-F. Wu, and F.-S. Xue. 2014. Effect of photoperiod and temperature on the intensity of pupal diapause in the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae). Bulletin of Entomological Research 104:12.

Chen, Y.-S., C. Chen, H.-M. He, Q.-W. Xia, and F.-S. Xue. 2013. Geographic variation in diapause induction and termination of the cotton bollworm, *Helicoverpa armigera* Hübner (Lepidoptera: Noctuidae). Journal of Insect Physiology 59:855–862.

Cherrill, A. J., and M. Begon. 1989. Timing of life cycles in a seasonal environment: the temperature-dependence of embryogenesis and diapause in a grasshopper (*Chorthippus brunneus* Thunberg). Oecologia 78:237–241.

Chino, H. 1958. Carbohydrate metabolism in the diapause egg of the silkworm, *Bombyx*

mori—II: Conversion of glycogen into sorbitol and glycerol during diapause. Journal of Insect

Physiology 2:1–12.

Dambroski, H. R., and J. L. Feder. 2007. Host plant and latitude-related diapause

variation in *Rhagoletis pomonella*: a test for multifaceted life history adaptation on different

stages of diapause development. Journal of Evolutionary Biology 20:2101–2112.

Damos, P., and M. Savopoulou-Soultani. 2012. Temperature-driven models for insect

development and vital thermal requirements. Psyche: A Journal of Entomology 2012:1–13.

Dean, R. L., and P. J. Chapman. 1973. Bionomics of the apple maggot in eastern New

York. Search Agriculture 3:1–64.

Denlinger, D. L. 2002. Regulation of diapause. Annual Review of Entomology 47:93–

122.

Denlinger, D. L. 2022. Insect Diapause. Cambridge University Press, Cambridge.

Denlinger, D. L., G. D. Yocum, and J. P. Rinehart. 2012. Hormonal Control of

Diapause. Pages 430–463 *in* L. I. Gilbert, ed. Insect Endocrinology. Academic Press, San Diego.

Dingle, H., T. A. Mousseau, and S. M. Scott. 1990. Altitudinal variation in life cycle

syndromes of California populations of the grasshopper, *Melanoplus sanguinipes* (F.). Oecologia

84:199–206.

Dowle, E. J., T. H. Q. Powell, M. M. Doellman, P. J. Meyers, M. B. Calvert, K. K. O.

Walden, H. M. Robertson, et al. 2020. Genome-wide variation and transcriptional changes in

diverse developmental processes underlie the rapid evolution of seasonal adaptation. Proceedings

of the National Academy of Sciences 117:23960–23969.

Dupuis, J. R., B. A. Mori, and F. A. Sperling. 2016. *Trogus* parasitoids of *Papilio*

butterflies undergo extended diapause in western Canada (Hymenoptera, Ichneumonidae).

Journal of Hymenoptera Research 50:179.

Emerson, K. J., W. E. Bradshaw, and C. M. Holzapfel. 2010. Microarrays reveal early

transcriptional events during the termination of larval diapause in natural populations of the

mosquito, *Wyeomyia smithii*. PLoS One 5:e9574.

Feder, J. L., T. A. Hunt, and L. Bush. 1993. The effects of climate, host plant phenology

and host fidelity on the genetics of apple and hawthorn infesting races of *Rhagoletis pomonella*.

Entomologia Experimentalis et Applicata 69:117–135.

Feder, J. L., U. Stolz, K. M. Lewis, W. Perry, J. B. Roethele, and A. Rogers. 1997. The

effects of winter length on the genetics of apple and hawthorn races of *Rhagoletis pomonella*

(Diptera: Tephritidae). Evolution 51:1862–1876.

Geisert, R. W., and L. J. Meinke. 2013. Frequency and distribution of extended diapause

in Nebraska populations of *Diabrotica barberi* (Coleoptera: Chrysomelidae). Journal of

Economic Entomology 106:1619–1627.

Green, D. A., and M. R. Kronforst. 2019. Monarch butterflies use an environmentally

sensitive, internal timer to control overwintering dynamics. Molecular Ecology 28:3642–3655.

Hahn, D. A., and D. L. Denlinger. 2011. Energetics of insect diapause. Annual Review

of Entomology 56:103–121.

Hand, S. C., D. L. Denlinger, J. E. Podrabsky, and R. Roy. 2016. Mechanisms of animal

diapause: recent developments from nematodes, crustaceans, insects, and fish. American Journal

of Physiology 310:R1193–R1211.

Hanski, I. 1988. Four kinds of extra long diapause in insects: a review of theory and

observations. Annales Zoologici Fennici 25:37–53.

Higaki, M., and M. Toyama. 2012. Evidence for reversible change in intensity of

prolonged diapause in the chestnut weevil *Curculio sikkimensis*. Journal of Insect Physiology

58:56–60.

Hilbert, D. W., J. A. Logan, and D. M. Swift. 1985. A unifying hypothesis of

temperature effects on egg development and diapause of the migratory grasshopper, *Melanoplus*

sanguinipes (Orthoptera: Acrididae). Journal of Theoretical Biology 112:827–838.

Hodek, I. 1996. Diapause development, diapause termination and the end of diapause.

European Journal of Entomology 93:475–487.

Hodek, I. 2002. Controversial aspects of diapause development. European Journal of

Entomology 99:163–173.

Irwin, J. T., V. A. Bennett, and R. E. Lee. 2001. Diapause development in frozen larvae

of the goldenrod gall fly, *Eurosta solidaginis* Fitch (Diptera: Tephritidae). Journal of

Comparative Physiology B 171:181–188.

Jenkins, J., J. A. Powell, J. A. Logan, and B. J. Bentz. 2001. Low seasonal temperatures

promote life cycle synchronization. Bulletin of Mathematical Biology 63:573–595.

Kankare, M., D. J. Parker, M. Merisalo, T. S. Salminen, and A. Hoikkala. 2016.

Transcriptional differences between diapausing and non-diapausing *D. montana* females reared

under the same photoperiod and temperature. PLoS One 11:e0161852.

Kharva, H., J. L. Feder, D. A. Hahn, and S. B. Olsson. 2022. Rapid brain development

and reduced neuromodulator titres correlate with host shifts in *Rhagoletis pomonella*. Royal

Society Open Science 9:220962.

Kim, M., R. M. Robich, J. P. Rinehart, and D. L. Denlinger. 2006. Upregulation of two

actin genes and redistribution of actin during diapause and cold stress in the northern house

mosquito, *Culex pipiens*. Journal of Insect Physiology 52:1226–1233.

King, A. M., and T. H. MacRae. 2012. The small heat shock protein p26 aids

development of encysting *Artemia* embryos, prevents spontaneous diapause termination and

protects against stress. PLoS One 7:e43723.

Kingsolver, J. G., G. J. Ragland, and J. G. Shlichta. 2004. Quantitative genetics of

continuous reaction norms: thermal sensitivity of caterpillar growth rates. Evolution 58:1521–

1529.

Kingsolver, J. G., and H. A. Woods. 2016. Beyond thermal performance curves:

modeling time-dependent effects of thermal stress on ectotherm growth rates. The American

Naturalist 187:283–294.

Kipyatkov, V. E., and E. B. Lopatina. 2010. Intraspecific variation of thermal reaction

norms for development in insects: new approaches and prospects. Entomological Review

90:163–184.

Koštál, V. 2006. Eco-physiological phases of insect diapause. Journal of Insect

Physiology 52:113–127.

Koštál, V., T. Štětina, R. Poupardin, J. Korbelová, and A. W. Bruce. 2017. Conceptual

framework of the eco-physiological phases of insect diapause development justified by

transcriptomic profiling. Proceedings of the National Academy of Sciences 114:8532–8537.

Koštál, V., M. Tollarová, and D. Doležel. 2008. Dynamism in physiology and gene

transcription during reproductive diapause in a heteropteran bug, *Pyrrhocoris apterus*. Journal of

Insect Physiology 54:77–88.

Kutcherov, D. 2020. Stagewise resolution of temperature-dependent embryonic and

postembryonic development in the cowpea seed beetle *Callosobruchus maculatus* (F.). BMC

Ecology 20:50.

Lehmann, P., P. Pruisscher, V. Koštál, M. Moos, P. Šimek, S. Nylin, R. Agren, et al.

2018. Metabolome dynamics of diapause in the butterfly *Pieris napi*: distinguishing

maintenance, termination and post-diapause phases. Journal of Experimental Biology

221:jeb169508.

Lehmann, P., W. Van Der Bijl, S. Nylin, C. W. Wheat, and K. Gotthard. 2017. Timing

of diapause termination in relation to variation in winter climate. Physiological Entomology

42:232–238.

Lindestad, O., L. von Schmalensee, P. Lehmann, and K. Gotthard. 2020. Variation in

butterfly diapause duration in relation to voltinism suggests adaptation to autumn warmth, not

winter cold. Functional Ecology 34:1029–1040.

Lopatina, E. B., D. A. Kutcherov, and S. V. Balashov. 2014. The influence of diet on the

duration and thermal sensitivity of development in the linden bug *Pyrrhocoris apterus* L.

(Heteroptera: Pyrrhocoridae). Physiological Entomology 39:208–216.

Lyons-Sobaski, S., and S. H. Berlocher. 2009. Life history phenology differences

between southern and northern populations of the apple maggot fly, *Rhagoletis pomonella*.

Entomologia Experimentalis et Applicata 130:149–159.

Marshall, K. E., K. Gotthard, and C. M. Williams. 2020. Evolutionary impacts of winter

climate change on insects. Current Opinion in Insect Science 41:54–62.

Masaki, S. 2002. Ecophysiological consequences of variability in diapause intensity.

European Journal of Entomology 99:143–154.

Meyers, P. J., T. H. Q. Powell, K. K. O. Walden, A. J. Schieferecke, J. L. Feder, D. A.

Hahn, H. M. Robertson, et al. 2016. Divergence of the diapause transcriptome in apple maggot

flies: winter regulation and post-winter transcriptional repression. Journal of Experimental

Biology 219:2613–2622.

Moraiti, C. A., C. T. Nakas, and N. T. Papadopoulos. 2014. Diapause termination of

Rhagoletis cerasi pupae is regulated by local adaptation and phenotypic plasticity: escape in time

through bet-hedging strategies. Journal of Evolutionary Biology 27:43–54.

Nielsen, M. E., P. Lehmann, and K. Gotthard. 2022. Longer and warmer prewinter

periods reduce post-winter fitness in a diapausing insect. Functional Ecology 36:1151–1162.

Nomura, M., and Y. Ishikawa. 2000. Biphasic effect of low temperature on completion

of winter diapause in the onion maggot, *Delia antiqua*. Journal of insect Physiology 46:373–377.

Papanastasiou, S. A., and N. T. Papadopoulos. 2014. Description of *Rhagoletis cerasi*

(Diptera: Tephritidae) pupal developmental stages: indications of prolonged diapause. Journal of

Insect Science 14:156.

Powell, J. A., and B. J. Bentz. 2009. Connecting phenological predictions with

population growth rates for mountain pine beetle, an outbreak insect. Landscape Ecology

24:657–672.

Powell, T. H. Q., A. Nguyen, Q. Xia, J. L. Feder, G. J. Ragland, and D. A. Hahn. 2020.

A rapidly evolved shift in life-history timing during ecological speciation is driven by the

transition between developmental phases. Journal of Evolutionary Biology 33:1371–1386.

Prokopy, R. J. 1968. Influence of photoperiod, temperature, and food on initiation of

diapause in the apple maggot. The Canadian Entomologist 100:318–329.

Pruisscher, P., P. Lehmann, S. Nylin, K. Gotthard, and C. W. Wheat. 2022. Extensive

transcriptomic profiling of pupal diapause in a butterfly reveals a dynamic phenotype. Molecular

Ecology 31:1269–1280.

Ragland, G. J., D. L. Denlinger, and D. A. Hahn. 2010. Mechanisms of suspended

animation are revealed by transcript profiling of diapause in the flesh fly. Proceedings of the

National Academy of Sciences 107:14909–14914.

Ragland, G. J., J. Fuller, J. L. Feder, and D. A. Hahn. 2009. Biphasic metabolic rate

trajectory of pupal diapause termination and post-diapause development in a tephritid fly.

Journal of Insect Physiology 55:344–350.

Ragland, G. J., and E. Keep. 2017. Comparative transcriptomics support evolutionary

convergence of diapause responses across Insecta. Physiological Entomology 42:246–256.

R Core Team. 2022. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rebaudo, F., and V.-B. Rabhi. 2018. Modeling temperature-dependent development rate

and phenology in insects: review of major developments, challenges, and future directions.

Entomologia Experimentalis et Applicata 166:607–617.

Régnière, J., J. Powell, B. Bentz, and V. Nealis. 2012. Effects of temperature on

development, survival and reproduction of insects: experimental design, data analysis and

modeling. Journal of Insect Physiology 58:634–647.

Reissig, W. H., J. Barnard, R. W. Weires, E. H. Glass, and R. W. Dean. 1979. Prediction

of apple maggot fly emergence from thermal unit accumulation. Environmental Entomology

8:51–54.

Reynolds, J. A., and S. C. Hand. 2009. Embryonic diapause highlighted by differential

expression of mRNAs for ecdysteroidogenesis, transcription and lipid sparing in the cricket

Allonemobius socius. Journal of Experimental Biology 212:2075–2084.

Rinehart, J. P., A. Li, G. D. Yocum, R. M. Robich, S. A. Hayward, and D. L. Denlinger.

2007. Up-regulation of heat shock proteins is essential for cold survival during insect diapause.

Proceedings of the National Academy of Sciences 104:11130–11137.

Roberts, K. T., N. E. Rank, E. P. Dahlhoff, J. H. Stillman, and C. M. Williams. 2021.

Snow modulates winter energy use and cold exposure across an elevation gradient in a montane

ectotherm. Global Change Biology 27:6103–6116.

Robich, R. M., and D. L. Denlinger. 2005. Diapause in the mosquito *Culex pipiens*

evokes a metabolic switch from blood feeding to sugar gluttony. Proceedings of the National

Academy of Sciences 102:15912–15917.

Rull, J., E. Tadeo, R. Lasa, and M. Aluja. 2016. The effect of winter length on survival

and duration of dormancy of four sympatric species of *Rhagoletis* exploiting plants with different

fruiting phenology. Bulletin of entomological research 106:818–826.

Salminen, T. S., L. Vesala, A. Laiho, M. Merisalo, A. Hoikkala, and M. Kankare. 2015.

Seasonal gene expression kinetics between diapause phases in *Drosophila virilis* group species

and overwintering differences between diapausing and non-diapausing females. Scientific

Reports 5.

Scranton, K., and P. Amarasekare. 2017. Predicting phenological shifts in a changing

climate. Proceedings of the National Academy of Sciences 114:13212–13217.

Sgolastra, F., J. Bosch, R. Molowny-Horas, S. Maini, and W. P. Kemp. 2010. Effect of

temperature regime on diapause intensity in an adult-wintering Hymenopteran with obligate

diapause. Journal of Insect Physiology 56:185–194.

Shi, P., F. Ge, Y. Sun, and C. Chen. 2011. A simple model for describing the effect of

temperature on insect developmental rate. Journal of Asia-Pacific Entomology 14:15–20.

Shingleton, A. W., G. C. Sisk, and D. L. Stern. 2003. Diapause in the pea aphid

(*Acyrthosiphon pisum*) is a slowing but not a cessation of development. BMC Developmental

Biology 3:7.

Sinclair, B. J. 2015. Linking energetics and overwintering in temperate insects. Journal

of Thermal Biology 54:5–11.

Sinclair, B. J., K. E. Marshall, M. A. Sewell, D. L. Levesque, C. S. Willett, S. Slotsbo,

Y. Dong, et al. 2016. Can we predict ectotherm responses to climate change using thermal

performance curves and body temperatures? Ecology Letters 19:1372–1385.

Sridhar, V., and P. V. R. Reddy. 2013. Use of degree days and plant phenology: a

reliable tool for predicting insect pest activity under climate change conditions. Pages 287–294

in H. C. P. Singh, N. K. S. Rao, and K. S. Shivashankar, eds. Climate-Resilient Horticulture:

Adaptation and Mitigation Strategies. Springer, India.

Tauber, M. J., and C. A. Tauber. 1972. Geographic variation in critical photoperiod and

in diapause intensity of *Chrysopa carnea* (Neuroptera). Journal of Insect Physiology 18:25–29.

Tauber, M. J., C. A. Tauber, and S. Masaki. 1986. Seasonal Adaptations of Insects.

Oxford University Press, New York.

Tauber, M., and C. Tauber. 1976. Insect seasonality: diapause maintenance, termination,

and postdiapause development. Annual Review of Entomology 21:81–107.

Taylor, F. 1981. Ecology and evolution of physiological time in insects. The American

Naturalist 117:1–23.

Toxopeus, J., L. Gadey, L. Andaloori, M. Sanaei, and G. J. Ragland. 2021. Costs of

averting or prematurely terminating diapause associated with slow decline of metabolic rates at

low temperature. Comparative Biochemistry and Physiology A 255:110920.

Trudgill, D. L., A. Honek, D. Li, and N. M. van Straalen. 2005. Thermal time - concepts

and utility. Annals of Applied Biology 146:1–14.

Urbanski, J. M., J. B. Benoit, M. R. Michaud, D. L. Denlinger, and P. Armbruster. 2010.

The molecular physiology of increased egg desiccation resistance during diapause in the invasive

mosquito, *Aedes albopictus*. Proceedings of the Royal Society B 277:2683–2692.

van Straalen, N. M. 1983. Physiological time and time-invariance. Journal of Theoretical

Biology 104:349–357.

Vargas, R. I., W. A. Walsh, E. B. Jang, J. W. Armstrong, and D. T. Kanehisa. 1996.

Survival and development of immature stages of four Hawaiian fruit flies (Diptera: Tephritidae)

reared at five constant temperatures. Annals of the Entomological Society of America 89:64–69.

Williams, C. M., H. A. Henry, and B. J. Sinclair. 2015. Cold truths: how winter drives

responses of terrestrial organisms to climate change. Biological Reviews 90:214–35.

Wilsterman, K., M. A. Ballinger, and C. M. Williams. 2021. A unifying, eco-

physiological framework for animal dormancy. Functional Ecology 35:11–31.

Xiao, H., S. Wu, C. Chen, and F. Xue. 2013. Optimal low temperature and chilling

period for both summer and winter diapause development in *Pieris melete*: based on a similar

mechanism. PLoS One 8:e56404.

Yurk, B. P., and J. A. Powell. 2010. Modeling the effects of developmental variation on

insect phenology. Bulletin of Mathematical Biology 72:1334–1360.

Zhu, D., Y. Yang, and Z. Liu. 2009. Reversible change in embryonic diapause intensity

by mild temperature in the Chinese rice grasshopper, *Oxya chinensis*. Entomologia

Experimentalis et Applicata 133:1–8.

1032

1033

 Figure Legends

Figure 1. Example durations of diapause for insects that typically enter diapause in late

summer or early autumn. (A) Some insects remain in diapause under warm (yellow) and cool

(green) conditions for most of the year. **(B)** Many insects complete diapause during winter when

it is cool (green), and enter a post-diapause dormancy called quiescence (blue), resuming post-

diapause development (morphogenesis; grey) immediately when temperatures become

permissive in the spring or early summer. **(C)** Warm summer or autumn conditions can cause

individuals to complete diapause rapidly, referred to as 'weak diapause' or 'shallow diapause' in

some species **(D)** Non-diapause individuals never enter diapause and may complete

morphogenesis prior to winter.

1044

Figure 2. Hypothesized effect of a single (A) chronological or (B – D) physiological diapause

timer on development duration under different combinations of chilling and warming.

Transition from cold to warm temperature indicated by dashed line. Development duration

includes diapause development under cold (green) and warm (yellow) conditions, and post-

diapause (e.g., morphogenic) development (grey) that can only occur under warm conditions. If

individuals remain chilled after diapause completes (arrows), cold-induced post-diapause

quiescence (dark blue) occurs until animals are returned to warm temperatures. Insets:

relationship between temperature (T) and diapause development rate (r) at a low (chilling) and

high (warming) temperatures based on an example thermal sensitivity (TS) of the diapause timer.

The equations that explain the relationships between development rates, chill duration, and

development time are fully explained in Appendix A. Multiple diapause timers may exist (e.g.,

for different stages of diapause), which would result in more complex relationships between

chilling duration and development duration (not shown).

1058

Figure 3. Diapause incidence and the effect of chilling on diapause completion in *Rhagoletis*

***pomonella*.** **(A)** Mass-specific metabolic rate after 10 or 45 days of constant warmth post-

pupariation was used to determine diapause phenotype: non-diapause (ND), weak diapause

(WD), and chill-dependent diapause (CD). Each open circle represents one individual. Filled

black circles represent the mean of each group. Error bars represent 95% confidence intervals;

small error bars are obscured by symbols. **(B)** The proportion of ND, WD, and CD pupae

inferred from the metabolic rate data in **(A)** and duration of total development (Fig. S2) in groups

that were not metabolically-phenotyped. **(C)** Each point represents the proportion of c. 100

pupae that completed diapause and post-diapause development (eclosed as adults) following

exposure to chilling (4°C) for a specific duration at time zero (10 d post-pupariation). Error bars

represent the standard error of proportion. Black line is a cumulative distribution fit with a mean

± standard deviation of 60 ± 15 d chilling.

1071

Figure 4. The effect of chilling on the time to complete diapause and post-diapause

development in *Rhagoletis pomonella*. Pupae were exposed to chilling (4°C; green) for a

specific duration at time zero (10 d post-pupariation) followed by warming (21°C). Each violin

plot represents the distribution of times to complete development from 10 d post-pupariation to

adult eclosion, with means represented by points, and error bars representing 95% confidence

intervals. Small error bars are obscured by symbols. Bars below the violin plots show the mean

duration of diapause development at 4°C (green – dictated by the chilling time) and 21°C

(yellow - estimated). We assumed a mean value of 30 days for post-diapause development

(morphogenesis; grey) at 21°C; grey bars are thus invariant across treatments.

1081

Figure 5. The effect of chilling at different ages on the on the time to complete diapause and

post-diapause development in *Rhagoletis pomonella*. Pupae were exposed to chilling (4°C) for

a specific duration at time zero (10 d post-pupariation), or 2 – 6 weeks after time zero followed

by warming (21°C). Each violin plot represents the distribution of times to complete from 10 d

post-pupariation to adult eclosion, with means represented by points, and error bars representing

95% confidence intervals. Small error bars are obscured by symbols. Bars below the violin plots

show the mean duration of diapause development at 4°C (green – dictated by chilling time) and

21°C (yellow - estimated), and assumed duration of morphogenesis at 21°C (grey).

1090

Figure 6. Summary of simulated diapause development in *Rhagoletis pomonella*.

Mean and standard deviation of diapause development rates in a simulated population of **(A)** 150

weak diapause (WD) pupae and **(B)** 850 chill-dependent diapause (CD) pupae under cold (green)

and warm (yellow) conditions during early and late diapause development. **(C)** Mean and

standard deviation of time to complete diapause in simulated WD pupae and early and late

diapause in simulated CD pupae at 4°C or 21°C. 0 indicates failure to complete the

developmental process. **(D-F)** Summary metrics of development time and proportion eclosion

phenotypes versus chill time for empirical data (symbols) and fitted simulation models (lines).

The full simulation model (black line, Eq'ns 1.1 – 2.4) is compared to models that only included

a single diapause phenotype (purple dotted line) or did not allow for ontogenetic shifts in thermal

sensitivity (TS) of diapause development rates (pink dashed line). Gray shading indicates the

95% confidence interval of the simulation summary metrics in (D) and (F). Error bars represent

standard error of the (D) mean or (F) proportion in empirical data. Small error bars are obscured

by symbols. CoV, coefficient of variation of mean development time.

1105

Appendix A: Simulation Model Building

Relationships between diapause development rates, temperature, and development duration

Development rates (R) are often expressed as proportion development per day (d^{-1}), and

the time (d) to complete development (Y) is simply equal to the inverse of that development rate

($Y = 1/R$) (Kipyatkov and Lopatina 2010; Shi et al. 2011; Damos and Savopoulou-Soultani

2012). The classic method to determine thermal sensitivity of a development rate is to rear

groups of individuals at different temperatures and calculate development rate at each

temperature as the inverse of developmental duration (Kipyatkov and Lopatina 2010). However,

we can assess the relative breadth of thermal sensitivity of diapause development, as well as

shifts in thermal sensitivity over time by using the following experimental design. Groups of

diapausing individuals are exposed to a cold temperature (e.g., 4°C) for different lengths of time

(X), followed by exposure to a warm temperature (e.g., 21°C) until evidence of post-diapause

morphogenesis is observed. These temperatures must be selected such that post-diapause

morphogenesis can only occur at the warm temperature (rate of morphogenesis at the cold

temperature = 0). We can represent R as the rate of diapause development at the cold

temperature, r as the rate of diapause development at the warm temperature, and r_g as the rate of

post-diapause morphogenesis at the warm temperature. Total time to complete development

following a range of chilling durations can then be modelled in several ways, depending on the

thermal sensitivity of diapause development.

1125

<u>A single diapause timer: narrow vs. broad thermal sensitivity</u>

If diapause development rate has a narrow thermal sensitivity such that diapause only

progresses at low temperatures (e.g., Fig. 2B), the rate of diapause development at the warm

temperature is zero, i.e. $r = 0$. Therefore, any cold exposure duration that is less than the time

required to complete diapause at 4°C ($1/R$) will result in a total development time (Y) of zero

because diapause is unable to complete (and no post-diapause morphogenesis is observed) at the

warm temperature:

$$Y = 0 \text{ when } X < \frac{1}{R} \qquad\qquad \text{(Eq'n A1.1)}$$

If an individual is kept at the cold temperature for a sufficient time to complete diapause ($1/R$),

post-diapause morphogenesis will resume when transferred to the permissive warm temperature,

and total development time will be:

$$Y = X + \frac{1}{r_g} \text{ when } X \geq \frac{1}{R} \qquad\qquad \text{(Eq'n A1.2)}$$

where $1/r_g$ is the time to complete post-diapause morphogenesis at the warm temperature. If an

individual is chilled for longer than is needed to complete diapause ($> 1/R$), that individual will

enter post-diapause quiescence, and any additional chilling will increase total development time

with a slope of 1.

1142

Conversely, we can model a broad thermal sensitivity of diapause development rate, i.e.

diapause can progress at both cold and warm temperatures (e.g., Fig. 2C and D). In this case, if

the individual is never chilled:

$$Y = \frac{1}{r} + \frac{1}{r_g} \text{ when } X = 0 \qquad\qquad \text{(Eq'n A2.1)}$$

where $1/r$ is the time to complete diapause at the warm temperature. Conversely, if the individual

is chilled for sufficient time to complete diapause development:

$$Y = X + \frac{1}{r_g} \text{ when } X \geq \frac{1}{R} \qquad\qquad \text{(Eq'n A2.2)}$$

If an individual is chilled for longer than is needed to complete diapause, that individual will

enter post-diapause quiescence, which will increase total development time with a slope of 1. If

the individual is chilled at the cold temperature for a duration $< 1/R$, some proportion of diapause

development $(X \times R)$ completes in the cold, and the remaining proportion of diapause

development $(1 - X \times R)$ completes during post-chill warming. The time to complete diapause

development is therefore:

$$Y = X + \frac{1-(X \times R)}{r} + \frac{1}{r_g} \text{ when } 0 \leq X < \frac{1}{R} \qquad \text{(Eq'n A2.3)}$$

which simplifies to a classic linear relationship $(y = mx + b)$ between X and Y:

$$Y = \frac{r-R}{r}X + \frac{1}{r} + \frac{1}{r_g} \text{ when } 0 \leq X < \frac{1}{R} \qquad \text{(Eq'n A2.4)}$$

Using this equation, we see that the slope will be negative if chilling accelerates diapause

development $(R > r$; Fig. 2C), and positive if warming accelerates diapause development $(r > R$;

Fig. 2D).

1162

<u>Two diapause timers: incorporating a shift in thermal sensitivity over time</u>

If thermal sensitivity of development can change during ontogeny, we must introduce

additional diapause development rates. R_e and r_e are the rates of diapause development at cold

and warm temperatures (respectively) early in diapause devleopment, and R_l and r_l are the rates

of diapause development at cold and warm temperatures (respectively) late in diapause

development. The relationship between chill time (X) and total development time (Y) in this

more complicated scenario can be broken down into early and late diapause sections.

1170

First, let's consider a narrow thermal sensitivity for our early diapause timer, such that

diapause only progresses at low temperatures. Any cold exposure duration that is less than the

time required to complete early diapause at 4°C ($1/R_e$) will result in a total development time (Y)

of zero because early diapause is unable to complete at the warm temperature (preventing any

further development), similar to Equation A1.1:

$$Y = 0 \text{ when } X < \frac{1}{R_e}$$ (Eq'n A3.1)

Conversely, we can consider a broad thermal sensitivity for our early diapause timer. For chill

times that are insufficient to complete early diapause, some proportion of early diapause

development ($X \times R_e$) completes in the cold, and the remaining proportion of early diapause

development ($1 - X \times R_e$) completes during post-chill warming. In this case, the total time to

complete diapause is:

$$Y = X + \frac{1-(X \times R_e)}{r_e} + \frac{1}{r_l} + \frac{1}{r_g} \text{ when } X < \frac{1}{R_e}$$ (Eq'n A3.2)

where X is the duration of partial early diapause development in the cold, $(1 - X \times R_e)/r_e$ is the

time to complete the rest of early diapause at the warm temperature, and $1/r_l$ is the time to

completed late diapause development at the warm temperature. This simplifies to a classic linear

relationship ($y = mx + b$) between X and Y:

$$Y = \frac{r_e - R_e}{r_e} X + \frac{1}{r_e} + \frac{1}{r_l} + \frac{1}{r_g} \text{ when } X < \frac{1}{R_e}$$ (Eq'n A3.3)

Using this equation, we see that the slope will be negative if chilling accelerates diapause

development ($R_e > r_e$; Fig. 2C), and positive if warming accelerates diapause development ($r_e >$

R_e; Fig. 2D).

1191

For the late diapause timer, we can consider the same two scenarios as for our early

diapause timer. If the late diapause timer has a narrow thermal sensitivity (diapause only

progresses at low temperatures), any cold exposure duration that is less than the time required to

complete early *and* late diapause at 4°C ($1/R_e + 1/R_l$) will result in a total development time (Y)

of zero because diapause is unable to complete at the warm temperature, similar to Equation

A1.1:

$$Y = 0 \text{ when } X < \frac{1}{R_e} + \frac{1}{R_l}$$ (Eq'n A3.4)

This would result in a development time pattern that is indistinguishable from a single diapause

timer that also has a cold-optimized narrow thermal sensitivity. However, if the late diapause

timer has a broad thermal sensitivity, chilling that is sufficient for completion of early diapause

but too short for completion of late diapause will result in the following:

$$Y = X + \frac{1 - ([X - \frac{1}{R_e}] \times R_l)}{r_l} + \frac{1}{r_g} \text{ when } \frac{1}{R_e} \leq X < \frac{1}{R_e} + \frac{1}{R_l}$$ (Eq'n A3.5)

where X is the duration of early diapause and partial late diapause development in the cold, and

$(1 - (X - 1/R_e) \times R_l)/r_l$ is the time to complete the rest of late diapause at the warm temperature.

This simplifies to a classic linear relationship ($y = mx + b$) between X and Y:

$$Y = \frac{r_l - R_l}{r_l} X + \frac{R_l + R_e}{R_e r_l} + \frac{1}{r_g} \text{ when } \frac{1}{R_e} \leq X < \frac{1}{R_e} + \frac{1}{R_l}$$ (Eq'n A3.6)

Using this equation, we see that the slope will be negative if chilling accelerates late diapause

development ($R_l > r_l$; Fig. 2C), and positive if warming accelerates diapause development ($r_l >$

R_l; Fig. 2D).

1211

When individuals are chilled for a sufficient time to complete diapause or for longer, total

development time will be expressed similar to Equations A1.2 and A2.2:

$$Y = X + \frac{1}{r_g} \text{ when } X \geq \frac{1}{R_e} + \frac{1}{R_l}$$ (Eq'n A3.7)

This relationship holds as long as both diapause processes can complete at 4°C, and does not

depend on whether the early and late diapause timers have a narrow or broad thermal sensitivity.

More complicated models are also possible (e.g., 3 or more diapause timers, each with their own

thermal sensitivity), but we were able to simulate a population closely predicting empirical

observations in *R. pomonella* using a maximum of two diapause timers per diapause phenotype.

1220

Simulation of Rhagoletis pomonella diapause development

We calculated total development time (since time zero) for each simulated individual

(and whether that individual would complete development) under a range of chill durations using

the assumptions in Table A1 and the equations in the main text. The rationale behind these

equations and assumptions (as well as tests of those assumptions) is explained in the subsections

below. Each individual in the population was simulated with its own diapause development rates

at 4°C and 21°C, and these rates were sampled from simulated distributions of diapause

development rates. The population distributions (mean and variation) of these simulated diapause

rates are summarized in Table A2.

1230

1231 **Table A1. Assumptions for development of simulated weak diapause and chill-dependent**
1232 **diapause pupae chilled at 4°C at time zero (10 d post-pupariation) for different durations,**
1233 **followed by warming at 21°C.**

	Weak diapause	**Chill-dependent diapause**
Thermal sensitivity	Diapause progresses across a wide range of temperatures and is accelerated by warming.	Early diapause only progresses well at low temperatures, while late diapause progresses across a wide range of temperatures and is accelerated by warming.
Variation in development rates	Diapause development rates at 4°C (R) and 21°C (r) have an inverse-Gaussian distribution.	The rate of early diapause development at 4°C (R_e) has an inverse-Gaussian distribution.
		Early diapause development cannot progress at 21°C ($r_e = 0$, or very close to 0).
	The ratio of the rate at 4°C to the rate at 21°C does not vary among individuals.	The rates of late diapause development at 4 °C (R_l) and 21 °C (r_l) are correlated with the rate of early diapause development.
	The rate of post-diapause morphogenesis at 21°C (r_g) is invariant.	The rate of post-diapause morphogenesis at 21°C (r_g) is invariant.
Mortality due to chilling	Pupae die if chilled for X_z days.	Pupae die if chilled for X_z days.

1234

1235

Table A2. Model parameters of a simulated population of 150 weak diapause and 850 chill-dependent diapause pupae. Development rates were modelled with inverse Gaussian distributions with mean μ and shape λ (Fig. S4). Corresponding mean ($\bar{x}$) and standard deviation (*s*) of times to complete diapause or a component of diapause (early and late) at a single temperature (4°C and 21°C, respectively) are shown in the same row. The effect of increasing or decreasing these rates can be seen in Fig. S7.

	Development rates (d^{-1})		Development times (d)	
	Parameter	Values	Parameter	Values
Weak diapausers (*N* = 150)				
At 21 °C	r	$\mu = 0.030,\ \lambda = 0.25$	$\dfrac{1}{r}$	$\bar{x} = 35.9, s = 11.9$
At 4 °C	R	$\mu = 0.017, \lambda = 0.1$	$\dfrac{1}{R}$	$\bar{x} = 65.7, s = 21.7$
Chill-dependent diapausers (*N* = 850)				
At 21 °C	r_e	$\mu = 0$	$\dfrac{1}{r_e}$	0 (no development)
	r_l	$\mu = 0.014,\ \lambda = 10$	$\dfrac{1}{r_l}$	$\bar{x} = 71.0, s = 2.7$
At 4 °C	R_e	$\mu = 0.0155,\ \lambda = 0.3$	$\dfrac{1}{R_e}$	$\bar{x} = 68.2, s = 16.4$
	R_l	$\mu = 0.0035, \lambda = 3$	$\dfrac{1}{R_l}$	$\bar{x} = 285.6, s = 10.6$

Thermal sensitivity, mortality and equations for simulated weak diapause

We simulated our **weak diapause (WD)** individuals with a broad thermal sensitivity of

diapause development (Table A1) for the following reasons. 1) Our empirical data (and that of

others; Dambroski and Feder 2007; Calvert et al. 2022) shows that WD *R. pomonella* can eclose

rapidly (35 – 65 days post-pupariation) with no chilling, suggesting diapause can complete at

warm temperatures. 2) The proportion of pupae that eclosed with little to no chilling (0 – 4

weeks) was constant at approximately 15% (Fig. 3C), and approximately 15 % of our population

consisted of WD pupae (Fig. 3B), suggesting that all flies that eclosed with 0 – 4 weeks chilling

were WD individuals. 3) During these first 4 weeks of chilling, mean total development time

1253   increased with chilling, but post-chill development time decreased (yellow + gray in Fig. 4),

1254   suggesting diapause could progress at 4°C as well as 21°C. Further, the positive slope of mean

1255   development time vs. chill time during these first 4 weeks of chilling (Fig. 4) suggested that WD

1256   pupal diapause was accelerated by warm temperatures (cf. Fig. 2D).

1257

To model total development time in our simulated WD pupae, we used Equation A2.4

(above), and Equations A2.5 – A2.6 (below) [Equations 1.1 – 1.3 in the main text]. We know

that WD pupae are killed by moderate chilling (Toxopeus et al. 2021), so we also incorporated a

mortality term into our final model. Equation A2.5 is a slightly modified Equation A2.2. that

incorporates this mortality:

$Y = X + \dfrac{1}{r_g}$ when $\dfrac{1}{R} \leq X < X_z$ (Eq'n A2.5)

Where X_z is the chill time that causes mortality in that WD individual. Chill times longer than X_z

resulted in a total development time of zero because no eclosion was observed:

$Y = 0$ when $X \geq X_z$ (Eq'n A2.6)

See Fig. S8 for the impacts of removing WD mortality from our models.

1268

Thermal sensitivity and equations for simulated chill-dependent diapause

We simulated our **chill-dependent diapause (CD)** individuals with two diapause timers:

an early diapause timer with a narrow thermal sensitivity and a late diapause timer with a broad

thermal sensitivity of diapause development (Table A1). Our reasoning for this was as follows.

1) For moderate chill times (6 – 14 weeks), as chill time increased proportion eclosion increased

in an approximately sigmoidal pattern (Fig. 3C). We moved forward with the likely assumption

that this trend in proportion eclosion was driven by CD pupae, as WD pupae seemed to exhibit a

proportion eclosion pattern that did not depend on chilling. Therefore, CD pupae seemed to

require chilling (e.g. due to a narrow thermal sensitivity of diapause development) early in

development. 2) We saw no evidence for post-diapause quiescence in the chilling treatments we

used in our study; even our longest chill treatment (29 weeks) resulted in post-chill development

time that took longer than the mean time to complete morphogenesis at 21°C (Fig. 4). Therefore,

although chilling was required early in diapause development, there seemed to be a later

diapause process with a broad thermal sensitivity – one that could complete at either 4°C or

21°C. Further, the positive slope of mean development time vs. chill time following prolonged

chilling (17 – 29 weeks) (Fig. 4) suggested that this late diapause process in CD pupae was

accelerated by warm temperatures (cf. Fig. 2D).

1286

To model total development time in our simulated CD pupae, we used Equations A3.1

and A3.6 (above), and Equations A3.8 – A3.9 (below) [Equations 2.1 – 2.4 in the main text]. We

saw increased mortality after prolonged chilling (Fig. S1) that was likely due to death of CD

pupae (because WD pupae likely die after moderate chilling). Equation A3.8 is a slightly

modified Equation A3.7. that incorporates mortality:

$$Y = X + \frac{1}{r_g} \text{ when } \frac{1}{R_e} + \frac{1}{R_l} \leq X < X_z \qquad \text{(Eq'n 3.8)}$$

where X_z is the chill time that causes CD pupa mortality. Chill times longer than X_z resulted in a

total development time of zero because no eclosion was observed:

$$Y = 0 \text{ when } X \geq X_z \qquad \text{(Eq'n 3.9)}$$

See Fig. S8 for the impacts of removing CD mortality from our models.

1297

 Variation in diapause development rates

1299   Because development time generally had a Gaussian distribution (Fig. S2), we simulated

1300   variation in diapause development rates (r, R, R_s, r_d, R_d) among individuals with inverse

1301   Gaussian distributions (Table A1). Normal distributions of diapause development rates did not

1302   substantially improve the simulation model fit to empirical data (Fig. S4).

1303

We assumed that different rates of diapause development (at 4°C vs. 21°C, pre- vs. post-

threshold) were correlated within an individual (Table A1). This assumption is based on

principles and examples from the morphogenesis development rate literature. For example,

development times across life history stages can be correlated within individuals such that some

individuals consistently develop faster than others at multiple temperatures. This can be seen in

yellow dung fly larvae vs. pupae (Blanckenhorn 1997) and grasshopper eggs early and late in

embryogenesis (Cherrill and Begon 1989). See Figs. S5 and S6 for the minimal impact of this

assumption about rate correlation within an individual on model fit.

1312

We allowed for a change in the magnitude of interindividual variation across

temperatures (e.g., variation in R [4°C] and r [21°C]), and across ontogeny (e.g., variation in Rs

[pre-threshold] and Rd [post-threshold] at 4°C). Variation in morphogenic development rates can

change across temperature; for example several fruit fly species shown less variation in pupal

development rate at high temperatures than low temperatures (Vargas et al. 1996). The

morphogenic development rate/time literature contains several examples of changes in the

variability of development rate across life stages, e.g., egg development time is less variable than

pupal development time in Mountain Pine Beetle at a given temperature (Yurk and Powell

2010). See Figs. S5 and S6 for the substantial impact of these assumptions about variation in

rates across temperature and ontogeny, respectively, on model fit.

1323

1324   We did not model interindividual variation in time to complete post-diapause

1325 morphogenesis (Table A1) because this post-diapause morphogenesis is minimally variant

1326 among individuals of *R. pomonella* at a given temperature, and variation in eclosion time is

1327 generally driven by variation in diapause duration in *R. pomonella* (Powell et al. 2020)

1328

Alternative thermal sensitivity patterns

While the simulation model outlined in Tables A1 and A2 effectively captured the

general trends in mean total development time, CoV development time, and proportion eclosion,

we considered whether alternative thermal sensitivity patterns could improve model fit. A small

proportion of CD *R. pomonella* can eclose without chilling, although this typically takes more

than 150 d (Dambroski and Feder 2007). We therefore considered whether CD pupae could be

modelled with a rate of pre-threshold diapause development at 21°C that was greater than zero

($r_e > 0$). When we calculated simulation summary metrics for our $r_e > 0$ model, the fit to

empirical data was very poor (Fig. S9A vs. B). However, in our empirical data experiment, we

did not track eclosion indefinitely, so we did not count any pupae that might have eclosed after a

prolonged time (e.g., > 100 days) post-chill. If we incorporated a 'cutoff' of 115 days into our

$r_e > 0$ model – i.e. if we did not 'count' any simulated eclosion that occurred after 115 days post-

chill – the fit of this model (Fig. S9C) was similar but not a dramatic improvement relative to our

base model ($r_e = 0$; Fig. S9A).

1343