**Title: Sleep and memory consolidation are linked by RNA processing genes in the *Drosophila* mushroom body**

Yongjun Li[1, 2, *], Nitin S. Chouhan[1, 4 *], Shirley Zhang[1, 5], Rebecca S. Moore[1], Sara B. Noya[1], Joy Shon[1], Zhifeng Yue[1], Amita Sehgal[1]

[1]Howard Hughes Medical Institute and Chronobiology and Sleep Institute, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA 19104, United States

[2]Department of Biology, University of Pennsylvania, PA 19104, United States

[3]Department of Pharmacology, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA 19104, United States

[4]Current address: Department of Biological Sciences, Tata Institute of Fundamental Research, Mumbai 400005, India

[5]Current address: Department of Cell Biology, Emory University School of Medicine, Atlanta GA 3032

*Equal contribution

Correspondence should be addressed to Amita Sehgal at amita@pennmedicine.upenn.edu

**Keywords**

Sleep, memory consolidation, translation, ribosomal RNA, mushroom body

**Abstract**

Memory consolidation in *Drosophila* can be sleep-dependent or sleep-independent, depending on the availability of food. Different regions of the mushroom body (MB) mediate these two mechanisms, with the anterior posterior (ap) alpha′/beta′ (α′/β′) neurons required for sleep-dependent memory consolidation in flies that are fed after training. These neurons are also involved in the increase of sleep after training, suggesting a coupling of sleep and memory. To better understand the mechanisms underlying sleep and memory consolidation initiation, we analyzed the transcriptome of ap α′/β′ neurons one hour after appetitive memory conditioning. A small number of genes were differentially expressed specifically in flies fed after training, but not in trained and starved flies or untrained flies. Knockdown of each of these differentially expressed genes in the ap α′/β′ neurons revealed multiple genes that affect sleep, with notable effects observed for Polr1F and Regnase-1, both of which decrease in expression after conditioning. Knockdown of Polr1F, a regulator of ribosome RNA transcription, in adult flies promotes sleep and increases pre-ribosome RNA expression as well as overall translation, supporting a function for Polr1F downregulation in memory consolidation. Conversely, knockdown of Regnase-1, an mRNA decay protein localized to the ribosome, reduces sleep. Given that Regnase-1 knockdown in ap α′/β′ neurons affects both sleep-dependent and sleep-independent memory, as well as short-term memory, Regnase-1 likely has an early role in the learning process, which may

obscure a later function for its downregulation during sleep-dependent memory. These

findings indicate that changes in RNA processing play a crucial role in triggering post-

training sleep and memory consolidation.

**Background**

Sleep is an optimized state for memory consolidation compared to wake (Rasch

and Born, 2013). Indeed, sleep is thought to reorganize and strengthen the neural

connections required for novel memory formation and long-term memory consolidation,

and sleep disruption leads to impaired memory consolidation (Roselli et al., 2021).

Beneficial effects of sleep on memory have also been attributed to post-learning

neuronal reactivation (Wagner et al., 2007; Dag et al., 2019).

We recently demonstrated that *Drosophila* switch between sleep-dependent and

sleep-independent memory consolidation based on food availability (Chouhan et al.,

2021). Flies that are fed after appetitive conditioning show sleep-dependent memory

consolidation, which is mediated by the anterior posterior (ap) α'/β' neurons of the

mushroom body. On the other hand, flies starved after training display sleep-

independent memory mediated by medial (m) neurons of the α'/β' lobes.

Neurotransmission from ap α'/β' neurons in the first 4 hours after training in the fed flies

is not only required for long term memory but also increased sleep post-training. Indeed,

activation of ap α'/β' neurons promotes baseline sleep, supporting the idea that post

training sleep is triggered by the same neurons that are required for sleep-dependent

memory consolidation. However, how sleep and memory consolidation are coordinated

in ap α′/β′ neurons in this time window is not understood.

Since both post-training sleep and memory consolidation occur rapidly within a

few hours after training, we investigated transient gene expression changes in the ap

α′/β′ neurons after training to elucidate the interplay of sleep and memory consolidation.

Here, we profiled the transcriptome of ap α′/β′ neurons 1 hour after flies were fed and

trained, as well as under control conditions in which flies were starved and trained or fed

and untrained. We knocked down the differentially expressed genes in trained and fed

flies and identified two RNA processing genes that affect sleep. Knockdown of one of

these, Polr1F, a regulator of ribosomal RNA synthesis, promotes sleep and translation,

both of which are required for consolidation of memory in trained and fed flies.

Knockdown of Polr1F does not affect memory, which is consistent with downregulation

of this gene during memory consolidation. Knockdown of Regnase-1, an mRNA decay

protein, reduces sleep and disrupts short- and long-term memory, suggesting an early

role, perhaps in learning. These results suggest that RNA processing is an important

mechanism linking sleep to memory.

## Results

### Transient transcriptome profiling of ap α′/β′ neurons after training

We previously demonstrated that flies fed after appetitive memory training exhibit increased sleep and form sleep-dependent memory, which is mediated by ap α′/β′ neurons of the mushroom body (MB) α′/β' lobes (Chouhan et al., 2021). To address the mechanisms that mediate increases in sleep and consolidation of memory in ap α′/β′ neurons, we assayed gene expression changes in ap α′/β′ neurons of flies fed after training in an appetitive conditioning paradigm. We crossed ap α′/β′ neuron driver *R35B12-Gal4* (BDSC #49822) with *UAS-nGFP* (BDSC #4775) to label all ap α′/β′ neurons with nuclear GFP and collected 5-7 day old F1 progeny, which were subjected to one of three different conditions: Trained-Fed, Trained-Starved, Untrained-Fed as illustrated (**Figure 1A**). The trained-starved flies served as controls for sleep-dependent changes, while flies that were fed but untrained served as controls for training-dependent changes. After one-hour, 50 mixed sex (25 for each sex) fly brains from each condition were dissected, and 500 GFP+ cells were sorted for bulk-RNA sequencing using the protocol described by Hongjie Li et al (Li et al., 2017).

Our analysis of the bulk RNA-seq data revealed that most genes are not altered in ap α′/β′ neurons after one hour, so there are no significant global changes observed among the three conditions. The correlation matrix calculated using the top 75% genes showed high similarity between samples, with most values exceeding 0.9 (**Figure 1 – source data**). Nonetheless, we did observe a small subset of genes that were rapidly responsive and differentially expressed between the groups (**Figure 1B**). Pathway Principal Component Analysis (PathwayPCA) of these differentially expressed genes

(DEGs), a statistical method to identify key patterns in gene expression data by

reducing data dimensionality, showed that genes contributing to PC1, which accounted

for 35% of the variance, largely encode proteins involved in transcription and

biosynthesis, including RNA biosynthesis processes (**Figure 1 – supplemental figure**

**1A**) (Odom et al., 2020). This suggests that the training and feeding paradigm

influenced transcription and RNA biosynthetic processes. Meanwhile, gene ontology

analysis of these DEGs, performed using DAVID Bioinformatics

([https://david.ncifcrf.gov/home.jsp](https://david.ncifcrf.gov/home.jsp)), revealed significant enrichment in processes such

as protein unfolding/refolding and the stress response (**Figure 1 – supplemental figure**

**1B**) (Sherman et al., 2022).

Of all the DEGs across the three groups, we focused on 59 DEGs that were

significantly different in TrainedStarved vs. TrainedFed and UntrainedFed vs.

TrainedFed; of these, 56 were downregulated and only three were upregulated in the

Trained-Fed condition compared to the control conditions (**Figure 1B, D, Figure 1 –**

**source data**). Gene ontology (GO) analysis of these 59 genes using FlyEnrichr

(https://maayanlab.cloud/FlyEnrichr/) indicated that they encode cellular components of

the 90S preribosome, Cajal body, DNA-directed RNA polymerase complex, nuclear

euchromatin, and condensed chromosome, consistent with the PCA enrichment (**Figure**

**1C**) (Chen et al., 2013). Our transcriptome results suggest that ribosome biosynthesis

and transcription are the initial changes in ap α′/β′ neurons of trained and fed flies.

**Two genes expressed differentially in ap α′/β′ neurons of trained and fed flies**

**predominately affect sleep.**

To investigate if any of the 59 DEGs identified in the ap α′/β′ neurons of trained

and fed flies affect baseline sleep, we knocked down each of the genes using UAS-

RNAi lines and screened for their potential effects on sleep. We used the ap α′/β′

neuron constitutive driver R35B12-Gal4 line to drive the RNAi constructs and compared

the sleep patterns of knockdown flies with those of R35B12-Gal4 and UAS-RNAi control

flies (**Figure 2A, B, Figure 2 – source data**). Using a cutoff of a 200 min change in

sleep relative to each control group, we identified two genes, Polr1F and Regnase-1,

that showed significant effects on baseline sleep. Knockdown of Polr1F, a component of

RNA polymerase I complex, led to an increase in sleep. Although Polr1F has not been

extensively studied in *Drosophila* (Marygold et al., 2020), its human ortholog hRPA43 is

part of the multi-subunit protein complex Pol I that regulates the transcription of

ribosomal RNA (Beckouët et al., 2011). Knockdown of Regnase-1, an RNA-binding

protein that binds to mRNA undergoing active translation and promotes mRNA decay

via its ribonuclease activity, in ap α′/β′ neurons resulted in a reduction of both nighttime

and daytime sleep (**Figure 2C-F**). Further analysis of sleep architecture revealed that

knockdown of Polr1F and Regnase-1 did not significantly impact the total activity of

flies, while knockdown of either Polr1F and Regnase-1 resulted in increased and

decreased average length of sleep episodes, respectively (**Figure 2 – supplemental**

**figure1A-F**).

Given that these two genes reduce expression rapidly in response to training

under fed conditions, we next used the inducible pan-neuronal GeneSwitch driver *nSyb-*

*GeneSwitch (GS)* to determine if restricting knockdown of these two genes to the adult

stage recapitulates changes in sleep and/or memory. RU486 (mifepristone) was added

to normal fly food and transgene expression is induced when flies are loaded into

*Drosophila* Activity Monitor (DAM) glass tubes (Robles-Murguia et al., 2019). We

crossed the *nSyb-GS* flies with Polr1F or Regnase-1 RNAi flies and transferred the

adult F1 progeny to RU486 tubes to induce expression of the RNAi 3-5 hours before

dusk, and continuously monitored sleep for 5 days. Knockdown of Polr1F resulted in

immediate inactivity and sleep (**Figure 3A-C**). However, with knockdown of Regnase-1,

immediate changes in sleep were not noted (**Figure 3 – supplemental figure 2A-C**).

This could be due to potential leakiness of the nSyb-GS, such that it reduced sleep

even without RU486 treatment, coupled with the fact that sleep before dusk is already

quite low, potentially hindering detection of further sleep reduction. We also analyzed

sleep for the next 3 consecutive days from day 3 to day 5 and found that constitutive

pan-neuronal Polr1F knockdown increased sleep while Regnase-1 knockdown seemed

to have no effect, again perhaps due to the leakiness of *nSyb-GS.* However, it is also

possible that Regnase-1 affects sleep during developmental stages or acts differently in

different brain areas (**Figure 3D, E; Figure 3 – supplemental figure 2D, E**). In general,

these results indicate that adult specific knockdown of Polr1F promotes sleep while

brain-wide adult-specific knockdown of Regnase-1 has limited effect on sleep.

To further assess the adult-specific sleep function of both genes in ap $\alpha'/\beta'$

neurons, we coupled the R35B12-Gal4 driver to tubulin-Gal80$^{ts}$, which allows

temperature-dependent expression of Gal4, and crossed it with the UAS-Polr1F$^{RNAi}$ and

UAS-Regnase-1$^{RNAi}$ lines. We confirmed that Polr1F RNAi promotes both transient and

chronic sleep when flies were transferred from permissive to restrictive temperatures

during the adult stages (**Figure 3 – supplemental figure 1**). Conversely, Regnase-1

RNAi showed no effect on sleep in the adult stage, consistent with our nSyb-GS

experiments, suggesting that the Regnase-1 RNAi sleep effect is likley developmental

(**Figure 3 – supplemental figure 3**). We also analyzed the daily peak activity of the

flies to determine if either Polr1F RNAi or Regnase-1 RNAi affected activity. We found

that neither gene affected peak activity, suggesting that both genes influence sleep but

not activity (**Figure 3 -supplemental figure4**).

**Knockdown of Regnase-1 affects memory**

We next evaluated the impact of Polr1F and Regnase-1 knockdown on memory

consolidation using our olfactory conditioning paradigm (**Figure 4A**). Starved flies were

subjected to training to associate an odor with a reward, and then post-training, they

were either kept on food vials for sleep-dependent memory consolidation or kept

starved to promote sleep-independent memory consolidation. Memory tests were

conducted 24 hours after training for starved flies, while fed flies were restarved for 42

hours before testing, as starvation is necessary for memory retrieval (Krashes and

Waddell, 2008). Efficacy of the RNAi line used above was verified by qPCR

experiments (**Figure 4 -supplemental figure 1A**). We observed that constitutive

knockdown of Polr1F in ap α′/β′ neurons did not affect sleep-dependent or sleep-

independent memory as memory performance was comparable to that of genetic

controls (**Figure 4B-C**). These results were consistent with the fact that Polr1F levels

typically decrease during memory consolidation. Monitoring of sleep from ZT8 to ZT12

after training at zeitgeber time (ZT) 6 showed that the post-training increase in sleep

was also not affected by Polr1F knockdown in ap α′/β′ neurons (**Figure 4E**), suggesting

that acute downregulation of Polr1F may not be essential to the post-training increased

sleep.

On the other hand, constitutive Regnase-1 knockdown in ap α′/β′ neurons

resulted in a significant decrease in long-term memory performance in both fed and

starved flies and eliminated the increase in post-training sleep (**Figure 4B-E**). These

findings suggested that Regnase-1 expression in ap α′/β′ neurons is necessary for both

sleep-dependent and sleep-independent memory consolidation. It is also possible that

disruption of Regnase-1 in ap α′/β′ neurons affects learning, short-term memory

formation, or appetitive memory retrieval (Aso et al., 2014; Shyu et al., 2019; Rutherford

et al., 2023). Indeed, short-term memory tests confirmed that Regnase-1 knockdown

flies performed significantly worse than control flies (**Figure 4F**), indicating that

Regnase-1 acts early in the learning/memory process, which might explain its effect on

sleep-dependent and sleep-independent memory. We also tested Regnase-1

overexpression for effects on sleep, and found that it does not affect sleep (**Figure 4 -**

**supplemental figure 1B**) (Zhu et al., 2019).

**Knockdown of Polr1F promotes translation**

Since Polr1F knockdown promotes sleep and a 22 amino acid peptide within

Polr1F inhibits ribosomal DNA transcription (Rothblum et al., 2014), we predicted that

Polr1F acted as a suppressor for ribosomal DNA transcription, and thus knockdown of

Polr1F would enhance the transcription and translation of ribosomal RNA and thereby

overall protein synthesis. As Regnase-1 is thought to promote decay of mRNAs

undergoing translation, its knockdown, or its downregulation following training, might

also be expected to promote translation, or at least the translation of its target mRNAs.

The role of protein synthesis in long-term memory consolidation is well-established

across organisms (Alberini and Kandel, 2015), and nascent rRNA synthesis was also

shown to be induced by training and required for memory consolidation in mouse (Allen

et al., 2018). We thus used real-time qPCR of total RNA extracted from vehicle control

and RU-treated fly brains to measure how precursor ribosomal RNA (Pre-rRNA) is

affected by knockdown of Polr1F. We found that the pre-rRNA level increased

significantly in the RU486 induction (RU+) group compared with vehicle control (RU-)

group for *nSyb-GS>polr1F RNAi* flies (**Figure 5A**), indicating Polr1F knockdown results

in higher levels of pre-rRNA, which is consistent with studies of Polr1F homologs in

yeast cells (Thuriaux et al., 1995; Rothblum et al., 2014).

The increasing ribosomal RNA should help translation and so we also used

incorporation of puromycin into newly synthesized peptides as an estimate of translation

after inducing pan-neuronal knockdown of Polr1F. We observed significantly higher

levels of anti-puromycin GFP fluorescence when Porl1F was knockdown pan-neuronally

(RU+) compared to the control (RU-) group (**Figure 5B**), indicating that Polr1F

suppresses translation, probably by suppressing ribosomal RNA transcription

(Rothblum et al., 2014). Thus, it may need to be downregulated after training to support

translation and memory. Given that knockdown of Porl1F enhances rRNA synthesis and

sleep, there is a question as to whether global alterations in rRNA synthesis impact

sleep. However, feeding of an rRNA inhibitor (CX-5461) failed to produce rapid sleep

phenotypes in fed or starved flies (**Figure 5 – supplemental figure 1**), suggesting that

rRNA synthesis may not directly influence sleep. Using puromycin to address effects of

Regnase-1 knockdown on translation revealed an insignificant slight increase in

translation (data not shown); given that Regnase-1 may specifically affect pre-existing

translationally active mRNA or may act only on specific target mRNAs, its effects on de

novo translation may not be obvious.

**Discussion**

The anterior-posterior (ap) α′/β′ neurons of the mushroom body make critical and

privileged contributions to sleep-dependent memory consolidation and post-training

sleep (Krashes et al., 2007; Chouhan et al., 2021), but the mechanisms that link

memory and sleep in these neurons are not known. To address this gap, we conducted

transcriptomic analysis of ap α′/β′ neurons from trained and fed flies to identify genes

that change rapidly under conditions that drive sleep-dependent memory. By uncovering

two RNA processing genes involved in memory and sleep, our transcriptome profiling of

ap α′/β′ neurons suggests that genes regulating rRNA transcription and translation are

altered in the context of sleep-dependent memory consolidation.

**RNA processing genes mediate sleep-dependent memory consolidation and**

**sleep.**

Many of the 59 DEGs we identified are implicated in RNA processing. Of these,

Polr1F and CG11920 affect ribosomal RNA processing, CG5654 is predicted to be part

of the 90S pre-ribosome and involved in endonucleoytic cleavages (Herold et al., 2009),

WDR79 encodes a small Cajal body specific RNA binding protein, Nup133 encodes a

component of nuclear pore complex, Regnase-1 degrades mRNA, and CG18011,

CG17568, Koko, CG11398 and Meics are all involved in RNA polymerase II-specific

transcription (Di Giorgio et al., 2017; Rogg et al., 2022). The enrichment of RNA

processing and translation genes is consistent with the notion that memory

consolidation requires transcription and translation (Seibt and Frank, 2012; Alberini and

Kandel, 2015). More importantly, our findings here indicate that alterations of some of

RNA processing genes may impact sleep as well.

**Polr1F regulates ribosome RNA synthesis and sleep**

Learning-induced changes in gene expression in memory-related neurons are

often critical for long-term memory formation (Cavallaro et al., 2002; Hoedjes et al.,

2015; Tadi et al., 2015). Our findings with Polr1F implicate changes in Pol I transcription

during sleep-dependent memory. Polr1F(Rpa43) is predicted to be part of the RNA

polymerase I complex and is involved in DNA-dependent RNA polymerase

activity/rDNA transcription in yeast, especially for the initiation of ribosomal RNA

(Beckouët et al., 2011; Marygold et al., 2020). As suggested by our data, Polr1F has an

inhibitory role in the RNA polymerase I complex, which is also consistent with the

aforementioned study that found a 22 amino acid peptide within Polr1F can inhibit rDNA

transcription (Rothblum et al., 2014). Based upon our finding that inducible knockdown

of Polr1F rapidly promotes translation, it is likely that the rapid and dramatic decline of

Polr1F after training in fed flies serves to increase *de novo* ribosome RNA synthesis.

This supports the report that ribosomal RNA is induced by learning and required for

memory consolidation in mice (Allen et al., 2018). While we do not know how

knockdown of Polr1F promotes sleep, an attractive possibility is that higher translation is

a result of elevated sleep. Sleep is thought to promote translation and its role in memory

consolidation in fed flies could be related to its effect on translation (Seibt and Frank,

2012; Chouhan et al., 2021). Alternatively, increased translation or rRNA synthesis

could promote sleep. However, translation is typically thought of as a consequence of

sleep rather than a cause (Zimmerman et al., 2006). Also, rRNA transcription rates

remain constant  throughout the day in the liver (Sinturel et al., 2017), although it is still

possible that these rates vary in particular regions of the brain and affect sleep. The role

of rRNA synthesis in *Drosophila* learning and memory has barely been explored, but our

work, together with that of Allen et al. (2018), indicates that the well-known requirement

for *de novo* protein synthesis during long-term memory consolidation (Jarome and

Helmstetter, 2014) includes increased synthesis of ribosomal RNA and protein.

**Regnase-1 inactivation may affect memory through effects on translation**

The role of RNA binding protein Regnase-1 in the innate immune response has

been extensively studied (Mino et al., 2015; Mao et al., 2017; Wei et al., 2019).

However, our study sheds light on a novel function of Regase-1 in ap α′/β′ neurons on

sleep and memory. Regnase-1 is an anti-inflammatory enzyme that inhibits mRNA

translation during acute inflammatory responses. It localizes to the ribosomes on the

surface of the endoplasmic reticulum (ER) and binds to translationally active mRNAs

with specialized stem-loop structures at the 3′UTR (Uehata et al., 2013; Mino et al.,

2015). When phosphorylated, Regnase-1 is released from the ER (Tanaka et al., 2019).

Because functional Regnase-1 binds and degrades its bound mRNA, Regnase-1

inactivation leads to an increase of its target mRNA (Uehata et al., 2013). The target

mRNAs of Regnase-1 in immune cells encode proinflammatory cytokines, which can

then be expressed when Regnase-1 is inactivated. However, Regnase-1 has also been

reported to modulate cytokines and neuronal injury in the microglia in rats (Liu et al.,

2016). Regulation of Regnase-1 is usually rapid and transient, and its rapid response to

microenvironmental changes, different pathological states and stress is critical for

cellular adaption (Mao et al., 2017).

Our study reveals that the expression of Regnase-1 changes after training, and

constitutive downregulation of Regnase-1 in ap α′/β′ neurons reduce sleep and causes

deficits in learning and memory consolidation. We suggest that downregulation of

Regnase-1 following training in fed flies may be crucial for the promotion of long-term

memory, perhaps by promoting translation of specific transcripts. For instance,

downregulation of Regnase-1 after training may release a pool of mRNAs that are

normally targeted by it for decay and that are important for consolidation of memory. On

the other hand, we find that constitutive loss of Regnase-1 impairs sleep-independent

memory and also short-term memory, suggesting that it is required early in the learning

and memory process, likely for learning. Interestingly, constitutive knockdown of

Regnase-1 also reduces sleep and prevents sleep increase after training. However, our

adult specific knockdown of Regnase-1 does not affect sleep, suggesting that

expression of Regnase-1 during development is important for both memory and sleep.

Given that ap α′/β′ neurons have a role in several aspects of memory, consolidation as

well as short-term appetitive memory retrieval, how Regnase-1 acts within these

neurons to regulate memory clearly requires more investigation.

Our study of local molecular changes in ap α′/β′ neurons after training highlights

how ribosomal RNA transcription and mRNA translation work in concert during the

consolidation of sleep-dependent memory. Neurons are regulated to generate short

bursts of boosted transcription and boosted translation, which are required for new

protein synthesis. How sleep is involved in this boosted protein synthesis process is

unclear, but we suggest that sleep affects several steps of the central dogma, and that

deleterious consequences of sleep deprivation are mediated, at least in part, through

effects on rRNA transcription and translation. Overall, our findings indicate the

importance of maintaining RNA homeostasis for sleep and memory.

**Figures**

**Figure1. Differential gene expression after training in mushroom ap α′/β′ neurons.**

(A) 5-7 day old mixed sex *wCS* flies were exposed to one of the following three

conditions: Trained-Fed, Trained-Starved and Untrained-Fed. Only Trained-Fed flies are

expected to increase sleep after treatment and thus form sleep-dependent memory

(Chouhan et al., 2021). Brain dissection, single cell suspension and cell sorting were

used to extract ap α′/β′ neurons in each of these three different conditions, and bulk-

sequencing of the sorted cells was conducted. (B) We sequenced four samples for each

condition and subjected them to differential gene analysis of pairwise comparison of the

three conditions. We found that 59 genes are significantly different in TrainedStarved

vs. TrainedFed and UntrainedFed vs. TrainedFed. (C) Gene ontology (GO) analysis of

these 59 genes using FlyEnrichr (https://maayanlab.cloud/FlyEnrichr/) revealed that

they encode cellular components of the 90S preribosome, Cajal body, DNA-directed

RNA polymerase complex, nuclear euchromatin, and condensed chromosome.(D)

Heatmap of the 59 DEGs including two genes labeled in red, CG13773 (Polr1F) and

Regnase-1, which affect sleep and are the focus of this study due to their impacts on

sleep. DEGs were identified by DESeq2 with the cutoff of FDR< 0.1 and fold

change >1.5. Heatmaps were plotted by using TPM values of genes for each sample;

data were log-transformed and scaled row-wise for visualization.

**Figure 2. The sleep screen of differentially expressed genes identifies Polr1F and Regnase-1 as sleep-regulating genes.** (A-B) Flies carrying the ap α′/β′ neuron driver R35B12-Gal4 were crossed with flies carrying UAS-RNAi constructs targeting DEGs identified from RNA-seq analysis. 5-7 days old female F1 progeny were loaded onto Trikinetics DAM monitors to measure their sleep in a 12-hour light: 12-hour dark (12:12

LD) cycle. Mean total sleep (A) and nighttime sleep (B) were calculated by Pysolo and

the difference between experimental flies and Gal4 and RNAi controls was calculated

separately for each independent experiment; average values comparing each

experimental to its Gal4 control (X-axis) and RNAi control (Y-axis) are shown in the

plots. Of all the lines screened, knockdown of Polr1F and Regnase-1 had strongest

effects on sleep, producing an increase and decrease in sleep respectively. (C-F) show

the representative sleep traces of *R35B12-Gal4>polr1F$^{RNAi}$* flies and *R35B12-*

*Gal4>regnase1$^{RNAi}$*. N = 23-32 per genotype from two independent replicates combined

are shown in E and F respectively, and bar graphs show mean + SEM. p values for

each comparison were calculated using the Kruskal-Wallis test with Dunn's multiple

comparisons test. **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

**Figure 3. Acute and chronic effects of pan-neuronal knockdown of Polr1F on sleep in adult flies.** (A) Schematic representation of transient and chronic sleep measurements in *nSyb-GS>polr1f^RNAi* flies. (B) Representative sleep traces and transient activity plot of flies expressing Polr1F RNAi under the control of an inducible pan-neuronal driver (*nSyb-GS>polr1f RNAi*) with and without RU treatment. (C) Quantification of sleep during the first 3 hours (ZT 9-12) after F1 progeny flies were loaded into RU- or RU+ DAM tubes at ZT8-T9. Sleep was measured starting at ZT9. N

= 39-40 individual flies per replicate with data from three independent replicates

combined. The Mann-Whitney test was used to compare RU+ group and RU- groups.

(D) Representative average sleep traces of *nSyb-GS>polr1f RNAi* in the RU- and RU+

DAM tubes for three consecutive days. Chronic sleep effects of pan-neuronal

knockdown Polr1F were measured based on sleep data from day 3 to day 5. (E)

Quantification of average total sleep of *nSyb-GS>polr1f^{RNAi}* and controls in the DAM

tubes from (D). Unpaired t-test was used to compare between RU- and RU+ groups.

****$p<0.0001$.

**Figure 4. Regnase-1 expression is essential for sleep-dependent and sleep-independent memory.** (A) Schematic representation of the memory test protocol. (B, C) Sleep-dependent and sleep-independent memory tests were conducted under fed and starved conditions, respectively. Knockdown of Regnase-1 significantly reduces long-term memory performance in both fed and starved flies. However, knockdown of Polr1F in ap α′/β′ neurons does not affect long-term memory performance. N≥6 biological replicates, each replicate containing 100-150 flies. (D, E) Fed UAS-regnase-1-RNAi/+ and R35B12/+ flies exhibit a significant increase in sleep after training, while *R35B12-Gal4>regnase-1 RNAi* flies fail to show a comparable increase in post-training sleep. The total sleep in the ZT8-ZT12 interval is shown in (E). Polr1F knockdown in ap α′/β′ neurons does not affect the post-training increase in sleep. N≥32. (F) Compared to R35B12-Gal4/+ and +/UAS-Regnase-1 RNAi flies, *R35B12-Gal4>reganse-1 RNAi* flies show a significant decrease in the performance index in short-term memory. N≥6 biological replicates, each containing 100-150 flies. ns=not significant, $p > 0.05$, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

**Figure 5. Knockdown of Polr1F results in high translation.** (A) The *nSyb-GS>polr1f*

*RNAi* flies exhibit a significant increase in pre-rRNA levels. (B) The ex-vivo puromycin

immunostaining assay was used to measure translation in dissected whole brains. The

results show that knockdown of Polr1F using the pan-neuronal nSyb-GeneSwitch (GS)

system increases translation relative to control flies that were not treated with RU. The

normalized mean grayscales from the RU- and RU+ groups are compared using an

unpaired t-test. The analysis includes data from 8-11 flies per group, with results from

two independent replicates combined. Scale bar: 100 μm. (C) The schematic model

illustrates the roles of Polr1F and Regnase-1 in memory consolidation. The genes

Polr1F and Regnase-1 are prominently downregulated during memory consolidation in

trained and fed flies, respectively. Polr1F is involved in regulating ribosomal RNA

synthesis, and its decrease in levels in trained and fed flies promotes sleep and

translation. In contrast, Regnase-1 is involved in mRNA decay, and its downregulation

during memory consolidation may contribute to the stabilization of mRNAs that encode

proteins important for long-term memory formation.

**Figure 1 – supplemental figure 1.** (A) PathwayPCA plot of all the differentially

expressed genes (DEGs) across the three different conditions. The plot shows that

genes responsible for PC1, which accounted for 35% of variance, largely encode

proteins involved in transcription and biosynthesis, including RNA biosynthesis

processes. (B) Top5 significant Gene Ontology (GO) biological process terms of all

differentially expressed genes (DEGs) from three pairwise comparisons are shown. The

x-axis represents the -log10 of the P-value, and the numbers on the bars indicate the

count of genes identified for each GO term.

**Figure 2 – supplemental figure 1. Effect of Polr1F and Regnase-1 knockdown on activity and sleep architecture.** (A-B) Total activity of flies is not altered by knockdown of Polr1F and Regnase-1. One-way ANOVA was used to calculate the p-values for each comparison. (C-D) Knockdown of Polr1F significantly increases the nighttime average sleep episode (SE) length in R35B12-Gal4>polr1F RNAi flies, while knockdown of

Regnase-1 reduces it in R35B12-Gal4>regnase1 RNAi flies. The p-values for each

comparison were calculated using the Kruskal–Wallis test with Dunn′s multiple

comparisons test. (E-F) Knockdown of Polr1F significantly reduces the nighttime

average sleep episode (SE) number in R35B12-Gal4>polr1FRNAi flies, but it is not

significant compared to control groups in R35B12-Gal4>regnase1RNAi flies. The p-

values for each comparison were calculated using the Kruskal–Wallis test with Dunn′s

multiple comparisons test. ns = not significant, $p > 0.05$, *$p < 0.05$, **$p < 0.01$, ***$p <$

$0.001$, ****$p < 0.0001$.

**Figure 3 – supplemental figure 1. Adult specific knockdown of Polr1F promotes sleep.** (A) Representative sleep traces of adult flies expressing Polr1F RNAi under the control of R35B12-Gal4/Tubulin-gal80ts. At permissive temperature, Gal80ts inhibits Gal4 activation of UAS; however, upon switching to restrictive temperature, Gal80ts is

deactivated, leading to downstream activation of the UAS-regulated gene. (B) Sleep

patterns were quantified over four consecutive days, with flies under permissive

temperature on the first and last days and restrictive temperature on the second and

third days. (C) Sleep quantification during the initial 3-hour period (ZT 0-3) after light

onset for each day is shown. N = 46-48 individual flies per replicate with data from three

independent replicates combined. The Kruskal-Wallis test with Dunn's multiple

comparisons test was used to compare three groups. ns = not significant, $p > 0.05$, *$p <$

$0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

**Figure 3 – supplemental figure 2. Acute and chronic effects of pan-neuronal knockdown of Regnase-1 on sleep in adult flies.** (A) Schematic representation of the protocol for transient and chronic sleep measurements in *nSyb-GS>regnase-1 RNAi* flies. (B) Representative sleep traces plot of flies expressing Rengase-1 RNAi under the control of an inducible pan-neuronal driver (*nSyb-GS>regnase-1 RNAi*) with and without RU treatment (C) Quantification of sleep during the first 3 hours (ZT 9-12) after F1

progeny flies were loaded into RU- or RU+ DAM tubes at ZT8-T9. Sleep was measured

starting at ZT9. N = 22-24 individual flies per replicate with data from two independent

replicates combined. The Mann-Whitney test was used to compare RU+ group and RU-

groups. (D) Representative average sleep traces of *nSyb-GS>regnase-1 RNAi* in the

RU- and RU+ DAM tubes for three consecutive days. Chronic sleep effects of

knockdown of Regnase-1 were measured based on sleep data from day 3 to day 5. (E)

Quantification of average total sleep of *nSyb-GS>regnase-1 RNAi* and controls in the

DAM tubes from (D). Unpaired t-test was used to compare between RU- and RU+

group. ns = not significant.

**Figure 3 – supplemental figure 3. Adult specific knockdown of Regnase-1 has no effect on sleep.** (A) Representative sleep traces of adult flies expressing Regnase-1 RNAi under the control of R35B12-Gal4/Tubulin-gal80ts. (B) Sleep was quantified over four consecutive days, with flies under permissive temperature on the first and last days and restrictive temperature on the second and third days. N = 46 individual flies per replicate with data from three independent replicates combined. The Kruskal-Wallis test with Dunn's multiple comparisons test was used to compare three groups. ns = not significant, $p > 0.05$, *$p < 0.05$, **$p < 0.01$.

**Figure 3 – supplemental figure 4. Adult specific knockdown of Polr1F or Regnase-1 has no effect on peak activity.** (A) Maximal activity per minute for R35B12-Gal4; TubGal80ts>polr1F RNAi was quantified over four consecutive days, with flies under permissive temperature on the first and last days and restrictive temperature on the second and third days. (B) Maximal activity per minute for R35B12-Gal4; TubGal80ts>Regnase-1 RNAi was quantified. N = 40-48 individual flies per replicate with data from three independent replicates combined. The Kruskal-Wallis test with Dunn's multiple comparisons test was used to compare three groups. ns = not significant, p > 0.05, *****p < 0.0001.

**Figure 4 – supplemental figure 1. Regnase-1 overexpression in the ap α′/β′**

**neurons does not affect sleep.** (A) The efficacy of the RNAi used in the experiments

was verified by real-time quantitative PCR. The different RNAi lines were crossed with

*nSyb-GS*, and adult progeny were placed in either RU- or RU+ tubes overnight before

being dissected for qPCR. Welch's test was used for statistical analysis across all three

RNAi, with *p<0.05 indicating significance. (B) Representative sleep traces and

quantification of adult flies expressing Regnase-1 under the control of R35B12-Gal4.

Sleep data were averaged over three consecutive days, with N = 31-32 individual flies

per replicate, and combined data from two independent replicates were analyzed.

**Figure 5 – supplemental figure 1. rRNA inhibitor (CX-5461) feeding does not affect**

**sleep.** Feeding of the ribosome RNA inhibitor CX-5461 at a concentration of 0.2mM

does not affect sleep in fed or starved flies. Statistical analysis shows that there is no

significant difference between CX-5461-fed and control flies. The sample size is N=32

per group from two independent replicates. ns, not significant).

**Reference:**

Alberini CM, Kandel ER (2015) The Regulation of Transcription in Memory Consolidation. Cold Spring Harbor Perspectives in Biology 7.

Allen KD, Regier MJ, Hsieh C, Tsokas P, Barnard M, Phatarpekar S, Wolk J, Sacktor TC, Fenton AA, Hernández AI (2018) Learning-induced ribosomal RNA is required for memory consolidation in mice—Evidence of differentially expressed rRNA variants in learning and memory. PLOS ONE 13:e0203374.

Aso Y et al. (2014) Mushroom body output neurons encode valence and guide memory-based action selection in Drosophila Luo L, ed. eLife 3:e04580.

Aviner R (2020) The science of puromycin: From studies of ribosome function to applications in biotechnology. Comput Struct Biotechnol J 18:1074–1083.

Beckouët F, Mariotte-Labarre S, Peyroche G, Nogi Y, Thuriaux P (2011) Rpa43 and its partners in the yeast RNA polymerase I transcription complex. FEBS Letters 585:3355–3359.

Cavallaro S, D'Agata V, Manickam P, Dufour F, Alkon DL (2002) Memory-specific temporal profiles of gene expression in the hippocampus. Proceedings of the National Academy of Sciences 99:16279–16284.

Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, Clark NR, Ma'ayan A (2013) Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics 14:128.

Chouhan NS, Griffith LC, Haynes P, Sehgal A (2021) Availability of food determines the need for sleep in memory consolidation. Nature 589:582–585.

Dag U, Lei Z, Le JQ, Wong A, Bushey D, Keleman K (2019) Neuronal reactivation during post- learning sleep consolidates long-term memory in drosophila. eLife 8.

Deliu LP, Ghosh A, Grewal SS (2017) Investigation of protein synthesis in Drosophila larvae using puromycin labelling. Biol Open 6:1229–1234.

Di Giorgio ML, Esposito A, Maccallini P, Micheli E, Bavasso F, Gallotta I, Vernì F, Feiguin F, Cacchione S, McCabe BD, Di Schiavi E, Raffa GD (2017) WDR79/TCAB1 plays a conserved role in the control of locomotion and ameliorates phenotypic defects in SMA models. Neurobiol Dis 105:42–50.

Diekelmann S, Born J (2010) The memory function of sleep. Nat Rev Neurosci 11:114–126.

Ge SX, Son EW, Yao R (2018) iDEP: an integrated web application for differential expression and pathway analysis of RNA-Seq data. BMC Bioinformatics 19:534.

Herold N, Will CL, Wolf E, Kastner B, Urlaub H, Lührmann R (2009) Conservation of the
Protein Composition and Electron Microscopy Structure of Drosophila
melanogaster and Human Spliceosomal Complexes. Mol Cell Biol 29:281–301.

Hoedjes KM, Smid HM, Schijlen EG, Vet LE, van Vugt JJ (2015) Learning-induced gene
expression in the heads of two Nasonia species that differ in long-term memory
formation. BMC Genomics 16:162.

Jarome TJ, Helmstetter FJ (2014) Protein degradation and protein synthesis in long-
term memory formation. Frontiers in Molecular Neuroscience 7 Available at:
https://www.frontiersin.org/articles/10.3389/fnmol.2014.00061 [Accessed
September 28, 2022].

Krashes MJ, Keene AC, Leung B, Armstrong JD, Waddell S (2007) Sequential use of
mushroom body neuron subsets during drosophila odor memory processing.
Neuron 53:103–115.

Krashes MJ, Waddell S (2008) Rapid Consolidation to a radish and Protein Synthesis-
Dependent Long-Term Memory after Single-Session Appetitive Olfactory
Conditioning in Drosophila. J Neurosci 28:3103–3113.

Li H, Horns F, Wu B, Xie Q, Li J, Li T, Luginbuhl DJ, Quake SR, Luo L (2017)
Classifying Drosophila Olfactory Projection Neuron Subtypes by Single-Cell RNA
Sequencing. Cell 171:1206-1220.e22.

Liu X-X, Wang C, Huang S-F, Chen Q, Hu Y-F, Zhou L, Gu Y (2016) Regnase-1 in
microglia negatively regulates high mobility group box 1-mediated inflammation
and neuronal injury. Sci Rep 6:24073.

Mao R, Yang R, Chen X, Harhaj EW, Wang X, Fan Y (2017) Regnase-1, a rapid
response ribonuclease regulating inflammation and stress responses. Cell Mol
Immunol 14:412–422.

Marygold SJ, Alic N, Gilmour DS, Grewal SS (2020) In silico identification of Drosophila
melanogaster genes encoding RNA polymerase subunits. MicroPubl Biol
2020:10.17912/micropub.biology.000320.

Mino T, Murakawa Y, Fukao A, Vandenbon A, Wessels H-H, Ori D, Uehata T, Tartey S,
Akira S, Suzuki Y, Vinuesa CG, Ohler U, Standley DM, Landthaler M, Fujiwara T,
Takeuchi O (2015) Regnase-1 and Roquin Regulate a Common Element in
Inflammatory mRNAs by Spatiotemporally Distinct Mechanisms. Cell 161:1058–
1073.

Odom GJ, Ban Y, Colaprico A, Liu L, Silva TC, Sun X, Pico AR, Zhang B, Wang L,
Chen X (2020) PathwayPCA: an R/Bioconductor Package for Pathway Based
Integrative Analysis of Multi-Omics Data. Proteomics 20:e1900409.

Rasch B, Born J (2013) About Sleep's Role in Memory. Physiological Reviews 93:681.

Robles-Murguia M, Hunt LC, Finkelstein D, Fan Y, Demontis F (2019) Tissue-specific
578           alteration of gene expression and function by RU486 and the GeneSwitch
579           system. npj Aging Mech Dis 5:1–5.

580    Rogg M, Maier JI, Ehle M, Sammarco A, Schilling O, Werner M, Schell C (2022)
581           NUP133 Controls Nuclear Pore Assembly, Transcriptome Composition, and
582           Cytoskeleton Regulation in Podocytes. Cells 11:1259.

583    Roselli C, Ramaswami M, Boto T, Cervantes-Sandoval I (2021) The Making of Long-
584           Lasting Memories: A Fruit Fly Perspective. Frontiers in Behavioral Neuroscience
585           15.

586    Rothblum K, Hu Q, Penrod Y, Rothblum LI (2014) Selective Inhibition of rDNA
587           Transcription by a Small-Molecule Peptide That Targets the Interface between
588           RNA Polymerase I and Rrn3. Molecular Cancer Research 12:1586–1596.

589    Rutherford S, Barkema P, Tso IF, Sripada C, Beckmann CF, Ruhe HG, Marquand AF
590           (2023) Evidence for embracing normative modeling Baker CI, Constable T,
591           Esteban O, eds. eLife 12:e85082.

592    Seibt J, Frank MG (2012) Translation regulation in sleep. Commun Integr Biol 5:491–
593           495.

594    Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, Imamichi T, Chang W
595           (2022) DAVID: a web server for functional enrichment analysis and functional
596           annotation of gene lists (2021 update). Nucleic Acids Res 50:W216–W221.

597    Shyu W-H, Lee W-P, Chiang M-H, Chang C-C, Fu T-F, Chiang H-C, Wu T, Wu C-L
598           (2019) Electrical synapses between mushroom body neurons are critical for
599           consolidated memory retrieval in Drosophila. PLoS Genet 15:e1008153.

600    Sinturel F, Gerber A, Mauvoisin D, Wang J, Gatfield D, Stubblefield JJ, Green CB,
601           Gachon F, Schibler U (2017) Diurnal Oscillations in Liver Mass and Cell Size
602           Accompany Ribosome Assembly Cycles. Cell 169:651-663.e14.

603    Tadi M, Allaman I, Lengacher S, Grenningloh G, Magistretti PJ (2015) Learning-Induced
604           Gene Expression in the Hippocampus Reveals a Role of Neuron -Astrocyte
605           Metabolic Coupling in Long Term Memory. PLoS One 10:e0141568.

606    Tanaka H, Arima Y, Kamimura D, Tanaka Y, Takahashi N, Uehata T, Maeda K, Satoh
607           T, Murakami M, Akira S (2019) Phosphorylation-dependent Regnase-1 release
608           from endoplasmic reticulum is critical in IL-17 response. Journal of Experimental
609           Medicine 216:1431–1449.

610    Thuriaux P, Mariotte S, Buhler J-M, Sentenac A, Vu L, Lee B-S, Nomura M (1995) Gene
611           RPA43 in Saccharomyces cerevisiae Encodes an Essential Subunit of RNA
612           Polymerase I (∗). Journal of Biological Chemistry 270:24252–24257.

Uehata T, Iwasaki H, Vandenbon A, Matsushita K, Hernandez-Cuellar E, Kuniyoshi K,
Satoh T, Mino T, Suzuki Y, Standley DM, Tsujimura T, Rakugi H, Isaka Y,
Takeuchi O, Akira S (2013) Malt1-Induced Cleavage of Regnase-1 in CD4+
Helper T Cells Regulates Immune Activation. Cell 153:1036–1049.

Wagner U, Kashyap N, Diekelmann S, Born J (2007) The impact of post-learning sleep
vs. wakefulness on recognition memory for faces with different facial
expressions. Neurobiology of learning and memory 87:679–687.

Wei J, Long L, Zheng W, Dhungana Y, Lim SA, Guy C, Wang Y, Wang Y-D, Qian C, Xu
B, Kc A, Saravia J, Huang H, Yu J, Doench JG, Geiger TL, Chi H (2019)
Targeting REGNASE-1 programs long-lived effector T cells for cancer therapy.
Nature 576:471–476.

Xu K, Zheng X, Sehgal A (2008) Regulation of Feeding and Metabolism by Neuronal
and Peripheral Clocks in Drosophila. Cell Metabolism 8:289–300.

Zhu L, Liao SE, Fukunaga R (2019) Drosophila Regnase-1 RNase is required for mRNA
and miRNA profile remodelling during larva-to-adult metamorphosis. RNA Biol
16:1386–1400.

Zimmerman JE, Rizzo W, Shockley KR, Raizen DM, Naidoo N, Mackiewicz M, Churchill
GA, Pack AI (2006) Multiple mechanisms limit the duration of wakefulness in
Drosophila brain. Physiological Genomics 27:337–350.

**Conflict of interest statement**

The authors declare no competing interests.

**Acknowledgments**

This work was supported by Howard Hughes Medical Institute (HHMI). We thank all members of the Sehgal lab for reagents, comments and support, especially rotation undergraduate Arielle Ketchum for help with molecular cloning. We thank Hongjie Li of Liqun Luo lab with technical assistance and protocol sharing from Stanford University. We thank Ryuya Fukunaga lab for kindly sharing with us the Regnase-1 overexpression line.

**Author contributions**

Conceptualization, Y.J.L., N.S.C. and A.S.; Methodology, Y.J.L., N.S.C., S.Z., S.B.N. and R.M. Software, Y.J.L; Validation, Y.J.L., N.S.C., S.Z., S.B.N., R.M., J.S., A.K. and Z.F.Y.; Formal analysis, Y.J.L., N.S.C. and R.M.; Investigation, Y.J.L. and N.S.C.; Resources, Y.J.L., N.S.C., S.Z., R.M., J.S., A.K. and Z.F.Y.; Data Curation, Y.J.L., N.S.C. and R.M.; Writing – Original Draft, Y.J.L. and N.S.C. Writing – Review & Editing, Y.J.L., N.S.C. and A.S. Visualization, Y.J.L., N.S.C. and R.M.; Supervision, Y.J.L., N.S.C. and A.S.; Project administration, A.S.; Funding acquisition, A.S.ss

**Materials and Methods**

**Contact for reagent and resource sharing.**

Amita Sehgal (amita@pennmedicine.upenn.edu)

**Key resource table**

See Appendix 1.

**Fly stock and maintenance**

All the stock information of the flies used in this project are listed in the key resource

table and flies were reared on the standard cornmeal vials or bottles at 25 °C with 12:12

hours light dark cycle in the preset incubator. The genetic background control used in

the paper is White-CantonS (wCS) unless specified.

**Behavior measurement in *Drosophila***

We have used both single beam and updated multibeam *Drosophila* activity monitoring

(DAM) system from Trikinetics (https://trikinetics.com/) in our experiments. Briefly, 5-7

days old female flies were loaded into 60/90 mm glass locomotor tubes for behavior

tests, using DAM2/5H *Drosophila* activity monitors from Trikinetics. 1/15 infrared beams

bisect each tube, providing movement (position in multibeam) information of the fly

across the tube. Locomotor tubes are loaded with 2% agar with 5% sucrose as fly food

on one side, and yarn is put on the other side to restrain the behavior of flies inside the

glass tubes. For experiments with the inducible Gene Switch system, 0.5mM RU486

(mifepristone) was added to the fly food to activate the expression of the transgenes

under the control of UAS. Three constitutive days of data were used for sleep analysis

by Pysolo (https://www.pysolo.net/).

**Appetitive conditioning**

~100 4-7 day old mixed-sexes flies were starved for 12 hours in *Drosophila* bottles with

water -soaked filtered paper and then trained at 25°C and 70% relative humidity to

associate sucrose with odor A for 2 minutes, and then a blank with odor B for 2 minutes

with 30-second clean air in between. The odors used were 4-Methylcyclohexanol (MCH)

and 3-Octanol (OCT) in paraffin oil and the concentration for both MCH and OCT in oil

was 1:10. After conditioning, flies were moved back to normal fly food or starved for 1

hour and dissected for subsequent ap cell sorting and RNA-sequencing. For the short-

term memory test, flies were tested immediately after conditioning in the same wheel for

2 mins.

To assess post-training sleep, flies were introduced in glass tubes containing 2% agar

and 5% sucrose through an aspirator without anesthesia and loaded into the DAM

system after training. For long-term memory assessment, trained flies were either kept

on food vials for 24 h or were further starved. Starved flies were tested for memory 24 h

after training, while fed flies were re-starved for 42 h before memory tests. Memory was

tested by giving flies a choice between odor A and odor B for 2 min in a T-maze.

Performance index (PI) was calculated as the number of flies selecting CS$^+$ odor minus

the number of flies selecting CS$^-$ odor divided by the total number of flies. Each PI is the

average of PIs from reciprocal experiments with two odors swapped to minimize non-

associative effects.

**Cell isolation and sorting**

Dissected brains are dissociated by following the protocol from (Li et al., 2017). Briefly,

brains are dissected in Schneider′s medium, and then are placed in a shaker and

dissociated in Papain solution, filtered through a 100 $\mu$m cell strainer, and re-suspended

in Schneider′s medium. 500 GFP+ cells from the same conditions were sorted into 96

well microplate with lysis buffer from Smart-seq2 HT kit and frozen. We dissected 50

brains for each group to ensure enough GFP+ cells. Cell sorting were conducted by

either BD FACSMelody or BD FACSAria (BD Biosciences), and dead cells were

excluded with 4′, 6-diamidino-2-phenylindole (DAPI). Doublets were excluded using and

forward scatter (FSC-H by FSC-W) and side scatter (SSC-H by SSC-W). Size of cells

was selected by FSC-A by FSC-A and validated for fly neurons using cells from flies

expressing nsyb-nGFP. Length of time from tissue harvest to cell collection

approximated 4 hours.

**RNA-seq and data analysis**

GFP+ cells were sorted and immediately frozen, then sent to Admera Health

(https://www.admerahealth.com/) for RNA extraction, RNA library construction, and

sequencing using the Smart-seq2 HT kit. To analyze the RNA-sequence data, we used

Hisat2 (http://daehwankimlab.github.io/hisat2/) to map the sequencing data FASTQ files

to the fly genome (BDSG6). The alignment results were then counted by LiBiNorm

(https://warwick.ac.uk/fac/sci/lifesci/research/libinorm/) using the GENCODE reference

genome. Raw count and TPM were used separately in further analysis. Raw count data

were analyzed by IDEP v0.95 to identify genes expressed differentially between three

conditions (Ge et al., 2018). We filtered out low-expressed genes using a cutoff of

CPM>0.5, at least detected in 3 independent samples, and treated missing values as

gene median. Regularized log transformation was used to transform raw count data for

clustering and PCA. Differentially expressed genes were identified using DESeq2 with

an FDR cutoff of 0.1 and minimum fold change of 2.

**Puromycin assay and imaging**

We developed a puromycin assay to measure the rate and localization of nascent

peptide synthesis in the fly brain, and similar method has been described in the fly

larvae (Deliu et al., 2017). Fly brains were dissected in Schneiders′ medium and

incubated with puromycin for 40 min *in vitro* to allow puromycin to incorporate into the

newly synthesized peptide. Subsequently, using an anti-puromycin antibody, standard

immunostaining protocols were applied to detect the number and position of newly

synthesized puromycin-tagged peptides (Aviner, 2020), which provided an estimate of

translation rate. The brains were then imaged with a 20X oil immersion confocal

microscope with a resolution of 1024*1024. The GFP intensity of the images was then

measured by Fiji software (ImageJ) and the average intensity of the samples was

analyzed for comparison.

**RNA polymerase I inhibitor II, CX-5461 protocol**

CX-5461 was mixed with 2% agar and 5% sucrose to make fly motor tubes at a final

concentration of 0.2 mM. Flies are loaded into these CX-5461 tubes or vehicle control

tubes 4 hours before light turns off and transient sleep changes are measured.

**Quantitative Real-time PCR (qPCR)**

10 flies′ brains from each group were dissected for RNA extraction using Qiagen's

RNeasy Plus Mini Kit. Total RNA was then reverse transcribed to cDNA by using

random hexamers and Superscript II from Invitrogen. qPCR was performed using

SYBR-green master mix and oligonucleotide information is provided in the Key

Resources table. Relative gene expression was analyzed using the ΔΔCt method.

**Statistical Analysis**

Fly sleep behavioral data extracted from Pysolo was analyzed by GraphPad Prism

(https://www.graphpad.com/). Data from different replicates were pooled directly and

first tested for normality using D'Agostino-Pearson and Shapiro-Wilk tests. For normally

distributed data, unpaired parametric Student's t-test is used for two-sample

experiments and one-way ANOVA with Turkey post hoc test for three-sample or more

experiments. Non-normally distributed data were analyzed using nonparametric tests,

the Mann-Whitney test for two-sample experiments and the Kruskal-Wallis test with

Dunn's multiple comparisons test for three or more samples experiments. For all

graphs, unless otherwise stated, data are presented as mean and standard error of the

mean (SEM) and statistical significance was accepted for $p$ value $< 0.05$.

**Data availability**

Sequencing data is available at NCBI BioProject with accession number PRJNA1132369. All other data for this study are included in the manuscript and supporting files.

**Supplemental information**

Supplemental information includes 8 figures and 1 table.