Title: Transposon mutagenesis screen in *Klebsiella pneumoniae* identifies genetic determinants required for growth in human urine and serum

Jessica Gray[1,2#], Von Vergel L Torres[1#], Emily CA Goodall[1], Karl A Dunne[2], Chris Icke[1], Jessica L Rooke[1], Thamarai Schneiders[3], Adam F Cunningham[4], Jeffrey A Cole[2], Ian R Henderson[1*]

1. Institute for Molecular Bioscience, University of Queensland, St. Lucia, Queensland, Australia
2. Institute of Microbiology and Infection, University of Birmingham, Birmingham, United Kingdom
3. Division of Infection Medicine, University of Edinburgh, Edinburgh, United Kingdom
4. Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, United Kingdom

# Authors contributed equally
*For correspondence: i.henderson@uq.edu.au

Abstract

Klebsiella pneumoniae is a global public health concern due to the rising myriad of hypervirulent and multi-drug resistant clones both alarmingly associated with high mortality. The molecular microbial genetics underpinning these recalcitrant *K. pneumoniae* infections is unclear, coupled with the emergence of lineages resistant to nearly all present day clinically important antimicrobials. In this study, we performed a genome-wide screen in *K. pneumoniae* ECL8, a member of the endemic K2-ST375 pathotype most often reported in Asia, to define genes essential for growth in a nutrient-rich laboratory medium (Luria-Bertani medium), human urine and serum. Through transposon directed insertion-site sequencing (TraDIS), a total of 427 genes were identified as essential for growth on LB agar, whereas transposon insertions in 11 and 144 genes decreased fitness for growth in either urine or serum, respectively. Genome-wide functional studies like these provide further knowledge on the genetics of this pathogen but also provide a strong impetus for discovering new antimicrobial targets to improve current therapeutic options for *K. pneumoniae* infections.

Introduction

Klebsiella pneumoniae are Gram-negative bacteria found ubiquitously in natural and man-made environments [1-3]. In immunocompromised patients, and those with co-morbidities such as diabetes or alcoholism, *K. pneumoniae* can colonize multiple body sites causing a diverse array of infections ranging from life-threatening pneumonia, urinary tract infections (UTIs), and bacteremia [4-8]. *K. pneumoniae* is a frequently reported multidrug-resistant (MDR) pathogen isolated from nosocomial settings. Such infections have generally higher mortality rates and require longer treatment regimens and hospital stays, placing a significant burden on healthcare providers [9-11]. Of most concern is resistance to β-lactam antibiotics such as penicillin, aztreonam, and extended-spectrum cephalosporins, by horizontally acquired plasmid-mediated β-lactamases [12]. While combined antibiotic therapies have been deployed in the clinic, the emergence of pandrug-resistant strains has created an urgency to develop new and efficient approaches to treating *K. pneumoniae* infections [13-15]. The identification of genetic determinants essential for growth and those required for survival of *K. pneumoniae in vivo* may help elucidate novel therapeutic targets to better treat such infections [16, 17].

Recent large scale phylogenetic investigations have suggested that distinct lineages of *K. pneumoniae* exist in clinical and environmental settings [18]. Furthermore, two clinical pathotypes of *K. pneumoniae* have emerged: classical and hypervirulent. While the full repertoire of genetic factors required for *K. pneumoniae* virulence remains unclear, various studies have highlighted the importance of genes involved in adhesion, iron acquisition, capsulation and cell envelope biogenesis [19]. Nevertheless, for both classical and hypervirulent *K. pneumoniae* strains, the ability to colonize the urinary tract and survive in

the bloodstream are essential virulence traits and are dependent on a large and diverse range

of virulence factors [20]. Indeed, previous studies demonstrated that lipopolysaccharide (O-)

(9 serotypes) and capsular polysaccharide (K-) antigens (79 serotypes) contribute to the

resistance of serum-mediated killing and antiphagocytosis [21, 22]. However, prior genetic

screens have revealed that serum-resistance in Gram-negative bacteria is an intricate

phenotype defined by a multitude of factors (the serum resistome) [23-27]. A recent study of

K. pneumoniae identified 93 different genes associated with serum resistance across four

distinct strains but found only three genes (*lpp*, *arnD*, and *rfaH*) played a role in all strains.

These data hint that multiple mechanisms of serum resistance may exist in *Klebsiella* lineages.

In contrast, the full repertoire of genes required for growth of *K. pneumoniae* in the urinary

tract (the urinome) has not been defined.

To define the genetic basis of the *K. pneumoniae* urinome and serum resistome, we used

Transposon Insertion Sequencing (TIS), also referred to as Transposon Directed Insertion-

site Sequencing (TraDIS). TraDIS is a genome-wide screening technique that has been widely

used to identify genes essential for bacterial growth and to discover conditionally essential

genes under physiologically relevant conditions [28-38]. In this study, we generated a highly

saturated transposon library within *K. pneumoniae* ECL8, a member of the K2-ST375

phylogenetic lineage that includes hypervirulent clones of epidemiological significance known

to cause infections in relatively healthy subjects predominantly in Asia [39]. We identified

the repertoire of genes essential for growth in laboratory conditions, in pooled human urine,

and in human serum. Selected fitness-genes were validated for growth in pooled urine or

serum by generating single-gene deletion mutants. Our study provides insight into the

molecular mechanisms that enable *K. pneumoniae* to survive *in vivo* and cause disease.

Methods

Bacterial strains, plasmids, and culturing conditions.

A complete list of bacterial strains, plasmids, and primers utilized in this study are listed in Table S1 and Table S2. To culture strains, scrapings from a frozen glycerol stock strain were plated onto solid Luria-Bertani (LB) (10 g tryptone, 5 g yeast extract, 10 g NaCl) supplemented with agar 1.5% (w/v) and were incubated overnight at 37°C. A single colony was isolated and used to inoculate 5 mL of liquid LB medium incubated overnight at 37°C with 180 RPM shaking. To prepare stocks, bacteria were grown to mid-exponential phase in LB medium and stored at -80°C with 25% (v/v) glycerol. When required, solid or liquid media were supplemented with appropriate antibiotics at the following concentrations: ampicillin (35 µg/mL); chloramphenicol (100 µg/mL); and kanamycin (100 µg/mL).

Generation of the *K. pneumoniae* ECL8 transposon mutant library

An overnight culture of *K. pneumoniae* ECL8 was inoculated into 2× YT broth at OD_{600} 0.05 supplemented with a final concentration of 0.7 mM EDTA and incubated at 37°C with 180 RPM shaking. Cells were harvested at 0.4 OD_{600} by centrifugation (4,000 *g*) at 4°C for 20 min. Cell pellets were washed 4 times with ice-cold 10% (v/v) glycerol. *K. pneumoniae* ECL8 electrocompetent cells were transformed with 0.2 µL of EZ-Tn5™ transposon (Epicentre) at 1.4 kV and recovered in brain heart infusion (BHI) broth for 2 h at 37°C with 180 RPM shaking. Cells were plated onto LB agar supplemented with 50 µg/mL kanamycin and incubated overnight at 37°C. More than 1 million transformants were pooled in 15% (v/v) LB-glycerol for storage at -80°C until required.

Transposon library screening in human urine

Human urine from 7 healthy male volunteers was sterilized using a vacuum filter (0.2 μM). Urine was pooled and stored at -80°C until required. Approximately 2×10^8 *K. pneumoniae* ECL8 TraDIS mutants were inoculated into 50 mL of urine and grown for 12 h at 37°C with shaking (P1) in biological duplicates in parallel with control samples that were instead inoculated into 50 mL of LB medium. A sample of P1 was inoculated into 50 mL of urine or LB medium, respectively, at an initial OD_{600} of 0.05 and grown for a subsequent 12 h (P2). Both control and test experiments were passaged and grown for a further 12 h (P3). A sample of culture from P3 was removed and normalized to an OD_{600} of 1 for subsequent genomic DNA extraction.

Transposon library screening in human serum

A 10 mL sample of human blood was collected from 8 healthy volunteers. Blood was pooled and centrifuged at 6,000 *g* for 20 min to separate sera from blood components. The pooled serum was sterilized, aliquoted and stored at -80°C until required. Approximately 2×10^8 *K. pneumoniae* ECL8 mutants were inoculated into 1 mL of pre-warmed sera, heat-inactivated sera (60°C for 1 h) or PBS in biological duplicates. Samples were incubated at 37°C for 90 min with 180 RPM shaking. Following this, cells were harvested by centrifugation at 6,000 *g* for 10 min and washed twice with PBS. Cells were then inoculated into 50 mL of LB for outgrowth at 37°C with 180 RPM shaking and harvested at an OD_{600} of 1 by centrifugation at 6,000 *g* for 10 min. A 1 mL sample of each culture was used for genomic DNA extraction.

Transposon library sequencing and data analysis.

Pooled kanamycin resistant *K. pneumoniae* ECL8 cells were prepared for sequencing following

an amended TraDIS protocol [28](21). Briefly, genomic DNA from the *K. pneumoniae* ECL8

library of pooled mutants was extracted from ~1×10^9 cells in biological duplicate, as per

manufacturer's instructions (Qiagen QIAamp DNA blood minikit). The DNA concentration of

replicates was determined using the Qubit 2.0 fluorometer (Thermo Fisher Scientific). The

DNA was sheared into ~200 bp fragments by ultra-sonication (Bioruptor Plus Diagenode).

Fragmented DNA samples were prepared for Illumina sequencing using the NEBNext Ultra I

DNA Library Prep Kit for Illumina according to manufacturer's instructions with the following

modifications: ends of fragmented DNA were repaired using the End Prep Enzyme Mix (NEB)

and an NEBNext adaptor for Illumina sequencing was ligated to the newly repaired ends.

During protocol optimization, samples were analyzed using a Tapestation 2200 (High

Sensitivity D5000) to determine DNA fragment sizes following adapter ligation. The uracil

within the adaptor hairpin loop was enzymically excised using the USER enzyme (NEB).

AMPure XP SPRI beads (Beckman Coulter) or custom-made SPRI beads were used to select

for DNA fragments ~250 bp in size.

To enrich for fragments containing the transposon a custom PCR step was introduced using a

forward primer annealing to the 3' end of the antibiotic marker of the transposon and a

reverse primer that annealed to the ligated adaptor. The PCR product was purified using SPRI

beads at a ratio of 0.9:1 (beads:sample). A further custom PCR step introduced sequencing

flow-cell adaptors (Illumina barcodes) and inline barcodes to allow for sample multiplexing

(Table S3). Samples were then stored at -20°C until sequencing. After qRT-PCR (KAPA Library

Quantification Kit Illumina® Platforms) to determine TraDIS library concentration, a 1.5 μL

sample of each library sample diluted to 8 nM was combined and mixed by pipetting to

generate a pooled amplified library (PAL) of samples for multiplexed sequencing. The PAL was

prepared for Illumina MiSeq sequencing according to manufacturer's instructions.

Bioinformatic analysis of sequencing reads was completed on the Cloud Infrastructure for

Microbial Bioinformatics (CLIMB) [40]. Raw fastQ files were first demultiplexed according to

Illumina barcodes automatically using the Illumina MiSeq software.Experimental replicates

were separated by custom inline barcodes using the FastX barcode splitter and the barcodes

were subsequently removed using the FastX trimmer (v0.0.13). Each read was checked for

the presence of the transposon sequence in two steps; first, reads that contained the first 25

bp of the transposon sequence (AGCTTCAGGGTTGAGATGTGTA), introduced by the TKK_F

primer, allowing for 3 mismatches were filtered and parsed. Parsed reads were subsequently

checked for the presence of the final 10 bp of the transposon sequence (TAAGAGACAG),

allowing for 1 mismatch. The transposon sequence and reads <20 bp were trimmed using

Trimmomatic (v0.39). Resulting reads were mapped to the *K. pneumoniae* ECL8 [HF536482.1

and HF536483] genome and plasmid using the Burrows-Wheeler Alignment tool (BWA-MEM).

Mapped reads were indexed using samtools (v1.8) and converted into bed format using the

Bam2bed tool (Bedtools suite v2.27.1). The resulting bed file was intersected against the

annotated CDS in the *K. pneumoniae* ECL8 gff file, which was generated using PROKKA

(v1.14.0). Transposon insertion sites and their corresponding location were quantified using

custom python scripts. Mapped transposon insertion sites were visualized using the Artemis

genome browser [41].

Statistical analysis of transposon insertion density

A modified geometric model was applied to identify the probability of finding k-1 insertion-

free bases followed by a transposon insertion as reported previously [28](21). In a string of

10,000 independent trials the probability of an insertion 'p' was calculated using the following

equation: P(k) = p(1-p)(k).

Identification of putative essential genes

In-house scripts kindly provided by Langridge and colleagues [42](34) were amended and

used for the prediction of essential genes. Briefly, the number of unique transposon insertion

sites for each gene was normalized for CDS length (number of UIPs/CDS length in bp), which

was denoted the insertion index score. The Freedman-Diaconis method was used to generate

a histogram of the insertion index scores for each gene of the *K. pneumoniae* ECL8 TraDIS

library. Distributions were fitted to the histogram using the R MASS library (v4.0.0).

An exponential distribution was applied to the 'essential' mode, situated on the left of the

histogram. A gamma distribution was applied to the 'non-essential' mode, situated on the

right of the histogram. The probability of a gene belonging to each mode was calculated and

the ratio of these values was denoted the log-likelihood score. Genes were classified as

'essential' if they were 12 times more likely to be situated in the left mode than the right

mode. Genes with log-likelihood scores between the upper and lower $\log_2(12)$ threshold

values of 3.6 and -3.6 respectively were classified as 'unclear'. Genes with log-likelihood

scores below the 12-fold threshold were classified as 'non-essential'.

Comparison of essential gene lists between ECL8, KPNIH1, RH201207 and ATC43816

Annotated genomes of *K. pneumoniae* KPNIH1 (Accession: CP009273.1), RH201207

(Accession: FR997879.1), and ATCC 43816 (Accession: CP009208.1) were downloaded from

NCBI. Nucleotide sequences of putative essential genes were extracted using the SeqKit

toolbox [43](33). *K. pneumoniae* ECL8 essential gene homologs were identified in KPNIH1,

RH201207 and ATC43816 using Galaxy AU NCBI BLAST+ wrapper blastn using the following

criteria (e-value > 1e-10, percent identity >90) and utilizing the top hit based on calculated e

value and bitscore [44].

BioTraDIS analysis for the identification of conditionally advantageous genes

To identify conditionally essential genes, we used the Bio-Tradis analysis pipeline [17], that

measures the read count $\log_2$ fold changes between each CDS. To ensure robust analysis, CDS

with <50 sequence reads in either the test or control condition were filtered and binned. CDS

with a $\log_2$ fold-change >2 (test/control) and a Q-value of >0.05 were classified as conferring

a fitness advantage under the conditions tested. To comparatively and analyze the impact of

transposon insertion-events in genomic regions in the serum-resistant dataset (180 min) to

the input, we used the AlbaTraDIS package (v1.0.1) with the following parameters: -a -c 10.0

-f 2 -m 0 [45].

Generation of knock-out strains

Mutants derived from *K. pneumoniae* ECL8 were constructed using the λ Red recombinase

system [46]. Briefly, the *K. pneumoniae* ECL8 strain was transformed with pACBSCSE, which

contains genes coding for the arabinose-induced λ Red recombinase system that permits

homologous recombination between dsDNA PCR products and target loci in the bacterial

genome. The recombination is based on short stretches of flanking homology arms (~65bp)

with the site of recombination. Using pKD4 as a template for the selectable kanamycin

cassette, the PCR products used for replacing the target genes were amplified, gel extracted,

and electroporated into electro-competent strain *K. pneumoniae* ECL8 harboring pACBSCSE

prepared in the presence of 0.5% (w/v) arabinose. Candidate mutants were screened using

antibiotics, verified by PCR using primers outside the region of recombination then followed

by SangerSanger sequencing.

Bacterial growth assay

A single colony of each bacterial strain was inoculated into 5 mL of LB medium and grown

overnight as previously described. Strains were normalized to an OD_{600} of 1.00 and washed

twice in PBS by centrifugation at 6,000 *g* for 10 min. Cell pellets were resuspended in 1 mL of

the growth medium required for the bacterial growth assay. Greiner Bio-One 96-well U-

bottom microtiter plates were inoculated with bacterial strains at an initial OD_{600} of 0.02 in a

final well medium volume of 150 µL and sealed with a Breathe-Easy® sealing membrane

(Sigma Aldrich). Inoculated plates were incubated at 37°C with 300 RPM shaking, OD_{600}

measurements were taken at 15 min over 24 h using a CLARIOstar® plate reader (BMG

LABTECH).

Urine co-culture competition studies between K. pneumoniae ECL8 and isogenic mutants

Overnight cultures of *K. pneumoniae* ECL8 and an isogenic mutant were normalized to an

OD_{600} of 1.00 and washed twice with PBS. Cell pellets were resuspended in 1 mL of urine. A

500 µL of either the WT or mutant culture were inoculated into 25 mL of urine and incubated

at 37°C with 180 RPM shaking for 12 h. The sample was then transferred to 25 mL of fresh

urine at an initial OD_{600} of 0.05 and grown for a further 12 h. The culture was passaged for a

further 12 h of growth. These cultures were serially diluted in PBS at 0, 6, 12, 24 and 36 h and

plated onto LB agar. Following overnight growth, agar plates were replica plated onto either

LB agar or LB agar supplemented with 100 µg/mL kanamycin grown overnight at 37°C then

enumerated. Relative fitness was calculated as a competitive index (CI), defined as ratio of

mutant:WT viable cells divided by the corresponding inoculum.

Lipopolysaccharide extraction

A single colony of ECL8 was inoculated into 5 mL of LB medium and incubated overnight at

37°C with 180 RPM shaking. Overnight cultures were normalized to an OD_{600} of 1.00 and

centrifuged at 14,000 g for 10 min. Cells were resuspended in 100 µL of cracking buffer (0.125

M Tris-HCl, 4% (w/v) SDS, 20% (v/v) Glycerol, 10% (v/v) 2-mercaptoethanol in dH2O). The cell

suspension was incubated at 100°C for 5 min and transferred to -80°C for 5 min. The cell

suspension was incubated at 100°C for a further 5 min and centrifuged at 14,000 g for 10 min.

An 80 µL sample of the supernatant was transferred to a new tube and incubated with 5 µL

of 5 mg/mL Proteinase K (QIAGEN) for 1 h at 60°C. Samples were diluted ×2 with Laemmli

sample buffer (Sigma-Aldrich) and incubated at 95°C for 5 min. Samples were loaded onto a

4-12% Bis-tris precast NuPAGE gel (Invitrogen) and run at 150 V for 1.5 h. LPS was visualized

by silver staining using the SilverQuest kit (Invitrogen) following manufacturer's instructions.

Serum bactericidal assay (SBA)

Pooled serum was thawed on ice and pre-warmed to 37°C. Overnight cultures of bacterial

strains were normalized to an OD_{600} of 1.00 in LB medium. Bacterial cells were washed twice

with PBS and resuspended in a final volume of 1 mL PBS. A 50 µL sample of the cell suspension

was inoculated into 50 µL of sera. Sera and cell samples were incubated at 37°C with 180 RPM

shaking. Viable bacterial cells were enumerated by plating 10 μL of cell and sera suspension

onto LB agar over a time-course: 0 min, 30 min, 60 min, 90 min, 180 min. Serum bactericidal

activity was plotted as the $\log_{10}$ change in colony forming units (CFU) relative to the initial

inoculum.

Results and discussion

Generation and sequencing of a *Klebsiella pneumoniae* strain ECL8 transposon input library

A *K. pneumoniae* ECL8 mini-Tn5 mutant library composed of >1 million mutants was

constructed and subsequently subjected to TraDIS. The resulting reads were mapped to the

reference genome that consists of a ~5.3 Mb chromosome and a single 205-kb plasmid (EMBL

Accessions: HF536482.1 and HF536483) (Figure 1A-B). The gene insertion index scores of the

two technical replicates of the *K. pneumoniae* ECL8 TraDIS library (KTL1 and KTL2) were highly

correlated with each other (R^2=0.955) demonstrating a high level of reproducibility (Figure

S1A). Therefore, sequence reads for both replicates were combined for downstream analysis

and essential gene prediction. Through iterative rounds of sequencing, we determined that

the library was sequenced to near saturation as the discovery of unique reads plateaued at

~2 million reads (Figure S1B). This library contained 554,834 unique genome-wide transposon

insertion sites and 499,919 of these insertions map within annotated coding sequence (CDS)

boundaries. This high level of transposon coverage equates to an average transposon

insertion every 9.93 bp throughout the genome, equivalent to an insertion approximately

every four codons. A de Brujin graph visualized in the genome assembly package Bandage

[47](38) revealed that the sequencing coverage of the plasmid was only 1.33 fold higher than

that of the chromosome indicating that the plasmid is likely present as a single copy (Figure

S2). Table 1 summarizes the data relating to the ECL8 libraries.

Table 1: Summary of transposon-containing sequence reads and unique insertion points

294 **(UIPs) mapped to the *K. pneumoniae* ECL8**

Sample	Mapped reads after QC	Genome-wide UIPs	CDS UIPs	Pearson Correlation of Replicates
ECL8 Input libraries (LB)				
KTL1	5,573,710	409,069	367,230	Figure S1A
KTL2	2,854,389	400,915	361,272	
KTL (Combined)	8,429,782	554,834	499,919	
Urine output libraries				
LB	4,324,125	4,324,125	403,381	Figure S7
Urine	4,839,277	4,839,277	389,251	
Serum output libraries				
Serum	4,729,032	115,328	101,348	Figure S9
Heat-inactivated serum	3,565,433	284,761	254,735	
180 min serum exposed	1,481,732	133,790	121,316	Figure S10

Essential chromosomal genes in *K. pneumoniae* ECL8

To classify a gene as essential or non-essential, the number of transposon insertions within each gene on the chromosome and plasmid was normalized for gene length and this value was denoted the gene insertion-index score (Figure 1C). Insertion index scores (IIS) for the *K. pneumoniae* genome followed a bimodal distribution, as has been previously described (Figure S3) [28, 42]. The probability of a gene being essential was calculated by determining the likelihood of each given insertion-index score as belonging to the essential or non-essential mode. The ratio of the IIS values was denoted the log-likelihood ratio (Log_2-LR), a metric previously used to categorize genes into essential or non-essential groupings [28, 42]. To limit the number of false-positive hits identified, genes were classified as essential only if they were 12-times more likely to be within the essential, rather than the non-essential mode. Based on the above criteria, 373 chromosomal genes were assigned a log_2-LR of less than -3.6 and were therefore classified as essential for growth on solid LB medium supplemented with kanamycin (Figure 1D). It should be noted that non-coding sequences such as tRNA and rRNA were excluded from this analysis. Most genes (4551) were classified as non-essential (Figure

1E). A subset of 241 genes were located between these two modes with their essentiality

mode was deemed 'unclear' (Figure 1F).

To understand the essential gene functions, we evaluated their COG (cluster of orthologous)

categories; a metric that predicts functional classification against a database of known

proteins. We applied a $\log_2$ COG enrichment index to identify COG categories that were

enriched among the essential genes in contrast to the wider genome eggNOG (v5.0) orthology

data [48] (Figure S4). We identified no essential genes within the COG categories for

chromatin structure and dynamics (B), cell motility (N) and secondary structure (Q) suggesting

genes within these categories are not required or redundant for *K. pneumoniae* growth in the

utilized nutrient-rich liquid medium. However, genes involved in signal transduction (T) and

inorganic ion transport/metabolism (P) were depleted, while genes involved in translation (J),

cell cycle control (D) and co-enzyme metabolism (H) were the most highly enriched. This

correlates with enriched COG categories shared among essential genes in other members of

the Proteobacteria such as *Salmonella enterica* Typhimurium, *S. enterica* Typhi and

Caulobacter crescentus and serves as an internal control for the validity of the library [49-51].

"Essential" plasmid genes

The identification of eleven genes on the *K. pneumoniae* ECL8 plasmid that met the criteria

to be defined as essential was an unexpected observation (Table 2). As previously noted, the

bacterial copy number of the plasmid was approximated to be one. The average IIS of a gene

located on the plasmid and the chromosome was comparable, 0.16 and 0.11, respectively.

Due to the presence of a single plasmid per bacterial cell, and an observed even distribution

of IISs for genes located on the plasmid, the lack of insertions in the 11 genes is unlikely to be

due to gene dosage effects arising from the presence of multiple copies of the plasmid.

Notably, the 11 plasmid-borne "essential" genes included *repB,* which is required for plasmid

replication. Therefore, these genes are likely to be required for plasmid replication and

stability and are unlikely to be essential for growth. This hypothesis is supported by previous

observations that *K. pneumoniae* remains viable when large indigenous virulence plasmids

are cured from the bacterium [52-54]. A complete list for all *K. pneumoniae* ECL8 bimodal

essentiality categorization and their

associated statistical significance metrics are listed in Dataset S1.

Table 2: *K. pneumoniae* ECL8 plasmid-borne genes computationally deemed essential.

Locus tag	Gene	Function[a]
ecl8_05075	*repB*_1	RCR (rolling circle replication) plasmid protein
ecl8_05092	-	SidC homolog
ecl8_05094	-	DUF2509 family protein
ecl8_05096	-	NB-ARC domain-containing protein
ecl8_05104	*rcsA*_2	DNA-binding transcriptional activator RcsA
ecl8_05105	*yedA*_2	putative transporter
ecl8_05106	-	DUF2695 domain-containing protein
ecl8_05170	-	DUF305 domain-containing protein
ecl8_05205	-	NADPH dependent preQ0 reductase
ecl8_05272	-	Transposase
ecl8_05279	*sopB*	Control of plasmid partitioning

[a]Function derived from the top hit using NCBI blastN (www.blast.ncbi.nlm.nih.gov/Blast.cgi)

Insertion free regions

Using a highly saturated transposon library it is possible to not only define essential genes, but also define chromosomal regions lacking transposons that are large enough not to have occurred by chance. Previously, we described a geometric model to predict statistically significant regions of the genome that were free from transposon insertions [28]. This model indicated that with the insertion frequency calculated for this library, for a genome of 5.3 Mb, an insertion-free region (IFR) of 145 bp or greater was statistically significant (Figure S5). The size of insertion-free regions that are statistically significant for two previously published TraDIS libraries are plotted for comparison [28, 42]. A total of 667 genomic regions with insertion-free regions ≥145 bp were identified, many which correspond to the 380 essential genes identified through the bimodal analysis described earlier.

The remaining insertion free regions can be explained in several ways. First, as noted

previously [28], genes that encode proteins with essential domains are often excluded from

the bimodal analysis described above as much of the gene contains transposon insertions.

For example, the *polA* gene was classified as "unclear" but contained, barring a single

transposon insertion event, an 826-bp insertion-free region. This IFR corresponds to 275

amino acids that are predicted to encode the 5'-3' exonuclease function of the protein (Figure

1F). This region has also been shown to be required for the growth of *Escherichia coli* in

laboratory conditions [55, 56]. In contrast, but as noted for *E. coli polA*, multiple transposon

insertions are present in the remaining portion of the gene, which confers the 3'-5'

proofreading and polymerase activity. Aside from those genes already identified by the

bimodal analyses described above, our analyses revealed 54 additional genes with statistically

significant IFRs, which indicates that they are likely essential for growth under the conditions

tested here. Thus, our data suggest *K. pneumoniae* contains at least 427 essential genes,

which we define as the ECL8 curated essential gene list.

An explanation for the remaining IFRs can be derived from traditional bioinformatic

approaches to genome annotation. Originally, genome annotation protocols overlooked

small ORFs because (i) they only consider genes that code for proteins >100 amino acids, and

(ii) define genes based on sequence homology to those that have been previously annotated

[57-59]. Interestingly, the methodology used here revealed 59 regions ≥145 bp that lacked

transposon insertion sites and did not map to within annotated CDS boundaries (Figure 1G).

The applied approach is conservative and does not include insertion free regions that overlap

with the 5' or 3' region of any annotated CDS. Forty-six of these regions contain potential

open reading frames that could encode proteins. These putative unannotated ORFs were not

characterized further within this study but might represent essential genes. This study

emphasizes the untapped potential of TraDIS datasets to identify previously uncharacterized,

essential, small coding sequences and promoter elements that are not immediately obvious

by traditional and computational genomic annotation methods. A complete list of IFRs, within

annotated ORFs or in intergenic regions, and an ECL8 curated essential gene list is found in

Dataset S2.

Directional insertion bias of Tn5 transposon into capsular polysaccharide (*cps*) operon

As reported previously [28, 60], due to the design of the transposon used in this study,

transcription of polycistronic operons can be affected by the insertion orientation of the

transposon (Figure 2A). Directional insertion bias of the transposon was noted within a large

genomic region of ~14 kb encoding the virulence-associated capsular polysaccharide (*cps)*

operon (Figure 2B). The bioinformatic web-tool 'Kaptive' determined with high confidence

the *K. pneumoniae* ECL8 *cps* operon encodes for a KL2/K2 capsule with 18/18 of the expected

genes (Dataset S6) [61].

Notably, transposon insertions were poorly tolerated in the forward Tn orientation of the

DNA strand. While bias for Tn insertions has previously been described in individual genes

(e.g. ribosomal RNA genes), this study is the first to describe such a large bias across an operon

[28, 49]. However, the basis for this bias is not fully understood. One potential explanation

for such bias might be the presence of a gene encoding a toxic small RNA (tsRNA). Typically,

tsRNAs are usually present in the 5' or 3' untranslated regions of genes and act as regulators

of gene expression at a post-transcriptional level, but due to their small size, typically <500

nucleotides, they are very difficult to predict [62, 63]. However, this hypothesis does not

explain why transposon insertions are uniformly absent for the reverse orientation of the

transposon and not isolated to 'hotspots' in the 5' and 3' untranslated regions. An alternative

hypothesis is that the reverse orientation expression of the capsular operon might result in

the accumulation of intracellular capsular intermediates that may be toxic if left to

accumulate in the periplasm. This finding may hint to a more cryptic regulation mechanism

of the *cps* operon with implications for other similar *Klebsiella* capsular serotypes and species.

Comparison of essential genes to other *K. pneumoniae* transposon libraries

Transposon libraries have previously been generated in other *K. pneumoniae* strains, however

these studies primarily focused on identifying genes that are required for a specific phenotype

i.e antibiotic stress or capsular mutants, and therefore they lack a definitive list of essential

genes for *in vitro* growth [64-68]. To benchmark our library, we compared our data to

previously published studies that did report *K. pneumoniae* essential gene lists: KPNIH1

(ST258), RH201207 (ST258) and ATCC 43816 (ST493) [31, 69]. As shown in Table 3, our

transposon mutant library is amongst the most saturated *K. pneumoniae* libraries reported to

date, permitting accurate determination of gene essentiality and delineating between real

and stochastic effects. The number of essential genes among the four strains ranged from 434

to 642; a complete list and comparison of our curated gene list against the reported gene lists

of KPNIH1, RH201207 and ATCC 43816 is found in Dataset S3. We speculate that the higher

number of essential genes predicted in the KPNIH1 gene list is due to a lower transposon

density, as this is associated with an increased likelihood of false-positive results [70]. This

might be due in part to differences in the techniques used to construct the libraries; here we

used a mini-Tn5 transposon which showed no bias in insertion whereas others have used the

TnSeq method relying on the Himar I Mariner transposon, which has a requirement for a TA

dinucleotide motif [71]. Differences in gene essentiality might also be due to inherent

genomic differences or due to differences in experimental methodology, computational

approaches, or the stringency of analysis used to categorize these genes. While further

validation experiments will be required to determine whether specific genes are essential,

57% of the essential genes identified in *K. pneumoniae* ECL8 were essential in the other three

strains and a further 13% were shared across at least two of the three strains. Thus, these

data indicate that our transposon library is sufficiently dense and representative of *K.*

pneumoniae to be used for further studies.

Table 3: Comparison of *K. pneumoniae* ECL8 essential gene list with other selected studies

Strain and ST	Chromosome length (bp)	UIPs	Total essential genes	Homolog essential in ECL8[a]	Essential, no homolog in ECL8	Reference
ECL8 (ST375)	5,307,154	499,919	427[b]	245[c]	27[d]	This study
KPNIH1 (ST258)	5,394,056	41,769	642	297	103	[31]
RH201207 (ST258)	5,475,790	510,834	608	172	227	[64, 69]
ATCC 43816 (ST493)	5,374,835	415,608	476	308	121	[65, 69]

[a] Listed as essential in the referenced study, but corresponding ECL8 homolog was deemed non-essential (E-value >1e-10,
nucleotide percent identity >90).
[b] Curated essential chromosomal gene number: bimodal analysis (n=373) + manual analysis of "unclear" genes with IFR
(n=54).
[c] A total of 245 genes deemed essential in ECL8 with homolog(s) present in the reported essential gene lists of KPNIH1,
RH201207, ATCC43816.
[d] A total of 27 essential genes were identified in ECL8 with no homolog in KPNIH1, RH201207, ATCC43816 (E-value >1e-10,
nucleotide percent identity >90)
450

Identification of genes required for growth in human urine

TIS has become a formidable tool for rapid genome-scale screens to link genotype to

phenotype. As the urinary tract is a major site for *K. pneumoniae* infection, we used our

transposon mutant library to identify genes required for growth in urine, thus enabling us to

define the urinome of *K. pneumoniae*. Cultures were grown until late stationary phase (12 h)

followed by two subsequent 12 h passages before sequencing (Figure 3A). This approach

eliminated mutants that were unable to grow in urine while allowing those capable of growth

to flourish. DNA from test and control samples was harvested, sequenced, and analyzed as

before (Figure S6). As the gene IISs for the library passaged in both LB and urine were highly

correlated with respective biological replicates of 0.8483 and 0.9078 R^2, replicates were

combined for downstream analyses (Figure S7). The data for all *K. pneumoniae* genes and

their associated statistical significance metrics are listed in Dataset S4.

Ten genes that satisfied a stringent threshold of a $\log_2$ fold change ($\log_2$FC) >-2 and a Q-value

≤0.05, relative to the LB control, were considered as advantageous for *K. pneumoniae* ECL8

growth in urine (Figure 3B). The transposon insertion profiles of these genes showed a marked

decrease in overall insertions following passaging in urine when compared to the LB control

(Figure 3C-F), suggesting loss of these genes confers a fitness defect. These included genes

encoding the superoxide dismutase SodA, the structural outer membrane protein OmpA (63),

the periplasmic chaperone Skp [72], and five proteins (*fepB*, *fepD*, *fepG*, *exbB* and *exbD*)

associated with iron acquisition [73]. The remaining two genes (*ytfL* and *argH*) play key roles

in controlling cytoplasmic cadaverine and putrescine concentrations and amino acid

biosynthesis, respectively [74, 75]. The proper regulation of cadaverine and putrescine

maintains intracellular pH concentrations and reduce oxidative damage to proteins and DNA

and may be in response to acidic conditions in urine known to be unconducive for colonization

of uropathogens [76, 77].

To experimentally validate our observations, seven independent mutants were constructed

by allelic replacement of the chromosomal gene with a kanamycin (*aph*) cassette. To

determine whether *K. pneumoniae* strains lacking the genes identified in this screen were

truly less fit in urine compared to the parent strain, the WT ECL8 and individual isogenic

mutant strains were inoculated into urine or LB medium at a 1:1 ratio. These cultures were

sequentially passaged, and the CFU/mL of both the WT and mutant strain were determined

over a time-course of three 12 h passages (Figure 3G). When compared to the wild-type, a

Δ*sodA*::*aph* mutant was less fit in both urine and LB, but the phenotype was exacerbated in

urine. In contrast, the fitness of the remaining mutants was comparable to the parent strain

when grown in LB medium and only less fit when grown in urine. While the relative

competitive index of the Δ*ytfL*::*aph* strain was 0.8 in urine, the *ompA*, *fepB*, *fepD*, *exbB* and

exbD mutants were dramatically less fit than the parent strain in urine (Figure 3G).

Based on these observations, we hypothesized that iron-levels were limiting growth in human

urine and that the ability to acquire iron was crucial for *K. pneumoniae* to survive and grow *in*

vivo [78, 79]. To test this hypothesis, urine was supplemented with exogenous ferrous iron

(100 μM $FeSO_4$) or an iron chelator (100 μM bipyridine/2,2-dipyridyl). No variance in optical

density was observed after passaging in LB medium (Figure 4A). However, the optical density

of *K. pneumoniae* cultures decreased after passage in urine (Figure 4B), and the restricted

growth was further exacerbated by the addition of the iron chelator 2,2-dipyridyl (Figure 4C).

In contrast, supplementation of urine with exogenous iron increased growth relative to non-

supplemented cultures with no difference in growth upon passage (Figure 4D). Surprisingly,

several siderophore synthesis genes did not confer a statistically significant fitness advantage

for growth in urine (i.e. *entACDF*, Figure 4E). In contrast, the highest Log_2FC increase (~7-fold)

in read counts mapped to the Ferric Uptake Regulator (*fur*) gene indicating its loss is beneficial

for growth in urine. In Gram-negative bacteria, Fur is a key regulator for iron homeostasis

functioning as a transcriptional repressor for siderophore synthesis genes in an iron

concentration-dependent manner [80-82]. Furthermore, the Fur regulon includes the

cognate ferric enterobactin uptake (Fep) genes, for which decreased Log_2FC read counts were

also noted (Figure 4F). Altogether, this suggests that the Fur regulon plays a pivotal role for *K.*

pneumoniae growth in urine by tightly regulating enterobactin synthesis and enterobactin-

mediated iron uptake.

Identification of genes required for resistance to complement-mediated killing

To identify genes that protect against complement-mediated killing (previously defined as the

serum resistome [23, 24], we first established the serum killing kinetics of *K. pneumoniae*

ECL8. Approximately, 2 x 10^8 cells were incubated with 100 μL of human serum for 180 min.

At specified time points, aliquots were plated to determine the number of viable bacteria. As

expected, no viable bacteria were recovered from the *E. coli* BW25113 control [83]. In

contrast, a 2 $\log_{10}$ decrease in the number of viable *K. pneumoniae* cells was observed after

90 min exposure to normal human serum, and a >4 $\log_{10}$ decrease after 180 min exposure. To

determine whether the observed reduction in viable cells was due to the presence of active

complement proteins, a heat-inactivated serum control was included (Figure S8).

Subsequently, 2 x 10^8 cells from the ECL8 TraDIS library were inoculated into 1 mL of human

serum or 1 mL of heat-inactivated human serum and incubated for 90 min. Following

exposure to serum, cells were grown in duplicate to an OD_{600} of 1 in LB medium to enrich for

viable mutants. DNA was harvested from each sample, and subsequently sequenced as

described previously (Figure 5A). The Pearson correlation coefficient (R^2) of gene insertion

index scores (IIS) for two sequenced biological replicates were calculated as 0.91 (normal

human serum) and 0.76 (heat inactivated serum) (Figure S9). To ensure robust data analysis,

thresholds were applied to the data where genes with less than 50 mapped reads in the input

library were removed from downstream analysis to minimize the effect of the stochastic loss

of mutants on downstream analysis. A total of 356 genes with a $\log_2FC$ ≤-2, P<0.05, and Q-

value of ≤0.05 between the serum and heat-inactivated control were classified as genes that

are advantageous for growth under complement-mediated killing conditions (Figure 5B). The

complete list of all genes analyzed for growth in human serum and their associated statistical

significance metrics are listed in Dataset S5. The operons encoding lipopolysaccharide (LPS)

core antigen, O-antigen, and enterobacterial common antigen (ECA) biosynthesis were noted

as containing several conditionally essential genes (Figure 5C). Mutants defective in these cell

envelope structures have previously been reported to play a role in membrane integrity and

resistance to serum-killing for a variety of Gram-negative bacteria [84-86]. Genes implicated

as important for serum resistance encode proteins with an important role in membrane

stability. They include GalE involved in LPS sugar sub-unit biosynthesis [87, 88], outer

membrane protein MlaA involved in phospholipid retrograde transport [89], cardiolipin

synthase ClsA [90], Bam complex component BamE [91], and proteins involved in LPS

biosynthesis (LpxC) and its regulator (YejM) [92].

To validate our observations, we selected the gene with the greatest Log_2FC decrease

between serum and a heat-inactivated control; a putative glycosyltransferase *espE_3*/*wbbY*

(inset, Figure 5B). Analyses using the 'Kaptive' webtool suggested that ECL8 encodes the O-

antigen serotype O1v2 with the operon having 98.98% nucleotide identity to the reference

genome (Dataset S6) [61]. The O-antigen of the O1v2 serotype features distal repeating D-

galactan I and D-galactan II sugar sub-units [93]. Despite its role in O-antigen biosynthesis,

wbbY is not located within the *rfb* LPS operon. A previous study reported that its deletion in

K. pneumoniae NTUH-K2044 (ST23) resulted in abrogated D-galactan II production. To

determine the role of *wbbY* in serum sensitivity, the gene was replaced with an *aph* cassette

and the resulting mutant was compared to its parent strain using a serum bactericidal assay.

A comparable growth profile was observed for *K. pneumoniae* ECL8 and the Δ*wbbY*::*aph*

mutant in LB medium (Figure 5D). However, the Δ*wbbY*::*aph* mutant was more sensitive to

human serum than the parent with no viable mutants surviving after 30 min (Figure 5E). To

investigate whether the *wbbY* deletion affected LPS production, crude LPS extracts were

visualized using silver staining (Figure 5F). The Δ*wbbY*::*aph* strain produced LPS with a lower

molecular-weight compared to the parent strain, suggesting that a truncated O-antigen was

expressed. Alterations in the amount or length of LPS produced are known to influence

susceptibility to serum mediated killing [85].

Identification of genes required for serum-resistance

Following the identification of genes that when mutated confer increased sensitivity to

complement-mediated killing (i.e. the serum resistome), the *K. pneumoniae* ECL8 TraDIS

library was screened for mutants that confer increased resistance to serum-killing.

Approximately 2 x 10^8 TraDIS mutants were incubated in 1 mL of human serum for 180 min.

The output pool was washed with PBS and plated onto LB agar supplemented with kanamycin

(Figure 6A). Following overnight growth, ~150,000 colonies were recovered and pooled for

sequencing confirming good correlation of replicates (Figure S10). We then used AlbaTraDIS

to identify mutations that conferred a fitness advantage or disadvantage for survival in serum.

In addition to identifying genes with an increase or decrease in reads between the condition

and control samples, AlbaTraDIS will also analyze changes in sequencing read depth of

insertions immediately upstream and downstream of a gene, as well as within-gene

transposon orientation biases. Such differences can be strong indicators of mutations that

result in gene expression or regulatory effects (For a detailed explanation please see [45]).

The data for insertions with a significant (>2-fold) change in read-depth following serum

exposure for 180 min are listed in Dataset S7. Genes with the greatest significance in

decreased insertions, suggesting decreased fitness, were O-antigen outer-core biosynthesis

gene espJ_2/*wabl* and O-antigen ligase rfaL/*waaL* (Figure 6B). Genes with significantly

increased insertions, suggesting increased fitness, included *hns*, *gnd* and *igaA* (Figure 6C).

Closer inspection of the transposon insertion profiles further confirmed the enrichment of

hns mutants when comparing the serum exposed test to the input control (Figure 6D). To

validate our observations, a defined *hns* mutant was constructed by allelic replacement with

an *aph* cassette in the same orientation as native gene transcription. A comparable growth

profile in LB was observed for the *hns* mutant relative to the parent (Figure 6C). To determine

whether the *hns* mutant was indeed more resistant to serum a sample of 1×10^8 cells was

incubated in serum for 360 min. At 30 min post-incubation the number of viable *hns*::aph cells

was decreased by ~2 log. However, no additional decrease in viable cell numbers was

observed at later time points and these strains proliferated in serum (Figure 6F). Ares and

colleagues demonstrated that *rcsA; galF; wzi;* and *manC* were derepressed in a Δ*hns K.*

pneumoniae mutant resulting in increased capsule production [94]. Similarly, up-regulation

of *rcsA* was noted in an *E. coli* Δ*hns* mutant. RcsA is known to positively regulate the capsular

locus [95, 96]. Notably, IgaA represses the Rcs regulon, preventing activation of RcsA. Mutants

in *igaA* are enriched in our experiments. These data suggest that activation of the Rcs regulon

confers resistance to serum killing in *K. pneumoniae*, as noted for other members of the

Enterobacteriaceae, by increasing capsule production [97].

The *gnd* gene encodes a 6-phosphogluconate dehydrogenase involved in the oxidative branch

of the pentose phosphate pathway. Interestingly, sequence reads mapping to *gnd* were

enriched for the forward orientation of the transposon only (Figure S11). Transposon

insertions in this orientation have transcriptional read-through driving expression of the

downstream neighboring gene *manC* and more distal O-antigen biosynthesis genes (i.e. *rfb*

cluster) [98]. The *manC* gene encodes mannose-1-phosphate guanylyltransferase that is

required for the synthesis of capsule, and as noted above, upregulation of *manC* results in

increased production of capsule.

Conclusion

For a variety of reasons, the determination of gene essentiality from TraDIS data is

challenging. Here, we used two different computational approaches to identify essential

genes ensuring that our essential gene dataset includes those genes that can tolerate

insertions in regions of the gene but where specific domains are essential for growth. The

observation that 57% of the essential genes identified in this study were found to be essential

in three of the four *K. pneumoniae* TraDIS libraries described to date provides reassurance

that this library would provide meaningful data for subsequent studies. Furthermore, the high

density of insertion sites achieved in our study ensured reliable identification of essential and

conditionally-essential genes and a low probability of false positive identification, which had

previously been noted for low complexity libraries [70].

The search for genetic determinants of bacterial survival and growth *in vivo* has been

augmented by whole genome approaches to infection, including transposon insertion site

sequencing techniques such as TraDIS and TnSeq [16]. These data have potential implications

for the design and development of novel drugs, vaccines and antivirulence strategies. In this

study, we employed TraDIS to identify genes which, when mutated, confer decreased or

increased fitness for growth in human urine and serum, two *in vivo* niches relevant to *K.*

pneumoniae infection. Surprisingly, our study revealed 11 genes significantly required for

growth in urine. These included genes encoding the outer membrane protein OmpA, its

chaperone Skp, the superoxide dismutase SodA, and genes required for iron acquisition by

the enterobactin uptake system. Notably, not all the genes for enterobactin synthesis, export

and uptake were essential (Fig. 4). The gene encoding FepB, the periplasmic chaperone for

the iron laden siderophore, was essential, as were *fepD* and *fepG*, which encode the inner

membrane uptake system. The TonB complex (TonB-ExbB-ExbD), which energizes the

translocation of iron laden enterobactin through the FepA outer membrane pore, was also

essential for growth in urine. In contrast, the genes encoding FepA, the inner membrane

ATPase FepC, and the enterobactin synthesis (Ent) and export (TolC) components were not

essential. These observations can be explained in two ways. First, as *K. pneumoniae* ECL8

contains paralogous copies of *fepA* (*fepA*_1 and *fepA*_2) and *fepC* (*fepC*_1 and *fepC*_2), the

non-essential nature of *fepA* and *fepC* might be the result of functional redundancy. Second,

as TraDIS is in principle a large-scale competition experiment between thousands of mutants,

mutants lacking enterobactin (or aerobactin) synthesis and export genes can "cheat" by

acquiring the siderophore released from siderophore-producing strains within the

population, a phenomenon described previously [99, 100].

In contrast to the limited number of genes in the urinome, the serum resistome consisted of

more than 144 genes. This number is significantly greater than reports of a recent study that

defined the serum resistome of four independent *K. pneumoniae* strains [23]. While that

study found 93 genes that were required for survival in one or more strains, remarkably only

three genes were common to all four strains: *rfaH*, *lpp* and *arnD*. Comparison with our dataset

revealed that in contrast to *rfaH, arnD* was not required for serum resistance in *K.*

pneumoniae ECL8. Surprisingly, comparison of our data with the 56 genes comprising the

serum resistome of the more distantly related multidrug resistant *E. coli* strain EC958 revealed

14 genes in common. The variation in the outcome of these experiments can be accounted

for in different ways. First, in all cases the genes encoding O-antigen and/or capsule synthesis

were required for growth in normal human serum; As the genes encoding these surface

structures vary with serotype, they are not identified as common genes, though ostensibly

they have the same function Second, the methodology appears at first glance to be identical

in each study (growth in the presence of serum for 90 min). However, not all strains exhibit

the same killing kinetics in normal human serum. Using fixed timepoints, strains with longer

killing kinetics are likely to have fewer genes identified in TraDIS experiments, whereas strains

with short killing windows will have more. Third, several of the genes identified in the serum

resistome of other strains were excluded from our downstream analysis as there were fewer

than 50 transposon insertions in the gene following exposure to heat-inactivated serum. A

case in point is the *lpp* gene. This observation indicates that *lpp* is required for *K. pneumoniae*

ECL8 growth in human serum but not resistance to complement dependent killing *per se*.

Fourth, the density of the library used, the level of comparative annotation, and the

thresholds to define significance, all have the potential to influence the final outcome of such

experiments. Finally, strain-specific features such as gene duplication events, functional

redundancy in biosynthetic pathways, and the variation in the genetic complement of each

strain, are likely to influence the contribution each gene makes to serum resistance.

Despite the intra and interspecies variation in the serum resistome noted above, comparison

of the urinome and serum resistome of *K. pneumoniae* ECL8 revealed that mutation of five

genes (*ompA*, *galE*, *exbB*, *exbD* and *sodA*) conferred reduced fitness for both growth in urine

and survival in serum. The role of *exbB* and *exbD* are described above but their detection in

both experimental conditions reinforces the importance of iron acquisition for infection. The

major outer membrane protein OmpA identified in this study encodes a functionally

pleiotropic outer membrane protein (85, 86). Notably, a role for OmpA in serum resistance

has previously been described for *E. coli*, however it was not identified by TraDIS experiments

investigating the serum resistome [24]. The molecular basis for this discrepancy remains

unresolved. GalE is known to play a key role in serum resistance for *E. coli* and *Salmonella*

spp. [24]. The gene *galE* encodes a UDP-glucose 4-epimerase catalyzing the interconversion

of UDP-D-galactose/UDP-D-glucose [87]. It has been previously reported that GalE plays a role

in D-galactan II synthesis, an O1 LPS component in *K. pneumoniae* [88, 101]. These data

suggest that loss of *galE* results in a modified LPS in *K. pneumoniae* resulting in increased

serum susceptibility. Subsequently, the manganese-dependent superoxide dismutase SodA

converts superoxide to hydrogen peroxide and oxygen to prevent DNA and other cellular

damage [102, 103]. A previous study in *K. pneumoniae* has reported the inducible expression

of *sodA* in a triumvirate system with *sodB* and *sodC*, expressed in response to anaerobic

challenge and high-oxygen concentration [104]. These transcriptional responses might very

well be present in both the microenvironments of blood vessels and the bladder, in the form

of nutrient limitation and oxygen exposure [105-107]. Hence, the observed importance of

sodA might be a consequence of elevated levels of reactive oxygen species (ROS) due to iron-

acquisition and could represent a specialized metabolic adaptation for growth in these

environments. Ultimately, these findings underline the interplay between bacterial iron

homeostasis and oxidative stress previously reported in Gram-negative bacteria [108-111].

However, these data shine a light on cross-niche (i.e. urine and serum) genes and associated

pathways as suitable drug development targets to treat *Klebsiella* infections. Finally, this

library was then used to identify genes conferring serum resistance. An interesting finding

was the observed transposon insertion orientation bias into *gnd.* As the downstream

transcriptional unit of *gnd* are O-antigen related genes (i.e. biosynthesis and export), this in

turn lends further credence to a transcriptional regulatory model where biosynthesis of O-

antigen and capsule is coupled for biophysical reasons as has been previously reported [65,

112]. Collectively, these data show that O-antigen biosynthesis genes (i.e *wabI* and *waaL*) play

a contributing role in serum resistance whilst *hns* mutants were shown to be more resistant

to serum killing than WT *K. pneumoniae* ECL8. However, the mechanism that underpins this

resistance is still largely unknown and requires further characterization. Despite the

previously described links between *hns* and the regulation of a selection of capsular genes

[65, 94], this is the first report of a role for *hns* in resisting complement-mediated killing in

human serum.

This genome-wide screen has identified a suite of genetic fitness-factors required for growth

of *K. pneumoniae* ECL8 in human urine and serum. The defined ECL8 urinome and serum

resistome gene sets from this study gene differ from other TraDIS studies, hinting that the

genotype of different pathogenic lineages might only be a piece in understanding the

underlying infection biology. Future studies like these but expanding to other isolates, and

the lesser studied clinically relevant *K. pneumoniae* species complex (KpSc), will be imperative

for the effective future development of targeted therapeutics towards this problematic

pathogen.

722
723

726
Figure 1 Overview of ECL8 TraDIS data: mapped insertions, insertion index profile and
example gene insertion plots. Insertions are illustrated on the (A) Chromosome and (B)
Plasmid, respectively. The outermost track displays the length of the ECL8 genome in base
pairs. The subsequent two inner tracks correspond to coding sequences (CDS) on the sense
(blue) and antisense (green) DNA strands, respectively. Putative essential CDSs are
highlighted in red. The inner-most track (black) corresponds to the location and read
frequency of transposon sequences mapped successfully to the *K. pneumoniae* ECL8 genome.
Plot generated using DNAPlotter. (C) Gene insertion index scores (IIS) of the *K. pneumoniae*
ECL8 TraDIS library mapped in order of genomic annotation of the *K. pneumoniae* ECL8 (left)
chromosome and (right) plasmid. Example transposon insertion profiles categorised into
essential, non-essential and unclear: (D) an essential gene – *bamD*, an essential outer
membrane factor for β-barrel protein assembly; (E) a non-essential gene – *int_*A1, a
redundant (several copies) integrase required for bacteriophage integration into the host
genome; (F) an "unclear" gene – *polA* an essential gene in prokaryotes required for DNA
replication but showed requirement for the N-terminal 5'-3' exonuclese domain; and (G) an
insertion free region suggestive of an unannotated ORF. Transposon insertion sites are
illustrated in black and capped at a maximum read depth of 1.
744

Figure 2 Directional insertion bias of transposon (Tn) into the *cps* operon. (A) Schematic
representing the Tn orientation and effect on downstream transcription. The utilized Tn
transposon is flanked by terminator sequences (purple) is shown inserted in gene A (yellow)
of a hypothetical two-gene transcription unit AB in the forward or reverse orientation. In
the forward orientation transcription of gene B (pink) is expected to occur from the
promoters of 5' of gene A or the internal Tn5 but polycistronic mRNA differs in length due to
the attenuation by the terminators. (B) Transposon insertions mapping to the K2 capsular
operon of *K. pneumoniae* ECL8. Transposon insertions are configured in the forward
orientation (green), and reverse orientation (blue) and insertion densities are capped at a
maximum read depth of 50. The operon structure of the K2 capsular genes consisting of
three promoters driving the expression of three unidirectional polycistronic transcripts is
depicted.

759
Figure 3 Overview and validation of ECL8 fitness-factors for growth in pooled human urine.
(A) Schematic of the experimental design used to identify genes that provide a fitness
advantage for *K. pneumoniae* ECL8 growth in pooled human urine. The *K. pneumoniae* ECL8
library was inoculated into either 50 mL of LB or 50 mL of urine and incubated at 37°C with
180 RPM shaking for 12 h. The library was passaged into 50 mL of fresh LB or pooled human
urine at an initial OD_{600} of 0.05 two subsequent times. A 1 mL sample normalized to an OD_{600}
of 1 from each culture was processed for genome extraction and multiplexed sequencing
using an Illumina MiSeq. (B) Log_2 fold change (Log_2FC) of the read count for each *K.*
pneumoniae ECL8 gene when passaged in urine relative to an LB control. Genes highlighted
in red satisfy a stringent applied threshold (Log2FC >-2, Q-value ≤0.05). The Q-value is the P-
value that has been adjusted for the false discovery rate for each gene. For brevity, only genes
with a Log2FC ≥0 are illustrated. Selected transposon insertion profiles of genes identified as
advantageous for growth in urine: (C) *fepB, fepD, fepG* (D) *exbB, exbD* (E) *sodA* and (F) *ompA*.
These genes exhibited a significant loss of transposon insertions following growth in urine
(red) in comparison to LB broth (blue). A 5-kb genomic region including the gene is illustrated.
Reads are capped at a maximum depth of 1. (G) The fitness of gene replacement mutants
relative to WT *K. pneumoniae* ECL8 in either LB medium or urine. The relative competitive
index of single-gene replacement mutants after 12 h passages ×3 in either LB medium or
urine. A relative fitness of one would indicate comparable fitness to WT. The mean is plotted
(± 1 SD).
780

781
Figure 4 Growth of the *K. pneumoniae* TraDIS library following passaging in LB and urine
and schematic diagrams of enterobactin synthesis, secretion and uptake. The OD_{600} of the
K. pneumoniae TraDIS library following 12 h of growth (P1) and two sequential 12 h passages
(P2 and P3). The library was passaged into fresh medium (A) LB or (B) urine to an initial OD_{600}
of 0.05. To determine the effect of iron supplementation and depletion, urine was
supplemented with exogenous iron (C) 100 μM $FeSO_4$ or an iron chelator (D) 100 μM 2,2-
dipyridyl. The average OD_{600} of three biological replicates for each time point is plotted (±) 1
SD. (E) Simplified schematic of the enterobactin synthesis pathway. YbdZ, a co-factor of EntF
for the terminal steps for enterobactin synthesis, depicted in light red had a Log_2FC sequence
read value of -1.58 suggesting this gene conferred an overall fitness advantage for growth in
urine. (F) Schematic representation of enterobactin secretion and uptake. The TonB transport
system is present in Gram-negative bacteria and is required to transport Fe-bound
enterobactin through the outer (OM) and inner membrane (IM) to the cytosol where it can
be utilized. Based on Log_2FC sequence read value, loss of TolC (blue) was beneficial for growth,
relative to an LB control. Loss of proteins, colored in red, had Log_2FC sequence read values <-
2 suggesting they confer a fitness advantage for grown in urine. Proteins depicted in grey
were genes that had Log_2FC values that ranged from -1 to 1 exposed to urine relative to an LB
control. Genes depicted in black were essential and had no determinable Log_2FC value.
800
801

Figure 5 Overview and validation of ECL8 fitness-factors for survival in pooled human serum.
(A) The experimental methodology utilized for screening the TraDIS library in human serum
and a heat-inactivated serum control. *K. pneumoniae* ECL8 ($2x10^8$ cells) of the mutant library
was inoculated into either 1 mL of human serum or 1 mL of heat-inactivated human serum
and incubated for 90 min. Following exposure to serum, cells were grown to an OD_{600} of 1 in
LB medium to enrich for viable mutants. A 1 mL sample normalized to an OD_{600} of 1 from each
culture was processed for genome extraction and multiplexed sequencing using an Illumina
MiSeq. (B) Log_2FC for each gene of the *K. pneumoniae* ECL8 TraDIS library when incubated in
pooled human serum relative to a heat-inactivated serum control. Selected genes highlighted
in red are amongst the total of 144 genes that satisfy a stringent applied threshold ($Log_2FC ≥-$
4, Q-value ≤0.05). For brevity, only genes with a $Log_2FC ≥0$ are illustrated. Inset: transposon
insertion profile of *wbbY*, gene with the highest fold Log_2FC, flanked by *wbbZ* and a
transposable element at its 3'. Transposon insertions following exposure to serum and a heat-
inactivated serum control are illustrated in red and blue, respectively. Transposon reads have
been capped at a maximum of 10. (C) Transposon insertion profiles of genes within the: LPS,
O-antigen and the ECA biosynthesis operons. Genes in red font had a significantly ($Log_2FC =$
≥-4, Q-Value = ≤0.05) decreased fitness when disrupted with a transposon following exposure
to serum for 90 minute (red), relative to a heat-inactivated serum control (blue). Operons are
not drawn to scale and reads capped at a maximum read depth of 1. (D) Growth profile of WT
K. pneumoniae ECL8 and Δ*wbby*::*aph* in LB broth. Mean is plotted (± 1 SD). (E) Serum killing
assay of WT *K. pneumoniae* ECL8 and Δ*wbby*::*aph.* Mean is plotted (± 1 SD). (F) LPS profiles of
WT *K. pneumoniae* ECL8 and Δ*wbby*::*aph*. Overnight cultures of each strain were normalised
to an OD_{600} of 1. The LPS was separated on 4-12% Bis-Tris gels and was visualized by silver
staining using the SilverQuest kit (Invitrogen).

828
Figure 6 Overview and validation of ECL8 genes that increase resistance to complement-
mediated killing. (A) The experimental methodology utilized for screening the TraDIS library
to identify genetic factors that increase resistance to human serum. *K. pneumoniae* ECL8
($2×10^8$ cells) of the mutant library was inoculated into 1 mL of human serum and incubated
for 180 min and compared to before serum exposure input control. The output pool was
washed with PBS and plated onto LB agar supplemented with kanamycin. Following overnight
growth, ~150,000 colonies were recovered and pooled for sequencing. Comparative analysis
using AlbaTraDIS software depicting genes with (B) decreased insertions suggesting a loss of
fitness or (C) increased insertions suggesting a gain of fitness to serum exposure. (D)
Transposon and read count insertion profiles of *hns* locus: red illustrating pooled mutant
serum exposed for 180 min and blue denoting the before serum exposure input control. (E)
Growth profile of WT K. pneumoniae ECL8 and Δ*hns*::*aph* in LB broth. Mean is plotted (± 1
SD), where n=3. (F) Serum killing assay of WT *K. pneumoniae ECL8* and Δ*hns*::*aph*. Mean is
plotted (± 1 SD), where n=3.

Supporting information captions

Figure S1: (A) Pearson correlation coefficient (R2) of gene insertion index scores (IIS) of two sequenced technical replicates of the Klebsiella pneumoniae ECL8 TraDIS library (KTL1 and KTL2). The IIS of genes located on the chromosome and plasmid are highlighted in blue and red respectively. (B) Number of unique reads identified (%) in the raw fastQ file of the *K. pneumoniae* TraDIS library in sequentially larger k-mer pools up to three million reads. Plot generated using BBTools: bbcountunique.sh in non-cumulative mode (https://jgi.doe.gov/data-and-tools/bbtools/)

Figure S2 Sequencing depth of the *K. pneumoniae* ECL8 plasmid (black) relative to the *K. pneumoniae* ECL8 chromosome (green). The plasmid has a read depth 1.33x that of the genome suggesting the plasmid has a copy number of one. Figure generated using Bandage (v3).

Figure S3 The frequency distribution of insertion index scores. The insertion index score for each coding sequence was calculated as the number of insertions per CDS divided by the CDS length in base pairs to normalize for gene length. An exponential distribution model was fitted to the left mode that includes essential genes, and a gamma distribution model was fitted to the right, nonessential mode (blue). For a given insertion index score, the probability of belonging to each mode was calculated, and the ratio of these values was the log likelihood score. A gene was classified as essential if its log likelihood score was less than $\log_2 12$ and was therefore 12 times more likely to belong to the red mode than the blue mode.

Figure S4 The COG Enrichment index comprising 373 genes classified as essential in *K. pneumoniae* ECL8. This index is calculated as the percentage of the essential genome made up of a COG divided by the percentage of the whole genome made up by the same COG. The $\log^2$ fold enrichment is displayed, and significant differences were calculated using the two-tailed Fisher's exact test. Annotations were computed using eggnog-mapper based on eggNOG orthology data (2)

Figure S5 Mathematical simulation (10^5 instances) of random transposon insertion events under the null model of random insertion previously described (3). The probability of at least one insertion free region (IFR) of length (l) occurring in a genome of 5.3 Mb containing 554,834 transposon insertions (blue). Genome length and no. of genome-wide insertions from TraDIS studies by Langridge *et al.,* 2012 and Goodall *et al.,* 2018 plotted for comparison (3, 4).

Figure S6 (A) Sequencing depth required to sample a given proportion of the K. pneumoniae TraDIS library. The following equation: UIPs = $s-s(\frac{s-1}{s})^n$ was applied to calculate the approximate number of sequence reads required to sample full library diversity i.e. 100% of Unique Insertion Points (UIPs). UIPs = Unique Insertion Points, n = number of mapped reads in millions (M) and s = sample size i.e. 499,919 CDS UIPs, as described in Chapter 3. (B) The approximate number of CDS UIPs represented by a given number of sequence reads (M) are shown as a percentage of the total CDS UIPs i.e. 499,919.

Figure S7 The Pearson correlation coefficient (R^2) of gene insertion index scores (IIS) for two sequenced biological replicates of the *K. pneumoniae* ECL8 TraDIS library following thee 12 h passages in (blue) LB broth or (red) pooled human urine. The inline barcode identifiers used to demultiplex and distinguish replicates are highlighted in brackets ()..

Figure S8 Serum killing assay of *K. pneumoniae* ECL8 and *E. coli* BW25113. A sample of 2×10^8 bacterial cells
were incubated in 100 μL of human serum (solid line) or heat-inactivated human serum (dashed line) for 180
min. Viable bacterial numbers (cfu/ml) were sampled at regular time points by plating onto LB agar, overnight
incubation at 37 °C and subsequent counting of the colonies. The $\log_{10}$ fold change in viable cell number was
measured. *E. coli* BW25113 lacks O-antigen and was used as a serum sensitive control. The mean of three
biological replicates is shown ± 1 SD.

Figure S9 The pearson correlation coefficient (R^2) of gene insertion index scores (IIS) for two sequenced
biological replicates of the *K. pneumoniae* ECL8 TraDIS library following 90 min exposure to (blue) Heat-
inactivated serum or (red) Serum. The inline barcodes used to demultiplex and distinguish replicates are given
in brackets ().

Figure S10 Pearson correlation coefficient (R2) of two biological replicates from the output TraDIS library
following exposure to human serum for 180 min.

Figure S11 Directional insertion bias of transposon (Tn) into *gnd*. Transposon insertions configured in the
forward orientation (green), reverse orientation (blue). Transposon insertion densities are capped at a
maximum read depth of 2000. Below – genomic context of *gnd* with a putative promoter for *manC* driving the
transcriptional unit (*rfb* cluster) for O-antigen biosynthesis.

Dataset S1 Essential gene table ECL8

Dataset S2 Insertion Free Regions (IFR) within ECL8 (LB)

Dataset S3 Essential Gene Comparison against ECL8

Dataset S4 Essential gene table ECL8 (Urine)

Dataset S5 Essential gene table ECL8 (Serum)

Dataset S6 Kaptive webserver results - ECL8 O-antigen K-antigen

Dataset S7 AlbaTraDIS ECL8 180 Serum exposure results

925
926
927
928

References

1. Bagley ST. Habitat association of Klebsiella species. Infect Control. 1985;6(2):52-8. doi: 10.1017/s0195941700062603. PubMed PMID: 3882590.

2. Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, Dance D, et al. Genomic analysis of diversity, population structure, virulence, and antimicrobial resistance in Klebsiella pneumoniae, an urgent threat to public health. Proc Natl Acad Sci U S A. 2015;112(27):E3574-81. Epub 20150622. doi: 10.1073/pnas.1501049112. PubMed PMID: 26100894; PubMed Central PMCID: PMCPMC4500264.

3. Podschun R, Ullmann U. Klebsiella spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. Clin Microbiol Rev. 1998;11(4):589-603. doi: 10.1128/CMR.11.4.589. PubMed PMID: 9767057; PubMed Central PMCID: PMCPMC88898.

4. Ko WC, Paterson DL, Sagnimeni AJ, Hansen DS, Von Gottberg A, Mohapatra S, et al. Community-acquired Klebsiella pneumoniae bacteremia: global differences in clinical patterns. Emerg Infect Dis. 2002;8(2):160-6. doi: 10.3201/eid0802.010025. PubMed PMID: 11897067; PubMed Central PMCID: PMCPMC2732457.

5. Goncalves Barbosa LC, Silva ESJA, Bordoni GP, Barbosa GO, Carneiro LC. Elevated Mortality Risk from CRKp Associated with Comorbidities: Systematic Review and Meta-Analysis. Antibiotics (Basel). 2022;11(7). Epub 20220629. doi: 10.3390/antibiotics11070874. PubMed PMID: 35884128; PubMed Central PMCID: PMCPMC9312274.

6. Tumbarello M, Trecarichi EM, De Rosa FG, Giannella M, Giacobbe DR, Bassetti M, et al. Infections caused by KPC-producing Klebsiella pneumoniae: differences in therapy and mortality in a multicentre study. J Antimicrob Chemother. 2015;70(7):2133-43. Epub 20150421. doi: 10.1093/jac/dkv086. PubMed PMID: 25900159.

7. Clegg S, Murphy CN. Epidemiology and Virulence of Klebsiella pneumoniae. Microbiol Spectr. 2016;4(1). doi: 10.1128/microbiolspec.UTI-0005-2012. PubMed PMID: 26999397.

8. Paczosa MK, Mecsas J. Klebsiella pneumoniae: Going on the Offense with a Strong Defense. Microbiol Mol Biol Rev. 2016;80(3):629-61. Epub 20160615. doi: 10.1128/MMBR.00078-15. PubMed PMID: 27307579; PubMed Central PMCID: PMCPMC4981674.

9. Ikuta KS, Swetschinski LR, Robles Aguilar G, Sharara F, Mestrovic T, Gray AP, et al. Global mortality associated with 33 bacterial pathogens in 2019: a systematic analysis for the Global Burden of Disease Study 2019. The Lancet. 2022;400(10369):2221-48. doi: 10.1016/S0140-6736(22)02185-7.

10. Pham MH, Hoi LT, Beale MA, Khokhar FA, Hoa NT, Musicha P, et al. Evidence of widespread endemic populations of highly multidrug resistant Klebsiella pneumoniae in hospital settings in Hanoi, Vietnam: a prospective cohort study. The Lancet Microbe. 2023;4(4):e255-e63. doi: 10.1016/S2666-5247(22)00338-X.

11. Mohd Asri NA, Ahmad S, Mohamud R, Mohd Hanafi N, Mohd Zaidi NF, Irekeola AA, et al. Global Prevalence of Nosocomial Multidrug-Resistant Klebsiella pneumoniae: A Systematic Review and Meta-Analysis. Antibiotics. 2021;10(12):1508. PubMed PMID: doi:10.3390/antibiotics10121508.

12. Navon-Venezia S, Kondratyeva K, Carattoli A. Klebsiella pneumoniae: a major worldwide source and shuttle for antibiotic resistance. FEMS Microbiol Rev. 2017;41(3):252-75. doi: 10.1093/femsre/fux013. PubMed PMID: 28521338.

13. Garbati MA, Al Godhair AI. The growing resistance of Klebsiella pneumoniae; the need to expand our antibiogram: case report and review of the literature. Afr J Infect Dis.

2013;7(1):8-10. doi: 10.4314/ajid.v7i1.2. PubMed PMID: 24381721; PubMed Central
PMCID: PMCPMC3647522.
14.   Campos AC, Albiero J, Ecker AB, Kuroda CM, Meirelles LE, Polato A, et al.
Outbreak of Klebsiella pneumoniae carbapenemase-producing K pneumoniae: A systematic
review. Am J Infect Control. 2016;44(11):1374-80. Epub 20160505. doi:
10.1016/j.ajic.2016.03.022. PubMed PMID: 27156198.
15.   Rojas LJ, Salim M, Cober E, Richter SS, Perez F, Salata RA, et al. Colistin Resistance
in Carbapenem-Resistant Klebsiella pneumoniae: Laboratory Detection and Impact on
Mortality. Clin Infect Dis. 2017;64(6):711-8. doi: 10.1093/cid/ciw805. PubMed PMID:
27940944; PubMed Central PMCID: PMCPMC5850634.
16.   Cain AK, Barquist L, Goodman AL, Paulsen IT, Parkhill J, van Opijnen T. A decade
of advances in transposon-insertion sequencing. Nat Rev Genet. 2020;21(9):526-40. Epub
20200612. doi: 10.1038/s41576-020-0244-x. PubMed PMID: 32533119; PubMed Central
PMCID: PMCPMC7291929.
17.   Barquist L, Mayho M, Cummins C, Cain AK, Boinett CJ, Page AJ, et al. The TraDIS
toolkit: sequencing and analysis for dense transposon mutant libraries. Bioinformatics.
2016;32(7):1109-11. Epub 20160121. doi: 10.1093/bioinformatics/btw022. PubMed PMID:
26794317; PubMed Central PMCID: PMCPMC4896371.
18.   Wyres KL, Lam MMC, Holt KE. Population genomics of Klebsiella pneumoniae. Nat
Rev Microbiol. 2020;18(6):344-59. Epub 20200213. doi: 10.1038/s41579-019-0315-1.
PubMed PMID: 32055025.
19.   Marques C, Menezes J, Belas A, Aboim C, Cavaco-Silva P, Trigueiro G, et al.
Klebsiella pneumoniae causing urinary tract infections in companion animals and humans:
population structure, antimicrobial resistance and virulence genes. J Antimicrob Chemother.
2019;74(3):594-602. doi: 10.1093/jac/dky499. PubMed PMID: 30535393.
20.   Bengoechea JA, Sa Pessoa J. Klebsiella pneumoniae infection biology: living to
counteract host defences. FEMS Microbiol Rev. 2019;43(2):123-44. doi:
10.1093/femsre/fuy043. PubMed PMID: 30452654; PubMed Central PMCID:
PMCPMC6435446.
21.   Choi M, Hegerle N, Nkeze J, Sen S, Jamindar S, Nasrin S, et al. The Diversity of
Lipopolysaccharide (O) and Capsular Polysaccharide (K) Antigens of Invasive Klebsiella
pneumoniae in a Multi-Country Collection. Front Microbiol. 2020;11:1249. Epub 20200612.
doi: 10.3389/fmicb.2020.01249. PubMed PMID: 32595624; PubMed Central PMCID:
PMCPMC7303279.
22.   Martin RM, Bachman MA. Colonization, Infection, and the Accessory Genome of
Klebsiella pneumoniae. Front Cell Infect Microbiol. 2018;8:4. Epub 20180122. doi:
10.3389/fcimb.2018.00004. PubMed PMID: 29404282; PubMed Central PMCID:
PMCPMC5786545.
23.   Short FL, Di Sario G, Reichmann NT, Kleanthous C, Parkhill J, Taylor PW. Genomic
Profiling Reveals Distinct Routes To Complement Resistance in Klebsiella pneumoniae.
Infect Immun. 2020;88(8). Epub 20200721. doi: 10.1128/IAI.00043-20. PubMed PMID:
32513855; PubMed Central PMCID: PMCPMC7375759.
24.   Phan MD, Peters KM, Sarkar S, Lukowski SW, Allsopp LP, Gomes Moriel D, et al.
The serum resistome of a globally disseminated multidrug resistant uropathogenic
Escherichia coli clone. PLoS Genet. 2013;9(10):e1003834. Epub 20131003. doi:
10.1371/journal.pgen.1003834. PubMed PMID: 24098145; PubMed Central PMCID:
PMCPMC3789825.
25.   McCarthy AJ, Taylor PW. Analysis of Escherichia coli K1 Virulence Genes by
Transposon-Directed Sequencing. Methods Mol Biol. 2022;2377:199-213. doi: 10.1007/978-
1-0716-1720-5_11. PubMed PMID: 34709618.

26. Ondari EM, Klemm EJ, Msefula CL, El Ghany MA, Heath JN, Pickard DJ, et al.
Rapid transcriptional responses to serum exposure are associated with sensitivity and
resistance to antibody-mediated complement killing in invasive Salmonella Typhimurium
ST313. Wellcome Open Res. 2019;4:74. Epub 20190425. doi:
10.12688/wellcomeopenres.15059.1. PubMed PMID: 31231691; PubMed Central PMCID:
PMCPMC6560496.
27. Subashchandrabose S, Smith S, DeOrnellas V, Crepin S, Kole M, Zahdeh C, et al.
Acinetobacter baumannii Genes Required for Bacterial Survival during Bloodstream
Infection. mSphere. 2016;1(1). Epub 20151104. doi: 10.1128/mSphere.00013-15. PubMed
PMID: 27303682; PubMed Central PMCID: PMCPMC4863628.
28. Goodall ECA, Robinson A, Johnston IG, Jabbari S, Turner KA, Cunningham AF, et
al. The Essential Genome of Escherichia coli K-12. mBio. 2018;9(1). Epub 20180220. doi:
10.1128/mBio.02096-17. PubMed PMID: 29463657; PubMed Central PMCID:
PMCPMC5821084.
29. Fabian BK, Foster C, Asher AJ, Elbourne LDH, Cain AK, Hassan KA, et al.
Elucidating Essential Genes in Plant-Associated Pseudomonas protegens Pf-5 Using
Transposon Insertion Sequencing. J Bacteriol. 2021;203(7). Epub 20210308. doi:
10.1128/JB.00432-20. PubMed PMID: 33257523; PubMed Central PMCID:
PMCPMC8088517.
30. Wong YC, Abd El Ghany M, Naeem R, Lee KW, Tan YC, Pain A, et al. Candidate
Essential Genes in Burkholderia cenocepacia J2315 Identified by Genome-Wide TraDIS.
Front Microbiol. 2016;7:1288. Epub 20160822. doi: 10.3389/fmicb.2016.01288. PubMed
PMID: 27597847; PubMed Central PMCID: PMCPMC4993015.
31. Ramage B, Erolin R, Held K, Gasper J, Weiss E, Brittnacher M, et al. Comprehensive
Arrayed Transposon Mutant Library of Klebsiella pneumoniae Outbreak Strain KPNIH1. J
Bacteriol. 2017;199(20). Epub 20170919. doi: 10.1128/JB.00352-17. PubMed PMID:
28760848; PubMed Central PMCID: PMCPMC5637181.
32. DeJesus MA, Gerrick ER, Xu W, Park SW, Long JE, Boutte CC, et al.
Comprehensive Essentiality Analysis of the Mycobacterium tuberculosis Genome via
Saturating Transposon Mutagenesis. mBio. 2017;8(1). Epub 20170117. doi:
10.1128/mBio.02133-16. PubMed PMID: 28096490; PubMed Central PMCID:
PMCPMC5241402.
33. Ebert M, Laass S, Burghartz M, Petersen J, Kossmehl S, Wohlbrand L, et al.
Transposon mutagenesis identified chromosomal and plasmid genes essential for adaptation
of the marine bacterium Dinoroseobacter shibae to anaerobic conditions. J Bacteriol.
2013;195(20):4769-77. Epub 20130823. doi: 10.1128/JB.00860-13. PubMed PMID:
23974024; PubMed Central PMCID: PMCPMC3807425.
34. Hardy L, Juan PA, Coupat-Goutaland B, Charpentier X. Transposon Insertion
Sequencing in a Clinical Isolate of Legionella pneumophila Identifies Essential Genes and
Determinants of Natural Transformation. J Bacteriol. 2021;203(3). Epub 20210111. doi:
10.1128/JB.00548-20. PubMed PMID: 33168636; PubMed Central PMCID:
PMCPMC7811196.
35. Armbruster CE, Forsyth-DeOrnellas V, Johnson AO, Smith SN, Zhao L, Wu W, et al.
Genome-wide transposon mutagenesis of Proteus mirabilis: Essential genes, fitness factors
for catheter-associated urinary tract infection, and the impact of polymicrobial infection on
fitness requirements. PLoS Pathog. 2017;13(6):e1006434. Epub 20170614. doi:
10.1371/journal.ppat.1006434. PubMed PMID: 28614382; PubMed Central PMCID:
PMCPMC5484520.
36. Roth M, Goodall ECA, Pullela K, Jaquet V, François P, Henderson IR, et al.
Transposon-Directed Insertion-Site Sequencing Reveals Glycolysis Gene gpmA as Part of the

H(2)O(2) Defense Mechanisms in Escherichia coli. Antioxidants (Basel). 2022;11(10). Epub
20221018. doi: 10.3390/antiox11102053. PubMed PMID: 36290776; PubMed Central
PMCID: PMCPMC9598634.
37. Winkle M, Hernández-Rocamora VM, Pullela K, Goodall ECA, Martorana AM, Gray
J, et al. DpaA Detaches Braun's Lipoprotein from Peptidoglycan. mBio. 2021;12(3). Epub
20210504. doi: 10.1128/mBio.00836-21. PubMed PMID: 33947763; PubMed Central
PMCID: PMCPMC8263019.
38. Goodall ECA, Isom GL, Rooke JL, Pullela K, Icke C, Yang Z, et al. Loss of YhcB
results in dysregulation of coordinated peptidoglycan, LPS and phospholipid synthesis during
Escherichia coli cell growth. PLOS Genetics. 2021;17(12):e1009586. doi:
10.1371/journal.pgen.1009586.
39. Hoashi K, Harada S, Ishii Y, Aoki K, Ishikawa S, Oshiro Y, et al. Community-
acquired liver abscess caused by capsular genotype K2-ST375 hypervirulent Klebsiella
pneumoniae isolates. IDCases. 2019;17:e00577. Epub 20190614. doi:
10.1016/j.idcr.2019.e00577. PubMed PMID: 31293894; PubMed Central PMCID:
PMCPMC6595073.
40. Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, et al.
CLIMB (the Cloud Infrastructure for Microbial Bioinformatics): an online resource for the
medical microbiology community. Microb Genom. 2016;2(9):e000086. Epub 20160920. doi:
10.1099/mgen.0.000086. PubMed PMID: 28785418; PubMed Central PMCID:
PMCPMC5537631.
41. Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, et al.
Artemis: sequence visualization and annotation. Bioinformatics. 2000;16(10):944-5. doi:
10.1093/bioinformatics/16.10.944. PubMed PMID: 11120685.
42. Langridge GC, Phan MD, Turner DJ, Perkins TT, Parts L, Haase J, et al.
Simultaneous assay of every Salmonella Typhi gene using one million transposon mutants.
Genome Res. 2009;19(12):2308-16. Epub 20091013. doi: 10.1101/gr.097097.109. PubMed
PMID: 19826075; PubMed Central PMCID: PMCPMC2792183.
43. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for
FASTA/Q File Manipulation. PLoS One. 2016;11(10):e0163962. Epub 20161005. doi:
10.1371/journal.pone.0163962. PubMed PMID: 27706213; PubMed Central PMCID:
PMCPMC5051824.
44. Cock PJ, Chilton JM, Gruning B, Johnson JE, Soranzo N. NCBI BLAST+ integrated
into Galaxy. Gigascience. 2015;4:39. Epub 20150825. doi: 10.1186/s13742-015-0080-7.
PubMed PMID: 26336600; PubMed Central PMCID: PMCPMC4557756.
45. Page AJ, Bastkowski S, Yasir M, Turner AK, Le Viet T, Savva GM, et al.
AlbaTraDIS: Comparative analysis of large datasets from parallel transposon mutagenesis
experiments. PLoS Comput Biol. 2020;16(7):e1007980. Epub 20200717. doi:
10.1371/journal.pcbi.1007980. PubMed PMID: 32678849; PubMed Central PMCID:
PMCPMC7390408.
46. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A. 2000;97(12):6640-5.
doi: 10.1073/pnas.120163297. PubMed PMID: 10829079; PubMed Central PMCID:
PMCPMC18686.
47. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de
novo genome assemblies. Bioinformatics. 2015;31(20):3350-2. Epub 20150622. doi:
10.1093/bioinformatics/btv383. PubMed PMID: 26099265; PubMed Central PMCID:
PMCPMC4595904.
48. Huerta-Cepas J, Szklarczyk D, Heller D, Hernandez-Plaza A, Forslund SK, Cook H,
et al. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology

resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res. 2019;47(D1):D309-D14. doi: 10.1093/nar/gky1085. PubMed PMID: 30418610; PubMed Central PMCID: PMCPMC6324079.

49.   Christen B, Abeliuk E, Collier JM, Kalogeraki VS, Passarelli B, Coller JA, et al. The essential genome of a bacterium. Mol Syst Biol. 2011;7:528. Epub 20110830. doi: 10.1038/msb.2011.58. PubMed PMID: 21878915; PubMed Central PMCID: PMCPMC3202797.

50.   Grazziotin AL, Vidal NM, Venancio TM. Uncovering major genomic features of essential genes in Bacteria and a methanogenic Archaea. FEBS J. 2015;282(17):3395-411. Epub 20150714. doi: 10.1111/febs.13350. PubMed PMID: 26084810; PubMed Central PMCID: PMCPMC4558319.

51.   Barquist L, Langridge GC, Turner DJ, Phan MD, Turner AK, Bateman A, et al. A comparison of dense transposon insertion libraries in the Salmonella serovars Typhi and Typhimurium. Nucleic Acids Res. 2013;41(8):4549-64. Epub 20130306. doi: 10.1093/nar/gkt148. PubMed PMID: 23470992; PubMed Central PMCID: PMCPMC3632133.

52.   Tang HL, Chiang MK, Liou WJ, Chen YT, Peng HL, Chiou CS, et al. Correlation between Klebsiella pneumoniae carrying pLVPK-derived loci and abscess formation. Eur J Clin Microbiol Infect Dis. 2010;29(6):689-98. Epub 20100411. doi: 10.1007/s10096-010-0915-1. PubMed PMID: 20383552.

53.   Rodriguez-Medina N, Martinez-Romero E, De la Cruz MA, Ares MA, Valdovinos-Torres H, Silva-Sanchez J, et al. A Klebsiella variicola Plasmid Confers Hypermucoviscosity-Like Phenotype and Alters Capsule Production and Virulence. Front Microbiol. 2020;11:579612. Epub 20201216. doi: 10.3389/fmicb.2020.579612. PubMed PMID: 33391198; PubMed Central PMCID: PMCPMC7772424.

54.   Yang X, Dong N, Liu X, Yang C, Ye L, Chan EW, et al. Co-conjugation of Virulence Plasmid and KPC Plasmid in a Clinical Klebsiella pneumoniae Strain. Front Microbiol. 2021;12:739461. Epub 20211108. doi: 10.3389/fmicb.2021.739461. PubMed PMID: 34819921; PubMed Central PMCID: PMCPMC8606748.

55.   Joyce CM, Grindley ND. Method for determining whether a gene of Escherichia coli is essential: application to the polA gene. J Bacteriol. 1984;158(2):636-43. doi: 10.1128/jb.158.2.636-643.1984. PubMed PMID: 6233260; PubMed Central PMCID: PMCPMC215477.

56.   Gallagher LA, Bailey J, Manoil C. Ranking essential bacterial processes by speed of mutant death. Proc Natl Acad Sci U S A. 2020;117(30):18010-7. Epub 20200714. doi: 10.1073/pnas.2001507117. PubMed PMID: 32665440; PubMed Central PMCID: PMCPMC7395459.

57.   Warren AS, Archuleta J, Feng WC, Setubal JC. Missing genes in the annotation of prokaryotic genomes. BMC Bioinformatics. 2010;11(1):131. Epub 20100315. doi: 10.1186/1471-2105-11-131. PubMed PMID: 20230630; PubMed Central PMCID: PMCPMC3098052.

58.   Mat-Sharani S, Firdaus-Raih M. Computational discovery and annotation of conserved small open reading frames in fungal genomes. BMC Bioinformatics. 2019;19(Suppl 13):551. Epub 20190204. doi: 10.1186/s12859-018-2550-2. PubMed PMID: 30717662; PubMed Central PMCID: PMCPMC7394265.

59.   Miravet-Verde S, Ferrar T, Espadas-Garcia G, Mazzolini R, Gharrab A, Sabido E, et al. Unraveling the hidden universe of small proteins in bacterial genomes. Mol Syst Biol. 2019;15(2):e8290. Epub 20190222. doi: 10.15252/msb.20188290. PubMed PMID: 30796087; PubMed Central PMCID: PMCPMC6385055.

60.   Hutchison CA, 3rd, Merryman C, Sun L, Assad-Garcia N, Richter RA, Smith HO, et
al. Polar Effects of Transposon Insertion into a Minimal Bacterial Genome. J Bacteriol.
2019;201(19). Epub 20190906. doi: 10.1128/JB.00185-19. PubMed PMID: 31262838;
PubMed Central PMCID: PMCPMC6755753.
61.   Lam MMC, Wick RR, Judd LM, Holt KE, Wyres KL. Kaptive 2.0: updated capsule
and lipopolysaccharide locus typing for the Klebsiella pneumoniae species complex. Microb
Genom. 2022;8(3). doi: 10.1099/mgen.0.000800. PubMed PMID: 35311639; PubMed
Central PMCID: PMCPMC9176290.
62.   Pain A, Ott A, Amine H, Rochat T, Bouloc P, Gautheret D. An assessment of
bacterial small RNA target prediction programs. RNA Biol. 2015;12(5):509-13. doi:
10.1080/15476286.2015.1020269. PubMed PMID: 25760244; PubMed Central PMCID:
PMCPMC4615726.
63.   Waters LS, Storz G. Regulatory RNAs in bacteria. Cell. 2009;136(4):615-28. doi:
10.1016/j.cell.2009.01.043. PubMed PMID: 19239884; PubMed Central PMCID:
PMCPMC3132550.
64.   Jana B, Cain AK, Doerrler WT, Boinett CJ, Fookes MC, Parkhill J, et al. The
secondary resistome of multidrug-resistant Klebsiella pneumoniae. Sci Rep. 2017;7:42483.
Epub 20170215. doi: 10.1038/srep42483. PubMed PMID: 28198411; PubMed Central
PMCID: PMCPMC5309761.
65.   Dorman MJ, Feltwell T, Goulding DA, Parkhill J, Short FL. The Capsule Regulatory
Network of Klebsiella pneumoniae Defined by density-TraDISort. mBio. 2018;9(6). Epub
20181120. doi: 10.1128/mBio.01863-18. PubMed PMID: 30459193; PubMed Central
PMCID: PMCPMC6247091.
66.   Jung HJ, Littmann ER, Seok R, Leiner IM, Taur Y, Peled J, et al. Genome-Wide
Screening for Enteric Colonization Factors in Carbapenem-Resistant ST258 Klebsiella
pneumoniae. mBio. 2019;10(2). Epub 20190312. doi: 10.1128/mBio.02663-18. PubMed
PMID: 30862751; PubMed Central PMCID: PMCPMC6414703.
67.   Paczosa MK, Silver RJ, McCabe AL, Tai AK, McLeish CH, Lazinski DW, et al.
Transposon Mutagenesis Screen of Klebsiella pneumoniae Identifies Multiple Genes
Important for Resisting Antimicrobial Activities of Neutrophils in Mice. Infect Immun.
2020;88(4). Epub 20200323. doi: 10.1128/IAI.00034-20. PubMed PMID: 31988174;
PubMed Central PMCID: PMCPMC7093148.
68.   Ma P, He LL, Pironti A, Laibinis HH, Ernst CM, Manson AL, et al. Genetic
determinants facilitating the evolution of resistance to carbapenem antibiotics. Elife. 2021;10.
Epub 20210419. doi: 10.7554/eLife.67310. PubMed PMID: 33871353; PubMed Central
PMCID: PMCPMC8079144.
69.   Bruchmann S, Feltwell T, Parkhill J, Short FL. Identifying virulence determinants of
multidrug-resistant Klebsiella pneumoniae in Galleria mellonella. Pathog Dis. 2021;79(3).
doi: 10.1093/femspd/ftab009. PubMed PMID: 33512418; PubMed Central PMCID:
PMCPMC7981267.
70.   Chao MC, Abel S, Davis BM, Waldor MK. The design and analysis of transposon
insertion sequencing experiments. Nat Rev Microbiol. 2016;14(2):119-28. doi:
10.1038/nrmicro.2015.7. PubMed PMID: 26775926; PubMed Central PMCID:
PMCPMC5099075.
71.   Lampe DJ, Churchill ME, Robertson HM. A purified mariner transposase is sufficient
to mediate transposition in vitro. EMBO J. 1996;15(19):5470-9. PubMed PMID: 8895590;
PubMed Central PMCID: PMCPMC452289.
72.   Mas G, Thoma J, Hiller S. The Periplasmic Chaperones Skp and SurA. Subcell
Biochem. 2019;92:169-86. doi: 10.1007/978-3-030-18768-2_6. PubMed PMID: 31214987.

73. Caza M, Kronstad JW. Shared and distinct mechanisms of iron acquisition by
bacterial and fungal pathogens of humans. Front Cell Infect Microbiol. 2013;3:80. Epub
20131119. doi: 10.3389/fcimb.2013.00080. PubMed PMID: 24312900; PubMed Central
PMCID: PMCPMC3832793.
74. Iwadate Y, Ramezanifard R, Golubeva YA, Fenlon LA, Slauch JM. PaeA (YtfL)
protects from cadaverine and putrescine stress in Salmonella Typhimurium and E. coli. Mol
Microbiol. 2021;115(6):1379-94. Epub 20210203. doi: 10.1111/mmi.14686. PubMed PMID:
33481283.
75. Xu Y, Liang Z, Legrain C, Ruger HJ, Glansdorff N. Evolution of arginine
biosynthesis in the bacterial domain: novel gene-enzyme relationships from psychrophilic
Moritella strains (Vibrionaceae) and evolutionary significance of N-alpha-acetyl ornithinase.
J Bacteriol. 2000;182(6):1609-15. doi: 10.1128/JB.182.6.1609-1615.2000. PubMed PMID:
10692366; PubMed Central PMCID: PMCPMC94458.
76. Chambers ST, Lever M. Betaines and urinary tract infections. Nephron. 1996;74(1):1-
10. doi: 10.1159/000189274. PubMed PMID: 8883013.
77. Chaturvedi KS, Hung CS, Crowley JR, Stapleton AE, Henderson JP. The siderophore
yersiniabactin binds copper to protect pathogens during infection. Nat Chem Biol.
2012;8(8):731-6. Epub 20120708. doi: 10.1038/nchembio.1020. PubMed PMID: 22772152;
PubMed Central PMCID: PMCPMC3600419.
78. Cassat JE, Skaar EP. Iron in infection and immunity. Cell Host Microbe.
2013;13(5):509-19. doi: 10.1016/j.chom.2013.04.010. PubMed PMID: 23684303; PubMed
Central PMCID: PMCPMC3676888.
79. Banerjee R, Weisenhorn E, Schwartz KJ, Myers KS, Glasner JD, Perna NT, et al.
Tailoring a Global Iron Regulon to a Uropathogen. mBio. 2020;11(2). Epub 20200324. doi:
10.1128/mBio.00351-20. PubMed PMID: 32209682; PubMed Central PMCID:
PMCPMC7157518.
80. Huang SH, Wang CK, Peng HL, Wu CC, Chen YT, Hong YM, et al. Role of the
small RNA RyhB in the Fur regulon in mediating the capsular polysaccharide biosynthesis
and iron acquisition systems in Klebsiella pneumoniae. BMC Microbiol. 2012;12(1):148.
Epub 20120724. doi: 10.1186/1471-2180-12-148. PubMed PMID: 22827802; PubMed
Central PMCID: PMCPMC3423075.
81. Troxell B, Hassan HM. Transcriptional regulation by Ferric Uptake Regulator (Fur) in
pathogenic bacteria. Front Cell Infect Microbiol. 2013;3:59. Epub 20131002. doi:
10.3389/fcimb.2013.00059. PubMed PMID: 24106689; PubMed Central PMCID:
PMCPMC3788343.
82. Seo SW, Kim D, Latif H, O'Brien EJ, Szubin R, Palsson BO. Deciphering Fur
transcriptional regulatory network highlights its complex role beyond iron metabolism in
Escherichia coli. Nat Commun. 2014;5(1):4910. Epub 20140915. doi: 10.1038/ncomms5910.
PubMed PMID: 25222563; PubMed Central PMCID: PMCPMC4167408.
83. Doorduijn DJ, Heesterbeek DAC, Ruyken M, de Haas CJC, Stapels DAC, Aerts PC,
et al. Polymerization of C9 enhances bacterial cell envelope damage and killing by membrane
attack complex pores. PLOS Pathogens. 2021;17(11):e1010051. doi:
10.1371/journal.ppat.1010051.
84. Onsare RS, Micoli F, Lanzilao L, Alfini R, Okoro CK, Muigai AW, et al.
Relationship between antibody susceptibility and lipopolysaccharide O-antigen
characteristics of invasive and gastrointestinal nontyphoidal Salmonellae isolates from
Kenya. PLoS Negl Trop Dis. 2015;9(3):e0003573. Epub 20150304. doi:
10.1371/journal.pntd.0003573. PubMed PMID: 25739091; PubMed Central PMCID:
PMCPMC4352093.

85.   Salamon H, Nissim-Eliraz E, Ardronai O, Nissan I, Shpigel NY. The role of O-
polysaccharide chain and complement resistance of Escherichia coli in mammary virulence.
Vet Res. 2020;51(1):77. Epub 20200615. doi: 10.1186/s13567-020-00804-x. PubMed PMID:
32539761; PubMed Central PMCID: PMCPMC7294653.
86.   Rai AK, Mitchell AM. Enterobacterial Common Antigen: Synthesis and Function of
an Enigmatic Molecule. mBio. 2020;11(4). Epub 20200811. doi: 10.1128/mBio.01914-20.
PubMed PMID: 32788387; PubMed Central PMCID: PMCPMC7439462.
87.   Frirdich E, Whitfield C. Characterization of Gla(KP), a UDP-galacturonic acid C4-
epimerase from Klebsiella pneumoniae with extended substrate specificity. J Bacteriol.
2005;187(12):4104-15. doi: 10.1128/JB.187.12.4104-4115.2005. PubMed PMID: 15937173;
PubMed Central PMCID: PMCPMC1151718.
88.   Ramos PI, Picao RC, Vespero EC, Pelisson M, Zuleta LF, Almeida LG, et al.
Pyrosequencing-based analysis reveals a novel capsular gene cluster in a KPC-producing
Klebsiella pneumoniae clinical isolate identified in Brazil. BMC Microbiol. 2012;12(1):173.
Epub 20120811. doi: 10.1186/1471-2180-12-173. PubMed PMID: 22882772; PubMed
Central PMCID: PMCPMC3438125.
89.   Yeow J, Tan KW, Holdbrook DA, Chong ZS, Marzinek JK, Bond PJ, et al. The
architecture of the OmpC-MlaA complex sheds light on the maintenance of outer membrane
lipid asymmetry in Escherichia coli. J Biol Chem. 2018;293(29):11325-40. Epub 20180530.
doi: 10.1074/jbc.RA118.002441. PubMed PMID: 29848551; PubMed Central PMCID:
PMCPMC6065193.
90.   Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CR, Guan Z. Discovery of a
cardiolipin synthase utilizing phosphatidylethanolamine and phosphatidylglycerol as
substrates. Proc Natl Acad Sci U S A. 2012;109(41):16504-9. Epub 20120917. doi:
10.1073/pnas.1212797109. PubMed PMID: 22988102; PubMed Central PMCID:
PMCPMC3478633.
91.   Rigel NW, Schwalm J, Ricci DP, Silhavy TJ. BamE modulates the Escherichia coli
beta-barrel assembly machine component BamA. J Bacteriol. 2012;194(5):1002-8. Epub
20111216. doi: 10.1128/JB.06426-11. PubMed PMID: 22178970; PubMed Central PMCID:
PMCPMC3294770.
92.   Nguyen D, Kelly K, Qiu N, Misra R. YejM Controls LpxC Levels by Regulating
Protease Activity of the FtsH/YciM Complex of Escherichia coli. J Bacteriol. 2020;202(18).
Epub 20200825. doi: 10.1128/JB.00303-20. PubMed PMID: 32540932; PubMed Central
PMCID: PMCPMC7925073.
93.   Hsieh PF, Wu MC, Yang FL, Chen CT, Lou TC, Chen YY, et al. D-galactan II is an
immunodominant antigen in O1 lipopolysaccharide and affects virulence in Klebsiella
pneumoniae: implication in vaccine design. Front Microbiol. 2014;5:608. Epub 20141119.
doi: 10.3389/fmicb.2014.00608. PubMed PMID: 25477867; PubMed Central PMCID:
PMCPMC4237132.
94.   Ares MA, Fernández-Vázquez JL, Rosales-Reyes R, Jarillo-Quijada MD, von Bargen
K, Torres J, et al. H-NS Nucleoid Protein Controls Virulence Features of Klebsiella
pneumoniae by Regulating the Expression of Type 3 Pili and the Capsule Polysaccharide.
Front Cell Infect Microbiol. 2016;6:13. Epub 20160209. doi: 10.3389/fcimb.2016.00013.
PubMed PMID: 26904512; PubMed Central PMCID: PMCPMC4746245.
95.   Sledjeski D, Gottesman S. A small RNA acts as an antisilencer of the H-NS-silenced
rcsA gene of Escherichia coli. Proc Natl Acad Sci U S A. 1995;92(6):2003-7. doi:
10.1073/pnas.92.6.2003. PubMed PMID: 7534408; PubMed Central PMCID:
PMCPMC42411.
96.   Ebel W, Trempy JE. Escherichia coli RcsA, a positive activator of colanic acid
capsular polysaccharide synthesis, functions To activate its own expression. J Bacteriol.

1999;181(2):577-84. doi: 10.1128/jb.181.2.577-584.1999. PubMed PMID: 9882673; PubMed
Central PMCID: PMCPMC93413.
97.   Meng J, Young G, Chen J. The Rcs System in Enterobacteriaceae: Envelope Stress
Responses and Virulence Regulation. Frontiers in Microbiology. 2021;12. doi:
10.3389/fmicb.2021.627104.
98.   Follador R, Heinz E, Wyres KL, Ellington MJ, Kowarik M, Holt KE, et al. The
diversity of Klebsiella pneumoniae surface polysaccharides. Microbial Genomics. 2016;2(8).
doi: https://doi.org/10.1099/mgen.0.000073.
99.   Butaite E, Baumgartner M, Wyder S, Kummerli R. Siderophore cheating and cheating
resistance shape competition for iron in soil and freshwater Pseudomonas communities. Nat
Commun. 2017;8(1):414. Epub 20170904. doi: 10.1038/s41467-017-00509-4. PubMed
PMID: 28871205; PubMed Central PMCID: PMCPMC5583256.
100.   Holden VI, Bachman MA. Diverging roles of bacterial siderophores during infection.
Metallomics. 2015;7(6):986-95. doi: 10.1039/c4mt00333k. PubMed PMID: 25745886.
101.   Clarke BR, Whitfield C. Molecular cloning of the rfb region of Klebsiella
pneumoniae serotype O1:K20: the rfb gene cluster is responsible for synthesis of the D-
galactan I O polysaccharide. J Bacteriol. 1992;174(14):4614-21. doi:
10.1128/jb.174.14.4614-4621.1992. PubMed PMID: 1378055; PubMed Central PMCID:
PMCPMC206256.
102.   Touati D. Cloning and mapping of the manganese superoxide dismutase gene (sodA)
of Escherichia coli K-12. J Bacteriol. 1983;155(3):1078-87. doi: 10.1128/jb.155.3.1078-
1087.1983. PubMed PMID: 6309739; PubMed Central PMCID: PMCPMC217801.
103.   Doukyu N, Taguchi K. Involvement of catalase and superoxide dismutase in
hydrophobic organic solvent tolerance of Escherichia coli. AMB Express. 2021;11(1):97.
Epub 20210629. doi: 10.1186/s13568-021-01258-w. PubMed PMID: 34189628; PubMed
Central PMCID: PMCPMC8241964.
104.   Najmuldeen H, Alghamdi R, Alghofaili F, Yesilkaya H. Functional assessment of
microbial superoxide dismutase isozymes suggests a differential role for each isozyme. Free
Radic Biol Med. 2019;134:215-28. Epub 20190115. doi:
10.1016/j.freeradbiomed.2019.01.018. PubMed PMID: 30658083.
105.   Flores-Mireles AL, Walker JN, Caparon M, Hultgren SJ. Urinary tract infections:
epidemiology, mechanisms of infection and treatment options. Nat Rev Microbiol.
2015;13(5):269-84. Epub 20150408. doi: 10.1038/nrmicro3432. PubMed PMID: 25853778;
PubMed Central PMCID: PMCPMC4457377.
106.   Murdoch CC, Skaar EP. Nutritional immunity: the battle for nutrient metals at the
host-pathogen interface. Nat Rev Microbiol. 2022;20(11):657-70. Epub 20220531. doi:
10.1038/s41579-022-00745-6. PubMed PMID: 35641670; PubMed Central PMCID:
PMCPMC9153222.
107.   Eberly AR, Floyd KA, Beebout CJ, Colling SJ, Fitzgerald MJ, Stratton CW, et al.
Biofilm Formation by Uropathogenic Escherichia coli Is Favored under Oxygen Conditions
That Mimic the Bladder Environment. Int J Mol Sci. 2017;18(10). Epub 20170930. doi:
10.3390/ijms18102077. PubMed PMID: 28973965; PubMed Central PMCID:
PMCPMC5666759.
108.   Cornelis P, Wei Q, Andrews SC, Vinckx T. Iron homeostasis and management of
oxidative stress response in bacteria. Metallomics. 2011;3(6):540-9. Epub 20110513. doi:
10.1039/c1mt00022e. PubMed PMID: 21566833.
109.   Adler C, Corbalan NS, Peralta DR, Pomares MF, de Cristobal RE, Vincent PA. The
alternative role of enterobactin as an oxidative stress protector allows Escherichia coli colony
development. PLoS One. 2014;9(1):e84734. Epub 20140102. doi:

10.1371/journal.pone.0084734. PubMed PMID: 24392154; PubMed Central PMCID:
PMCPMC3879343.
110.   Peralta DR, Adler C, Corbalan NS, Paz Garcia EC, Pomares MF, Vincent PA.
Enterobactin as Part of the Oxidative Stress Response Repertoire. PLoS One.
2016;11(6):e0157799. Epub 20160616. doi: 10.1371/journal.pone.0157799. PubMed PMID:
27310257; PubMed Central PMCID: PMCPMC4911079.
111.   Guest RL, Court EA, Waldon JL, Schock KA, Raivio TL. Impaired Efflux of the
Siderophore Enterobactin Induces Envelope Stress in Escherichia coli. Front Microbiol.
2019;10:2776. Epub 20191206. doi: 10.3389/fmicb.2019.02776. PubMed PMID: 31866967;
PubMed Central PMCID: PMCPMC6908949.
112.   Singh S, Wilksch JJ, Dunstan RA, Mularski A, Wang N, Hocking D, et al. LPS O
Antigen Plays a Key Role in Klebsiella pneumoniae Capsule Retention. Microbiology
Spectrum. 2022;10(4):e01517-21. doi: doi:10.1128/spectrum.01517-21.
1389
1390