Neuronal expression of human amyloid-β and Tau drives global phenotypic and multi-omic changes in *C. elegans*

Angelina Holcom[1,2,#], Matias Fuentealba[1,#], Renuka Sivapatham[1], Christina D. King[1], Hadley Osman[1], Anna Foulger[1], Dipa Bhaumik[1], Birgit Schilling[1], David Furman[1], Julie K. Andersen[1*], Gordon J. Lithgow[1*]

[1]Buck Institute for Research on Aging; Novato, California, 94945 USA

[2]Leonard Davis School of Gerontology, University of Southern California. Los Angeles, California, 90089, USA

*Corresponding author

#Co-first authors

Corresponding authors:

glithgow@buckinstitute.org

jandersen@buckinstitute.org

Abstract

Alzheimer's disease (AD) and Alzheimer's related diseases (ADRD) are prevalent age-related neurodegenerative disorders characterized by the accumulation of amyloid-β (Aβ) plaques and Tau neurofibrillary tangles. The nematode *Caenorhabditis elegans* (*C. elegans*) serves as an invaluable model organism in diseases of old age-due to its rapid aging. Here we performed an unbiased systems analysis of a *C. elegans* strain expressing both Aβ and Tau proteins within neurons. We set out to determine if there was a phenotypic interaction between Aβ and Tau. In addition, we were interested in determining the temporal order of the phenotypic and multi-omic (geromic) outcomes. At an early stage of adulthood, we observed reproductive impairments and mitochondrial dysfunction consistent with disruptions in mRNA transcript abundance, protein solubility, and metabolite levels. Notably, the expression of these neurotoxic proteins exhibited a synergistic effect, leading to accelerated aging. Our findings shed light on the close relationship between normal aging and ADRD. Specifically, we demonstrate alterations to metabolic functions preceding age-related neurotoxicity, offering a resource for the development of new therapeutic strategies.

Main

Alzheimer's disease (AD) and Alzheimer's related diseases (ADRD) represent the most prevalent forms of dementia, affecting millions of individuals worldwide[1]. The greatest risk factor for ADRD is aging. Pathological hallmarks of these disorders include the formation of amyloid-β (Aβ) plaques, neurofibrillary tangles composed of hyperphosphorylated Tau protein, glial cell activation, and impairment of the blood-brain barrier.

The interaction between Aβ and Tau in AD has been closely linked to hypometabolism in the posterior cingulate cortex, which precedes memory decline[2]. Aβ presence accelerates Tau oligomerization, hyperphosphorylation, and propagation[2–4]. In mouse models, co-expression of Aβ and Tau leads to early synapse loss and synergistic effects on defects in mitochondrial function and turnover[2]. Patients with both Aβ and Tau, as determined by cerebrospinal fluid (CSF) analysis and MRI/PET methods, experience an amplified rate of cognitive decline compared to individuals with only Aβ, only Tau, or neither[5].

Extensive research has focused on finding disease-modifying treatments for AD, including inhibition of Aβ production by targeting the cleavage of APP by β and γ secretase, along with immunotherapy, and inhibitors of Aβ aggregation. However, these approaches have not succeeded in halting disease progression[6]. Consequently, there is need to further explore therapeutic targets, such as targeting multiple factors of the disease concurrently. A potential target could be centered around interactions between Aβ and Tau, instead of focusing on the individual proteins associated with the disease process.

We investigated the multi-omic (geromic) consequences of pan-neuronal, simultaneous
expression of human Aβ and Tau in the model organism *C. elegans* leveraging its short
20-day lifespan. In previous studies transgenic *C. elegans* models have expressed
human Aβ in the body muscle or neurons, resulting in a range of phenotypes[7–9].
Similarly, investigations of Tau expression in *C. elegans* have focused on expression of
human Tau containing mutations associated with frontotemporal dementia (FTD), a
disorder related to AD[10]. Recently, there have been two models that have combined
human Aβ and Tau into a single *C. elegans* model and observed decreased lifespan
and progeny viability along with increased neuron loss, motor dysfunction and protein
aggregation[11,12]. In addition, there were alterations on a transcriptomic level including
changes in UDP-glucuronosyltransferase, protein phosphorylation, and aging genes[11].
These models all express the human form of the toxic proteins due to them not being
endogenously found within *C. elegans*. Despite these proteins not being endogenously
expressed these transgenic models have each successfully reproduced certain aspects
of AD-like phenotypes, including insoluble Tau accumulation, age-related uncoordinated
movement, and neurodegeneration[10].

To examine the impact of simultaneous expression of Aβ and Tau, we crossed *C.*
elegans strains expressing pan-neuronal Aβ and pan-neuronal 4R1N Tau with V337M
mutation, which allows us to further examine the interaction of the proteins but with the
addition of the V337M mutation to accelerate Tau aggregation[13,14]. This model allowed
us to explore the impact of these AD-related proteins on whole-organism protein
insolubility, transcriptome dynamics, and metabolomics across different stages of life,
including young, middle, and old age. Significantly, this model allowed us to uncover
insights into potential early-life effects on reproductive functions and the intricate
relationship between normal aging and the development of chronic age-related
pathologies.

98 **Results**

99 **Systemic phenotypes observed with expression of Tau or Aβ.**

To investigate the impact of expression of WT human Aβ and 4R1N Tau containing the
V337M mutation on protein aggregation[13,14], we generated a *C. elegans* strain with
simultaneous pan-neuronal expression of both proteins (Fig. 1A). This was achieved by
crossing the existing pan-neuronal Aβ strain, CL2355, with the pan-neuronal Tau strain,
CK10. The CL2355 strain uses the *snb-1* promoter to drive expression of Aβ in the
nerve ring, ventral cord, dorsal cord and neuronal cell bodies[15]. While the CK10 strain
uses the *aex-3* promoter to drive expression of Tau in nearly all the neurons including
the major ganglia next to the nerve ring, ventral cord and tail ganglia[16]. This newly
generated strain was first used to determine whether the expression of Aβ and Tau
leads to enhanced phenotypic dysfunction. We initially focused on testing for a
phenotype seen in AD, alteration in mitochondrial function. We tested mitochondrial

function by measuring the worm's ability to move and the rate of oxygen consumption of
the mitochondria. Movement patterns of the worms were quantitated by measuring the
number of body bends (thrashing rate) in a given time. The Tau;Aβ strain exhibited a
significantly decreased thrashing rate compared to the WT, Tau, and Aβ strains (Fig.
1B). Mitochondrial function was quantitated by measuring the rate of oxygen
consumption using a Seahorse XFe96 Analyzer. The Tau;Aβ strain displayed increased
average oxygen consumption rates (OCR) compared to the Aβ strain while the Tau
strain had increased average OCR compared to WT and Aβ strains, suggesting
heightened mitochondrial activity (Fig. 1C). These findings imply that the expression of
Tau and Aβ have a synergistic effect on the body movement phenotype while Tau
appears to have a stronger effect on mitochondrial OCR.

In the maintenance of the Tau;Aβ strain, we observed altered hermaphrodite self-
fertility. This observation drove us to characterize this possible toxicity phenotype. The
number of eggs laid (fecundity) and the number of eggs that hatched (an early
indication of fertility) were measured until the exhaustion of egg laying. The Tau and
Tau;Aβ strains exhibited a decreased average total number of eggs laid, while the Aβ
and Tau;Aβ strains showed a significantly decreased total progeny (Fig. 1D).

As a measure of overall health, we performed a survival assay on adult hermaphrodites,
revealing that the Tau;Aβ expressing strain had a decreased median and maximum
lifespan compared to the other three strains (Fig. 1E&F). This may represent
accelerated aging.

These results indicate that the pan-neuronal expression of Aβ and Tau exerts systemic
effects on the worms, with Aβ affecting progeny production, Tau influencing
mitochondrial OCR, and the combined expression of both proteins leading to reduced
body movement and lifespan. These observations underscore the impact of Tau and Aβ
interactions on the phenotypic characteristics of the worms.

Synergistic effects of Aβ and Tau expression on metabolic profiles in young
animals.

To study the systemic consequences of expressing human neurotoxic proteins in *C.*
elegans, we conducted global metabolomic analysis of the WT, Tau, Aβ, and Tau;Aβ
expressing strains on day 1, 3, 5 and 8 of worm adulthood. At day 1 of adulthood the
Tau and Aβ strains exhibited relatively minor differences in metabolite profiles
compared to the WT strain (Fig. 2A). However, the Tau;Aβ expressing strain displayed
extensive alterations in metabolite profiles compared to the WT strain (Fig. 2A), with the
Aβ expressing strain resembling this trend based on correlation analysis (Fig. 2B).

Further examination of the metabolites altered in the day 1 Tau;Aβ expressing strain
revealed contrasting abundance compared to the WT strain (Fig. 2C). The metabolites
significantly increased in the Tau;Aβ strain primarily belonged to the categories of lipids
and amino acids, specifically metabolites involved in spermidine and spermine
biosynthesis, biotin metabolism and butyrate metabolism (Fig. 2C). In contrast the
metabolites significantly decreased were in the categories of nucleotides and amino

acids involved in folate metabolism and glycine and serine metabolite(Fig. 2C). These
changes could connect back to the mitochondrial phenotype as we see changes in
mitochondrial-related metabolites including spermidine and spermine biosynthesis along
with glycine and serine metabolism.

In summary, our metabolomic analysis highlighted the synergistic effects of Aβ and Tau
expression on the metabolic profiles in young adult animals. The Tau;Aβ expressing
strain exhibited substantial alterations in metabolites, particularly in lipid and amino acid
metabolism.

160 Expression of Tau and Aβ disrupts metabolism throughout life.

To gain deeper insights into the metabolic changes associated with aging in the WT,
Tau, Aβ, and Tau;Aβ strains, we examined day 3, 5, and 8 metabolomic data. When
looking at the number of metabolites that significantly change on day 3, 5 or 8 when
compared to day 1 we observed that significant changes were primarily evident on day
8 for the WT and Tau expressing strains (Fig. 3A). The Aβ expressing strain exhibited a
slight increase in significantly altered metabolites on day 5, which became more
prominent on day 8. In contrast, the Tau;Aβ expressing strain showed the most
significant changes on all three days, indicating a potential additive and/or synergistic
relationship resulting from the simultaneous expression of both proteins in neurons.

Comparing the metabolite levels on day 8 compared to day 1 for each strain, we
identified an overlap in metabolites altered in abundance on day 8 across all four
strains. However, the Tau;Aβ strain had a greater number of unique lower abundance
metabolites that did not overlap with the other strains (Supplementary Fig. 1A&B).
When comparing the AD strains to the WT on day 3, 5, and 8, the Tau;Aβ strain
exhibited the most significant alterations in metabolites on day 5 (Supplementary Fig.
1C). This was also visualized in Supplementary Figs. 1D, E and F where the log fold
change of the metabolites in AD strains vs WT on the different days are depicted. The
Tau;Aβ strain were shown to have a slight increase in log fold change on day 3 and a
larger change on day 5 in comparison to the other AD strains. On day 8 the AD strains
appear to have an increase in metabolite fold change with the Tau;Aβ strain still having
the largest change (Supplementary Fig. 1F).

We performed a weighted correlation network analysis (WGCNA) to cluster metaboliets
with similar abundance profiles into different modules (Supplementary Fig. 2A&B). One
of the modules, labeled "green," displayed relatively consistent levels of clustered
metabolites in the WT strain with age (Fig. 3B). The Aβ, Tau and Tau;Aβ strains
showed elevated metabolite abundance levels on day 1 and decreasing abundance
from day 3 to day 8. The Tau;Aβ strain exhibited a synergistic effect with a more
significant decrease in metabolite abundance with age compared to Aβ and Tau (Fig.
3B). This trend can also be seen in the heatmap of the green module metabolites (Fig.
3C). The metabolites in this green module consist mostly of lipids, such as
monoacylglycerols which are glycerolipids that are found to be elevated in AD brains
and alterations are found to occur in early stages of the disease[17].

In addition to the "green" WGCNA module, we also identified additional modules labeled
"salmon", "pink", and "grey60". The "salmon" module represents a cluster of metabolites
that are elevated in all ages in the Tau;Aβ strain, while they increased with age in the
WT, Tau and Aβ strains (Supplementary Fig.2C). These metabolites are associated with
glycerophospholipid metabolism and purine metabolism (Supplementary Fig.2D). The
"pink" module represents a cluster of metabolites that are elevated on day 1 in the
Tau;Aβ strain and decreased with age, while having constant levels in the Tau and Aβ
strains. (Supplementary Fig.2E). These metabolites are associated with biosynthesis of
unsaturated fatty acids (Supplementary Fig.2F). The "grey60" module represents a
cluster of metabolites that are elevated on day 1 in the Tau;Aβ strain and decreased
with age, while in the WT, Aβ and Tau strains the metabolites are decreased on day 1
and increased with age (Supplementary Fig.2G). These metabolites are associated with
glycolysis/gluconeogenesis and pentose phosphate pathway (Supplementary Fig.2H).

Elevated pathways in the Tau;Aβ strain, including energy and fat, suggests that the
expression of Tau and Aβ leads to disruptions in energy metabolism, lipid metabolism,
and mitochondrial function, which may contribute to the observed changes in the
mitochondrial, body movement, fertility and fecundity phenotypes associated with the
Tau;Aβ strain.

Expression of Aβ and Tau drives insoluble protein accumulation in young
animals.

To further study the systemic effects of the expression of Aβ and Tau, we were
interested in determining if their expression leads to the aggregation of other proteins as
we had previously observed in *C. elegans* with aging, via induction by other stressors,
such as iron[18–20]. We collected insoluble proteins from Day 1 adult worms of WT, Tau,
Aβ, and Tau;Aβ strains, followed by mass spectrometric analysis (see Supplementary
Table 1-4) using data-independent acquisition[21,22].

A comparison of the insoluble proteins in Tau vs WT, Aβ vs WT, and Tau;Aβ vs WT
strains demonstrated an overlap among all three AD strains, with the Aβ expression
alone significantly increased the abundance of insoluble proteins compared to WT
worms. The Tau;Aβ expressing strain showed the most substantial alterations similar to
the metabolomics data, where we saw changes in serine, glycine metabolism which
plays a role in protein synthesis which may have a role in this proteomic finding
(Fig.4A).

To gain insights into the functional interactions of the insoluble proteins in the AD
strains, we constructed a protein-protein interaction network. This network revealed
associations with germ cells and cell physiology, oxidative phosphorylation, and
organism development (Fig. 4A).

Notably, the expression of either Tau or Aβ alone led to increased insolubility of proteins
associated with germ cell morphology and oxidative phosphorylation with potential
relevance to the observed fertility defects and mitochondrial functional changes.
Perturbation in mitochondrial function is also observed in human AD, suggesting that

even at an early stage of adulthood, the Tau;Aβ expressing strain exhibits specific
aspects relevant to AD[23,24].

Increased protein insolubility induced by Tau and Aβ expression in middle-aged
animals.

Building upon our observations of increased protein insolubility in the Tau;Aβ
expressing strain in young animals, we investigated this molecular phenotype during
aging. We collected insoluble proteins from day 3, 5, and 8 adult worms of the WT, Tau,
Aβ, and Tau;Aβ expressing strains, followed by mass spectrometric analysis (see
Supplementary Table 1-4). Our findings revealed that comparing insoluble proteins
significantly elevated on day 3, 5, and 8 in the WT, Tau, Aβ, and Tau;Aβ strains
compared to day 1 adults, we observed overlaps on day 3, 5 and 8 adults with day 1
adults for all four strains (Fig. 5A). The abundance of insoluble proteins in the WT strain
increased with age (Fig. 5A), consistent with previous reports[18,25,26]. On day 5, the Tau
expressing strain showed an increase in insoluble protein abundance compared to day
1. On day 8, the Aβ and Tau;Aβ strains exhibited an increase in insoluble proteins
compared day 1. In contrast, the Aβ and Tau;Aβ strains exhibited a lower amount of
insoluble proteins on day 3 and 5 (Fig. 5A).

Enrichment analysis revealed an enrichment of proteins involved in ion and small
molecule binding in the WT strain on day 8 compared to day 1 (Fig. 5B). Similarly,
enrichment analysis of the Tau strain on day 5 compared to day 1 highlighted an
increase in proteins associated with ribonucleotide binding (Fig. 5C). Enrichment
analysis of the Aβ strain on day 8 compared to day 1 revealed an increase in proteins
associated with cytoskeleton structure and ribonucleoprotein granules (Fig. 5D).
Furthermore, enrichment analysis of the Tau;Aβ expressing strain on day 8 compared to
day 1 showed an increase in proteins involved in ribonucleotide binding (Fig. 5E). The
increase in insolubility of ribonucleotide binding proteins may connect back to our
previously mentioned observation of decreased nucleotide metabolites in our day 1
metabolomics data.

Supplementary Figure 4A demonstrates that the insoluble proteome of the Tau;Aβ
strain was increased compared to the WT strain on day 3, 5, and 8, in comparison to
Tau or Aβ alone. Enrichment analysis of the insoluble proteome on day 3 and day 5 of
the Tau;Aβ strain compared to WT highlighted numerous ribosomal proteins
(Supplementary Fig. 4B). When comparing day 3, 5, and 8 of the Tau;Aβ strain to day 1,
the insoluble proteins observed on day 3 and 5 were also present on day 8
(Supplementary Fig. 4C). There was an overlap of insoluble proteins identified on day 1,
3, 5, and 8 of the Tau;Aβ strain compared to WT, with each day also featuring proteins
not observed on the other days (Supplementary Fig. 4D).

The aging insoluble proteome analysis revealed that the Tau;Aβ expressing strain
exhibited an early increase in insoluble proteins on day 1 (Fig. 4), which decreased after
day 1 to then increase again on day 8. The aging insoluble proteome of the WT, Tau,
and Tau;Aβ strains consisted of proteins involved in binding a range of small molecules.

Taken in total, our data suggests that the expression of Tau and/or Aβ leads to a
complex remodeling of protein-wide insolubility. The non-linear aging response
observed may reflect compensatory mechanisms aimed at preventing or removing
insoluble proteins.

279   Tau and Aβ expression in young adult worms resemble WT aging at insoluble
280   proteome and metabolic levels.

Comparing insoluble proteome and metabolomic changes in the day 1 Tau;Aβ
expressing strain to the aged WT strain revealed similarities between the two
conditions. Proteins that were more abundant in the insoluble proteome of day 8 WT
compared to day 1 WT overlapped with the majority of insoluble proteins in day 1
Tau;Aβ compared to day 1 WT (Fig. 6A). These overlapping proteins are involved in
lipid transport and ribosome function. Similarly, some of the metabolites that were
altered in day 8 WT compared to day 1 WT also overlapped with metabolites altered in
day 1 Tau;Aβ compared to day 1 WT (Fig.6B). These overlapping metabolites fall
mostly in the categories of lipids and amino acids (Fig.6C). In the Tau;Aβ strain, these
metabolites had elevated abundance levels on day 1, resembling the levels observed in
day 8 WT (Fig. 6C).

This suggests that the expression of Tau and Aβ in young adult worms resembles the
aging-associated changes observed in the insoluble proteome and metabolic pathways
of WT worms. The simultaneous expression of Tau and Aβ in young adult worms
appears to accelerate the aging process, particularly those affecting lipid pathways. The
observed similarities between the insoluble proteome and metabolic changes in the
Tau;Aβ strain on day 1 and the aged WT strain suggest that the presence of Tau and
Aβ proteins leads to accelerated aging on a proteomic and metabolomic level,
resembling the changes typically seen in older organisms. These alterations in lipid
pathways may contribute to the dysregulation of cellular processes and potentially
exacerbate the detrimental effects associated with aging.

303   Discussion

We developed a *C. elegans* model of Alzheimer's disease and Alzheimer's related
disease (ADRD) proteotoxicity by expressing both human Aβ and human 4R1N Tau
containing the V337 mutation in neurons. The generation of this new dual strain
facilitated an investigation into the combined effects of Aβ and Tau expression on
disease progression through phenotypic and geromic analysis.

The simultaneous expression of Tau and Aβ has been demonstrated to influence
various phenotypes encompassing progeny viability, body movement, lifespan, and
neurol loss[11,12]. In our Tau;Aβ strain, we observed a synergistic effect on health (body
movement and lifespan). Intriguingly, mitochondrial function and fertility phenotypes
were predominantly influenced by either Tau or Aβ alone. This dichotomy suggests that

the combined expression of Tau and Aβ may have exert distinct effects on different
phenotypes.

Collectively, these findings provide compelling evidence for the intricate relationship
between Aβ and Tau expression and their associated phenotypic outcomes,
underscoring the significance of their interplay in the context of neurodegenerative
diseases.

The phenotypic findings give rise to further questions, specifically whether Tau and Aβ
drive different aspects of the disease in humans or if all aspects of the disease result
from a combined effect. Additionally, there is a pressing need to delve deeper to
elucidate the mechanisms involved and navigating whether Tau and Aβ are directly
interacting. These avenues of inquiry are pivotal for comprehensive understanding of
the molecular intricacies underpinning ADRD pathogenesis.

To delve further into the intricate effects of Tau and Aβ we conducted a geromic
analysis, including transcriptomics, metabolomics, and insoluble proteomics. In addition,
we were interested in determining the temporal order of the phenotypic versus the multi-
omic changes. Transcriptomic analysis of the young adult Tau;Aβ strain highlighted
significant changes in mRNAs related to reproduction and mitochondrial function,
potentially linking to our observed phenotypes (Supplementary Fig. 5). Notably, we
identified similarities with the transcriptomic data published by Wang et al., showing
shared upregulated mRNAs and genes associated with protein phosphorylation and
UDP-glucuronosyl transferase[11]. However, differences were noted in downregulated
genes possibly attributed to the variation in the Tau strain used in creating the Tau;Aβ
strain. This underscores the importance of employing multiple models with different
protein isoforms and mutations to comprehensively understand the range of
transcriptomic changes in the disease.

Metabolomics analysis revealed early-life increases in metabolite levels for the Tau;Aβ
strain, particularly lipids and amino acids. Elevated amino acid levels can be attributed
to higher energy demand, prompting amino acid oxidation as an alternative energy
source[27]. Furthermore, elevated amino acid levels may be connected to increased
protein synthesis and activation of mTOR inhibiting autophagy[27]. This underscores the
importance of studying a range of omics, as a synergistic effect between Tau and Aβ
was specifically observed at the metabolomic level, emphasizing need to explore
various omics to avoid potential oversights.

The insoluble proteome analysis indicated an increase in insoluble proteins in the young
adult Tau;Aβ strain compared to the WT strain, associated with germ cell morphology
and oxidative phosphorylation, which may also impact our observed phenotypes. These
changes align with the increase in amino acids observed in our metabolomic data and
its connection to protein synthesis. Additionally, alterations in mRNAs related to
ribonucleotide binding and ribosome structure which could associated with the increase
in insoluble proteins.

Thus far we have observed that phenotypic changes and multi-omic changes both occur
in early life with simultaneous expression of Tau and Aβ. Given these early life changes
we observed we were keen on examining whether these alterations persist as the
organisms age. Our observations revealed a progressive increase in metabolite
alterations associated with energy and lipid metabolism in the Tau;Aβ strain, mirroring
changes observed in AD[23,24,28–31]. These findings imply that the simultaneous
expression of Tau and Aβ induces disruptions in energy metabolism, lipid metabolism,
and mitochondrial function, manifesting early in life and that persists with age.
Significantly, this finding prompts consideration of whether metabolomic dysfunction
represents one of the initial indicators of the disease.

Additionally, aging in the Tau;Aβ strain was accompanied by an increase in insoluble
proteins associated with ribonucleotide and small molecule binding. Dysfunction in
these binding proteins may prevent proper binding, leading to their aggregation with
age, a phenomenon observed in the initial stages of AD[32]. Notably, in the young adults
Aβ appears to drive protein insolubility, yet this trend does not persist with age. The
observed plateau and subsequent increase in insolubility at day 8 could be attributed to
the induction of Aβ at the late larval stage, resulting in increased protein insolubility at
day 1. This may represent an acute stress response that does not have a prolonged
effect beyond the first 24 hours of expression. However, this initial stress may have
lasting effects, becoming evident in the increased insoluble proteome at the advanced
age of day 8.

Comparative analysis of the insoluble proteome and metabolomic changes between the
day 1 Tau;Aβ strain and the aged WT strain unveiled striking similarities, suggesting a
convergence of molecular alterations. The overlapping insoluble proteins predominantly
associated with lipid transport and ribosome function, along with overlapping
metabolites already elevated in the day 1 Tau;Aβ strain, became more abundant with
age in the WT strain. The simultaneous expression of Tau and Aβ in young adult worms
appears to accelerate and intensify aging processes, particularly those impacting lipid
pathways. These disruptions in lipid pathways may contribute to the dysregulation of
cellular processes, potentially exacerbating the detrimental effects associated with
aging.

The Tau;Aβ strain, when juxtaposed with the findings from the prior studies by Wang et
al. and Benbow et al., collectively illustrates that the concurrent expression of both Aβ
and Tau gives rise to distinct neuronal and systemic phenotypes[11,12]. This integrated
model not only advances our understanding of the individual contributions of Tau and
Aβ but also illuminates their synergistic effects, providing a comprehensive perspective
on their impact on AD progression from early life. It facilitates the examination of early
systemic changes in AD, and aids in the identification of potential therapeutic targets.

Crucially, the Tau;Aβ strain contributes valuable insights into AD by unveiling
transcriptomic changes related to reproduction and mitochondrial function, significant
shifts in metabolomics, and alterations in the insoluble proteome. These findings not
only align with, but also expand upon previous studies, underscoring the intricate

relationship between Aβ and Tau expression and their consequential impact on
neurodegenerative processes. Furthermore, our research accentuates the convergence
of molecular alterations between the Tau;Aβ strain and aging in the WT strain,
particularly within lipid pathways.

In summary, the Tau;Aβ strain, in conjunction with earlier studies, provides a holistic
portrayal of early systemic changes in Alzheimer's disease. Our study introduces
additional transcriptomic timepoints, complemented by metabolomic and insoluble
proteomic data, contributing to the existing field of Tau and Aβ co-expression *C.*
elegans models. Moreover, our results validate prior studies that observed AD-
associated proteins affected reproduction, lifespan, and body movement in a *C. elegans*
model expressing both Tau and Aβ[11,12]. Our study yields valuable insights into potential
therapeutic targets and serves as a resource model for studying AD pathogenesis. By
uncovering changes at the metabolomic and insoluble proteomic levels, our research
introduces novel perspectives to the field, providing geromic datasets that can be mined
to enhance our understanding of ADRD progression.

Methods

C. elegans strains

The transgenic strains utilized in this study include non-transgenic wild-type N2, CK10
(P_{aex-3}::h4R1N Tau V337M; P_{myo-2}::GFP)[14], CL2355(*smg-1(cc546)*;P_{snb-1}::Aβ$_{1-42}$ + P_{mtl-}
$_2$::GFP)[13], and the simultaneous expression strain GL405 (P_{aex-3}::h4R1N Tau V337M;
P_{myo-2}::GFP; *smg-1(cc546)*;P_{snb-1}::Aβ$_{1-42}$ + P_{mtl-2}::GFP). CL2355 was obtained from the
Caenorhabditis Genetic Center and CK10 was obtained from the lab of Dr. Brian
Kraemer at the University of Washington. N2, CK10, CL2355, GL405 strains are
maintained at 15°C due to the *smg-1(cc546)* gene, which allows for the temperature-
dependent expression of Aβ. Once the worms reached the fourth larval stage (L4), they
were shifted to 25°C to induce Aβ expression. Throughout the study, the worms were
cultured on nematode growth media (NGM) (containing: Becton Dickinson and
Company Difco Agar, Bacteriological, ref:214510, Gibco Bacto Peptone Enzymatic
Digest of Protein ref: 211820, Sigma-Aldrich Sodium Chloride S9888-1KG, Sigma-
Aldrich Magnesium Sulfate M7506-500G lot#SLBR0877V, Sigma-Aldrich Calcium
Chloride C1016-500G lot#SLCC1966) plates (VWR, petri dish polystyrene disposable
sterilized size: 60 x 15mm, cat no.25384-090) containing *E. coli* OP50 and passaged
once a week.

Referenced Name	Strain name	Genotype	Growth condition	Experiment condition
WT	N2	wildtype	15°C	25°C
Tau	CK10	P_{aex-3}::h4R1N Tau V337M; P_{myo-2}::GFP	15°C	25°C
Aβ	CL2355	*smg-1(cc546)*;P_{snb-1}::Aβ$_{1-42}$ + P_{mtl-2}::GFP	15°C	25°C
Tau;Aβ, double transgenic, simultaneous expression	GL405	*smg-1(cc546)*;P_{snb-1}::Aβ$_{1-42}$ + P_{mtl-2}::GFP x P_{aex-3}::h4R1N Tau V337M; P_{myo-2}::GFP	15°C	25°C

Hermaphrodite self-fertility and fecundity assay

Egg lays were conducted at 15°C for each strain. Once worms develop to the L4 stage,
10 hermaphrodite worms were moved to individual small NGM plates (Gen Clone,
tissue culture dishes 38x13mm, cat# 25-200) and shifted to 25°C. On day 1 of the adult
stage, the number of eggs laid was counted to determine fecundity and the number of
hatched larvae was counted to determine fertility. The worms were then transferred to
fresh NGM plates. The number of eggs laid, and hatched larvae were counted each day
until the end of their reproductive cycle. To determine if there were significant
differences in fertility and fecundity among the strains, statistical analysis was
performed using ordinary one-way ANOVA with Tukey's multiple comparisons test and
the data consisted of 3 biological replicates.

Thrashing assay

Following egg lays conducted at 15°C, the worms were allowed to develop to the L4
stage and then transferred to 10 μg/mL FUdR plates, followed by a shift to 25°C. On
day 1 of the adult stage, a droplet of S-basal solution (containing: Sigma-Aldrich
Potassium Phosphate Monobasic P5379-1KG lot#SLCG1814, Sigma-Aldrich Potassium
Phosphate Dibasic P3786-1KG lot#SLCC4706, and Sigma-Aldrich Sodium Chloride
S9888-1KG) was placed on a glass slide, and a single worm was placed into the droplet
to acclimate for 30 seconds. Subsequently, the body bends of the worm were counted
for another 30 seconds. This process was repeated for each strain, with 10-20 worms
included in each of the 3 biological replicate. To determine if there were significant
differences in the number of body bends between the strains, statistical analysis was
performed using ordinary one-way ANOVA with Tukey's multiple comparisons test and
the data consisted of 3 biological replicates.

Mitochondrial oxygen consumption rate assay

To evaluate mitochondrial function, a mitochondrial oxygen consumption rate (OCR) assay was performed. Over 200 worms were mass cultured on 100 mm NGM agar plates (petri dish polystyrene disposable sterilized size: 100 x 15mm, cat no. 25384-342) at 15°C. Age synchronization was achieved by isolating eggs through hypochlorite treatment of gravid adults. Once the worms reached the L4 stage, they were transferred to NGM plates containing 10 μg/mL FUdR and shifted to a temperature of 25°C. Approximately 200 worms were collected on day 1 of the adult stage for further analysis.

An Agilent Seahorse XFe96 Analyzer was utilized to measure the OCR using a previously described protocol[33]. Briefly, for each strain, approximately 20 hermaphrodite worms were loaded into each well of a 96-well microtiter plate (Agilent Seahorse XFe96/XF Pro FluxPak, ref#103792-100), with 8 technical replicates per strain. Basal OCR was measured for 5 cycles, where each cycle consisted of 2 minutes of mixing in the well, 4 minutes of waiting, and 2 minutes of OCR measurement in the well.

To determine if there were significant differences in OCR between the strains, statistical analysis was performed using ordinary one-way ANOVA, followed with Tukey's multiple comparisons test and the data consisted of 3 biological replicates.

Lifespan assay

To assess the lifespan of the worms, egg lays were conducted at 15°C. Once the worms reached the L4 stage, approximately 40-60 worms per strain were transferred to NGM plates containing 10 μg/mL FUdR and maintained at a temperature of 25°C. Three technical replicates were performed for each strain to ensure robustness and reproducibility of the results. The worms were monitored every two days for touched provoked movement. This was achieved by gently poking the worms with a platinum wire and observing their response. The absence of movement upon stimulation was considered an endpoint, indicating mortality.

Statistical analysis was conducted using survival representation (Kaplan-Meier) in GraphPad Prism™ software. Survival curves were generated, and the lifespan of each strain was compared using the Log-rank (Mantel-Cox) test. This analysis allowed for the assessment of significant differences in survival between strains and the data consisted of 3 biological replicates.

3' Tag RNA-seq

For the 3' Tag RNA-seq analysis, worms were mass cultured at 15°C, and age-synchronization was achieved by isolating eggs through hypochlorite treatment of gravid adults. Once the worms reached the L4 stage, they were transferred to FUdR NGM plates and maintained at a temperature of 25°C.

Approximately 1000 worms and 3 biological replicates per strain were collected on the first day of the adult stage and stored in a 1.5 mL Eppendorf tube containing 300-500 μL of RNA lysis buffer from the Zymo Research RNA extraction kit (cat#R1055). The samples were immediately placed at -80°C for storage. To facilitate RNA extraction, the

samples underwent three cycles of freeze-thaw, alternating between room temperature
and -80°C. Subsequently, the samples were incubated on a heat block at 55°C for 0.5-1
hour.

RNA extraction was performed using a Zymo Research RNA extraction kit according to
the provided protocol. The protocol included a DNAse treatment step to remove any
genomic DNA contamination. The elution step resulted in approximately 40 µL of
purified RNA.

The prepared RNA samples were then sent out on dry ice for 3' Tag RNA-Seq analysis.
The sequencing was conducted at the DNA Technologies and Expression Analysis
Cores at the UC Davis Genome Center, with support from NIH Shared Instrumentation
Grant 1S10OD010786-01.

Binarized transcriptomic aging (BiT age) clock calculation

Read counts from the RNA-seq data were transformed into counts per million using
edgeR [34]. Using this file and the regression coefficients in https://github.com/Meyer-
DH/AgingClock/blob/main/Data/Predictor_Genes.csv we calculated the biological age of
each sample from day 1 using the Python script https://github.com/Meyer-
DH/AgingClock/blob/main/src/biological_age_prediction.py

Metabolomics

For metabolomic analysis, worms were mass cultured at 15°C and synchronized by
isolating eggs using hypochlorite treatment. Upon reaching the L4 stage, the worms
were transferred to FUdR 100mm NGM plates at 25°C. At the adult stage day 1,3,5,
and 8, approximately 250µl of worms and 3 biological replicates were collected by
washing off with S-basal solution and immediately snap freezing [35,36]. These samples
were then submitted to Metabolon Inc. (Morrisville, NC, USA) for global metabolic
profiling.

Sample preparation was conducted using the automated MicroLab STAR® system from
the Hamilton Company. Prior to the extraction process, several recovery standards
were added to the samples for quality control purposes. The extracted samples were
analyzed using a Waters ACQUITY ultra-performance liquid chromatography (UPLC)
system and a Thermo Scientific Q-Exactive high resolution/accurate mass spectrometer
equipped with a heated electrospray ionization (HESI-II) source and Orbitrap mass
analyzer operating at a mass resolution of 35,000.

Raw data obtained from the analysis was processed by Metabolon's hardware and
software. This involved extraction of the data, identification of peaks, and quality control
processing. Each metabolite was corrected by registering the medians to a value of one
(1.00) and normalizing each data point proportionately. In certain cases, metabolite data
was further normalized to total protein levels determined by Bradford assay to account
for differences in metabolite levels due to variations in the amount of biological material
present in each sample.

A total of 767 biochemical features were identified, including 695 metabolites with
known composition and 72 compounds with unknown structural identity. Statistical

analysis was performed using two-way ANOVA to identify biochemicals that exhibited
significant differences between experimental groups. Additionally, a false discovery rate
(q-value) was estimated to account for multiple comparisons.

<u>Insoluble proteome</u>

For the insoluble proteome analysis, which is based on [25], worms were mass cultured at
15°C and age-synchronized by isolating eggs using hypochlorite treatment of gravid
adults. Upon reaching the L4 stage, the worms were transferred to FUdR NGM plates at
25°C. Approximately 3,000 worms at the adult stage day 1,3,5 and 8 were collected in
S-basal solution and stored at -80°C.

To extract the insoluble protein fraction, the samples were thawed and vortexed with
worm lysis buffer containing 20 mM Tris base (pH 7.4), 100 mM NaCl, 1 mM $MgCl_2$, and
EDTA-free protease inhibitor. The lysate was then sonicated for 10 cycles and
centrifuged at 3,000 x g for 4 minutes in a cold room. The resulting supernatant,
containing the aqueous-soluble protein fraction, was transferred to new 1.5 mL
Eppendorf tubes, and the protein concentration was quantified using a BCA assay.

The protein lysate was further processed by centrifuging it for 15 minutes at 20,000 x g
in 4ºC. The resulting supernatant was saved as the SDS-soluble fraction. The pellet
obtained from this step was washed with 500μl of worm lysis buffer containing 1% SDS
and centrifuged at 20,000 x g for 15 minutes. The supernatant was removed, and this
washing step was repeated two more times to remove the SDS-soluble fraction. The
remaining pellet, known as the 1% SDS-insoluble protein fraction, was resuspended in
60μl of 7% formic acid (Thermo Scientific, ref: 85178) and vortexed to dissolve the
pellet. Subsequently, the sample was sonicated for 30 minutes and dried in a vacuum
concentrator for 1 hour to remove the formic acid solution.

The total amount of 1% SDS-insoluble protein, per sample, was prepared for trypsin
digestion. Each sample was reduced using 20 mM dithiothreitol in 50 mM
triethylammonium bicarbonate buffer (TEAB) at 50°C for 10 min, cooled to room
temperature (RT) and held at RT for 10 min, and alkylated using 40 mM iodoacetamide
in 50 mM TEAB at RT in the dark for 30 min. Samples were acidified with 12%
phosphoric acid to obtain a final concentration of 1.2% phosphoric acid. S-Trap buffer
consisting of 90% methanol in 100 mM TEAB at pH ~7.1, was added and samples were
loaded onto the S-Trap mini spin columns. The entire sample volume was spun through
the S-Trap mini spin columns at 4,000 × g and RT, binding the proteins to the mini spin
columns. Subsequently, S-Trap mini spin columns were washed twice with S-Trap
buffer at 4,000 × g at RT and placed into clean elution tubes. Samples were incubated
for one hour at 47 oC with sequencing grade trypsin (Promega, San Luis Obispo, CA)
dissolved in 50 mM TEAB at a 1:25 (w/w) enzyme:protein ratio. Afterwards, trypsin
solution was added again at the same ratio, and proteins were digested overnight at
37°C.

Peptides were sequentially eluted from mini S-Trap spin columns with 50 mM TEAB,
0.5% formic acid (FA) in water, and 50% acetonitrile (ACN) in 0.5% FA. After centrifugal
evaporation, samples were resuspended in 0.2% FA in water and desalted with C18
Ziptips (MilliporeSigma, Burlington, MA). Desalted peptides were then subjected to an

additional round of centrifugal evaporation and re-suspended in 20 μL of 0.2% FA in
water and 1 μL of indexed Retention Time Standard (iRT, Biognosys, Schlieren,
Switzerland). Samples were then subjected to mass spectrometric analysis using a
high-performance liquid chromatography (HPLC) system combined with a chip-based
HPLC system (Eksigent nano-LC) directly connected to a quadrupole time-of-flight
mass spectrometer (TripleTOF 5600, a QqTOF instrument) as detailed in a step-by-step
protocol by Xie et al [25].

Each sample was acquired in data-dependent acquisition (DDA) mode to build peptide
spectral libraries, as described in Xie et al [25]. Data-Independent Acquisition
(DIA)/SWATH data was processed in Spectronaut (version 14.10.201222.47784) using
DIA. Data extraction parameters were set as dynamic and non-linear iRT calibration
with precision iRT was selected. DIA data was matched against an in-house
Caenorhabditis elegans spectral library that provides quantitative DIA assays for 3,651
C. elegans peptides corresponding to 910 protein groups and supplemented with
scrambled decoys (library size fraction of 0.01), using dynamic mass tolerances and
dynamic extraction windows. DIA/SWATH data was processed for relative quantification
comparing peptide peak areas from different days. Identification was performed using
1% precursor and protein q-value. Quantification was based on the peak areas of
extracted ion chromatograms (XICs) of 3 – 6 MS2 fragment ions, specifically b- and y-
ions, with q-value sparse data filtering and iRT profiling being applied (Supplementary
Table 1-4). For this sample set, local normalization was not implemented. Differential
protein expression analyses for all comparisons were performed using a paired t-test,
and p-values were corrected for multiple testing, using the Storey method [37].
Specifically, group wise testing corrections were applied to obtain q-values. Protein
groups with at least two unique peptides, q-value ≤ 0.01, and absolute Log2(fold-
change) > 0.58 are significantly altered.

Protein-protein interaction network

We extracted 62 proteins with increased abundance (q-value < 0.1) in the insoluble
fraction compared to N2 controls at day 1. We loaded this list of proteins into Cytoscape
[38] and generated a protein-protein interaction network using the stringApp plugin [39].
Then, we performed enrichment analysis for Biological Processes (Gene ontology),
KEGG pathways, and WormBase phenotypes. We colored the nodes of proteins from
the top-enriched processes in the network.

Geromic analysis

We conducted several omic analyses to gain a comprehensive geromic understanding
of the molecular changes associated with the studied conditions. The following
approaches were employed:

Enrichment analysis:

For the RNAseq and insoluble proteome data, we performed enrichment analysis using
g:Profiler. The custom background used for each specific comparison included all genes
or all proteins measured in that particular analysis. The enrichment analysis was carried
out with a user threshold of 0.05 and significance was determined using the Benjamini-

Hochberg false discovery rate (FDR) correction. Additionally, WormBase Enrichment
Analysis was utilized for some of the RNAseq enrichment, applying a q-value threshold
of 0.05 and the previously mentioned custom background. Metaboanalyst was utilized
for enrichment analysis of metabolomics data, employing a custom background
consisting of all measured metabolites.

WGCNA (weighted gene co-expression network analysis):

To identify modules of metabolites that exhibited correlated patterns across samples,
we employed WGCNA. This analysis involved using the abundance data of metabolites
and allowed us to identify groups or modules of metabolites that displayed similar
expression profiles, providing insights into potential coordinated metabolic pathways or
regulatory mechanisms[40].

Generation of heatmap and dot plots:

RStudio was utilized to generate heatmaps and dot plots to provide a comprehensive
visual overview of observed molecular changes and enriched pathways. Code and
referenced files can be found at https://github.com/Holcom-AM/multi-omic-analysis and
https://data.mendeley.com/datasets/zc4s3dszf5/1,

Resource Availability

Lead contact

647 Further information and requests for resources and reagents should be directed to and
648 will be fulfilled by the lead contact, Gordon J. Lithgow (glithgow@buckinstitute.org).

Materials availability

650 The Tau;Aβ strain generated in this study is available from the lead contact upon
651 request.

Data and code availability

Raw data and complete MS data sets have been uploaded to the Mass Spectrometry
Interactive Virtual Environment (MassIVE) repository, developed by the Center for
Computational Mass Spectrometry at the University of California San Diego, and can be
downloaded using the following link: https://massive.ucsd.edu/ProteoSAFe/private-
dataset.jsp?task=e2c3bbad2f214e88bc70ea487d71a3ed (MassIVE ID number:
MSV000092031; ProteomeXchange ID: PXD042484. Enter the username and
password in the upper right corner of the page: Username: MSV000092031_reviewer;
Password: winter. Transcriptomics and metabolomics data can be found at
https://github.com/Holcom-AM/multi-omic-analysis and
https://data.mendeley.com/datasets/zc4s3dszf5/1.

References

1. Rostagno, A. A. Pathogenesis of Alzheimer's Disease. *Int. J. Mol. Sci.* **24**, 107 (2022).

2. Busche, M. A. & Hyman, B. T. Synergy between amyloid-β and tau in Alzheimer's disease. *Nat. Neurosci.* **23**, 1183–1193 (2020).

3. Lee, W. J. *et al.* Regional Aβ-tau interactions promote onset and acceleration of Alzheimer's disease tau spreading. *Neuron* **110**, 1932-1943.e5 (2022).

4. Zhang, H. *et al.* Interaction between Aβ and Tau in the Pathogenesis of Alzheimer's Disease. *Int. J. Biol. Sci.* **17**, 2181–2192 (2021).

5. Pascoal, T. A. *et al.* Synergistic interaction between amyloid and tau predicts the progression to dementia. *Alzheimers Dement.* **13**, 644–653 (2017).

6. Vegh, C. *et al.* A Bird's-Eye View of the Multiple Biochemical Mechanisms that Propel Pathology of Alzheimer's Disease: Recent Advances and Mechanistic Perspectives on How to Halt the Disease Progression Targeting Multiple Pathways. *J. Alzheimers Dis. JAD* **69**, 631–649 (2019).

7. Chen, X., Barclay, J. W., Burgoyne, R. D. & Morgan, A. Using C. elegans to discover therapeutic compounds for ageing-associated neurodegenerative diseases. *Chem. Cent. J.* **9**, (2015).

8. Teschendorf, D. & Link, C. D. What have worm models told us about the mechanisms of neuronal dysfunction in human neurodegenerative diseases? *Mol. Neurodegener.* **4**, 38 (2009).

9. Sin, O., Michels, H. & Nollen, E. A. A. Genetic screens in Caenorhabditis elegans models for neurodegenerative diseases. *Biochim. Biophys. Acta BBA - Mol. Basis Dis.* **1842**, 1951–1959 (2014).

10. Alexander, A. G., Marfil, V. & Li, C. Use of Caenorhabditis elegans as a model to

study Alzheimer's disease and other neurodegenerative diseases. *Front. Genet.* **5**,

(2014).

11. Wang, C., Saar, V., Leung, K. L., Chen, L. & Wong, G. Human amyloid β peptide

and tau co-expression impairs behavior and causes specific gene expression

changes in Caenorhabditis elegans. *Neurobiol. Dis.* **109**, 88–101 (2018).

12. Benbow, S. J., Strovas, T. J., Darvas, M., Saxton, A. & Kraemer, B. C. Synergistic

toxicity between tau and amyloid drives neuronal dysfunction and

neurodegeneration in transgenic C. elegans. *Hum. Mol. Genet.* **29**, 495–505 (2020).

13. Wu, Y. *et al.* Amyloid-β-Induced Pathological Behaviors Are Suppressed by Ginkgo

biloba Extract EGb 761 and Ginkgolides in Transgenic Caenorhabditis elegans. *J.*

Neurosci. **26**, 13102–13113 (2006).

14. Kraemer, B. C. *et al.* Neurodegeneration and defective neurotransmission in a

Caenorhabditis elegans model of tauopathy. *Proc. Natl. Acad. Sci.* **100**, 9980–9985

(2003).

15. Nonet, M. L., Saifee, O., Zhao, H., Rand, J. B. & Wei, L. Synaptic Transmission

Deficits in Caenorhabditis elegansSynaptobrevin Mutants. *J. Neurosci.* **18**, 70–80

(1998).

16. Iwasaki, K., Staunton, J., Saifee, O., Nonet, M. & Thomas, J. H. aex-3 Encodes a

Novel Regulator of Presynaptic Activity in C. elegans. *Neuron* **18**, 613–622 (1997).

17. Yin, F. Lipid metabolism and Alzheimer's disease: clinical evidence, mechanistic link

and therapeutic promise. *FEBS J.* **290**, 1420–1453 (2023).

18. Reis-Rodrigues, P. *et al.* Proteomic analysis of age-dependent changes in protein

solubility identifies genes that modulate lifespan. *Aging Cell* **11**, 120–127 (2012).

19. Klang, I. M. *et al.* Iron promotes protein insolubility and aging in C. elegans. *Aging* **6**,

975–991 (2014).

20. Mark, K. A. *et al.* Vitamin D Promotes Protein Homeostasis and Longevity via the

Stress Response Pathway Genes SKN-1, IRE-1, and XBP-1. *Cell Rep.* **17**, 1227–

1237 (2016).

21. Gillet, L. C. *et al.* Targeted data extraction of the MS/MS spectra generated by data-

independent acquisition: a new concept for consistent and accurate proteome

analysis. *Mol. Cell. Proteomics MCP* **11**, O111.016717 (2012).

22. Collins, B. C. *et al.* Multi-laboratory assessment of reproducibility, qualitative and

quantitative performance of SWATH-mass spectrometry. *Nat. Commun.* **8**, 291

(2017).

23. Poddar, M. K., Banerjee, S., Chakraborty, A. & Dutta, D. Metabolic disorder in

Alzheimer's disease. *Metab. Brain Dis.* **36**, 781–813 (2021).

24. Patro, S. *et al.* ATP Synthase and Mitochondrial Bioenergetics Dysfunction in

Alzheimer's Disease. *Int. J. Mol. Sci.* **22**, 11185 (2021).

25. Xie, X. *et al.* Quantification of Insoluble Protein Aggregation in Caenorhabditis

elegans during Aging with a Novel Data-Independent Acquisition Workflow. *J. Vis.*

Exp. JoVE 10.3791/61366 (2020) doi:10.3791/61366.

26. Huang, C. *et al.* Intrinsically aggregation-prone proteins form amyloid-like

aggregates and contribute to tissue aging in Caenorhabditis elegans. *eLife* **8**,

e43059 (2019).

27. Li, Q. & Hoppe, T. Role of amino acid metabolism in mitochondrial homeostasis.

Front. Cell Dev. Biol. **11**, 1127618 (2023).

28. Wilson, D. M. *et al.* Hallmarks of neurodegenerative diseases. *Cell* **186**, 693–714

(2023).

29. François, M. *et al.* Multi-Omics, an Integrated Approach to Identify Novel Blood

Biomarkers of Alzheimer's Disease. *Metabolites* **12**, 949 (2022).

30. Chen, Z. & Zhong, C. Decoding Alzheimer's disease from perturbed cerebral

glucose metabolism: Implications for diagnostic and therapeutic strategies. *Prog.*

Neurobiol. **108**, 21–43 (2013).

31. Olloquequi, J. *et al.* Impact of New Drugs for Therapeutic Intervention in Alzheimer's

Disease. *Front. Biosci. Landmark Ed.* **27**, 146 (2022).

32. Ghosh, A. *et al.* Alzheimer's disease-related dysregulation of mRNA translation

causes key pathological features with ageing. *Transl. Psychiatry* **10**, 1–18 (2020).

33. Koopman, M. *et al.* A screening-based platform for the assessment of cellular

respiration in Caenorhabditis elegans. *Nat. Protoc.* **11**, 1798–1816 (2016).

34. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. *Bioinforma. Oxf.*

Engl. **26**, 139–140 (2010).

35. Murphy, J. T. *et al.* Simple nutrients bypass the requirement for HLH-30 in coupling

lysosomal nutrient sensing to survival. *PLOS Biol.* **17**, e3000245 (2019).

36. Wang, W. *et al.* Comparative Metabolomic Profiling Reveals That Dysregulated

Glycolysis Stemming from Lack of Salvage NAD+ Biosynthesis Impairs

Reproductive Development in Caenorhabditis elegans. *J. Biol. Chem.* **290**, 26163–

26179 (2015).

37. Burger, T. Gentle Introduction to the Statistical Foundations of False Discovery Rate

in Quantitative Proteomics. *J. Proteome Res.* **17**, 12–22 (2018).

38. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of

biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

39. Doncheva, N. T., Morris, J. H., Gorodkin, J. & Jensen, L. J. Cytoscape StringApp:

Network Analysis and Visualization of Proteomics Data. *J. Proteome Res.* **18**, 623–

632 (2019).

40. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation

network analysis. *BMC Bioinformatics* **9**, 559 (2008).

41. Meyer, D. H. & Schumacher, B. BiT age: A transcriptome-based aging clock near

the theoretical limit of accuracy. *Aging Cell* **20**, e13320 (2021).

Acknowledgements

We would like to thank all members of the Lithgow, Andersen, Schilling, and Furman
labs. This work was supported by the Larry L. Hillblom Foundation, NIH Shared
Instrumentation Grant 1S10OD010786-01, 1S10OD016281-01 (to Buck Institute for the
TripleTOF mass spectrometry system), NIA grants RF1AG057358-01 and
R01AG029631-10. This work was also supported by U54 AG075932 (PI Campisi/ MPI
Schilling) and P01AI153559 (to Buck Bioinformatics Core). Traineeship supported by
NIA T32 AG052374 awarded to AH. The authors thank Dr. Akos Gerencser and former
Brand lab members for assistance with the Seahorse XFe96. The Aβ strain CL2355 is
from the lab of Dr. Chris Link and the Tau strain CK10 is from the lab of Dr. Brian
Kraemer. DeepVenn was used for venn diagrams and Biorender was used for the Fig
1A.

Author contributions

Conceptualization: R.S., J.K.A, G.J.L.; Phenotypic analysis: R.S., A.H.; Transcriptomics:
A.H.; Metabolomics: A.H.; Insoluble proteome: R.S., C.D.K., H.O., A.F., D.B., B.S.; Data

analysis: M.F., A.H., C.D.K.; Writing: A.H. wrote the manuscript, with input from all
authors; Supervision: J.K.A., G.J.L., D.F., B.S. M.F is co-first author.

Declaration of interests

The authors declare no competing interests.

Figure 1

A

Aβ Strain (CL2355)
$smg-1(cc546)$;P$_{snb-1}$::Aβ$_{1-42}$ + P$_{mtl-2}$::GFP

Tau =
Aβ =

Tau Strain (CK10)
P$_{aex-3}$::h4R1N Tau V337M; P$_{myo-2}$::GFP

×

Tau;Aβ Strain (GL405)

B

Day 1 Thrashing Rate

Body bends/30 seconds

Strains: WT, Tau, Aβ, Tau;Aβ

*** *** ns *** ns ns **

C

Mitochondrial Basal Oxygen Consumption Rate Day 1

Average oxygen consumption rate (pmol/min/worm)

Strains: WT, Tau, Aβ, Tau;Aβ

ns ns ns * *** ***

D

Average Total Number of Eggs Laid

Average number of eggs laid

Strains: WT, Tau, Aβ, Tau;Aβ

* ns ns ns ns ns

Average Total Progeny

Average number of progeny

Strains: WT, Tau, Aβ, Tau;Aβ

*** *** *** *** *** ns

Progeny Viability

Percent progeny viability

Strains: WT, Tau, Aβ, Tau;Aβ

*** *** *** ns *** ns

E

Survival (%) vs Day

WT, Tau, Aβ, Tau;Aβ

F

Strains	Average Max Lifespan	Average Median Lifespan	Log-rank (Mantel-Cox) test P-value *(compared to WT)*
WT	21	12	
Tau	18	11	0.0003
Aβ	25	11	0.1464
Tau;Aβ	17	10	<0.0001

Figure 1: Systemic phenotypes observed in the context of expression of Tau and Aβ. (A) Diagram of the cross used to create the dual model simultaneously expressing both human WT Aβ and mutant Tau. The newly generated strain expresses pan-neuronal human WT Aβ and 4R Tau containing the FTDP-17 mutation V337M as well as both pharyngeal and intestinal GFP. (B) Average thrashing rates of the 4 strains on day 1 of adulthood. 3 biological replicates with 10-20 worms per replicate. Error bars represent mean ± SEM. (C) Average basal mitochondrial oxygen consumption rate (OCR) of day 1 adult WT, Tau, Aβ and Tau;Aβ strains. 3 biological replicates with 8 technical replicates, which had around 20 worms in each technical replicate. Error bars represent mean ± SEM. (D) The average total number of eggs laid in the 4 strains with 3 biological replicates, 10 worms per replicate. Error bars represent mean ± SEM. The total progeny in the 4 strains with 3 biological replicates, 10 worms per replicate.. Error bars represent mean ± SEM. The percent of progeny viable was determined by the total number of progeny hatched and total eggs laid. Ordinary one-way ANOVA with Tukey's multiple comparisons test. Error bars represent mean ± SEM. (E) Average lifespan for WT, Tau, Aβ and Tau;Aβ strains. Includes table of average max and median lifespan and log-rank test p-value of AD strains compared to WT. 3 biological replicates with 3 technical replicates that had 40-60 worms each. Error bars represent mean ± SEM.

Figure 2

Figure 2: Expression of Tau and Aβ synergistically alters metabolite levels. (A)
Bar graph of fold metabolite change in day 1 Tau, Aβ and Tau;Aβ compared to WT. (B)
Pearson's product-moment correlation of fold metabolite change between Aβ vs WT
and Tau vs WT, and Tau;Aβ vs WT. (C) Heatmap of metabolites and their super
pathways, which are significantly increased in Tau;Aβ compared to WT, Tau and Aβ.
(D) Heatmap of metabolite and their super pathways, which are significantly decreased
in Tau;Aβ compared to WT, Tau and Aβ.

Figure 3

Figure 3: Expression of Tau and Aβ enhances metabolic changes with age. (A)
Bar graphs of the number of metabolites significantly changing in day 3, day 5 and day
8 vs day 1 in all 4 strains. (B) Graph of the WGCNA green module with all 4 strains from
day 1 to day 8 with 3 biological replicates. The y-axis value represents the eigengene
value, which is the summarized abundance of the metabolites in the green module. (C)
Heatmap of metabolites and their super pathways in the green module for all strains on
day 1, 3, 5 and 8.

Figure 4

Figure 4: Expression of Tau and Aβ increase insoluble proteins early in life. (A)
Upset plot of the number of insoluble proteins significantly increased in the AD strains
compared to WT on day 1 of adulthood. The intersections are the number of proteins
shared between the different comparisons, which are represented by a line connecting
dots. q ≤ 0.05. (B) GO analysis of insoluble proteins increased in Tau;Aβ strain found
significant increase of lipid transport genes, heatmap shows the abundance of those
lipid transport genes (C) Protein-protein interaction network of insoluble proteins
increased in the AD mutant strains compared to WT. 3 biological replicates, q ≤ 0.1.

Figure 5

Figure 5: Expression of Tau and Aβ increases day 8 insoluble proteome. (A) Upset
plot of the number of insoluble proteins significantly increased in the AD strains
compared to WT on day 3 vs day 1, day 5 vs day 1, and day 8 vs day 1 of adulthood
and the number of proteins shared between the different comparisons on each day. q ≤
0.05. (B) g:Profiler GO molecular function of proteins with increased insolubility on day 8
vs day 1 in WT. (C) g:Profiler GO molecular function of proteins with increased
insolubility on day 5 vs day 1 Tau. (D) Wormbase Gene ontology enrichment of proteins
with increased insolubility on day 8 vs day 1 in Aβ. (E) g:Profiler GO molecular function
of proteins with increased insolubility on day 8 vs day 1 in Tau;Aβ.

935   **Figure 6**

Figure 6: Tau and Aβ expression in young adult worms resembles WT aging at insoluble proteome and metabolomic levels. (A) Venn diagram of insoluble proteins that are increased on day 1 Tau;Aβ vs day 1 WT compared to day 8 WT vs day 1 WT and GO biological process of the insoluble proteins shared between the day 8 WT and day 1 Tau;Aβ. (B) Venn diagram of the metabolites in day 1 Tau;Aβ vs day 1 WT compared to day 8 WT vs day 1 WT and (C) a heatmap of the shared metabolites.

Extended Data Figure 1

Extended Data Figure 1: Enhanced Age-Related Metabolomic Changes in Tau;Aβ.
(A) Venn diagram of downregulated metabolites on day 8 of adulthood in Tau, Aβ, and Tau;Aβ compared to day 1, significance was based on q ≤0.05. (B) Venn diagram of metabolites with increased abundance on day 8 of adulthood in Tau, Aβ, and Tau;Aβ compared to day 1, significance was based on q ≤0.05. (C) Table of number of significantly increased and decreased metabolites on day 3, 5 and 8 of adulthood in Tau, Aβ, and Tau;Aβ compared to WT, significance was based on q ≤0.05. (D) Bar graph of fold metabolite change in day 3 Tau, Aβ and Tau;Aβ compared to WT. (E) Bar graph of fold metabolite change in day 5 Tau, Aβ and Tau;Aβ compared to WT. (F) Bar graph of fold metabolite change in day 8 Tau, Aβ and Tau;Aβ compared to WT.

Extended Data Figure 2

Extended Data Figure 2: WGCNA of metabolomic changes. (A) Clustering
dendrogram of metabolites with assigned module colors. (B) Module-trait relationship
plot with rows representing the modules and column the trait of interest, which is age.
Each cell contains the correlation and below is the p-value in parenthesis. The table's
cells are color-coded based on correlation, as indicated by the color legend. (C)
WGCNA eigengene plot of salmon module with all 4 strains from day 1 to day 8. (D) Dot
plot of MetaboAnalyst KEGG enrichment analysis of salmon module. (E) WGCNA
eigengene plot of pink module with all 4 strains from day 1 to day 8. (F) Dot plot of
MetaboAnalyst KEGG enrichment analysis of pink module. (G) WGCNA eigengene plot
of grey60 module with all 4 strains from day 1 to day 8. (H) Dot plot of MetaboAnalyst
KEGG enrichment analysis of grey60 module.

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034 **Extended Data Figure 3**

A

Tau vs WT	Aβ vs WT	Tau;Aβ vs WT
CELE_Y62H9A.6	vit-1	CELE_ZK180.6
CELE_D1054.10	CELE_ZC373.2	CELE_F10D11.6
CELE_ZK813.3	ttr-14	far-1
cbd-1	CELE_D1054.11	vit-1
perm-4	CELE_D1054.10	CELE_ZC373.2
	mup-4	col-184
	CELE_ZK813.3	tct-1
	CELE_Y62H9A.3	tbb-1
	vit-6	mup-4
	CELE_F48E3.4	ttr-14
	CELE_Y62H9A.6	CELE_D1054.11
	cht-3	F46H5.3
	perm-4	mua-3
	skpo-1	CELE_Y62H9A.5
	vit-2	grd-14
	vha-11	CELE_Y62H9A.3
	CELE_F55B11.2	pck-2
	cbd-1	cht-3
	efa-6	vha-13
	rpl-22	CELE_F48E3.4
	H28O16.1	vit-6
	CELE_Y47G6A.15	CELE_D1054.10
	let-805	vit-5
		fipr-21
		CELE_Y62H9A.6
		skpo-1
		rme-1
		car-1
		vit-2
		uba-1
		cbd-1
		perm-4
		rpl-4
		tba-1
		ant-1.1
		rpl-22
		pqn-22

1035

1036 **Extended Data Figure 3: Increased abundance of insoluble proteome with Aβ**
1037 **expression.** (A) List of insoluble proteins with increased abundance in Tau vs WT, Aβ
1038 vs WT and Tau;Aβ vs WT, significance was based on q ≤0.05.

1039

Extended Data Figure 4

A

Day 3 Increased Insoluble Proteins

Day 5 Increased Insoluble Proteins

Day 8 Increased Insoluble Proteins

B

Strain Comparison	Pathway Source	Term	-Log10(p-adj)	Gene Count	Proteins
Day 3 Tau;Aβ vs WT	WP	Cytoplasmic ribosomal proteins	1.392401	31	RPL-30,RPS-12,RPS-1,RPS-20,RPS-10,RPS-3,RPS-18,RPL-10,RPL-24.1,RPS-17,RPL-16,RPS-6,RPS-16,RPS-9,RPS-2,RPL-14,RPL-11.1,RPL-27,RPL-18,RPL-17,RPS-8,RPL-4,RPS-0,RPL-25.1,RPL-9,RPL-22,RPL-7A,RPS-4,RPL-5,RPL-6,RPL-3
Day 5 Tau;Aβ vs WT	GO:CC	Ribosome	1.422177869	25	RPL-17,RPS-5,RPS-19,RPS-7,RPS-12,RPS-1,RPS-3,RPL-16,RPL-9,RPS-8,RPS-6,RPL-7,RPS-16,RPL-5,RPL-12,RPL-36,RPL-22,RACK-1,RPS-9,RPL-4,RPS-2,RPL-19,RPS-4,RPS-0,RPL-6

C

Tau;Aβ Increased Insoluble Proteins

D

Tau;Aβ vs WT Increased Insoluble Proteins

Extended Data Figure 4: Alterations in aged insoluble proteome with enrichment for ribosomal proteins in Tau;Aβ strain. (A) Venn diagrams of insoluble proteins with increased abundance on day 3, 5 and 8 of adulthood in Tau, Aβ, and Tau;Aβ compared to WT, significance was based on q ≤0.05. (B) Table of g:Profiler enrichment of day 3 Tau;Aβ vs WT and day 5 Tau;Aβ vs WT. (C) Venn diagram of insoluble proteins that are increased on day 3, 5 and 8 of adulthood in Tau;Aβ compared to day 1 Tau;Aβ, significance was based on q ≤0.05. (D) Venn diagram of insoluble proteins increased on day 1, 3, 5 and 8 of adulthood in Tau;Aβ compared to WT, significance was based on q ≤0.05.

Extended Data Figure 5

Extended Data Figure 5: Aβ expression drives transcriptomic changes in the dual
Tau;Aβ strain. (A) Volcano plots of fold gene expression change in Tau compared to
WT, Aβ compared to WT and Tau;Aβ compared to WT on day 1 of adulthood. (B) Venn
diagram of significant genes downregulated and (C) upregulated in Tau, Aβ and Tau;Aβ
compared to WT on day 1 of adulthood. (D) Wormbase tissue enrichment and GO
enrichment of the genes downregulated in day 1 Tau;Aβ strain compared to WT. (E)
Wormbase tissue enrichment and GO enrichment of the genes upregulated in day 1
Tau;Aβ strain compared to WT. (F) Based on transcriptomic aging clock from [41], the
predicted biological age of Tau, Aβ and Tau;Aβ are increased compared to WT.

1083

1084

1085