T:B cell communication in ectopic lymphoid follicles in CNS autoimmunity

Anna Kolz[1], Isabel Bauer[1], Clara de la Rosa del Val[1], Vladyslav Kavaka[1], Anna Sophie Thomann[1], Rosa Schmitz[1], Martin Kerschensteiner[1], Eduardo Beltran[1], Naoto Kawakami[1], Anneli Peters[1]

[1] Institute of Clinical Neuroimmunology, University Hospital and Biomedical Center of the LMU Munich, Germany

Correspondence to: Anneli.Peters@med.uni-muenchen.de

Abstract

Meningeal ectopic lymphoid follicle-like structures (eLFs) have been described in multiple sclerosis (MS) and its animal model experimental autoimmune encephalomyelitis (EAE), but the cellular processes within and their role in CNS autoimmunity is not clearly understood. Here, we employed the Th17 adoptive transfer EAE model, which features formation of large, numerous eLFs, to analyze the cellular processes within these structures. We show that clusters of activated B cells and B1/Marginal Zone-like B cells are overrepresented in the CNS and identified B cells poised for undergoing antigen-driven germinal center (GC) reactions and clonal expansion in the CNS. Furthermore, we found evidence for enhanced capacity for antigen presentation and immunological synapse formation in CNS B cells. To visualize Th17:B cell communication in eLFs, we labeled Th17 cells with a ratiometric calcium sensor, allowing us to study their interactions with tdTomato-labeled B cells in real-time using intravital microscopy of the CNS. Thus, we demonstrate for the first time that there is extensive communication and long-lasting contacts between T and B cells in meningeal eLFs, and that B cells are able to reactivate T cells. Consistent with these findings, we show that CNS T cells depend on CNS B cells to maintain a highly pro-inflammatory cytokine profile. Our data suggest that interaction of T and B cells in meningeal eLFs in our model not only promotes differentiation and clonal expansion of B cells, but also leads to reactivation of CNS T cells and thereby supports smoldering inflammatory processes within the CNS. Thus, our results may provide a direction for future research into the function of eLFs in MS.

Introduction

Multiple sclerosis (MS) is an autoimmune disease of the central nervous system (CNS) characterized by chronic inflammation and demyelination, resulting in neuronal damage and progressive paralysis. Autopathogenic T cells are thought to be important disease drivers and, to our current knowledge, they are primed in the periphery before they invade the CNS, where they presumably become reactivated by contact with the autoantigen, and subsequently orchestrate the inflammatory cascade. Both Th1 and Th17 cells contribute to pathogenesis in MS and its animal model experimental autoimmune encephalomyelitis (EAE), and during the past two decades, compelling evidence has also suggested a pivotal role of B cells. However, it is not clear how T cells and B cells cooperate to induce and

propagate disease. In MS, ectopic lymphoid follicle-like structures (eLFs) were found in the leptomeninges of patients with a chronic progressive disease course (Magliozzi et al., 2007; Serafini et al., 2004), but also in earlier stages of relapsing-remitting MS (Lucchinetti et al., 2011). Initially, they were thought to be restricted to brain meninges, where they are localized mainly in sulci, but a recent study showed that they can also form in spinal cord meninges (Reali et al., 2020). These aggregates show varying degrees of maturation, ranging from loose clusters of cells to highly organized follicle structures, which contain plasma cells, proliferating B cells, T cells and FDCs, and therefore are suggestive of germinal center formation (Mitsdoerffer and Peters, 2016). Clinical data show a correlation between the presence of eLFs and a more severe disease course – younger age at disease onset, irreversible disability, and death, as well as a more pronounced cortical pathology in the tissue underneath eLFs (Howell et al., 2011; Magliozzi *et al.*, 2007; Magliozzi et al., 2010; Serafini et al., 2007; Serafini *et al.*, 2004). Therefore, these follicles have been postulated to maintain or exacerbate chronic inflammation directly in the CNS. Pathogenic functions of eLFs could be due to both *de novo* generation and maturation of autoreactive B and T cell clones, as well as to reactivation and expansion of pre-existing autoreactive T and B cells recruited from the periphery. In addition, eLFs may be a source of pathogenic soluble factors such as auto-antibodies and pro-inflammatory cytokines that affect the underlying regions. However, they could also harbor regulatory processes aimed at shutting down excessive inflammation. Since data from human eLFs is based on and largely restricted to histological analysis of post-mortem CNS material, our knowledge of cellular processes going on in eLFs is still limited.

To study eLFs in more detail, here we are employing the Th17 adoptive transfer EAE model, which features formation of eLFs within the meninges (Jager et al., 2009; Peters et al., 2011). Meningeal eLFs were also identified in other EAE models differing regarding genetic background, mode of EAE induction and auto-antigen (Dang et al., 2015; Kuerten et al., 2012; Magliozzi et al., 2004; Pikor et al., 2015), but the Th17 adoptive transfer EAE model exhibits the largest and most numerous eLFs. Due to the relatively short and severe disease course, the Th17 EAE model is not suitable to study the clinical effects of eLFs, but is rather a model for the early stages of eLF formation. Importantly, the large number and size of the eLFs and the massive recruitment of B cells allow for in depth phenotyping via single cell RNA sequencing and even for studying cellular interactions via live imaging.

Across all EAE models, eLFs are characterized by the presence of B cell zones, whereas T cells appear to be more scattered. In addition, meningeal eLFs were reported to be positive for germinal center markers (Batoulis et al., 2015; Kuerten et al., 2012; Lehmann-Horn et al., 2016; Peters et al., 2011) and to recapitulate some features of SLOs (Mitsdoerffer and Peters, 2016). However, a detailed characterization of the cellular components of meningeal eLFs on a transcriptional or protein-level is still lacking. Using adoptive transfer EAE, Th17 cells, but not other Th cell subsets, were demonstrated to induce the formation of eLFs (Peters et al., 2011). Indeed, several lines of evidence speak in favor of a special relationship between Th17 cells and B cells. On the one hand, Th17 cells produce IL-21, which promotes B cell proliferation, maturation and isotype switching, and they are excellent B cell helpers in SLO GC reactions (Mitsdoerffer et al., 2010). In the Th17 adoptive transfer EAE model, transferred T cells were shown to express markers of Tfh cells, as well as GC markers in the CNS (Peters et al., 2011), suggesting that they may provide help to the B cells on site. On the other hand, B cells are efficient APCs for the activation of T cells with the same antigen specificity (Lanzavecchia, 1990), and they can also produce IL-6 (Barr et al., 2012), an

important cytokine for pathogenic Th17 cell differentiation in EAE (Molnarfi et al., 2013).
Since B cells are well positioned in meningeal eLFs to interact with infiltrating T cells, they
are a good candidate as APCs for the reactivation of T cells in the CNS, which is crucial for
disease development and progression. However, whether B cells fulfill this function in eLFs
has not been investigated.

The aim of this study was to investigate the unique relationship of Th17 cells and B cells in
EAE development, and, in particular, to assess the nature of their interaction in eLFs.
Combining histological and flow cytometric analyses with single cell transcriptomics and
intravital imaging of meningeal eLFs, we have gathered evidence that eLFs have primarily
pro-inflammatory functions in the Th17 adoptive transfer EAE model.

Results
Th17 cells induce formation of ectopic lymphoid follicles in CNS meninges

To be able to study cellular processes in eLFs, we first established adoptive transfer EAE
according to Jäger et al. [72], a model in which disease is induced by transfer of *in vitro*
differentiated pure MOG-specific Th1 or Th17 cell subsets into immunocompetent recipient
mice (Figure S1A+B). As previously shown, Th1 and Th17 cells induced EAE independently
of each other with comparable clinical disease courses and similar severity (Figure S1C). In
the inflamed CNS, transferred Th1 and Th17 cells maintained stable expression of their
signature cytokine IFNγ and IL-17A, respectively, and Th17 cells can additionally adopt IFNγ
production (Figure S1D+E). To assess B cell recruitment to the CNS, we compared B cell
frequencies among CNS infiltrating cells between Th1 and Th17 recipient mice, which had
been on a high score for several days to allow for continued immune cell infiltration. We
detected a significant increase in both frequency and absolute numbers of B cells in the CNS
of Th17 recipients compared to Th1 recipients (Figure 1A-C). We next performed a detailed
immunohistochemical analysis of CNS lesions to investigate the localization, structural
organization, and composition of cellular infiltrates in Th17-EAE mice. Following identification
of regions with lymphocytic infiltrates by Giemsa staining (Figure S2A), neighboring sections
were stained for T cells (CD3), B cells (B220) and laminin, which visualizes the basal lamina
of blood vessels and the pia mater. In line with previous observations in Th17 recipient mice
(Peters et al., 2011; Pikor et al., 2015), we found eLFs distributed without a preferential
location throughout the meninges of the spinal cord (Figure S2C). eLFs consisted of
accumulated T and B cells and showed varying degrees of organization and size: while some
contained loose aggregates of a few cells, others were composed of large amounts of T and
B cells and extended over wide distances of up to several millimeters (Figure 1D+E). Based
on their unique morphology and expression of CD11b, we also identified macrophages in
close vicinity, but also within the follicle structures (Figure S2B). In addition to spinal cord
meninges, enormous eLFs were also found in many different areas of the brain, including the
subarachnoid space at the base of the brain (Figure 1F) and the cortex (Figure 1G), and
appeared to originate from the meninges and invade the underlying parenchyma. eLFs often
exhibited separate T and B cell zones in our model, thereby mimicking the structural
organization of SLOs. Notably, all eLFs were characterized by prominent B cell clustering,
with B cells being almost exclusively confined to these meningeal aggregates. In contrast, T
cells were more widely distributed and detected both inside eLFs in large numbers, as well
as in the underlying subpial parenchymal tissues (Figure 1D-G). Staining for Ki67 revealed

eLFs to be sites of T and B cell proliferation (Figure 1H). To evaluate the degree of eLF
maturation, we determined expression of GC markers on B and T cells. In many eLFs, only
few scattered B cells expressed the GC markers PNA or AID (Figure 1I+J), however, some
eLFs exhibited clusters of AID$^+$ B cells, indicative of GC formation (Figure 1K). In addition,
PNA$^+$ B cells (Figure S3A), isotype-switched IgM$^-$ IgD$^-$ B cells (Figure S3B), and a small
population of plasma cells (Figure S3C) were detected among CNS-infiltrating cells via flow
cytometry. Furthermore, a fraction of CNS-infiltrating T cells expressed the GC and activation
marker GL7 with transferred 2D2 Th17 cells showing higher frequency of GL7$^+$ cells than
endogenous T cells from the recipient (Figure S3D). Taken together, we confirmed that the
Th17 adoptive transfer EAE model prominently features formation of eLFs in both spinal cord
and brain meninges. Considering the high frequency of recruited B cells, the high abundance
and large size of eLFs, as well as our initial evidence that at least some of the eLFs are
mature enough to harbor GC B and T cells, our model is well suited to study the cellular
processes in CNS eLFs in more detail.

152 CNS-infiltrating B cells are highly activated and poised for undergoing
153 GC reactions

To develop a clearer picture of properties and functions of B cells in meningeal eLFs, we
characterized the transcriptome of CNS-infiltrating B cells from Th17 recipients by single-cell
RNA sequencing and compared to peripheral B cells from spleen and cervical lymph nodes
(cLN) of the same animal. Unbiased clustering of the single cell transcriptomes revealed ten
individual B cell clusters varying substantially in size and containing B cells of all three
compartments. Interestingly, CNS B cells showed a distinct distribution of B cells into the
different clusters, indicating them to be phenotypically different from peripheral B cells
(Figure 2A, Figure S4A+B). Thus, cluster 2 and especially cluster 6-9 were highly enriched
for CNS B cells, most pronounced for cluster 6, in which 93.5 % of cells belonged to the CNS
fraction (Figure 2B). Gene set enrichment analysis against the GO biological processes
database revealed that cluster 6 B cells are metabolically highly active and have a high
energy demand as they showed upregulation of pathways associated with respiration, ATP
biosynthesis, glycolysis, fatty acid metabolism and oxidative phosphorylation (Fig. 2C, Fig.
S4C). In line with this, they showed highly increased expression of many ribosomal proteins
(Fig. S4A and Table S1), which have been shown to be upregulated in early GC B cells in
response to mTOR signaling in order to prepare for the increased translational demand of an
imminent GC reaction (Ersching et al., 2017). Notably, both mTOR and Myc signaling, which
precede entry into a GC reaction, were highest in cluster 2, 9, 6 and 7 B cells, and always
higher in CNS B cells compared to peripheral B cells within each cluster (Fig. S4C). Pathway
analysis showed that cluster 7 B cells are highly activated / differentiated and
transcriptionally very active (Fig. 2D). Thus, *Nr4a1*, *Egr1, Egr2* and *Egr3*, which are quickly
induced upon B cell activation through the BCR, are among the top 10 up-regulated genes in
cluster 7 (Fig. S4A and Table S1). Cluster 9 cells showed upregulation of pathways
associated with cell cycle, mitosis, DNA replication, and proliferation and accordingly *Mki67*
was among the top 10 upregulated genes in this cluster (Fig. 2E, Fig. S4A and Table S1).
Importantly, the CNS-enriched clusters 2, 6, 7 and 9 showed increased expression of genes
associated with a light zone (LZ) germinal center B cell signature (Victora et al., 2010), with
higher expression in CNS B cells compared to peripheral B cells within each cluster (Figure
2F). Thus, for example the LZ GC markers *Cd83* and *Fas*, as well as the cyclin *Ccnd2* and
the transcription factor *Bhlhe40*, which have been associated with LZ GCs (Rauschmeier et

al., 2022; Victora et al., 2010), were among the top 50 upregulated genes in both cluster 6
and 7 (Table S1). These data indicate that the phenotype of B cells especially in cluster 6
and 7, which are highly overrepresented in the CNS, is compatible with LZ GC B cells,
suggesting that a considerable fraction (about 19.5 %) of CNS B cells is poised to undergo
GC reactions. In contrast, only Cluster 9 B cells showed elevated expression of genes
associated with dark zone (DZ) germinal center B cells (Victora et al., 2010) (Figure 2G),
including for example *Stmn1, Cenpa,* and the cyclin *Ccnb2*. This is in line with the increased
proliferative capacity of these cells suggested by the pathway analysis (Fig. 2E) and the high
level of oxidative phosphorylation (Fig. S4C), suggesting that with cluster 9 only a very small
fraction (1.4 %) of B cells in eLFs has progressed to DZ GC B cells to begin clonal
expansion.

CNS-infiltrating B cells express markers associated with B1a / MZ B cells and undergo clonal expansion

In addition to the clusters described above, also cluster 8 was highly enriched for CNS B
cells. A closer look at the upregulated genes in this cluster, such as *Mzb1*, *Fcrl5* and *Cd9*
suggested similarities to marginal zone (MZ) and B1 cells (Table S1). Indeed, comparison
with publicly available gene signatures from B1a cells from the spleen and peritoneum and
MZ B cells from the spleen confirmed that cluster 8 represents a population of B1a / MZ-like
B cells in the CNS (Figure 3A). The only other cluster that showed elevated expression of
B1a and MZ markers was cluster 1, which consists mainly of peripheral B cells and contains
the MZ B cells in the spleen (Fig. 3A). In addition to the transcriptional analysis, we
confirmed the presence of MZ-like B cells in the CNS by analyzing CD1d protein levels via
flow cytometry (Figure 3B). In Th17-EAE mice, CNS-infiltrating cells were analyzed for the
presence of IgM^{high} IgD^{low} CD21^{high} CD1d$^+$ MZ B cells. In six out of seven mice, a small and
dim MZ-like B cell population was detected (Figure 3B, left panel), whereas one mouse
contained a more substantial fraction of MZ-like B cells in the CNS (26 % of IgM^{high} IgD^{low} B
cells) (Figure 3B, right panel). In addition, spinal cord cryosections of Th17-EAE mice were
stained for T cells (CD3), B cells (B220) and CD1d for confocal microscopy, thereby
confirming the presence of scattered CD1d$^+$ B cells in eLFs in the CNS (Figure 3C).
Considering that CD1d is not necessarily expressed on B1a cells, this gating and staining
strategy may not have included all the cells in cluster 8.

Analysis of the BCR repertoire revealed the existence of expanded B cell clones, and
interestingly, the clones with the highest expansion factors preferentially belonged to cluster
8, especially in the CNS (Figure 3D, S5A+B). Importantly, 55.5 % of expanded cells
represented B cells from the CNS, whereas 28.8 % and 15.5 % were derived from spleen
and cLN, respectively (Figure 3E). This preferential localization of expanded clones may
indicate that infiltrating B cells undergo clonal expansion in the inflamed CNS. The clones
with the highest expansion factors in the CNS were predominantly localized in cluster 8, but
we also found clones with lower expansion factors in the CNS in cluster 9 and 4 (Figure 3F).
Since B cells were sorted based on their expression of CD19, plasma cells, which do not
express CD19 anymore, were not included in this analysis. Therefore, the expanded CNS B
cells could be in the process of differentiating into plasmablasts. Indeed, only cluster 8 and
cluster 9 B cells showed increased expression of plasmablast-signature genes (Figure 3G).
Considering that we identified cluster 9 B cells in the CNS to carry features of DZ GC B cells
(Fig. 2E, G), these data suggest that cluster 9 B cells are starting to undergo clonal

expansion with still low expansion factors and are also upregulating the plasmablast
differentiation programme at the beginning of an antigen-driven GC reaction. In contrast,
cluster 8 B cells have much higher expansion factors but show lower expression of GC
markers, indicating that they may differentiate into plasmablasts independent of classical GC
reactions. In addition, cluster 8 B cells showed upregulation of pathways associated with
regulation of immune responses (Figure S5C) and expressed for example *Ctla4* and *Apoe*
among the top 50 upregulated genes Table S1). In addition, by comparing the transcriptome
of non-expanded and expanded cells in cluster 8, the expanded cells were found to
upregulate the expression of genes associated with plasma cell differentiation and MZ B cells
or B1 B cells, for example *Fam46c*, *Emb* and *Gpx4* (Figure S5D).

As we observed that the CNS in Th17 recipient mice supports clonal expansion of B cells
and harbors B cells poised to undergo GC reactions, we next investigated whether
development and expansion of MOG-specific B cells from the polyclonal repertoire occurs in
our model. We first determined the presence of a MOG-specific IgG1 response in the serum
of Th17 and Th1 recipients via ELISA (Figure S6A). At late peak of clinical disease, many
Th17 recipients showed high levels of MOG-specific IgG1 antibodies compared to very low
levels before EAE onset. Some Th1 recipients also exhibited elevated antibody titers at the
peak of disease compared to very low levels in mice, which were analyzed at an early
disease stage before B cells are starting to infiltrate into the CNS. However, on average
MOG-specific IgG1 titers in Th17 recipients were about 3-fold higher than in Th1 recipients.
To determine whether CNS eLFs contain MOG-specific B cells, we optimized detection of
MOG-specific B cells via IHC by staining CNS cryosections with a fluorescently labeled
tetramerized MOG-protein, which was validated on LN cryosections from IgH^{MOG} mice
carrying MOG-specific B cells (Figure S6B). Staining of CNS cryosections from Th17-EAE
mice revealed that most eLFs contained only very few MOG tetramer+ B cells (Figure S6C,
left panel). However, occasionally some eLFs displayed a massive accumulation of MOG
tetramer+ B cells (Figure S6C, right panel). Together, these data indicate that transferred
MOG-specific Th17 cells recruit B cells with the same antigen specificity from the
endogenous pool and that maturation and/or expansion of MOG-specific B cells may at least
partially happen in meningeal eLFs.

CNS-infiltrating B cells respond to T cell cytokines and show enhanced APC function

Our transcriptomic analysis suggests that many of our CNS B cells in Th17 adoptive transfer
EAE are highly activated, poised for entering GC reactions and undergoing clonal expansion
(Fig. 2 and 3). In addition to cellular interactions, phenotype and function can also be shaped
by cytokines. Thus, we evaluated whether we could find evidence of cytokine-induced
signaling in CNS B cells. B cells are known to strongly respond to cues from both type I and
type II interferons and thus we compared gene expression in our clusters to publicly available
gene signatures induced by IFNy and IFNa. Although, some IFNy is produced by transferred
Th17 cells as well as by endogenous T cells in our model, primarily cluster 5 B cells showed
evidence of IFNy-driven signaling (Figure 4A) including expression of *Stat1* and interferon
response factors such as *Irf7* (Fig. S4A and Table S1), and this was slightly enhanced in
CNS B cells compared to peripheral B cells. Cluster 5 B cells were also the only cluster
showing upregulation of IFNa-induced genes, however, here CNS B cells did not show an
enhanced response compared to peripheral B cells (Figure S7A), suggesting that IFNa is not

part of the CNS-specific cytokine milieu in our model. In contrast, all of the CNS-enriched B
cell clusters exhibited elevated expression of genes induced by IL-21, and expression was
always higher in CNS B cells compared to peripheral B cells from the same cluster (Fig. 4B).
Since IL-21 is produced by Th17 cells and is also a T helper cell-derived cytokine important
for B cell responses in GC reactions, this speaks for T cells providing help to B cells to
initiate GC reactions in the CNS. Indeed, expression levels of IL-21 induced genes were
highest in cluster 2, 7 and 9 (Figure 4B), which were shown to express markers of LZ and DZ
GC B cells, respectively (Figure 2F,G). In addition, CNS B cells strongly responded to TNFα,
while peripheral B cells did not (Figure 4C). This suggests that TNFα, which is produced by T
cells but can also be released by myeloid cells and B cells themselves, is an important
component of the CNS-specific cytokine milieu in our model. The highest levels of TNFα-
induced gene expression were detected in cluster 7 B cells, which also showed high levels of
Nr4a1 expression indicating that TNFα may even be a driver of B cell activation in CNS B
cells. In line with this, cluster 7 followed by cluster 6 and 9 displayed increased expression of
B cell activation signature genes (Figure 4D), as well as genes associated with APC function
and immunological synapse (IS) formation (Figure 4E). We confirmed the upregulation of
molecules involved in B cell activation, APC function and IS formation on CNS B cells by flow
cytometry. Thus, we found that the frequency of activated CD69$^+$ and CD83$^+$ B cells was
strongly increased in CNS B cells compared to splenic B cells (Figure 4F + S7B). In addition,
CNS B cells showed highly elevated expression levels of MHC-II, as well as elevated
expression levels of the costimulatory molecule CD40, and increased frequencies of the
costimulatory molecule CD86, which together with MHC-II form the central supramolecular
activation complex (SMAC) of the immunological synapse. Furthermore, CNS B cells also
showed increased levels of ICAM-1, which binds to LFA-1 on T cells and forms the
peripheral SMAC of the IS. In line with the increased response to IL-21 (Fig. 4B), CNS B
cells also expressed higher levels of IL-21R than peripheral B cells (Figure 4G + S7C).
Taken together, our transcriptomic and flow cytometric analysis suggests that CNS B cells
show enhanced capacity for antigen presentation, co-stimulation and formation of
immunological synapses.

T cells form long-lasting contacts with B cells in meningeal eLFs

Meningeal eLFs have been proposed to fuel chronic inflammation directly in the CNS by
providing a good microenvironment for the interaction of T and B cells and our phenotypic
analyses of CNS B cells show their enhanced capacity for interaction with T cells. However,
to our knowledge, it has never been demonstrated whether and to what effect T and B cells
interact in meningeal eLFs. Therefore, we employed intravital two-photon microscopy to
visualize the behavior of T and B cells in meningeal eLFs *in situ* and, in particular, to
determine whether B cells are able to reactivate T cells and thereby support inflammatory
processes within the CNS. To fluorescently label T cells, we used the ratiometric calcium
sensor Twitch-2B, which allows for studying T cell activation in real-time by visualizing the
increase of the intracellular calcium concentration upon TCR ligation (Kawakami, 2016). An
overview of the experimental set-up is given in Figure 5A. During their differentiation into
Th17 cells, MOG-specific 2D2 T cells were transduced with a retrovirus containing Twitch-
2B, and adoptively transferred into Mb1.Cre x ROSA26 tdTomato mice, which express
tdTomato specifically in B cells under control of the *Cd79a* promoter. EAE developed after 8
to 15 days, and using mice, which had been on a high score for several days to allow
maximal infiltration of B cells and eLF formation, intravital two-photon microscopy of spinal

cord eLFs was performed to visualize interactions of transferred Twitch-2B+ Th17 cells and
tdTomato+ B cells and subsequent activation events of the Th17 cells. First, we determined
the optimal time point for the retroviral transduction during the Th17 differentiation process
and found that the transduction efficiency was considerably higher during the secondary
phase (restimulation with anti-CD3 and anti-CD28) with up to 60 % Twitch-2B-expressing
Th17 cells (Figure S8A). Next, we confirmed that the transduced Th17 cells survived well *in*
vivo and infiltrated the CNS, as indicated by almost identical fractions of Twitch-2B+ cells in
the CNS compared to the *in vitro* culture (Figure S8B). Moreover, transduced T cells
appeared to be fully functional *in vivo*, since they showed the same cytokine profile as
Twitch-2B- T cells in the CNS (Figure S8C). In addition, tdTomato expression in CNS-
infiltrating B cells in Mb1.Cre x ROSA26 tdTomato Th17-EAE mice was detected by flow
cytometry as well as confocal microscopy of CNS cryosections stained for T cells (CD3) and
B cells (B220) (Figure S8D). In contrast, other CNS-infiltrating APCs such as CD11b+
macrophages showed frequencies of tdTomato+ cells below 4 % (not shown). Intravital
microscopy of spinal cord eLFs showed massive accumulation of Twitch-2B+ T cells and
tdTomato+ B cells in the meninges of all mice analyzed, revealing eLFs of substantial size
and structure. A representative image of the dorsal view onto a large eLF is shown in Figure
5B. In the central area, there was strong clustering of B cells with infiltrating T cells, whereas
in the outer region, there were mainly T cell aggregates. Since in this experiment, the
transduction rate of T cells ranked at 58 %, it can be assumed that the accumulation of
transferred T cells was even underestimated in this image. Zooming inside these eLFs
revealed extensive T:B cell contacts within these structures (Figure 5C). To compare the T
cell response of the EAE-inducing MOG-specific 2D2 Th17 cells to non-encephalitogenic
Th17 cells, a set of control experiments was performed, where ovalbumin (OVA)-specific OT-
II Th17 cells were transduced with the Twitch-2B sensor and transferred together with non-
transduced 2D2 Th17 cells into Mb1.Cre x ROSA26 tdTomato mice. Both 2D2 and OT-II T
cells were investigated regarding their interaction with B cells by monitoring the movement of
each individual cell in the 3D follicle structure during the entire length of the acquired videos.
Intriguingly, 2D2 T cells interacted more intensely with B cells than OT-II T cells. Around half
of the 2D2 T cells formed contacts with B cells compared to only 15 % of OT-II T cells
(Figure 5D). 2D2 T cells and B cells engaged in very long contacts, which often lasted during
the entire observation period of 35 minutes or in some cases even up to 50 min. In contrast,
OT-II T cells formed B cell contacts of much shorter duration (Figure 5E). Setting a threshold
of five minutes to define short and long contacts, as was suggested by measuring the contact
length of B cells and Tfh cells in regular GC reactions (Jacobsen et al., 2021), we found that
the majority of 2D2 T cells (82 %) engaged in long T:B cell contacts compared to only a third
of OT-II T cells (Figure 5F). Accordingly, 2D2 T cells engaged in T:B cell contacts also
showed a reduced velocity (mean 1.63 µm/min) compared to 2D2 T cells without B cell
contact (mean 2.47 µm/min). In contrast, OT-II T cells exhibited significantly higher motility
(mean 3.00 µm/min) than both 2D2 T cell groups (Figure 5G), indicating that the OT-II T cells
are being arrested less often due to lack of cognate interaction with B cells and other APCs
and therefore scan their surroundings faster in search of antigen.

B cells reactivate T cells in meningeal eLFs

367 To investigate whether 2D2 T cells become activated during B cell contact, we made use of
368 the ratiometric calcium sensor Twitch-2B, which represents a powerful tool to visualize
369 intracellular calcium signaling in activated T cells. **Error! Reference source not found.** 6A

depicts a single time point of a representative video, with the left and right part showing a fluorescence overlay of Twitch-2B$^+$ 2D2 T cells and tdTomato$^+$ B cells and a pseudocolor calcium ratio image indicating calcium signaling in T cells, respectively. Importantly, many 2D2 T cells showed elevated calcium levels and, in addition, appeared very blastic, indicating that they are highly activated (see also Movie 1). Quantitative analysis of the calcium ratios revealed that 2D2 Th17 cells exhibited highly increased calcium levels compared to OT-II Th17 cells. To correctly interpret increased intracellular calcium concentrations as calcium signaling, we set the threshold for the calcium ratio to the maximal level reached spontaneously by 95 % of OT-II T cells (0.534). For 2D2 T cells, 75.0 % of calcium ratios in T cells with B cell contact exceeded the threshold compared to 47.2 % of calcium ratios in T cells without B cell contact, showing that T cells exhibit more pronounced calcium signaling during T:B cell contact (Figure 6B). This quantitative analysis suggests that T cells can become reactivated by interacting with B cells in eLFs. Notably, the majority of 2D2 T cells (81.7 %) exhibited calcium spikes compared to less than a third of OT-II T cells (27.1 %) (Figure 6C). 2D2 T cells displayed long-lasting calcium signaling, which could often be observed during the entire observation period of up to 35 min, whereas OT-II T cells primarily showed calcium signaling of short duration (Figure 6D). To define short- and longterm calcium spikes, we used a cut-off at two minutes since calcium spikes longer than two minutes require antigen-dependent T cell activation (Kyratsous et al., 2017). We found that about 60% of 2D2 T cells showed long-term signaling compared to less than 7 % of OT-II T cells. 2D2 T cells also showed more short-term signaling than OT-II cells, whereas more than 70% of OT-II cells but less than 20% of 2D2 T cells showed no calcium signaling at all (Figure 6E). For better visualization of T:B cell interaction and the ensuing T cell response, Figure S8E depicts snapshots of representative 2D2 and OT-II T cells during B cell contact. The indicated 2D2 T cell formed a long-lasting contact with one and later two B cells. While the cell appeared to be arrested and only underwent minor changes in cell shape, it displayed high calcium signaling for over 30 min. In the second example, the 2D2 T cell showed low calcium levels before contacting the B cell (0 min) and increasing calcium levels upon contact formation (1.8 min - 26.2 min). In contrast, the indicated OT-II T cell did not show a difference in calcium levels before (0 min) and during contact with a B cell (1.4 min - 14.8 min). The difference in T cell response upon B cell contact became even more evident by comparing single cell tracks of 2D2 and OT-II T cells. The four representative 2D2 T cells formed B cell contact during the entire or a part of the observation period and exhibited pronounced calcium signaling during B cell contact (Figure 6F). In contrast, the calcium ratios of the two representative OT-II T cells remained unchanged during B cell contact (Figure 6G). Together, these data indicate that B cells in eLFs provide a reactivation stimulus to 2D2 T cells, but not to OT-II T cells.

T cells depend on CNS B cells to sustain pro-inflammatory effector functions in the CNS

Our intravital imaging showed intense T:B cell communication in eLFs that lead to increased Ca^{2+} signaling indicative of T cell reactivation. To investigate the functional consequences of these interactions, we transferred 2D2 Th17 cells into B cell-deficient Mb1-KO mice. At the peak of disease, infiltrating cells were recovered from the CNS, and T cells were analyzed for their cytokine profile. Transferred T cells could be distinguished from endogenous T cells due to their expression of the transgenic MOG-specific Vα3.2 TCR chain. Interestingly, the frequency of IL-17A$^+$, IFNγ$^+$ and IL-17A$^+$IFNγ$^+$ producers was significantly decreased among

417   transferred T cells in Mb1-KO mice compared to WT mice. Similarly, endogenous T cells,
418   which mainly produced IFNγ, showed lower levels of IL-17A$^+$, IFNγ$^+$ and IL-17A$^+$IFNγ$^+$ cells in
419   Mb1-KO mice (Figure 7A). However, in the spleen we did not find decreased T cell effector
420   cytokines in the absence of B cells (Figure 7B), strengthening the importance of the
421   reactivation stimulus provided by B cells in the eLFs in the CNS rather than in the periphery.

Discussion

In our study, we combined flow cytometric, transcriptomic and microscopic analyses to investigate the unique relationship of Th17 cells and B cells and, in particular, to shed light on their communication in meningeal eLFs. In our Th17 adoptive transfer EAE model, which features formation of large eLFs within the meninges of brain and spinal cord, these follicles contain clusters of proliferating T and B cells, show expression of germinal center markers, and are reminiscent of eLFs in MS patients. Our single-cell RNA sequencing analysis revealed that many CNS-infiltrating B cells are highly activated, poised for entering GC reactions and undergoing clonal expansion. In addition, CNS B cells respond to T cell cytokines and show enhanced capacity for antigen presentation, co-stimulation and formation of immunological synapses. Correspondingly, our intravital imaging provided the first real-time observation of intense T:B cell communication in eLFs and demonstrated that T and B cells form long-lived contacts, whereby B cells can provide a reactivation stimulus to T cells. Consequently, we showed that without this reactivation stimulus from B cells CNS T cells are not able to maintain their highly pro-inflammatory cytokine profile.

The histological analyses in our Th17 adoptive transfer EAE model suggest that eLFs appear in different stages of maturation, also within the same animal, and that many eLFs are not fully matured at the time point of investigation (Figure 1). Our single cell transcriptomic analysis of CNS B cells largely confirmed this observation, as we identified several clusters of substantial size of B cells expressing markers of activation and markers associated with LZ GC B cells, but only very few DZ GC B cells (Figure 2). Therefore, we suggest that our model features mostly eLFs harboring very early stages of GC reactions and this is probably due to the rather short disease duration in our model: since B cells only start accumulating in the CNS at the beginning of clinical symptoms, they had only 4-7 days to arrive and mature in eLFs until the recipients were analyzed. We speculate that the GC reaction in CNS eLFs would progress to more advanced stages with time, however, due to ethical and practical reasons this is not possible in our model. Nonetheless, we found elevated titers of MOG-specific antibodies in the serum at the late peak of disease compared to very low levels before disease onset (Figure S6A) and, additionally, rare eLFs containing massive amounts of MOG-specific B cells (Figure S6C). This shows the capacity of eLFs to support *de novo* generation or - at the very least -expansion of autoreactive B cells that were first primed in the periphery and then migrated to eLFs to expand. This speaks for a pathogenic role of eLFs since CNS-reactive autoantibodies are known to be pathogenic in CNS autoimmunity (Mader et al., 2020). In addition, eLFs have been hypothesized to mediate cortical damage in the underlying parenchyma by releasing soluble pro-inflammatory factors, which could include CNS-reactive antibodies but also cytokines. In fact, we found CNS B cells to respond strongly to TNFα (Figure 4C) and IL-21 (Figure 4B), an important Th17 effector cytokine, suggesting that the eLF cytokine milieu is enriched for those cytokines. Especially TNFα has pleiotropic effects on microglia, neurons, astrocytes and oligodendrocytes (Raffaele et al., 2020), and is therefore an interesting candidate for further investigation regarding the association of eLFs and cortical damage. Although IL-21 is primarily known to act on immune cells, IL-21 receptor is also expressed by cortical neurons and expression is up-regulated in MS patients (Tzartos et al., 2011). In a stroke model, IL-21 signaling in neurons was associated with increased autophagy (Clarkson et al., 2014), suggesting that IL-21 may also be an interesting candidate for mediating cortical damage underneath eLFs. The effect of other Th17-derived cytokines on B cells such as IL-17A was not investigated, since there are no gene expression signatures available. Surprisingly, IFNγ-driven signaling seemed less important for CNS B cells (Figure 4A), although some IFNγ is produced by transferred Th17 cells, as well as by endogenous T cells in the CNS (Figure S1E). However, the effect of IFNγ

on CNS B cells may be more pronounced in case of Th1 adoptive transfer EAE, where the
CNS T cells secrete massive amounts of IFNγ (Figure S1D) but much fewer eLFs are
formed.

Notably, we found a population of B1a / MZ-like B cells in the CNS (Figure 3A-C), which
could result from migration of B1a / MZ B cells from the spleen or peritoneum to the CNS.
However, it is also possible that the inflamed CNS provides a microenvironment, which
induces infiltrating B cells to adopt a B1a / MZ-like phenotype. In several autoimmune
settings, MZ-like B cells infiltrate the target tissue, e.g. the thyroid gland (Segundo et al.,
2001), the pancreas (Mariño et al., 2008) or the salivary glands (Groom et al., 2002).
Interestingly, a MZ-like population was also detected in CSF from MS patients suggesting
that a human counterpart of our cluster 8 B1a / MZ-like B cells may also exist in MS patients
(Lee-Chang et al., 2011). Our B1a / MZ-like B cells expand clonally (Figure 3D+F), but show
less evidence of T cell help-induced GC reaction (Figure 2F+G), which is in line with the
property of regular B1a / MZ B cells to operate more independently of T cells. Considering
the high expansion factors and expression levels of plasmablast-associated genes (Figure
3F+G), they are probably a rich source of antibodies, but it remains to be determined
whether these are autoreactive. It is important to note that normal B1a and MZ B cells can
also perform regulatory functions (Zhang, 2013). They are more potent IL-10 producers
compared to other splenic B cell subsets (Barr et al., 2012) and may play a protective role in
autoimmunity (Lenert et al., 2005; Matsushita et al., 2018). In fact, several upregulated genes
in cluster 8, such as *Fcrl5*, *Zbtb20*, *Cd9* and *Apoe* (Table S1), were recently defined as
markers of Bregs by scRNA-seq of B cells from different murine organs (Yang et al., 2021).
Therefore, it is possible that they play a regulatory function in the CNS, especially in later
disease stages and when differentiating into plasmas cells.

The importance of B cells as APCs in neuroinflammation has been reported in several EAE
studies (Bettelli et al., 2006; Krishnamoorthy et al., 2006; Molnarfi *et al.*, 2013; Parker Harp et
al., 2015; Weber et al., 2010). In our model, we demonstrated that CNS B cells reactivate
MOG-specific T cells in eLFs, as shown by our intravital microscopy (Figure 6B+F) and
adoptive transfer into B cell-deficient Mb1-KO mice (Figure 7). However, considering that the
B cell repertoire in our WT recipient mice is polyclonal and only few eLFs were positive for
MOG-specific B cells by MOG$_{tet}$-staining (Figure S6), B cells in eLFs cannot all be MOG-
specific. Thus, we speculate that antigen bound to MHC-II is transferred to B cells from other
APCs, which are also present in eLFs (Figure S2). Although, the mechanism requires further
investigations, this could happen either actively for example by trogocytosis (Schriek et al.,
2022), or passively by extracellular vesicles (Becker et al., 2021) derived from rare MOG-
specific B cells or from ubiquitous APCs including macrophages and DCs. Due to the
extensive tissue damage at this late peak of disease the area must be saturated with
released CNS autoantigens ready to be taken up and interestingly, extracellular vesicles
containing MOG were also shown to be increased in the CSF of MS patients during disease
activity (Galazka et al., 2018). Another point of consideration is that the specificity of the 2D2
TCR may not be completely restricted to MOG, as it was shown to also respond to
neurofilament (Krishnamoorthy et al., 2009). Nonetheless, we demonstrated that OT-II T
cells do not become reactivated by B cells in eLFs even though they are also differentiated
into Th17 cells (Figure 6B-E, G), indicating that the reactivation stimulus provided by the B
cells to the 2D2 T cells must be CNS antigen-dependent.

Taken together, our data suggest that interaction of T and B cells in meningeal eLFs not only
promotes differentiation and clonal expansion of B cells, but also leads to reactivation of CNS
T cells. Thus, in our model eLFs seem to primarily have pro-inflammatory function. Still, the
clinical significance of eLFs requires further investigation. In MS, formation of eLFs is
associated with a stronger disease course and subpial cortical as well as spinal cord

pathology (Howell et al., 2011; Magliozzi et al., 2007; Magliozzi et al., 2010; Reali et al.,
2020; Serafini et al., 2007; Serafini et al., 2004). In fact, the neuronal loss in the cortex
displays a gradient from the pia mater to the white matter (Magliozzi et al., 2010; Mainero et
al., 2015), suggesting that eLFs contribute to the pathogenesis of cortical damage. Due to
the short and severe disease course, our Th17-transfer EAE model is not suitable to study
the clinical impact of eLFs, but in the spontaneous OSE model, which features both
genetically modified MOG-specific T and B cells and a milder disease course that allows for
longer observation periods, a correlation between eLF presence and a more severe disease
course was found (Dang et al., 2015). In the same model, a recent study suggested an
immunoregulatory function of eLFs since eLF abrogation resulted in a higher disease burden
(Mitsdorffer et al., 2021). It is possible that eLFs predominantly form in severe disease cases
as a counter-regulatory process in an attempt to contain excessive inflammation, and this
would involve presence of regulatory T_{FR} cells and/or regulatory B cells producing anti-
inflammatory cytokines. Although, Foxp3[+] regulatory T_{FR} cells were not detected in eLFs in
post-mortem CNS tissue from MS patients (Bell et al., 2019), Mitsdörffer and colleagues
identified Bregs producing the immunoregulatory cytokines IL-10 and IL-35 in eLFs in the
OSE model (Mitsdorffer et al., 2021). Although we did not detect IL-10 or IL-35 in our
transcriptomic dataset, we also found markers of regulatory B cells expressed on B1a / MZ-
like cluster 8 B cells. Thus, it is likely that eLFs may provide a supportive environment for
Bregs and this may become more relevant at later time points, as differentiation to plasma
cells, which are a known source of anti-inflammatory cytokines, progresses further.

In contrast to the OSE model, Th17-mediated EAE employs non-transgenic B cells with an
endogenous repertoire and therefore, the B cell response in our model can be assumed to
better reflect B cell-related disease aspects in MS. The possibility that eLFs support *de novo*
generation and/or expansion of CNS-reactive B cell clones has clear immunopathological
implications and the increased MOG-specific antibody titers and rare MOG_{tet}-positive eLFs in
our model point in that direction, although more research is required to confirm eLFs as
sources of CNS-reactive B cells. Overall, it is possible that eLFs with an overall pro-
inflammatory and an overall regulatory impact co-exist in MS either in different subsets of
patients or in different phases of the disease. Therefore, both models are important to study
the role of eLFs in CNS autoimmunity, as each of them may represent one end of the
spectrum of what can happen within eLFs, ranging from immunoregulation to
immunopathology.

Materials and methods

Mice

C57BL/6 mice were purchased from Janvier Labs. C57BL/6 mice with a transgenic TCR specific for the MOG_{35-55} peptide, here referred to as 2D2 mice, have been described previously (Bettelli et al., 2003) and were purchased from The Jackson Laboratory. IgH^{MOG} mice on a C57BL/6 background, here referred to as Th mice, carrying an Ig heavy chain knock-in of the rearranged VDJ gene from the MOG-specific monoclonal antibody 8.18C5, have been described previously (Litzenburger et al., 1998) and were kindly provided by Prof. Hartmut Wekerle. Mb1-Cre mice on a C57BL/6 background carry a codon optimized Cre recombinase, replacing exon 2 and 3 of the *Cd79a* gene, which encodes the Ig-α subunit of the BCR and is exclusively expressed in B cells. This knock-in leads to abolishment of endogenous *Cd79a* gene function and expression of Cre recombinase under control of the *Cd79a* promoter (Hobeika et al., 2006). Mb1-Cre mice were purchased from The Jackson Laboratory. Homozygous mice are B cell-deficient and hereafter referred to as Mb1-KO mice. Mb1-Cre mice were crossed to ROSA26 tdTomato mice to obtain mice that are heterozygous for both Mb1-Cre and ROSA26 tdTomato, thereby expressing tdTomato specifically in B cells for intravital microscopy experiments. ROSA26 tdTomato mice on a C57BL/6 background carry a *loxP*-flanked STOP cassette and *tdTomat*o gene, both inserted into the ROSA26 locus. Upon Cre-mediated recombination, tdTomato fluorescence is robustly expressed in these mice (Madisen et al., 2010). ROSA26 tdTomato mice were purchased from The Jackson Laboratory. OT-II mice, carrying a transgenic TCR specific for the $ovalbumin_{323-339}$ peptide on a C57BL/6 background (Barnden et al., 1998), were purchased from The Jackson Laboratory.

All animals used in this study were housed and bred in specific pathogen-free conditions in the Core Facility Animal Models at the Biomedical Center of the Ludwig-Maximilians-Universität München. Animal experiments were designed and performed in accordance with regulations of the animal welfare acts and under approval by the animal ethics committee of the state of Bavaria (Regierung von Oberbayern) in accordance with European guidelines.

Isolation of cells from spleen and lymph nodes

Spleens were mashed and centrifuged at 300 g for 5 minutes (min) at 4 °C. To lyse erythrocytes, the cells were resuspended in ACK buffer (150 mM NH4Cl (Merck), 10 mM KHCO3 (Merck), 0.1 mM EDTA (Sigma)) and incubated for 1.5 min at RT. The reaction was stopped by adding 9 ml T cell medium (RPMI 1640 (Sigma) supplemented with 10 % heat-inactivated FBS, 1 % penicillin-streptomycin, 10 mM HEPES, 2 mM L-glutamine, 1 % non-essential amino acids, 1 mM sodium pyruvate and 50 µM β-mercaptoethanol), and the cell suspension was passed through a 70 µm cell strainer. After centrifugation, the cells were resuspended in T cell medium. Lymph nodes (LNs) were cut into pieces, mashed and filtered through a 70 µm cell strainer. Then, the cells were centrifuged at 300 g for 5 min at 4 °C and resuspended in T cell medium.

In vitro differentiation of T effector cells

Naïve CD4[+] T cells were purified from spleen and lymph nodes of 6-15 week old 2D2 mice or OT-II mice using the naïve CD4[+] T cell isolation Kit (Miltenyi Biotec) according to the

manufacturer`s instructions. After isolation of naïve CD4$^+$ T cells the rest of the cells (flow-
through from T cell purification) was irradiated at 35 Gy and used as APCs for costimulation
of differentiating T cells. Naïve T cells were cultured at a concentration of 1.5-2x10^6 ml^{-1} in T
cell medium in the presence of 7.5-10x10^6 ml^{-1} irradiated APCs (ratio T cells : APCs is 1 : 5)
and 2.5 µg/ml soluble anti-CD3 antibody (clone 145-2C11, BioXCell). Th1 cells were
generated by addition of IL-12 at a concentration of 10 ng/ml and anti-IL-4 antibody (clone
11B11, BioXCell) at a concentration of 10 µg/ml into the culture. For the generation of Th17
cells, naïve T cells were cultured with IL-6 at a concentration of 30 ng/ml, TGFβ at a
concentration of 3 ng/ml, IL-1β at a concentration of 20 ng/ml, and anti-IFNγ (clone XMG1.2,
BioXCell) and anti-IL-4 antibody (clone 11B11, BioXCell) at a concentration of 10 µg/ml. After
48h, Th1 cells and Th17 cells were split with medium containing 10 ng/ml of IL-2 and
medium containing 10 ng/ml of IL-23, respectively. All cytokines were purchased from
Biolegend except IL-23 (Miltenyi Biotec). The different T cell subsets were analyzed for
cytokine production after 4 days by intracellular cytokine staining and subsequent flow
cytometry. After 5-8 days of primary culture, T cells reached a resting state and were
restimulated at a concentration of 2x10^6 ml^{-1} for 48 h in the presence of plate-bound anti-CD3
(clone 145-2C11, BioXCell) and anti-CD28 (clone PV-1; BioXCell) antibodies both at 2 µg/ml
in fresh medium without any cytokines.

Retrovirus production

Phoenix-eco cells (Pear et al., 1993) were routinely cultured in TCM medium (DMEM
(Sigma) containing 10 % heat-inactivated FBS, 1 % penicillin-streptomycin, 2 mM L-
glutamine, 360 mg/l asparagine, 1 % non-essential amino acids, 1 mM sodium pyruvate and
57.2 µM β-mercaptoethanol). The cells were kept in a subconfluent state by splitting them
every 2-3 days. For production of retroviral particles, the cells were plated at 1.5-2x10^6 per
10 cm dish in 10ml TCM medium and allowed to attach for 18-24 hours. First, chloroquine
diphosphate (Sigma) was added at a concentration of 25 µM. Per dish, 12 µg pMSCV-Δneo-
Twitch-2B plasmid (kindly provided by Dr. Naoto Kawakami) and 3.5 µg pCL-Eco packaging
plasmid (kindly provided by Dr. Gurumoorthy Krishnamoorthy) were prepared in 438 µl H$_2$O.
Subsequently, 62 µl 2 M CaCl$_2$ (Riedel de Haën) were added followed by dropwise addition
of 500 µl 2X BES (50 mM N,N-bis(2hydroxyethyl)-2-aminoethanesulfonic acid (Sigma) + 280
mM NaCl (Sigma) + 1.5 mM Na$_2$HPO$_4$ (Roth) in H$_2$O) while vortexing. This mixture was
incubated for 20 min at 37 °C and then added dropwise to the cells. On the next day, the
medium was replaced with 8 ml fresh and warm TCM medium. After another 24 hours, the
cell culture supernatant was collected using a 10 ml syringe, filtered through a 0.45 µm filter
and kept at 4 °C. To every dish, 4 ml fresh and warm TCM medium was added and the cells
were incubated for another day. The supernatant was collected again, filtered through a 0.45
µm filter and pooled with the supernatant from the day before. Then, the supernatant was
concentrated by centrifugation at 1000 g at 4 °C through Amicon® Ultra 15 mL Centrifugal
Filters (100 K; Merck). Finally, the concentrated supernatant was snap frozen and stored at -
80 °C.

Retroviral transduction of Th17 cells

Retroviral transduction during primary phase: After 48 h of primary culture, Th17 cells were
split (usually 1:3) and transduced by adding 1:80-diluted retroviral supernatant and 4 µg/ml
polybrene (Sigma) in addition to 10 ng/ml IL-23 and 10 µg/ml anti-IFNγ antibody (all diluted in

pre-warmed T cell medium and calculated for culture volume). Then, cells were spin-infected
at 480 g for 90 min at 30 °C. On the next day, half of the medium was replaced with fresh
and warm T cell medium containing 10 ng/ml IL-23 and 10 µg/ml anti-IFNγ antibody. 48 h
after transduction, Twitch-2B expression was analyzed via GFP-signal by flow cytometry.

Retroviral transduction during secondary phase: 24 h after restimulation with anti-CD3/CD28
antibodies, half of the medium was taken up and Th17 cells were transduced by adding 1:80-
diluted retroviral supernatant and 4 µg/ml polybrene (Sigma) (all diluted in pre-warmed T cell
medium and calculated for culture volume). Then, cells were spin-infected at 480 g for 90
min at 30 °C. On the next day, half of the medium was replaced with fresh and warm T cell
medium. If cells were used for EAE experiments, cells were collected and washed for
adoptive transfer, while an aliquot of cells, with half of the medium being replaced with fresh
and warm T cell medium, was kept for determining the transduction efficacy. 48 h after
transduction, Twitch-2B expression was analyzed via GFP-signal by flow cytometry.

Adoptive transfer EAE

Naïve MOG-specific CD4$^+$ T cells, isolated from spleen and lymph nodes of 2D2 mice, were
differentiated into Th1 or Th17 cells and restimulated with anti-CD3 and anti-CD28
antibodies. 48 h after restimulation, cells were collected and washed three times with PBS. In
Th1-EAE, 1.7-3.2x10^6 IFNγ-producing cells were injected intraperitoneally (i.p.) into C57BL/6
recipients. In Th17-EAE, 4x10^6 IL-17A-producing cells were injected i.p. or intravenously (i.v.)
into C57BL/6 or Mb1-KO or Mb1Cre x ROSA26 tdTomato recipients. In case of 2D2 and OT-
II T cell co-transfer, 3x10^6 IL-17A-producing 2D2 cells and 2x10^6 IL-17A-producing OT-II cells
were injected together i.p. into Mb1Cre x ROSA26 tdTomato recipients. Animals of both
genders were used as donors, whereby male cells were injected only into male recipients.
Clinical disease developed 8-15 days post transfer. Mice were monitored daily for the
development of clinical signs of EAE, including classical and atypical disease signs,
according to the following criteria: 0, no disease; 0.5, decreased tail tonus; 1, limp tail or mild
balance defects; 1.5, limp tail and ataxia; 2, hind limb weakness, or severe balance defects
that cause spontaneous falling over; 2.5, partial hind limb paralysis; 3, complete hind limb
paralysis or very severe balance defects that prevent walking; 3.5, complete hind limb
paralysis and partial front limb paralysis; 4, complete front and hind limb paralysis or inability
to move body weight into a different position; 5, moribund state.

Isolation of infiltrating mononuclear cells from the CNS

Recipient mice were sacrificed at indicated time points and perfused through the left cardiac
ventricle with PBS. Brains were dissected by removing the skull using forceps and scissors,
while spinal cords were flushed out from the bone column using a 21 gauge needle attached
to a syringe filled with PBS. In addition, the vertebral column was opened to collect parts of
meninges, which were still connected to the vertebrae. Then, brains and spinal cords were
cut into pieces and digested for 30 min at 37 °C with DNase I (Sigma-Aldrich or Roche) at a
concentration of 1 mg/ml and collagenase D (Roche) at a concentration of 3.75 mg/ml.
Subsequently, the tissues were mashed and passed through a 70 µm cell strainer to prepare
a single-cell suspension, followed by centrifugation at 350 g for 10 min at 4 °C. The cell pellet
was resuspended in 5 ml 70 % Percoll (Cytiva), which was prepared by mixing 30 ml Percoll
with 18 ml Percoll Mix Solution (mixture of 90 ml 10X PBS and 264 ml H$_2$O). Then, the cell

suspension was slowly and carefully overlaid with 5 ml 37 % Percoll, which was prepared by
diluting the 70 % Percoll solution with 1X PBS, followed by centrifugation at 650 g for 30 min
at 21 °C. Then, mononuclear cells were collected from the interface between the 70 % and
37 % layer and washed twice with T cell medium to remove residual Percoll. Finally, cells
were either stimulated for intracellular cytokine staining or stained directly for flowcytometric
analysis.

Production of MOG tetramer

Mouse MOG_{1-125} protein (mMOG) was produced in human embryonic kidney cells expressing
the Epstein-Barr virus nuclear antigen-1 (HEK-EBNA1) according to previously published
protocols (Perera et al., 2013), a cell line designed for large-scale production of recombinant
proteins (Tom et al., 2008). Monomer production and tetramerization was performed as
previously described (Salvador et al., 2022). Briefly, purified mMOG monomers were
biotinylated using BirA ligase to enable multimerization. The biotin-binding protein
streptavidin was used to form stable tetramer structures consisting of four biotinylated
mMOG monomers. As the streptavidin was directly conjugated to AF 488, staining with MOG
tetramer could be used to identify MOG-specific B cells by flow cytometry or confocal
microscopy.

Flow cytometry and cell sorting

For intracellular cytokine staining, cultured or isolated cells were stimulated with 50 ng/ml
PMA (Sigma) and 500 ng/ml ionomycin (Sigma) in the presence of 0.7 µL/ml monensin
(GolgiStop; BD Biosciences) for 4 h. For surface stainings, the cells were washed with PBS.
To exclude dead cells and debris, the samples were stained with the Zombie Violet™ or
UV™ fixable viability kit (Biolegend) diluted 1:600 in PBS for 20 min at RT. Then, cells were
washed twice with FACS buffer (PBS containing 2 % FBS and 0.05 % sodium azide), and
unspecific binding of antibodies to Fc receptors was blocked by incubating the cells with
TruStain FcX™ (Biolegend) diluted 1:50 in FACS buffer for 15 min at 4 °C. The last step was
omitted when *in vitro* differentiated T cells were analyzed. Afterwards, cells were stained with
fluorophore-conjugated antibodies against surface markers diluted in FACS buffer for 20-30
min at 4 °C. If biotinylated antibodies were used, the samples were washed and
subsequently stained with fluorophore-conjugated streptavidin (SA) diluted in FACS buffer for
20 min at 4 °C. After two washing steps, cells were resuspended in FACS buffer for analysis.
For intracellular cytokine staining, cells were fixed for 30 min at 4°C with 0.4 %
paraformaldehyde (PFA, Merck KGaA) and permeabilized by washing twice with
permeabilization buffer (PBS containing 2 % FBS and 0.1 % saponin (Sigma)). Then,
samples were incubated with antibodies against intracellular markers diluted in
permeabilization buffer for 30 minutes at 4 °C, washed twice and resuspended in FACS
buffer for analysis. The following antibodies/reagents were used for flow cytometry: anti-
CD19 (1D3 or 6D5), anti-CD1d (1B1), anti-CD4 (RM4-5), anti-TCR Vα3.2 (RR3-16), anti-
B220 (RA3-6B2), anti-CD11b (M1/70), anti-CD21/CD35 (7E9), anti-CD11c (N418), anti-
CD69 (H1.2F3), anti-CD83 (Michel-19), anti-CD40 (HM40-3), anti-IL-21R (4A9), anti-ICAM-1
(YN1/1.7.4), anti-IFNγ (XMG1.2), anti-IL-10 (JES5-16E3), anti-IL-17A (TC11-18H10.1, all
Biolegend), anti-CD138 (REA104, Miltenyi), anti-GL7 (GL7), anti-IgD (11-26c.2a, all BD
Pharmingen), anti-CD45 (30-F11), anti-IgM (II/41), anti-CD86 (GL1), anti-MHC-II (NIMR-4),
anti-IL-17A (eBio17B7, all eBioscience), anti-TCR Vß5.1/5.2 (MR9-4, Invitrogen), PNA

(Vector Laboratories). Data were acquired on a BD LSRFortessa^{TM}, BD FACSverse^{TM} flow
cytometer or Beckman Coulter CytoFLEX S flow cytometer and analyzed using Flowjo
software (version 10.6.2; TreeStar). For sorting of B cells, surface staining of 1x10^6 cells from
spleen and cLN or 2.7-3.7x10^6 cells from CNS samples was performed in T cell medium for
20 min at 4 °C. After two washing steps, cells were resuspended in T cell medium containing
2 mM EDTA (Sigma), filtered through a 35 µm cell strainer and kept on ice. CD45^{high}
CD11b^{low} CD4$^-$ CD19$^+$ were sorted using a BD FACSAria^{TM} Fusion No1.

Immunohistochemistry (IHC) on cryosections

To increase the chances for finding eLFs in the CNS, IHC was performed using Th17-EAE
mice that had a high score for several days. The mice were perfused through the left cardiac
ventricle with 10 ml PBS and 10 ml 4 % PFA. Then, the brain was dissected by carefully
removing the skull using forceps and scissors. After removal of the outer muscle layer, the
spinal cord was dissected by snapping each vertebra open with forceps to carefully remove it
from the vertebral column in one piece leaving the leptomeninges as intact as possible.
Spleens and lymph nodes were also dissected as control tissue. After post-fixation in PBS
containing 4 % PFA for 2 h at 4 °C, the organs were dehydrated in PBS containing 30 %
sucrose for 24 h at 4 °C. Then, they were embedded in O.C.T. compound medium (Sakura
or VWR Chemicals), frozen using dry ice and stored at -80 °C. Subsequently, the brain was
cut into 20 µm coronal sections on a Leica CM1850 cryostat. From spinal cord, spleen and
lymph nodes, 10 µm sections (longitudinal for spinal cord and spleen) were prepared and
placed on slides (Superfrost Plus^{TM}, Thermo Scientific) and frozen at -80 °C.

Giemsa staining

To screen for eLFs, every 15th section of brain and spinal cord was fixed in 10 % methanol for
10 min, air-dried for 20-30 min and stained with Giemsa-stain (AppliChem), diluted 1:20 in
deionized water, for 45 min. After washing with deionized water, the sections were dried
overnight and mounted using Eukitt mounting medium (Sigma). Sections were screened for
immune cell infiltrates by light microscopy using a Leica DM2500 microscope with DMC2900
CMOS camera.

Immunofluorescence staining

Following identification of eLF-containing sections via Giemsa staining, neighboring sections
were stained with fluorescently labeled antibodies. After thawing, sections were fixed in cold
acetone for 10 min and air-dried for 20-30 min. Then, sections were outlined with a Pap-pen
and O.C.T. medium was rinsed off twice with PBS. Afterwards, sections were blocked with
PBS containing 1 % bovine serum albumin (BSA) and 1:300-diluted TruStain FcX^{TM}
(Biolegend) for 1 h at RT, and incubated with primary antibodies diluted in PBS containing 1
% BSA in a chamber humidified with PBS for 1 h at RT, followed by three washing steps with
PBS containing 0.1 % Tween20 for 7 min. Then, sections were stained with secondary
antibodies diluted in PBS containing 1 % BSA in a dark, humidified chamber for 1 h at RT.
When intracellular markers, such as activation-induced cytidine deaminase (AID), were
stained, the sections were incubated with antibodies diluted in PBS containing 1 % BSA and
0.1 % saponin to permeabilize the tissue. Following three washing steps with PBS containing
0.1 % Tween20 for 5 min, sections were mounted with VECTASHIELD Antifade Mounting
Medium with DAPI (Vector Laboratories) and analyzed by confocal microscopy using a Leica
SP8X WLL microscope. The following antibodies/reagents were used for
immunofluorescence staining: Syrian hamster anti-mouse CD3e (500A2), rat anti-mouse

B220 (RA3-6B2), mouse anti-mouse/human Ki67 647 (B56, all BD Pharmingen), rabbit anti-
mouse/human laminin (Abcam), rat anti-mouse/human CD11b AF 647 (M1/70), rat anti-
mouse CD1d (1B1, both BioLegend), rat anti-mouse/human AID (mAID-2, eBioscience),
PNA (biotinylated, Vector Laboratories), goat anti-hamster AF 568, goat anti-hamster AF 647
(both Invitrogen), goat anti-rabbit AF 568, goat anti-rabbit AF 647, goat anti-rabbit Cy3, goat
anti-rat AF 633, streptavidin AF 488 (all Life technologies).

MOG tetramer staining
Sections were fixed in cold acetone for 5 min and O.C.T. medium was rinsed off three times
with PBS for 5 min. Then, sections were outlined with a Pap-pen and blocked with PBS
containing 4 % BSA, 4 % goat serum and 1:300-diluted TruStain FcX™ (Biolegend) for 1 h at
RT in a humidified dark chamber. After washing three times with PBS for 5 min, sections
were incubated with rat anti-mouse B220 (1:200, clone RA3-6B2, BD Pharmingen) and
mMOGtet-SA488 (1:1000, housemade) diluted in PBS containing 4 % BSA and 1 % goat
serum overnight at 4 °C. Subsequently, sections were washed three times with washing
buffer (PBS containing 0.3 % Triton and 1 % goat serum) for 5 min and fixed in 4 % PFA for
30 min at RT. After three washing steps in washing buffer, sections were incubated with
rabbit anti-SA488 (1:10000, Life technologies) for 3 h and goat anti-rat AF 633 (1:200, Life
technologies) for 1 h diluted in PBS containing 4 % BSA and 1 % goat serum at RT in a
humidified dark chamber. After three washing steps in washing buffer, sections were
incubated with goat anti-rabbit Cy3 (1:1000, Life technologies) diluted in PBS containing 4 %
BSA and 1 % goat serum for 3 h at RT in a humidified dark chamber, followed by three
washing steps in washing buffer. Then, sections were mounted with VECTASHIELD Antifade
Mounting Medium with DAPI (Vector Laboratories) and analyzed by confocal microscopy
using a Leica SP8X WLL microscope.

812 Enzyme-linked immunosorbent assay (ELISA)

Anti-MOG IgG1 antibodies were measured in the serum of Th17- and Th1-EAE mice by
ELISA. 96 well high affinity, protein-binding plates (Nunc Maxisorp) were coated with rMOG
(housemade) at a concentration of 10 µg/ml in PBS and incubated overnight at 4 °C. After
washing four times with wash buffer (PBS containing 0.05 % Tween20), the plate was
blocked with blocking solution (PBS containing 10 % FBS) for 1-2 h at RT, followed by four
washing steps. Then, the serum samples and the antibody standard (mouse anti-MOG IgG1;
clone 8.18c5; housemade) were diluted in blocking solution, added to the plate and
incubated overnight at 4 °C. Following extensive washing, the plate was incubated with
biotin-labeled detection antibody rat anti-mouse IgG1 (clone A85-1; BD Pharmingen) at a
concentration of 1 µg/ml diluted in blocking solution for 1 h at RT. Then, the plate was
washed again and incubated with avidin-horseradish peroxidase (Av-HRP; eBioscience)
diluted 1:1000-1:2000 in blocking solution for 30 min at RT. After extensive washing, TMB
Substrate (BioLegend) was used to induce the Av-HRP-dependent color reaction, which was
stopped by addition of TMB Stop Solution (BioLegend), and the optical density (OD) was
measured at 450 nm. Samples were tested in duplicates and results were reported as mean
concentrations according to the standard curve.

Single-cell RNA sequencing (scRNA-seq)

Library construction and sequencing

For each mouse, B cells from CNS, spleen and cLNs were pooled (90.000 B cells in total), centrifuged and, after a washing step, resuspended in PBS containing 0.04 % BSA at a concentration of 1000 cells/µl. The viability of cells was confirmed by staining with trypan blue solution. Further processing of samples was performed using the 10x Chromium Single Cell 5` Solution (Chromium Next GEM Single Cell V(D)J v1.1 with Feature Barcoding Technology for Cell Surface Protein, 10x Genomics) according to the manufacturer`s instructions. Prior to loading on the chip, 5000 cells of each compartment were labeled with anti-mouse hashtag oligonucleotide (HTO) antibodies (TotalSeq™-C0301/C0302/C0303, Biolegend) and pooled together. Next, the B cells were processed with enzymes and gel beads, followed by partitioning in oil suspension to create gel beads in emulsion (GEMs) containing single cells. In GEMs, the cells` RNA was reverse-transcribed into cDNA, whereby all the cDNA from the same cell was labeled with a shared barcode. Then, the cDNA was pooled, purified and amplified according to the manufacturer`s protocol. The scRNA-seq, scBCR-seq and cell hashing libraries were constructed, followed by quality control using a Bioanalyzer 2100 (Agilent Technologies) and quantification using the Qubit dsDNA HS kit (Life Technologies). Next-generation sequencing (NGS) was performed on the Illumina HiSeq 1500 using read lengths of 28 base pairs (bp) read 1, 8 bp i7 index and 91 bp read 2 at the Gene Center Munich.

Data processing

To demultiplex samples, align the reads to the reference mouse mm10 genome, and process the raw data, we used the Cell Ranger Software (10X Genomics, v. 3.1.0). Following this, the unique molecule identifiers (UMIs) were summarized and the single-cells were filtered by the number of UMI counts identified. Then, we processed the gene matrices with R (v. 4.0.5) and the R package Seurat (v. 4.0.6 and 4.9.9) (Butler et al., 2018; Stuart et al., 2019). Cell hashing raw counts were normalized using centered log-ratio (CLR) transformation by comparing the counts of single cells to the geometric mean of the HTOs signal with subsequent logarithmic transformation. Cells with missing HTO signals or containing two different HTO-identifiers were removed from the further analysis. Further, the scRNA-seq count matrices were log-normalized and scaled by passing through only the cells containing more than 200 and less than 5000 genes with mitochondrial gene fractions below 5 %. Subsequently, the highly variable genes were identified with the vst method, and cells from each compartment were integrated using the canonical correlation analysis (CCA) method. The aligned BCR sequencing information was added into the metadata file of the Seurat object with subsequent removal of cells with missing BCR sequences. The cells were considered as expanded if at least 2 cells belonged to the same B cell clonotype. In the next step, 18413 cells were recovered and the corresponding PCA was computed using the previously identified highly variable genes. Based on the first 15 PCA dimensions, we constructed the Shared Nearest Neighbor (SNN) graph and clustered the cells by using the FindNeighbours and FindClusters functions of the Seurat R package. The following dimensional reduction was performed with the help of the Uniform Manifold Approximation and Projection (UMAP) method. One cluster, cluster 11, with markedly lower quality compared to the other clusters based on nFeature and nCount, was removed from further analysis. In addition, cluster 10 was identified as expressing macrophage/microglia markers and was therefore also excluded (based on AddModuleScore from Immgen Monocytes MHCIIhi blood vs Fo B cells spleen and Microglia CNS vs FoB form spleen).

For identification of differentially upregulated genes per cluster, compartment, or expansion condition we used the FindMarkers function based on the log-transformed count matrices of each single cell by applying the Wilcoxon test and subsequent p-value adjustment with the Bonferroni correction, with logfc.threshold = 0.

Pathway network analysis

For each cluster, GSEA (Mootha et al., 2003; Subramanian et al., 2005) PreRanked files were generated with the gene and avg_log2FC columns. GSEA v4.2.3 was run with the mouse GO BP database m5.go.bp.v2023.1.Mm.symbols (https://data.broadinstitute.org/gsea-msigdb/msigdb/release/2023.1.Mm/), without collapsing gene symbols, scoring_scheme classic, set_max 500, set_min 2, and otherwise default parameters. The results were exported to Cytoscape v3.9.1 (Shannon et al., 2003), where only the nodes with NES > 0, fdr_qvalue < 0.1 and fwer_qvalue < 0.5 were kept. AutoAnnotate from the EnrichmentMap app collection (Merico et al., 2010) was used to cluster and label the pathway networks, with Cluster Algorithm = MCL cluster, Label Column = Node Names, Max Words Per Label = 4, Minimum Word Occurrence = 1, Adjacent Word Bonus = 8, setting numbers and "GOBP" as extra Excluded Words.

B cell functional signatures

Functional signatures expressed by B cells were calculated with the AddModuleScore function from Seurat from manually curated gene lists based on ImmGen (https://www.immgen.org/), GO terms (https://www.informatics.jax.org/vocab/gene_ontology) and literature search.

APC/synapse signature: this signature was assembled by combining the GO gene set "lymphocyte co-stimulation" (GO:0031294, without T cell specific genes) and the GO gene set "immunological synapse" (GO:0001772, without T cell specific genes), and then adding the core APC signature described by (Jordão et al., 2019), the list of molecules described to be present in the immunological synapse formed by primary murine B cells by Cunha et al. (https://www.biorxiv.org/content/10.1101/2023.02.23.529674v1.full.pdf), and manually adding Icosl, Tnfsf9, Timd4, Pvr, and Il21r.

B1a signature: this gene set is derived from Immgen by population comparison using the ULI RNAsequencing datasets. B1a cells from spleen and B1a cells from the peritoneal cavity were compared against follicular B cells from spleen, and Nfatc1 and Mzb1 were added manually.

MZ signature: this gene set is derived from Immgen by population comparison using the ULI RNAsequencing datasets. Here, marginal zone B cells from spleen were compared to follicular B cells from spleen and Nfatc1 and Mzb1 were added manually.

LZ-GC and DZ-GC signature: these signatures were described by (Victora et al., 2010).

mTOR signaling: Hallmark_mtorc1_signaling (mouse) gene set from the MSigDB database.

Myc signaling: Hallmark_myc_targets_v1 (mouse) gene set from the MSigDB database.

OXPHOS: Hallmark_oxidative phosphorylation (mouse) gene set from the MSigDB database.

PB signature: this gene set is derived from Immgen by population comparison using the ULI RNAsequencing datasets. Here, plasmablasts and plasma cells from spleen and plasma cells from bone marrow were compared against follicular B cells from spleen.

IFNa response: Hallmark_interferon_alpha_response (mouse) gene set from the MSigDB database.

IFNg response: Hallmark_interferon_gamma_response (mouse) gene set from the MSigDB database.

<u>TNFa signaling:</u> Hallmark_TNFa_signaling_via_nfkb (mouse) gene set from the MSigDB
database.
<u>IL-21 response:</u> this signature is described on Immgen and includes all genes upregulated in
follicular B cells at least 2-fold by IL-21 relative to PBS treatment.
<u>B cell activation:</u> this signature represents the GO gene set "B cell activation involved in
immune response" (GO:0002312).

Data and code availability
All sequencing data (fastq files, processed data files and Seurat Object) from this study is
available upon request. Custom code used for analysis is available at Zenodo under the
same doi as the published manuscript.

Intravital two-photon microscopy
Animal preparation
Prior to surgery, the mice were anesthetized with MMF anesthesia consisting of midazolam
(5 mg/kg bodyweight (bw)), medetomidine (500 µg/kg bw) and fentanyl (50 µg/kg bw).
Subsequently, they were tracheotomized and ventilated with 1.5-2 % isoflurane, which was
supplied throughout the surgery and the entire imaging session. Thereby, the mice were
placed on a custom-made microscope stage, while their body temperature was stabilized at
37.5 °C using a heated pad. During imaging, breathing parameters, including concentrations
of inspiratory and expiratory gases as well as ventilation pressure, and electrocardiograms
were constantly monitored. The experimental procedure of spinal cord imaging has been
described previously (Bartholomäus et al., 2009; Mues et al., 2013)). In brief, a small imaging
window at level Th12/L1 of the spinal cord was prepared. First, a midline incision of about 2
cm was placed to open the skin and, subsequently, to detach the paravertebral musculature.
To reduce artefacts caused by breathing and thereby ensure stable imaging conditions, a
custom-made stage was used in order to fix three spines. Then, a laminectomy on the
central vertebra was performed using a dental drill (Foredom). Finally, low-melting agarose
was used to build a ring-shaped surrounding, which was filled with PBS to keep the tissue of
the opened spine hydrated and to allow using the water immersion objective.

Image acquisition
Time-lapse two-photon laser-scanning microscopy was performed using a SP8 multi-photon
microscope (Leica). The SP8 system was equipped with a pulsed InSight DS+ laser (Spectra
Physics). The emission wavelength was tuned to 841 nm and the fluorescent signal was
collected using a water immersion objective (25x, NA 1, Leica) and detected with external,
non-descanned hybrid photo detectors (HyDs) featuring 483/32 nm, 535/30 nm, 585/40 nm
and 650/50 nm bandpass filters. Images were recorded using a 1-2x zoom, whereby areas of
221x221 µm, 295x295 µm or 443x443 µm were scanned, and 49-115 µm z-stacks with 2.5-
4.3 µm step size were acquired with an acquisition rate of 21-22 s per z-stack.

Image processing and analysis
Time-lapse images were acquired using the LAS X software (Leica), followed by processing
and analysis using ImageJ (NIH). Before the stacks were z-projected with maximum intensity
to obtain two-dimensional movies, a Gaussian blur filter was applied. For Twitch-2B-
expressing cells, the cpVenus^{CD} (FRET) channel was divided by the mCerulean3 (CFP)
channel and, by changing to a fire lookup table, ratiometric pseudocolor images were
generated. In each time frame of the two-dimensional maximum intensity projection, the cell

shapes were outlined manually to create a region of interest (ROI) for analyzing the calcium
ratios. Since the bleed-through of CFP into the YFP channel was determined to account for
44 %, the FRET signal was corrected as follows: YFP = FRET − 0.44 x CFP. In order to
calculate the calcium ratio in every time frame, the average signal intensities of all pixels in
each ROI were used. Cell coordinates over time were used to calculate motility parameters
by using ImageJ. T:B cell contact analysis was performed by monitoring the movement of
each Twitch-2B[+] T cell and its interaction with tdTomato[+] B cells in the 3D follicle structure
using the LAS X software.

Statistical analysis

Statistical analysis was performed using GraphPad Prism software (version 7). The statistical
test chosen is described in the figure captions. For comparison of two groups, statistical
differences were calculated using the unpaired Student`s *t*-test for normally distributed data,
including Welch`s correction in case of unequal variances, or the Mann-Whitney U test if
Gaussian distribution could not be determined. For comparison of three groups, the one-way
ANOVA with Tukey`s test for multiple comparisons was applied.

Declaration of interests

987 The authors declare no conflict of interest.

Acknowledgements

We thank the Core Facility Flow Cytometry and the Core facility Bioimaging at the
Biomedical Center, Ludwig-Maximilians University Munich, for providing equipment, services
and expertise. We thank Isabel Gök and Klaus Hämäläinen for technical assistance. The
study was supported by the German Research Foundation (DFG) through the Emmy
Noether programme PE-2681/1−1 (to A.P.), through SFB TRR 128 (project B08 to A.P.;
project B10 to N.K.), through CRC TRR 152 (project P27; to A.P. and M.K.), through the
Heisenberg programme (KA2951/3-1 and KA2951/3-2 (264119140) to N.K. and through
research grant KA2951/2-2 (246754395) to N.K.

References

Barnden, M.J., Allison, J., Heath, W.R., and Carbone, F.R. (1998). Defective TCR expression
in transgenic mice constructed using cDNA-based alpha- and beta-chain genes under the
control of heterologous regulatory elements. Immunol Cell Biol *76*, 34-40. 10.1046/j.1440-
1711.1998.00709.x.
Barr, T.A., Shen, P., Brown, S., Lampropoulou, V., Roch, T., Lawrie, S., Fan, B., O'Connor,
R.A., Anderton, S.M., Bar-Or, A., et al. (2012). B cell depletion therapy ameliorates
autoimmune disease through ablation of IL-6-producing B cells. The Journal of experimental
medicine *209*, 1001-1010. 10.1084/jem.20111675.
Bartholomäus, I., Kawakami, N., Odoardi, F., Schläger, C., Miljkovic, D., Ellwart, J.W.,
Klinkert, W.E., Flügel-Koch, C., Issekutz, T.B., Wekerle, H., and Flügel, A. (2009). Effector T
cell interactions with meningeal vascular structures in nascent autoimmune CNS lesions.
Nature *462*, 94-98. 10.1038/nature08478.
Batoulis, H., Wunsch, M., Birkenheier, J., Rottlaender, A., Gorboulev, V., and Kuerten, S.
(2015). Central nervous system infiltrates are characterized by features of ongoing B cell-
related immune activity in MP4-induced experimental autoimmune encephalomyelitis.
Clinical immunology *158*, 47-58. 10.1016/j.clim.2015.03.009.
Becker, P.D., Ratnasothy, K., Sen, M., Peng, Q., Romano, M., Bazoer, J., Suvitra, E., Stout,
A., Hylton, S.G., Dorling, A., et al. (2021). B lymphocytes contribute to indirect pathway T cell
sensitization via acquisition of extracellular vesicles. Am J Transplant *21*, 1415-1426.
10.1111/ajt.16088.
Bell, L., Lenhart, A., Rosenwald, A., Monoranu, C.M., and Berberich-Siebelt, F. (2019).
Lymphoid Aggregates in the CNS of Progressive Multiple Sclerosis Patients Lack Regulatory
T Cells. Frontiers in immunology *10*, 3090. 10.3389/fimmu.2019.03090.
Bettelli, E., Baeten, D., Jager, A., Sobel, R.A., and Kuchroo, V.K. (2006). Myelin
oligodendrocyte glycoprotein-specific T and B cells cooperate to induce a Devic-like disease
in mice. The Journal of clinical investigation *116*, 2393-2402. 10.1172/JCI28334.
Bettelli, E., Pagany, M., Weiner, H.L., Linington, C., Sobel, R.A., and Kuchroo, V.K. (2003).
Myelin oligodendrocyte glycoprotein-specific T cell receptor transgenic mice develop
spontaneous autoimmune optic neuritis. The Journal of experimental medicine *197*, 1073-
1081. 10.1084/jem.20021603.
Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell
transcriptomic data across different conditions, technologies, and species. Nat Biotechnol *36*,
411-420. 10.1038/nbt.4096.
Clarkson, B.D., Ling, C., Shi, Y., Harris, M.G., Rayasam, A., Sun, D., Salamat, M.S.,
Kuchroo, V., Lambris, J.D., Sandor, M., and Fabry, Z. (2014). T cell-derived interleukin (IL)-
21 promotes brain injury following stroke in mice. The Journal of experimental medicine *211*,
595-604. 10.1084/jem.20131377.
Dang, A.K., Tesfagiorgis, Y., Jain, R.W., Craig, H.C., and Kerfoot, S.M. (2015). Meningeal
Infiltration of the Spinal Cord by Non-Classically Activated B Cells is Associated with Chronic
Disease Course in a Spontaneous B Cell-Dependent Model of CNS Autoimmune Disease.
Frontiers in immunology *6*, 470. 10.3389/fimmu.2015.00470.
Ersching, J., Efeyan, A., Mesin, L., Jacobsen, J.T., Pasqual, G., Grabiner, B.C., Dominguez-
Sola, D., Sabatini, D.M., and Victora, G.D. (2017). Germinal Center Selection and Affinity
Maturation Require Dynamic Regulation of mTORC1 Kinase. Immunity *46*, 1045-
1058.e1046. 10.1016/j.immuni.2017.06.005.
Galazka, G., Mycko, M.P., Selmaj, I., Raine, C.S., and Selmaj, K.W. (2018). Multiple
sclerosis: Serum-derived exosomes express myelin proteins. Multiple sclerosis (Houndmills,
Basingstoke, England) *24*, 449-458. 10.1177/1352458517696597.
Groom, J., Kalled, S.L., Cutler, A.H., Olson, C., Woodcock, S.A., Schneider, P., Tschopp, J.,
Cachero, T.G., Batten, M., Wheway, J., et al. (2002). Association of BAFF/BLyS
overexpression and altered B cell differentiation with Sjögren's syndrome. The Journal of
clinical investigation *109*, 59-68. 10.1172/jci14121.
Hobeika, E., Thiemann, S., Storch, B., Jumaa, H., Nielsen, P.J., Pelanda, R., and Reth, M.
(2006). Testing gene function early in the B cell lineage in mb1-cre mice. Proceedings of the

National Academy of Sciences of the United States of America *103*, 13789-13794. 10.1073/pnas.0605944103.

Howell, O.W., Reeves, C.A., Nicholas, R., Carassiti, D., Radotra, B., Gentleman, S.M., Serafini, B., Aloisi, F., Roncaroli, F., Magliozzi, R., and Reynolds, R. (2011). Meningeal inflammation is widespread and linked to cortical pathology in multiple sclerosis. Brain : a journal of neurology *134*, 2755-2771. 10.1093/brain/awr182.

Jacobsen, J.T., Hu, W., TB, R.C., Solem, S., Galante, A., Lin, Z., Allon, S.J., Mesin, L., Bilate, A.M., Schiepers, A., et al. (2021). Expression of Foxp3 by T follicular helper cells in end-stage germinal centers. Science *373*. 10.1126/science.abe5146.

Jager, A., Dardalhon, V., Sobel, R.A., Bettelli, E., and Kuchroo, V.K. (2009). Th1, Th17, and Th9 effector cells induce experimental autoimmune encephalomyelitis with different pathological phenotypes. Journal of immunology *183*, 7169-7177. 10.4049/jimmunol.0901906.

Jordão, M.J.C., Sankowski, R., Brendecke, S.M., Sagar, Locatelli, G., Tai, Y.H., Tay, T.L., Schramm, E., Armbruster, S., Hagemeyer, N., et al. (2019). Single-cell profiling identifies myeloid cell subsets with distinct fates during neuroinflammation. Science *363*. 10.1126/science.aat7554.

Kawakami, N. (2016). In vivo imaging in autoimmune diseases in the central nervous system. Allergology international : official journal of the Japanese Society of Allergology *65*, 235-242. 10.1016/j.alit.2016.02.001.

Krishnamoorthy, G., Lassmann, H., Wekerle, H., and Holz, A. (2006). Spontaneous opticospinal encephalomyelitis in a double-transgenic mouse model of autoimmune T cell/B cell cooperation. The Journal of clinical investigation *116*, 2385-2392. 10.1172/jci28330.

Krishnamoorthy, G., Saxena, A., Mars, L.T., Domingues, H.S., Mentele, R., Ben-Nun, A., Lassmann, H., Dornmair, K., Kurschus, F.C., Liblau, R.S., and Wekerle, H. (2009). Myelin-specific T cells also recognize neuronal autoantigen in a transgenic mouse model of multiple sclerosis. Nature medicine *15*, 626-632. 10.1038/nm.1975.

Kuerten, S., Schickel, A., Kerkloh, C., Recks, M.S., Addicks, K., Ruddle, N.H., and Lehmann, P.V. (2012). Tertiary lymphoid organ development coincides with determinant spreading of the myelin-specific T cell response. Acta Neuropathol *124*, 861-873. 10.1007/s00401-012-1023-3.

Kyratsous, N.I., Bauer, I.J., Zhang, G., Pesic, M., Bartholomaus, I., Mues, M., Fang, P., Worner, M., Everts, S., Ellwart, J.W., et al. (2017). Visualizing context-dependent calcium signaling in encephalitogenic T cells in vivo by two-photon microscopy. Proceedings of the National Academy of Sciences of the United States of America *114*, E6381-e6389. 10.1073/pnas.1701806114.

Lanzavecchia, A. (1990). Receptor-mediated antigen uptake and its effect on antigen presentation to class II-restricted T lymphocytes. Annual review of immunology *8*, 773-793. 10.1146/annurev.iy.08.040190.004013.

Lee-Chang, C., Zéphir, H., Top, I., Dubucquoi, S., Trauet, J., Prin, L., and Vermersch, P. (2011). B-cell subsets up-regulate α4 integrin and accumulate in the cerebrospinal fluid in clinically isolated syndrome suggestive of multiple sclerosis onset. Neurosci Lett *487*, 273-277. 10.1016/j.neulet.2010.10.036.

Lehmann-Horn, K., Wang, S.Z., Sagan, S.A., Zamvil, S.S., and von Budingen, H.C. (2016). B cell repertoire expansion occurs in meningeal ectopic lymphoid tissue. JCI insight *1*, e87234. 10.1172/jci.insight.87234.

Lenert, P., Brummel, R., Field, E.H., and Ashman, R.F. (2005). TLR-9 activation of marginal zone B cells in lupus mice regulates immunity through increased IL-10 production. J Clin Immunol *25*, 29-40. 10.1007/s10875-005-0355-6.

Litzenburger, T., Fassler, R., Bauer, J., Lassmann, H., Linington, C., Wekerle, H., and Iglesias, A. (1998). B lymphocytes producing demyelinating autoantibodies: development and function in gene-targeted transgenic mice. The Journal of experimental medicine *188*, 169-180.

Lucchinetti, C.F., Popescu, B.F.G., Bunyan, R.F., Moll, N.M., Roemer, S.F., Lassmann, H., Brück, W., Parisi, J.E., Scheithauer, B.W., Giannini, C., et al. (2011). Inflammatory Cortical

Demyelination in Early Multiple Sclerosis. New England Journal of Medicine *365*, 2188-2197. 10.1056/NEJMoa1100648.

Mader, S., Kümpfel, T., and Meinl, E. (2020). Novel insights into pathophysiology and therapeutic possibilities reveal further differences between AQP4-IgG- and MOG-IgG-associated diseases. Curr Opin Neurol *33*, 362-371. 10.1097/wco.0000000000000813.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat Neurosci *13*, 133-140. 10.1038/nn.2467.

Magliozzi, R., Columba-Cabezas, S., Serafini, B., and Aloisi, F. (2004). Intracerebral expression of CXCL13 and BAFF is accompanied by formation of lymphoid follicle-like structures in the meninges of mice with relapsing experimental autoimmune encephalomyelitis. Journal of neuroimmunology *148*, 11-23. 10.1016/j.jneuroim.2003.10.056.

Magliozzi, R., Howell, O., Vora, A., Serafini, B., Nicholas, R., Puopolo, M., Reynolds, R., and Aloisi, F. (2007). Meningeal B-cell follicles in secondary progressive multiple sclerosis associate with early onset of disease and severe cortical pathology. Brain : a journal of neurology *130*, 1089-1104. 10.1093/brain/awm038.

Magliozzi, R., Howell, O.W., Reeves, C., Roncaroli, F., Nicholas, R., Serafini, B., Aloisi, F., and Reynolds, R. (2010). A Gradient of neuronal loss and meningeal inflammation in multiple sclerosis. Annals of neurology *68*, 477-493. 10.1002/ana.22230.

Mainero, C., Louapre, C., Govindarajan, S.T., Giannì, C., Nielsen, A.S., Cohen-Adad, J., Sloane, J., and Kinkel, R.P. (2015). A gradient in cortical pathology in multiple sclerosis by in vivo quantitative 7 T imaging. Brain : a journal of neurology *138*, 932-945. 10.1093/brain/awv011.

Mariño, E., Batten, M., Groom, J., Walters, S., Liuwantara, D., Mackay, F., and Grey, S.T. (2008). Marginal-zone B-cells of nonobese diabetic mice expand with diabetes onset, invade the pancreatic lymph nodes, and present autoantigen to diabetogenic T-cells. Diabetes *57*, 395-404. 10.2337/db07-0589.

Matsushita, T., Kobayashi, T., Mizumaki, K., Kano, M., Sawada, T., Tennichi, M., Okamura, A., Hamaguchi, Y., Iwakura, Y., Hasegawa, M., et al. (2018). BAFF inhibition attenuates fibrosis in scleroderma by modulating the regulatory and effector B cell balance. Sci Adv *4*, eaas9944. 10.1126/sciadv.aas9944.

Merico, D., Isserlin, R., Stueker, O., Emili, A., and Bader, G.D. (2010). Enrichment map: a network-based method for gene-set enrichment visualization and interpretation. PLoS One *5*, e13984. 10.1371/journal.pone.0013984.

Mitsdoerffer, M., Lee, Y., Jager, A., Kim, H.J., Korn, T., Kolls, J.K., Cantor, H., Bettelli, E., and Kuchroo, V.K. (2010). Proinflammatory T helper type 17 cells are effective B-cell helpers. Proceedings of the National Academy of Sciences of the United States of America *107*, 14292-14297. 10.1073/pnas.1009234107.

Mitsdoerffer, M., and Peters, A. (2016). Tertiary Lymphoid Organs in Central Nervous System Autoimmunity. Frontiers in immunology *7*, 451. 10.3389/fimmu.2016.00451.

Mitsdorffer, M., Di Liberto, G., Dotsch, S., Sie, C., Wagner, I., Pfaller, M., Kreutzfeldt, M., Frassle, S., Aly, L., Knier, B., et al. (2021). Formation and immunomodulatory function of meningeal B-cell aggregates in progressive CNS autoimmunity. Brain : a journal of neurology. 10.1093/brain/awab093.

Molnarfi, N., Schulze-Topphoff, U., Weber, M.S., Patarroyo, J.C., Prod'homme, T., Varrin-Doyer, M., Shetty, A., Linington, C., Slavin, A.J., Hidalgo, J., et al. (2013). MHC class II-dependent B cell APC function is required for induction of CNS autoimmunity independent of myelin-specific antibodies. The Journal of experimental medicine *210*, 2921-2937. 10.1084/jem.20130699.

Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat Genet *34*, 267-273. 10.1038/ng1180.

Mues, M., Bartholomaus, I., Thestrup, T., Griesbeck, O., Wekerle, H., Kawakami, N., and Krishnamoorthy, G. (2013). Real-time in vivo analysis of T cell activation in the central

nervous system using a genetically encoded calcium indicator. Nature medicine *19*, 778-783. 10.1038/nm.3180.

Parker Harp, C.R., Archambault, A.S., Sim, J., Ferris, S.T., Mikesell, R.J., Koni, P.A., Shimoda, M., Linington, C., Russell, J.H., and Wu, G.F. (2015). B cell antigen presentation is sufficient to drive neuroinflammation in an animal model of multiple sclerosis. Journal of immunology *194*, 5077-5084. 10.4049/jimmunol.1402236.

Pear, W.S., Nolan, G.P., Scott, M.L., and Baltimore, D. (1993). Production of high-titer helper-free retroviruses by transient transfection. Proceedings of the National Academy of Sciences of the United States of America *90*, 8392-8396. 10.1073/pnas.90.18.8392.

Perera, N.C., Wiesmüller, K.H., Larsen, M.T., Schacher, B., Eickholz, P., Borregaard, N., and Jenne, D.E. (2013). NSP4 is stored in azurophil granules and released by activated neutrophils as active endoprotease with restricted specificity. Journal of immunology *191*, 2700-2707. 10.4049/jimmunol.1301293.

Peters, A., Pitcher, L.A., Sullivan, J.M., Mitsdoerffer, M., Acton, S.E., Franz, B., Wucherpfennig, K., Turley, S., Carroll, M.C., Sobel, R.A., et al. (2011). Th17 cells induce ectopic lymphoid follicles in central nervous system tissue inflammation. Immunity *35*, 986-996. 10.1016/j.immuni.2011.10.015.

Pikor, N.B., Astarita, J.L., Summers-Deluca, L., Galicia, G., Qu, J., Ward, L.A., Armstrong, S., Dominguez, C.X., Malhotra, D., Heiden, B., et al. (2015). Integration of Th17- and Lymphotoxin-Derived Signals Initiates Meningeal-Resident Stromal Cell Remodeling to Propagate Neuroinflammation. Immunity *43*, 1160-1173. 10.1016/j.immuni.2015.11.010.

Raffaele, S., Lombardi, M., Verderio, C., and Fumagalli, M. (2020). TNF Production and Release from Microglia via Extracellular Vesicles: Impact on Brain Functions. Cells *9*. 10.3390/cells9102145.

Rauschmeier, R., Reinhardt, A., Gustafsson, C., Glaros, V., Artemov, A.V., Dunst, J., Taneja, R., Adameyko, I., Månsson, R., Busslinger, M., and Kreslavsky, T. (2022). Bhlhe40 function in activated B and TFH cells restrains the GC reaction and prevents lymphomagenesis. The Journal of experimental medicine *219*. 10.1084/jem.20211406.

Reali, C., Magliozzi, R., Roncaroli, F., Nicholas, R., Howell, O.W., and Reynolds, R. (2020). B cell rich meningeal inflammation associates with increased spinal cord pathology in multiple sclerosis. Brain Pathology *30*, 779-793. https://doi.org/10.1111/bpa.12841.

Salvador, F., Deramoudt, L., Leprêtre, F., Figeac, M., Guerrier, T., Boucher, J., Bas, M., Journiac, N., Peters, A., Mars, L.T., and Zéphir, H. (2022). A Spontaneous Model of Experimental Autoimmune Encephalomyelitis Provides Evidence of MOG-Specific B Cell Recruitment and Clonal Expansion. Frontiers in immunology *13*, 755900. 10.3389/fimmu.2022.755900.

Schriek, P., Ching, A.C., Moily, N.S., Moffat, J., Beattie, L., Steiner, T.M., Hosking, L.M., Thurman, J.M., Holers, V.M., Ishido, S., et al. (2022). Marginal zone B cells acquire dendritic cell functions by trogocytosis. Science *375*, eabf7470. 10.1126/science.abf7470.

Segundo, C., Rodríguez, C., García-Poley, A., Aguilar, M., Gavilán, I., Bellas, C., and Brieva, J.A. (2001). Thyroid-infiltrating B lymphocytes in Graves' disease are related to marginal zone and memory B cell compartments. Thyroid *11*, 525-530. 10.1089/105072501750302813.

Serafini, B., Rosicarelli, B., Franciotta, D., Magliozzi, R., Reynolds, R., Cinque, P., Andreoni, L., Trivedi, P., Salvetti, M., Faggioni, A., and Aloisi, F. (2007). Dysregulated Epstein-Barr virus infection in the multiple sclerosis brain. The Journal of experimental medicine *204*, 2899-2912. 10.1084/jem.20071030.

Serafini, B., Rosicarelli, B., Magliozzi, R., Stigliano, E., and Aloisi, F. (2004). Detection of ectopic B-cell follicles with germinal centers in the meninges of patients with secondary progressive multiple sclerosis. Brain pathology (Zurich, Switzerland) *14*, 164-174.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res *13*, 2498-2504. 10.1101/gr.1239303.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. Cell *177*, 1888-1902.e1821. 10.1016/j.cell.2019.05.031.
Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. Proceedings of the National Academy of Sciences of the United States of America
102, 15545-15550. 10.1073/pnas.0506580102.
Tom, R., Bisson, L., and Durocher, Y. (2008). Culture of HEK293-EBNA1 Cells for
Production of Recombinant Proteins. CSH Protoc *2008*, pdb.prot4976.
10.1101/pdb.prot4976.
Tzartos, J.S., Craner, M.J., Friese, M.A., Jakobsen, K.B., Newcombe, J., Esiri, M.M., and
Fugger, L. (2011). IL-21 and IL-21 receptor expression in lymphocytes and neurons in
multiple sclerosis brain. Am J Pathol *178*, 794-802. 10.1016/j.ajpath.2010.10.043.
Victora, G.D., Schwickert, T.A., Fooksman, D.R., Kamphorst, A.O., Meyer-Hermann, M.,
Dustin, M.L., and Nussenzweig, M.C. (2010). Germinal center dynamics revealed by
multiphoton microscopy with a photoactivatable fluorescent reporter. Cell *143*, 592-605.
10.1016/j.cell.2010.10.032.
Weber, M.S., Prod'homme, T., Patarroyo, J.C., Molnarfi, N., Karnezis, T., Lehmann-Horn, K.,
Danilenko, D.M., Eastham-Anderson, J., Slavin, A.J., Linington, C., et al. (2010). B-cell
activation influences T-cell polarization and outcome of anti-CD20 B-cell depletion in central
nervous system autoimmunity. Annals of neurology *68*, 369-383. 10.1002/ana.22081.
Yang, S.Y., Long, J., Huang, M.X., Luo, P.Y., Bian, Z.H., Xu, Y.F., Wang, C.B., Yang, S.H.,
Li, L., Selmi, C., et al. (2021). Characterization of Organ-Specific Regulatory B Cells Using
Single-Cell RNA Sequencing. Frontiers in immunology *12*, 711980.
10.3389/fimmu.2021.711980.
Zhang, X. (2013). Regulatory functions of innate-like B cells. Cell Mol Immunol *10*, 113-121.
10.1038/cmi.2012.63.

1247

1248

Figure legends

Figure 1. Th17 cells induce formation of large ectopic lymphoid follicle-like structures expressing germinal center markers in association with the meninges of spinal cord and brain. (A-C) At the peak of disease, cells were isolated from the CNS of Th1 and Th17-EAE mice and analyzed for the presence of CD19$^+$ B cells. (**A**) Representative flow cytometry plots (pre-gated on CD45^{high} CD11b^{low} cells) and quantification of the (**B**) frequency of CD19$^+$ B cells in CD45^{high} CD11b^{low} cells and (**C**) absolute numbers of infiltrating CD19$^+$ B cells in the CNS. Graphs show cumulative data from three (Th1-EAE) or eight (Th17-EAE) independent experiments. Mann-Whitney U test. *p < 0.05; **p < 0.01. Horizontal lines indicate means. Dots represent individual mice. (**D-G**) At the peak of disease, (**D, E**) longitudinal spinal cord and (**F, G**) coronal brain cryosections of Th17-EAE mice were prepared and stained for T cells (CD3, red), B cells (B220, green) and laminin (blue). (**E**) Zoom of red highlighted area in D. (**F**) Subarachnoid space at the base of the brain. (**G**) Cortex. Scale bars: 200 µm. (**H-K**) CNS cryosections of Th17-EAE mice at the peak of disease were stained for (**H**) T cells (CD3, red), B cells (B220, green) and the proliferation marker Ki67 (blue); (**I**) B cells (B220, magenta) and the germinal center marker peanut agglutinin (PNA, green); (**J, K**) T cells (CD3, red), B cells (B220, green) and the enzyme activation-induced cytidine deaminase (AID, magenta). For better visualization, the AID staining is shown separately in K.

Figure 2. CNS-infiltrating B cells are phenotypically different from peripheral B cells. At the peak of disease, B cells were sorted from indicated organs of Th17 recipients (n = 3 mice) based on their expression of CD19, and their gene expression was characterized by single-cell RNA sequencing. (**A**) Single cell transcriptomes are depicted in UMAP plots representing ten color-coded cell clusters, shown separately for CNS, cervical lymph node (cLN) and spleen. (**B**) For each cluster, the fraction of cells belonging to CNS, cLN or spleen was calculated. (**C**-**E**) Pathway network analysis of upregulated pathways in (**C**) cluster 6, (**D**) cluster 7 and (**E**) cluster 9. (**F, G**) Violin plots showing expression level of genes associated with (**F**) light zone (LZ) and (**G**) dark zone (DZ) germinal center B cells.

Figure 3. CNS-infiltrating B cells express markers associated with B1a / marginal zone (MZ) B cells and undergo clonal expansion in the CNS. Violin plots showing expression levels of gene sets associated with (**A**) B1a B cells and MZ B cells. (**B**) Cells were isolated from the CNS of Th17 recipients and analyzed for expression of CD21 and CD1d by flow cytometry. A pre-gate on CD19$^+$ IgM^{high} IgD^{low} B cells was applied. (Left panel) One representative plot of two independent experiments (n = 6 mice). (Right panel) One plot from a mouse with substantial CD21^{high} CD1d$^+$ population (n = 1 mouse). (**C**) Spinal cord cryosections of Th17 recipients were stained for T cells (CD3, blue), B cells (B220, red) and CD1d (green). (**D**) At the peak of disease, B cells were sorted from indicated organs of Th17 recipients (n = 3 mice) based on their expression of CD19, and characterized by BCR repertoire analysis. Feature plots depicting distribution of nonexpanded (grey) and expanded (colored) B cells among B cell clusters, plotted separately for CNS, cLN and spleen. The color code indicates the number of cells per clone. (**E**) Fraction of expanded cells belonging to CNS, spleen and cLN. (**F**) Percentage of clones with expansion factors ≥3 for each cluster in the CNS. (**G**) Violin plots showing expression levels of genes associated with plasmablasts (PB).

Figure 4. CNS-infiltrating B cells show elevated expression of genes associated with cytokine signaling, B cell activation and antigen presentation/immune synapse. (**A-E**) Violin plots showing expression levels of gene signatures induced by (**A**) IFNγ, (**B**) IL-21 and (**C**) TNFα, (**D**) gene sets associated with B cell activation and (**E**) antigen presentation/immune synapse. (**F-G**) At the peak of disease, cells were isolated from CNS or spleen and B cells were analyzed for expression of (**F**) CD69, CD83, CD86 and (**G**) MHC-II, CD40, ICAM-1, and IL-21R (pre-gated on CD19$^+$ cells) by flow cytometry. Graphs show the frequency of (**F**) CD69$^+$, CD83$^+$ or CD86$^+$ cells and (**G**) the geometric mean fluorescence intensity (gMFI) of MHC-II, CD40, ICAM-1 or IL-21R. Paired t-test. *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001. Dots represent individual mice.

Figure 5. 2D2 T cells form long-lasting contacts with B cells compared to OT-II T cells. Intravital microscopy of Twitch-2B$^+$ T cells and tdTomato$^+$ B cells in spinal cord eLFs of Th17-EAE mice. (**A**) Experimental outline. MOG-specific T cells from 2D2 mice are retrovirally transduced with the Twitch-2B construct during their differentiation into Th17 cells. Upon adoptive transfer in Mb1.Cre x ROSA26 tdTomato mice, intravital two-photon microscopy of spinal cord eLFs in EAE mice can be performed to visualize interactions of Th17 cells (green) and B cells (red), as well as subsequent activation events of Th17 cells. (**B**) Dorsal view onto eLF containing Twitch-2B$^+$ T cells (green) and tdTomato$^+$ B cells (red). Scale bar: 100μm. (**C**) Exemplary images of T:B cell contacts. (**D**) Proportion of 2D2 (n = 232 cells from six independent experiments) and OT-II T cells (n = 107 cells from four independent experiments) forming contact with B cells. (**E**) Quantification of T:B cell contact duration. For both experimental groups, videos of 15-50 min length were acquired. Dots represent individual contacts. Horizontal lines indicate means. Mann-Whitney U test. ****p < 0.0001. (**F**) Proportion of long (> 5 min) and short (< 5 min) T:B cell contacts. (**G**) Quantification of the velocity of 2D2 T cells without B cell contact (n = 10938 velocities), 2D2 T cells with B cell contact (n = 8314 velocities) and OT-II T cells (n = 9413 velocities). One-way ANOVA with Tukey post-test. ****p < 0.0001. Horizontal lines indicate means. Graph shows cumulative data from (2D2) six or (OT-II) four independent experiments.

Figure 6. 2D2 T cells, but not OT-II T cells, become reactivated during B cell contact. Intravital microscopy of Twitch-2B$^+$ T cells and tdTomato$^+$ B cells in spinal cord eLFs of Th17-EAE mice. (**A**) (Left) Fluorescence overlay of Twitch-2B$^+$ T cells (green) and tdTomato$^+$ B cells (red) and (right) pseudocolor calcium ratio image showing calcium levels of T cells. (**B**) Quantification of the calcium indicator ratio (YFP/CFP) of 2D2 T cells without B cell contact (n = 4988 ratios), 2D2 T cells with B cell contact (n = 3653 ratios) and OT-II T cells (n = 9531 ratios). One-way ANOVA with Tukey post-test. ****p < 0.0001. The black dashed line indicates the ratio threshold (0.534). Horizontal lines indicate means. The graph shows cumulative data from (2D2) three or (OT-II) four independent experiments. (**C**) Proportion of 2D2 and OT-II T cells exhibiting calcium signaling. (**D**) Duration of calcium signaling (min) of 2D2 and OT-II T cells. The dotted line at 2 min indicates the cut-off between short-term and long-term calcium signaling. (**E**) Proportion of T cells with short-term (< 2 min), long-term (> 2 min) or no calcium signaling. (**C-E**) Cumulative data of 2D2 (n = 104 cells from three independent experiments) and OT-II T cells (n = 107 cells from four independent experiments). (**F, G**) Representative tracks of (**F**) 2D2 T cells (n = 104 cells from three independent experiments) and (**G**) OT-II T cells (n = 107 cells from four independent experiments) showing calcium indicator ratio (YFP/CFP, red) and velocity (black) in each

1343 time frame. The red dashed line indicates the ratio threshold (0.534). Grey shading highlights
1344 time frames with B cell contact.
1345

Figure 7. T cells depend on B cells to maintain a highly pro-inflammatory cytokine
profile in the CNS. Comparison of T cell responses in Th17-mediated EAE in B cell-deficient
Mb1-KO mice and WT mice. At the peak of disease, cells were isolated from (**A**) CNS or (**B**)
spleen and T cells were analyzed for IL-17A and IFNγ expression by intracellular cytokine
staining. Transferred T cells were identified by their expression of the transgenic MOG-
specific Vα3.2 TCR. Representative flow cytometry plots and quantification of IL-17A$^+$, IFNγ$^+$
and IL-17A$^+$IFNγ$^+$ producers of (upper panel) transferred (Vα3.2$^+$) and (lower panel)
endogenous (Vα3.2$^-$) T cells. Graphs show cumulative data from two independent
experiments. (**A**, upper panel) unpaired *t*-test (left), Welch`s t-test (middle), Mann-Whitney U
test (right). (**A**, lower panel) Mann-Whitney U test (left and right), Welch`s t-test (middle). (**B**,
upper panel) unpaired *t*-test (left), Mann-Whitney U test (middle), Welch`s t-test (right). (**B**,
lower panel) unpaired *t*-test (left and right), Welch`s t-test (middle). *$p < 0.05$; **$p < 0.01$; ***p
< 0.001. Mean ± SEM. Dots represent individual mice.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

T cells
B cells

Low Ca²⁺ High

B

ratio [YFP/CFP]

**** 47.2 % 75.0 % 5.0 %

without B cell contact
with B cell contact
OT-II T cells

2D2 T cells

C

Proportion of T cells (%)

with calcium spikes
without calcium spikes

2D2 T cells
OT-II T cells

E

Proportion of T cells (%)

2D2 T cells
OT-II T cells

Long-term   Short-term   no signaling

D

Duration of calcium signaling [min]

2D2 T cells
OT-II T cells

F

2D2 T cells

ratio [YFP/CFP] — velocity [µm/min]
time (min)

G

OT-II T cells

ratio [YFP/CFP] — velocity [µm/min]
time (min)

Figure 6

Figure 7