# T:B cell cooperation in ectopic lymphoid follicles propagates CNS autoimmunity

Anna Kolz[1,2], Isabel Julia Bauer[1,2], Clara de la Rosa[1,2], Vladyslav Kavaka[1,2], Anna Sophie Thomann[1,2], Rosa Schmitz[1,2], Martin Kerschensteiner[1,2,3], Eduardo Beltran[1,2,3], Naoto Kawakami[1,2], Anneli Peters[1,2]

[1] Institute of Clinical Neuroimmunology, LMU University Hospital, LMU Munich, Germany
[2] Biomedical Center (BMC), Faculty of Medicine, LMU Munich, Germany
[3] Munich Cluster for Systems Neurology (SyNergy), Munich, Germany

Correspondence to: Anneli.Peters@med.uni-muenchen.de

## Abstract

Meningeal ectopic lymphoid follicle-like structures (eLFs) have been described in multiple sclerosis (MS) and its animal model experimental autoimmune encephalomyelitis (EAE), but their role in CNS autoimmunity is unclear. To analyze the cellular phenotypes and interactions within these structures, we employed a Th17 adoptive transfer EAE model featuring formation of large, numerous eLFs. Single-cell transcriptomic analysis revealed that clusters of activated B cells and B1/Marginal Zone-like B cells are overrepresented in the CNS and identified B cells poised for undergoing antigen-driven germinal center (GC) reactions and clonal expansion in the CNS. Furthermore, CNS B cells showed enhanced capacity for antigen presentation and immunological synapse formation compared to peripheral B cells. To directly visualize Th17:B cell cooperation in eLFs, we labeled Th17 cells with a ratiometric calcium sensor, and tracked their interactions with tdTomato-labeled B cells in real-time. Thereby, we demonstrated for the first time that T and B cells form long-lasting antigen-specific contacts in meningeal eLFs that result in reactivation of autoreactive T cells. Consistent with these findings, autoreactive T cells depended on CNS B cells to maintain a pro-inflammatory cytokine profile in the CNS. Collectively, our study reveals that extensive T:B cell cooperation occurs in meningeal eLFs in our model promoting differentiation and clonal expansion of B cells, as well as reactivation of CNS T cells and thereby supporting smoldering inflammatory processes within the CNS compartment. Our results provide valuable insights into the function of eLFs and may provide a direction for future research in MS.

## Introduction

In multiple sclerosis (MS) autoreactive T cells recognizing central nervous system (CNS)-specific antigens are thought to be important drivers of disease. To our current knowledge, they are primed in the periphery before they invade the CNS, where they presumably become reactivated by contact with the autoantigen, and subsequently orchestrate the inflammatory cascade. Both Th1 and Th17 cells contribute to pathogenesis in MS and its animal model experimental autoimmune encephalomyelitis (EAE), and during the past two decades, compelling evidence has also suggested a pivotal role of B cells (Comi et al.,

2021). However, it is not clear how T cells and B cells cooperate to induce and propagate
disease. In MS, ectopic lymphoid follicle-like structures (eLFs) were found in the
leptomeninges of patients with progressive disease course (Magliozzi et al., 2007; Serafini et
al., 2004), but also in earlier stages of relapsing-remitting MS (Lucchinetti et al., 2011).
Initially, they were thought to be restricted to brain meninges, where they are localized mainly
in sulci, but a recent study showed that they can also form in spinal cord meninges (Reali et
al., 2020). These aggregates show varying degrees of maturation, ranging from loose
clusters of cells to highly organized follicle structures, which contain plasma cells,
proliferating B cells, T cells and follicular dendritic cells (FDCs), and therefore are suggestive
of germinal center formation (Mitsdoerffer and Peters, 2016). Clinical data show a correlation
between the presence of eLFs and a more severe disease course – younger age at disease
onset, irreversible disability, and death, as well as a more pronounced cortical pathology in
the tissue underneath eLFs (Howell et al., 2011; Magliozzi *et al.*, 2007; Magliozzi et al., 2010;
Serafini et al., 2007; Serafini *et al.*, 2004). Therefore, these follicles have been postulated to
maintain or exacerbate chronic inflammation directly in the CNS. Pathogenic functions of
eLFs could include both initial priming and maturation of autoreactive B and T cell clones, as
well as reactivation and expansion of pre-existing autoreactive T and B cells recruited from
the periphery. In addition, eLFs may be a source of pathogenic soluble factors such as auto-
antibodies and pro-inflammatory cytokines that affect the underlying regions. However, eLFs
might also harbor regulatory processes aimed at dampening excessive inflammation and it is
possible that both eLFs with disease promoting and eLFs with regulatory function exist in MS
either in different subsets of patients or in different phases of the disease. Since data from
human eLFs is based on and largely restricted to histological analysis of post-mortem CNS
material, our knowledge of cellular processes within eLFs remains limited.

To study eLFs in more detail, here we employed a Th17 adoptive transfer EAE model, which
features formation of eLFs within the meninges (Jager et al., 2009; Peters et al., 2011). While
meningeal eLFs were also identified in other EAE models differing regarding genetic
background, mode of EAE induction and auto-antigen (Dang et al., 2015; Kuerten et al.,
2012; Magliozzi et al., 2004; Pikor et al., 2015), we chose the Th17 adoptive transfer EAE
model for our study as it exhibits robust formation of the largest and most numerous eLFs.
Due to the relatively short and severe disease course, the Th17 EAE model is not suitable to
study the clinical impact of eLFs, however, it can provide valuable insights into the early
stages of eLF formation. Specifically, the large number and size of the eLFs and the
substantial recruitment of B cells facilitate in depth phenotyping via single-cell RNA
sequencing and even enable us to study cellular interactions via live imaging.

The aim of this study was to investigate the unique relationship of Th17 cells and B cells in
EAE development, and, in particular, to assess the nature of their interaction in eLFs.
Combining histological and flow cytometric analyses with single-cell transcriptomics and
intravital imaging, we show that meningeal eLFs are sites of extensive T:B cell cooperation
resulting in both development of germinal center B cells and reactivation of autoreactive T
cells and can thereby perpetuate autoimmune reactions in the CNS compartment.

**Results**
**Th17 cells induce formation of ectopic lymphoid follicles in CNS**
**meninges**

To be able to study cellular processes in eLFs, we first established adoptive transfer EAE
according to Jäger et al. (Jager *et al.*, 2009), a model in which disease is induced by transfer
of *in vitro* differentiated pure MOG-specific Th1 or Th17 cells into immunocompetent recipient
mice (Figure 1A, S1A). As previously shown, Th1 and Th17 cells induced EAE independently
of each other with comparable clinical disease courses and similar severity (Figure S1B). In
the inflamed CNS, transferred Th1 and Th17 cells maintained stable expression of their
signature cytokine IFNγ and IL-17A, respectively, and Th17 cells can additionally adopt IFNγ
production (Figure S1C, D). To assess B cell recruitment to the CNS, we compared B cell
frequencies among CNS-infiltrating cells between Th1 and Th17 recipient mice at late peak
of disease. We detected a significant increase in both frequency and absolute numbers of B
cells in the CNS of Th17 recipients compared to Th1 recipients (Figure 1B-D). We next
performed a detailed immunohistochemical analysis of CNS lesions to investigate the
localization, structural organization, and composition of cellular infiltrates in Th17-EAE mice.
Following identification of regions with lymphocytic infiltrates by Giemsa staining (Figure
S2A), neighboring sections were stained for T cells, B cells and laminin, which visualizes the
basal lamina of blood vessels and the pia mater. In line with previous observations in Th17
recipient mice (Peters *et al.*, 2011; Pikor *et al.*, 2015), we found numerous eLFs within the
meninges of the spinal cord. eLFs consisted of accumulated T and B cells and showed
varying degrees of organization and size: while some contained loose aggregates of a few
cells, others were composed of large amounts of T and B cells and extended over wide
distances of up to several millimeters (Figure 1E, S2B). Based on their unique morphology
and expression of CD11b, we also identified macrophages in close vicinity, but also within
the follicle structures (Figure S2D). Large eLFs were also found in many different areas of
the brain, including for example the subarachnoid space at the base of the brain (Figure
S2C) and the cortex (Figure 1F). Similarly, we did not observe a preferential location for eLFs
within spinal meninges but rather a widespread distribution along the cord (Figure 1G). eLFs
appeared to originate from the meninges and invade the underlying parenchyma, and often
exhibited separate T and B cell zones in our model, thereby mimicking the structural
organization of SLOs. Notably, all eLFs were characterized by prominent B cell clustering,
with B cells being almost exclusively confined to these meningeal aggregates. In contrast, T
cells were more widely distributed and detected both inside eLFs in large numbers, as well
as in the underlying subpial parenchymal tissues (Figure 1E, F, S2B, C). Similar to eLFs in
human CNS tissue (Magliozzi *et al.*, 2007), staining for Ki67 revealed eLFs in our model to
be sites of T and B cell proliferation (Figure 1H). To evaluate the degree of eLF maturation,
we assessed expression of GC markers on B and T cells. In many eLFs, only few scattered
B cells expressed the GC markers PNA or AID (Figure 1I, J), however, some eLFs exhibited
clusters of AID$^+$ B cells (Figure 1K), which have also been detected in human eLFs (Serafini
*et al.*, 2007). In addition, PNA$^+$ B cells (Figure S3A), isotype-switched IgM$^-$ IgD$^-$ B cells
(Figure S3B), and a small population of plasma cells (Figure S3C) were detected among
CNS-infiltrating cells via flow cytometry. Furthermore, a fraction of CNS-infiltrating T cells
expressed the GC and activation marker GL7 with transferred 2D2 Th17 cells showing higher
frequency of GL7$^+$ cells than endogenous T cells from the recipient (Figure S3D).

Taken together, we confirmed that the Th17 adoptive transfer EAE model prominently
features formation of eLFs reminiscent of eLFs in MS patients in both spinal cord and brain
meninges. Considering the high frequency of recruited B cells, the high abundance and large
size of eLFs, as well as our initial evidence that at least some of the eLFs are mature enough
to harbor GC B and T cells, our model is well suited to study the cellular processes in CNS
eLFs in more detail.

## CNS-infiltrating B cells are highly activated and poised for undergoing germinal center reactions

To obtain a more refined understanding of the properties and effector functions of B cells in meningeal eLFs, we characterized the transcriptome of CNS-infiltrating B cells from Th17 recipients by single-cell RNA sequencing, and compared it to the transcriptome of peripheral B cells from spleen and cervical lymph nodes (cLN) of the same animal. Unbiased clustering of the single-cell transcriptomes revealed ten individual B cell clusters varying substantially in size and containing B cells of all three compartments. Interestingly, CNS B cells showed a distinct distribution of B cells across the different clusters, indicating them to be phenotypically different from peripheral B cells (Figure 2A, Figure S4A+B). Among the identified clusters, cluster 6-9 were highly enriched for CNS B cells, most pronounced for cluster 6, in which 93.5 % of cells belonged to the CNS fraction (Figure 2B). Gene set enrichment analysis against the GO biological processes database revealed that cluster 6 B cells are metabolically highly active and have a high energy demand as they showed upregulation of pathways associated with respiration, ATP biosynthesis, glycolysis, fatty acid metabolism and oxidative phosphorylation (Figure S4C, S5A). In line with this, they showed highly increased expression of many ribosomal proteins (Figure S4A and Table S1), which have been shown to be upregulated in early GC B cells in response to mTOR signaling in order to prepare for the increased translational demand of an imminent GC reaction (Ersching et al., 2017). Notably, both mTOR and Myc signaling, which precede entry into a GC reaction, were high in cluster 9, 6 and 7 B cells, and always higher in CNS B cells compared to peripheral B cells within each cluster (Fig. S4C). Pathway analysis showed that cluster 7 B cells are highly activated / differentiated and transcriptionally very active (Fig. S5B). Thus, *Nr4a1*, *Egr1*, *Egr2* and *Egr3*, which are promptly induced upon B cell activation through the BCR, are among the top 10 up-regulated genes in cluster 7 (Fig. S4A and Table S1). Cluster 9 cells showed upregulation of pathways associated with cell cycle, mitosis, DNA replication, and proliferation and accordingly *Mki67* was among the top 10 upregulated genes in this cluster (Fig. S5C, Fig. S4A and Table S1). Importantly, the CNS-enriched clusters 6 and 7 showed highly significantly increased expression of genes associated with a light zone (LZ) germinal center B cell signature (Victora et al., 2010), compared to other CNS B cells as well as compared to peripheral B cells within the same cluster (Figure 2C). Thus, for example the LZ GC markers *Cd83* and *Fas*, as well as the cyclin *Ccnd2* and the transcription factor *Bhlhe40*, which have been associated with LZ GCs (Rauschmeier et al., 2022; Victora *et al.*, 2010), were among the top 50 upregulated genes in both cluster 6 and 7 (Table S1). These data indicate that the phenotype of B cells especially in cluster 6 and 7, which are highly overrepresented in the CNS, is compatible with LZ GC B cells, suggesting that a considerable fraction (about 19.5 %) of CNS B cells is poised to undergo GC reactions. In contrast, only cluster 9 B cells showed highly elevated expression of genes associated with dark zone (DZ) germinal center B cells compared to other CNS B cells (Victora *et al.*, 2010) (Figure 2D), including for example *Stmn1, Cenpa,* and the cyclin *Ccnb2*. This is in line with the increased proliferative capacity of these cells indicated by the pathway analysis (Fig. S5C) and the high level of oxidative phosphorylation (Fig. S4C), suggesting that with cluster 9 only a very small fraction (1.4 %) of B cells in eLFs has progressed to DZ GC B cells to begin clonal expansion.

## B1a / MZ-like B cells infiltrate the CNS

In addition to the clusters described above, also cluster 8 was highly enriched for CNS B cells. Further examination of the upregulated genes in this cluster, such as *Mzb1*, *Fcrl5* and *Cd9* suggested similarities to marginal zone (MZ) and B1 cells (Fig. S4A, Table S1). This observation was validated by comparing the cluster 8 transcriptome with publicly available gene signatures of B1a cells from spleen and peritoneum and splenic MZ B cells confirming that cluster 8 represents a population of B1a / MZ-like B cells in the CNS (Figure 2E, F). The only other cluster that showed highly elevated expression of B1a and MZ markers including *Cd9* and *Fcrl5* was cluster 1, which consists mainly of peripheral B cells and contains the MZ B cells in the spleen (Fig. 2G, H, Fig. S4A, Table S1). As regular MZ B cells are often identified via expression of CD1d and CD21, we analyzed CNS-infiltrating cells for the presence of $IgM^{high}$ $IgD^{low}$ $CD21^{high}$ $CD1d^+$ B cells via flow cytometry, and detected a small and relatively dim population (Figure S5D, E). Additionally, we confirmed presence of scattered $CD1d^+$ B cells in eLFs in the CNS via confocal microscopy (Figure S5F). Considering that CD1d is not necessarily expressed on B1a cells, this gating and staining strategy may not have included all the cells in cluster 8.

## CNS-infiltrating B cells undergo clonal expansion

Analysis of the BCR repertoire via comparison of the sequence of Ig variable regions revealed the existence of expanded B cell clones, and interestingly, the clones with the highest expansion factors predominantly belonged to cluster 8, especially in the CNS (Figure 3A, S5G, H). Importantly, 55.5 % of expanded cells represented B cells from the CNS, while 28.8 % and 15.5 % were derived from spleen and cLN, respectively (Figure 3B). This preferential localization of expanded clones may indicate that infiltrating B cells undergo clonal expansion in the inflamed CNS. The clones with the highest expansion factors in the CNS were predominantly localized in cluster 8, but we also found clones with lower expansion factors in the CNS in cluster 9 and 4 (Figure 3C). Since B cells were sorted based on their expression of CD19, plasma cells, which no longer express CD19, were not included in this analysis. Therefore, the expanded CNS B cells could be in the process of differentiating into plasmablasts. Indeed, only cluster 8 and cluster 9 B cells showed significantly increased expression of plasmablast-signature genes compared to other CNS B cells (Figure 3D). Considering that cluster 9 B cells in the CNS displayed features of DZ GC B cells (Fig. 2D), these data suggest that cluster 9 B cells are starting to undergo clonal expansion with still low expansion factors and are also upregulating the plasmablast differentiation program at the beginning of an antigen-driven GC reaction. In contrast, cluster 8 B cells have much higher expansion factors but show lower expression of GC markers, indicating that they may differentiate into plasmablasts independent of classical T cell-driven GC reactions. This is in line with the known properties of regular B1 and MZ B cells to be able to operate more independently of T cell help and give rise to extrafollicular plasmablasts (Cyster and Allen, 2019). A comparison between the transcriptome of non-expanded and expanded cells in cluster 8 revealed that the expanded cells upregulated genes associated with plasma cell differentiation and MZ B cells or B1 B cells, including *Fam46c*, *Emb* and *Gpx4* (Figure S5J). Additionally, cluster 8 B cells showed upregulation of pathways associated with regulation of immune responses (Figure S5I) and expressed for example *Ctla4* and *Apoe* among the top 50 upregulated genes (Fig. S4A, Table S1).

As we observed that the CNS in Th17 recipient mice supports clonal expansion of B cells
and harbors B cells poised to undergo GC reactions, we next investigated whether
development and expansion of MOG-specific B cells from the polyclonal repertoire occurs in
our model. We first determined the presence of a MOG-specific IgG1 response in the serum
of Th17 and Th1 recipients via ELISA. At late peak of clinical disease, many Th17 recipients
showed high levels of MOG-specific IgG1 antibodies compared to very low levels before EAE
onset. Some Th1 recipients also exhibited elevated antibody titers at the peak of disease
compared to very low levels in mice, which were analyzed at an early disease stage before B
cells are starting to infiltrate into the CNS. However, on average MOG-specific IgG1 titers in
Th17 recipients were about 3-fold higher than in Th1 recipients (Figure S6A). To determine
whether CNS eLFs contain MOG-specific B cells, we optimized detection of MOG-specific B
cells via IHC by staining CNS cryosections with a fluorescently labeled tetramerized MOG-
protein (MOG$_{tet}$), which was validated on LN cryosections from IgH$^{MOG}$ mice carrying MOG-
specific B cells (Figure S6B). Staining of CNS cryosections from Th17-EAE mice revealed
that most eLFs contained only very few MOG$_{tet}^+$ B cells (Figure S6C, left panel). However,
occasionally some eLFs displayed a massive accumulation of MOG$_{tet}^+$ B cells (Figure S6C,
right panel). Together, these data indicate that transferred MOG-specific Th17 cells recruit B
cells with the same antigen specificity from the endogenous pool and that maturation and/or
expansion of MOG-specific B cells may at least partially happen in meningeal eLFs.

## CNS-infiltrating B cells respond to T cell cytokines

Our transcriptomic analyses suggest that many of our CNS B cells in Th17 adoptive transfer
EAE are highly activated, poised for entering GC reactions and undergoing clonal expansion
(Fig. 2 and 3). In addition to cellular interactions, phenotype and function can also be shaped
by cytokines. Thus, we evaluated whether we could find evidence of cytokine-induced
signaling in CNS B cells. B cells are known to strongly respond to cues from both type I and
type II interferons and thus we compared gene expression in our clusters to publicly available
gene signatures induced by IFNγ and IFNα. Although, some IFNγ is produced by transferred
Th17 cells as well as by endogenous T cells in our model, among the CNS-specific clusters
only cluster 6 and 7 showed evidence of IFNγ-driven signaling compared to the other CNS B
cells, and this was enhanced in CNS B cells compared to peripheral B cells within the same
cluster (Figure 4A). In addition, cluster 5 B cells, where only 32 % of cells are derived from
CNS, express IFNγ-driven genes including *Stat1* and *Irf7* (Fig. S4A and Table S1). In
contrast, the CNS-specific clusters did not show an enhanced response to IFNα neither
compared to other CNS B cells nor compared to peripheral B cells within the same cluster
(Figure S7A), suggesting that IFNα is not a relevant component of the CNS-specific cytokine
milieu in our model. Among the CNS-enriched B cell clusters, cluster 7 and 9 exhibited
elevated expression of genes induced by IL-21, and for all CNS-specific clusters expression
was always higher in CNS B cells compared to peripheral B cells from the same cluster (Fig.
4B). Since IL-21 is produced by Th17 cells and is also a T helper cell-derived cytokine
important for B cell responses in GC reactions, this speaks for T cells providing help to B
cells to initiate GC reactions in the CNS. Indeed, expression levels of IL-21 induced genes
were highest in cluster 7 and 9 (Figure 4B), which were shown to express markers of LZ and
DZ GC B cells, respectively (Figure 2C, D). In line with the increased response to IL-21, flow
cytometric analysis showed that CNS B cells also expressed higher levels of IL-21R than
peripheral B cells (Figure 4C + S7C). CNS B cells also strongly responded to TNFα, while
peripheral B cells did not (Figure 4D). This suggests that TNFα, which is produced by T cells

but can also be released by myeloid cells and B cells themselves, is an important component of the CNS-specific cytokine milieu in our model. The highest levels of TNFα-induced gene expression were detected in cluster 7 B cells, which also showed very high levels of *Nr4a1* expression indicating that TNFα may even be a driver of B cell activation in CNS B cells. In line with this, we confirmed via flow cytometry that the frequency of activated CD69+ and CD83+ B cells was strongly increased in CNS B cells compared to splenic B cells (Figure 4E, F, S7B).

## CNS B cells show enhanced APC function

Since we detected subpopulations of activated B cells that show features of GC B cells and respond to T cell cytokines, it is likely that eLF B cells closely interact with T cells in our model. To determine, whether CNS B cells not only respond to T cell help but also act on the T cells we determined the expression level of genes associated with APC function and immunological synapse (IS) formation in the CNS-specific clusters (Figure 4G). Notably, cluster 6, 7, and 9, which also showed the highest expression of the LZ and DZ GC B cell-associated genes, displayed highly significantly upregulated expression of APC/IS genes compared to other CNS B cells (Figure 4G). In line with this, CNS B cells showed highly elevated protein expression levels of MHC-II (Figure 4H, S7C), as well as elevated expression levels of the costimulatory molecule CD40 (Figure 4I, S7C), and increased frequencies of the costimulatory molecule CD86 (Figure 4J, S7B), which together with MHC-II form the central supramolecular activation complex (SMAC) of the immunological synapse. Furthermore, CNS B cells also showed increased levels of ICAM-1 (Figure 4K, S7C), which binds to LFA-1 on T cells and forms the peripheral SMAC of the IS. Taken together, our transcriptomic and flow cytometric analysis suggests that CNS B cells show enhanced capacity for antigen presentation, co-stimulation and formation of immunological synapses.

## T cells form long-lasting contacts with B cells in meningeal eLFs

Meningeal eLFs have been proposed to perpetuate chronic inflammation in the CNS compartment by providing a favorable environment for the interaction of T and B cells. This view is supported by our phenotypic analyses of CNS B cells that revealed their enhanced capacity for interaction with T cells. However, to our knowledge, it has never been demonstrated whether and to what effect T and B cells interact in meningeal eLFs. Therefore, we employed intravital two-photon microscopy to visualize the behavior of T and B cells in meningeal eLFs *in situ* and, in particular, to determine whether B cells are able to reactivate T cells (Figure 5A). To fluorescently label T cells, we used the ratiometric calcium sensor Twitch-2B, which allows for studying T cell activation in real-time by visualizing the increase of the intracellular calcium concentration upon TCR ligation (Thestrup et al., 2014). MOG-specific Th17 cells were transduced with a retrovirus containing Twitch-2B during differentiation, and we found that transduction efficiency was highest after restimulation with up to 60 % Twitch-2B-expressing Th17 cells (Figure S8A). Transduced Th17 cells were transferred into Mb1.Cre x ROSA26 tdTomato mice, which express tdTomato specifically in B cells (Figure 5A), and we confirmed that the transduced Th17 cells survived well *in vivo,* infiltrated the CNS, and showed stable expression of both Twitch-2B+ (Figure S8B) and effector cytokines in the CNS (Figure S8C). Furthermore, CNS-infiltrating B cells in Mb1.Cre x ROSA26 tdTomato Th17-EAE mice faithfully expressed tdTomato detected by flow

cytometry as well as confocal microscopy of CNS cryosections (Figure S8D). In contrast,
other CNS-infiltrating APCs such as CD11b$^+$ macrophages showed frequencies of tdTomato$^+$
cells below 4 % (not shown). In all diseased Th17 recipients, intravital microscopy of spinal
cord showed massive accumulation of Twitch-2B$^+$ T cells and clustered tdTomato$^+$ B cells in
the meninges, revealing eLFs of substantial size and structure (Figure 5B). Since in this
experiment, the transduction rate of T cells ranked at 58 %, it can be assumed that the
accumulation of transferred T cells is even underestimated in this image. Zooming inside
eLFs revealed extensive T:B cell contacts within these structures (Figure 5C). To compare
the T cell response of the MOG-specific 2D2 Th17 cells to non-encephalitogenic Th17 cells,
a set of control experiments was performed, where ovalbumin (OVA)-specific OT-II Th17
cells were transduced with the Twitch-2B sensor and transferred together with non-
transduced 2D2 Th17 cells into Mb1.Cre x ROSA26 tdTomato mice. Both 2D2 and OT-II T
cells were investigated regarding their interaction with B cells by monitoring the movement of
individual cells in the 3D follicle structure over time. Intriguingly, 2D2 T cells interacted more
intensely with B cells than OT-II T cells. Around half of the 2D2 T cells formed contacts with
B cells compared to only 15 % of OT-II T cells (Figure 5D). 2D2 T cells and B cells engaged
in very long contacts, which often lasted during an entire observation period of 35 min. In
contrast, OT-II T cells formed B cell contacts of much shorter duration (Figure 5E). Setting a
threshold of five minutes to define short and long contacts, as was suggested by measuring
the contact length of B cells and Tfh cells in regular GC reactions (Jacobsen et al., 2021), we
found that the majority of 2D2 T cells (82 %) engaged in long T:B cell contacts compared to
only a third of OT-II T cells (Figure 5F). Accordingly, 2D2 T cells engaged in T:B cell contacts
also showed a reduced velocity (mean 1.63 µm/min) compared to 2D2 T cells without B cell
contact (mean 2.47 µm/min), while OT-II T cells exhibited significantly higher motility (mean
3.00 µm/min) than both 2D2 T cell groups (Figure 5G). Taken together, this indicates that
MOG-specific 2D2 T cells are being arrested due to cognate interaction with B cells.

## B cells reactivate T cells in meningeal eLFs

To investigate whether 2D2 T cells become activated during B cell contact, we made use of
the ratiometric calcium sensor Twitch-2B, which represents a powerful tool to visualize
intracellular calcium signaling in activated T cells. Importantly, many 2D2 T cells showed
elevated calcium levels and, in addition, appeared very blastic, indicating that they are highly
activated (Figure 6A, Movie 1). Quantitative analysis of the calcium ratios revealed that 2D2
Th17 cells exhibited highly increased calcium levels compared to OT-II Th17 cells. To
correctly interpret increased intracellular calcium concentrations as calcium signaling, we set
a threshold at the maximal level reached spontaneously by 95 % of OT-II T cells. For 2D2 T
cells, 75.0 % of calcium ratios in T cells with B cell contact exceeded the threshold compared
to 47.2 % of calcium ratios in T cells without B cell contact, showing that T cells exhibit more
pronounced calcium signaling during T:B cell contact (Figure 6B). Notably, the majority of
2D2 T cells (81.7 %) exhibited calcium spikes compared to less than a third of OT-II T cells
(Figure 6C). 2D2 T cells displayed long-lasting calcium signaling, which could often be
observed during an entire observation period of 35 min, whereas OT-II T cells primarily
showed calcium signaling of short duration (Figure S8E). To define short- and longterm
calcium spikes, we used a cut-off at two minutes since calcium spikes longer than two
minutes require antigen-dependent T cell activation (Kyratsous et al., 2017). We found that
about 60% of 2D2 T cells showed long-term signaling compared to less than 7 % of OT-II T
cells. 2D2 T cells also showed more short-term signaling than OT-II cells, whereas the

majority of OT-II cells showed no calcium signaling at all (Figure 6D). The difference in T cell
response upon B cell contact became even more evident by comparing single cell tracks of
2D2 and OT-II T cells. 2D2 T cells exhibited pronounced calcium signaling during B cell
contact (Figure 6E, S8F), whereas calcium ratios of OT-II T cells remained unchanged during
B cell contact (Figure 6F, S8G). For better visualization of T:B cell interaction and the
ensuing T cell response, Figure 6G depicts snapshots of representative 2D2 and OT-II T
cells during B cell contact. The first 2D2 T cell formed a long-lasting contact with one and
later two B cells. While the cell appeared to be arrested and only underwent minor changes
in cell shape, it displayed high calcium signaling for over 30 min. In the second example, the
2D2 T cell showed low calcium levels before contacting the B cell and increasing calcium
levels upon contact formation. In contrast, the OT-II T cell did not show a difference in
calcium levels before and during contact with a B cell (1.4 min - 14.8 min).Together, these
data indicate that B cells in eLFs provide a reactivation stimulus to 2D2 T cells, but not to
OT-II T cells.

## T cells depend on CNS B cells to sustain pro-inflammatory effector functions in the CNS

Intravital imaging showed long-lasting T:B cell contacts in eLFs that lead to increased $Ca^{2+}$
signaling indicative of T cell reactivation. To investigate the functional consequences of these
interactions, we transferred 2D2 Th17 cells into B cell-deficient Mb1-KO mice. At the peak of
disease, infiltrating cells were recovered from the CNS, and T cells were analyzed for their
cytokine profile. Transferred T cells could be distinguished from endogenous T cells due to
their expression of the transgenic MOG-specific $V\alpha3.2$ TCR chain. Interestingly, the
frequency of IL-17A$^+$, IFN$\gamma^+$ and IL-17A$^+$IFN$\gamma^+$ producers was significantly decreased among
transferred T cells in Mb1-KO mice compared to WT mice. Similarly, endogenous T cells,
which mainly produced IFN$\gamma$, showed lower levels of IL-17A$^+$, IFN$\gamma^+$ and IL-17A$^+$IFN$\gamma^+$ cells in
Mb1-KO mice (Figure 7A). However, in the spleen we did not find decreased T cell effector
cytokines in the absence of B cells (Figure 7B), strengthening the importance of the
reactivation stimulus provided by B cells in the eLFs in the CNS rather than in the periphery.

## Discussion

In our study, we combined flow cytometric, transcriptomic and imaging analyses to investigate the unique relationship of Th17 cells and B cells and, in particular, to shed light on their cooperation in meningeal eLFs. In our Th17 adoptive transfer EAE model, which features formation of large eLFs within the meninges of brain and spinal cord, these follicles contain clusters of proliferating T and B cells, show expression of germinal center markers, and are reminiscent of eLFs in MS patients. Through single-cell RNA sequencing and flow cytometric analyses we discovered that substantial subsets of CNS-infiltrating B cells are highly activated, poised for entering GC reactions and undergoing clonal expansion. In addition, CNS B cells respond to T cell cytokines and show enhanced capacity for antigen presentation, co-stimulation and formation of immunological synapses. Correspondingly, intravital imaging provided the first real-time observation of intense T:B cell cooperation in meningeal eLFs and demonstrated that T and B cells form antigen-dependent long-lived contacts, whereby B cells provide a reactivation stimulus to T cells. Consequently, we showed that this reactivation stimulus from B cells is required for CNS T cells to maintain their highly pro-inflammatory cytokine profile.

Histological analyses in the Th17 adoptive transfer EAE model suggest that eLFs appear in different stages of maturation, even within the same animal, and that many eLFs are not fully matured at the time point of investigation (Figure 1). Single-cell transcriptomic analysis of CNS B cells largely confirmed this observation, as we identified several clusters of substantial size of B cells expressing markers of activation and markers associated with light zone GC B cells, but only very few dark zone GC B cells (Figure 2). Therefore, we suggest that our model features mostly eLFs harboring very early stages of GC reactions and this is probably due to the rather short disease duration in our model: since B cells only start accumulating in the CNS at the beginning of clinical symptoms, they had only 4-7 days to arrive and mature in eLFs until the recipients were analyzed. We speculate that the GC reaction in CNS eLFs would progress to more advanced stages with time, however, due to the severe disease course these later stages cannot be studied in our model. Nonetheless, we found elevated titers of MOG-specific antibodies in the serum at the late peak of disease compared to very low levels before disease onset (Figure S6A) and some eLFs containing massive amounts of MOG-specific B cells (Figure S6C). This shows the capacity of eLFs to support initial priming and maturation or - at the very least - expansion of autoreactive B cells that may have been primed in the periphery and then migrated to eLFs to expand. This speaks for an active disease-promoting role of eLFs since CNS-reactive autoantibodies are known to be pathogenic in CNS autoimmunity (Mader et al., 2020). In addition, eLFs have been suspected to mediate cortical damage in the underlying parenchyma by releasing soluble pro-inflammatory factors, which may include CNS-reactive antibodies but also cytokines. In fact, we found CNS B cells to respond strongly to TNFα (Figure 4D) and IL-21 (Figure 4B), an important Th17 effector cytokine, suggesting that the eLF cytokine milieu is enriched for those cytokines in comparison to the periphery. Especially TNFα has pleiotropic effects on microglia, neurons, astrocytes and oligodendrocytes (Raffaele et al., 2020), is increased in the CSF of MS patients with high levels of meningeal inflammation and grey matter damage (Magliozzi et al., 2018), and is therefore a prime candidate for further investigation regarding the association of eLFs and cortical damage. Although IL-21 is primarily known to act on immune cells, IL-21 receptor is also expressed by cortical neurons and expression is up-regulated in MS patients (Tzartos et al., 2011). In a stroke model, IL-21 signaling in neurons was associated with increased autophagy (Clarkson et al., 2014), suggesting that IL-21 may also be an interesting candidate for mediating cortical damage underneath eLFs. The effect of other Th17-derived cytokines on B cells such as IL-17A was

not investigated, since there are no gene expression signatures available. Surprisingly, IFNγ-
driven signaling seemed less important for CNS B cells (Figure 4A), although some IFNγ is
produced by transferred Th17 cells, as well as by host-derived endogenous T cells in the
CNS (Figure S1D). However, the effect of IFNγ on CNS B cells may be more pronounced in
case of Th1 adoptive transfer EAE, where the CNS T cells secrete massive amounts of IFNγ
(Figure S1C) but much fewer eLFs are formed.

Notably, our study revealed the presence of B1a / MZ-like B cells in the CNS (Figure 2E-H,
S5D-F), which may result from migration of B1a / MZ B cells from the spleen or peritoneum
to the CNS. However, it is also possible that the inflamed CNS provides a microenvironment,
which induces infiltrating B cells to adopt a B1a / MZ-like phenotype. In several autoimmune
settings, MZ-like B cells infiltrate the target tissue, e.g. the thyroid gland (Segundo et al.,
2001), or the salivary glands (Groom et al., 2002). Interestingly, a MZ-like population was
also detected in CSF from MS patients suggesting that a human counterpart of the cluster 8
B1a / MZ-like B cells may also exist in MS patients (Lee-Chang et al., 2011). Our B1a / MZ-
like B cells expand clonally (Figure 3A, C), but show less evidence of a T cell help-induced
GC reaction (Figure 2C, D), which is in line with the property of regular B1a / MZ B cells to
operate more independently of T cells. Considering the high expansion factors and
expression levels of plasmablast-associated genes (Figure 3C, D), they are probably a rich
source of antibodies, but it remains to be determined whether these are autoreactive. It is
important to note that normal B1a and MZ B cells can also perform regulatory functions
(Zhang, 2013). They are more potent IL-10 producers compared to other splenic B cell
subsets (Barr et al., 2012) and may play a protective role in autoimmunity (Lenert et al.,
2005; Matsushita et al., 2018). In fact, several upregulated genes in cluster 8, such as *Fcrl5*,
*Zbtb20*, *Cd9* and *Apoe* (Table S1), were recently defined as markers of regulatory B cells by
single-cell RNA sequencing of B cells from different murine organs (Yang et al., 2021).
Therefore, it is possible that they play a regulatory function in the CNS, especially in later
disease stages and when differentiating into plasma cells.

The importance of B cells as APCs in neuroinflammation has been reported in several EAE
studies (Bettelli et al., 2006; Krishnamoorthy et al., 2006; Molnarfi et al., 2013; Parker Harp et
al., 2015; Weber et al., 2010). In our model, we demonstrated that CNS B cells reactivate
MOG-specific T cells in eLFs, as shown by intravital microscopy (Figure 6B, E, G) and
adoptive transfer into B cell-deficient Mb1-KO mice (Figure 7). However, considering that the
B cell repertoire in our WT recipient mice is polyclonal and only few eLFs were positive for
MOG-specific B cells by MOG$_{tet}$-staining (Figure S6), B cells in eLFs cannot all be MOG-
specific. Thus, we speculate that antigen bound to MHC-II is transferred to B cells from other
APCs, which are also present in eLFs (Figure S2). Although, the mechanism requires further
investigations, this could happen for example by trogocytosis (Schriek et al., 2022), or via
extracellular vesicles (Becker et al., 2021) derived from rare MOG-specific B cells or from
ubiquitous APCs including macrophages and DCs. Due to the extensive tissue damage at
this late peak of disease the area must be saturated with released CNS autoantigens ready
to be taken up and interestingly, extracellular vesicles containing MOG were also shown to
be increased in the CSF of MS patients during disease activity (Galazka et al., 2018).
Another point of consideration is that the 2D2 TCR does not exclusively recognize MOG, as
it was shown to also respond to neurofilament (Krishnamoorthy et al., 2009). Nonetheless,
we demonstrated that OT-II T cells do not become reactivated by B cells in eLFs even
though they are also differentiated into Th17 cells (Figure 6B-D, F, G), indicating that the
reactivation stimulus provided by the B cells to the 2D2 T cells must be CNS antigen-
dependent.

The clinical significance of eLFs requires further investigation. In MS, formation of eLFs is
associated with a more severe disease course and subpial cortical as well as spinal cord

pathology (Howell *et al.*, 2011; Magliozzi *et al.*, 2007; Magliozzi *et al.*, 2010; Reali *et al.*, 2020; Serafini *et al.*, 2007; Serafini *et al.*, 2004). Due to the short and severe disease course, our Th17-transfer EAE model is not suitable to study the clinical impact of eLFs, but in the spontaneous opticospinal EAE model, which features both genetically modified MOG-specific T and B cells and a milder disease course that allows for longer observation periods, a correlation between eLF presence and a more severe disease course was found (Dang *et al.*, 2015). In the same model, a recent study suggested an immunoregulatory function of eLFs since eLF abrogation resulted in a higher disease burden (Mitsdorffer et al., 2021). It is possible that eLFs predominantly form in severe disease cases in an attempt to contain excessive inflammation, and this would involve presence of regulatory $T_{FR}$ cells and/or regulatory B cells producing anti-inflammatory cytokines. Although, $Foxp3^+$ regulatory $T_{FR}$ cells were not detected in eLFs in post-mortem CNS tissue from MS patients (Bell et al., 2019), Mitsdörffer and colleagues identified B cells producing the immunoregulatory cytokines IL-10 and IL-35 in eLFs in the opticospinal EAE model (Mitsdorffer *et al.*, 2021). Although we did not detect IL-10 or IL-35 in our transcriptomic dataset, we also found markers of regulatory B cells expressed in the B1a / MZ-like cluster 8 B cells. Thus, eLFs may provide a supportive environment for regulatory B cells and this may become more relevant at later time points, as differentiation to plasma cells, which are a known source of anti-inflammatory cytokines, progresses further. eLFs with disease-promoting and regulatory impact may co-exist in MS either in different subsets of patients or in different phases of the disease. Therefore, both models are important to study the role of eLFs in CNS autoimmunity, as each of them may represent one end of the spectrum of what can happen within eLFs, ranging from immunoregulation to immunopathology.

In contrast to the opticospinal EAE model, Th17-mediated EAE employs non-transgenic B cells with an endogenous repertoire and therefore, the B cell response in our model can be assumed to better reflect B cell-related disease aspects in MS. The possibility that eLFs support initial priming and maturation and/or expansion of CNS-reactive B cell clones has clear immunopathological implications and the increased MOG-specific antibody titers and rare $MOG_{tet}$-positive eLFs in our model point in that direction, although more research is required to confirm eLFs as sources of CNS-reactive B cells. In addition, our data suggests that - maybe even independent of their own specificity - B cells in eLFs excel at presenting CNS antigen to interacting T cells and reactivating them to maintain pro-inflammatory effector functions. Taken together, we conclude that the bilateral interactions of T and B cells in meningeal eLFs in our model support smoldering inflammatory processes that could drive CNS pathology even behind a closed blood brain barrier, and propose that further research investigating whether T:B cell cooperation in eLFs can be therapeutically targeted is warranted.

## Materials and methods

### Mice

C57BL/6 mice were purchased from Janvier Labs. C57BL/6 mice with a transgenic TCR specific for the $MOG_{35-55}$ peptide, here referred to as 2D2 mice, have been described previously (Bettelli et al., 2003) and were purchased from The Jackson Laboratory. $IgH^{MOG}$ mice on a C57BL/6 background, also known as Th mice, carrying an Ig heavy chain knock-in of the rearranged VDJ gene from the MOG-specific monoclonal antibody 8.18C5, have been described previously (Litzenburger et al., 1998) and were kindly provided by Prof. Hartmut Wekerle. Mb1-Cre mice on a C57BL/6 background carry a codon optimized Cre recombinase, replacing exon 2 and 3 of the *Cd79a* gene, which encodes the Ig-α subunit of the BCR and is exclusively expressed in B cells. This knock-in leads to abolishment of endogenous *Cd79a* gene function and expression of Cre recombinase under control of the *Cd79a* promoter (Hobeika et al., 2006). Mb1-Cre mice were purchased from The Jackson Laboratory. Homozygous mice are B cell-deficient and hereafter referred to as Mb1-KO mice. Mb1-Cre mice were crossed to ROSA26 tdTomato mice to obtain mice that are heterozygous for both Mb1-Cre and ROSA26 tdTomato, thereby expressing tdTomato specifically in B cells for intravital microscopy experiments. ROSA26 tdTomato mice on a C57BL/6 background carry a *loxP*-flanked STOP cassette and *tdTomat*o gene, both inserted into the ROSA26 locus. Upon Cre-mediated recombination, tdTomato fluorescence is robustly expressed in these mice (Madisen et al., 2010). ROSA26 tdTomato mice were purchased from The Jackson Laboratory. OT-II mice, carrying a transgenic TCR specific for the $ovalbumin_{323-339}$ peptide on a C57BL/6 background (Barnden et al., 1998), were purchased from The Jackson Laboratory.

All animals used in this study were housed and bred in specific pathogen-free conditions in the Core Facility Animal Models at the Biomedical Center of the Ludwig-Maximilians-Universität München. Animal experiments were designed and performed in accordance with regulations of the animal welfare acts and under approval by the animal ethics committee of the state of Bavaria (Regierung von Oberbayern) in accordance with European guidelines.

### Isolation of cells from spleen and lymph nodes

Spleens were mashed and centrifuged at 300 g for 5 minutes (min) at 4 °C. To lyse erythrocytes, the cells were resuspended in ACK buffer (150 mM NH4Cl (Merck), 10 mM KHCO3 (Merck), 0.1 mM EDTA (Sigma)) and incubated for 1.5 min at RT. The reaction was stopped by adding 9 ml T cell medium (RPMI 1640 (Sigma) supplemented with 10 % heat-inactivated FBS, 1 % penicillin-streptomycin, 10 mM HEPES, 2 mM L-glutamine, 1 % non-essential amino acids, 1 mM sodium pyruvate and 50 µM β-mercaptoethanol), and the cell suspension was passed through a 70 µm cell strainer. After centrifugation, the cells were resuspended in T cell medium. Lymph nodes (LNs) were cut into pieces, mashed and filtered through a 70 µm cell strainer. Then, the cells were centrifuged at 300 g for 5 min at 4 °C and resuspended in T cell medium.

### *In vitro* differentiation of T effector cells

Naïve CD4[+] T cells were purified from spleen and lymph nodes of 6-15 week old 2D2 mice or OT-II mice using the naïve CD4[+] T cell isolation Kit (Miltenyi Biotec) according to the

manufacturer`s instructions. After isolation of naïve CD4$^+$ T cells the rest of the cells (flow-
through from T cell purification) was irradiated at 35 Gy and used as APCs for costimulation
of differentiating T cells. Naïve T cells were cultured at a concentration of 1.5-2x10$^6$ ml$^{-1}$ in T
cell medium in the presence of 7.5-10x10$^6$ ml$^{-1}$ irradiated APCs (ratio T cells : APCs is 1 : 5)
and 2.5 µg/ml soluble anti-CD3 antibody (clone 145-2C11, BioXCell). Th1 cells were
generated by addition of IL-12 at a concentration of 10 ng/ml and anti-IL-4 antibody (clone
11B11, BioXCell) at a concentration of 10 µg/ml into the culture. For the generation of Th17
cells, naïve T cells were cultured with IL-6 at a concentration of 30 ng/ml, TGFβ at a
concentration of 3 ng/ml, IL-1β at a concentration of 20 ng/ml, and anti-IFNγ (clone XMG1.2,
BioXCell) and anti-IL-4 antibody (clone 11B11, BioXCell) at a concentration of 10 µg/ml. After
48h, Th1 cells and Th17 cells were split with medium containing 10 ng/ml of IL-2 and
medium containing 10 ng/ml of IL-23, respectively. All cytokines were purchased from
Biolegend except IL-23 (Miltenyi Biotec). The different T cell subsets were analyzed for
cytokine production after 4 days by intracellular cytokine staining and subsequent flow
cytometry. After 5-8 days of primary culture, T cells reached a resting state and were
restimulated at a concentration of 2x10$^6$ ml$^{-1}$ for 48 h in the presence of plate-bound anti-CD3
(clone 145-2C11, BioXCell) and anti-CD28 (clone PV-1; BioXCell) antibodies both at 2 µg/ml
in fresh medium without any cytokines.

## Retrovirus production

Phoenix-eco cells (Pear et al., 1993) were routinely cultured in TCM medium (DMEM
(Sigma) containing 10 % heat-inactivated FBS, 1 % penicillin-streptomycin, 2 mM L-
glutamine, 360 mg/l asparagine, 1 % non-essential amino acids, 1 mM sodium pyruvate and
57.2 µM β-mercaptoethanol). The cells were kept in a subconfluent state by splitting them
every 2-3 days. For production of retroviral particles, the cells were plated at 1.5-2x10$^6$ per
10 cm dish in 10 ml TCM medium and allowed to attach for 18-24 hours. First, chloroquine
diphosphate (Sigma) was added at a concentration of 25 µM. Per dish, 12 µg pMSCV-Δneo-
Twitch-2B plasmid (kindly provided by Dr. Naoto Kawakami) and 3.5 µg pCL-Eco packaging
plasmid (kindly provided by Dr. Gurumoorthy Krishnamoorthy) were prepared in 438 µl H$_2$O.
Subsequently, 62 µl 2 M CaCl$_2$ (Riedel de Haën) were added followed by dropwise addition
of 500 µl 2X BES (50 mM N,N-bis(2hydroxyethyl)-2-aminoethanesulfonic acid (Sigma) + 280
mM NaCl (Sigma) + 1.5 mM Na$_2$HPO$_4$ (Roth) in H$_2$O) while vortexing. This mixture was
incubated for 20 min at 37 °C and then added dropwise to the cells. On the next day, the
medium was replaced with 8 ml fresh and warm TCM medium. After another 24 hours, the
cell culture supernatant was collected using a 10 ml syringe, filtered through a 0.45 µm filter
and kept at 4 °C. To every dish, 4 ml fresh and warm TCM medium was added and the cells
were incubated for another day. The supernatant was collected again, filtered through a 0.45
µm filter and pooled with the supernatant from the day before. Then, the supernatant was
concentrated by centrifugation at 1000 g at 4 °C through Amicon® Ultra 15 mL Centrifugal
Filters (100 K; Merck). Finally, the concentrated supernatant was snap frozen and stored at -
80 °C.

## Retroviral transduction of Th17 cells

*Retroviral transduction during primary phase*: After 48 h of primary culture, Th17 cells were
split (usually 1:3) and transduced by adding 1:80-diluted retroviral supernatant and 4 µg/ml
polybrene (Sigma) in addition to 10 ng/ml IL-23 and 10 µg/ml anti-IFNγ antibody (all diluted in

pre-warmed T cell medium and calculated for culture volume). Then, cells were spin-infected
at 480 g for 90 min at 30 °C. On the next day, half of the medium was replaced with fresh
and warm T cell medium containing 10 ng/ml IL-23 and 10 µg/ml anti-IFNγ antibody. 48 h
after transduction, Twitch-2B expression was analyzed via GFP-signal by flow cytometry.

*Retroviral transduction during secondary phase*: 24 h after restimulation with anti-CD3/CD28
antibodies, half of the medium was taken up and Th17 cells were transduced by adding 1:80-
diluted retroviral supernatant and 4 µg/ml polybrene (Sigma) (all diluted in pre-warmed T cell
medium and calculated for culture volume). Then, cells were spin-infected at 480 g for 90
min at 30 °C. On the next day, half of the medium was replaced with fresh and warm T cell
medium. If cells were used for EAE experiments, cells were collected and washed for
adoptive transfer, while an aliquot of cells, with half of the medium being replaced with fresh
and warm T cell medium, was kept for determining the transduction efficacy. 48 h after
transduction, Twitch-2B expression was analyzed via GFP-signal by flow cytometry.

## Adoptive transfer EAE
Naïve MOG-specific CD4$^+$ T cells, isolated from spleen and lymph nodes of 2D2 mice, were
differentiated into Th1 or Th17 cells and restimulated with anti-CD3 and anti-CD28
antibodies. 48 h after restimulation, cells were collected and washed three times with PBS. In
Th1-EAE, 1.7-3.2x10$^6$ IFNγ-producing cells were injected intraperitoneally (i.p.) into C57BL/6
recipients. In Th17-EAE, 4x10$^6$ IL-17A-producing cells were injected i.p. or intravenously (i.v.)
into C57BL/6 or Mb1-KO or Mb1Cre x ROSA26 tdTomato recipients. In case of 2D2 and OT-
II T cell co-transfer, 3x10$^6$ IL-17A-producing 2D2 cells and 2x10$^6$ IL-17A-producing OT-II cells
were injected together i.p. into Mb1Cre x ROSA26 tdTomato recipients. Animals of both
genders were used as donors, whereby male cells were injected only into male recipients.
Clinical disease developed 8-15 days post transfer. Mice were monitored daily for the
development of clinical signs of EAE, including classical and atypical disease signs,
according to the following criteria: 0, no disease; 0.5, decreased tail tonus; 1, limp tail or mild
balance defects; 1.5, limp tail and ataxia; 2, hind limb weakness, or severe balance defects
that cause spontaneous falling over; 2.5, partial hind limb paralysis; 3, complete hind limb
paralysis or very severe balance defects that prevent walking; 3.5, complete hind limb
paralysis and partial front limb paralysis; 4, complete front and hind limb paralysis or inability
to move body weight into a different position; 5, moribund state.

## Isolation of infiltrating mononuclear cells from the CNS
Recipient mice, which had been on a high score for several days to allow for continued
immune cell infiltration, were sacrificed and perfused through the left cardiac ventricle with
PBS. Brains were dissected by removing the skull using forceps and scissors, while spinal
cords were flushed out from the bone column using a 21 gauge needle attached to a syringe
filled with PBS. In addition, the vertebral column was opened to collect parts of meninges,
which were still connected to the vertebrae. Then, brains and spinal cords were cut into
pieces and digested for 30 min at 37 °C with DNase I (Sigma-Aldrich or Roche) at a
concentration of 1 mg/ml and collagenase D (Roche) at a concentration of 3.75 mg/ml.
Subsequently, the tissues were mashed and passed through a 70 µm cell strainer to prepare
a single-cell suspension, followed by centrifugation at 350 g for 10 min at 4 °C. The cell pellet
was resuspended in 5 ml 70 % Percoll (Cytiva), which was prepared by mixing 30 ml Percoll

with 18 ml Percoll Mix Solution (mixture of 90 ml 10X PBS and 264 ml $H_2O$). Then, the cell
suspension was slowly and carefully overlaid with 5 ml 37 % Percoll, which was prepared by
diluting the 70 % Percoll solution with 1X PBS, followed by centrifugation at 650 g for 30 min
at 21 °C. Then, mononuclear cells were collected from the interface between the 70 % and
37 % layer and washed twice with T cell medium to remove residual Percoll. Finally, cells
were either stimulated for intracellular cytokine staining or stained directly for flowcytometric
analysis.

## Production of MOG tetramer

Mouse $MOG_{1-125}$ protein (mMOG) was produced in human embryonic kidney cells expressing
the Epstein-Barr virus nuclear antigen-1 (HEK-EBNA1) according to previously published
protocols (Perera et al., 2013), a cell line designed for large-scale production of recombinant
proteins (Tom et al., 2008). Monomer production and tetramerization was performed as
previously described (Salvador et al., 2022). Briefly, purified mMOG monomers were
biotinylated using BirA ligase to enable multimerization. The biotin-binding protein
streptavidin was used to form stable tetramer structures consisting of four biotinylated
mMOG monomers. As the streptavidin was directly conjugated to AF 488, staining with MOG
tetramer could be used to identify MOG-specific B cells by flow cytometry or confocal
microscopy.

## Flow cytometry and cell sorting

For intracellular cytokine staining, cultured or isolated cells were stimulated with 50 ng/ml
PMA (Sigma) and 500 ng/ml ionomycin (Sigma) in the presence of 0.7 µL/ml monensin
(GolgiStop; BD Biosciences) for 4 h. For surface stainings, the cells were washed with PBS.
To exclude dead cells and debris, the samples were stained with the Zombie Violet™ or
UV™ fixable viability kit (Biolegend) diluted 1:600 in PBS for 20 min at RT. Then, cells were
washed twice with FACS buffer (PBS containing 2 % FBS and 0.05 % sodium azide), and
unspecific binding of antibodies to Fc receptors was blocked by incubating the cells with
TruStain FcX™ (Biolegend) diluted 1:50 in FACS buffer for 15 min at 4 °C. The last step was
omitted when *in vitro* differentiated T cells were analyzed. Afterwards, cells were stained with
fluorophore-conjugated antibodies against surface markers diluted in FACS buffer for 20-30
min at 4 °C. If biotinylated antibodies were used, the samples were washed and
subsequently stained with fluorophore-conjugated streptavidin (SA) diluted in FACS buffer for
20 min at 4 °C. After two washing steps, cells were resuspended in FACS buffer for analysis.
For intracellular cytokine staining, cells were fixed for 30 min at 4°C with 0.4 %
paraformaldehyde (PFA, Merck KGaA) and permeabilized by washing twice with
permeabilization buffer (PBS containing 2 % FBS and 0.1 % saponin (Sigma)). Then,
samples were incubated with antibodies against intracellular markers diluted in
permeabilization buffer for 30 minutes at 4 °C, washed twice and resuspended in FACS
buffer for analysis. The following antibodies/reagents were used for flow cytometry: anti-
CD19 (1D3 or 6D5), anti-CD1d (1B1), anti-CD4 (RM4-5), anti-TCR Vα3.2 (RR3-16), anti-
B220 (RA3-6B2), anti-CD11b (M1/70), anti-CD21/CD35 (7E9), anti-CD11c (N418), anti-
CD69 (H1.2F3), anti-CD83 (Michel-19), anti-CD40 (HM40-3), anti-IL-21R (4A9), anti-ICAM-1
(YN1/1.7.4), anti-IFNγ (XMG1.2), anti-IL-10 (JES5-16E3), anti-IL-17A (TC11-18H10.1, all
Biolegend), anti-CD138 (REA104, Miltenyi), anti-GL7 (GL7), anti-IgD (11-26c.2a, all BD
Pharmingen), anti-CD45 (30-F11), anti-IgM (II/41), anti-CD86 (GL1), anti-MHC-II (NIMR-4),

anti-IL-17A (eBio17B7, all eBioscience), anti-TCR Vß5.1/5.2 (MR9-4, Invitrogen), PNA
(Vector Laboratories). Data were acquired on a BD LSRFortessa[TM], BD FACSverse[TM] flow
cytometer or Beckman Coulter CytoFLEX S flow cytometer and analyzed using Flowjo
software (version 10.6.2; TreeStar). For sorting of B cells, surface staining of $1x10^6$ cells from
spleen and cLN or $2.7\text{-}3.7x10^6$ cells from CNS samples was performed in T cell medium for
20 min at 4 °C. After two washing steps, cells were resuspended in T cell medium containing
2 mM EDTA (Sigma), filtered through a 35 μm cell strainer and kept on ice. $CD45^{high}$
$CD11b^{low}$ $CD4^-$ $CD19^+$ were sorted using a BD FACSAria™ Fusion No1.

## Immunohistochemistry (IHC) on cryosections

To increase the chances for finding eLFs in the CNS, IHC was performed using Th17-EAE
mice that had a high score for several days. The mice were perfused through the left cardiac
ventricle with 10 ml PBS and 10 ml 4 % PFA. Then, the brain was dissected by carefully
removing the skull using forceps and scissors. After removal of the outer muscle layer, the
spinal cord was dissected by snapping each vertebra open with forceps to carefully remove it
from the vertebral column in one piece leaving the leptomeninges as intact as possible.
Spleens and lymph nodes were also dissected as control tissue. After post-fixation in PBS
containing 4 % PFA for 2 h at 4 °C, the organs were dehydrated in PBS containing 30 %
sucrose for 24 h at 4 °C. Then, they were embedded in O.C.T. compound medium (Sakura
or VWR Chemicals), frozen using dry ice and stored at -80 °C. Subsequently, the brain was
cut into 20 μm coronal sections on a Leica CM1850 cryostat. From spinal cord, spleen and
lymph nodes, 10 μm sections (longitudinal for spinal cord and spleen) were prepared and
placed on slides (Superfrost Plus[TM], Thermo Scientific) and frozen at -80 °C.

### Giemsa staining

To screen for eLFs, every 15[th] section of brain and spinal cord was fixed in 10 % methanol for
10 min, air-dried for 20-30 min and stained with Giemsa-stain (AppliChem), diluted 1:20 in
deionized water, for 45 min. After washing with deionized water, the sections were dried
overnight and mounted using Eukitt mounting medium (Sigma). Sections were screened for
immune cell infiltrates by light microscopy using a Leica DM2500 microscope with DMC2900
CMOS camera.

### Immunofluorescence staining

Following identification of eLF-containing sections via Giemsa staining, neighboring sections
were stained with fluorescently labeled antibodies. After thawing, sections were fixed in cold
acetone for 10 min and air-dried for 20-30 min. Then, sections were outlined with a Pap-pen
and O.C.T. medium was rinsed off twice with PBS. Afterwards, sections were blocked with
PBS containing 1 % bovine serum albumin (BSA) and 1:300-diluted TruStain FcX™
(Biolegend) for 1 h at RT, and incubated with primary antibodies diluted in PBS containing 1
% BSA in a chamber humidified with PBS for 1 h at RT, followed by three washing steps with
PBS containing 0.1 % Tween20 for 7 min. Then, sections were stained with secondary
antibodies diluted in PBS containing 1 % BSA in a dark, humidified chamber for 1 h at RT.
When intracellular markers, such as activation-induced cytidine deaminase (AID), were
stained, the sections were incubated with antibodies diluted in PBS containing 1 % BSA and
0.1 % saponin to permeabilize the tissue. Following three washing steps with PBS containing
0.1 % Tween20 for 5 min, sections were mounted with VECTASHIELD Antifade Mounting
Medium with DAPI (Vector Laboratories) and analyzed by confocal microscopy using a Leica
SP8X WLL microscope. The following antibodies/reagents were used for

immunofluorescence staining: Syrian hamster anti-mouse CD3e (500A2), rat anti-mouse
B220 (RA3-6B2), mouse anti-mouse/human Ki67 647 (B56, all BD Pharmingen), rabbit anti-
mouse/human laminin (Abcam), rat anti-mouse/human CD11b AF 647 (M1/70), rat anti-
mouse CD1d (1B1, both BioLegend), rat anti-mouse/human AID (mAID-2, eBioscience),
PNA (biotinylated, Vector Laboratories), goat anti-hamster AF 568, goat anti-hamster AF 647
(both Invitrogen), goat anti-rabbit AF 568, goat anti-rabbit AF 647, goat anti-rabbit Cy3, goat
anti-rat AF 633, streptavidin AF 488 (all Life technologies).

*MOG tetramer staining*
Sections were fixed in cold acetone for 5 min and O.C.T. medium was rinsed off three times
with PBS for 5 min. Then, sections were outlined with a Pap-pen and blocked with PBS
containing 4 % BSA, 4 % goat serum and 1:300-diluted TruStain FcX™ (Biolegend) for 1 h at
RT in a humidified dark chamber. After washing three times with PBS for 5 min, sections
were incubated with rat anti-mouse B220 (1:200, clone RA3-6B2, BD Pharmingen) and
mMOG$_{tet}$-SA488 (1:1000, housemade) diluted in PBS containing 4 % BSA and 1 % goat
serum overnight at 4 °C. Subsequently, sections were washed three times with washing
buffer (PBS containing 0.3 % Triton and 1 % goat serum) for 5 min and fixed in 4 % PFA for
30 min at RT. After three washing steps in washing buffer, sections were incubated with
rabbit anti-SA488 (1:10000, Life technologies) for 3 h and goat anti-rat AF 633 (1:200, Life
technologies) for 1 h diluted in PBS containing 4 % BSA and 1 % goat serum at RT in a
humidified dark chamber. After three washing steps in washing buffer, sections were
incubated with goat anti-rabbit Cy3 (1:1000, Life technologies) diluted in PBS containing 4 %
BSA and 1 % goat serum for 3 h at RT in a humidified dark chamber, followed by three
washing steps in washing buffer. Then, sections were mounted with VECTASHIELD Antifade
Mounting Medium with DAPI (Vector Laboratories) and analyzed by confocal microscopy
using a Leica SP8X WLL microscope.

**Enzyme-linked immunosorbent assay (ELISA)**
Anti-MOG IgG1 antibodies were measured in the serum of Th17- and Th1-EAE mice by
ELISA. 96 well high affinity, protein-binding plates (Nunc Maxisorp) were coated with rMOG
(housemade) at a concentration of 10 µg/ml in PBS and incubated overnight at 4 °C. After
washing four times with wash buffer (PBS containing 0.05 % Tween20), the plate was
blocked with blocking solution (PBS containing 10 % FBS) for 1-2 h at RT, followed by four
washing steps. Then, the serum samples and the antibody standard (mouse anti-MOG IgG1;
clone 8.18c5; housemade) were diluted in blocking solution, added to the plate and
incubated overnight at 4 °C. Following extensive washing, the plate was incubated with
biotin-labeled detection antibody rat anti-mouse IgG1 (clone A85-1; BD Pharmingen) at a
concentration of 1 µg/ml diluted in blocking solution for 1 h at RT. Then, the plate was
washed again and incubated with avidin-horseradish peroxidase (Av-HRP; eBioscience)
diluted 1:1000-1:2000 in blocking solution for 30 min at RT. After extensive washing, TMB
Substrate (BioLegend) was used to induce the Av-HRP-dependent color reaction, which was
stopped by addition of TMB Stop Solution (BioLegend), and the optical density (OD) was
measured at 450 nm. Samples were tested in duplicates and results were reported as mean
concentrations according to the standard curve.

## Single-cell RNA sequencing (scRNA-seq)

### *Library construction and sequencing*

For each mouse, B cells from CNS, spleen and cLNs were pooled (90.000 B cells in total), centrifuged and, after a washing step, resuspended in PBS containing 0.04 % BSA at a concentration of 1000 cells/μl. The viability of cells was confirmed by staining with trypan blue solution. Further processing of samples was performed using the 10x Chromium Single Cell 5` Solution (Chromium Next GEM Single Cell V(D)J v1.1 with Feature Barcoding Technology for Cell Surface Protein, 10x Genomics) according to the manufacturer`s instructions. Prior to loading on the chip, 5000 cells of each compartment were labeled with anti-mouse hashtag oligonucleotide (HTO) antibodies (TotalSeq™-C0301/C0302/C0303, Biolegend) and pooled together. Next, the B cells were processed with enzymes and gel beads, followed by partitioning in oil suspension to create gel beads in emulsion (GEMs) containing single cells. In GEMs, the cells` RNA was reverse-transcribed into cDNA, whereby all the cDNA from the same cell was labeled with a shared barcode. scRNA-seq, scBCR-seq and cell hashing libraries were multiplexed using individual Chromium i7 Sample Indices. Quality controls were performed with Qubit dsDNA HS kit (Life Technologies) to quantify the libraries, and Bioanalyzer 2100 (Agilent Technologies) to confirm fragment length. Then, the resulting cDNA libraries were pooled and send for sequencing. scRNA-seq and scBCR-seq libraries were sequenced on NovaSeq S4 Flowcells using 150-bp paired-end reads and 8□bp for the i7 index, aiming for 50,000 reads per cell for gene expression and 5,000 reads per cell for BCR enrichment. Cell hashing libraries were sequenced on NovaSeq S1 Flowcells using 50-bp paired-end reads and 8□bp for the i7 index, aiming for 10,000 reads per cell.

### *Data processing*

To demultiplex samples, align the reads to the reference mouse mm10 genome, and process the raw data, we used the Cell Ranger Software (10X Genomics, v. 3.1.0). Following this, the unique molecule identifiers (UMIs) were summarized and the single-cells were filtered by the number of UMI counts identified. Then, we processed the gene matrices with R (v. 4.0.5) and the R package Seurat (v. 4.0.6 and 4.9.9) (Butler et al., 2018; Stuart et al., 2019). Cell hashing raw counts were normalized using centered log-ratio (CLR) transformation by comparing the counts of single cells to the geometric mean of the HTOs signal with subsequent logarithmic transformation. Cells with missing HTO signals or containing two different HTO-identifiers were removed from the further analysis. Further, the scRNA-seq count matrices were log-normalized and scaled by passing through only the cells containing more than 200 and less than 5000 genes with mitochondrial gene fractions below 5 %. Subsequently, the highly variable genes were identified with the vst method, and cells from each compartment were integrated using the canonical correlation analysis (CCA) method. The aligned BCR sequencing information was added into the metadata file of the Seurat object with subsequent removal of cells with missing BCR sequences. The cells were considered as expanded if at least 2 cells belonged to the same B cell clonotype. In the next step, 18413 cells were recovered and the corresponding PCA was computed using the previously identified highly variable genes. Based on the first 15 PCA dimensions, we constructed the Shared Nearest Neighbor (SNN) graph and clustered the cells by using the FindNeighbours and FindClusters functions of the Seurat R package. The following dimensional reduction was performed with the help of the Uniform Manifold Approximation and Projection (UMAP) method. One cluster, cluster 11, with markedly lower quality compared to the other clusters based on nFeature and nCount, was removed from further analysis. In addition, cluster 10 was identified as expressing macrophage/microglia markers

and was therefore also excluded (based on AddModuleScore from Immgen Monocytes
MHCIIhi blood vs Fo B cells spleen and Microglia CNS vs FoB form spleen).

For identification of differentially upregulated genes per cluster, compartment, or expansion
condition we used the FindMarkers function based on the log-transformed count matrices of
each single cell by applying the Wilcoxon test and subsequent p-value adjustment with the
Bonferroni correction, with logfc.threshold = 0.

*Pathway network analysis*
For each cluster, GSEA (Mootha et al., 2003; Subramanian et al., 2005) PreRanked files
were generated with the gene and avg_log2FC columns. GSEA v4.2.3 was run with the
mouse GO BP database m5.go.bp.v2023.1.Mm.symbols
(https://data.broadinstitute.org/gsea-msigdb/msigdb/release/2023.1.Mm/), without collapsing
gene symbols, scoring_scheme classic, set_max 500, set_min 2, and otherwise default
parameters. The results were exported to Cytoscape v3.9.1 (Shannon et al., 2003), where
only the nodes with NES > 0, fdr_qvalue < 0.1 and fwer_qvalue < 0.5 were kept.
AutoAnnotate from the EnrichmentMap app collection (Merico et al., 2010) was used to
cluster and label the pathway networks, with Cluster Algorithm = MCL cluster, Label Column
= Node Names, Max Words Per Label = 4, Minimum Word Occurrence = 1, Adjacent Word
Bonus = 8, setting numbers and "GOBP" as extra Excluded Words.

*B cell functional signatures*
Functional signatures expressed by B cells were calculated with the AddModuleScore
function from Seurat from manually curated gene lists based on ImmGen
(https://www.immgen.org/), GO terms (https://www.informatics.jax.org/vocab/gene_ontology)
and literature search.

APC/synapse signature: this signature was assembled by combining the GO gene set
"lymphocyte co-stimulation" (GO:0031294, without T cell specific genes) and the GO gene
set "immunological synapse" (GO:0001772, without T cell specific genes), and then adding
the core APC signature described by (Jordão et al., 2019), the list of molecules described to
be present in the immunological synapse formed by primary murine B cells by Cunha et al.
(https://www.biorxiv.org/content/10.1101/2023.02.23.529674v1.full.pdf), and manually adding
Icosl, Tnfsf9, Timd4, Pvr, and Il21r.
B1a signature: this gene set is derived from Immgen by population comparison using the ULI
RNAsequencing datasets. B1a cells from spleen and B1a cells from the peritoneal cavity
were compared against follicular B cells from spleen, and Nfatc1 and Mzb1 were added
manually.
MZ signature: this gene set is derived from Immgen by population comparison using the ULI
RNAsequencing datasets. Here, marginal zone B cells from spleen were compared to
follicular B cells from spleen and Nfatc1 and Mzb1 were added manually.
LZ-GC and DZ-GC signature: these signatures were described by (Victora *et al.*, 2010).
mTOR signaling: Hallmark_mtorc1_signaling (mouse) gene set from the MSigDB database.
Myc signaling: Hallmark_myc_targets_v1 (mouse) gene set from the MSigDB database.
OXPHOS: Hallmark_oxidative phosphorylation (mouse) gene set from the MSigDB
database.
PB signature: this gene set is derived from Immgen by population comparison using the ULI
RNAsequencing datasets. Here, plasmablasts and plasma cells from spleen and plasma
cells from bone marrow were compared against follicular B cells from spleen.

<u>IFNα response:</u> Hallmark_interferon_alpha_response (mouse) gene set from the MSigDB
database.
<u>IFNγ response:</u> Hallmark_interferon_gamma_response (mouse) gene set from the MSigDB
database.
<u>TNFα signaling:</u> Hallmark_TNFa_signaling_via_nfkb (mouse) gene set from the MSigDB
database.
<u>IL-21 response:</u> this signature is described on Immgen and includes all genes upregulated in
follicular B cells at least 2-fold by IL-21 relative to PBS treatment.
<u>B cell activation:</u> this signature represents the GO gene set "B cell activation involved in
immune response" (GO:0002312).

***Data and code availability***
All sequencing data (fastq files, processed data files and Seurat Object) from this study is
available upon request. Custom code used for analysis is available at Zenodo under the
same doi as the published manuscript.

**Intravital two-photon microscopy**
***Animal preparation***
Th17 recipients, which had been on a high score for several days to allow for continued
immune cell infiltration and maturation of eLFs, were selected for live imaging analysis. Prior
to surgery, the mice were anesthetized with MMF anesthesia consisting of midazolam (5
mg/kg bodyweight (bw)), medetomidine (500 µg/kg bw) and fentanyl (50 µg/kg bw).
Subsequently, they were tracheotomized and ventilated with 1.5-2 % isoflurane, which was
supplied throughout the surgery and the entire imaging session. Thereby, the mice were
placed on a custom-made microscope stage, while their body temperature was stabilized at
37.5 °C using a heated pad. During imaging, breathing parameters, including concentrations
of inspiratory and expiratory gases as well as ventilation pressure, and electrocardiograms
were constantly monitored. The experimental procedure of spinal cord imaging has been
described previously (Bartholomäus et al., 2009; Mues et al., 2013)). In brief, a small imaging
window at level Th12/L1 of the spinal cord was prepared. First, a midline incision of about 2
cm was placed to open the skin and, subsequently, to detach the paravertebral musculature.
To reduce artefacts caused by breathing and thereby ensure stable imaging conditions, a
custom-made stage was used in order to fix three spines. Then, a laminectomy on the
central vertebra was performed using a dental drill (Foredom). Finally, low-melting agarose
was used to build a ring-shaped surrounding, which was filled with PBS to keep the tissue of
the opened spine hydrated and to allow using the water immersion objective.

***Image acquisition***
Time-lapse two-photon laser-scanning microscopy was performed using a SP8 multi-photon
microscope (Leica). The SP8 system was equipped with a pulsed InSight DS+ laser (Spectra
Physics). The emission wavelength was tuned to 841 nm and the fluorescent signal was
collected using a water immersion objective (25x, NA 1, Leica) and detected with external,
non-descanned hybrid photo detectors (HyDs) featuring 483/32 nm, 535/30 nm, 585/40 nm
and 650/50 nm bandpass filters. Images were recorded using a 1-2x zoom, whereby areas of
221x221 µm, 295x295 µm or 443x443 µm were scanned, and 49-115 µm z-stacks with 2.5-
4.3 µm step size were acquired with an acquisition rate of 21-22 s per z-stack.

***Image processing and analysis***
Time-lapse images were acquired using the LAS X software (Leica), followed by processing
and analysis using ImageJ (NIH). Before the stacks were z-projected with maximum intensity
to obtain two-dimensional movies, a Gaussian blur filter was applied. For Twitch-2B-
expressing cells, the cpVenus$^{CD}$ Fluorescence resonance energy transfer (FRET) channel
was divided by the mCerulean3 (CFP) channel and, by changing to a fire lookup table,
ratiometric pseudocolor images were generated. In each time frame of the two-dimensional
maximm intensity projection, the cell shapes were outlined manually to create a region of
interest (ROI) for analyzing the calcium ratios. Since the bleed-through of CFP into the YFP
channel was determined to account for 44 %, the FRET signal was corrected as follows: YFP
= FRET – 0.44 x CFP. In order to calculate the calcium ratio in every time frame, the average
signal intensities of all pixels in each ROI were used. Cell coordinates over time were used to
calculate motility parameters by using ImageJ. T:B cell contact analysis was performed by
monitoring the movement of each Twitch-2B$^+$ T cell and its interaction with tdTomato$^+$ B cells
in the 3D follicle structure using the LAS X software.

## Statistical analysis

Statistical analysis was performed using GraphPad Prism software (version 7). The statistical
test chosen is described in the figure captions. For comparison of two groups, statistical
differences were calculated using the unpaired Student`s *t*-test for normally distributed data,
including Welch`s correction in case of unequal variances, or the Mann-Whitney U test if
Gaussian distribution could not be determined. For comparison of three groups, the one-way
ANOVA with Tukey`s test for multiple comparisons was applied.

## Declaration of interests

966 The authors declare no conflict of interest.

## Acknowledgements

We thank the Core Facility Flow Cytometry and the Core facility Bioimaging at the
Biomedical Center, Ludwig-Maximilians University Munich, for providing equipment, services
and expertise. We are grateful to the staff from CCGA Kiel for their assistance with
sequencing and data transfer. We thank Isabel Gök and Klaus Hämäläinen for technical
assistance. The study was supported by the German Research Foundation (DFG) through
the Emmy Noether programme PE-2681/1–1 (to A.P.), through SFB TRR 128 (project B08 to
A.P.; project B10 to N.K. and M.K.), through CRC TRR 152 (ID 239283807; project P27; to
A.P. and M.K.), through the Heisenberg programme (KA2951/3-1 and KA2951/3-2
(264119140) to N.K. and through research grant KA2951/2-2 (246754395) to N.K.

# References

Barnden, M.J., Allison, J., Heath, W.R., and Carbone, F.R. (1998). Defective TCR expression in transgenic mice constructed using cDNA-based alpha- and beta-chain genes under the control of heterologous regulatory elements. Immunol Cell Biol *76*, 34-40. 10.1046/j.1440-1711.1998.00709.x.

Barr, T.A., Shen, P., Brown, S., Lampropoulou, V., Roch, T., Lawrie, S., Fan, B., O'Connor, R.A., Anderton, S.M., Bar-Or, A., et al. (2012). B cell depletion therapy ameliorates autoimmune disease through ablation of IL-6-producing B cells. The Journal of experimental medicine *209*, 1001-1010. 10.1084/jem.20111675.

Bartholomäus, I., Kawakami, N., Odoardi, F., Schläger, C., Miljkovic, D., Ellwart, J.W., Klinkert, W.E., Flügel-Koch, C., Issekutz, T.B., Wekerle, H., and Flügel, A. (2009). Effector T cell interactions with meningeal vascular structures in nascent autoimmune CNS lesions. Nature *462*, 94-98. 10.1038/nature08478.

Becker, P.D., Ratnasothy, K., Sen, M., Peng, Q., Romano, M., Bazoer, J., Suvitra, E., Stout, A., Hylton, S.G., Dorling, A., et al. (2021). B lymphocytes contribute to indirect pathway T cell sensitization via acquisition of extracellular vesicles. Am J Transplant *21*, 1415-1426. 10.1111/ajt.16088.

Bell, L., Lenhart, A., Rosenwald, A., Monoranu, C.M., and Berberich-Siebelt, F. (2019). Lymphoid Aggregates in the CNS of Progressive Multiple Sclerosis Patients Lack Regulatory T Cells. Frontiers in immunology *10*, 3090. 10.3389/fimmu.2019.03090.

Bettelli, E., Baeten, D., Jager, A., Sobel, R.A., and Kuchroo, V.K. (2006). Myelin oligodendrocyte glycoprotein-specific T and B cells cooperate to induce a Devic-like disease in mice. The Journal of clinical investigation *116*, 2393-2402. 10.1172/JCI28334.

Bettelli, E., Pagany, M., Weiner, H.L., Linington, C., Sobel, R.A., and Kuchroo, V.K. (2003). Myelin oligodendrocyte glycoprotein-specific T cell receptor transgenic mice develop spontaneous autoimmune optic neuritis. The Journal of experimental medicine *197*, 1073-1081. 10.1084/jem.20021603.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nat Biotechnol *36*, 411-420. 10.1038/nbt.4096.

Clarkson, B.D., Ling, C., Shi, Y., Harris, M.G., Rayasam, A., Sun, D., Salamat, M.S., Kuchroo, V., Lambris, J.D., Sandor, M., and Fabry, Z. (2014). T cell-derived interleukin (IL)-21 promotes brain injury following stroke in mice. The Journal of experimental medicine *211*, 595-604. 10.1084/jem.20131377.

Comi, G., Bar-Or, A., Lassmann, H., Uccelli, A., Hartung, H.P., Montalban, X., Sørensen, P.S., Hohlfeld, R., and Hauser, S.L. (2021). Role of B Cells in Multiple Sclerosis and Related Disorders. Annals of neurology *89*, 13-23. 10.1002/ana.25927.

Cyster, J.G., and Allen, C.D.C. (2019). B Cell Responses: Cell Interaction Dynamics and Decisions. Cell *177*, 524-540. 10.1016/j.cell.2019.03.016.

Dang, A.K., Tesfagiorgis, Y., Jain, R.W., Craig, H.C., and Kerfoot, S.M. (2015). Meningeal Infiltration of the Spinal Cord by Non-Classically Activated B Cells is Associated with Chronic Disease Course in a Spontaneous B Cell-Dependent Model of CNS Autoimmune Disease. Frontiers in immunology *6*, 470. 10.3389/fimmu.2015.00470.

Ersching, J., Efeyan, A., Mesin, L., Jacobsen, J.T., Pasqual, G., Grabiner, B.C., Dominguez-Sola, D., Sabatini, D.M., and Victora, G.D. (2017). Germinal Center Selection and Affinity Maturation Require Dynamic Regulation of mTORC1 Kinase. Immunity *46*, 1045-1058.e1046. 10.1016/j.immuni.2017.06.005.

Galazka, G., Mycko, M.P., Selmaj, I., Raine, C.S., and Selmaj, K.W. (2018). Multiple sclerosis: Serum-derived exosomes express myelin proteins. Multiple sclerosis (Houndmills, Basingstoke, England) *24*, 449-458. 10.1177/1352458517696597.

Groom, J., Kalled, S.L., Cutler, A.H., Olson, C., Woodcock, S.A., Schneider, P., Tschopp, J., Cachero, T.G., Batten, M., Wheway, J., et al. (2002). Association of BAFF/BLyS overexpression and altered B cell differentiation with Sjögren's syndrome. The Journal of clinical investigation *109*, 59-68. 10.1172/jci14121.

Hobeika, E., Thiemann, S., Storch, B., Jumaa, H., Nielsen, P.J., Pelanda, R., and Reth, M. (2006). Testing gene function early in the B cell lineage in mb1-cre mice. Proceedings of the National Academy of Sciences of the United States of America *103*, 13789-13794. 10.1073/pnas.0605944103.

Howell, O.W., Reeves, C.A., Nicholas, R., Carassiti, D., Radotra, B., Gentleman, S.M., Serafini, B., Aloisi, F., Roncaroli, F., Magliozzi, R., and Reynolds, R. (2011). Meningeal inflammation is widespread and linked to cortical pathology in multiple sclerosis. Brain : a journal of neurology *134*, 2755-2771. 10.1093/brain/awr182.

Jacobsen, J.T., Hu, W., TB, R.C., Solem, S., Galante, A., Lin, Z., Allon, S.J., Mesin, L., Bilate, A.M., Schiepers, A., et al. (2021). Expression of Foxp3 by T follicular helper cells in end-stage germinal centers. Science *373*. 10.1126/science.abe5146.

Jager, A., Dardalhon, V., Sobel, R.A., Bettelli, E., and Kuchroo, V.K. (2009). Th1, Th17, and Th9 effector cells induce experimental autoimmune encephalomyelitis with different pathological phenotypes. Journal of immunology *183*, 7169-7177. 10.4049/jimmunol.0901906.

Jordão, M.J.C., Sankowski, R., Brendecke, S.M., Sagar, Locatelli, G., Tai, Y.H., Tay, T.L., Schramm, E., Armbruster, S., Hagemeyer, N., et al. (2019). Single-cell profiling identifies myeloid cell subsets with distinct fates during neuroinflammation. Science *363*. 10.1126/science.aat7554.

Krishnamoorthy, G., Lassmann, H., Wekerle, H., and Holz, A. (2006). Spontaneous opticospinal encephalomyelitis in a double-transgenic mouse model of autoimmune T cell/B cell cooperation. The Journal of clinical investigation *116*, 2385-2392. 10.1172/jci28330.

Krishnamoorthy, G., Saxena, A., Mars, L.T., Domingues, H.S., Mentele, R., Ben-Nun, A., Lassmann, H., Dornmair, K., Kurschus, F.C., Liblau, R.S., and Wekerle, H. (2009). Myelin-specific T cells also recognize neuronal autoantigen in a transgenic mouse model of multiple sclerosis. Nature medicine *15*, 626-632. 10.1038/nm.1975.

Kuerten, S., Schickel, A., Kerkloh, C., Recks, M.S., Addicks, K., Ruddle, N.H., and Lehmann, P.V. (2012). Tertiary lymphoid organ development coincides with determinant spreading of the myelin-specific T cell response. Acta Neuropathol *124*, 861-873. 10.1007/s00401-012-1023-3.

Kyratsous, N.I., Bauer, I.J., Zhang, G., Pesic, M., Bartholomaus, I., Mues, M., Fang, P., Worner, M., Everts, S., Ellwart, J.W., et al. (2017). Visualizing context-dependent calcium signaling in encephalitogenic T cells in vivo by two-photon microscopy. Proceedings of the National Academy of Sciences of the United States of America *114*, E6381-e6389. 10.1073/pnas.1701806114.

Lee-Chang, C., Zéphir, H., Top, I., Dubucquoi, S., Trauet, J., Prin, L., and Vermersch, P. (2011). B-cell subsets up-regulate α4 integrin and accumulate in the cerebrospinal fluid in clinically isolated syndrome suggestive of multiple sclerosis onset. Neurosci Lett *487*, 273-277. 10.1016/j.neulet.2010.10.036.

Lenert, P., Brummel, R., Field, E.H., and Ashman, R.F. (2005). TLR-9 activation of marginal zone B cells in lupus mice regulates immunity through increased IL-10 production. J Clin Immunol *25*, 29-40. 10.1007/s10875-005-0355-6.

Litzenburger, T., Fassler, R., Bauer, J., Lassmann, H., Linington, C., Wekerle, H., and Iglesias, A. (1998). B lymphocytes producing demyelinating autoantibodies: development and function in gene-targeted transgenic mice. The Journal of experimental medicine *188*, 169-180.

Lucchinetti, C.F., Popescu, B.F.G., Bunyan, R.F., Moll, N.M., Roemer, S.F., Lassmann, H., Brück, W., Parisi, J.E., Scheithauer, B.W., Giannini, C., et al. (2011). Inflammatory Cortical Demyelination in Early Multiple Sclerosis. New England Journal of Medicine *365*, 2188-2197. 10.1056/NEJMoa1100648.

Mader, S., Kümpfel, T., and Meinl, E. (2020). Novel insights into pathophysiology and therapeutic possibilities reveal further differences between AQP4-IgG- and MOG-IgG-associated diseases. Curr Opin Neurol *33*, 362-371. 10.1097/wco.0000000000000813.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-throughput Cre

reporting and characterization system for the whole mouse brain. Nat Neurosci *13*, 133-140.
10.1038/nn.2467.
Magliozzi, R., Columba-Cabezas, S., Serafini, B., and Aloisi, F. (2004). Intracerebral
expression of CXCL13 and BAFF is accompanied by formation of lymphoid follicle-like
structures in the meninges of mice with relapsing experimental autoimmune
encephalomyelitis. Journal of neuroimmunology *148*, 11-23. 10.1016/j.jneuroim.2003.10.056.
Magliozzi, R., Howell, O., Vora, A., Serafini, B., Nicholas, R., Puopolo, M., Reynolds, R., and
Aloisi, F. (2007). Meningeal B-cell follicles in secondary progressive multiple sclerosis
associate with early onset of disease and severe cortical pathology. Brain : a journal of
neurology *130*, 1089-1104. 10.1093/brain/awm038.
Magliozzi, R., Howell, O.W., Nicholas, R., Cruciani, C., Castellaro, M., Romualdi, C., Rossi,
S., Pitteri, M., Benedetti, M.D., Gajofatto, A., et al. (2018). Inflammatory intrathecal profiles
and cortical damage in multiple sclerosis. Annals of neurology *83*, 739-755.
10.1002/ana.25197.
Magliozzi, R., Howell, O.W., Reeves, C., Roncaroli, F., Nicholas, R., Serafini, B., Aloisi, F.,
and Reynolds, R. (2010). A Gradient of neuronal loss and meningeal inflammation in multiple
sclerosis. Annals of neurology *68*, 477-493. 10.1002/ana.22230.
Matsushita, T., Kobayashi, T., Mizumaki, K., Kano, M., Sawada, T., Tennichi, M., Okamura,
A., Hamaguchi, Y., Iwakura, Y., Hasegawa, M., et al. (2018). BAFF inhibition attenuates
fibrosis in scleroderma by modulating the regulatory and effector B cell balance. Sci Adv *4*,
eaas9944. 10.1126/sciadv.aas9944.
Merico, D., Isserlin, R., Stueker, O., Emili, A., and Bader, G.D. (2010). Enrichment map: a
network-based method for gene-set enrichment visualization and interpretation. PLoS One *5*,
e13984. 10.1371/journal.pone.0013984.
Mitsdoerffer, M., and Peters, A. (2016). Tertiary Lymphoid Organs in Central Nervous
System Autoimmunity. Frontiers in immunology *7*, 451. 10.3389/fimmu.2016.00451.
Mitsdorffer, M., Di Liberto, G., Dotsch, S., Sie, C., Wagner, I., Pfaller, M., Kreutzfeldt, M.,
Frassle, S., Aly, L., Knier, B., et al. (2021). Formation and immunomodulatory function of
meningeal B-cell aggregates in progressive CNS autoimmunity. Brain : a journal of
neurology. 10.1093/brain/awab093.
Molnarfi, N., Schulze-Topphoff, U., Weber, M.S., Patarroyo, J.C., Prod'homme, T., Varrin-
Doyer, M., Shetty, A., Linington, C., Slavin, A.J., Hidalgo, J., et al. (2013). MHC class II-
dependent B cell APC function is required for induction of CNS autoimmunity independent of
myelin-specific antibodies. The Journal of experimental medicine *210*, 2921-2937.
10.1084/jem.20130699.
Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J.,
Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-1alpha-
responsive genes involved in oxidative phosphorylation are coordinately downregulated in
human diabetes. Nat Genet *34*, 267-273. 10.1038/ng1180.
Mues, M., Bartholomaus, I., Thestrup, T., Griesbeck, O., Wekerle, H., Kawakami, N., and
Krishnamoorthy, G. (2013). Real-time in vivo analysis of T cell activation in the central
nervous system using a genetically encoded calcium indicator. Nature medicine *19*, 778-783.
10.1038/nm.3180.
Parker Harp, C.R., Archambault, A.S., Sim, J., Ferris, S.T., Mikesell, R.J., Koni, P.A.,
Shimoda, M., Linington, C., Russell, J.H., and Wu, G.F. (2015). B cell antigen presentation is
sufficient to drive neuroinflammation in an animal model of multiple sclerosis. Journal of
immunology *194*, 5077-5084. 10.4049/jimmunol.1402236.
Pear, W.S., Nolan, G.P., Scott, M.L., and Baltimore, D. (1993). Production of high-titer
helper-free retroviruses by transient transfection. Proceedings of the National Academy of
Sciences of the United States of America *90*, 8392-8396. 10.1073/pnas.90.18.8392.
Perera, N.C., Wiesmüller, K.H., Larsen, M.T., Schacher, B., Eickholz, P., Borregaard, N., and
Jenne, D.E. (2013). NSP4 is stored in azurophil granules and released by activated
neutrophils as active endoprotease with restricted specificity. Journal of immunology *191*,
2700-2707. 10.4049/jimmunol.1301293.
Peters, A., Pitcher, L.A., Sullivan, J.M., Mitsdoerffer, M., Acton, S.E., Franz, B.,
Wucherpfennig, K., Turley, S., Carroll, M.C., Sobel, R.A., et al. (2011). Th17 cells induce

ectopic lymphoid follicles in central nervous system tissue inflammation. Immunity *35*, 986-
996. 10.1016/j.immuni.2011.10.015.

Pikor, N.B., Astarita, J.L., Summers-Deluca, L., Galicia, G., Qu, J., Ward, L.A., Armstrong, S.,
Dominguez, C.X., Malhotra, D., Heiden, B., et al. (2015). Integration of Th17- and
Lymphotoxin-Derived Signals Initiates Meningeal-Resident Stromal Cell Remodeling to
Propagate Neuroinflammation. Immunity *43*, 1160-1173. 10.1016/j.immuni.2015.11.010.

Raffaele, S., Lombardi, M., Verderio, C., and Fumagalli, M. (2020). TNF Production and
Release from Microglia via Extracellular Vesicles: Impact on Brain Functions. Cells *9*.
10.3390/cells9102145.

Rauschmeier, R., Reinhardt, A., Gustafsson, C., Glaros, V., Artemov, A.V., Dunst, J., Taneja,
R., Adameyko, I., Månsson, R., Busslinger, M., and Kreslavsky, T. (2022). Bhlhe40 function
in activated B and TFH cells restrains the GC reaction and prevents lymphomagenesis. The
Journal of experimental medicine *219*. 10.1084/jem.20211406.

Reali, C., Magliozzi, R., Roncaroli, F., Nicholas, R., Howell, O.W., and Reynolds, R. (2020).
B cell rich meningeal inflammation associates with increased spinal cord pathology in
multiple sclerosis. Brain Pathology *30*, 779-793. https://doi.org/10.1111/bpa.12841.

Salvador, F., Deramoudt, L., Leprêtre, F., Figeac, M., Guerrier, T., Boucher, J., Bas, M.,
Journiac, N., Peters, A., Mars, L.T., and Zéphir, H. (2022). A Spontaneous Model of
Experimental Autoimmune Encephalomyelitis Provides Evidence of MOG-Specific B Cell
Recruitment and Clonal Expansion. Frontiers in immunology *13*, 755900.
10.3389/fimmu.2022.755900.

Schriek, P., Ching, A.C., Moily, N.S., Moffat, J., Beattie, L., Steiner, T.M., Hosking, L.M.,
Thurman, J.M., Holers, V.M., Ishido, S., et al. (2022). Marginal zone B cells acquire dendritic
cell functions by trogocytosis. Science *375*, eabf7470. 10.1126/science.abf7470.

Segundo, C., Rodríguez, C., García-Poley, A., Aguilar, M., Gavilán, I., Bellas, C., and Brieva,
J.A. (2001). Thyroid-infiltrating B lymphocytes in Graves' disease are related to marginal
zone and memory B cell compartments. Thyroid *11*, 525-530.
10.1089/105072501750302813.

Serafini, B., Rosicarelli, B., Franciotta, D., Magliozzi, R., Reynolds, R., Cinque, P., Andreoni,
L., Trivedi, P., Salvetti, M., Faggioni, A., and Aloisi, F. (2007). Dysregulated Epstein-Barr
virus infection in the multiple sclerosis brain. The Journal of experimental medicine *204*,
2899-2912. 10.1084/jem.20071030.

Serafini, B., Rosicarelli, B., Magliozzi, R., Stigliano, E., and Aloisi, F. (2004). Detection of
ectopic B-cell follicles with germinal centers in the meninges of patients with secondary
progressive multiple sclerosis. Brain pathology (Zurich, Switzerland) *14*, 164-174.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated
models of biomolecular interaction networks. Genome Res *13*, 2498-2504.
10.1101/gr.1239303.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. Cell *177*, 1888-1902.e1821. 10.1016/j.cell.2019.05.031.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. Proceedings of the National Academy of Sciences of the United States of America
*102*, 15545-15550. 10.1073/pnas.0506580102.

Thestrup, T., Litzlbauer, J., Bartholomäus, I., Mues, M., Russo, L., Dana, H., Kovalchuk, Y.,
Liang, Y., Kalamakis, G., Laukat, Y., et al. (2014). Optimized ratiometric calcium sensors for
functional in vivo imaging of neurons and T lymphocytes. Nat Methods *11*, 175-182.
10.1038/nmeth.2773.

Tom, R., Bisson, L., and Durocher, Y. (2008). Culture of HEK293-EBNA1 Cells for
Production of Recombinant Proteins. CSH Protoc *2008*, pdb.prot4976.
10.1101/pdb.prot4976.

Tzartos, J.S., Craner, M.J., Friese, M.A., Jakobsen, K.B., Newcombe, J., Esiri, M.M., and
Fugger, L. (2011). IL-21 and IL-21 receptor expression in lymphocytes and neurons in
multiple sclerosis brain. Am J Pathol *178*, 794-802. 10.1016/j.ajpath.2010.10.043.
Victora, G.D., Schwickert, T.A., Fooksman, D.R., Kamphorst, A.O., Meyer-Hermann, M.,
Dustin, M.L., and Nussenzweig, M.C. (2010). Germinal center dynamics revealed by
multiphoton microscopy with a photoactivatable fluorescent reporter. Cell *143*, 592-605.
10.1016/j.cell.2010.10.032.
Weber, M.S., Prod'homme, T., Patarroyo, J.C., Molnarfi, N., Karnezis, T., Lehmann-Horn, K.,
Danilenko, D.M., Eastham-Anderson, J., Slavin, A.J., Linington, C., et al. (2010). B-cell
activation influences T-cell polarization and outcome of anti-CD20 B-cell depletion in central
nervous system autoimmunity. Annals of neurology *68*, 369-383. 10.1002/ana.22081.
Yang, S.Y., Long, J., Huang, M.X., Luo, P.Y., Bian, Z.H., Xu, Y.F., Wang, C.B., Yang, S.H.,
Li, L., Selmi, C., et al. (2021). Characterization of Organ-Specific Regulatory B Cells Using
Single-Cell RNA Sequencing. Frontiers in immunology *12*, 711980.
10.3389/fimmu.2021.711980.
Zhang, X. (2013). Regulatory functions of innate-like B cells. Cell Mol Immunol *10*, 113-121.
10.1038/cmi.2012.63.

1215

1216

## Figure legends

**Figure 1. Th17 cells induce formation of large ectopic lymphoid follicle-like structures expressing germinal center markers in association with the meninges of spinal cord and brain.** (**A**) Experimental outline. Naïve MOG-specific T cells isolated from 2D2 mice are differentiated with polarizing cytokines into Th1 or Th17 cells in the presence of anti-CD3 and APCs. After 5-8 days, the cells are restimulated with anti-CD3 and anti-CD28 for 2 days before they are transferred i.v. or i.p. into recipient animals. Clinical disease develops after 8-15 days post transfer. (**B-D**) At the peak of disease, cells were isolated from the CNS of Th1 and Th17-EAE mice and analyzed for the presence of CD19$^+$ B cells. (**B**) Representative flow cytometry plots (pre-gated on CD45$^{high}$ CD11b$^{low}$ cells) and quantification of the (**C**) frequency of CD19$^+$ B cells in CD45$^{high}$ CD11b$^{low}$ cells and (**D**) absolute numbers of infiltrating CD19$^+$ B cells in the CNS. Graphs show cumulative data from three (Th1-EAE) or eight (Th17-EAE) independent experiments. Mann-Whitney U test. *p < 0.05; **p < 0.01. Horizontal lines indicate means. Dots represent individual mice. (**E, F**) At the peak of disease, (**E**) longitudinal spinal cord and (**F**) coronal brain cryosections of Th17-EAE mice were prepared and stained for T cells (CD3, red), B cells (B220, green) and laminin (blue). (**E**) Zoom of red highlighted area in Figure S2B. (**F**) Cortex. Scale bars: 200 µm. (**G**) Scheme shows the spatial distribution of the largest ectopic lymphoid aggregates (red stars) in the spinal cord of the five mice analyzed. *V, ventral; D, dorsal.* (**H-K**) CNS cryosections of Th17-EAE mice at the peak of disease were stained for (**H**) T cells (CD3, red), B cells (B220, green) and the proliferation marker Ki67 (blue) (white arrows indicate Ki67$^+$ T and B cells); (**I**) B cells (B220, magenta) and the germinal center marker peanut agglutinin (PNA, green); (**J**, **K**) T cells (CD3, red), B cells (B220, green) and the enzyme activation-induced cytidine deaminase (AID, magenta). For better visualization, the AID staining is shown separately in K. Scale bars: 50 µm.

**Figure 2. CNS-infiltrating B cells are phenotypically different from peripheral B cells.** At the peak of disease, B cells were sorted from indicated organs of Th17 recipients (n = 3 mice) based on their expression of CD19, and their gene expression was characterized by single-cell RNA sequencing. (**A**) Single cell transcriptomes are depicted in UMAP plots representing ten color-coded cell clusters, shown separately for CNS, cervical lymph node (cLN) and spleen. (**B**) For each cluster, the fraction of cells belonging to CNS, cLN or spleen was calculated. (**C, D**) Violin plots showing expression levels of gene sets associated with (**C**) light zone and (**D**) dark zone germinal center (GC) B cells. (**E, F**) Violin plots showing expression levels of gene sets associated with (**E**) B1a B cells and (**F**) MZ B cells. (**G**, **H**) Feature plots depicting distribution of (**G**) Cd9 or (**H**) Fcrl5-expressing B cells among B cell clusters. Two-sided *t*-test with FDR adjusted p-values. *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001. Significantly downregulated gene sets are marked in grey.

**Figure 3. CNS-infiltrating B cells undergo clonal expansion in the CNS.** (**A**) At the peak of disease, B cells were sorted from indicated organs of Th17 recipients (n = 3 mice) based on their expression of CD19, and characterized by BCR repertoire analysis. Feature plots depicting distribution of nonexpanded (grey) and expanded (colored) B cells among B cell clusters, plotted separately for CNS, cLN and spleen. The color code indicates the number of cells per clone. (**B**) Fraction of expanded cells belonging to CNS, spleen and cLN. (**C**) Percentage of clones with expansion factors ≥3 for each cluster in the CNS. (**D**) Violin plots showing expression levels of genes associated with plasmablasts (PB). Two-sided *t*-test with FDR adjusted p-values. *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001.

**Figure 4. CNS-infiltrating B cells show elevated expression of genes associated with cytokine signaling, B cell activation and antigen presentation/immune synapse.** (**A**, **B**, **D**, **G**) Violin plots showing expression levels of gene signatures induced by (**A**) IFNγ, (**B**) IL-21 and (**D**) TNFα, and (**G**) a gene set associated with antigen presentation/immune synapse. Two-sided *t*-test with FDR adjusted p-values. *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001. Significantly downregulated gene sets are marked in grey. (**C**, **E**, **F**, **H-K**) At the peak of disease, cells were isolated from CNS or spleen and B cells were analyzed for expression of (**C**) IL-21R, (**E**) CD69, (**F**) CD83, (**H**) MHC-II, (**I**) CD40, (**J**) CD86 and (**K**) ICAM-1 (pre-gated on CD19$^+$ cells) by flow cytometry. Graphs show the geometric mean fluorescence intensity (gMFI) of (**C**), IL-21R, (**H**) MHC-II, (**I**) CD40 or (**K**) ICAM-1 and the frequency of (**E**) CD69$^+$, (**F**) CD83$^+$ or (**J**) CD86$^+$ cells. Paired *t*-test. *p < 0.05; **p < 0.01; ***p < 0.001, ****p < 0.0001. Dots represent individual mice.

**Figure 5. 2D2 T cells form long-lasting contacts with B cells compared to OT-II T cells.** Intravital microscopy of Twitch-2B$^+$ T cells and tdTomato$^+$ B cells in spinal cord eLFs of Th17-EAE mice. (**A**) Experimental outline. MOG-specific T cells from 2D2 mice are retrovirally transduced with the Twitch-2B construct during their differentiation into Th17 cells. Upon adoptive transfer in Mb1.Cre x ROSA26 tdTomato mice, intravital two-photon microscopy of spinal cord eLFs in EAE mice can be performed to visualize interactions of Th17 cells (green) and B cells (red), as well as subsequent activation events of Th17 cells. (**B**) Dorsal view onto eLF containing Twitch-2B$^+$ T cells (green) and tdTomato$^+$ B cells (red). Scale bar: 100μm. (**C**) Exemplary images of T:B cell contacts. (**D**) Proportion of 2D2 (n = 232 cells from six independent experiments) and OT-II T cells (n = 107 cells from four independent experiments) forming contact with B cells. (**E**) Quantification of T:B cell contact duration. For both experimental groups, videos of 15-50 min length were acquired. Dots represent individual contacts. Horizontal lines indicate means. Mann-Whitney U test. ****p < 0.0001. (**F**) Proportion of long (> 5 min) and short (< 5 min) T:B cell contacts. (**G**) Quantification of the velocity of 2D2 T cells without B cell contact (n = 10938 velocities), 2D2 T cells with B cell contact (n = 8314 velocities) and OT-II T cells (n = 9413 velocities). One-way ANOVA with Tukey post-test. ****p < 0.0001. Horizontal lines indicate means. Graph shows cumulative data from (2D2) six or (OT-II) four independent experiments.

**Figure 6. 2D2 T cells, but not OT-II T cells, become reactivated during B cell contact.** Intravital microscopy of Twitch-2B$^+$ T cells and tdTomato$^+$ B cells in spinal cord eLFs of Th17-EAE mice. (**A**) Single time point of a representative video. (Left) Fluorescence overlay of Twitch-2B$^+$ T cells (green) and tdTomato$^+$ B cells (red) and (right) pseudocolor calcium ratio image showing calcium levels of T cells. Scale bar: 50 μm. (**B**) Quantification of the calcium indicator ratio (YFP/CFP) of 2D2 T cells without B cell contact (n = 4988 ratios), 2D2 T cells with B cell contact (n = 3653 ratios) and OT-II T cells (n = 9531 ratios). One-way ANOVA with Tukey post-test. ****p < 0.0001. The black dashed line indicates the ratio threshold (0.534). Horizontal lines indicate means. The graph shows cumulative data from (2D2) three or (OT-II) four independent experiments. (**C**) Proportion of 2D2 and OT-II T cells exhibiting calcium signaling. (**D**) Proportion of T cells with short-term (< 2 min), long-term (> 2 min) or no calcium spikes. (**C**, **D**) Cumulative data of 2D2 (n = 104 cells from three independent experiments) and OT-II T cells (n = 107 cells from four independent experiments). (**E, F**) Representative track of (**E**) 2D2 T cells (n = 104 cells from three independent experiments) and (**F**) OT-II T cells (n = 107 cells from four independent experiments) showing calcium

indicator ratio (YFP/CFP, red) and velocity (black) in each time frame. The red dashed line
indicates the ratio threshold (0.534). Grey shading highlights time frames with B cell contact.
(**G**) Exemplary microscopic images of (left and middle panel) two representative 2D2 and
(right panel) a representative OT-II T cell. (Left) Fluorescence overlay of Twitch-2B$^+$ T cells
(green) and tdTomato$^+$ B cells (red) and (right) pseudocolor calcium ratio image showing
calcium levels of T cells. White arrow indicates cell of interest. (Left panel) 2D2 T cell has B
cell contact at all indicated time points. (Middle panel) 0 min: before B cell contact; 1.8 min –
26.2 min: with B cell contact. (Right panel) 0 min: before B cell contact; 1.4 min – 14.8 min:
with B cell contact.
1322
**Figure 7. T cells depend on B cells to maintain a highly pro-inflammatory cytokine**
**profile in the CNS.** Comparison of T cell responses in Th17-mediated EAE in B cell-deficient
Mb1-KO mice and WT mice. At the peak of disease, cells were isolated from (**A**) CNS or (**B**)
spleen and T cells were analyzed for IL-17A and IFNγ expression by intracellular cytokine
staining. Transferred T cells were identified by their expression of the transgenic MOG-
specific Vα3.2 TCR. Representative flow cytometry plots and quantification of IL-17A$^+$, IFNγ$^+$
and IL-17A$^+$IFNγ$^+$ producers of (upper panel) transferred (Vα3.2$^+$) and (lower panel)
endogenous (Vα3.2$^-$) T cells. Graphs show cumulative data from two independent
experiments. (**A**, upper panel) unpaired *t*-test (left), Welch`s t-test (middle), Mann-Whitney U
test (right). (**A**, lower panel) Mann-Whitney U test (left and right), Welch`s t-test (middle). (**B**,
upper panel) unpaired *t*-test (left), Mann-Whitney U test (middle), Welch`s t-test (right). (**B**,
lower panel) unpaired *t*-test (left and right), Welch`s t-test (middle). *p < 0.05; **p < 0.01; ***p
< 0.001. Mean ± SEM. Dots represent individual mice.

Figure 1

Figure 2

Figure 3

Figure 5

**A**

T cells
B cells

Low ⬛ Ca²⁺ High

**B**

ratio [YFP/CFP]

47.2 %  75.0 %  5.0 %

****
****
****

without B cell contact    with B cell contact    OT-II T cells

2D2 T cells

**C**

Proportion of T cells (%)

■ with calcium spikes
■ without calcium spikes

2D2 T cells    OT-II T cells

**D**

Proportion of T cells (%)

■ 2D2 T cells
■ OT-II T cells

Long-term    Short-term    no calcium spikes

**E**

2D2 T cells

ratio [YFP/CFP]    velocity [µm/min]

time (min)

**F**

OT-II T cells

ratio [YFP/CFP]    velocity [µm/min]

time (min)

**G**

2D2 T cell    OT-II T cell

0 min    0 min    0 min

5.0 min    1.8 min    1.4 min

11.1 min    6.8 min    4.9 min

16.1 min    11.5 min    8.5 min

30.5 min    26.2 min    14.8 min

Low ⬛ Ca²⁺ High

T cells
B cells

Figure 6

Figure 7