Inhibition of BCR::ABL1 tyrosine kinase activity Aids in the Generation of Stable Chronic Myeloid Leukemia Induced Pluripotent Stem Cells

Esther Sathya Bama Benjamin[1, 2], Dinesh Babu[3*], Gaurav Joshi[1,2*], Bharathi M Rajamani[1], Krittika Nandy[3], Sonam Rani[3], Senthamizhselvi Anandan[1], Chitra Premkumar[3], Madhavi Maddali[1], Aby Abraham[1], Shaji R Velayudhan[1,3 #], Poonkuzhali Balasubramanian[1#]

[1] Department of Haematology, Christian Medical College- Vellore
[2] Sree Chitra Tirunal Institute for Medical Sciences & Technology (SCTIMST), Thiruvananthapuram
[3] Centre for Stem Cell Research (a unit of INSTEM, Bengaluru), Christian Medical College, Vellore
Corresponding authors
* Equal contribution

Address for correspondence
Dr. Shaji R Velayudhan
rvshaji@cmcvellore.ac.in
and
Dr. Poonkuzhali Balasubramanian
bpoonkuzhali@cmcvellore.ac.in

Department of Haematology Christian Medical College Vellore-632004, and
Centre for Stem Cell Research (a unit of INSTEM, Bengaluru)

Acknowledgments:
We acknowledge the support of the core imaging and flow cytometry facilities at the Centre for Stem Cell Research (a unit of inStem, Bengaluru, Christian Medical College Campus, Bagayam, Vellore). We thank Ms. Sangeetha for her help in collecting and processing primary CML samples. We thank our clinical colleagues in the Department of Haematology for managing the patients with CML and providing the data on response to therapy.

Funding:
This study was supported by the Department of Biotechnology, Government of India, Centre of Excellence grant [BT/COE/34/SP13432/2015] to Poonkuzhali Balasubramanian; the Department of Science and Technology Core Research Grant [CRG/2022/006590] and [BT/PR17316/MED/31/326/2015] to RVS; the institutional FLUID research grant to BMR. SRV and PB are supported by the DBT/Wellcome Trust India Alliance [IA/S/17/1/503118; IA/S/15/1/501842]. ESB is supported by the Department of Science and Technology INSPIRE fellowship; GJ by the University Grants Commission; and BMR by the Indian Council of Medical Research.

Abstract

Induced pluripotent stem cells (iPSCs) generated from patients with chronic

myeloid leukemia (CML) have the potential for disease modeling to study disease

pathogenesis and screening therapeutic interventions. In this study, we aimed to

generate iPSCs from $CD34^+$ hematopoietic progenitors of CML patients with varying

responses to tyrosine kinase inhibitor (TKI) therapy. The generated CML-CD34-iPSC

colonies displayed atypical "dome-shaped" morphology and underwent spontaneous

differentiation in a few days. However, supplementation with imatinib (IM), the most

widely used TKI to treat CML patients, in the culture medium improved the stability

and maintenance of all isolated CML-CD34-iPSC colonies, allowing them to be

maintained for more than 20 passages without significant differentiation. In contrast to

previous studies, our results indicate that suppressing the BCR::ABL1 oncogenic

pathway is essential for efficiently generating stable CML-iPSC colonies. Furthermore,

we successfully differentiated these iPSCs to $CD34^+$ hematopoietic progenitors both in

the presence and absence of IM. This robust protocol for generating CML-iPSCs

provides a valuable resource for disease modelling. The generated iPSCs will be a

valuable tool for investigating CML pathophysiology, drug resistance mechanisms, and

drug screening to identify novel and effective therapies for this disease.

Introduction

Chronic myeloid leukemia (CML) is a myeloproliferative neoplasm that arises

from hematopoietic stem cells (HSCs) transformed by BCR::ABL1 oncogenic fusion

protein with constitutive serine/threonine kinase activity expressed in these cells.

BCR::ABL1   triggers   various   downstream   signaling   pathways,   including

RAS/RAF/MEK/ERK, JAK2/STAT, and PI3K/AKT/mTOR pathways, leading to the

accumulation of granulocytes at various stages of myeloid maturation (1,2). Although

targeted therapy with tyrosine kinase inhibitors (TKIs), such as imatinib (IM), dasatinib

(DA), and nilotinib (NIL), has dramatically improved the long-term survival of CML

patients, only around 50% of patients achieve treatment-free remission (TFR) (3). The

persistence of leukemic stem cells (LSCs) is one of the primary causes of disease

recurrence (4). Due to the rarity of the LSC population, it is challenging to obtain

sufficient cells for research to understand the mechanisms of TKI resistance of these
cells and screen drugs to eradicate them (4,5).

Induced pluripotent stem cells (iPSCs) derived from patients with genetic
diseases or cells with genetic alterations can be differentiated into scalable quantities
of disease-relevant differentiated cells. Thus, they are a powerful tool for studying
disease pathogenesis and drug screening (5). iPSCs generated from patients with
leukemias help understand how oncogenes and patient-specific chromosomal
abnormalities influence the development of leukemia-like phenotypes *in-vitro* (6).
CML-iPSCs generated from the same patient at various stages of the disease and from
patients with different responses to TKI therapy are valuable resources for
investigating molecular and epigenetic mechanisms of CML progression and drug
resistance, leading to the identification of novel therapeutic targets and drug
candidates for the treatment of CML (7,8). Moreover, iPSCs can be efficiently
genetically manipulated by gene editing methods to identify specific genes involved in
disease pathogenesis (9).

Previous studies have demonstrated the successful generation of CML-iPSCs
from peripheral blood mononuclear cells (10) and CD34$^+$ hematopoietic stem and
progenitor cells (HSPCs) isolated from CML patients (7,11–13). CML-iPSCs exhibit
resistance to IM like CML-LSCs, so they are considered suitable for investigating the
cellular mechanisms underlying IM resistance (12,14,15). CML-iPSCs could be
differentiated into hematopoietic progenitors that recapitulated some of the
pathophysiologic features of the disease (7,10,12–14). However, it remains unclear
how *BCR::ABL1* expression and TKI treatment impact the efficiency and quality of iPSC
generation from CML cells.

In this study, we describe the generation of iPSCs from the CD34$^+$ cells of CML
patients who exhibited varying responses to TKI therapy for disease modeling and
investigating the effect of BCR::ABL1 oncoprotein expression on the maintenance of
CML-iPSCs. In contrast to previous studies, we found that the expression and activity
of BCR::ABL1 affect the maintenance of CML-CD34-iPSCs, and it was essential to
suppress the TKI activity of BCR::ABL1 fusion protein by TKI supplementation for their
survival and maintenance. The CML-iPSCs generated using our protocol could be
differentiated into BCR::ABL1 positive hematopoietic progenitors. The patient-specific

CML-iPSCs and the differentiated hematopoietic cells generated by this method have

a wide range of applications, including high throughput CRISPR-Cas9 screening, to

identify novel pathways involved in CML pathogenesis and drug resistance.

Materials and methods

Ethical statement: This study was approved by the institutional review board and the

institutional bio-safety committee of Christian Medical College, Vellore.

Isolation and expansion of CD34$^+$ cells: Peripheral blood samples were obtained

from a healthy donor (N=1), who underwent mobilization of CD34$^+$ cells for

hematopoietic transplantation, and newly diagnosed chronic phase-CML (CP-CML)

patients (N=4) after obtaining written informed consents. Peripheral blood

mononuclear cells (PBMNCs) were isolated using Ficol density gradient centrifugation.

The CD34$^+$ cells were purified from PBMNCs by the magnetic separation method

(EasySep, Stem Cell Technologies) and were cryopreserved. The cryopreserved

CD34$^+$ cells of the CML patients and the normal donor were thawed and cultured in

CD34$^+$ cell expansion medium constituting StemSpan™ SFEM II medium (Stem Cell

Technologies) supplemented with CD34$^+$ Expansion Supplement and UM729 (Stem

Cell Technologies). The cells were cultured for 6-7 days with half medium change

every alternative day to assess the cell proliferation.

Reprogramming of CML and normal CD34$^+$ cells: The CD34$^+$ cells were cultured in

the expansion medium for 3-days and then nucleofected with episomal

reprogramming plasmids (pCXLE-hOCT3/4-shp53 (Addgene: 27077), pCXLE-hSK

(Addgene 27078) and pCXLE-hUL (Addgene: 27080) (kind gift from Shinya

Yamanaka), as described previously (16). Electroporation was carried out using Neon

Transfection System (ThermoFisher) at 1600V, 10ms, and 3 pulses. After five days of

culture, the nucleofected cells were transferred to Matrigel-coated plates containing

Essential-8 (E8) medium (ThermoFisher Scientific). Individual colonies were hand-

picked between day 21 and day 32 and were passaged and maintained in mTeSR Plus

medium (StemCell Technologies) supplemented with 10µM IM (Sigma Aldrich) to

generate stable iPSC clones.

Western Blot: The iPSCs were dissociated using 1mM EDTA (ThermoFisher

Scientific), and the protein lysates were prepared in radioimmunoprecipitation assay

buffer supplemented with a protease inhibitor mixture (Roche Applied Science, IN, USA) and 2mM phenylmethylsulfonyl fluoride (Sigma-Aldrich). About 30µg of whole cell lysate was loaded in 10% SDS-polyacrylamide gel, and the proteins were transferred to a polyvinylidene difluoride membrane. The membrane was blocked with 10% non-fat dry milk powder in tris-buffered saline containing 0.1% Tween 20 (TBST) for two hours and then incubated with an anti-phosphoCRKL antibody (Cell Signaling Technology) and Anti β –Actin antibody (ThermoFisher Scientific). The proteins were detected using the ECL chemiluminescence kit (Thermo Scientific, Pierce), and the images were captured using the FluorChem E system (Protein Simple).

Analysis of *BCR::ABL1* fusion transcripts: RNA was extracted from patients' PBMNCs and iPSCs using TRI Reagent (Sigma-Aldrich). cDNAs were prepared from 2 µg RNA with random hexamers using the High-Capacity cDNA Synthesis Kit (Thermo Scientific), followed by RT-PCR to identify the *BCR::ABL1* fusion transcripts using a previously described protocol (17). The *BCR::ABL1* mRNA copies were assessed using an in-house ddPCR assay (Datari et. al., manuscript under preparation). iPSC colonies were screened for BCR::ABL1 kinase domain mutations using a previously described protocol (18).

Alkaline phosphatase staining: Alkaline Phosphatase staining was performed using the Alkaline Phosphatase Detection Kit (Sigma-Aldrich) following the manufacturer's instructions.

Flow cytometry: To analyze the expression of TRA-1-60 in iPSCs, the colonies were dissociated using TrypLE (Thermo Fisher Scientific) and then stained using a PE-conjugated TRA-1-60 antibody (Thermo Fisher Scientific) for 20 mins. The cells were washed with mTeSR Plus (Stem Cell Technologies) containing Revitacell (Thermo Fisher Scientific), and the expression was analyzed using Navios (Beckman Coulter) or FACS Aria (BD Biosciences) flow cytometers. To measure the purity of CD34+ cells after their magnetic separation from PBMNCs, the cells were stained with an APC-conjugated anti-CD34 antibody (BioLegend).

Immunofluorescence: Immunofluorescence for pluripotency markers, such as SOX2, TRA-1-81, OCT4A, SSEA-4, NANOG and TRA-1-60, was performed as previously described (19).

Trilineage differentiation: Trilineage differentiation of iPSCs to ectoderm, endoderm and mesoderm lineages was carried out using STEMdiff tri-lineage differentiation Kit (Stem Cell Technologies) as per the manufacturer's protocols.

Hematopoietic differentiation of iPSCs: For the hematopoietic differentiation of iPSCs, we used the STEMdiff™ Hematopoietic Kit (Stem Cell Technologies). Briefly, iPSCs were either single-cell sorted or seeded as small aggregates in mTeSR Plus (Stem Cell Technologies). Subsequently, the culture medium was changed to STEMdiff™ A medium, with or without IM. After three days, 70,000 cells were reseeded in STEMdiff™ A medium and maintained for another three days. Following this, the medium was switched to STEMdiff™ B. After 12 days of culture, the cells present in suspension were harvested and analyzed by flow cytometry for the expression of hematopoietic surface markers.

Colony-Forming Assay: In each experiment, 7,000 hematopoietic progenitors generated from iPSCs were plated onto MethoCult GF H4636 medium (Stem Cell Technologies). The number of colonies in each experiment was scored on day 14 following the manufacturer's instructions.

Results

Generation of iPSCs from normal and CML CD34+ cells:

As CD34+ cells of CML patients are known to constitute the leukemia stem/progenitor cells (LSPCs) that are associated with disease recurrence (4,20), we aimed to reprogram CML-CD34+ cells from two patients who achieved complete cytogenetic response and one who achieved suboptimal response (>1% *BCR::ABL1* transcripts detected in peripheral blood cells after 12months) after TKI therapy (**Suppl Table 1**) and a healthy volunteer as described (**Figure 1A**). CD34+ cells purified from CML patients and those from a healthy donor were cultured in CD34+ expansion medium. The number of CD34+ cells increased by 15 to 20-fold (Mean: 20.2±:3.34) without reducing the percentage of CD34+ cells (**Figure S1A** and **S1B**).

We nucleofected proliferating CD34+ cells from day 3 of the culture with episomal reprogramming plasmids (13). The normal CD34+ cells gave rise to iPSC colonies with typical morphology between day 14 and day 18 (**Figure 1B**). These

colonies were hand-picked between day 20 and day 24 to generate stable iPSC

clones. The normal iPSC clones maintained typical iPSC morphology without

differentiation for 12 passages and expressed high levels of pluripotency markers

(**Figures 1C** and **1D**). In contrast, CD34$^+$ cells from TKI-responsive CML patients did

not form typical iPSCs but formed several "dome-shaped" colonies after 30 days

(**Figures 1B and C**). These colonies were surrounded by adherent cells (**Figure 1C)**

that resembled those formed by the spontaneous differentiation of iPSCs. We isolated

11 "dome-shaped" colonies from 3 patients and cultured them for 4 to 5 passages.

They also expressed TRA-1-60 in the initial passages (**Figure 1D**). During subsequent

passages, the "dome-shaped" colonies peeled off from the bottom of the plates and

floated in the medium, leaving the differentiated cells attached to the plate.

IM supplementation aids in the maintenance, stability, and pluripotency of CML

iPSCs

Previous studies showed that CML-iPSCs express *BCR::ABL1* transcripts

present in the donor cells used for reprogramming (10–12,14). In our study, the

"dome-shaped" CML-CD34-iPSC colonies also expressed the same types of

*BCR::ABL*1 fusion transcripts (e13a2 and e14a2) as those present in the respective

patients' peripheral blood cells used for reprogramming (**Figure S2A** and **S2B**).

Additionally, karyotyping confirmed the presence of t(9;22) in all iPSC clones (**Figure**

S2C), and none of the clones had mutations in the *BCR::ABL1* kinase domain.

As it has been reported that TKI treatment enhances pluripotency marker

expression in BCR::ABL1+ iPSC clones (15), we hypothesized that the atypical

morphology and high rate of spontaneous differentiation observed in our CML-CD34-

iPSC colonies were due to the constitutive expression of BCR::ABL1 fusion protein

and its tyrosine kinase (TK) activity. To inhibit the TK activity of the BCR::ABL1 fusion

protein, we supplemented the medium with 10μM IM and monitored the colonies for

changes in their morphology characteristics. We observed that within two days of IM

supplementation, the colonies gained the typical iPSC morphology (**Figure 2A**) with

an increase in the TRA-1-60+ cell population (**Figures 2B** and **2C**). All the clones were

positive for pluripotency marker expression (**Figure 2D**). They also could be

differentiated into three germ layers, ectoderm, endoderm, and mesoderm (**Figure**

S3). When IM was withdrawn, the IM-supplemented CML-CD34-iPSC colonies

reverted to the "dome-shaped" morphology within 2-4 days (**Figure 3A**), with a

reduction in cell viability (**Figure 3B**) and TRA-1-60 expression (**Figure 3C**). In

contrast, when the control iPSCs were treated with IM, there was a reduction in the

colony number, suggesting that IM supplementation specifically enhances the

maintenance of CML-CD34-iPSCs (**Figure S4**).

CRK-like proto-oncogene (CRKL) is a downstream target of BCR::ABL1 and is

phosphorylated by BCR::ABL1 fusion protein (20,21). As p-CRKL levels can be used

to determine the status of BCR::ABL1 kinase activity, the expression of p-CRKL was

analyzed in CML-CD34-iPSCs before and after the IM treatment (**Figure 3D**). It was

found that p-CRKL expression was decreased in CML-CD34-iPSCs after IM treatment,

suggesting that suppression of BCR::ABL1 mediated tyrosine phosphorylation resulted

in the stable maintenance of CML-CD34-iPSCs.

The effect of IM in reprogramming of CML-CD34+ cells

As IM supplementation inhibits BCR::ABL1 kinase activity and enhances the

stability and pluripotency of CML-iPSCs, we investigated the effect of IM on the

reprogramming of CML-CD34$^+$ cells. After nucleofection of CD34+ cells from two IM-

responsive patients, the cells were cultured in the reprogramming medium

supplemented with 10µM IM (**Figure 4A**). Reprogramming without IM supplementation

generated "dome-shaped" colonies after 3 weeks, and they gained typical iPSC

morphology after IM supplementation with increased TRA-1-60 expression (**Figure**

4B). However, continuous IM supplementation throughout reprogramming did not

generate "dome-shaped" colonies or colonies with typical morphology. These findings

suggest that while IM enhances the stability of CML-iPSCs by inhibiting *BCR::ABL1*

kinase activity, it impedes the reprogramming of CML-CD34$^+$ cells. Overall, our results

demonstrate that IM supplementation transforms the "dome-shaped" colonies formed

during the reprogramming of CML-CD34$^+$ cells into iPSCs with typical flat morphology.

Furthermore, IM improves the pluripotency of CML-CD34-iPSCs by inhibiting the

BCR::ABL1-mediated TK activity.

Reprogramming of CD34$^+$ cells from an IM refractory CML patient

The HPCs derived from the iPSCs generated from IM refractory CML patients

offer a valuable tool to investigate the mechanisms underlying IM therapy resistance

and develop a more effective treatment for those who do not respond well to IM.

Therefore, we reprogrammed CD34$^+$ cells from an IM refractory patient, with and

without IM supplementation, during the reprograming process (**Figure 4A**). In

contrast to our observation in the IM responders' cells, we observed an early

emergence of iPSC colonies from the IM-refractory patient's cells (~3 weeks Vs. ~4

weeks). However, reprogramming of the cells from the IM refractory patient-generated

more colonies with continuous IM supplementation than without IM (**Figure 4C**). We

isolated 9 colonies, 8 of which exhibited "dome-shaped" morphology (**Figure 1C**),

while 1 showed typical iPSC morphology (**Figure S5A**). Karyotyping analysis of the

"dome-shaped" colonies generated from the IM refractory patient demonstrated the

presence of t(9;22), whereas the one clone with the typical iPSC morphology showed

a normal karyotype (**Figure S5A** and **S5B**). These results suggest that the "dome-

shaped" morphology of the CML-CD34-iPSCs is because of BCR::ABL1 expression in

these cells.

Second and third generation TKIs also aid in the maintenance of CML iPSCs.

Second generation TKIs, DA and NIL, and third generation TKI ponatinib were

approved for patients who develop resistance to or are intolerant to first or second-

generation TKIs, respectively. These TKIs target specific mutations in the BCR::ABL1

fusion protein, providing more effective and durable responses in CML patients. We

explored the effects of second and third generation TKIs, DA, NIL, and ponatinib, in

the maintenance of CML-CD34-iPSCs. We replaced IM with different concentrations

of DA, NIL, and ponatinib and evaluated their effects on CML-CD34-iPSCs. CML-

CD34-iPSCs treated with these TKIs generated stable the CML-CD34-iPSCs with 2nM

DA, 40nM NIL, and 2nM ponatinib, while at lower concentrations, resulted in "dome-

shaped" colonies (**Figure 5**). Together, our results suggest that the inhibition of

BCR::ABL1 TK activity promotes the pluripotency of CML-iPSCs.

Hematopoietic differentiation of CML iPSCs

Previous studies have demonstrated that CML-iPSCs can be successfully

differentiated into hematopoietic progenitor cells (HPCs), despite the expression of

BCR::ABL1 oncoprotein and its TK activity in these cells (7,14,22). However, the

efficiency of hematopoietic differentiation of CML-iPSCs has been reported to be lower

than normal iPSCs (7,10), and the resulting HPCs exhibited a partial recovery of TKI

sensitivity (13). Considering our finding that IM supplementation is necessary for the

maintenance of CML-CD34-iPSCs, we investigated whether IM supplementation is

also required for the differentiation of CML-iPSCs into HPCs.

To address this question, we initially performed single-cell sorting to obtain

iPSC colonies with an equal number of cells, which were subsequently subjected to

hematopoietic differentiation (**Figures 6A and B**). We incorporated IM

supplementation at different stages of the differentiation process (**Figure 6B**). We

differentiated CML-CD34-iPSCs using two different concentrations of IM (1µM and

2µM) as well as without any IM supplementation. We could obtain suspension cells

constituting of HPCs both in the presence and absence of IM (**Figure S6**). Interestingly,

CML-CD34-iPSCs differentiated without IM before and after mesoderm induction (E1)

(**Figure 6B**) showed a 5-fold increase in the yield of CD34$^+$CD45$^+$ HPCs and the total

number of suspension cells compared to those differentiated in the presence of IM

throughout the entire differentiation (E3 and E5) (**Figure 6D**). Flow cytometry analysis

of hematopoietic markers revealed that approximately 41% of the suspension cells

were CD34$^+$CD45$^+$ HPCs (**Figure 6C**). Utilizing single-cell sorting of iPSCs, we were

able to generate a sufficient number of HPCs for a drug screening experiment.

Based on the results of this experiment, we seeded iPSCs as small aggregates

and then subjected differentiation with and without IM supplementation after

mesoderm induction (**Figure 7A**). Notably, the CML-CD34-iPSCs differentiated

without IM yielded approximately 30 times more HPCs (**Figure 7C**) compared to

differentiation with IM supplementation. The decline in HPC differentiation observed in

the presence of IM can be attributed to the sensitivity of CML-iPSC-HPCs to IM. Flow

cytometry analysis revealed that around 50% of the suspension cells constituted the

CD34$^+$CD45$^+$ population of HSPCs (**Figure 7B**). Furthermore, hematopoietic colony

formation showed that the HPCs generated without IM had more myeloid progenitors

compared to those derived from a normal iPSC clone (**Figure 7D**). To further validate

these findings, we differentiated CML-CD34-iPSCs derived from two additional patients **(Figure 7E)**. Flow cytometry analysis showed 40% to 70% CD34$^+$CD45$^+$ cells in the absence of IM from these two CML-CD34-iPSCs (**Figure 7G**). Consistently, it was observed that differentiation in the absence of IM resulted in more HPCs compared to the other experimental groups (**Figure 7F**).

Discussion

Eliminating CML-LSCs is a critical objective in the treatment of CML, as it can prevent disease recurrence, extend survival, improve quality of life, and enable curative treatments (3,4,23). However, the rarity of these cells poses a challenge in studying the molecular mechanisms that contribute to their survival. Generating iPSCs from CML CD34$^+$ cells can help overcome this hurdle. These cells can be expanded and redifferentiated into hematopoietic cells, providing a continuous supply of cells for extensive proteome, epigenome, transcriptome, and drug screening studies.

Several groups have generated CML iPSCs from PBMNCs (10) and CD34$^+$ cells (7,11,12,14) using lentiviral (14,15), retroviral (12,13), Sendai (11), or episomal vectors (7,10). However, these studies did not address the atypical morphology and low reprogramming efficiency of CML iPSCs. Additionally, the effect of BCR::ABL expression on the morphology and stabilization of CML-iPSC colonies (9) was not investigated.

We observed that the reprogramming efficiency of CML CD34$^+$ was approximately 50 times lower than the normal CD34$^+$ cells, as previously reported (7,11). One possible explanation for this low efficiency could be the presence of the BCR::ABL1 oncogene, which may disrupt the reprogramming process of CML CD34$^+$ cells (24). Furthermore, CMLCD34$^+$ cells formed colonies later than normal CD34$^+$ cells, and they exhibited atypical colony morphology (14). Bedel et. al. reported that CML iPSCs had a distinct morphology resembling that of cell aggregates similar to mouse embryonic stem cells (12,14).

The instability and differentiation-prone nature of CML-CD34-iPSCs may be due to the presence of *BCR::ABL1* oncogene, which can interfere with the reprogramming factors required to generate stable iPSC colonies. Ito *et al*. demonstrated that the use

of molecularly targeted drugs such as IM, which can suppress the activity of the BCR-ABL oncogene, can improve the transcriptional regulation of reprogramming factors and enhance pluripotency marker expression (24). This study also demonstrated that the expression of reprogramming factors in the BCR::ABL+ K562 cells increased pluripotency marker expression after IM supplementation, confirming that oncogene suppression improves the reprogramming of cancer cells. Our study confirmed these findings and showed that IM supplementation was necessary to generate stable CML iPSCs without spontaneous differentiation. IM supplementation improved the stability and maintenance of all isolated CML-CD34-iPSC colonies, allowing them to be cultured for over 20 passages without significant differentiation. The CML-CD34-iPSCs cultured with IM showed increased proliferation, while IM withdrawal decreased the proliferation and TRA-1-60 expression. These results were consistent with a previous study that reported faster growth of CML-iPSCs in the presence of TKI (14,15).

The phosphorylation of CRKL is decreased in IM-treated primary CD34$^+$ HPCs isolated from CML patients (25) and CML iPSCs (**Figure 3D**), suggesting that the BCR-ABL TK activity in hematopoietic cells and CML iPSCs may involve similar pathways. The fact that IM could stabilize the pluripotency of CML-iPSCs uncovers a novel mechanism of pluripotency maintenance. Both CML-iPSCs and LSCs express BCR-ABL fusion protein but are IM resistant. Therefore, CML-iPSCs created using our strategy and the HPCs obtained from these iPSCs can be used for CRISPR-Cas9 screening to study the mechanisms of TKI resistance exhibited by these cells and discover drugs that can effectively eliminate CML-LSCs.

CML-iPSCs generated in the presence of IM could be successfully differentiated into HPCs. The iPSC-derived HPCs obtained from multiple patients will help investigate the survival mechanisms of BCR-ABL+ hematopoietic cells in TKI-resistant patients.

Reference

1.   Bhatia R, Holtz M, Niu N, Gray R, Snyder DS, Sawyers CL, et al. Persistence of malignant hematopoietic progenitors in chronic myelogenous leukemia patients in complete cytogenetic remission following imatinib mesylate treatment. Blood. 2003 Jun;101(12):4701–7.

2. Imagawa J, Tanaka H, Okada M, Nakamae H, Hino M, Murai K, et al.
Discontinuation of dasatinib in patients with chronic myeloid leukaemia who
have maintained deep molecular response for longer than 1 year (DADI trial): a
multicentre phase 2 trial. Lancet Haematol. 2015 Dec;2(12):e528–35.

3. Etienne G, Guilhot J, Rea D, Rigal-Huguet F, Nicolini F, Charbonnier A, et al.
Long-term follow-up of the French Stop Imatinib (STIM1) study in patients with
chronic myeloid leukemia. J Clin Oncol. 2017 Jan 20;35(3):298–305.

4. Corbin AS, Agarwal A, Loriaux M, Cortes J, Deininger MW, Druker BJ. Human
chronic myeloid leukemia stem cells are insensitive to imatinib despite
inhibition of BCR-ABL activity. J Clin Invest [Internet]. 2011 Jan 4 [cited 2023
Jan 7];121(1):396–409. Available from: http://www.jci.org

5. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al.
Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined
Factors. Cell [Internet]. 2007 Nov 30 [cited 2023 Apr 11];131(5):861–72.
Available from: http://www.cell.com/article/S0092867407014717/fulltext

6. Papapetrou EP. Modeling Leukemia with Human Induced Pluripotent Stem
Cells. Cold Spring Harb Perspect Med [Internet]. 2019 Dec 1 [cited 2023 Apr
19];9(12). Available from: /pmc/articles/PMC6886454/

7. Miyauchi M, Koya J, Arai S, Yamazaki S, Honda A, Kataoka K, et al. ADAM8 Is
an Antigen of Tyrosine Kinase Inhibitor-Resistant Chronic Myeloid Leukemia
Cells Identified by Patient-Derived Induced Pluripotent Stem Cells. Stem Cell
Reports. 2018 Mar;10(3):1115–30.

8. Corbin AS, Agarwal A, Loriaux M, Cortes J, Deininger MW, Druker BJ. Human
chronic myeloid leukemia stem cells are insensitive to imatinib despite
inhibition of BCR-ABL activity. J Clin Invest. 2011 Jan;121(1):396–409.

9. Imeri J, Desterke C, Marcoux P, Telliam G, Sanekli S, Barreau S, et al.
Modeling Blast Crisis Using Mutagenized Chronic Myeloid Leukemia-Derived
Induced Pluripotent Stem Cells (iPSCs). Cells [Internet]. 2023 Feb 1 [cited
2023 Apr 11];12(4):598. Available from: https://www.mdpi.com/2073-
4409/12/4/598/htm

10. Hu K, Yu J, Suknuntha K, Tian S, Montgomery K, Choi KD, et al. Efficient
generation of transgene-free induced pluripotent stem cells from normal and

neoplastic bone marrow and cord blood mononuclear cells. Blood.

2011;117(14):109–19.

11. Telliam G, Féraud O, Griscelli F, Opolon P, Divers D, Bennaceur-Griscelli A, et

al. Generation of an induced pluripotent stem cell line from a patient with

chronic myeloid leukemia (CML) resistant to targeted therapies. Stem Cell Res.

2016;17(2):235–7.

12. Kumano K, Arai S, Hosoi M, Taoka K, Takayama N, Otsu M, et al. Generation of

induced pluripotent stem cells from primary chronic myelogenous leukemia

patient samples. Blood. 2012 Jun;119(26):6234–42.

13. Hosoi M, Kumano K, Taoka K, Arai S, Kataoka K, Ueda K, et al. Generation of

induced pluripotent stem cells derived from primary and secondary

myelofibrosis patient samples. Exp Hematol. 2014;42(9):816–25.

14. Bedel A, Pasquet JM, Lippert É, Taillepierre M, Lagarde V, Dabernat S, et al.

Variable Behavior of iPSCs Derived from CML Patients for Response to TKI

and Hematopoietic Differentiation. PLoS One. 2013;8(8).

15. Charaf L, Mahon FX, Lamrissi-Garcia I, Moranvillier I, Beliveau F, Cardinaud B,

et al. Effect of tyrosine kinase inhibitors on stemness in normal and chronic

myeloid leukemia cells. Leukemia. 2017;31(1):65–74.

16. Manian K V., Bharathan SP, Maddali M, Srivastava VM, Srivastava A,

Velayudhan SR. Generation of an integration-free iPSC line (CSCRi005-A) from

erythroid progenitor cells of a healthy Indian male individual. Stem Cell Res.

2018 May 1;29:148–51.

17. Hamilton A, Helgason GV, Schemionek M, Zhang B, Myssina S, Allan EK, et al.

Chronic myeloid leukemia stem cells are not dependent on Bcr-Abl kinase

activity for their survival. Blood [Internet]. 2012 Feb 9 [cited 2023 Jan

7];119(6):1501–10. Available from:

https://ashpublications.org/blood/article/119/6/1501/30195/Chronic-myeloid-

leukemia-stem-cells-are-not

18. Markose P, Chendamarai E, Balasubramanian P, Velayudhan SR, Srivastava

VM, Mathews V, et al. Spectrum of BCR-ABL kinase domain mutations in

patients with chronic myeloid leukemia from India with suspected resistance to

imatinib-mutations are rare and have different distributions.

https://doi.org/103109/10428190903332486 [Internet]. 2009 Dec [cited 2023

Apr 10];50(12):2092–5. Available from:

https://www.tandfonline.com/doi/abs/10.3109/10428190903332486

19.   Bharathan SP, Nandy K, Palani D, Janet A NB, Natarajan K, George B, et al.

Generation of an induced pluripotent stem cell line that mimics the disease

phenotypes from a patient with Fanconi anemia by conditional

complementation. Stem Cell Res. 2017 Apr;20:54–7.

20.   Hamilton A, Elrick L, Myssina S, Copland M, Jørgensen H, Melo J V., et al.

BCR-ABL activity and its response to drugs can be determined in CD34+ CML

stem cells by CrkL phosphorylation status using flow cytometry. Leuk 2006 206

[Internet]. 2006 Mar 30 [cited 2022 Dec 20];20(6):1035–9. Available from:

https://www.nature.com/articles/2404189

21.   Nichols G, Raines M, Vera J, Lacomis L, Tempst P, Golde D. Identification of

CRKL as the Constitutively Phosphorylated 39-kD Tyrosine Phosphoprotein in

Chronic Myelogenous Leukemia Cells. Blood. 1994 Nov 1;84(9):2912–8.

22.   Kumano K, Arai S, Hosoi M, Taoka K, Takayama N, Otsu M, et al. Generation of

induced pluripotent stem cells from primary chronic myelogenous leukemia

patient samples. Blood [Internet]. 2012 Jun 28 [cited 2022 Dec

28];119(26):6234–42. Available from:

https://ashpublications.org/blood/article/119/26/6234/105596/Generation-of-

induced-pluripotent-stem-cells-from

23.   Mahon FX, Etienne G. Deep molecular response in chronic myeloid leukemia:

The new goal of therapy? Clin Cancer Res [Internet]. 2014 Jan 15 [cited 2023

Mar 27];20(2):310–22. Available from:

https://aacrjournals.org/clincancerres/article/20/2/310/78489/Deep-Molecular-

Response-in-Chronic-Myeloid

24.   Ito K, Nagata K, Ohta S, Matsuda Y, Ukai T, Yasuda I, et al. The oncogene-

dependent resistance to reprogramming unveils cancer therapeutic targets.

Cell Rep. 2022 Apr;39(4):110721.

25.   Suknuntha K, Ishii Y, Tao L, Hu K, McIntosh BE, Yang D, et al. Discovery of

survival factor for primitive chronic myeloid leukemia cells using induced

pluripotent stem cells. Stem Cell Res. 2015 Nov 1;15(3):678–93.

https://cmcv-
my.sharepoint.com/personal/esther_sathya_cmcvellore_ac_in/_layouts/15/onedrive.a
spx?id=%2Fpersonal%2Frvshaji%5Fcmcvellore%5Fac%5Fin%2FDocuments%2FStu
dents%2FEsther%2FManuscript%2FFinal%20CML%2DiPSCs%20manuscript%2Fbio
rvix%2FSupplementary%20data1%2Epdf&listurl=%2Fpersonal%2Frvshaji%5Fcmcvel
lore%5Fac%5Fin%2FDocuments&parentview=1

Figure 1

A

Expansion of CD34+ cells

Nucleofection and reprogramming

Isolation and maintenance of iPSCs

Seed CD34+ cells in HEM

Nucleofection

Medium change

Seeded in Matrigel-coated plates

1:1 (E8 : HEM)

100% E8 medium

Appearance of iPSC colonies from normal CD34+ cells

Appearance of dome-shaped colonies from CML CD34+ cells

Day -3   0   2   5   7   10   14   18   20   24   31   35

B

Normal-CD34-iPSCs

CML-CD34-iPSCs

C

CML-CD34-iPSCs

G13   G15

G17   G18

Normal-CD34- iPSCs

D

Normal-CD34-iPSCs

CML-CD34-iPSCs

Clone 1

TRA-1-60

TRA-1-60

Clone 2

A

CML-CD34-iPSCs

| | G13 | G15 | G17 | G18 |

IM -

IM +

B

Unstained

IM -   19.8%

IM +   57.4%

TRA-1-60

C

DAPI   TRA-1-60   DAPI+TRA-1-60

D

| | SSEA-4 | NANOG | TRA-1-60 | OCT4A | TRA-1-81 | SOX2 |

CML-CD34-iPSC Clone 1

CML-CD34-iPSC Clone 2

Normal iPSC Clone 1

Figure 3

Figure 4

A

Expansion of CD34+ cells

Seed CD34+ cells

Nucleofection

Medium change

Seed the cells for reprogramming

50% E8 Medium

100% E8 Medium

+IM

Dome-shaped colonies → flat colonies

No colony formation

Day   -3   0   2   5   7   10   14   18   24   31

+IM

B

DAPI   TRA-1-60   DAPI-TRA-1-60

C

IM-   IM+

Figure 5

IM+ CML-CD34-iPSCs

Figure 6

A

CML-CD34-iPSCs → Single cell-sorting → iPSCs → Mesoderm Differentiation → Emerging HSPCs

96-well plate

B

-5   0   3   7   12

iPSCs Expansion | Mesoderm induction | Endothelial to hematopoietic transition

E1   -IM

E2   - IM   +2µM IM

E3   +2µM IM

E4   +2µM IM   -IM

E5   +1µM IM

C

E1   34.3%
E2   26.9%
E3   47.2%
E4   43.3%
E5   32.4%

CD45 / CD34

D

No of cells (x10⁶)

Total cells
CD34+CD45+ cells

E1   E2   E3   E4   E5

Figure 7

Figure S1

A

B

Figure S2

A

Patient ID	Transcript Type
G17	e13a2
G15	e13a2
G18	e13a2
G13	e14a2

B

Patient ID	Clones	Transcript Type
G17	C1,C2 and C3	e13a2
G15	C1,C2 and C3	e13a2
G13	C1,C2 and C3	e14a2
G18	C1,C2,C4,C5,C6,C8 and C9	e13a2
	C7	Nil

C

TKI–R CML-CD34-iPSCs

G13

G15

TKI SOR CML-CD34-iPSC

G17

IM-Refractory CML-CD34-iPSCs

G18

Normal CD34-iPSCs

Ph-

Figure S3:

Ectoderm

Mesoderm

Endoderm

Figure S4

| Control | 5µM IM | 10µM IM | 20µM IM |

Figure S5

A

IM-Refractory iPSC Ph- clone G18
Morphology

B

IM-Refractory iPSC Ph- clone G18
Karyotype

Figure S6

Table S1

Sample ID	Age/Gender	TKI/Dose	Phase	Response status
G13	58/M	DA/50mg	CP	CCYR*
G15	53/M	DA/50mg	CP	SOR**
G17	28/M	DA/50mg	CP	CCYR*
G18	51/F	IM/400mg	CP	REFRACTORY***

Complete cytogenetic response - <1% BCR::ABL1 cells
**Sub-optimal response ->1% BCR::ABL1 cells*
***No hematological response at 3 month evaluation*