Isolation by environment is more important than isolation by distance along a tropical

gradient in direct developing frogs

Ruth Percino-Daniel[1,2], Kara Jones[3], Thomas A. Maigret[3], David W. Weisrock[3] and Daniel

Piñero[1]

[1]Departamento de Ecología Evolutiva, Instituto de Ecología, Universidad Nacional

Autónoma de México, Ciudad Universitaria, Coyoacán, Mexico City, Mexico CP 04510.

[2]Posgrado en Ciencias Biológicas, Universidad Nacional Autónoma de México, Coyoacán,

Mexico City, 04510, Mexico

[3]Department of Biology, University of Kentucky, Lexington, KY 40506, United States

Keywords: elevation gradient, *Craugastor,* environmental heterogeneity, isolation by

environment, genomics, RADseq

Running title: Landscape genomics of frogs across a tropical gradient

Abstract

To identify which abiotic factors are most important for defining population structure, both

spatiotemporal scales and species-specific life history attributes and ecology must be

accounted for. Our study aim is to quantify the extent to which divergence is driven by

abiotic factors such as temperature, precipitation and elevation. To do so, we used a direct-

developing frog species, *Craugastor loki* that occurs along a steep elevation gradient. Using

restriction-site associated DNA sequencing (RADseq) from individuals collected from 100

m to 2250 m of elevation at 13 localities at Sierra Madre de Chiapas in southern Mexico we

described population structure using a variety of model-based clustering and landscape

genomics approaches. We found that populations sampled at higher elevation correspond to

an undescribed new species of *Craugastor*, and that populations from *Craugastor loki*

between 120 m and 1500 m are clustered in two different genetic groups: a Pacific slope

group and a Central Depression slope group. We found that a null model of isolation-by-

distance was not supported and divergence is more likely influenced by the interaction

between elevation and precipitation, but also temperature. Our results underscore the

importance of isolation by environment and highlight that spatial analyses can illuminate

fine scale population structure.

Introduction

Biotic and abiotic factors modulate the balance between gene flow and population divergence [1] and can yield a wide range of patterns of genetic and phenotypic variation across the landscape [2]. For example, biotic factors related to life history traits and ecological interactions (e.g., dispersal abilities and generational time) have been shown to determine population structure [3,4]. Likewise abiotic factors, such as geography and environment (e.g., geographic barriers or environmental heterogeneity) also play a key role in population connectivity across a given landscape [2].

Restriction of gene flow across a landscape can often be explained by the ubiquitous pattern of isolation by distance (IBD; [5], where genetic differentiation increases with the increase in geographic differences. However, IBD assumes a uniform landscape and does not account for landscape heterogeneity. In contrast, a model of isolation by resistance (IBR) consider how characteristics of the landscape result in spatial genetic structure [6]. An IBR model predicts a positive relationship between genetic differentiation and resistance to gene flow. Gene flow can also be limited in environmentally heterogeneous landscapes due to divergent selection driving genetic differentiation, leading to isolation by environment (IBE). IBE can be identified by assessing the contribution of environmental factors on genetic variation while controlling for geographic distance [7,8]. IBD, IBR and IBE can easily be confounded with one another and disentangling which of these processes is more important is key to understanding how the landscape influences genetic variation.

Recent studies in amphibians have shown that different climatic variables (e.g., annual mean temperature, isothermality, temperature range and precipitation) function as drivers of population divergence, thus producing a pattern of IBE [9,10]. Thermal

differences can act as an environmental pressure on phenotypic traits through divergent

selection and increase population differentiation [10]. Likewise, seasonal variation in

temperature influences the movement and behavior of frogs [11] resulting in limited gene

flow. Indeed, candidate single nucleotide polymorphisms (SNPs) and structural variants

associated with temperature in frogs have been identified and suggest a potential functional

role in life history strategies [9,12]. Precipitation can limit amphibian distribution at both

the regional and local scales [13]. In dry seasons, frogs tend to move less because their

ability to absorb water from the ground decreases when soil water potential decreases

[14,15]. Despite an increasing number of studies that explore the contribution of

environment on population divergence in ectotherm species, much remains to be explored

in the way in which environmental variables shape gene flow and genetic diversity at a fine

scale. This is particularly true in tropical forests, where the potential for biodiversity loss is

higher.

At small scales, tropical mountains are highly heterogeneous in temperature,

precipitation, and humidity [16]. Thus, they are ideal for assessing the potential for IBR and

IBE to limit gene flow [17]. Elevation gradients can also be an ideal model to study which

factors can lead the adaptive differentiation and identify local adaptation [18,19].

The Sierra Madre de Chiapas is a physiographic region with an ample variety of

habitats, where lowland and intermediate habitats are influenced by the orientation of its

slopes: the Pacific and the Central Depression. The Pacific slope is more humid in

comparison to the Central Depression, due to a continental influence and a rain shadow

effect. Furthermore, the region offers a natural elevation gradient in both slopes. Given the

natural differences of temperature and precipitation, we can expect that IBE could

contribute more than IBD to population structuring, especially for ectothermic organisms

with low vagility and high site fidelity [20].

Here, we use the amphibian species *Craugastor loki*, a direct-developing frog

species found in North and Central America, inhabiting a variety of humid habitats in

Chiapas, Mexico. This group of frogs has a wide elevational distribution range extending

from sea level up to 2200 meters above sea level (masl), particularly in southern Mexico

[21,22], and is often locally abundant. Consequently, this species is particularly suitable for

investigating fine-scale genetic differentiation patterns across landscape-level gradients. Its

continuous distribution allows for testing abiotic factors, such as the environment, as

contributors to genetic differentiation and increased population structure. Our study aims to

(1) quantify both fine-scale genetic differentiation and more divergent evolutionary

relationships between populations of *C. loki* in the Sierra Madre de Chiapas, (2) test

whether patterns of genetic variation in *C. loki* are best explained by geographic distance or

environmental conditions, and (3) identify which specific environmental variables might be

associated with patterns of genetic differentiation in *C. loki*.

Materials and Methods

Study area and sampling

We collected 90 tissue samples of the direct-developing frog *C. loki* from 13 different

localities in the Sierra Madre de Chiapas of southern Mexico (Figure 1, Table S1). Our

sampling regime was designed to capture environmental variation along an elevation

gradient, with sampling sites ranging from 120 to 2200 masl [23]. We collected mostly toe

and, in some cases, liver tissues. Sampling was performed in the rainy season (June to

November in 2017). The number of samples varied from five to eight individuals per site.

We included *Craugastor pygmaeus* as an outgroup taxon, as it is closely related to the

members of *Craugastor rhodopis* group, which includes *C. loki*. We also included

Craugastor rupinius, which inhabits the Sierra Madre de Chiapas, is distantly related to *C.*

loki, and is nested within the *Craugastor punctariolus* species group [24].

DNA extraction and library preparation

Genomic DNA extractions were obtained using a QIAamp DNA Mini kit. Quality of

extractions were checked with agarose gel electrophoresis and the quantity of extractions

(20ng/µl) using a Qubit 3 Fluorometric Quantitation Kit (Invitrogen). We sequenced 94 *C.*

loki samples using a restriction-site associated DNA sequencing (RAD-seq) method,

employing the restriction enzyme *SbfI*. Sequencing libraries were run on two lanes of an

Ilumina HiSeq 2000. Library preparation and sequencing were performed at Floragenex Inc

(Eugene, Oregon, USA).

Preprocessing and data assembly

We trimmed adaptors, filtered out low-quality data and demultiplexed raw data of the 94

samples using process_radtags in Stacks v.2.5. [25]. We subsequently assembled the data

using Ipyrad v.0.7.30 [26], exploring parameter optimization with a focus on the maximum

allowed divergence among allelic variants and the number of samples required per locus.

We made two datasets, the first (dataset 1) considered all samples from lowlands

and intermediate elevation (120 to 1500 masl), the highlands (samples above to 2000 masl),

plus the two outgroup samples of *C. pygmaeus* and *C. rupinius.* The second dataset (dataset

2) only included samples from lowlands to intermediate elevation (100 to 1500 masl).

Samples from higher elevations likely represent a cryptic species. For dataset 1, assembly

and variant calling was performed using a clustering threshold of 0.85 and a minimum

number of samples per locus of 35. For dataset 2, we used 0.85 as clustering threshold and

30 as the minimum number of samples per locus. We filtered one SNP for each RAD-seq

locus using VCFTOOLS v.4.2 [27] and PLINK v.1.0.7 [28], and converted the SNP data to

VCF format for landscape genomic analysis. Finally, we filtered dataset 2 allowing for at

least 50% missing data, a minimum mean read depth of 26, and a minor allele frequency

(MAF) of 0.01.

Phylogenetic analyses

We reconstructed phylogenetic relationships among individuals of *C. loki* along the

elevation gradient and outgroup species using dataset 1 and two tree-building methods.

First, we used FastTree v.2.1.11 [29] using nearest-neighbor interchange for the topology

and subtree-prune and regraft moves (NNI + SPR) for branch length optimization. We used

a general time-reversible nucleotide substitution model with a single rate per site (GTR +

CAT). Second, we performed a maximum likelihood analysis in RAxML-HPC v.8.2.12

[30] using a GTR + GAMMA nucleotide substitution model and saving the best-scoring

tree. For node support we ran 100 bootstrap samples for both analyses. Trees were rooted

with our choice of outgroup. Analyses were performed on the CIPRES Science Gateway

v.3.3 [31]

Non-spatial population structure analyses

We used two non-spatially aware methods to evaluate population structure. First, we used

the non-parametric Discriminant Analysis of Principal Components (DAPC) with the

Adegenet package v.2.1.1 in R v.3.6.1 [32] to analyze both dataset 1 and dataset 2. To

determine the number of the groups, we used the find.clusters function and selected the

optimal K using the lowest BIC value exploring K 1 to 10. The optimal number of PCs to

retain was selected using a cross-validation procedure [33]. Second, we used the model-

based method Admixture v.13 (Alexander et al., 2009) to analyze dataset 2. We ran K from

1 to 5, with 10 iterations per K and selected the K with the lowest cross-validation error.

Spatially informed population structure analyses

We used multiple approaches to investigate the relationship between geographic and

genetic information. First, we inferred the strength of an IBD pattern and compared a

spatial model with a non-spatial model using the R package conStruct v.1.3 (Bradburd et

al., 2018) implemented in R. conStruct uses an explicit spatial component and models

genetic variation in genotyped individuals as partitioned within or admixed specific number

of discrete layers, within each layer the relatedness decays as a parameter function of the

distance between samples [34]. We analyzed K from 1 to 5, each with 10 replicates with

1000 MCMC iterations. We also used a cross-validation function to determine the

statistical support for models with and without a spatial component. Because conStruct is

sensitive to missing data, we collapsed some closely related individuals from the same

population into a single, multi-individual samples

(https://github.com/gbradburd/conStruct.git) and ran the cross-validation test using 35

multi-individual samples from 10 localities.

Next, we studied spatial and genomic variation across our sampled elevation

gradient using a spatial Principal Component Analysis (sPCA; [35] of dataset 2

implemented in Adegenet v.2.1.1. sPCA relies on an ordination approach for analyzing

spatial and genetic patterns; using Moran's *I* index, it identifies eigenvectors which

maximize genetic variation and spatial autocorrelation and maps selected eigenvectors onto

geographic space.

Finally, we used the ResistanceGA v.4.0.14 software [36] to associate specific

landscape attributes with levels of gene flow. The landscape attributes we used were

elevation, temperature, and precipitation (during the rainy season from July to November).

ResistanceGA uses pairwise genetic dissimilarity and a genetic algorithm to optimize

resistance surfaces. We employed least cost paths to estimate the effective distances across

the landscape [36] and used a linear mixed effects model with maximum likelihood to test

for landscape effects, where pairwise genetic distance was the response variable and the

optimized effective distance was the predictor variable. We used both approaches for each

individual variable and for all combinations of the individual surface and null model [36].

Isolation by environment (IBE)

We tested for a signature of IBE using both temperature and precipitation variables, factors

highly relevant to amphibian activity. Analyses were performed in the BEDASSLE v1.5 R

package [37], which models the covariance of allele frequencies as a Gaussian process and

uses a Bayesian model to estimate the contribution of environmental and geographic

variables. For environmental variables, we obtained bioclimatic variables available to

Mexico from Cuervo-Robayo et al. (2014) interpolated to ~90m resolution. We used annual

temperature, annual precipitation, temperature seasonality, and precipitation seasonality,

variables that have been associated with genetic differentiation in studies of amphibians

[9,10,12]. Pairwise Euclidean distances were generated between sampling sites for

environmental variables. Pairwise great-circle geographic distances were generated

between sampling sites using the function rdist.earth with the package field in R. We

standardized both distance matrices by dividing values by their standard deviation

constants. We performed two replicate Markov chain Monte Carlo runs to ensure

convergence of the parameters using the beta-binomial model, running one analysis for 5

million generations, and the second for 8 million generations, sampling every 1000

generations in both runs. Performance of the model was assessed by visualizing plot

acceptance rates and parameter trace plots. We discarded the first 50% of samples as the

burn-in and estimated the contribution of the environmental distance versus geographic

distance to genetic differentiation using the ratio (αE/αD).

Results

Sequence data and bioinformatics

We obtained ~661 million reads for the 94 samples, with 6.9 million mean reads per

sample. After filtering, we obtained a total of 848 loci for dataset 1(Table 2S) and 1801 loci

for dataset 2 (Table 3S). The assembled RADseq data set are available on Dryad repository

as VCF files.

Phylogenetics and population structure

Both FastTree and RAxML phylogenetically placed samples from the highlands group

(localities from elevation above 2000 masl, Figure 1) in a separate clade with 100%

bootstrap support (blue shaded clade in Figure 2a), suggesting that they may belong to a

different species of *Craugastor*. Some samples from intermediate elevations (samples from

1000 – 1200 masl) clustered with the outgroup samples: ArrNeg8 clustered with *C.*

pygmaeus and the samples NBra1_1 and NBra1_4 grouped with *C. rupinius*. This is likely

explained by the difficulty in differentiating adult members of *C. pygmaeus* from *C. loki*,

and the difficulty in differentiating juveniles of *C. rupinius* from *C. loki*. Samples from the

low-elevation localities and the majority of samples from intermediate-elevation localities

were recovered as a clade of *Craugastor loki* in both analyses (orange and green shaded

groups in Figure 2a).

The DAPC results for dataset 1 identified a $K = 3$ as the best-fitting model (Figure

S1a). The first axis accounted for 39.1% of the variation and identified the greatest

separation between a cluster containing samples from the highlands, the outgroup samples

(*C. pygmaeus* and *C. rupinius*), and the intermediate-elevation samples associated with

those clades in the phylogenetic trees, and two clusters containing samples from the

lowland and intermediate elevations (Figure 2b). Interestingly, four additional samples

(GnNB2_4 LaFlor2, LaLoma1, LaLoma7) from intermediate elevations were placed closer

to the "highland" cluster in ordination space. To verify the identity of these four samples,

we reviewed fieldnotes and confirmed uncertainty in our taxonomic identification; thus, we

omitted these individuals from dataset 2.

DAPC analysis of dataset 2 identified three groups, one of which is represented by

the orange cluster that corresponds to the Central Depression slope (Figure S1b). The other

two groups (light green and dark green) identified in discriminant space are samples from

the lowlands and intermediate elevations of the Pacific slope (Figure S1b), accounting for

16.26% of the conserved variance. BIC scores stabilized at values of $K = 3$ (Figure S1c).

Admixture analysis of dataset 2 identifying two groups with cross-validation

support for $K = 2$ as the best fit model but additional structure was identified at $K = 3$

(Figure 2c). One genetic cluster recovered samples from the Central Depression slope and

the other clustered the lowland and intermediate samples from the Pacific slope (Figure 2c).

conStruct analyses recovered two genetic clusters (Figure 2c): Central Depression

slope (orange color) and Pacific slope (green color) including samples from low and

intermediate elevations. However, there was no difference between non-spatial model and

spatial model of conStruct (Figure S4).

Landscape genomics

The sPCA detected a significant pattern both globally (P = 0.003) and locally (P =

0.002). The first axis identified a break between the clusters (Figure S5) associated with the

Central Depression slope and the Pacific slope (Figure 3). However, some sampled

localities fall at the boundary of the genetic break, corresponding to a lower elevation area.

The resistance surface analysis with ResistanceGA found the highest support for a

model containing rainy season precipitation and elevation, although a competing model

containing only temperature was nearly as well-supported (Table 1). Little to no support

was found for a null model of no geographic structuring or for isolation-by-distance.

Isolation by environment

Bayesian analysis of the relative contribution of IBE and IBD to genetic

differentiation in BEDASSLE produced evidence of IBE when elevation was considered.

Both temperature seasonality and precipitation seasonality show an important contribution

on genetic differentiation across elevation more than the geographic distance (Table 2,

Figure S6). The annual mean temperature and annual precipitation never reached

convergence despite running for 8 million MCMC iterations. The mean of the α_E/α_D ratio

can be interpreted as the relative effect of changing one degree Celsius or one millimeter

precipitation with one meter elevation distance between populations when the IBD has the

same effect size of IBE. That is, for elevation, we would expect that 1°C change of the

variation of temperature to have a similar effect on genetic divergence as a shift of 857.98

m.

Discussion

Disentangling which abiotic factors have an important role in population

connectivity and limiting gene flow can be challenging, particularly given that abiotic

environmental variables such as temperature and precipitation are usually autocorrelated

with geography [37], and frequently also with elevation [38]. Here, we found significant

genetic differentiation between populations of the direct-developing frog *Craugastor loki,*

and further found that patterns of gene flow were best explained by the interaction between

temperature and precipitation, with elevation suggesting a pattern of isolation by

environment.

We found that *Craugastor loki* inhabits areas from sea level to ~1500 masl.

Previous work on the evolutionary relationship of the *Craugastor rhodopis* group

(including *C. rhodopis, C. occidentalis* and *C. loki*) using mitochondrial DNA suggested

three clades of *C. loki* occurring from Mexico to Central America, with the southern clade

inhabiting mainly lowland areas [22]. Here we confirm that *C. loki* occupies mainly

lowland and intermediate elevations and that samples above 1500 masl possibly correspond

to another species. This group of frogs are highly polytypic and field identification is

challenging due to a lack of diagnostic characters [39,40]. Here, both phylogenetic and

clustering results of our genomic data show that the members of *Craugastor* found above

1500 masl are relatively distinct from lower elevation populations of *C. loki*, indicating that

they either belong to another species that we did not include in the analysis (e.g., *C.*

matudai, *C. montanus, C. greggi*) or possibly are an unidentified species. Our sampling

covers different localities that occur near the type locality of these species, like Cerro

Ovando (Figure 1), the type locality where *C. matudai* [41] and *C. montanus* occur [42].

Both species were described by museum specimens as *C. loki*. *Craugastor greggi* [43] is

also supposed to be present in this area, inhabiting cloud forest like the highland ecosystem

where we sampled. Even if all those species represent different groups in the taxonomical

classification of *Craugastor*, in the field these frogs could not to be distinguished from each

other and the original descriptions did not describe the vivid coloration of live individuals.

Further work and a geographic more extensive sampling are needed to clarify the

evolutionary relationships among these taxa.

Within *Craugastor loki*, we identified two main groups: one cluster from the Central

Depression slope and the other from the Pacific slope. Spatial analyses (e.g., sPCA) show a

discontinuity in the area sampled that matches a valley that breaks the range of the Sierra

Madre de Chiapas (Figure 3). It may be that the dispersal of frogs along the Sierra Madre

de Chiapas could be restricted by this break in elevation. Frogs often have low dispersal

capacity and limited movement up and down elevation gradients [44]. In addition, it seems

that the break identified by our spatial analysis is reflected in local forest communities. To

the southeast of where the range breaks the local conditions become more humid. In

contrast, to the northwest the conditions of the Sierra Madre de Chiapas are usually drier

and more influenced by the Central Depression, resulting in a continuous dry forest habitat.

Our landscape resistance and IBE analysis both suggest an effect of climatic

variables on spatial genetic differentiation. Precipitation and temperature are important

factors influencing the natural history and ecology of these direct-developing frogs,

characterized by the lack of a water-dwelling larval phase and their use of substrate

humidity for reproduction [44]. The Central Depression slope cluster presents a thermal

landscape different from the Pacific slope, where the former is characterized mainly by

temperature seasonality [23]. Hence, our results suggest that the abiotic variables

considered in our study, temperature and precipitation, have an important role in the pattern

of IBE.

Several studies on amphibians have documented patterns of genetic differentiation

across elevation gradients [17,45] and have found evidence of phenotypic divergence

associated with temperature and elevation, explained by a pattern of IBE [9]. In addition,

Medina et al. (2021) found a clinal pattern of genomic differentiation associated with

temperature and identified some candidate SNPs associated with temperature and body size

across the clinal gradient. Our results suggest a pattern of IBR and IBE. When we used

specific layers of temperature and precipitation, we found that the seasonality of both

parameters show signatures of IBE when elevation is taken into account. Both

environmental variables are intrinsically associated with elevation and the orientation of the

slope. In a previous study, Percino et al. (2021), studied some physiological traits from the

same localities of this study. The frogs that inhabit at the Central Depression slope showed

a different thermal sensitivity compared with the populations from Pacific slope. First, they

exhibit high thermal accuracy, meaning that their body temperature is close to the

microenvironmental temperature, and high thermal quality, that is, the thermal landscape

offers suitable microhabitats with temperature close to body temperature. Thus, it is

suggested that the landscape plays an important role on frog populations and eventually

could drive phenotypic and probably genotypic divergence as a result of potential local

adaptation. Further work is necessary to specifically study the role of local adaptation or

thermal plasticity coupled with evapotranspiration rates across the elevation of the Sierra

Madre de Chiapas.

Acknowledgements

We are very grateful to local authorities and landowners in Chiapas, who helped us and

allowed us to work in their properties, and also for their invaluable hospitality. Thanks to

undergrad students of the UNICACH for their invaluable help in the fieldwork, especially

to J. M. Contreras-Lopez. We are grateful to F. Mendez de la Cruz (IB-UNAM), for

extended his collected permits to RPD, which were issued by Secretaria del Medio

Ambiente y Recursos Naturales (SEMARNAT), no. SGPA/DGVS/002490/18 (license no.

FAUT-0074). We thank O. Godinez (CONABIO) for helping with environmental data and

A. Cardenas for advice in some R scripts. We are grateful to the University of Kentucky

Information Technology Department and the Center for Computational Sciences for

granting access to the Lipscomb High Performance Computing Cluster resources,

especially to Vikram Gazula. We also thank computer resources group from the CONABIO

cluster, especially E. Campos for his advice.

Competing interests

The authors declare no conflicts of interests exist.

Funding

RP-D received a fellowship from Consejo Nacional de Ciencia y Tecnologia (CONACYT

no. 434471) for her doctoral studies and grants for visiting scholar at University of

Kentucky (CONACYT-Beca de Movilidad 2019) with additional support of the Programa

de Apoyo a Estudios de Posgrado of UNAM (PAEP – UNAM 2019). This work was

supported by the Instituto de Ecología, Universidad Nacional Autónoma de México to DP

(IE-UNAM 2016-2019).

References

1. Coyne JA, Orr HA. 2004 *Speciation*. Sinauer Sunderland.
2. Cushman SA, McRae BH, McGaridal K. 2015 Basic of landscape ecology: An Introduction to landscape and population processes for landscape geneticist. In *Landscape Genetics: Concepts, Methods, Applications*, p. 257. Wiley Blackwell.
3. Steele CA, Baumsteiger J, Storfer A. 2009 Influence of life-history variation on the genetic structure of two sympatric salamander taxa. *Molecular Ecology* **18**, 1629–1639. (doi:10.1111/j.1365-294X.2009.04135.x)
4. Travis J. 1994 Ecological Genetics of Life-History Traits: Variation and Its Evolutionary Significance. In *Ecological Genetics* (ed LA Real), pp. 171–204. Princeton University Press.
5. Wright S. 1943 Isolation by Distance. *Genetics* **2**, 114–138.
6. McRae BH. 2006 Isolation by resistance. *Evolution* **60**, 1551–1561. (doi:10.1111/j.0014-3820.2006.tb00500.x)
7. Wang IJ. 2013 Examining the full effects of landscape heterogeneity on spatial genetic variation: A multiple matrix regression approach for quantifying geographic and ecological isolation. *Evolution* **67**, 3403–3411. (doi:10.1111/evo.12134)
8. Wang IJ, Bradburd GS. 2014 Isolation by environment. *Molecular Ecology* **23**, 5649–5662. (doi:10.1111/mec.12938)
9. Medina R, Wogan GOU, Bi K, Termignoni-García F, Bernal MH, Jaramillo-Correa JP, Wang IJ, Vázquez-Domínguez E. 2021 Phenotypic and genomic diversification with isolation by environment along elevational gradients in a neotropical treefrog. *Molecular Ecology* **30**, 4062–4076. (doi:10.1111/mec.16035)
10. Páez-Vacas MI, Trumbo DR, Funk WC. 2022 Contrasting environmental drivers of genetic and phenotypic divergence in an Andean poison frog (Epipedobates anthonyi). *Heredity* **128**, 33–44. (doi:10.1038/s41437-021-00481-2)
11. Seebacher F, Holmes S, Roosen NJ, Nouvian M, Wilson RS, Ward AJW. 2012 Capacity for thermal acclimation differs between populations and phylogenetic lineages within a species. *Functional Ecology* **26**, 1418–1428. (doi:10.1111/j.1365-2435.2012.02052.x)
12. Cayuela H *et al.* 2021 Genomic signatures of thermal adaptation are associated with clinal shifts of life history in a broadly distributed frog. *Journal of Animal Ecology* , 1365-2656.13545. (doi:10.1111/1365-2656.13545)
13. Ochoa☐Ochoa LM, Mejía☐Domínguez NR, Velasco JA, Marske KA, Rahbek C. 2019 Amphibian functional diversity is related to high annual precipitation and low precipitation seasonality in the New World. *Global Ecol Biogeogr* **28**, 1219–1229. (doi:10.1111/geb.12926)
14. Vitt LJ, Caldwell JP. 2014 *Herpetology: An Introductory Biology of Amphibians and Reptiles*. San Diego California, USA: Elsevier, Inc.
15. Wells KD. 2007 *The Ecology and Behavior of Amphibians*. The University of Chicago Press.
16. Stevens GC. 1992 The Elevational Gradient in Altitudinal Range☐: An Extension of Rapoport ' s Latitudinal Rule to Altitude Author (s): George C . Stevens Reviewed work (s): Source☐: The American Naturalist , Vol . 140 , No . 6 (Dec ., 1992), pp . 893-911 Published b. *American Society of Naturalists* **140**, 893–911.
17. Funk WC, Murphy MA, Hoke KL, Muths E, Amburgey SM, Lemmon EM,

Lemmon AR. 2016 Elevational speciation in action? Restricted gene flow associated with adaptive divergence across an altitudinal gradient. *Journal of Evolutionary Biology* **29**, 241–252. (doi:10.1111/jeb.12760)

18.   Luquet E, Léna JP, Miaud C, Plénet S. 2015 Phenotypic divergence of the common toad (Bufo bufo) along an altitudinal gradient: Evidence for local adaptation. *Heredity* **114**, 69–79. (doi:10.1038/hdy.2014.71)

19.   Wake DB. 1987 Adaptive radiation of salamanders in Middle American cloud forests. *Annals of the Missouri Botanical Garden* **74**, 242–264.

20.   Zamudio KR, Bell RC, Nali RC, Haddad CFB, Prado CPA. 2016 Polyandry, Predation, and the Evolution of Frog Reproductive Modes. *The American Naturalist* **188**, S41–S61. (doi:10.1086/687547)

21.   Frost D. 2020 Amphibian Species of the World: an Online Reference. Version 6.0. (doi:doi.org/10.5531/db.vz.0001)

22.   Streicher JW, García-Vázquez UO, Ponce-Campos P, Flores-Villela O, Campbell JA, Smith EN. 2014 Evolutionary relationships amongst polymorphic direct-developing frogs in the Craugastor rhodopis Species group (Anura: Craugastoridae). *Systematics and Biodiversity* **12**, 1–22. (doi:10.1080/14772000.2014.882428)

23.   Percino-Daniel R, Contreras López JM, Téllez-Valdés O, Méndez de la Cruz FR, Gonzalez-Voyer A, Piñero D. 2021 Environmental heterogeneity shapes physiological traits in tropical direct-developing frogs. *Ecology and Evolution* **11**, 6688–6702. (doi:10.1002/ece3.7521)

24.   Hedges SB, Duellman WE, Heinicke MP, World N, Molecular T. 2008 New World direct-developing frogs (Anura: Terrarana)... *Zootaxa* **1737**, 1–182.

25.   Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. 2013 StackCatchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool set for population genomics. Molecular Ecology, 22(11), 3124–40. http://doi.org/10.1111/mec.12354s: an analysis tool set for population genomics. *Molecular ecology* **22**, 3124–40. (doi:10.1111/mec.12354.Stacks)

26.   Eaton DAR, Overcast I. 2020 Ipyrad: Interactive assembly and analysis of RADseq datasets. *Bioinformatics* **36**, 2592–2594. (doi:10.1093/bioinformatics/btz966)

27.   Danecek P *et al.* 2011 The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158. (doi:10.1093/bioinformatics/btr330)

28.   Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. 2015 Second-generation PLINK: Rising to the challenge of larger and richer datasets. *GigaScience* **4**, 1–16. (doi:10.1186/s13742-015-0047-8)

29.   Price MN, Dehal PS, Arkin AP. 2009 Fasttree: Computing large minimum evolution trees with profiles instead of a distance matrix. *Molecular Biology and Evolution* **26**, 1641–1650. (doi:10.1093/molbev/msp077)

30.   Stamatakis A. 2014 RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313. (doi:10.1093/bioinformatics/btu033)

31.   Miller MA, Pfeiffer W, Schwartz T. 2010 Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *2010 Gateway Computing Environments Workshop, GCE 2010* (doi:10.1109/GCE.2010.5676129)

32.   Jombart T, Ahmed I. 2011 adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics* **27**, 3070–3071. (doi:10.1093/bioinformatics/btr521)

33.   Jombart T, Devillard S, Balloux F. 2010 Discriminant analysis of principal

components: a new method for the analysis of genetically structured populations. *BMC*
Genetics **11**. (doi:10.1371/journal.pcbi.1000455)
34.   Bradburd GS, Coop GM, Ralph PL. 2018 Inferring continuous and discrete
population genetic structure across space. *Genetics* **210**, 33–52.
(doi:10.1534/genetics.118.301333)
35.   Jombart T. 2008 Adegenet: A R package for the multivariate analysis of genetic
markers. *Bioinformatics* **24**, 1403–1405. (doi:10.1093/bioinformatics/btn129)
36.   Peterman WE. 2018 ResistanceGA: An R package for the optimization of resistance
surfaces using genetic algorithms. *Methods in Ecology and Evolution* **9**, 1638–1647.
(doi:10.1111/2041-210X.12984)
37.   Bradburd GS, Ralph PL, Coop GM. 2013 Disentangling the effects of geographic
and ecological isolation on genetic differentiation. *Evolution* **67**, 3258–3273.
(doi:10.1111/evo.12193)
38.   Barry RG. 2008 *Mountain Weather and Climate*. Cambridge University Press.
(doi:https://doi.org/10.1017/CBO9780511754753)
39.   Campbell JA, Savage JM. 2000 Taxonomic Reconsideration of Middle American
Frogs of the Eleutherodactylus rugulosus Group (Anura: Leptodactylidae): A
Reconnaissance of Subtle Nuances among Frogs. *Herpetological Monographs* **14**, 186.
(doi:10.2307/1467048)
40.   Crawford AJ, Smith EN. 2005 Cenozoic biogeography and evolution in direct-
developing frogs of Central America (Leptodactylidae: Eleutherodactylus) as inferred from
a phylogenetic analysis of nuclear and mitochondrial genes. *Molecular Phylogenetics and*
Evolution **35**, 536–555. (doi:10.1016/j.ympev.2005.03.006)
41.   Taylor EH. 1941 New Amphibians from the Hobart M. Smith Mexican Collections.
University of Kansas Science Bulletin **27**, 141–167.
42.   Taylor EH. 1942 New Tailless Amphibian from Mexico. *University of Kansas*
Science Bulletin **38**, 67–89.
43.   Lynch JD. 1965 A new species of Eleutherodactylus (Leptodactylidae: Anura) from
southeastern Chiapas, Mexico. *Natural History Miscellanea* **181**, 1–6.
44.   Duellman WE, Trueb L. 1994 *Biology of Amphibians*. The Johns Hopkins
University Press.
45.   Funk WC, Blouin MS, Corn PS, Maxell BA, Pilliod DS, Amish S, Allendorf FW.
2005 Population structure of Columbia spotted frogs (Rana luteiventris) is strongly affected
by the landscape. *Molecular Ecology* **14**, 483–496. (doi:10.1111/j.1365-
294X.2005.02426.x)
503

Data Accessibility

The data are accessible to the repository in Dryad (XXXXXXX). The repository contains

the data in vcf format.

Author Contributions

RPD and DP designed the research. RPD performed the fieldwork and lab work. RPD, KJ,

TM analyzed the data. DP, DW contributed with analytical tools. RPD lead the manuscript

writing with input of all authors.

Table 1. Summary of model selection for the generalized linear mixed-effects models carried out in ResistanceGA. The null model assumes no geographic structure and the alternative models performed with environmental surfaces using temperature, precipitation, and elevation, where the interaction between elevation and precipitation is well fitted to the model.

Layer	k	AIC	AICc	R2m	R2c	LL	delta.AICc	weight
Elevation and Precipitation	7	-7893.07	-7891.07	0.0516	0.1276	3953.53	0	0.3965
Temperature	4	-7891.19	-7890.51	0.0425	0.1129	3949.60	0.5546	0.3005
Precipitation and temperature	7	-7891.14	-7889.14	0.0487	0.1247	3952.57	1.9287	0.1512
Elevation and temperature	7	-7890.51	-7888.51	0.0486	0.1213	3952.25	2.5596	0.1103
Precipitation	4	-7885.98	-7885.30	0.0513	0.1389	3946.99	5.7669	0.0222
Distance	2	-7884.71	-7884.51	0.0260	0.0950	3944.35	6.5607	0.0150
Elevation, Precipitation, and temperature	10	-7885.42	-7881.27	0.0494	0.1256	3952.71	9.7999	0.0030
Elevation	4	-7880.71	-7880.02	0.0260	0.0949	3944.35	11.0420	0.0016
Null	1	-7836.83	-7836.77	0	0.0630	3919.41	54.3006	6.41E-13

Table 2. Results of the BEDASSLE analysis. We ran two replicates for each analysis. The first runs compared the relative effect of the precipitation seasonality (αE) and geographical distance (αD) on genetic differentiation. The third and fourth runs compared the relative effect of the temperature seasonality (αE) and geographical distance (αD). The fifth and sixth runs compared the relative effect of the precipitation seasonality (αE) and elevation distance (αD) on genetic differentiation, while the last two runs compared the relative effect of the precipitation seasonality (αE) and elevational distance (αD).

	Ratio (αE/αD)	95% Confidence Intervals
Precipitation seasonality and geographical distance	0.428	(0.212 – 0.695)
Precipitation seasonality (replicate)	0.390	(0.192 – 0.621)
Temperature seasonality and geographical distance	0.483	(0.052 – 1.196)
Temperature seasonality (replicate)	0.436	(0.214 – 0.710)
Precipitation seasonality and elevation distance	12.674	(1.192 – 29.929)
Precipitation seasonality (replicate)	10.323	(1.119 – 25.548)
Temperature seasonality and elevation distance	857.980	(446.588 – 1629.37)
Temperature seasonality (replicate)	932.536	(453.102 – 1826.62)

Figure 1. Sampled localities of *Craugastor* in Chiapas, Mexico. The orange and green circles correspond to the two genetic clusters of *C. loki* identified by Admixture and conStruct analysis, as seen in Figure 2. Blue circles correspond to the localities of *Craugastor* sp. from the highlands (see Supplementary material). Black circles with white outline are visited localities where frogs were not found. Triangle and square symbols correspond to the type localities of the *C. montanus* and *C. matudai,* respectively.

Figure 2. (a) Results from phylogenetic analysis using maximum likelihood performed in RAXML and (b) discriminant analysis of principle components of the entire genomic data set. The individuals from higher elevation populations may correspond to other species of *Craugastor*. Colors correspond to the individuals sampled in the different localities shown in Figure 1. (c) Genetic assignment of the sampled populations using Admixture (from K = 2 to K = 3) and non-spatial model performed in conStruct. Individuals are ordered from lower to higher elevation. Elevation (in meters) is represented by horizontal bars below the assignment plots. Central Depression cluster and Pacific slope are shown in orange and green respectively. Photos in Fig. 1 by R. Percino-Daniel (*C. rupinius* & C. sp) and J. E. Pérez Sanchez (*C. pygmaeus*).

Figure 3. Interpolated spatial genetic structure based on sPCA superimposed over elevation map. The bar at left represents the interpolated vector scores, which corresponds to the degree of differentiation between individuals. The white color represents the break of the mountains edge of the highlands at Sierra Madre de Chiapas (lower elevation area). Black dots represent sampled sites.