Diversity and evolutionary growth of biomineralized columns in early Cambrian phosphatic-shelled brachiopods

Zhiliang Zhang[1,*], Zhifei Zhang[2], Lars E. Holmer[3], Timothy P. Topper[2,4], Bing Pan[1], Guoxiang Li[1]

[1] State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China; [2] State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life & Environments, Department of Geology, Northwest University, Xi'an, 710069, China; [3] Institute of Earth Sciences, Palaeobiology, Uppsala University, SE-752 36 Uppsala, Sweden; [4] Department of Palaeobiology, Swedish Museum of Natural History, Box 50007, SE-104 05, Stockholm, Sweden

Abstract Biologically controlled mineral crystals producing organic-inorganic composites (hard skeletons) by metazoan biomineralizers has been an evolutionary innovation for more than half billion years on Earth. Among them, linguliform brachiopods are the only invertebrates that secrete phosphate to build their skeletons. One of the most distinct shell structures is the organic-phosphatic columns applied exclusively by phosphatic-shelled brachiopods. However, the complexity, diversity and biomineralization process of these microscopic columns are far from clear in brachiopod ancestors. Here, exquisitely well preserved columnar structures are discovered for the first time in the earliest eoobolids. The hierarchical shell architectures, epithelial cell moulds, and the shape and size of cylindrical columns are scrutinised in *Latusobolus xiaoyangbaensis* gen. et sp. nov. and *Eoobolus acutulus* sp. nov from the Cambrian Series 2 Shuijingtuo Formation of South China. The secretion and construction of the stacked sandwich model of columnar shell, which played a significant role in the evolution of linguliforms, is highly biologically controlled and organic-matrix mediated. Furthermore, a continuous transformation of anatomic features resulting

28   from the growth of columns is revealed between Eoobolidae, Lingulellotretidae and

29   Acrotretida, shedding new light on the evolutionary growth and adaptive innovation of

30   stacked sandwich columns among early phosphatic-shelled brachiopods during the

31   Cambrian explosion.

34   Introduction

The early Cambrian witnessed a great burst of animal body plans and biomineralized shell

architectures around half billion years ago (Briggs, 2015; Erwin, 2015, 2020; Budd and

Jackson, 2016; Murdock, 2020; Zhang and Shu, 2021; Yun et al., 2021; Z.L. Zhang et al.,

2021). The novel biomineralizations, biologically controlled mineral crystals producing

organic-inorganic composites (hard skeletons), in these animals have played a vital role in

the survival and fitness of early species (Balthasar and Cusack, 2015; Cuif et al., 2010; Li et

al., 2022; Skovsted et al., 2008; Yun et al., 2022), and in turn build the fundamental bricks of

complex marine ecosystems (Bicknell and Paterson, 2018; Buatois et al., 2020; Chen et al.,

2022; Zhang et al., 2010; Z. Zhang et al., 2020). Since the early Cambrian to the present day,

this adaptive evolution has been excellently demonstrated and continuously preserved in

brachiopods, one of the key members of Cambrian Evolutionary Fauna (Carlson, 2016;

Harper et al., 2017a, 2021; Sepkoski, 1984). Among them, the phosphatic-shelled

brachiopods are some of the most common animals in early Cambrian faunas. With high

fidelity preservation and high abundance of biomineralized shells in the fossil record, their

morphological disparity, diversity of shell structures and growth patterns, and ecology

complexity are preserved in great detail (Chen et al., 2021; Claybourn et al., 2020; Zhang,

2018; Z. L. Zhang et al., 2020a, 2021b).

Studying the processes by which organisms form biomineral materials has been the

focuse at the interface between earth and life sciences (Leadbeater and Riding, 1986;

Simonet Roda, 2021). Brachiopods are very unique animals in having the ability to secrete

two different minerals, calcium phosphate and calcium carbonate, making them the ideal

group to further explore the processes of biomineralization (Simonet Roda et al., 2021a;

Williams, 1977). Hard tissues composed of calcium phosphate with an organic matrix are

also largely present in vertebrates, which have remarkably shaped the evolutionary

trajectory of life on Earth (Ruben and Bennett, 1987; Wood and Zhuravlev, 2012). The origin

of phosphate biomineralization in the evolutionary distant invertebrate brachiopods and

vertebrates is still a big mystery in animal evolution (Luo et al., 2015; Neary et al., 2011;

Simonet Roda, 2021). Thus, more research is needed in order to understand how to relate

hierarchical structure to function in the very early examples of calcium phosphate-based

biomineralization process (Kallaste et al., 2004; Weiner, 2008). It is noteworthy, that South

China has been considered as one of centres for the origination and early dispersal of

phosphatic-shelled brachiopods (Z. L. Zhang et al., 2021a). It provides a great opportunity to

explore the unique biomineralization process and consequent adaptive evolution of their

earliest representatives during the Cambrian radiation of life.

The shell forming process of brachiopods have long represented a conundrum, which is

crucial to unravel their poorly resolved phylogeny and early evolution (Carlson, 2016;

Cusack and Williams, 2007; L. Holmer et al., 2008; Murdock, 2020; Streng et al., 2007;

Temereva, 2022; Williams et al., 2004; Williams and Cusack, 1999). Extensive studies have

been conducted on living and fossil shells, but most of them are focused on articulated or

carbonate-shelled representatives (Cusack et al., 2010; Griesshaber et al., 2007; Simonet

Roda et al., 2021b; Simonet Roda, 2021; Ye et al., 2021). By contrast, the linguliform

brachiopods, which are composed of an organic matrix and apatite minerals that show

extremely intricate architectures and permit exquisite preservation, are less well understood,

and their shell structural complexity and diversity, especially of their fossil representatives

require further studies (Cusack et al., 1999; Streng et al., 2007; Williams and Holmer, 1992;

Zhang et al., 2017). The building of shells by microscopic cylindrical columns is a unique

feature, which is restricted to the phosphatic-shelled brachiopods and their assumed

ancestors (L. Holmer et al., 2008; Holmer, 1989; Holmer et al., 2002; Skovsted and Holmer,

2003; Williams and Holmer, 2002a). These type of columns were previously believed to be

exclusively restricted to micromorphic acrotretide brachiopods, a group which demonstrate

more complex hierarchical architectures and graded structures compared to simple lamella

shell structure in older lingulides (Holmer, 1989; Williams and Holmer, 1992). As a result of

further studies, diverse columnar shell structures have been recognised in more brachiopod

groups, including stem group taxa *Mickwitzia* and *Micrina* (Skovsted and Holmer, 2003),

lingulellotretid *Lingulellotreta* (L. Holmer et al., 2008), eoobolid *Eoobolus* (Streng et al., 2007;

Z. L. Zhang et al., 2021a), and the possible paterinide *Bistramia* (L. Holmer et al., 2008).

However, the columnar shells among the oldest linguliforms and their evolutionary variations

have not been scrutinised.

For the first time, exquisitely well preserved columnar shell structures are described from

the oldest known eoobolid brachiopods. *Latusobolus xiaoyangbaensis* gen. et sp. nov., and

Eoobolus acutulus sp. nov. are described here, based on new specimens from the Cambrian

Series 2 Shuijingtuo Formation of southern Shaanxi and western Hubei in South China. In

this study, the shell architectures, epithelial cell moulds, and the shape and size of cylindrical

columns are minutely examined, shedding new light for understanding the architecture

intricacy, biomineralization process and evolutionary fitness of early phosphatic-shelled

brachiopods.

Results

Systematic palaeontology

Brachiopoda Duméril, 1806

Linguliformea Williams, Carlson, Brunton, Holmer and Popov, 1996

Lingulata Gorjansky and Popov, 1985

Lingulida Waagen, 1885

Linguloidea, Menke, 1828

Eoobolidae, Holmer, Popov and Wrona, 1996

Genus *Latusobolus* Zhang, Zhang and Holmer gen. nov.

Type species. *Latusobolus xiaoyangbaensis* sp. nov., here designated.

Etymology. From the Latin '*latus*' (wide) with the ending '*obolus*', to indicate the

transversely oval outline of both ventral and dorsal valves, morphologically similar to *Obolus*.

The gender is masculine.

Diagnosis. For a full description and discussion of *Latusobolus* gen. nov., refer to

Appendix 1.

Latusobolus xiaoyangbaensis Zhang, Zhang and Holmer sp. nov.

Figure 1 and *Appendix 2—figures 1-4*, *Appendix 3—table 1*.

Etymology. After the occurrence at the Xiaoyangba section in southern Shaanxi, China.

Type material. Holotypes, ELI-XYB S5-1 BR09 (*Appendix 2—figure 1M–P*), ventral

valve and ELI-XYB S4-2 BO11 (*Appendix 2—figure 2M–P*), dorsal valve, from Cambrian

Series 2 Shuijingtuo Formation at the Xiaoyangba section (Z. L. Zhang et al., 2021a) near

Xiaoyang Village in Zhenba County, southern Shaanxi Province, China.

Diagnosis. As for the genus.

Description. For a full description and discussion of *Latusobolus xiaoyangbaensis* gen.

et sp. nov., refer to Appendix 1.

Genus *Eoobolus* Matthew, 1902

Type species. *Obolus* (*Eoobolus*) *triparilis* Matthew, 1902 (selected by Rowell, 1965)

Diagnosis. See Holmer et al. (p. 41) (Holmer et al., 1996).

Eoobolus acutulus Zhang, Zhang and Holmer sp. nov.

Figure 2 and *Appendix 2—figures 5–7*, *Appendix 3—table 2*.

Etymology. From the Latin '*acutulus*' (somewhat pointed), to indicate the slightly

acuminate ventral valve with an acute apical angle. The gender is masculine.

Type material. Holotypes, ELI-AJH S05 BT11 (*Appendix 2—figure 5E–H*), ventral

valve, and ELI-AJH S05 1-5-07 (*Appendix 2—figure 5M*), dorsal valve, from Cambrian

Series 2 Shuijingtuo Formation at the Aijiahe section (Z. L. Zhang et al., 2016) near Aijiahe

Village in Zigui County, north-western Hubei Province, China.

Diagnosis. For a full description and discussion of *Eoobolus acutulus* sp. nov., refer to

Appendix 1.

Discussion

Complex linguliform brachiopod shells

Although the supposed living fossil *Lingula* has long been considered to virtually lack

morphological evolutionary changes (Schopf, 1984), more recent studies have shown that it

actually has experienced dramatic modifications in many aspects (Liang et al., 2023),

including internal organs (Zhang et al., 2008a), life mode (Topper et al., 2015), shell

structures (Cusack et al., 1999), and even genome (Goto et al., 2022; Luo et al., 2015). The

complexity and diversity of linguliform shell architecture was increasingly recognised in the

pioneering study of Cusack, Williams and Holmer (Cusack et al., 1999; Holmer, 1989;

Williams and Cusack, 1999; Williams and Holmer, 1992). What's more, such complex

architectures have a wide distribution in closely related brachiopod groups at the beginning

of Cambrian, when they made their first appearance. In connection with an ongoing

comprehensive scrutiny of well-preserved linguliform shell architectures from the early

Cambrian limestones of South China, their complexity and diversity hidden in the

conservative oval shape is becoming more and more intriguing. However, compared to their

living representatives, the shell structures are relatively simple, revealing profound

modifications during the long evolutionary history (Cusack et al., 1999; L. Holmer et al., 2008;

Williams, 1977; Williams and Cusack, 1999).

In general, the shells of organo-phosphatic brachiopods are stratiform, composed of an

outer periostracum and inner rhythmically-disposed succession of biomineralized lamellae or

laminae (Holmer, 1989; Williams, 1977). The organic periostracum, serving as a rheological

coat to the underneath shell, is scarcely fossilized. However, its wrinkling and vesicular

features have largely been preserved as superficial imprints (pits, pustules, fila, grooves,

ridges, rugellae, drapes, reticulate networks, spines) on the surface of the primary layer, and

are important characters in understanding brachiopod phylogeny (Holmer 1989; Holmer *et al.*

1996; Cusack *et al.* 1999). By contrast, the structures preserved in the secondary layer have

been characterised in fossil and living groups by three ancient fabrics –columnar, baculate,

laminated – all of which persists in living shells except for the columnar fabric (Cusack et al.,

1999).

The primary layer commonly consists of one heavily biomineralized compact lamina

composed of apatite granules, with a thickness from 2 μm to 20 μm (**Figures 1B, C** and **2G**).

Usually, the concentric growth lines are regularly distributed on primary layer surface of post-

metamorphic shell. By contrast, the surface ornamentation tends to be less ordered in

distribution, demonstrating a strong phylogenetic differentiation. Superficial pustules are one

of the most distinct patterns and are readily recognised in one of the oldest brachiopod

group, the Eoobolidae (Holmer et al., 1996). The pustules are roughly round in outline,

composed of apatite aggregates, and range from 2 µm to 20 µm in diameter (*Figure 1G*).

Such pustules are also found on early obolid, zhanatellid and acrotheloid shells with

relatively wide size variations from 5 µm to 30 µm. Although different in size, the similar

pattern may indicate a same secretion regime that originated as vesicles during the very

early stages in periostracum secretion (Cusack et al., 1999). The thickness of the

underneath secondary layer varies greatly in different brachiopod groups, depending on

shell component and fabric type. Columnar, baculate and laminated fabrics are incorporated

into the basic lamination component to form the diverse stratiform successions of secondary

shell layer.

The fossil record reveals that the columnar shell structure is generally preserved in most

early linguliforms (L. Holmer et al., 2008; Streng et al., 2007; Williams, 1977; Zhang, 2018; Z.

L. Zhang et al., 2020a, 2021a). It is a multi-stacked sandwich architecture (multi columnar

laminae are stacked in a vertical direction), which is initially developed in the earliest

linguliform *Eoobolus* as an early stage (Z. L. Zhang et al., 2021a). One to three stacked

sandwich units with short columns can also be found in *Latusobolus xiaoyangbaensis* gen.

et sp. nov. (*Figure 1*) and *E. incipiens* (*Figure 4A*), while the number of stacked sandwich

unit increased in later eoobolids (*Figure 2*), obolids and lingulellotretids (*Figure 4B–D*).

Eventually, the acrotretides developed a more complex columnar layer with multiple stacked

sandwich units (*Figure 4F–J*). Moreover, the columnar shell structures were also found in

stem group brachiopods, e.g., *Mickwitzia*, *Setatella* and *Micrina*, but with different column

size and numbers of lamination (L. Holmer et al., 2008; Holmer et al., 2002; Skovsted et al.,

2010; Williams and Holmer, 2002a). It is assumed here that the columnar layer maybe

plesiomorphic in linguliform brachiopods, and was inherited from stem group brachiopods.

Biomineralization process of organo-phosphatic columnar shell

Metazoans are known for secretion of very different types of biominerals through the

biological mineralization process, which linked the soft organic tissues of life with minerals of

solid earth in processes that would revolutionised the Earth's fossil record (Addadi and

Weiner, 2014; Lowenstam and Weiner, 1989; Simonet Roda, 2021; Wood and Zhuravlev,

2012). Because of the fine nature of organo-phosphatic biomineralization in linguliforms

(Cusack et al., 1999; Williams and Cusack, 1999), they have an exquisitely finely preserved

shell architectures (*Figures 1-2*, *3A*, *B* and *4*), including the epithelium cell moulds (*Figure*

3C and *F–I)* among the earliest eoobolids, lingulellotretids and acrotretids. This permits us to

reconstruct the biomineralization process of their apatitic cylindrical columns and address

key questions how these hierarchical structures relate to mechanical functions. Although the

biomineralization process of living brachiopods at the cellular level is not well known yet,

many biochemical experiments (Cusack et al., 1999, 1992; Lévêque et al., 2004; Williams

and Cusack, 1999) reveal the high possibility of the biologically controlled and organic matrix

mediated extracellular mineralization during brachiopod shell secretion, which can be

compared to the hard tissue forming process of mollusc shells and vertebrate teeth (Golub,

2011; Neary et al., 2011; Simonet Roda, 2021).

Many generally polygonal structures(*Figure 3C* and *F*), preserved on the internal

surface of successively alternating laminae (*Figure 3G–I*), have generally been considered

to represent the moulds of epithelial cells (McClean, 1988). The average size of epithelial

cells is 20 μm, ranging from 5 to 30 μm. The rheological vesicle environment may cause the

shape variation of epithelia, while different secreting rate could result in different sizes;

usually smaller epithelial cells had high secretion rates (Z. L. Zhang et al., 2016). As the

active secretion activity of outer epithelial cells, they can be easily embedded in the newly

formed inner and outer bounding surfaces of shell laminae (McClean, 1988; Winrow and

Sutton, 2012), which left shallow grooves between epithelial margins as intercellular

boundaries *(Figure 3F)*. Epithelial cells are preserved as moulds – presumably by the

phosphatization of smooth organic sheets (*Figure 4G* and *H*) – that had been secreted

relatively slow by the outer plasmalemma of the outer mantle (Cusack et al., 1999).

These active cells are responsible for the secretion of linguliform periostracum with

rhemorphic wrinkling features (Cusack et al., 1999), which left typical microtopography on

the external surface of primary layer beneath (*Figure 3A* and *B*). On the other hand, they

also secreted linguliform biomineral in a rheological extracellular environment, inferred from

living *Lingula* as apatite growing from an amorphous calcium phosphate precursor, which

forms the basic crystals of apatite around 5 nm in diameter (Lévêque et al., 2004; Williams

and Cusack, 1999). These nanoscale crystals were packaged into apatite granules with an

average size of 100 nm (*Figures 1I* and *2K*), which acted as the fundamental component

building up the hierarchical stratiform columnar shells, including the primary layer and

secondary layer (*Figure 5*). The apatite granules were probably coated or saturated with

organic compounds to form granule aggregations or clusters (spherular mosaics) as irregular

spherules or rods of about 500 nm (*Figures 1I* and *2H; Appendix 2—figure 7J* and *K*),

usually leaving gaps between the aggregation boundaries in fossils after the degradation of

organic counterparts (*Figure 2G; Appendix 2—figure 4K*). These apatite spherules are

aggregated in the planar orientation as a compact thin lamella less than 4 µm. Several thin

lamellae are closely compacted as the primary layer (*Figure 1B* and *C*). Some less obvious

gaps were preserved, and they may indicate the existence of organic matrix intercalated

randomly (*Figures 1D*, *I* and *2F*; *Appendix 2—figures 4I*, *J* and *7J*). On the other hand,

similar nanometre scale networks of spherules are aggregated to be linearly organised as an

orthogonal column perpendicular to the layer surface. Each columnar composite unit

consists of numerous columns disposed orthogonally between a pair of compact lamella

(Cusack et al., 1999; Holmer, 1989). Multi columnar laminae are stacked in a vertical

direction from exterior to interior to form the secondary layer, applying a stacked sandwich

model (*Figure 5*), which differs from the layer cake model. A very thin gap, commonly less

than 1 µm, between each pair of stacked column lamina is very obvious in well preserved

specimens (*Figures 1C*, *D*, *I*, *2C*, *F*, and *4C; Appendix 2—figures 4I* and *7J*), probably

indicating one organic membrane acts as an extracellular matrix with functions of a template

guiding mineral nucleation (Addadi and Weiner, 2014; Cusack et al., 1999; Lévêque et al.,

2004). This is also supported by the symmetrical nature of the stacked columnar architecture,

revealing a homogeneous organic substrate responsible for the succeeding rhythmic

sequence. Although newly secreted columnar unit may succeed the older one

unconformably with overlap (*Figure 2C*) and be involved in lateral facies changes (*Figure*

2K and *4C*), it is supposed to reflect intracellular deviation with the same secretory cycle with

the outer mantle as a whole as in living lingulids (Cusack et al., 1999; Williams et al., 1992).

The secreting and building process of the early Cambrian phosphatic-shelled brachiopod

columnar shell, secreted by the underlying outer epithelium cells of the mantle lobe, is

demonstrated in *Figure 5*.

It is worth noting that nanoscale openings are permeated on the terminal end of the

columns (*Figures 1C*, *4E* and *H*) and surface of columnar laminae (*Figures 1H, 2H*, *3E* and

4G) on well preserved specimens. The openings are round in shape with a mean size of 600

nm. Observation from some natural fractures of shells, some canals can be traced

continuously through several columnar sequences (*Figures 2G* and *4E*), while in most other

poorly preserved fossils, they are filled with secondarily phosphatised spherules or mosaics

(*Figures 1I* and *2G*), occasionally leaving random gaps (*Figures 1J*, *2F* and *4I*). These

canals are very likely empty space of degraded organic material, which is confirmed by a

different taphonomic process that preserves canals in contemporaneous Cambrian

siliciclastic rocks (Duan et al., 2021). The regular arrangement of the canal systems and

closely related columns revealed that the process is organic matrix-mediated, and moreover

the even disposition of the organic matrix in columns indicate the rheological and center

areas, on which biologically controlled biomineralization took place that the nucleation,

growth and aggregation of the deposited amorphous calcium phosphate is directed by the

same group of epithelial cells (Cusack et al., 1999; Pérez-Huerta et al., 2018; Simonet Roda,

2021; Weiner and Dove, 2003). In the stacked columnar units, the chambers between each

columns delineated by a pair of compact lamellae (**Figure 4F** and **J**) would be originally filled

with glycosaminoglycans (GAGs) as in living *Lingula* (Cusack et al., 1999; Williams et al.,

1994). The chambers are often filled with coarse spherular mosaics when being secondarily

phosphatised, resulting in being indistinguishable between columns, paired compact

lamellae and organic membrane (**Figures 1D** and **2F**). At the margin of mature shells,

especially the ventral pseudointerarea where the vertical component of growth vector

becomes increasingly important, the short columns are replaced by relatively higher columns,

resulting in the transformation from two stacked columnar units into one unit during growing

anteriorly (**Figures 2C**, **K**, **3I** and **4C**), which may demonstrate allometric growth of shell

(Cusack et al., 1999).

The most intriguing and enigmatic phenomenon of skeletal biomineralization is the

evolutionary selection of calcium carbonate and calcium phosphate in invertebrates and

vertebrates, respectively (Lévêque et al., 2004; Luo et al., 2015). However, the Brachiopoda

is a very unique phylum that utilises both. For the subphylum Linguliformea, the appearance

of apatite as a shell biomineral dates back as far as the early Cambrian (Topper et al., 2013;

Ushatinskaya, 2002) and persists to the present (Carlson, 2016). Calcium phosphate can

build a relatively less soluble skeletal component as compared with calcium carbonate shells,

but with the disadvantage of a greater energetical and physiological cost (Wood and

Zhuravlev, 2012). The acquisition of this specific biomineral in phosphatic-shelled

brachiopods has been considered an ecological consequence of the globally elevated

phosphorous levels during phosphogenic event in the calcite seas with a low Mg:Ca ratio

and/or high CO_2 pressure (Balthasar and Cusack, 2015; Brasier, 1990; Cook and Shergold,

1984; Wood and Zhuravlev, 2012). In such situations, linguliforms were likely to utilise the

sufficient phosphorous in ambient waters by chance from a brachiopod ancestor with an

unmineralized shell coated with detrital grains, like *Yuganotheca* found in the Chengjiang

Lagerstätte (Zhang et al., 2014), possibly as an evolutionary response of prey to an

escalation in predation pressure during the Cambrian explosion of metazoans (Cook and

Shergold, 1984; Wood and Zhuravlev, 2012). Consequently, organic-rich biomineral

composites of linguliform brachiopod shells possessed innovative mechanical functions,

providing competitive superiority and adaptation on Cambrian soft substrate as well as

reducing susceptibility to predation (Wood and Zhuravlev, 2012). Thus, even with the

physiological cost of biomineralization in demanding metabolism keeps increasing,

linguliforms keep their phosphate shells during dramatically oscillations of sea water

chemistry and temperature over 520 million years.

Given the long history of this subphylum, the possession of a phosphatic shell must have

had numerous advantages. The innovative columnar fabric can economically increase the

thickness and strength of the shell by the presence of numerous, stacked thinner laminae,

comparable with the laminated fabric seen in obolids (Cusack et al., 1999; Z.-F. Zhang et al.,

2016). Furthermore, the sandwich architecture also increases the toughness, flexibility and

the ability to resist crack propagation by filling the space between the compact lamellae with

organic material, comparable with the baculate fabric (Lévêque et al., 2004; Merkel et al.,

2009). Thus, the stacked sandwich model of the columnar shell possesses a greater

advantage of mechanical functions and adaptation with a superior combination of strength,

toughness and flexibility, resembling the colonnaded and reinforced concrete constructions

of our buildings. This type of more efficient and economic shell may also have been

responsible for the early diversity of linguliform brachiopods during the Cambrian explosion,

resulting in this group becoming a significant component of the Cambrian Evolutionary

Fauna (Bassett et al., 1999; Sepkoski, 1984; Zhang et al., 2008b; Z. Zhang et al., 2020; Z. L.

Zhang et al., 2021b). New data from nuclear magnetic resonance spectroscopy and X-ray

diffraction reveals that apatite in brachiopod shells is highly ordered and thermodynamically

stable crystalline that is more robust in the extremes of moisture, ambient osmotic potential

and temperature, unlike the poorly ordered crystal of vertebrate bone (Neary et al., 2011).

Evolution of stacked sandwich columns in early brachiopod clades

Evolutionary transformations have repeatedly modified the organo-phosphatic

architecture consisting of various aggregates of spherular apatite, held together by a

scaffolding of glycosaminoglycan complexes, fibrous proteinaceous struts and chitinous

platforms, in linguliform brachiopod shells since the early Cambrian (Cusack et al., 1999). As

one of the oldest type of brachiopod shell architectures, the columnar shell has previously

been regarded as an unique character of acrotretide brachiopods (Cusack et al., 1999;

Holmer, 1989), however, more recent discoveries of columnar shell in a diversity of early

Cambrian stem group brachiopods have revealed that the same biomineralization strategy is

also utilised in a wider range of early brachiopod and brachiopod-like groups (L. Holmer et

al., 2008; Holmer et al., 1996; Skovsted et al., 2010; Streng et al., 2007; Ushatinskaya and

Korovnikov, 2014; Zhang, 2018; Z. L. Zhang et al., 2021a). This increases the need for a

better understanding of the origin and adaptive modification of stacked sandwich columns in

the early lophophorate evolution.

The Eoobolidae is presently considered to be the oldest known linguliform brachiopods

(Z. L. Zhang et al., 2021a). The biomineralized columns in *Eoobolus incipiens* from

Cambrian Stage 3 probably represents an early and simple shell structure type with a poorly

developed secondary columnar layer (***Figure 4A***). The columns are relatively small with a

mean diameter of 1.8 μm ranging from 0.6 to 3.0 μm, and a mean height of 4.1 μm

(***Appendix 3—table 3***). Furthermore, the complete secondary layer is just composed of two

to three stacked columnar units, resulting in a shell thickness of about 30 μm. Such simple

shell structure is also developed in slightly younger *Latusobolus xiaoyangbaensis* gen. et sp.

nov. (***Figure 1***), but with a slightly higher column of about 6.2 μm. From Cambrian Age 4,

the unit number of layers increases rapidly in the Eoobolidae, growing as many as 10

stacked columnar units in *Eoobolus acutulus* sp. nov. (***Figure 2***), and as many as 10 in

Eoobolus? aff. *priscus*, and the shell thickness increases to 50 μm. However, the size of

individual columns keeps within a very stable range, around 4 μm in height and 2 μm in

diameter. Compared with Eoobolidae, the Lingulellotretidae, demonstrates a more

developed columnar shell, which has a relatively larger number of columnar units up to 20,

effectively increasing the shell thickness to 70 μm. The column size is very similar in both

Lingulellotreta malongensis and *L. ergalievi* (**Figure 4B** and **D**), and matches that of

contemporaneous *Eoobolus*. But the slightly younger *L. ergalievi* has more columnar units,

resulting in a thicker shell than the former.

Among all early Cambrian linguliforms with columnar shells, the acrotretides have

developed the most complex shell structure (**Figure 4F–J**). Furthermore, the successive

fossil record in Cambrian unveiled an obvious pattern of the increasing growth of stacked

columns in acrotretides from a very simple type in *Palaeotreta shannanensis* like *E. incipiens*

to slightly developed *Palaeotreta zhujiahensis* like *L. malongensis*, and to the most

developed *Eohadrotreta zhenbaensis* onward (**Figure 6**). The diameter of column increases

about two times in acrotretides from linguliforms, whereas the height of column increases to

10 μm in *Eohadrotreta zhenbaensis* and 29 μm in *Hadrotreta primaeva*, which is about 10

times as high as in *Eoobolus variabilis*. The number of columnar units has also increased to

30, which collectively increased the shell thickness to a maximal value of more than 300 μm.

The homology and continuity of the columnar shell in early linguliforms outlines a clear

picture, most likely representing an evolutionarily continuous transformation between

Eoobolidae, Lingulellotretidae and Acrotretida, which represent the major components of

early Cambrian benthic communities (Chen et al., 2021; Claybourn et al., 2020; Topper et al.,

2015; Zhang et al., 2008a; Zhang, 2018; Z. L. Zhang et al., 2020a). However, the possible

occurrence of this shell fabric within the family Obolidae cannot be discussed here, as

detailed information of the possible columnar shell structures in *Kyrshabaktella* and

Experilingula are too poorly known (Cusack et al., 1999; Streng et al., 2007; Williams, 1977).

The small size and simple patterns of stacked columnar shells remain very stable in the

Eoobolidae since its origin, which likely limits both the shell thickness and overall body size,

with ventral and dorsal valves of the family below 6 mm until Miaolingian (middle Cambrian).

Stacked sandwich columns are a character of the *Lingulellotreta* shell structure as well;

however more columnar units are developed to slightly increase the shell thickness and

species reach body size twice that of Eoobolidae (Zhang et al., 2007). A continuous

transformation of anatomic features can be deduced from the growth of columns between

the two clades. Firstly, the stacked columns are markedly developed at the pseudointerarea

area of ventral valve, which causes the great elevation of the pseudointerarea above the

shell floor (**Figure 2A**). It leaves a large amount of space for the posterior extension of the

digestive system, which are well protected by the covering mineralized ventral

pseudointerarea. This is evidenced by the discovery of a curved gut under the

pseudointerarea in the soft tissue preserved Chengjiang Lagerstätte (Zhang et al., 2007).

Secondly, with the continuous growth of the ventral pseudointerarea from both lateral sides,

the opening obolide-like pedicle groove is sealed off, resulting in a unique pedicle opening

exclusively observed in the lingulellotretid brachiopods. Thus, the pedicle protruding opening

between the ventral and dorsal valves of Lingulida is transformed to a new body plan where

the pedicle opening is restricted in ventral valve. It supports the scenario that the columnar

shell structure is monophyletic in at least Linguloidea, and that the younger lingulellotretid

columnar shell was derived from an *Eoobolus*-like ancestor during late Cambrian Age 3.

In another evolutionary direction, acrotretide brachiopods fully utilise the columnar fabric

including the derived camerate fabric as shell structure in the whole clade (Streng and

Holmer, 2006). The similarity and gradually evolutionary transformations of columns from

simple forms in lingulides to complex types in acrotretides suggests the stacked sandwich

columns did not evolve independently in actrotretides. In terms of the derived camerate

fabric, more mineralized material is utilised compared to its precursor, the columnar fabric

(Streng et al., 2007). The column size, including height and diameter, and the number of

stacked columnar units are constantly increasing to about 10 to 20 times greater in

acrotretides than in *E. incipiens and L. xiaoyangbaensis*, since late Cambrian Age 3 (**Figure**

6). Despite the increase in size and number, the whole body size of acrotretides is restricted

in millimetre size (Holmer, 1989; Williams et al., 2000). A continuous transformation of

anatomic features and shell structure functions can be deduced from the evolutionary growth

of columns in early acrotretides. Firstly, the stacked columns were markedly developed at

the pseudointerarea area of the ventral valve, which resulted in a greater elevation of the

pseudointerarea above the valve floor, compared to that of *Lingulellotreta*. Two

transformations subsequently changed an obolid-like flat ventral valve (*Palaeotreta*

shannanensis) to a cap shape vale (*P. zhujiahensis*) (Z. L. Zhang et al., 2020b), and a

conical shape (*Eohadrotreta zhenbaensis*) (Zhang et al., 2018), and eventually to a tubular

shape (*Acrotreta*) (Holmer and Popov, 1994). During this time, dorsal valves constantly kept

a flat shape. The obolid-like ventral pseudointerarea changed from orthocline to catacline

and eventually to procline with strongly reduced propareas, while ventral muscular system

moved to the elevated posterior floor, resulting in forming the new apical process (Popov,

1992; Zhang et al., 2018). With the increasing growth of the individual columns, the thick

organo-phosphatic shell with increased strength might provide more mechanical support to

the conical or tubular valve in a fluid environment.

Based on the similar ontogenetic transformation with the increasing growth of the shell in

Lingulellotretidae and Acrotretida, the forming of the ventral pedicle foramen is very likely

homologous, modified from a lingulide-like pedicle groove between the two valves.

Furthermore, the similarity and continuity in increasing the number and size of the stacked

columns suggested that columnar shell structure is synapomorphy in Linguloidea and

Acrotretida. However, the phylogenetic puzzle of whether the columnar fabric is paraphyletic

with the baculate fabric in Linguliformea, or even in Lingulata hangs on two pieces of

important fossil evidence. The shell structure of the earliest linguliform brachiopods needs to

be discovered on more well-preserved fossils globally. Moreover, the shell structure in widely

distributed early Cambrian stem group brachiopods needs to be scrutinized comprehensively.

The complex shell in stem group taxa *Micrina* and *Mickwitzia*, demonstrating advanced

columnar features in specimens yonger than *Eoobolus incipiens*, might reveal the

plesiomorphy of the columnar shell in Linguliformea (L. E. Holmer et al., 2008; Skovsted and

Holmer, 2003; Williams and Holmer, 2002b). But this assumption depends on future work

and whether the columnar shell is preserved in older stem groups. In another scenario,

baculate, laminated and columnar fabric might have originated independently from an

unmineralized ancestor like the agglutinated *Yuganotheca* during the early Cambrian

(Cusack et al., 1999; Zhang et al., 2014). No matter what scenario is true in nature, the

origin of the innovative columnar shell with a stacked sandwich model has played a

significant role in the evolution of linguliform brachiopods. The evolutionary growth of

stacked columns matches the increasing diversity of phosphatic-shelled brachiopods during

the Cambrian explosion. Among them, the micromorphic acrotretides demonstrated the

superb application of columnar shell combined with its innovative conical shape and possible

exploitation of secondary tiering niches (Topper et al., 2015; Wang et al., 2012; Zhang et al.,

2008a, 2018; Z. L. Zhang et al., 2021b), having flourished during the Great Ordovician

Biodiversification Event, thriving and playing an important role in marine benthic

communities for more than 100 million years (Bassett et al., 1999; Brunton et al., 2001;

Congreve et al., 2021; Harper et al., 2017b, 2015; Z. L. Zhang et al., 2020b). The fitness of

the diminutive body size of acrotretides is likely a trade-off between the increasing metabolic

demand of phosphate biomineralization after the Cambrian phosphogenic event and the

increased chance of evolutionary survival and adaptation of producing a high mechanical

protect skeleton for protection in the shallow water environment (Cook and Shergold, 1984;

Garbelli et al., 2017; Lévêque et al., 2004; Neary et al., 2011; Simonet Roda, 2021; Wood

and Zhuravlev, 2012).Their relatively large surface/volume ratio mechanically requires a

strong support with the composition of stacked sandwich columns and a relatively lower

density of the shell by organic biomineralized material for the secondary tiering life. Such

adaptive innovations may account for the flourish of phosphatic-shelled acrotretides in the

latter half of the Cambrian, continuing to the Great Ordovician Biodiversification Event,

thriving and playing an important role in marine benthic communities for more than 100

million years

Material and Methods

The brachiopod material studied here was collected from the Cambrian Series 2 Xihaoping

Member of the Dengying Formation and the Shuijingtuo Formation at the Xiaoyangba

section of southern Shaanxi (Z. L. Zhang et al., 2021a), and the Shuijingtuo Formation at the

Aijiahe section and Wangjiaping section of western Hubei (Z. L. Zhang et al., 2016). All

specimens are recovered through maceration of limestones by acetic acid (~10%) in lab,

and deposited in the Early Life Institute (ELI), Northwest University, China. Selected

specimens were coated and studied further using Fei Quanta 400-FEG SEM at Northwest

University, Zeiss Supra 35 VP field emission at Uppsala University and JEOL JSM 7100F-

FESEM at Macquarie University. Measurements of length, width and angle of different parts

of *Latusobolus xiaoyangbaensis* gen. et sp. nov. and *Eoobolus acutulus* sp. nov. are

performed on SEM images of well-preserved specimens by TpsDig2 v. 2.16.

Acknowledgements

We would like to thank Prof. G. Brock, Y. Cai and C.Y. Cai for insightful discussion, and Q.C. Feng

and J.P. Zhai for sample preparation in Xi'an. Thanks to S. Lindsay and C. Shen at Microscopy Unit at

Macquarie University, M. Streng at Uppsala University and Y.L. Pang at Northwest University for

assistance with SEM imaging.

Additional information

Funding

This research has been supported by the National Key R&D Program of China (grant no.

2022YFF0802700), Chinese Academy of Sciences (grant no. 202200020), National Natural Science

Foundation of China (grant nos.41720104002, 42072003), Swedish Research Council (VR Project no.

2017-05183, 2018-03390, 2021-04295) and Zhongjian Yang Scholarship from the Department of

Geology, Northwest University, Xi'an.

Authors contributions

Zhiliang Zhang, Conceptualization, Investigation, Methodology, Resources, Visualization, Validation,

Funding acquisition, Writing – original draft, Writing – review and editing; Zhifei Zhang, Investigation,

Resources, Funding acquisition, Writing – review and editing; Lars Holmer, Investigation, Writing –

review and editing; Timothy Topper, Investigation, Writing – review and editing; Bing Pan, Writing –

review and editing; Guoxiang Li, Investigation, Writing – review and editing.

Competing interests. The authors declare no competing interests.

References

Addadi L, Weiner S. 2014. Biomineralization: mineral formation by organisms. *Phys Scr* **89**:098003.

doi:10.1088/0031-8949/89/9/098003

Balthasar U, Cusack M. 2015. Aragonite-calcite seas—Quantifying the gray area. *Geology* **43**:99–102.

doi:10.1130/G36293.1

Bassett MG, Popov LE, Holmer LE. 1999. Organophosphatic brachiopods:Patterns of

biodiversification and extinction in the Early Palaeozoic. *Geobios* **32**:145–163.

doi:10.1016/S0016-6995(99)80026-6

Bicknell RDC, Paterson JR. 2018. Reappraising the early evidence of durophagy and drilling predation

in the fossil record: implications for escalation and the Cambrian Explosion: Reappraising the

early evidence of predation. *Biol Rev* **93**:754–784. doi:10.1111/brv.12365

Brasier MD. 1990. Phosphogenic events and skeletal preservation across the Precambrian-Cambrian

boundary interval. *Geological Society, London, Special Publications* **52**:289–303.

Briggs DEG. 2015. The Cambrian explosion. *Current Biology* **25**:R864–R868.

doi:10.1016/j.cub.2015.04.047

Brunton H, Cocks LRM, Long SL. 2001. Brachiopods. CRC Press.

Buatois LA, Mángano MG, Minter NJ, Zhou K, Wisshak M, Wilson MA, Olea RA. 2020. Quantifying

ecospace utilization and ecosystem engineering during the early Phanerozoic—The role of

bioturbation and bioerosion. *Science Advances* **6**:eabb0618. doi:10.1126/sciadv.abb0618

Budd GE, Jackson ISC. 2016. Ecological innovations in the Cambrian and the origins of the crown

group phyla. *Phil Trans R Soc B* **371**:20150287. doi:10.1098/rstb.2015.0287

Carlson SJ. 2016. The evolution of Brachiopoda. *Annual Review of Earth and Planetary Sciences*

44:409–438.

Chen F, Brock GA, Zhang Z, Laing B, Ren X, Zhang Z. 2021. Brachiopod-dominated communities and

depositional environment of the Guanshan Konservat-Lagersttte, eastern Yunnan, China.

Journal of the Geological Society **178**. doi:10.1144/jgs2020-043

Chen F, Topper TP, Skovsted CB, Strotz LC, Shen J, Zhang Z. 2022. Cambrian ecological complexities:

Perspectives from the earliest brachiopod – supported benthic communities in the early

Cambrian Guanshan Lagerstätte. *Gondwana Research* **107**:30–41.

doi:10.1016/j.gr.2022.02.008

Claybourn TM, Skovsted CB, Holmer LE, Pan B, Myrow PM, Topper TP, Brock GA. 2020. Brachiopods

from the Byrd Group (Cambrian Series 2, Stage 4) Central Transantarctic Mountains, East

Antarctica: biostratigraphy, phylogeny and systematics. *Papers in Palaeontology* **n/a**.

doi:10.1002/spp2.1295

Congreve CR, Patzkowsky ME, Wagner PJ. 2021. An early burst in brachiopod evolution

corresponding with significant climatic shifts during the Great Ordovician Biodiversification

Event. *Proceedings of the Royal Society B: Biological Sciences* **288**:8.

doi:doi.org/10.1098/rspb.2021.1450

Cook PJ, Shergold JH. 1984. Phosphorus, phosphorites and skeletal evolution at the Precambrian—

Cambrian boundary. *Nature* **308**:231–236. doi:10.1038/308231a0

Cuif J-P, Dauphin Y, Sorauf JE. 2010. Biominerals and Fossils Through Time. Cambridge University

Press.

Cusack M, Chung P, Dauphin Y. 2010. Brachiopod primary layer crystallography and nanostructure.

Palaeontology **84**:99–105.

Cusack M, Curry G, Clegg H, Abbott G. 1992. An intracrystalline chromoprotein from red brachiopod

shells: Implications for the process of biomineralization. *Comparative Biochemistry and*

Physiology Part B: Comparative Biochemistry **102**:93–95. doi:10.1016/0305-0491(92)90278-Y

Cusack M, Williams A. 2007. Chemicostructural diversity of the brachiopod shell. 2396-2521 in

Selden, PA. *Treatise on Invertebrate Paleontology, Part H, Brachiopoda, revised* **6**.

Cusack M, Williams A, Buckman JO. 1999. Chemico-structural evolution of linguloid brachiopod

shells. *Palaeontology* **42**:799–840. doi:10.1111/1475-4983.00098

Duan X, Betts MJ, Holmer LE, Chen Y, Liu F, Liang Y, Zhang Z. 2021. Early Cambrian (Stage 4)

brachiopods from the Shipai Formation in the Three Gorges area of South China. *Journal of*

Paleontology **95**:497–526. doi:10.1017/jpa.2020.117

Erwin DH. 2020. The origin of animal body plans: a view from fossil evidence and the regulatory

genome. *Development* **147**:dev182899. doi:10.1242/dev.182899

Erwin DH. 2015. Early metazoan life: divergence, environment and ecology. *Phil Trans R Soc B*

370:20150036. doi:10.1098/rstb.2015.0036

Garbelli C, Angiolini L, Shen S. 2017. Biomineralization and global change: A new perspective for

understanding the end-Permian extinction. *Geology* **45**:19–22. doi:10.1130/G38430.1

Golub EE. 2011. Biomineralization and matrix vesicles in biology and pathology. *Semin*

Immunopathol **33**:409–417. doi:10.1007/s00281-010-0230-z

Goto R, Takano T, Seike K, Yamashita M, Paulay G, Rodgers KS, Hunter CL, Tongkerd P, Sato S, Hong

J-S, Endo K. 2022. Stasis and diversity in living fossils: Species delimitation and evolution of

lingulid brachiopods. *Molecular Phylogenetics and Evolution* **175**:107460.

doi:10.1016/j.ympev.2022.107460

Griesshaber E, Schmahl WW, Neuser R, Pettke T, Blum M, Mutterlose J, Brand U. 2007.

Crystallographic texture and microstructure of terebratulide brachiopod shell calcite: An

optimized materials design with hierarchical architecture. *American Mineralogist* **92**:722–

734. doi:10.2138/am.2007.2220

Harper DAT, Popov LE, Holmer LE. 2017a. Brachiopods: origin and early history. *Palaeontology*

60:609–631. doi:10.1111/pala.12307

Harper DAT, Popov LE, Holmer LE. 2017b. Brachiopods: origin and early history. *Palaeontology*

60:609–631. doi:10.1111/pala.12307

Harper DAT, Stouge S, Christiansen JL, Topper TP, Alwmark C, Richoz S, Ahlberg P. 2021. Early

Cambrian brachiopod-dominated shell concentrations from North-East Greenland:

Environmental and taphonomic implications. *Global and Planetary Change* **204**:103560.

doi:10.1016/j.gloplacha.2021.103560

Harper DAT, Zhan R-B, Jin J. 2015. The Great Ordovician Biodiversification Event: Reviewing two

decades of research on diversity's big bang illustrated by mainly brachiopod data.

Palaeoworld, Geologic and biotic events and their relationships during the Early to Middle

Paleozoic **24**:75–85. doi:10.1016/j.palwor.2015.03.003

Holmer L, Popov L, Streng M. 2008. Organophosphatic stem group brachiopods - implications for the
phylogeny of the Subphylum Linguliformea. *Fossils and Strata* **54**:3–11.

Holmer LE. 1989. Middle Ordovician phosphatic inarticulate brachiopods from Vastergotland and
Dalarna, Sweden. *Fossils and Strata* **26**:1–172.

Holmer LE, Popov LE. 1994. Revision of the type species of Acrotreta and related lingulate
brachiopods. *Journal of Paleontology* **68**:433–450. doi:10.1017/S002233600002583X

Holmer LE, Popov LE, Wrona R. 1996. Early Cambrian lingulate brachiopods from glacial erratics of
King George Island, (South Shetland Islands), Antarctica. *Palaeontologia Polonica* **55**:37–50.

Holmer LE, Skovsted CB, Brock GA, Valentine JL, Paterson JR. 2008. The Early Cambrian tommotiid
Micrina, a sessile bivalved stem group brachiopod. *Biology Letters* **4**:724–728.
doi:10.1098/rsbl.2008.0277

Holmer LE, Skovsted CB, Williams A. 2002. A Stem Group Brachiopod From The Lower Cambrian:
Support For A Micrina (Halkieriid) Ancestry. *Palaeontology* **45**:875–882. doi:10.1111/1475-
4983.00265

Kallaste T, Kurvits T, Nemliher J, Puura I. 2004. Apatite varieties in the shell of the Cambrian lingulate
brachiopod Obolus apollinis Eichwald. *Proceedings of the Estonian Academy of Sciences*
Geology **53**:246. doi:10.3176/geol.2004.4.02

Leadbeater BSC, Riding R. 1986. Biomineralization in lower plants and animals. *The systematics*
association 1–401.

Lévêque I, Cusack M, Davis SA, Mann S. 2004. Promotion of Fluorapatite Crystallization by Soluble-
Matrix Proteins from Lingula Anatina Shells. *Angewandte Chemie International Edition*
43:885–888. doi:10.1002/anie.200353115

Li L, Skovsted CB, Topper TP. 2022. Deep origin of the crossed-lamellar microstructure in early
Cambrian molluscs. *Palaeontology* **65**:e12620. doi:10.1111/pala.12620

Liang Y, Strotz LC, Topper TP, Holmer LE, Budd GE, Chen Y, Fang R, Hu Y, Zhang Z. 2023. Evolutionary
contingency in lingulid brachiopods across mass extinctions. *Current Biology*.
doi:10.1016/j.cub.2023.02.038

Lowenstam HA, Weiner S. 1989. On biomineralization. New York: Oxford University Press.

Luo Y-J, Takeuchi T, Koyanagi R, Yamada L, Kanda M, Khalturina M, Fujie M, Yamasaki S, Endo K,
Satoh N. 2015. The Lingula genome provides insights into brachiopod evolution and the
origin of phosphate biomineralization. *Nat Commun* **6**:8301. doi:10.1038/ncomms9301

McClean AE. 1988. Epithelial moulds from some Upper Ordovician acrotretide brachiopods of Ireland
- McCLEAN - 1988 - Lethaia - Wiley Online Library. *Lethaia* **21**:43–50.

Merkel C, Deuschle J, Griesshaber E, Enders S, Steinhauser E, Hochleitner R, Brand U, Schmahl WW.
2009. Mechanical properties of modern calcite- (Mergerlia truncata) and phosphate-shelled
brachiopods (Discradisca stella and Lingula anatina) determined by nanoindentation. *Journal*
of Structural Biology **168**:396–408. doi:10.1016/j.jsb.2009.08.014

Murdock DJE. 2020. The 'biomineralization toolkit' and the origin of animal skeletons. *Biological*
Reviews **95**:1372–1392. doi:10.1111/brv.12614

Neary MT, Reid DG, Mason MJ, Friščić T, Duer MJ, Cusack M. 2011. Contrasts between organic
participation in apatite biomineralization in brachiopod shell and vertebrate bone identified
by nuclear magnetic resonance spectroscopy. *J R Soc Interface* **8**:282–288.
doi:10.1098/rsif.2010.0238

Pérez-Huerta A, Coronado I, Hegna TA. 2018. Understanding biomineralization in the fossil record.
Earth-Science Reviews **179**:95–122. doi:10.1016/j.earscirev.2018.02.015

Popov LE. 1992. The Cambrian Radiation of BrachiopodsOrigin and Early Evolution of the Metazoa.
Springer US. pp. 399–432.

Ruben JA, Bennett AA. 1987. The evolution of bone. *Evolution* **41**:1187–1197.

Schopf TJM. 1984. Rates of Evolution and the Notion of "Living Fossils." *Annual Review of Earth and*
Planetary Sciences **12**:245–292. doi:10.1146/annurev.ea.12.050184.001333

Sepkoski JJ. 1984. A kinetic model of Phanerozoic taxonomic diversity. III. Post-Paleozoic families and
mass extinctions. *Paleobiology* **10**:246–267.

Simonet Roda M del M. 2021. Pathways of biomineralization and microstructure evolution in
brachiopod shells (Text.PhDThesis). Ludwig-Maximilians-Universität München.

Simonet Roda M, Griesshaber E, Angiolini L, Rollion-Bard C, Harper EM, Bitner MA, Milner Garcia S,
Ye F, Henkel D, Häussermann V, Eisenhauer A, Gnägi H, Brand U, Logan A, Schmahl WW.
2021a. The architecture of Recent brachiopod shells: diversity of biocrystal and biopolymer
assemblages in rhynchonellide, terebratulide, thecideide and craniide shells. *Mar Biol* **169**:4.
doi:10.1007/s00227-021-03962-4

Simonet Roda M, Griesshaber E, Angiolini L, Rollion-Bard C, Harper EM, Bitner MA, Milner Garcia S,
Ye F, Henkel D, Häussermann V, Eisenhauer A, Gnägi H, Brand U, Logan A, Schmahl WW.
2021b. The architecture of Recent brachiopod shells: diversity of biocrystal and biopolymer
assemblages in rhynchonellide, terebratulide, thecideide and craniide shells. *Mar Biol* **169**:4.
doi:10.1007/s00227-021-03962-4

Skovsted CB, Brock GA, Paterson JR, Holmer LE, Budd GE. 2008. The scleritome of Eccentrotheca
from the Lower Cambrian of South Australia: Lophophorate affinities and implications for
tommotiid phylogeny. *Geol* **36**:171. doi:10.1130/G24385A.1

Skovsted CB, Holmer LE. 2003. The Early Cambrian (Botomian) stem group brachiopod Mickwitzia
from Northeast Greenland. *Acta Palaeontologica Polonica* 20.

Skovsted CB, Streng M, Knight I, Holmer LE. 2010. Setatella significans, a new name for mickwitziid
stem group brachiopods from the lower Cambrian of Greenland and Labrador. *GFF* **132**:117–
122. doi:10.1080/11035897.2010.490878

Streng M, Holmer L, Popov L, Budd G. 2007. Columnar shell structures in early linguloid brachiopods
- New data from the Middle Cambrian of Sweden. *Earth and Environmental Science
Transactions of the Royal Society of Edinburgh* **98**:221–232.
doi:10.1017/S1755691007079820

Streng M, Holmer LE. 2006. New and poorly known acrotretid brachiopods (Class Lingulata) from
the Cedaria-Crepicephalus zone (late Middle Cambrian) of the Great Basin, USA. *Geobios*
39:125–154. doi:10.1016/j.geobios.2004.10.004

Temereva E. 2022. First Modern Data on the Lophophore Nervous System in Adult Novocrania
anomala and a Current Assessment of Brachiopod Phylogeny. *Biology* **11**:406.
doi:10.3390/biology11030406

Topper TP, Holmer LE, Skovsted CB, Brock GA, Balthasar U, Larsson CM, Stolk SP, Harper DAT. 2013.
The Oldest Brachiopods from the Lower Cambrian of South Australia. *acpp* **58**:93–109.
doi:10.4202/app.2011.0146

Topper TP, Strotz LC, Holmer LE, Caron J-B. 2015. Survival on a soft seafloor: life strategies of
brachiopods from the Cambrian Burgess Shale. *Earth-Science Reviews* **151**:266–287.
doi:10.1016/j.earscirev.2015.10.015

Ushatinskaya GT. 2002. Genus Micrina (Small Shelly Problematics) from the Lower Cambrian of
South Australia: Morphology, Microstructures, and Possible Relation to Halkieriids.
Paleontologicheskii Zhurnal 20–21.

Ushatinskaya GT, Korovnikov IV. 2014. Revision of the Early-Middle Cambrian Lingulida (Brachiopoda)
from the Siberian Platform. *Paleontol J* **48**:26–40. doi:10.1134/S0031030114010158

Wang H, Zhang Z, Holmer LE, Hu S, Wang X, Li G. 2012. Peduncular attached secondary tiering
acrotretoid brachiopods from the Chengjiang fauna: Implications for the ecological
expansion of brachiopods during the Cambrian explosion. *Palaeogeography,
Palaeoclimatology, Palaeoecology* **323**:60–67. doi:10.1016/j.palaeo.2012.01.027

Weiner S. 2008. Biomineralization: A structural perspective. *Journal of Structural Biology*, JSB 50th
Anniversary Issue **163**:229–234. doi:10.1016/j.jsb.2008.02.001

Weiner S, Dove PM. 2003. An Overview of Biomineralization Processes and the Problem of the Vital
Effect. *Reviews in Mineralogy and Geochemistry* **54**:1–29. doi:10.2113/0540001

Williams A. 1977. Shell StructureTreatise on Invertebrate Palaeontology, Part H, Brachiopoda.
Lawrence: Geological Society of America, Boulder and University of Kansas. pp. 267–320.

Williams A, Brunton C, Mackinno I. 2000. MorphologyTreatise on Invertebrate Paleontology, Part H,
Brachiopoda (Revised). Lawrence: Geological Society of America, Boulder and University of
Kansas. pp. 321–440.

Williams A, Cusack M. 1999. Evolution of a Rhythmic Lamination in the Organophosphatic Shells of
Brachiopods. *Journal of Structural Biology* **126**:227–240. doi:10.1006/jsbi.1999.4117

Williams A, Cusack M, Mackay S. 1994. Collagenous Chitinophosphatic Shell of the Brachiopod
Lingula. *Philosophical Transactions: Biological Sciences* **346**:223–266.

Williams A, Holmer LE. 2002a. Shell Structure And Inferred Growth, Functions And Affinities Of The
Sclerites Of The Problematic Micrina. *Palaeontology* **45**:845–873. doi:10.1111/1475-
4983.00264

Williams A, Holmer LE. 2002b. Shell Structure And Inferred Growth, Functions And Affinities Of The
Sclerites Of The Problematic Micrina. *Palaeontology* **45**:845–873. doi:10.1111/1475-
4983.00264

Williams A, Holmer LE. 1992. Ornamentation and shell structure of acrotretoid brachiopods.
Palaeontology **35**:657–692.

Williams A, Holmer LE, Cusack M. 2004. Chemico-structure of the organophosphatic shells of
siphonotretide brachiopods. *Palaeontology* **47**:1313–1337. doi:10.1111/j.0031-
0239.2004.00404.x

Williams A, Mackay S, Cusack M. 1992. Structure of the organo-phosphatic shell of the brachiopod
Discina. *Philosophical Transactions of the Royal Society of London Series B: Biological
Sciences* **337**:83–104. doi:10.1098/rstb.1992.0086

Winrow P, Sutton MD. 2012. Epithelial cell moulds in acrotretoid brachiopods. *Historical Biology*
24:557–565. doi:10.1080/08912963.2012.657633

Wood R, Zhuravlev AYu. 2012. Escalation and ecological selectively of mineralogy in the Cambrian
Radiation of skeletons. *Earth-Science Reviews* **115**:249–261.
doi:10.1016/j.earscirev.2012.10.002

Ye F, Garbelli C, Shen S, Angiolini L. 2021. The shell fabric of Palaeozoic brachiopods: patterns and
trends. *Lethaia* **54**:419–439. doi:10.1111/let.12412

Yun H, Luo C, Chang C, Li L, Reitner J, Zhang X. 2022. Adaptive specialization of a unique sponge body
from the Cambrian Qingjiang biota. *Proceedings of the Royal Society B: Biological Sciences*
289:20220804. doi:10.1098/rspb.2022.0804

Yun H, Zhang X, Brock GA, Li L, Li G. 2021. Biomineralization of the Cambrian chancelloriids. *Geology*
49:623–628. doi:10.1130/G48428.1

Zhang X, Shu D. 2021. Current understanding on the Cambrian Explosion: questions and answers.
PalZ. doi:10.1007/s12542-021-00568-5

Zhang Z, Han J, Wang Y, Emig CC, Shu D. 2010. Epibionts on the lingulate brachiopod Diandongia
from the Early Cambrian Chengjiang Lagerstätte, South China. *Proceedings of the Royal*
Society B: Biological Sciences **277**:175–181. doi:10.1098/rspb.2009.0618

Zhang Z, Han J, Zhang X, Liu J, Guo J, Shu D. 2007. Note on the gut preserved in the Lower Cambrian
Lingulellotreta (Lingulata, Brachiopoda) from southern China. *Acta Zoologica* **88**:65–70.
doi:10.1111/j.1463-6395.2007.00252.x

Zhang Z, Robson SP, Emig C, Shu D. 2008a. Early Cambrian radiation of brachiopods: A perspective
from South China. *Gondwana Research*, Snowball Earth to Cambrian Explosion **14**:241–254.
doi:10.1016/j.gr.2007.08.001

Zhang Z, Robson SP, Emig C, Shu D. 2008b. Early Cambrian radiation of brachiopods: A perspective
from South China. *Gondwana Research*, Snowball Earth to Cambrian Explosion **14**:241–254.
doi:10.1016/j.gr.2007.08.001

Zhang Z-F, Li G-X, Holmer LE, Brock GA, Balthasar U, Skovsted CB, Fu D-J, Zhang X-L, Wang H-Z, Butler
A, Zhang Z-L, Cao C-Q, Han J, Liu J-N, Shu D-G. 2014. An early Cambrian agglutinated tubular
lophophorate with brachiopod characters. *Sci Rep* **4**:1–8. doi:10.1038/srep04682

Zhang Z-F, Zhang Z-L, Li G-X, Holmer LE. 2016. The Cambrian brachiopod fauna from the first-trilobite
age Shuijingtuo Formation in the Three Gorges area of China. *Palaeoworld* **25**:333–355.

Zhang Z, Strotz LC, Topper TP, Chen F, Chen Y, Liang Y, Zhang Zhiliang, Skovsted CB, Brock GA. 2020.
An encrusting kleptoparasite-host interaction from the early Cambrian. *Nat Commun*
11:2625. doi:10.1038/s41467-020-16332-3

Zhang ZL. 2018. Early Cambrian phosphatic-shelled brachiopods from South China (Doctoral
dissertation). Northwest University.

Zhang ZL, Chen FY, Zhang ZF. 2020a. The earliest phosphatic-shelled brachiopods from the
carbonates of South China-their diversification, ontogeny and distribution. *Earth Science*
Frontiers **27**:127–151. doi:10.13745/j.esf.sf.2020.6.4

Zhang Z. L., Ghobadi Pour M, Popov LE, Holmer LE, Chen F, Chen Y, Brock GA, Zhang Z. 2021a. The
oldest Cambrian trilobite – brachiopod association in South China. *Gondwana Research*
89:147–167. doi:10.1016/j.gr.2020.08.009

Zhang ZL, Holmer LE, Chen F, Brock GA. 2020b. Ontogeny and evolutionary significance of a new
acrotretide brachiopod genus from Cambrian Series 2 of South China. *Journal of Systematic*
Palaeontology **18**:1569–1588. doi:10.1080/14772019.2020.1794991

Zhang Z. L., Topper TP, Chen Y, Strotz LC, Chen F, Holmer LE, Brock GA, Zhang Z. 2021b. Go large or
go conical: allometric trajectory of an early Cambrian acrotretide brachiopod. *Palaeontology*
64:727–741. doi:10.1111/pala.12568

Zhang ZL, Zhang Z, Holmer LE. 2017. Studies on the shell ultrastructure and ontogeny of the oldest
acrotretid brachiopods from South China. *Acta Palaeontologica Sinica* **56**:483–503.

Zhang ZL, Zhang Z, Holmer LE, Chen F. 2018. Post-metamorphic allometry in the earliest acrotretoid
brachiopods from the lower Cambrian (Series 2) of South China, and its implications.
Palaeontology **61**:183–207. doi:10.1111/pala.12333

Zhang Z.L., Zhang Z, Ma J, Taylor PD, Strotz LC, Jacquet SM, Skovsted CB, Chen F, Han J, Brock GA.
2021. Fossil evidence unveils an early Cambrian origin for Bryozoa. *Nature* **599**:251–255.
doi:10.1038/s41586-021-04033-w

Zhang ZL, Zhang ZF, Wang H. 2016. Epithelial cell moulds preserved in the earliest acrotretid
brachiopods from the Cambrian (Series 2) of the Three Gorges area, China. *GFF* **138**:455–466.
doi:10.1080/11035897.2016.1143528

【Figure captions】

Figure 1. Shell ultrastructure of *Latusobolus xiaoyangbaensis* gen. et sp. nov. from

the Cambrian Series 2 Shuijingtuo formation in southern Shaanxi. **A-C**, ELI-XYB S5-

1 BS01. **A**, cross section of a ventral lateral margin, note post-metamorphosis

pustules by arrow and shell layers 1-7, box indicates area in **B**. **B**, enlarged view of

A, showing primary-secondary layer boundary by dashed line. **C,** enlargement of

shell layers 1-4 of **A**, note organic canals of columns (arrows), organic layer (tailed

arrows) between two stratiform lamellae by dashed lines. **D**, poorly phosphatised

columns of ventral valve, note organic canals by tailed arrows, ELI-XYB S5-1 BR06.

E, columns of dorsal valve, ELI-XYB S5-1 BS17. **F**, cross section of a ventral lateral

margin, show developed post-metamorphosis pustules, box indicates the area in **G**,

ELI-XYB S4-2 BO06. **G**, enlarged primary layer pustules and secondary layer

columns underneath. **H**, one layer of dorsal columnar shell with the exfoliation of

above primary layer, noting column canals on the layer surface by arrows, ELI-XYB

S4-2 BO08. **I**, apatite spherules of granule aggregations of ventral columnar shell,

note granule rods by arrow and organic membrane by tailed arrow, ELI-XYB S4-2

BO06. Scale bars: **A**, 50 μm; **B**, **E**, **G**, 20 μm; **C**, **H**, 10 μm; **D**, 5 μm; **F**, 100 μm; **I**, 1

μm.

Figure 2. Ultrastructure of ventral *Eoobolus acutulus* sp. nov. from the Cambrian

Series 2 Shuijingtuo formation in Three Gorges areas. **A-C**, ELI-AJH S05 BT12. **A**,

cross section of a ventral lateral margin, note stacked columnar units 2-11, box

indicates area in **B**. **B**, enlarged view of **A**, showing shell layers 2-11. **C**, enlarged

view of **B**, show thin organic layer (tailed arrow) between two stratiform lamellae by

dashed lines, the fusion of two columnar layers by arrow. **D-F**, ELI-AJH 8-2-3 BT02.

D, cross section of shell margin, box indicates area in **E**. **E**, different preservation

condition of columns. **F**, poorly phosphatised columns, note the opening of organic

canals along the organic lamellae by arrows, and space between two stratiform

laminae by tailed arrows. **G-H**, ELI-AJH 8-2-3 BT03. **G**, note organic canals on the

cross section and surface of columnar layer by arrows, and partly exfoliated primary

layer by tailed arrow. **H**, magnified columns composed of granule spherules with

canal in **G** by arrow. **I**, cross section of shell margin, box indicates area in (**J**), ELI-

AJH 8-2-3 BT04. **J**, enlarged short columns. **K**, imbricated columnar layers (arrows),

ELI-AJH S05 BT12. Scale bars: **A**, **E**, 50 µm; **B**, **I**, 20 µm; **C**, 5 µm; **D**, 200 µm; **F**, **G**,

J, **K**, 10 µm; **H**, 1 µm.

Figure 3. Shell ornamentation, ultrastructure and epithelial cell moulds of Cambrian

Series 2 brachiopods. **A**, post-metamorphic pustules of *Latusobolus*

xiaoyangbaensis gen. et sp. nov., ELI-XYB S4-3 AU11. **B-D**, *Eoobolus acutulus* sp.

nov. **B**, metamorphic hemispherical pits, ELI-AJH 8-2-2 Lin01. **C**, epithelial cell

moulds, note column openings on layer surface beneath by arrow, ELI-AJH S05 N31.

D, enlarged column openings on layer surface, ELI-WJP 7 CE05. **E-I**, *Eohadrotreta*

zhenbaensis. **E**, partly broken columns, note organic canals by arrows, ELI-AJH F36.

F, polygonal epithelial cell moulds on valve floor, dash lines indicate margin of one

epithelial cell, ELI-WJP 6 R79. **G-I**, ELI-AJH Acro 053. **G**, epithelial cell moulds on

median septum with columns between. **H**, enlarged view of **G**, note rudiment of

columns by tailed arrows and one column on epithelial cell margin by arrow. **I,** three

layers of epithelial cell moulds on cardinal muscles with columns between (arrow).

Scale bars: **A**, **D**, 50 µm; **B**, **E**, **F**, **H**, 10 µm; **C**, **G**, **I**, 20 µm.

Figure 4. Columnar layers of Cambrian Series 2 brachiopods. **A**, *Eoobolus incipiens*,

AJXM-267.5 DT12. **B-D**, *Lingulellotreta ergalievi*, ELI-AJH 8-2-3 CI11. **B**, cross

section of shell margin, box indicates area in **C**, note stacked columnar units 2-22,

and raised pseudointerarea (tailed arrow). **C**, enlarged view of thin organic layer

(tailed arrow) between two stratiform lamellae by dashed lines, the fusion of two

columnar units into one by arrow. **D**, imbricated growth pattern of columnar layers. **E**,

Palaeotreta zhujiahensis, note column openings (arrow) on eight successive

columnar units by tailed arrow, ELI-AJH 8-2-1 AE09. **F-J**, *Eohadrotreta zhenbaensis*.

F, relatively higher columns (ca. 20 µm), ELI-AJH 8-2-1 acro16. **G**, Apatite spherules

of granule aggregations in columnar layer, note column openings (arrows) on both

stratiform lamella surface, ELI-AJH S05 E18. **H**, cross section show column

openings on 4 successive units by tailed arrow, ELI-WJP 7 AB98. **I**, poorly

phosphatised columns (arrows), note openings of organic canals on stratiform

lamella by tailed arrows, ELI-AJH S05 I76. **J**, stacked columnar layers in imbricated

pattern, ELI-WJP 6 R47. Scale bars: **A**, **E**, **I**, 10 µm; **B**, 100 µm; **C**, **F**, 20 µm; **G**, 2

µm; **H**, 5 µm; **J**, 50 µm.

853 **Figure 5.** Biomineralization process of typical columnar layer by the conveyor-belt

854 system of phosphatic-shelled brachiopods. Abbreviation: D=Diameter; H=height;

855 T=Thickness. (modified from (Williams and Holmer, 1992; Z. L. Zhang et al., 2016))

Figure 6. The evolutionary growth of stacked columnar shells in early Eoobolidae

Eoobolus incipiens, *Latusobolus xiaoyangbaensis* gen. et sp. nov., *Eoobolus*

acutulus sp. nov., *Eoobolus variabilis*, *Eoobolus*? aff. *priscus*, Lingulellotretidae

Lingulellotreta malongensis, *L. ergelievi*, Acrotretida *Palaeotreta shannanensis*,

Palaeotreta zhujiahensis, *Eoohadrotreta zhenbaensis*, *Canthylotreta crista* and stem

group *Mickwitzia* cf. *occidens*. The height and diameter of column is from the

literature (Skovsted and Holmer, 2003; Streng et al., 2007; Streng and Holmer, 2006;

Ushatinskaya and Korovnikov, 2014; Z. L. Zhang et al., 2020a, 2020b, 2016).