Predicting drug response from single-cell expression profiles of tumours

AUTHORS
Simona Pellecchia[1,4,†], Gaetano Viscido[1,2,†], Melania Franchini[1,3], Gennaro Gambardella[1,#]

AFFILIATIONS
[1]Telethon Institute of Genetics and Medicine, Naples, Italy
[2]University of Naples Federico II, Department of Chemical, Materials and Industrial Engineering, Naples, Italy
[3]University of Naples Federico II, Department of Electrical Engineering and Information Technology, Naples, Italy
[4]Scuola Superiore Meridionale, Genomics and Experimental Medicine Program, Naples, Italy

[†]**These authors contributed equally to the work.**

[#]**CORRESPONDENCE: gambardella@tigem.it**

ABSTRACT
Drug response prediction at the single cell level is an emerging field of research that aims to improve the efficacy and precision of cancer treatments. Here, we introduce DREEP (Drug Response Estimation from single-cell Expression Profiles), a computational method that leverages publicly available pharmacogenomic screens and functional enrichment analysis to predict single cell drug sensitivity from transcriptomic data. We validated DREEP extensively *in vitro* using several independent single-cell datasets with over 200 cancer cell lines and showed its accuracy and robustness. Additionally, we also applied DREEP to molecularly barcoded breast cancer cells and identified drugs that can selectively target specific cell populations. DREEP provides an in-silico framework to prioritize drugs from single-cell transcriptional profiles of tumours and thus helps in designing personalized treatment strategies and accelerate drug repurposing studies. DREEP is available at https://github.com/gambalab/DREEP.

Introduction

Intra-tumour heterogeneity (ITH) is a recurrent issue in designing effective treatment strategies in clinical practice. It refers to heterogeneous cancer cell populations present in the same tumour that exhibit high cellular plasticity and phenotypic heterogeneity. Accumulating evidence from independent studies shows that ITH gives rise to rare sub-populations of malignant cells characterized by a different epigenetic and transcriptional state that renders them refractory to a variety of anticancer drugs, protecting the whole population from complete eradication (1–4).

Single-cell RNA sequencing (scRNA-seq) has now become a powerful tool to explore ITH at the transcriptional level (5) and thus offers an unprecedented chance to tackle it therapeutically (4, 6, 7). In addition, large-scale pharmacogenomic screens have been performed across hundreds of molecularly characterized cancer cell lines (8–14). The resulting data have provided a valuable resource to link drug sensitivity with genomic and transcriptomic features. Indeed, in the last few years, several machine-learning models have leveraged these datasets to identify molecular markers linked to *in vitro* drug sensitivity in order to predict drug response from -omics profiles of tumour biopsies (8, 15, 16). However, these approaches do not explicitly account for ITH with the result that

genomic and non-genomic biomarkers of drug response have been found for only a restricted number of small molecules. Nonetheless, expression-based biomarkers measured from bulk cell populations have emerged as the most powerful predictors of drug response *in vitro* for several cytotoxic and targeted drugs, with far more predictive power than other -omics profiles (8, 10). To overcome these limitations, a few recent studies like scDRUG (17, 18), scDEAL (19), beyondCell (20), chemCPA (21), and NCI-DOE (22) have attempted to leverage scRNA-seq and publicly available drug sensitivity profiles to predict drug response at the single cell level. However, these studies often exhibit certain drawbacks. They may lack comprehensive *in vitro* validation, fail to provide user-friendly software packages, rely on complex models requiring parameter fine-tuning before practical use, or are confined to predicting the efficacy of a limited number of drugs.

To address these challenges, we introduce DREEP (Drug Response Estimation from single-cell Expression Profiles), a novel bioinformatics tool designed to augment the precision and efficacy of therapeutic strategies by addressing the innate heterogeneity of cancer cell populations. DREEP's primary function is to assess the complexity of tumour therapeutics and recommend specific drugs tailored to target distinct cell subpopulations. DREEP can predict a cell's vulnerability to more than 2,000 drugs by simplifying the whole process with enrichment analysis, eliminating the need for complex parameter adjustments and thus making this type of analysis accessible to a broad audience.

We validate DREEP performance extensively using *in vitro* drug response data of over 200 cancer cell lines covering 22 distinct cancer types from publicly available single-cell and drug viability datasets. We show that our method can effectively identify recurring drug vulnerabilities across cells of diverse tumour types and establishes a clear relationship between drug response and the functional status of the cells. We show that DREEP can detect changes in drug sensitivity over time in the breast cancer cell line MCF7 and predict the emergence of drug resistance. Additionally, using molecular barcoded MDA-MB-468 cells (4), we show that our method can recognize innate drug-tolerant cell subpopulations and predict drug co-treatments that will inhibit all of them. Finally, we demonstrate that DREEP can accurately predict drug sensitivity within heterogeneous cancer cell populations of patient-derived cultures (PDCs) containing a mix of primary and metastatic cells derived from individuals with head and neck squamous cell carcinomas (23). In conclusion, our results demonstrate that it is possible to detect differences in drug response among individual cells within the same tumour. DREEP is implemented as an open-source R package, making it easily accessible and user-friendly for the broader scientific community. It is publicly available on GitHub at the following address https://github.com/gambalab/DREEP.

RESULTS

1. Drug Response Estimation from single-cell Expression Profiles.

DREEP (Drug Response Estimation from single-cell Expression Profiles) is a bioinformatics tool that leverages results from large-scale cell-line viability screens and enrichment analysis to predict drug vulnerability from the transcriptional state of a cell. It requires a *pre-defined* collection of Genomic Profiles of Drug Sensitivity (GPDS); these are lists of genes ranked according to their contribution in correctly predicting the effect of

a small molecule. To build GPDS, we integrated several publicly available RNA-seq
datasets and cell-line viability screens, including the following datasets: GDSC (8), CTRP2
(13, 24) and PRISM (25) (Figure 1A).

GPDS were computed by correlating the *in vitro* response to a small molecule from the
above studies with the expression of all messenger RNA (mRNA) across all the cancer cell
lines for which the drug potency was measured (Figure 1A and methods). In these studies,
RNA-seq represents the basal gene expression of a cell line, while the potency of a small
molecule in a cell line is evaluated as the Area Under the Curve (AUC) of the
corresponding dose-response curve. The AUC value reflects cell growth inhibition after 72
hours of treatment, where lower values indicate sensitivity while higher values imply
resistance to the tested drug. Thus, a gene whose expression positively correlates with the
AUC of a drug across many cell lines can be considered predictive of drug resistance (i.e.,
the more expressed the gene, the higher the concentration needed to inhibit cell growth)
(24). *Vice versa*, a negatively correlated gene can be considered predictive of drug
sensitivity (*i.e.* the more expressed the gene, the lower the concentration needed to inhibit
cell growth) (24).

Using this approach, we generated a collection of 2,434 GPDS signatures (i.e. 478 from
GDSC, 511 from CTRP, and 1,445 from PRISM) and used them to predict the single-cell-
level drug response with the strategy depicted in Figure 1B. Briefly, single cell RNA
sequencing data from a sample was first normalized with the gf-icf (Gene Frequency –
Inverse Cell Frequency) pipeline to rank genes in each cell according to their relevance
(26–28). Then, for each cell we used the most relevant genes as input for Gene Set
Enrichment Analysis (GSEA) (29) against each GPDS ranked-list. Since resistance
biomarkers are at the top and sensitivity biomarkers are at the bottom of GPDS ranked lists,
a positive value of the enrichment score (ES) returned by DREEP implies that the cell
expresses genes associated with drug resistance. *Vice versa*, a negative ES indicates the
cell expresses genes correlated with drug sensitivity. At the end of this process, it is thus
possible to estimate the overall effect of each drug on a cell population by estimating the
median DREEP enrichment score on that population and compute the percentage of cells
sensitive to the drug by simply counting the number of cells in the sample that are
associated with a significant ES less than zero (methods).

We found 90 small molecules shared among GDSC, CTRP2, and PRISM datasets. To
assess the consistency of GPDS ranked lists, we calculated Spearman Correlation
Coefficients (SCC) between GPDS representations of the same drug but constructed from
distinct drug viability datasets. Supplementary Figure 01 shows a substantial similarity in
GPDS rankings of the same drug even if derived from two distinct datasets, surpassing
random comparisons by several folds (methods).

2. Estimation of the predictive performance of DREEP.
To validate the GPDS ranked list and demonstrate the reliability of our method as well as
measure its predictive performance on *in vitro* data, we applied it to the Kinker et al. dataset
(30) comprising 53,513 single cell transcriptional profiles from 198 tumorigenic cell lines
and spanning 22 distinct cancer types. To evaluate DREEP performance, we first converted
predictions from the single-cell level to the cell-line level by computing the median
enrichment score for each drug (i.e. GPDS) across the cells of each individual cell line.

Then, to minimize potential confounders, we evaluated the performance of the DREEP
method for each sensitivity dataset independently (Methods). We also searched for the
optimal number of relevant genes to use for enrichment analysis and predict the sensitivity
of a cell (*i.e.*, 50, 100, 250, 500, and 1,000 genes). Figure 1C shows the method's overall
performance in terms of Precision-Recall and ROC (Receiver Operating Characteristic)
curves across 198 cancer cell lines for each drug sensitivity dataset independently (i.e.,
CTRP2, GDSC and PRISM). In this analysis, predicted drug/cell-line pairs are ranked
according to the median enrichment score estimated by DREEP and the performance is
evaluated against the corresponding gold standard (methods). Figure 1D reports the ROC
curve's AUC (Area Under the Curve) for all drugs and all cell lines individually. Next, to
investigate whether DREEP exhibited a prediction bias toward a class of drugs
characterized by specific Mechanism of Action (MoA), we categorized the ROC curve's
AUC of each drug in Figure 1D (left panels) into two distinct groups: (i) Accurately
Predicted (defined as ROC-AUC $\geq$ 0.75) and (ii) Erroneously Predicted drugs (defined as
ROC-AUC $\leq$ 0.5, indicating random performance). This analysis yielded a total of 639
drugs whose efficacy was accurately predicted (209 from GDSC, 147 from CTRP2, and
282 from PRISM; Supplementary Table 01) and 135 drugs whose efficacy was erroneously
predicted by DREEP (21 from GDSC, 23 from CTRP2, and 91 from PRISM;
Supplementary Table 01). Notably, accurately predicted drugs exhibited a significant
enrichment (31) (FDR < 10%) across a broad spectrum of MoAs (methods), spanning
various biological mechanisms (Supplementary Table 02). In contrast, the 135 drugs
predicted with low accuracy showed enrichment for a limited number of MoAs but
overlapped with the MoAs of accurately predicted drugs (Supplementary Table 02), except
for prostanoid and glucocorticoid agonist molecules. This analysis indicates that DREEP
did not have a prediction bias toward a specific class of drugs, but it proved incapable of
predicting the effects of these 135 small molecules, leading us to exclude them from the
final version of the tool. Lastly, to assess DREEP's ability to generalize across different
cancer types, we grouped the ROC curve's AUC values for individual cell lines (as reported
in Figure 1D, right panels) according to their respective cancer types. Supplementary
Figure 02 illustrates that DREEP does not reveal any significant performance drop
associated with specific cancer types except for Neuroblastoma where average AUC is
lower than in other cancer types. Finally, we conducted an additional assessment of
DREEP's predictive performance where we constructed GPDS ranked lists using the drug's
IC50 values instead of the AUC metric. As depicted in Supplementary Figure 03, DREEP's
performance noticeably decreased across all three drug viability datasets when employing
this alternate approach.

To test DREEP's performance on an additional independent dataset, we applied it to a
single-cell breast cancer dataset (5). This dataset comprises 35,276 individual cells from
32 breast cancer cell lines covering all the main breast cancer tumour subtypes
(Luminal/Her2-positive/Basal Like). Figure 1E,F show DREEP's performance in this
additional dataset.

Next, we tested the prediction performance of the DREEP method on both pan-cancer and
breast cancer datasets using absolute gene expression levels instead of gf-icf normalized
data. We observed a drastic reduction of the predictive performance of the method
(Supplementary Figure 04) using the absolute gene expression level to estimate its
relevance in a cell, showing the importance of data pre-processing prior to the application

of a drug prediction algorithm. As a good compromise between the required computational
time and prediction accuracy, we decided to use N = 500 genes as a default value for the
DREEP sensitivity predictions in all subsequent analyses in the manuscript.

Finally, we used both the single-cell pan-cancer and breast cancer datasets to conduct a
comparative analysis of DREEP's performance with four other widely used single-cell
drug prediction methods, including scDRUG (17, 18), scDEAL (19), beyondCell (20)
(methods). We evaluated the performance of these methods based on the shared drugs, as
depicted in Supplementary Figure 05A. As demonstrated in Supplementary Figures 05B-
F, DREEP consistently outperformed the other methods across both datasets.

Taken together, these results collectively demonstrate the validity of our GPDS ranked lists
for predicting the sensitivity of cells to drugs. Furthermore, our comprehensive evaluation
showcases that DREEP consistently outperforms random predictions across various
scenarios and outperforms other state-of-the-art methods.

198 3. The DREEP method reconstructs the therapeutic landscape of the pan-cancer
199 dataset.
Next, we used the DREEP method to reconstruct the therapeutic landscape of the 22 cancer
types in the cell line pan-cancer dataset of Kinker et al. (30). As shown in Figure 2A,B (and
Supplementary Table 03), we first used DREEP to convert each cell line into a list of small
molecules predicted to be effective for at least 90% of profiled cells. We then grouped the
cells and the drugs into four clusters each, so that cell lines that respond similarly across
the tested drugs are grouped together, and analogously drugs that have an effect on the
same cell lines are grouped together (i.e., cell line clusters C1, C2 C3, C4 and drug clusters
D1, D2, D3 and D4).

These analyses revealed interesting associations between cancer vulnerabilities and drug
response, highlighting numerous cell lines and cancer types predicted to be sensitive to
small molecules with shared mechanisms of action (MoA) (31). Indeed, as shown in Figure
2A,B (and Supplementary Table 03 and Supplementary Table 04), cell lines in clusters C1
and C2 comprised 87% (13 out of 15) of head and neck cell lines, 100% of colorectal cell
lines (10 out of 10), 83% of bladder cell lines (5 out of 6), 73% of breast cell lines (11 out
of 15), 73% of pancreatic cell lines (8 out of 11), and 33% of lung cancer cell lines (13 out
of 39). These cancers all have in common Epidermal Grow Factor (EGF) receptor
overexpression/amplification, such as EGFR and ERBB2 (32). Indeed, this is one of the
major alterations in colorectal (CRC, 60-80%) (33), lung (NSCLC, 40-80%) (34), head and
neck (HNSCC, 80-90%) (35, 36), bladder (37, 38), pancreatic (PDAC, 30-89%) (39), and
breast cancers (40). For these tumour types, therapies targeting EGFR, ERBB2, or their
downstream effectors such as MEK or PI3K, have shown some degree of success (36, 41–
43). As depicted in Figure 2A,B, cell-line clusters C1 and C2 exhibit sensitivity to drugs
in the D1 cluster that is enriched for drugs whose mode of action (methods and
Supplementary Table 04) is related to the inhibition of EGFR and MEK signalling
pathways.

Cluster C3 includes 69% of ovarian cancer cell lines (9 out of 13) and 70% of
endometrial/uterine ones (7 out of 10). Cell lines in this cluster are predicted to be sensitive
to drugs in the D2 cluster, which is enriched for topoisomerase, AURORA, HDAC, and
CDK inhibitors. Interestingly, ovarian and uterine cancers are usually due to inefficient

DNA repair (44). The standard of care for these cancer types is mainly based on
neoadjuvant chemotherapy and several assays on cell lines identified drugs targeting
apoptosis or cell cycle regulators as being effective (45, 46, 55–57, 47–54).

Cluster C4 includes all skin cancer cell lines (16 out of 16), all kidney (6 out of 6), all brain
(12 out of 12), and 13 lung cancer cell lines. Cell lines in cluster C4 are predicted to be
sensitive to multiple families of drugs, including RAF, KIT, PDGFR, and PI3K/mTOR
inhibitors. This agrees with current clinical practice which uses mTOR and BRAF
inhibitors as first-line therapy for kidney cancer and melanoma, respectively (58, 59), since
the c-Met/mTOR axis is frequently altered in kidney cancer (60, 61) while BRAF mutations
are the main genetic drivers of melanoma (59). Finally, RTKs are frequently altered in
glioblastoma (62, 63).

To further demonstrate DREEP's ability to identify recurring drug vulnerabilities across
cancer cells of different tumour types with shared genetic backgrounds, we applied our
method to the dataset from McFarland et al. (64). This dataset encompasses 3,630 untreated
and 4,175 nutlin-treated cells, culminating in an extensive collection of 7,805 cells (Figure
2C). Cells were sampled at two time points, 6 and 24 hours post-treatment, and originated
from 24 distinct cancer cell lines covering 14 different tumour types. Nutlin, a selective
MDM2 inhibitor, is a negative regulator of the tumour-suppressor gene TP53, triggering
apoptosis and cell cycle arrest exclusively in cancer cells retaining the wild-type TP53 form
(65). As depicted in Figure 2C, seven of the 24 cell lines within this dataset harbour a
missense mutation in the TP53 gene, while all other cell lines retain the wild-type (WT)
TP53 form. Indeed, as demonstrated in Supplementary Figure 06, DREEP accurately
predicted that most TP53 wild-type cells would exhibit sensitivity to nutlin treatment, while
cells bearing TP53 mutations would display resistance. To validate these predictions, we
first aggregated them at the cell line level by computing the median enrichment score of
nutlin estimated by DREEP across individual cells within each respective cell line and then
correlated these scores with the experimentally observed nutlin response from McFarland
et al. (64) (methods). As depicted in Figure 2D,E, this correlation held true for both
untreated and nutlin-perturbed cells and across the two time points. These findings further
underscore the predictive accuracy of our method in effectively characterizing drug
responses.

Altogether, these results show that DREEP can find recurrent drug sensitivity patterns
shared by cell lines from multiple cancer types, as well as the relationship between drug
response and functional cell status.

4. The DREEP method recapitulates cell population drug response over time.
Next, to evaluate DREEP's ability to recognize adaptation to drug treatment over time, we
applied it to the Ben-David et al. dataset (66). As Figure 3A shows, this dataset comprises
7,440 cells from a single cell-derived clone of the MCF7 cell line exposed for two days to
500 nM of bortezomib, a 26S proteasome inhibitor. Specifically, cells were sequenced
before treatment (t0), at 12 hrs and 48 hrs following drug exposure (t12, t48), and 96 hrs
following drug washout and recovery (t96).

As shown in Figure 3B, the distribution of Bortezomib ES scores estimated by DREEP at
time t0, t12 and t48 exhibits a bimodal pattern, indicating the presence of a heterogeneous
cell population comprising both bortezomib-tolerant and bortezomib-sensitive cells

(Figure 2E,F). By contrast, DREEP identified that at the final time point (t96), the
distribution of bortezomib ES scores is unimodal, with a homogeneous cell population
where most cells underwent transcriptome rewiring characterized by the expression of
biomarker genes associated with bortezomib resistance (Figure 3C,D). Finally, we were
interested in evaluating DREEP's ability to detect distinct drug-response cell
subpopulations before and after bortezomib treatment. To this end, we used drug response
profiles estimated by DREEP instead of transcriptional profiles to build a new cell-
embedding space (Methods). Figure 3E shows that the cell-embedding space built from
DREEP sensitivity predictions is better than the mixed cell groups identified using
transcriptional profiles (Figure 3A) at separating MCF7 cells into two main clusters
resembling the drug exposure over time.

In summary, these analyses demonstrate DREEP's ability to recapitulate drug response and
variability over time.

5. The DREEP method identifies small molecules that selectively inhibit cancer cell
subpopulations.
Next, we evaluated DREEP's ability to identify innate drug-resistant cell populations and
to suggest drug co-treatments. To this end, we used single cell data of the untreated
barcoded MDA-MB-468 breast cancer cell line that we recently generated (4). This dataset
includes 1,541 single cells from the untreated MDA-MB-468 cell line, each associated with
a molecularly expressed barcode (i.e., lineage). In the original study, the purpose of these
barcodes was to enable single-cell lineage tracing within the MDA-MB-468 cell
population. Subsequently, these cells underwent exposure to increasing concentrations of
Afatinib, a tyrosine kinase inhibitor that targets EGFR and HER2 receptors, over an
extended period exceeding 40 days. A bulk NGS approach was then used to identify the
lineages (i.e., barcodes) that survived this prolonged Afatinib exposure. Finally, the
surviving lineages were retrospectively mapped onto the single-cell sequencing data of the
untreated barcoded MDA-MB-468 cells. This approach, known as "retrospective lineage
tracing", enabled the categorization of untreated MDA-MB-468 cells as either sensitive or
tolerant to Afatinib and anti-EGFR treatments (4) (Figure 3F).

We applied DREEP to each of the two untreated MDA-MB-468 cell subpopulations (i.e.
afatinib-tolerant and afatinib-sensitive) and predicted drugs able to selectively inhibit the
growth of either. We thus found 156 drugs predicted to preferentially inhibit the afatinib-
tolerant subpopulation and 80 drugs (FDR < 10% and Supplementary Table 05) for the
afatinib-sensitive subpopulation (Figure 3G), which included 11 out of 12 EGFR inhibitors
(Supplementary Table 05).

Interestingly, the most overrepresented classes among the drugs predicted to selectively
inhibit the afatinib-tolerant subpopulation were IGF1R, HDAC, and AKT/MTOR/PI3K
inhibitors (Figure 3H). We have already shown that afatinib and IGF1R inhibitors have a
strong synergistic effect on these cells (4), thus validating the DREEP predictions that
IGF1R inhibitors should deplete the cell population that do not respond to afatinib in this
cell line.

To experimentally validate these predictions further, we selected three HDAC inhibitors
(Tenovin-6, bellinostat, and AR-42) and three AKT/MTOR/PI3K inhibitors (Uprosertib,
PIK93, and buparlisib) from the list of drugs predicted to selectively inhibit the afatinib-

tolerant subpopulation. We then measured experimentally if these drugs had an additive,
synergistic, or antagonistic effect on MDA-MB-468 cells in combination with afatinib
(Supplementary Table 06). As shown in Figure 3I, the median synergy score of each of the
six selected drugs tested in combination with afatinib is compatible with an additive effect
(67), suggesting that the two drugs work independently on different subpopulations of cells.
Finally, as Figure 3J shows, when cells are embedded into an embedding space
reconstructed from DREEP sensitivity profiles (Methods), two main cell clusters appear
that are enriched for either afatinib-tolerant or afatinib-sensitive lineages, in contrast to the
mixed cell groups identified using transcriptional profiles (Figure 3F). These results show
how the DREEP method alone can recognise that there are two different cell populations
in this cell line characterized by a distinct drug response profile.

Altogether, these results show that DREEP can predict drug sensitivity at the single cell
level and identify drugs able to selectively inhibit a subpopulation of cells that can be then
combined to find drug combinations.

6. The DREEP method accurately predicts drug response in heterogeneous cancer
patient cells.
Cancer cell lines often display restricted transcriptomic heterogeneity, failing to mirror the
full complexity of tumours. Therefore, to demonstrate the clinical applicability of DREEP,
we utilized our methodology on a series of patient-derived cultures (PDCs) from both
primary and metastatic sites of individuals suffering from head and neck squamous cell
carcinoma (23). These models have already been proven to effectively replicate the
complexity of the corresponding matching tumour (68), with the advantage that sensitivity
measurements could be consistently assessed across multiple drugs (68). Specifically, this
dataset comprises 1,027 single cell transcriptional profiles of PDCs obtained from five head
and neck cancer patients whose drug response was measured in triplicate for five different
small molecules at two different concentrations (18).

As illustrated in Figure 4A, the UMAP representation of single-cell transcriptomic profiles
confirmed the presence of intra-patient transcriptomic heterogeneity within PDCs,
particularly when juxtaposed with inter-patient heterogeneity. Notably, in line with
observations previously reported in (23, 68), patients HN120 and HN137 displayed a
significant degree of heterogeneity, where primary and lymph node metastatic tumour cells
delineate two transcriptionally distinct cell subpopulations.

Next, to assess the predictive performance of DREEP on this dataset, we first applied it to
the single-cell profiles of each patient and then converted these predictions from the single-
cell level to the patient level by computing the median enrichment score for each drug
across the cells within each individual patient. As depicted in Figure 4B, our analysis
uncovered a significant correlation (Spearman's correlation coefficient = 0.634, p-value =
3.6e-09) between DREEP predictions and the experimentally observed drug responses for
five specific drugs: Docetaxel, Doxorubicin, Epothilone B, Gefitinib, and Vorinostat (refer
to methods) (18). Patients' cell viability was determined using each drug's median IC50
values (18), which were threefold lower than those estimated from ATCC head and neck
cancer cell lines present in the GDSC portal (methods). Finally, Figure 4C illustrates
Spearman's correlation coefficient (SCC) between DREEP predictions and the
experimentally observed drug responses within each independent GPDS dataset. Notably,

GDSC exhibited the strongest performance with an SCC of 0.749 (p-value = 1.6e-05),
followed by CTRP2 with an SCC of 0.591 (p-value = 0.006), and PRISM with an SCC of
0.512 (p-value = 0.008). These findings remained consistent even when we employed
patient cell viability estimates derived from drug concentrations matching the estimated
IC50 values, as shown in Supplementary Figure 07.

Discussion
Intra-tumour heterogeneity (ITH) presents a significant challenge in developing effective
cancer treatments and achieving precision oncology. ITH refers to the variation of tumour
cells both within and between tumours. As such, it is crucial to have methods to accurately
identify the drug sensitivities of different tumour clones and determine the most effective
treatment for each cancer type and the individual patient. To address this challenge, we
introduced DREEP, a novel computational method that predicts drug sensitivity at the
single cell level and can propose cell population-specific treatments. To achieve this,
DREEP uses a set of drug-sensitivity signatures obtained by correlating gene expression
and drug sensitivity profiles across hundreds of tumorigenic cell lines. It then uses these
signatures to compute an enrichment score reflecting the degree of sensitivity of a cell to a
specific drug.
To test the predictive performance and reliability of DREEP, we applied it to two
independent datasets: (i) the pan-cancer cell line dataset from Kinker et al. comprising 198
tumorigenic cell lines from 22 distinct cancer types (30) and (ii) the single cell breast cancer
atlas from Gambardella et al. that includes 32 breast cancer cell lines (5). We observed that
DREEP successfully identifies the most effective drugs for each cell line with an overall
performance that is several fold better than random on both datasets. We also successfully
used DREEP to reconstruct the single-cell variability in drug response across the 198 cell
lines of the pan-cancer dataset and to identify the functional cellular activities linked to
common drug response patterns among cancer types and cell lines. Our approach can offer
valuable insights for understanding the complex interplay between genetic backgrounds
and drug sensitivity, paving the way for more personalized and effective cancer treatments.
In addition, when used to analyse single-cell data collected from the breast cancer cell line
MCF7 exposed to bortezomib over time (66), DREEP easily reconstructed the interchange
between the sensitive and tolerant cell populations. Finally, using single cell data of the
molecularly labelled MDA-MB-468 cell line (4), we showed that DREEP can identify
innate drug-resistant cell populations and predict drug co-treatments able to deplete these
cell populations. Overall, we show that DREEP produces an accurate reconstruction of the
variability of drug response in cancer cell lines from single cell transcriptomics. The
DREEP R package is available at https://github.com/gambalab/DREEP.
Our tool relies heavily on the quality and predictive power of GPDS, which were trained
by linking gene expression to drug response data from pharmacogenomic screens.
However, these screens may possess inherent biases and limitations in covering the broad
spectrum of cancer types. We applied several evaluation metrics on multiple showcases to
assess DREEP's predictive accuracy and ability to generalize across different cancer types,
encompassing approximately one hundred thousand cells originating from more than 20
distinct cancer types.
408

While our results have demonstrated DREEP's ability to generalize across various cancer
types effectively, we recognize that potential biases in our training data could limit
DREEP's applicability to underrepresented cancer types. Therefore, it is crucial to interpret
DREEP's performance within the context of the available training data. We advocate for
ongoing refinement and expansion of pharmacogenomic screens to improve the diversity
and coverage of cancer types. Addressing these biases in future data collection efforts can
enhance DREEP's utility in advancing personalized cancer treatment strategies. Moreover,
in the future, it would be crucial to evaluate how the integration of other single-cell -omics
measurements, such as genomic, proteomic, and epigenetic cell states, can improve the
identification of variations in drug responses among cancer cell populations. Finally, since
tumour heterogeneity is linked to clonal evolution processes, it would be also important to
consider the dynamics of clonal populations under therapy pressure to improve the
performance of future computational methods that will be developed for personalized
diagnosis and treatment of cancer patients.

Figure & Legends

Figure 1 – Drug Response Estimation from single-cell Expression Profiles (DREEP). (**A**) Construction of Genomic Profiles of Drug Sensitivity (GPDS). The potency profile of each drug in terms of AUC (Area Under the Curve) is correlated with the expression of all messenger RNA across the cell lines for which the drug potency was measured. (**B**) Bioinformatics pipeline for the prediction of single-cell drug response. Raw UMI counts are first normalized with the gf-icf pipeline and then the most relevant genes in a single cell are used as input for a GSEA against the GPDS ranked-list to predict its drug sensitivity. (**C**) Performance of DREEP drug sensitivity on 198 cell lines for each set of GPDS signatures used and different numbers of relevant genes to predict the effect of a drug. DREEP's performance is estimated using either Precision-recall curve (upper) or ROC curve (bottom). Max AUC of ROC curves are CTRP2 = 0.691, GDSC = 0.728 and PRISM = 0.661. (**D**) Left column: Each point is a drug whose y-coordinate is AUC of the corresponding ROC curve across the 198 cell lines and computed by sorting cell lines from the most to the least sensitive to the drug. Each panel reports the results using different numbers of relevant genes to predict the effect of the drug. Right column: Each point is a cell line whose y-coordinate is the AUC of the corresponding ROC curve obtained by sorting drug predictions from the most to the least sensitive. Each panel reports the results using different numbers of relevant genes to predict the effect of a drug. (**E**) Same as (C) but for the 32 breast cancer cell lines published in Gambardella et al. (5). Max AUC of ROC curves are CTRP2 = 0.721, GDSC = 0.699 and PRISM = 0.637. (**F**) Same as (D) but for 32 breast cancer cell lines published in Gambardella et al. (5).

445
Figure 2 – DREEP captures single cell variability in drug response. (**A**) Heatmap depicting both cell-line
clusters (rows) and cancer-specific drug clusters (columns). Cell lines are colour-coded according to the
cancer type while drugs are colour-coded according to the dataset to which they belong. Both rows and
columns of the heatmap were grouped in four clusters named C1-C4 for cell line (rows) and D1-D4 for drugs
(columns). (**B**) Bar-plot depicting enriched MoAs in each cancer-specific drug cluster. Only the top ten
significant MoAs (FDR < 10%) for each cluster are shown. (**C**) UMAP plot of 7,805 cells of 24 cell lines
either untreated or exposed to nutlin treatment. (**D**) The Spearman correlation coefficient (SCC) between
observed and predicted sensitivity scores for the XX untreated (DMSO) cell lines in (C). In the upper plot
the SCC is computed using cells sequenced after 6 hours, while in the bottom plot SCC is computed using
cells collected after 24 hours. (**E**) Same as in (D) but for nutlin-treated cells.

457
Figure 3 – DREEP identifies drugs that can selectively inhibit a subpopulation of cells. (**A**) UMAP plot
of 7,440 cells of the MCF7 cell line exposed to 500 nM bortezomib. (**B**) Bortezomib DREEP enrichment
score (ES) distribution at each timepoint for the 7,440 cells of the MCF7 cell line in (A). A positive or
negative score means that the cell is predicted to be resistant or sensitive to bortezomib, respectively. (**C**)
Bortezomib DREEP's median enrichment score of MCF7 cells at each time-point. (**D**) Percentage of
predicted bortezomib sensitive or tolerant cells for each time point. (**E**) UMAP plot of the same cells in (A)
but using the drug profile estimated by DREEP instead of its corresponding transcriptional profile for each
cell. In the left panel, cells are colour-coded according to the sequencing timepoint, while in the right panel
cells are colour-coded according to the bortezomib score predicted by DREEP. A positive or negative score
means that the cell is predicted to be resistant or sensitive to bortezomib, respectively. (**F**) UMAP plot of
1,541 MDA-MB-468 cells from Pellecchia et al. (4). Cells are colour-coded according to whether they belong
to an afatinib-sensitive or tolerant lineage. (**G**) Each point represents a drug whose coordinates are the median
enrichment score predicted by DREEP for the drug in afatinib-tolerant cells [x-axis] and afatinib-sensitive
cells [y-axis]. The more negative the value of the enrichment score, the more potent the predicted effect of
the drug. The drugs specifically inhibiting at least one of the two subpopulations are indicated in red. (**H**)
Heatmap showing enriched drug classes predicted to specifically inhibit afatinib-tolerant or afatinib-sensitive
cells. Each row of the heatmap represents a drug coloured according to its median enrichment score predicted
by DREEP across cells of the same lineage (i.e. afatinib-tolerant or afatinib-sensitive). (**I**) Median synergy
score of drugs predicted by DREEP to selectively inhibit afatinib-tolerant lineages in MDA-MB-468 cells.
Each drug was tested in combination with afatinib, and the synergy score computed using the Loewe
statistical model. A positive score represents a synergism between the two tested drugs. (**J**) UMAP plot of
the cells in (A) but using the drug profile estimated by DREEP instead of the corresponding transcriptional
profile for each cell. Cells are colour-coded according to whether they belong to an afatinib-sensitive or
tolerant lineage.

483
484

485 **Figure 4 4 Drug response prediction in heterogenous patient-derived cultures (PDCs).** (**A**) UMAP plot
486 depicting 1,027 cells of PDCs from both primary and metastatic sites in five individuals with head and neck
487 squamous cell carcinoma. (**B**) The Spearman correlation coefficient (SCC) between observed and predicted
488 sensitivity scores for the five PDCs using five different drugs at lower concentrations. All three GDPS dataset
489 are used. (**C**) A breakdown of (B) by GDPS datasets.

490
491

Methods

scRNA-seq datasets processing: The scRNA-seq data of the cell line pan-cancer dataset from Kinker et al. (30) was obtained from the Broad Institute's single cell portal (SCP542). Specifically, CPM count matrix was downloaded. Before being processed, gene symbols were converted into ensemble id and only ensemble id associated with no more than one gene were retained. At the end of this pre-processing step, the final cell line pan-cancer matrix contained 16,558 genes and 53,513 cells. Raw scRNA-seq data of the MCF7 cell line exposed to bortezomib (66) was obtained from GEO (GSE114461) and only cells with at least 5,000 UMI were retained. The single cell breast cancer atlas dataset was obtained from figshare at the following address https://doi.org/10.6084/m9.figshare.15022698. scRNA-seq data of barcoded MDA-MB-468 cells were obtained from GEO (GSE228154). Raw UMI count data of nutlin treated and untreated cells was downloaded from figshare at https://figshare.com/s/139f64b495dea9d88c70 and only cells with at least 5,000 UMI and with less than 10% of UMI counts in mitochondrial genes were used. PDCs TPM counts were instead retrieved from supplementary data of the original work (18).

Bulk RNA-seq dataset processing: Raw counts and associated metadata of the bulk RNA-seq data of cancer cell lines used to build GPDS signature were obtained from the depmap portal (https://depmap.org). Cell lines associated with haematopoietic tumours were excluded. Raw counts were normalised with edgeR (69) and transformed into log10 (CPM+1). Poorly expressed genes and genes whose entropy values were in the fifth percentile were discarded. Entropy values were estimated by discretizing the expression of each gene in ten equally spaced bins using the functions included in the R package named *entropy*. After these filtering steps, the final expression matrix comprised 1,117 cell lines and 13,849 genes.

Drug sensitivity dataset processing: The drug sensitivity dataset of CTRP2 (Cancer Therapeutic Response Portal), GDSC (Genomics of Drug Sensitivity in Cancer) and PRISM (Profiling Relative Inhibition in Mixtures) were all obtained from the depmap portal (https://depmap.org). Only drugs for which the in vitro response was available for at least 100 cell lines out of the 1,117 for which gene expression data was available were retained. After this filtering, the final drug sensitivity dataset comprised 478 drugs from the GDSC dataset, 511 drugs from the CTRP2 dataset, and 1,445 drugs from the PRISM dataset. For all three cell viability datasets, drug potency was expressed in terms of AUC (Area Under the Curve) of the corresponding dose response graph.

Construction of GPDS ranked-lists: Genomic Profiles of Drug Sensitivity (GPDS) ranked-lists were built by computing the Pearson correlation coefficient (PCC) between the expression of each gene and the potency of a drug across the cell lines for which drug potency was measured. At the end of this process, each GPDS is a ranked list of 13,849 expression-based biomarker genes whose degree of PCC reflects their importance in predicting the effect of a small molecule. This is because the AUC value of a drug reflects its cell line growth inhibition, with lower values indicating sensitivity and higher values implying tolerance to the drug. Thus, a gene whose expression positively correlates with the AUC of a drug is predictive of resistance to that drug, conversely a negatively correlated gene is predictive of sensitivity to the drug.

Drug Response Estimation from single-cell Expression Profiles (DREEP): To predict drug response at the single-cell level, we first used gf-icf normalization (https://github.com/gambalab/gficf) (27) to extract the top relevant genes from each cell. We then used these gene sets as input for Gene Set Enrichment Analysis (GSEA) (29) against each GDPS ranked-list to predict the sensitivity of a cell to a specific drug. Since GPDS ranked-lists contain predictive biomarkers of resistance at the top and predictive biomarkers of sensitivity at the bottom for each drug, a positive enrichment score implies that the cell expresses genes that render it tolerant to the drug, while a negative enrichment score indicates the cell expresses genes that render it sensitive to the drug. Finally, by estimating the median enrichment score on a cell population and counting the number of cells with a significant enrichment score less than zero, it is possible to estimate the overall effect of a drug on that population and the percentage of cells sensitive to it. The DREEP package is implemented in R and is available on github at the following address https://github.com/gambalab/DREEP. Enrichment Scores (ES) and associated p-values were computed using the *fgsea* package (70). The *fgsea* package utilizes an adaptive and efficient multi-level split Monte Carlo approach (71) to estimate the p-value of an ES. Subsequently, the p-values for each drug across the cells are corrected for the false discovery rate (FDR) using the Benjamini-Hochberg correction method, implemented within the function *p.adjust* in the

547 *base* package of the R statistical environment. Finally, only ES values with an FDR < 0.1 were deemed
548 significant (unless otherwise stated) and used to predict the effect of a drug on a single cell. The percentage
549 of sensitive and resistant cells in a sample are finally computed using only significant ES scores.
550
Estimation of DREEP performance: To estimate DREEP's precision and sensitivity in predicting drug
response from scRNA-seq data of cancer cell lines, we used the three cell-viability screening datasets from
CTRP2, GDSC, and PRISM described above. For each cell-viability screening dataset, the corresponding
"gold standard" is composed of pairs of cell lines and drugs for which the corresponding cell line is sensitive.
To build the "gold standard" of each cell-viability screening dataset, we first computed the z-score percentiles
from the AUC of each drug across the cell lines for which AUC was available and then defined a cell line
sensitive to a drug only if its z-score was in the 5% percentile. Next, we applied DREEP to scRNA-seq data
of each cell line and computed its median enrichment score for each drug. Finally, Precision-Recall and ROC
curves on each cell-viability screening dataset depicted in Figure 1C-F were computed with the *precrec* R
package (https://github.com/evalclass/precrec) using the corresponding "gold standard" and sorting DREEP
drug/cell line pairs according to the estimated median cell line enrichment score multiplied by minus one in
order to have drugs predicted to be sensitive at the top.
563
Cell embedding space from DREEP predictions: DREEP output can be arranged in a drug response matrix
E of dimensions $n \times m$ where n is equal to the number of cells and m equal to the number of GPDS used.
Each element $E_{i,j}$ of matrix E ranges within the $[-1, +1]$ interval and it is different from zero only if the
enrichment score between the relevant genes of the cell i and the GPDS ranked-list j is significant (i.e. FDR
< 0.1). Specifically, $E_{i,j} > 0$ indicates the cell i is predicted resistant to the drug j, while $E_{i,j} < 0$ indicates
the cell i is predicted sensitive to the drug j, and $E_{i,j} = 0$ indicates that there is no supporting evidence to
decide if cell i is sensitive or not to the drug j. Next, we computed the similarity of the predicted drug
response profile between two cells as the distance between two cells x and y with *Fuzzy Jaccard Distance*
defined as follow:

$$FJD_{x,y} = 1 - \frac{\sum_{j=1}^{m} |sign(E_{x,j}) + sign(E_{y,j})|}{2 * m}$$

FJD is a non-negative symmetric distance matrix of dimensions $n \times n$ whose elements range within the $[0,1]$
interval. Specifically, each element $FJD_{x,y}$ of FJD quantifies how many of the drugs to which the cell x is
sensitive are shared with the cell y and how many of the drugs the cell x is resistant to being shared with the
cell y. The more two cells share drugs that are predicted to have the same effect, the closer the $FJD_{x,y}$ value
is to 0, *and vice versa* for two cells with opposite drug predictions, $FJD_{x,y}$ equals 1. It can be easily shown
that the Fuzzy Jaccard Distance we implemented satisfies all the conditions to be defined as a metric,
including the triangle inequality constraint. Finally, the cell embedding space was built using the
precomputed cell-to-cell distance matrix FJD as input of the *umap* function of the uwot R package
(https://github.com/jlmelville/uwot). Cell embedding from DREEP predictions has been implemented in the
function *runDrugReduction* of the DREEP package.
584
The therapeutic landscape of the cell line pan-cancer dataset (Figure 2A): We first used DREEP to
predict the effect of the 2,434 drugs on each of the 53,513 single-cell transcriptional profiles of the Kinker et
al. dataset (30). Then, we converted predictions from the single-cell level to the cell-line level by counting
the fraction of cells predicted to be sensitive to each drug (i.e. DREEP enrichment score < 0 and P < 0.05).
At the end of this process, we obtained a matrix $A_{198 \times 2,434}$ where $A_{i,j}$ contains the percentage of cells of the
cell line i predicted to be sensitive to the drug j. Next, we set elements of A less than 0.9 as being equal to 0
while all elements of A greater or equal to 0.9 were set as equal to 1. Then, we hierarchically clustered rows
(i.e. cell lines) and columns (i.e. drugs) of A using the Jaccard distance. Jaccard distances between row pairs
and column pairs of the matrix A were computed using the *dist* function of the R statistical environment.
Clustering of both, cell lines and drugs, was performed using the *hclust* function and resulting dendrograms
cut with the *cutree* function of R statistical environment.
596
597 **Differential drug sensitivity analysis:** We used DREEP to predict the effects of 2,434 drugs on 1,541
598 sequenced cells of the barcoded MDA-MB-468 cell line (4). Then, we performed a Mann-Whitney test and

corrected p-values using the Benjamini-Hochberg method to identify drugs with differential sensitivity between afatinib-sensitive and afatinib-tolerant subpopulations. Finally, a drug was considered specific for a subpopulation if its FDR was less than 0.1 and the median enrichment score across the other subpopulation was greater than zero. This analysis has been implemented in the *runDiffDrugAnalysis* function of the DREEP package.

Drug combinations assay: 4×10^4 MDA-MB-468 cells were seeded in a 96-well plate and incubated overnight at 37 °C. The two drugs to test were prepared in different dilutions and then combined in all possible drug pairs to generate a 5 x 5 combination matrix. Then, cells were exposed to either single agent drug or to the drug pairs of the drug combination matrix, while negative controls were treated with DMSO (each treatment was performed in triplicate). Following 72 hr incubation at 37 °C, cell viability was measured with CellTiter (Promega) and the absorbance was read at 490 nm with the plate reader GloMax® Discover instrument. The drug interactions and expected drug responses were calculated with the Combenefit tool (67) using the Loewe additivity model.

Drug Set Enrichment Analysis (DSEA): DSEA was employed by using the online drugEnrichr tool (31) using the Drug Repurposing Hub Mechanism of Action table resource.

Comparison with other state-of-the-art methods: All methods were run using default parameters and when required single cell expression was normalized using the Seurat package. Common drugs between DREEP and each method were found by simply intersecting the drug names

PDCs drug response data: PDCs drug response data were retrieved from supplementary data of the original work (18). Specifically, in the original work, eight drugs were tested on these five PDCs. However, we only used the five out of eight drugs for which cell viability was measured at least in triplicate. Raw viability measures we used can be found in Supplementary Table 07.

Code availability: The DREEP package is available on github at the following address https://github.com/gambalab/DREEP.

Data availability: All the processed datasets and associated metadata, including scRNA-seq, bulk RNA-seq, and sensitivity profiles used in this work have been deposited on figshare at the following address https://doi.org/10.6084/m9.figshare.23261234.

Funding: This work was supported by the My First AIRC grant 23162.

Author Contribution: SP and GV performed drug combination experiments. MF helped with bioinformatics analysis and in writing the manuscript. GG supervised the work, wrote the manuscript, and conceived the original idea.

Competing interests: The authors declare no competing interests.

Acknowledgement: We express our gratitude to Dr. Cathal Wilson for meticulously proofreading our manuscript. His invaluable efforts have significantly contributed to enhancing the clarity and accuracy of our work.

References

1. Shaffer,S.M., Dunagin,M.C., Torborg,S.R., Torre,E.A., Emert,B., Krepler,C., Beqiri,M., Sproesser,K., Brafford,P.A., Xiao,M., *et al.* (2017) Rare cell variability and drug-induced reprogramming as a mode of cancer drug resistance. *Nature*, **546**, 431–435.

2. Sharma,S. V., Lee,D.Y., Li,B., Quinlan,M.P., Takahashi,F., Maheswaran,S., McDermott,U., Azizian,N., Zou,L., Fischbach,M.A., *et al.* (2010) A Chromatin-Mediated Reversible Drug-Tolerant State in Cancer Cell Subpopulations. *Cell*, **141**, 69–80.

3. Tirosh,I., Izar,B., Prakadan,S.M., Wadsworth,M.H., Treacy,D., Trombetta,J.J., Rotem,A., Rodman,C., Lian,C., Murphy,G., *et al.* (2016) Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science (80-.).*, **352**, 189–196.

4. Pellecchia,S., Franchini,M., Viscido,G., Arnese,R. and Gambardella,G. (2023) Single cell lineage tracing reveals subclonal dynamics of anti-EGFR therapy resistance in triple negative breast cancer. *bioRxiv*, 10.1101/2023.04.04.535588.

5. Gambardella,G., Viscido,G., Tumaini,B., Isacchi,A., Bosotti,R. and di Bernardo,D. (2022) A single-cell analysis of breast cancer cell lines to study tumour heterogeneity and drug response. *Nat. Commun.*, **13**, 1714.

6. Li,X., Francies,H.E., Secrier,M., Perner,J., Miremadi,A., Galeano-Dalmau,N., Barendt,W.J., Letchford,L., Leyden,G.M., Goffin,E.K., *et al.* (2018) Organoid cultures recapitulate esophageal adenocarcinoma heterogeneity providing a model for clonality studies and precision therapeutics. *Nat. Commun.*, **9**, 2983.

7. Saeed,K., Ojamies,P., Pellinen,T., Eldfors,S., Turkki,R., Lundin,J., Järvinen,P., Nisen,H., Taari,K., af Hällström,T.M., *et al.* (2019) Clonal heterogeneity influences drug responsiveness in renal cancer assessed by ex vivo drug testing of multiple patient-derived cancer cells. *Int. J. Cancer*, **144**, 1356–1366.

8. Iorio,F., Knijnenburg,T.A., Vis,D.J., Bignell,G.R., Menden,M.P., Schubert,M., Aben,N., Gon??alves,E., Barthorpe,S., Lightfoot,H., *et al.* (2016) A Landscape of Pharmacogenomic Interactions in Cancer. *Cell*, **166**, 740–754.

9. Rees,M.G., Seashore-Ludlow,B., Cheah,J.H., Adams,D.J., Price,E. V, Gill,S., Javaid,S., Coletti,M.E., Jones,V.L., Bodycombe,N.E., *et al.* (2016) Correlating chemical sensitivity and basal gene expression reveals mechanism of action. *Nat Chem Biol*, **12**, 109–116.

10. Costello,J.C., Heiser,L.M., Georgii,E., Gonen,M., Menden,M.P., Wang,N.J., Bansal,M., Ammad-ud-din,M., Hintsanen,P., Khan,S.A., *et al.* (2014) A community effort to assess and improve drug sensitivity prediction algorithms. *Nat Biotechnol*, **32**, 1202–1212.

11. Daemen,A., Griffith,O.L., Heiser,L.M., Wang,N.J., Enache,O.M., Sanborn,Z., Pepin,F., Durinck,S., Korkola,J.E., Griffith,M., *et al.* (2013) Modeling precision treatment of breast cancer. *Genome Biol*, **14**, R110.

12. Heiser,L.M., Sadanandam,A., Kuo,W.L., Benz,S.C., Goldstein,T.C., Ng,S., Gibb,W.J., Wang,N.J., Ziyad,S., Tong,F., *et al.* (2012) Subtype and pathway specific responses to anticancer compounds in breast cancer. *Proc Natl Acad Sci U S A*, **109**, 2724–2729.

13. Rees,M.G., Seashore-Ludlow,B., Cheah,J.H., Adams,D.J., Price,E. V, Gill,S., Javaid,S., Coletti,M.E., Jones,V.L., Bodycombe,N.E., *et al.* (2015) Correlating chemical sensitivity and basal gene expression reveals mechanism of action. *Nat. Chem. Biol.*, **12**, 1–10.

14. Trastulla,L., Noorbakhsh,J., Vazquez,F., McFarland,J. and Iorio,F. (2022) Computational estimation of quality and clinical relevance of cancer cell lines. *Mol. Syst. Biol.*, **18**, e11017.

15. Wu,L., Wen,Y., Leng,D., Zhang,Q., Dai,C., Wang,Z., Liu,Z., Yan,B., Zhang,Y., Wang,J., *et al.* (2022) Machine learning methods, databases and tools for drug combination prediction. *Brief. Bioinform.*, **23**, bbab355.

16. Adam,G., Rampášek,L., Safikhani,Z., Smirnov,P., Haibe-Kains,B. and Goldenberg,A. (2020) Machine learning approaches to drug response prediction: challenges and recent progress. *npj Precis. Oncol.*, **4**, 19.

17. Hsieh,C.-Y., Wen,J.-H., Lin,S.-M., Tseng,T.-Y., Huang,J.-H., Huang,H.-C. and Juan,H.-F. (2023) scDrug: From single-cell RNA-seq to drug response prediction. *Comput. Struct. Biotechnol. J.*, **21**, 150–157.

18. Suphavilai,C., Chia,S., Sharma,A., Tu,L., Da Silva,R.P., Mongia,A., DasGupta,R. and Nagarajan,N. (2021) Predicting heterogeneity in clone-specific therapeutic vulnerabilities using single-cell transcriptomic signatures. *Genome Med.*, **13**, 189.

19. Chen,J., Wang,X., Ma,A., Wang,Q.-E., Liu,B., Li,L., Xu,D. and Ma,Q. (2022) Deep transfer learning of cancer drug responses by integrating bulk and single-cell RNA-seq data. *Nat. Commun.*, **13**, 6494.

20. Fustero-Torre,C., Jiménez-Santos,M.J., García-Martín,S., Carretero-Puche,C., García-Jimeno,L., Ivanchuk,V., Di Domenico,T., Gómez-López,G. and Al-Shahrour,F. (2021) Beyondcell: targeting cancer therapeutic heterogeneity in single-cell RNA-seq data. *Genome Med.*, **13**, 187.

21. Hetzel,L., Böhm,S., Kilbertus,N., Günnemann,S., Lotfollahi,M. and Theis,F. (2022) Predicting Cellular Responses to Novel Drug Perturbations at a Single-Cell Resolution. *Adv. Neural Inf. Process. Syst.*, **35**, 1–19.

22. Xia,F., Allen,J., Balaprakash,P., Brettin,T., Garcia-Cardona,C., Clyde,A., Cohn,J., Doroshow,J., Duan,X., Dubinkina,V., *et al.* (2022) A cross-study analysis of drug response prediction in cancer cell lines. *Brief. Bioinform.*, **23**, bbab356.

23. Sharma,A., Cao,E.Y., Kumar,V., Zhang,X., Leong,H.S., Wong,A.M.L., Ramakrishnan,N., Hakimullah,M., Teo,H.M.V., Chong,F.T., *et al.* (2018) Longitudinal single-cell RNA sequencing of patient-derived primary cells reveals drug-induced infidelity in stem cell hierarchy. *Nat. Commun.*, **9**, 4931.

24. Rees,M.G., Brenan,L., do Carmo,M., Duggan,P., Bajrami,B., Arciprete,M., Boghossian,A., Vaimberg,E., Ferrara,S.J., Lewis,T.A., *et al.* (2022) Systematic identification of biomarker-driven drug combinations to overcome resistance. *Nat. Chem. Biol.*, **18**, 615–624.

25. Corsello,S.M., Nagari,R.T., Spangler,R.D., Rossen,J., Kocak,M., Bryan,J.G., Humeidi,R., Peck,D., Wu,X., Tang,A.A., *et al.* (2020) Discovering the anticancer potential of non-oncology drugs by systematic viability profiling. *Nat. Cancer*, **1**, 235–248.

26. Gambardella,G. and di Bernardo,D. (2019) A Tool for Visualization and Analysis of Single-Cell RNA-Seq Data Based on Text Mining. *Front. Genet.*, **10**.

27. Franchini,M., Pellecchia,S., Viscido,G. and Gambardella,G. (2023) Single-cell gene set enrichment analysis and transfer learning for functional annotation of scRNA-seq data. *NAR Genomics Bioinforma.*, **5**, lqad024.

28. Slovin,S., Carissimo,A., Panariello,F., Grimaldi,A., Bouché,V., Gambardella,G. and Cacchiarelli,D. (2021) Single-Cell RNA Sequencing Analysis: A Step-by-Step Overview BT - RNA Bioinformatics. In Picardi,E. (ed). Springer US, New York, NY, pp. 343–365.

29. Subramanian,A., Tamayo,P., Mootha,V.K., Mukherjee,S., Ebert,B.L., Gillette,M.A., Paulovich,A., Pomeroy,S.L., Golub,T.R., Lander,E.S., *et al.* (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.*, **102**, 15545–50.

30. Kinker,G.S., Greenwald,A.C., Tal,R., Orlova,Z., Cuoco,M.S., McFarland,J.M., Warren,A., Rodman,C., Roth,J.A., Bender,S.A., *et al.* (2020) Pan-cancer single-cell RNA-seq identifies recurring programs of cellular heterogeneity. *Nat. Genet.*, **52**, 1208–1218.

31. Kropiwnicki,E., Evangelista,J.E., Stein,D.J., Clarke,D.J.B., Lachmann,A., Kuleshov,M. V, Jeon,M., Jagodnik,K.M. and Ma'ayan,A. (2021) Drugmonizome and Drugmonizome-ML: integration and abstraction of small molecule attributes for drug enrichment analysis and machine learning. *Database*, **2021**, baab017.

32. Tajadura-Ortega,V., Gambardella,G., Skinner,A., Halim,A., Van Coillie,J., Schjoldager,K.T.-B.G., Beatson,R., Graham,R., Achkova,D., Taylor-Papadimitriou,J., *et al.* (2020) O-linked mucin-type glycosylation regulates the transcriptional programme downstream of EGFR. *Glycobiology*, 10.1093/glycob/cwaa075.

33. Pabla,B., Bissonnette,M. and Konda,V.J. (2015) Colon cancer and the epidermal growth factor receptor: Current treatment paradigms, the importance of diet, and the role of chemoprevention. *World J. Clin. Oncol.*, **6**, 133–141.

34. Prabhakar,C.N. (2015) Epidermal growth factor receptor in non-small cell lung cancer. *Transl. Lung Cancer Res. Vol 4, No 2 (April 19, 2015) Transl. Lung Cancer Res. (Molecular Genet. Lung Cancer)¹*

35. Stransky,N., Egloff,A.M., Tward,A.D., Kostic,A.D., Cibulskis,K., Sivachenko,A., Kryukov,G. V, Lawrence,M.S., Sougnez,C., McKenna,A., *et al.* (2011) The Mutational Landscape of Head and Neck Squamous Cell Carcinoma. *Science (80-.).*, **333**, 1157–1160.

36. Lawrence,M.S., Sougnez,C., Lichtenstein,L., Cibulskis,K., Lander,E., Gabriel,S.B., Getz,G., Ally,A., Balasundaram,M., Birol,I., *et al.* (2015) Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*, **517**, 576–582.

37. Chaux,A., Cohen,J.S., Schultz,L., Albadine,R., Jadallah,S., Murphy,K.M., Sharma,R., Schoenberg,M.P. and Netto,G.J. (2012) High epidermal growth factor receptor immunohistochemical expression in urothelial carcinoma of the bladder is not associated with EGFR mutations in exons 19 and 21: a study using formalin-fixed, paraffin-embedded archival tissues. *Hum. Pathol.*, **43**, 1590–1595.

38. Røtterud,R., Nesland,J.M., Berner,A. and Fosså,S.D. (2005) Expression of the epidermal growth factor receptor family in normal and malignant urothelium. *BJU Int.*, **95**, 1344–1350.

39. Nedaeinia,R., Avan,A., Manian,M., Salehi,R. and Ghayour-Mobarhan,M. (2014) EGFR as a potential target for the treatment of pancreatic cancer: dilemma and controversies. *Curr. Drug Targets*, **15**, 1293–1301.

40. Harbeck,N., Penault-Llorca,F., Cortes,J., Gnant,M., Houssami,N., Poortmans,P., Ruddy,K., Tsang,J. and Cardoso,F. (2019) Breast cancer.

41. Han,J., Liu,Y., Yang,S., Wu,X., Li,H. and Wang,Q. (2021) MEK inhibitors for the treatment of non-small cell lung cancer. *J. Hematol. Oncol.*, **14**, 1.

42. Xie,Y.-H., Chen,Y.-X. and Fang,J.-Y. (2020) Comprehensive review of targeted therapy for colorectal cancer. *Signal Transduct. Target. Ther.*, **5**, 22.

43. Johnson,D.E., Burtness,B., Leemans,C.R., Lui,V.W.Y., Bauman,J.E. and Grandis,J.R. (2020) Head and neck squamous cell carcinoma. *Nat. Rev. Dis. Prim.*, **6**, 92.

44. Mirza-Aghazadeh-Attari,M., Ostadian,C., Saei,A.A., Mihanfar,A., Darband,S.G., Sadighparvar,S., Kaviani,M., Samadi Kafil,H., Yousefi,B. and Majidinia,M. (2019) DNA damage response and repair in ovarian cancer: Potential targets for therapeutic strategies. *DNA Repair (Amst).*, **80**, 59–84.

45. Al-Alem,L.F., Baker,A.T., Pandya,U.M., Eisenhauer,E.L. and Rueda,B.R. (2019) Understanding and Targeting Apoptotic Pathways in Ovarian Cancer. *Cancers* , **11**.

46. Zervantonakis,I.K., Iavarone,C., Chen,H.-Y., Selfors,L.M., Palakurthi,S., Liu,J.F., Drapkin,R., Matulonis,U., Leverson,J.D., Sampath,D., *et al.* (2017) Systems analysis of apoptotic priming in ovarian cancer identifies vulnerabilities and predictors of drug response. *Nat. Commun.*, **8**, 365.

47. Shen,Q., Li,J., Mai,J., Zhang,Z., Fisher,A., Wu,X., Li,Z., Ramirez,M.R., Chen,S. and Shen,H. (2018) Sensitizing non-small cell lung cancer to BCL-xL-targeted apoptosis. *Cell Death Dis.*, **9**, 986.

48. Han,Z., Liang,J., Li,Y. and He,J. (2019) Drugs and Clinical Approaches Targeting the Antiapoptotic Protein: A Review. *Biomed Res. Int.*, **2019**, 1212369.

49. Du,R., Huang,C., Liu,K., Li,X. and Dong,Z. (2021) Targeting AURKA in Cancer: molecular mechanisms and opportunities for Cancer therapy. *Mol. Cancer*, **20**, 15.

50. Umene,K., Yanokura,M., Banno,K., Irie,H., Adachi,M., Iida,M., Nakamura,K., Nogami,Y., Masuda,K., Kobayashi,Y., *et al.* (2015) Aurora kinase A has a significant role as a therapeutic target and clinical biomarker in endometrial cancer. *Int. J. Oncol.*, **46**, 1498–1506.

51. Do,T.-V., Xiao,F., Bickel,L.E., Klein-Szanto,A.J., Pathak,H.B., Hua,X., Howe,C., O'Brien,S.W., Maglaty,M., Ecsedy,J.A., *et al.* (2014) Aurora kinase A mediates epithelial ovarian cancer cell migration and adhesion. *Oncogene*, **33**, 539–549.

52. Pérez-Fidalgo,J.A., Gambardella,V., Pineda,B., Burgues,O., Piñero,O. and Cervantes,A. (2020) Aurora kinases in ovarian cancer. *ESMO Open*, **5**, e000718.

53. Shah,K.N., Bhatt,R., Rotow,J., Rohrberg,J., Olivas,V., Wang,V.E., Hemmati,G., Martins,M.M., Maynard,A., Kuhn,J., *et al.* (2019) Aurora kinase A drives the evolution of resistance to third-generation EGFR inhibitors in lung cancer. *Nat. Med.*, **25**, 111–118.

54. Takai,N., Desmond,J.C., Kumagai,T., Gui,D., Said,J.W., Whittaker,S., Miyakawa,I. and Koeffler,H.P. (2004) Histone Deacetylase Inhibitors Have a Profound Antigrowth Activity in Endometrial Cancer Cells. *Clin. Cancer Res.*, **10**, 1141–1149.

55. Psilopatis,I., Pergaris,A., Giaginis,C. and Theocharis,S. (2021) Histone Deacetylase Inhibitors: A Promising Therapeutic Alternative for Endometrial Carcinoma. *Dis. Markers*, **2021**, 7850688.

56. Moufarrij,S., Srivastava,A., Gomez,S., Hadley,M., Palmer,E., Austin,P.T., Chisholm,S., Diab,N., Roche,K., Yu,A., *et al.* (2020) Combining DNMT and HDAC6 inhibitors increases anti-tumor immune signaling and decreases tumor burden in ovarian cancer. *Sci. Rep.*, **10**, 3470.

57. Yang,H., Sun,B., Xu,K., He,Y., Zhang,T., Hall,S.R.R., Tan,S.T., Schmid,R.A., Peng,R.-W., Hu,G., *et al.* (2021) Pharmaco-transcriptomic correlation analysis reveals novel responsive signatures to HDAC inhibitors and identifies Dasatinib as a synergistic interactor in small-cell lung cancer. *eBioMedicine*, **69**.

58. Leonardi C.,G., Falzone,L., Salemi,R., Zanghì,A., Spandidos A.,J., Mccubrey A.,J., Candido,S. and Libra,M. (2018) Cutaneous melanoma: From pathogenesis to therapy (Review). *Int J Oncol*, **52**, 1071–1080.

59. Hodis,E., Watson,I.R., Kryukov,G.V., Arold,S.T., Imielinski,M., Theurillat,J.-P., Nickerson,E., Auclair,D., Li,L., Place,C., *et al.* (2012) A Landscape of Driver Mutations in Melanoma. *Cell*, **150**, 251–263.

60. Ghosh,A.P. and Sudarshan,S. (2016) Genetics of renal cancer: focus on MTOR. *Aging (Albany. NY).*, **8**, 421–422.

61. Miricescu,D., Balan Gabriela,D., Tulin,A., Stiru,O., Vacaroiu Adela,I., Mihai Andrada,D., Popa Constantin,C., Papacocea
Ioana,R., Enyedi,M., Sorin Andrei,N., *et al.* (2021) PI3K/AKT/mTOR signalling pathway involvement in renal cell carcinoma
pathogenesis (Review). *Exp Ther Med*, **21**, 540.

62. Qin,A., Musket,A., Musich,P.R., Schweitzer,J.B. and Xie,Q. (2021) Receptor tyrosine kinases as druggable targets in
glioblastoma: Do signaling pathways matter? *Neuro-Oncology Adv.*, **3**, vdab133.

63. Alexandru,O., Horescu,C., Sevastre,A.-S., Cioc,C., Baloi,C., Oprita,A. and Dricu,A. (2020) Receptor tyrosine kinase targeting in
glioblastoma: performance, limitations and future approaches. *Contemp. Oncol. Onkol.*, **24**, 55–66.

64. McFarland,J.M., Paolella,B.R., Warren,A., Geiger-Schuller,K., Shibue,T., Rothberg,M., Kuksenko,O., Colgan,W.N., Jones,A.,
Chambers,E., *et al.* (2020) Multiplexed single-cell transcriptional response profiling to define cancer vulnerabilities and
therapeutic mechanism of action. *Nat. Commun.*, **11**, 4296.

65. Vassilev,L.T., Vu,B.T., Graves,B., Carvajal,D., Podlaski,F., Filipovic,Z., Kong,N., Kammlott,U., Lukacs,C., Klein,C., *et al.*
(2004) In Vivo Activation of the p53 Pathway by Small-Molecule Antagonists of MDM2. *Science (80-.).*, **303**, 844–848.

66. Ben-David,U., Siranosian,B., Ha,G., Tang,H., Oren,Y., Hinohara,K., Strathdee,C.A., Dempster,J., Lyons,N.J., Burns,R., *et al.*
(2018) Genetic and transcriptional evolution alters cancer cell line drug response. *Nature*, **560**, 325–330.

67. Di Veroli,G.Y., Fornari,C., Wang,D., Mollard,S., Bramhall,J.L., Richards,F.M. and Jodrell,D.I. (2016) Combenefit: an interactive
platform for the analysis and visualization of drug combinations. *Bioinformatics*, **32**, 2866–2868.

68. Chia,S., Low,J.-L., Zhang,X., Kwang,X.-L., Chong,F.-T., Sharma,A., Bertrand,D., Toh,S.Y., Leong,H.-S., Thangavelu,M.T., *et al.*
(2017) Phenotype-driven precision oncology as a guide for clinical decisions one patient at a time. *Nat. Commun.*, **8**, 435.

69. Robinson,M.D., McCarthy,D.J. and Smyth,G.K. (2010) edgeR: a Bioconductor package for differential expression analysis of
digital gene expression data. *Bioinformatics*, **26**.

70. Korotkevich,G., Sukhov,V., Budin,N., Shpak,B., Artyomov,M.N. and Sergushichev,A. (2021) Fast gene set enrichment analysis.
bioRxiv, 10.1101/060012.

71. Botev,Z.I. and Kroese,D.P. (2008) An Efficient Algorithm for Rare-event Probability Estimation, Combinatorial Optimization, and
Counting. *Methodol. Comput. Appl. Probab.*, **10**, 471–505.