Demonstration that sublinear dendrites enable linearly non-separable computations

Romain D. Cazé[1,*], **Alexandra Tran-Van-Minh**[2], **Boris S. Gutkin**[3,*,***], **and David A. DiGregorio**[2,*,***]

[1]UMR 8520 CNRS, IEMN, Villeneuve d'Asqu, 59650, France
[2]Institut Pasteur, Université Paris Cité, Synapse and Circuit Dynamics Laboratory, CNRS UMR 3571, Paris, 75015, France
[3]Group for Neural Theory, Laboratoire des Neurosciences Cognitives et Computationelles INSERM U960, Ecole Normale Superieure PSL* University, Paris France; CCDM, ICN, NRU Higher School of Economics, Moscow, Russia
[*]Corresponding authors: romain.caze@univ-lille.fr, david.digregorio@pasteur.fr, boris.gutkin@ens.fr
[***]joint senior authors

ABSTRACT

Theory predicts that nonlinear summation of synaptic potentials within dendrites allows neurons to perform linearly non-separable computations (LNSCs). Using Boolean analysis approaches, we predicted that both supralinear and sublinear synaptic summation could allow single neurons to implement a type of LNSC, the feature binding problem (FBP), which does not require inhibition contrary to the XOR. Notably, sublinear dendritic operations enable LNSCs when scattered synaptic activation generates increased somatic spike output. However, experimental demonstrations of scatter-sensitive neuronal computations have not yet been described. Using glutamate uncaging onto cerebellar molecular layer interneurons, we show that scattered synaptic-like activation of dendrites evoked larger compound EPSPs than clustered synaptic activation, generating a higher output spiking probability. Moreover, we also demonstrate that single interneurons can indeed implement the FBP. Since sublinear synaptic summation is a property of passive dendrites we expect that many different neuron types can implement LNSCs.

Introduction

Dendritic filtering influences the shape and amplitude of postsynaptic potentials depending on the synaptic conductance, dendritic morphology, synapse location and the expression of voltage-gated channels[1]. Dendritic properties also influence how multiple synaptic potentials sum, either linearly (if the response amplitude equals the arithmetic sum of the individual synaptic responses), sub-linearly (response amplitude below the sum of the individual responses), or supra-linearly (response amplitude above their sum)[2]. This dendrite-specific arithmetic can greatly enhance a neuron's computational abilities[3] and is thought to be prominent in human neurons[4]. Supra-linear summation is thought to underlie active whisker sensation[5], generate orientation selectivity[6], grid cell activity[7], and sensory perception[8]. In contrast, little is known about the computational advantages imparted by sub-linear dendritic integration[9, 10]. Recent computational studies suggest that sub-linear integration in cortical fast-spiking interneurons could contribute to memory encoding[11].

Our previous theoretical work showed that the class of computations performed by neurons with sub-linear dendrites was on the order of that for supra-linear dendrites and is vastly larger than those accessible for excitatory synapses onto neurons with only linearly summating dendrites[12]. These additional computations are, by definition, a part of the class of linearly non-separable functions .

While LNSCs cannot be implemented by a classic neuron model, a neuron with sublinear dendrites can easily implement an LNSC called the feature binding problem (FBP) (Fig 1). The FBP computation allows a neuron to represent multiple distinct objects, each comprised of independent features. An example of FBP is given in Figure 1 where the goal is to detect a yellow circle or a green triangle separately by giving an output of 1, but not respond to various other different conjunctions of features such as a yellow triangle or a green circle (Fig 1). We note that this distinction cannot be done by linearly separating the inputs based on their summed amplitudes. Because the FBP can be implemented with only excitatory synapses, we used glutamate uncaging at single synapses to demonstrate the FBP experimentally.

To implement the FBP, supralinear dendrites require that the correct feature conjunction for an object cluster onto the same dendritic compartment, while another feature conjunction that corresponds to a different object would innervate a second

Figure 1. The Feature Binding Problem: A linearly inseparable computation and its implementation in a scatter sensitive neuron.(A) A truth table showing the implementation of the FBP. There are four input neurons carrying four different features Circle (Ci), Triangle (Tr), Yellow (Ye), and Green (Gr). Neurons that are active are idnidcated with a "1". The "Int" column describes the resulting integration for each dendrite. Both dendrites must receive a 1 for the soma to reach its threshold of 2, thereby firing a spike. Because Ci+Gr and Tr+Ye target the same dendrite, the resulting integration fails to reach the depolarization necessary to spike and thus cannot reach the threshold at the soma. Yet for Ci+Ye and Tr+Gr, each dendrite generates a one and the soma surpasses its threshold. Panel C indicates that features activating the same dendrites do not reach threshold and cannot generate a spike, but those features scattered onto the two dendrites generate a somatic spike.

dendrite. However, sublinear dendrites allow that the features of a single object be "scattered" onto different dendrites to drive a somatic spike since the compound EPSP produced by innervation of the same dendrites would be sublinear and below the somatic threshold[1, 12]. Thus the case where scattered inputs fire the cell and clustered inputs do not is essential for solving the FBP using sublinear dendrites (Fig 1C). Here we used glutamate uncaging and biophysical modeling to show that the sublinear dendrites of cerebellar molecular layer interneurons render them scatter-sensitive, thereby enabling them to implement the FBP.

Results

Cerebellar stellate cells are scatter sensitive

Dendrites of cerebellar stellate cells are thin and integrate synaptic conductances passively[9]. The high-impedance dendrites generate large local depolarization upon excitatory synaptic activation that reduce the driving force for synaptic currents, resulting in a sublinear summation of postsynaptic potentials when activating multiple synapses[13]. Therefore, when synapses on the same sublinear dendrite are activated simultaneously, they "interact" and thus summate less effectively than if synapses were activated on two different dendrites, due to a lack of effective electrical interaction. In other words, the somatic depolarization is larger and more sensitive to synapse activation patterns that are "scattered" across the dendritic tree.

We used scanning two-photon glutamate uncaging[13, 14] in parasagittal cerebellar brain slices to test the hypothesis that neurons with sublinear dendrites generate larger excitatory potentials when synaptic activation is distributed across dendrites rather than synaptic activation within a dendrite (Fig 2). We identified stellate cells by patching somata located in the outer third of the molecular layer. To mimic clustered activation, we uncaged glutamate using 0.2 ms 720 nm laser pulses in four locations within putative stellate cell dendritic trees in any one trial (slices were superfused wth 2 mM MNI-glutamate) and recorded uncaging-evoked excitatory postsynaptic potentials (uEPSPs) using somatic current clamp patch recordings. Current was injected into the soma to maintain a resting potential near -70mV. Either all uncaging locations were clustered in the same dendrite, or two uncaging locations were placed on two different primary dendrites (see Fig 2A). We used pairs of uncaging sites to increase the local depolarization. Compound subthreshold uEPSPs were generated from four sites ranging from 4-22 mV across all experiments (Fig 2C). We observed that compound uEPSPs evoked on the same dendritic branch were systematically smaller than their linear sum, as expected for sublinear integration (Fig 2B, C). However, compound uEPSPs generated from uncaging locations scattered onto two different dendrites were not significantly different than their linear sum and thus consistent with scatter-sensitive subthreshold PSPs. Because of the inertia of galvanometer mirrors, we could more easily perform simultaneous activation of three dendrites if one was stimulated using an extracellular electrode (see Methods). Results were similar for the activation of effectively six uncaging sites across three dendrites (Fig 2D-F). These results confirm that compound synaptic potentials in neurons with sublinear dendrites were larger if the inputs were scattered across dendrites rather than clustered on a single dendrite.

Unlike most studies that focus on the characterization of subthreshold dendritic operations, we set out to verify that the scatter-sensitive subthreshold behavior of stellate cell dendrites also translated into scatter-sensitive spiking probability. We applied the same uncaging protocol described for Fig 2, but adjusted the holding current to maintain the resting potential around -60 mV, to facilitate spiking. Indeed, we found that uncaging locations that were located on the same dendrite produced a significantly lower spiking probability than when uncaging locations were on separate dendrites (Fig 3). These data demonstrate

Figure 2. Scattered inputs sum more linearly than clustered inputs. (A, C) 2PLSM image of a cerebellar stellate cell filled with Alexa Fluor 594. Colored spots indicate dendritic locations stimulated using glutamate uncaging (branches 1 and 2, A and C) or electrical stimulation (branch 3, C). (B, D) Example traces of somatic current-clamp recordings in response to stimulation of a group of clustered (left) of scattered (right) inputs (blue traces), and traces obtained as the sum of EPSPs recorded in response to stimulation of individual locations (black traces). (E, F) Peak amplitudes of observed and expected EPSPs in response to stimulation of clustered or scattered inputs.

that stellate cells are more likely to fire when inputs are distributed over their dendritic tree than when they cluster on specific dendrites. We also extended this study to three-branch stimulation as described above and observed a robust increased firing probability if the stimulation was distributed across the dendrites (Fig 3C, D, and F). Thus, the neuronal computation (as represented by somatic spiking) of cerebellar stellate cells is scatter-sensitive. It remains to be demonstrated that the FBP can be computed by the neuron with scatter-sensitive spike production.

Figure 3. Multi-branch glutamate uncaging reveals a higher firing probability for scattered versus clustered stimulation. (A) 2PLSM image of a cerebellar stellate cell filled with Alexa Fluor 594. Colored spots indicate locations stimulated using glutamate uncaging (branch 1 and 2) or electrical stimulation (branch 3). (B-C) Example traces of somatic current-clamp recordings of EPSPs and action potentials in response to stimulation of a group of clustered or scattered inputs on 2 (B) or 3 (C) branches. The combinations of spots stimulated by glutamate uncaging are indicated above the trace, with clustered combinations in orange and scattered combinations in gray. Left, trials for which the neuron's somatic response corresponds to a linearly non-separable function. Right, trials for which the neuron did not compute a linearly non-separable function. (D-E) Spike probabilities in response to clustered and scattered synaptic inputs.

Since we established that stellate cells are scatter-sensitive, we next examined whether such scatter sensitivity would confer them the ability to perform linearly non-separable computations. Specifically, we set out to show they could implement the FBP (see Fig 4). Glutamate was uncaged at four distinct locations in six different patterns. There were four scattered cases (uncaging sites on different dendrites) and two clustered cases (uncaging sites on the same dendrite). To observe an FBP we examined whether a neuron remained silent (no action potential) in the two clustered cases and fired in two scattered cases with non-overlapping uncaging sites. Let us consider each uncaging location as a synaptic input encoding an object's feature, then combinations of uncaging locations that trigger a somatic spike effectively encode the object. If we observed that the neuron fires for specific combinations of non-overlapping uncaging sites and stays silent for others, then we can claim that it implements the FBP. To give a proof-of-principle demonstration, we chose to label each uncaging location with a unique symbol shape or color: triangle or a circle shape, and green vs. yellow color. Given such labeling, we would conclude that the feature binding problem was correctly implemented if a green triangle and yellow circle were encoded by the soma. In our experiments,

we observed multiple trials in which the results were consistent with the FBP, while in others they were not (either scattered synaptic activation did not produce a spike or the clustered case did) (see Fig 4 lower right, red shading). The cell shown in Figure 4 implemented the FBP in nearly half of the trials (n=15), i.e. it fired for two objects with two specific disjoint features (e.g. green triangle or yellow circle) and stayed silent for two other objects made up of other disjoint feature combinations (the green circle and yellow triangle). We also observed, in this cell, trials where a false positive spike was generated (red shading Fig 4). Because of the costly experimental setup we managed to perform suprathreshold experiments only on four cells, where we stimulated a couple of dendrites. Two cells were capable of implementing the FBP 50% of the time. To strengthen this experimental work and explain cases where cells fail to implement the FBP we used a realistic biophysical model.

Figure 4. Cerebellar stellate cells might implement the feature binding problem (FBP). (A) 2PLSM image of a cerebellar stellate cell filled with Alexa Fluor 594. Colored spots indicate the uncaging locations, on this cell we uncaged glutamate in four locations, each pair corresponds to one feature (Purple:"Green" on branch 1, "Triangle" on branch 2; Teal:"Yellow" on branch 2, "Circle" on branch 1). The color code is in the top of panel B.and indicates the two objects to be encoded (yellow circle, green triangle). (B) Examples of somatic voltage traces of the same stellate cell in two distinct trials. In the top trial, the neuron computes a feature binding problem, and in the bottom trial it fails to do it because of a false positive (red case). We discarded the two cases when inputs are coming from the shape or the color inputs only as they do not affect the result.

Scatter sensitive neurons can compute the feature binding problem

To understand better the origin of the failed FBP trials, We implemented a biophysical model of a cerebellar stellate cell. We built a multi-compartment model with a realistic dendritic tree morphology (using a reconstruction of cerebellar stellate cell[15]) that was capable of reproducing the uEPSP amplitudes observed in the subthreshold protocol(see Fig 5, 4 pS peak synaptic conductance). The dendrites were modeled as passive and showed sublinear EPSP summation with clustered inputs and more linear summation with inputs scattered on different dendrites (Fig 5A and B). We considered two principal factors that could underlie trial-to-trial variability as well as cell variability in the ability to implement the FBP: EPSP size relative to the threshold and membrane fluctuations. To correctly implement the FBP, the voltage differential from rest to threshold must be such that the single uEPSP and the sublinear compound uEPSP (two-site activation within the same dendrite) are subthreshold, but the linearly summed uEPSPs (two-site activation on different dendrites) are suprathreshold. We thus varied the resting membrane voltage and examined if simulated compound uEPSPs (adjusted to 10 pS per site to ensure spikes) produced false positive or negative spikes (compare Fig 5 to Fig 4). For the depolarized resting membrane potential of -64mV the model reaches the threshold not only in all the scattered stimulation cases but also in one of the clustered cases creating a false positive. When the membrane potential was decreased to -68mV the neuron reached the threshold only in the scattered case, thus correctly implementing the FBP. Finally, if the resting membrane potential was set too low (-72mV), the neuron spiked only in one of the scattered cases. Together, these three simulations exemplify that variations in the resting membrane voltage can influence the successful computation of the FBP. Thus, the EPSP size relative to the threshold must be tuned correctly to implement the FBP.

We also recognized that membrane potential fluctuations could generate false positives and false negatives. We, therefore, modeled the voltage fluctuations using a Gaussian noise distribution that matched experimental observations (mode at -68.4 mV with a standard deviation equal to 0.9 mV vs. resting potential mean for the cell in (Fig 6) was -68.5 mV with an SD of 0.8 mV (Fig 5D)). We, therefore, systematically varied the noise level around this experimental value as well as varying the resting membrane potential of the model from -72mV and -64mV (Fig 5E). To calculate the FBP probability we performed 1000 trials for each simulation condition. As expected from above, when the resting membrane potential is too high the neuron generated false positive spikes, and when it is too low, the neuron generated false negative errors. The simulations indicated

Figure 5. **Simulations exploring synaptic and cell parameters necessary to perform FBP** (A) Morphologically detailed model of a cerebellar stellate cell, with colored circles indicating input locations corresponding to the two combinations of features (teal:yellow+circle;purple:green+triangle. (B) compound uEPSPs were simulated by placing two 4 pS synaptic conductance (max value) with a $\tau = 1ms$ time constant on the same (clustered) or different dendrites (scattered). (C) Membrane potential responses for compound uEPSP responses when stimulating pairs of uncaging locations as indicated in the diagram. The dashed line is the the threshold for spike generation. Note that only a resting membrane potential of -68 mV allows for a correct implementation of the FBP, the two other voltages lead to errors in the FBP implementation- being due to respectively a false positive (extraneous spike) and a false negative (absence of a spike). (D) The experimental distribution of the voltage (black, μ=-68.5 mV, σ=0.8 mV). We reproduced this distribution using the same mean and standard deviation normal distribution (light blue superimposed). (E) Probability of implementing the FBP for different resting membrane potentials and noise levels (SD), calculated from 1000 simulations for each condition.

an optimum around a resting membrane potential of -68mV, but the FBP probability degraded within increasing membrane potential fluctuations.

We were able to show that by adjusting the synaptic conductance in the model to match the uEPSP amplitudes for each individual uncaging site, as well as the membrane potential fluctuations, we could reproduce the probability of implementing (around 50%), see (Fig 5). We, therefore, conclude that scatter-sensitive neurons can perform the FBP, but the relative amplitude of the EPSP versus the voltage-to-threshold and the membrane fluctuations need to be tuned to reliably compute the FBP.

Based on the analysis above, we can now demonstrate that a cerebellar stellate cell can indeed perform the FBP even in noisy experimental conditions, as we can see the cell (Fig 6A-B top) performs the FBP in approximately 50 percent of the trials as is also shown by our model (Fig 6A bottom). We note that the model further matches the error trials in the neuron giving spurious spikes at the inputs encoding the yellow triangle (Fig 6B). This enabled us to computationally reproduce the variation observed experimentally (Fig 6C) and observe the neuron implementing the FBP in seven trials over fifteen, thereby matching the experimentally observed variability in FBP implementation. More generally, out of the 13 neurons we subjected to this protocol, we observed two cells that robustly implemented the FBP with a probability of approx 0.5 (cell shown 7/15 trials, cell not shown 6/11 trials).

In Fig 6 we used the same biophysical model as described above with four synaptic conductances placed as indicated (Fig 5) with peak amplitudes of Branch 1 (Br1) teal:7 pS, purple:3 pS; Branch 2 (Br2) teal:20 pS, and purple:10 pS, to match as close as possible the experimental data. The two first stimulation points have a weaker value than the two last as the second uEPSP is

Figure 6. **Model simulations reproduce measured FBP probability**. (A) Voltage distribution of the first trial for the first 100ms of simulation (B) Top, somatic voltage trace recorded experimentally. Here the cell was stimulated by uncaging at two pairs of locations per dendrite and two dendrites, giving a total of 4 location pairs and each corresponding to a feature (7 and 3 pS on the first dendrite resp. Circle, Green; 10 and 20 pS for the 2nd standing for Triangle, Yellow). Locations were co-uncaged to provide inputs with distinct feature-pair conjunctions. We observe spiking when we uncage at the locations tagged as yellow-circle and green-triangle conjunctions. In this way the cell implements the FBP for yellow circle and green triangle objects. we also see that the cell gives falls positive spikes on several trials when the yellow-triangle conjunction is stimulated, but not on the green-circle. Bottom: Model simulations with -68.5 mV resting potential and a noise level of $\sigma = 0.6mV$. EPSP amplitudes were adjusted to match experimental values (C) Top, the obtained experimental result from the cell in Fig. 4 with all trials plotted. Trials in black are those in which the FBP was correctly implemented. Bottom, are 15 simulated trials showing a similar FBP probability.

larger than the first. The third input with a high value guarantees that the neuron will fire in the third scattered case (20+3pS) while the sum is lower in the second scattered case (10+7pS) keeping the neuron silent in this situation.

Discussion

It is well known that dendritic nonlinearities can theoretically increase the computational capacity of neurons because they enable multi-layered information processing. In particular, while there has been fast-growing theoretical literature arguing for universal computing power of dendritic neurons[3, 11, 12, 16] (bolstered by experimental results on subthreshold dendritic integration[1, 2, 4, 5] and the potential role of dendritic processes in neuronal tuning properties[8, 17–19]), experimental validation of an actual implementation of LNSC by a neuron endowed with nonlinear dendrites, to our knowledge, has not been performed. In this study, we demonstrated using multi-point glutamate uncaging that the sublinear dendritic integration of cerebellar stellate cells leads to larger EPSPs and spiking probabilities when synaptic activation is spread across dendrites (scatter-sensitive) as compared to when the synapses are activated within the same dendritic branch. We also showed that SCs can implement the feature binding problem (as previously predicted in[12]). Of course, sublinear dendrites cannot implement all LNSCs. The XOR, for example, can only be implemented with inhibition or with a non-monotone transfer function. Although, unlike the XOR, the FBP can be computed using only excitatory synapses. The FBP is of particular interest since neurons implementing it are able to bind multiple feature combinations together into an object. Modeling dendritic integration of the thin passive dendrites of cerebellar stellate cells allowed us to identify the biophysical conditions under which a neuron could implement the FBP.

The FBP is of particular computational interest since a neuron implementing it can detect multiple objects made of multiple features whereas a neuron lacking non-linear dendrites can only detect a single object. We have shown here that in order to implement the FBP using excitatory synapses and sublinear dendrites, the two features of a common object must innervate two different dendrites such that their compound EPSP, upon simultaneous activation, is maximized and generates the large spike probability. Otherwise, if two synapses are simultaneously activated within the same dendritic compartment, and their summed local depolarization is large enough to decrease the driving force for ionic currents, the net somatic depolarization will be smaller than the arithmetic sum. Thus clustered synaptic activation produces less depolarization than when the synapses are electronically independent. For a neuron to bind two (or more) features of an object, the features must be distributed across the dendritic tree to maximize the spike probability associated with the object, a computation termed "scatter-sensitive". However, an important condition is that the scattered activation produces a somatic spike, whereas the equivalent clustered activation must be subthreshold for spike generation. This fine-tuning requires a specific relation between the single synapse EPSP amplitude, the number of synapses associated with the different features, and the difference between the resting membrane potential and the spike threshold. EPSPs that are too strong generate false positives, whereas EPSPs that are too weak generate false negatives spiking representations of objects (Fig 5). Stronger background membrane potential fluctuations will also generate more false

positives and negatives on a trial-to-trial basis (see Fig 4 and Fig 6). Synaptic plasticity and intrinsic dendritic modulation may help to train a neuron to implement the FBP, a suggestion that could be tested in future studies.

In this study, we emphasized scatter-sensitivity of the neuron as a key factor for the implementation of linearly non-separable computations with passive dendrites. Passive thin dendrites have high input impedances and are good candidates for large local depolarizations that generate sublinear summation. Activation of voltage-gated potassium conductances can also generate sublinear integration[10]. Moreover, it is possible that for small numbers of active synapses, their compound EPSP can be large enough to decrease the driving force locally in the dendrite, but still too small to activate nonlinear conductances[20]. In that window of synaptic depolarization, a scatter-sensitive computation can be performed. Moreover, for dendrites that exhibit robust calcium spikes[9], once the dendritic spike has been generated, additional synaptic depolarization is ineffective in increasing spike probability. Synaptic contacts scattered onto other dendrites must be activated to generate additional somatic depolarization and increased spike probability. Thus both interneurons and principal neurons could exhibit scatter sensitivity and thus use sublinear summation to compute the FBP[11], albeit under specific regimes of synaptic and intrinsic cell membrane parameters. Given that scatter sensitivity responses on passive cable properties of dendritic trees, and is not catastrophically deteriorated by active dendritic processes (albeit under different synaptic and intrinsic regimes), it might be a widely generic property of information processing in wide classes of neurons.

Methods

Animals and slice preparation

Animal experiments were performed in accordance with the guidelines of Institut Pasteur, France, and all protocols were approved by the ethics committee CEEA-Ile de France-Paris 1. Cerebellar acute slices were prepared from CB6F1 mice (F1 cross of BalbC and C57Bl/6J) of postnatal day P60 to 90. The mice were killed by rapid decapitation, after which the brains were quickly removed and placed in an ice-cold solution containing (in mM): 2.5 KCl, 0.5 CaCl2, 4 MgCl2, 1.25 NaH2PO4, 24 NaHCO3, 25 glucose, 230 sucrose, and 0.5 ascorbic acid bubbled with 95% O2 and 5% CO2. Parasagittal slices (20 0μm thick) were prepared from the dissected cerebellar vermis using a Leica VT1200S vibratome. After preparation, the slices were incubated at 32 degree Celsius for 30 minutes in the following solution (in mM): 85 NaCl, 2.5 KCl, 0.5 CaCl2, 4 MgCl2, 1.25 NaH2PO4, 24 NaHCO3, 25 glucose, 75 sucrose and 0.5 ascorbic acid. Slices were then transferred to an external recording solution containing (in mM): 125 NaCl, 2.5 KCl, 2 CaCl2, 1 MgCl2, 1.25 NaH2PO4, 25 NaHCO3, 25 glucose and 0.5 ascorbic acid, bubbled with 95% O2 and 5% CO2, and maintained at room temperature for up to 7 hours.

Slice electrophysiology and imaging

Whole-cell current-clamp recordings were performed from stellate cells (SCs) (33°C – 36°C) located in the outer third of the molecular layer, using a Multiclamp 700B amplifier (Molecular Devices), and fire-polished thick-walled glass patch-electrodes (tip resistances of 4-6 MΩ). The pipettes were backfilled with an internal solution containing (in mM): 110 K-MeSO3, 40 HEPES, 1 EGTA, 4.5 MgCl2, 0.49 CaCl2, 10 Na-pyruvic acid, 0.3 NaGTP, 4 NaATP, 10 Tris phosphocreatine and 0.04 Alexa Fluor 594 and adjusted to 305 mOsm and pH 7.3. Synaptic responses were filtered at 10 kHz, and digitized at 100 kHz using an analogue-to-digital converter (model NI USB 6259, National Instruments, USA) and acquired with NClamp (www.neuromatic.thinkrandom.com), running in the Igor Pro environment (Wavemetrics). Current was injected to maintain the membrane potential between -70mV and -90mV (after correcting for liquid junction potentials, calculated to be -7 mV using JPCalcW (Barry, 1994; J. Neurosci. Method., 51: 107-116)), and series resistance was compensated by balancing the bridge and compensating pipette capacitance.

Unless otherwise stated, the external solution included 10 μM SR-95531 to block GABAA receptors to avoid confounding results due to partial blockade of GABAA receptors by MNI glutamate, and 50 μM D-AP5 to avoid stimulation of extra-synaptic NMDA receptors.

A pulsed Ti:Sapphire laser (DeepSee, Spectra-Physics) beam tuned at 810 nm was scanned on the preparation using an Ultima microscope (Bruker Fluorescence Microscopy) mounted on an Olympus BX61WI microscope and equipped with a 60x (1.1 NA) water-immersion objective. Simultaneous two-photon fluorescence and Dodt contrast imaging (Luigs and Neumann, Germany) were used to position extracellular stimulating electrodes and uncaging points along spatially isolated dendrites of Alexa Fluor-594-filled SCs, using a transmitted light PMT mounted after the Dodt tube to acquire a laser-illuminated contrast image simultaneously with the 2PLSM image. Alexa Fluor 594 fluorescence was filtered using 640/100 nm bandpass filters (Chroma) and detected using side-on multi-alkali PMTs (3896, Hamamatsu Photonics). In addition to the light collected through the objective, the transmitted infrared light was collected through a 1.4 NA oil-immersion condenser (Olympus), and reflected on a set of substage photomultiplier tubes (PMTs).

Chemicals

D-AP5 (D-(-)-2-Amino-5-phosphonopentanoic acid) and SR 95531 (2-(3-Carboxypropyl)-3-amino-6-(4 methoxyphenyl) pyridazinium bromide) were purchased from Abcam, UK. MNI-glutamate (4 methoxy-7-nitroindolinyl caged L-glutamate) was purchased from Tocris Bioscience, UK. Alexa Fluor 594 was from Life Technologies, USA. All other chemicals were purchased from Sigma-Aldrich, France.

Glutamate uncaging

MNI-glutamate was bath applied at a final concentration of 2 mM in ACSF, and recycled. The solution was kept protected from light and any lamps used for ambient light and microscope trans-illumination used during approach of the patch-clamp electrode were covered with a UV yellow filter to prevent undesired photolysis of MNI-glutamate. The preparation was illuminated through a second set of galvanometer-based scan mirrors, allowing independent and rapid positioning of the photolysis beam. The photolysis laser was either a 1P laser at 405 nm diode laser (Omicron) or a pulsed Ti:Sapphire laser (DeepSee, Spectra-Physics) tuned at 720nm. The outputs of the two lasers were independently modulated to combine uncaging of MNI-glutamate and imaging morphology. The imaging laser beam was modulated using a Pockels cell (350-50-BK 02, Conoptics, Danbury, CT). For 2P uncaging, the intensity and duration (200-300 μs) of the photolysis pulse was modulated using an acousto-optic modulator (MT110-B50A1.5-IR-Hk, AA Opto-Electronic, France). For 1P uncaging, the intensity and duration of the photolysis pulse were. A telescope placed on the path of each uncaging beam (Thorlabs) was used to adjust the convergence angle to both backfill the objective and match the focal plane of the two-photon excitation for imaging. Parfocality of the two beams was verified using bleached spots on a microscope slide coated with fluorescent ink. Photolysis laser powers, estimated at the exit of the objective were <1 mW for 1P uncaging and <20 mW for 2P uncaging. We uncaged near simultaneously at up to 6 spots, each 3-6 μm apart, by rapidly switching the focal spot to each new location along the dendrite at 200 μs intervals. For multi-dendrites stimulation, the minimum displacement time of uncaging mirrors between two uncaging locations constrained the distance between the stimulated dendrites (typically in the order of 100-150 μm between the most distant uncaging locations).

Distribution of the resting membrane voltage

The voltage membrane is sampled at 1kHz. We looked at all the 15 trials and used the first 100ms to plot an histogram of the voltage distribution. We have overlapped surrogate data (n=150000 points) generated from a normal distribution using the observed μ=68.48 and σ=0.90. We have used a visual verification to test the normality of the distribution as usual statistical test (Shapiro or Kolmogorov) were not adapted for the extended number of data points and as we approximated the distribution as a normal.

Biophysical model

All simulations were performed with Brian 2[21] and we used a detailed reconstruction from[15]. The code needed to reproduce the last figure is available under an open-source licence. We used a conductance-based model with the reconstructed dendritic arbor. The axial resistance between compartments equals 150Ω. Every compartment contained the following passive currents:

$$C_m \frac{dV}{dt} = g_L(V - E_L) + I_s \tag{1}$$

V is the membrane potentials, $C_m = 1\,\mu\mathrm{F\,cm^{-2}}$ is the membrane capacitance, g_L is the leak capacitance equal to $5e-5$siemens which is equivalent to a $R_i = 20000\Omega$ input resistance.

The synaptic current I_s is described by

$$I_s = g_s(E_s - V) \tag{2}$$

with E_s being the synaptic reversal potential and g_s the synaptic conductance. This conductance jumps up instantaneously for each incoming input and decays exponentially with time constant $\tau_s = 1\,\mathrm{ms}$ otherwise:

$$\frac{dg_s}{dt} = -\frac{g_s}{\tau_s} \tag{3}$$

The neuron "spikes" when it reaches -50mV from below and we signal a spike using a vertical bar. We set the resting membrane voltage to -68mV.

Simulations to match the experimental observation

In Fig 6 we used the same biophysical model as described above with four synaptic conductances placed as indicated (Fig 5) with peak amplitudes of Br1 teal:7 pS, purple:3 pS; Br2 teal:20 pS, and purple:10 pS exponentially decaying with a 1 ms time constant, to match as close as possible the experimental data (uEPSP amplitude or spike probability). The two first stimulation points have a weaker value than the two last as the second uEPSP is larger than the first. The third input with a high value guarantees that the neuron will fire in the third scattered case (20+3pS) while the sum is lower in the second scattered case (10+7pS) keeping the neuron silent in this situation.

To mimic the observed variation in the membrane resting voltage we used two conductances based processes, one inhibitory and one excitatory of 0.2pS targeting the soma. To create randomly distributed membrane voltage in the model we used the following procedure. We randomly picked 400 integers between 0 and 2000 each with the same probability. Of these 200 corresponded to the times at which an excitatory synapse activates (with the reversal potential of 0mV) and the other 200 to the activation times of an inhibitory synapse (reversal potential -140mV). Both of these surrogate synapses target the soma and have a conductance strengths of 0.2nS. Each synapse was modeled as an instantaneous increase preceding an exponential decay with a 1ms time constant. The goal was to obtain a resting potential distribution that matched the experimentally observed one (see Fig 6A).

The python code to entirely reproduce the two last figures and subpanels is available on zenodo[22].

References

1. Tran-Van-Minh, A. *et al.* Contribution of sublinear and supralinear dendritic integration to neuronal computations. *Front. Cell. Neurosci.* **9**, DOI: 10.3389/fncel.2015.00067 (2015).

2. Polsky, A., Mel, B. W. & Schiller, J. Computational subunits in thin dendrites of pyramidal cells. *Nat. Neurosci.* **7**, 621–627, DOI: 10.1038/nn1253 (2004).

3. Poirazi, P., Brannon, T. & Mel, B. W. Pyramidal Neuron as Two-Layer Neural Network. *Neuron* **37**, 989–999, DOI: https://doi.org/10.1016/S0896-6273(03)00149-1 (2003).

4. Gidon, A. *et al.* Dendritic action potentials and computation in human layer 2/3 cortical neurons. *Science* **367**, 83–87, DOI: 10.1126/science.aax6239 (2020).

5. Xu, N.-l. *et al.* Nonlinear dendritic integration of sensory and motor input during an active sensing task. *Nature* **492**, 247–251, DOI: 10.1038/nature11601 (2012).

6. Wilson, D. E., Whitney, D. E., Scholl, B. & Fitzpatrick, D. Orientation selectivity and the functional clustering of synaptic inputs in primary visual cortex. *Nat. Neurosci.* **19**, 1003–1009, DOI: 10.1038/nn.4323 (2016).

7. Schmidt-Hieber, C. & Nolan, M. F. Synaptic integrative mechanisms for spatial cognition. *Nat. Neurosci.* **20**, 1483–1492, DOI: 10.1038/nn.4652 (2017).

8. Takahashi, N., Oertner, T. G., Hegemann, P. & Larkum, M. E. Active cortical dendrites modulate perception. *Science* **354**, 1587–1590, DOI: 10.1126/science.aah6066 (2016).

9. Abrahamsson, T., Cathala, L., Matsui, K., Shigemoto, R. & DiGregorio, D. A. Thin Dendrites of Cerebellar Interneurons Confer Sublinear Synaptic Integration and a Gradient of Short-Term Plasticity. *Neuron* **73**, 1159–1172, DOI: 10.1016/j.neuron.2012.01.027 (2012).

10. Hu, H., Martina, M. & Jonas, P. Dendritic Mechanisms Underlying Rapid Synaptic Activation of Fast-Spiking Hippocampal Interneurons. *Science* **327**, 52–58, DOI: 10.1126/science.1177876 (2010).

11. Tzilivaki, A., Kastellakis, G. & Poirazi, P. Challenging the point neuron dogma: FS basket cells as 2-stage nonlinear integrators. *Nat. Commun.* **10**, 3664, DOI: 10.1038/s41467-019-11537-7 (2019).

12. Cazé, R. D., Humphries, M. & Gutkin, B. Passive Dendrites Enable Single Neurons to Compute Linearly Non-separable Functions. *PLOS Comput. Biol.* **9**, e1002867, DOI: 10.1371/journal.pcbi.1002867 (2013).

13. Tran-Van-Minh, A., Abrahamsson, T., Cathala, L. & DiGregorio, D. A. Differential Dendritic Integration of Synaptic Potentials and Calcium in Cerebellar Interneurons. *Neuron* **91**, 837–850, DOI: 10.1016/j.neuron.2016.07.029 (2016).

14. Tran-Van-Minh, A., Rebola, N., Hoehne, A. & DiGregorio, D. A. Two-Photon Neurotransmitter Uncaging for the Study of Dendritic Integration. In Hartveit, E. (ed.) *Multiphoton Microscopy*, Neuromethods, 33–64, DOI: 10.1007/978-1-4939-9702-2_3 (Springer, New York, NY, 2019).

15. Scorcioni, R., Polavaram, S. & Ascoli, G. A. L-Measure: a web-accessible tool for the analysis, comparison and search of digital reconstructions of neuronal morphologies. *Nat. Protoc.* **3**, 866–876, DOI: 10.1038/nprot.2008.51 (2008).

16. Ujfalussy, B. B., Makara, J. K., Lengyel, M. & Branco, T. Global and Multiplexed Dendritic Computations under In Vivo-like Conditions. *Neuron* **100**, 579–592.e5, DOI: 10.1016/j.neuron.2018.08.032 (2018).

17. Smith, S. L., Smith, I. T., Branco, T. & Häusser, M. Dendritic spikes enhance stimulus selectivity in cortical neurons in vivo. *Nature* **503**, 115–120, DOI: 10.1038/nature12600 (2013).

18. Palmer, L. M. *et al.* NMDA spikes enhance action potential generation during sensory input. *Nat. Neurosci.* **17**, 383–390, DOI: 10.1038/nn.3646 (2014).

19. Fischer, L. *et al.* Dendritic Mechanisms for In Vivo Neural Computations and Behavior. *J. Neurosci.* **42**, 8460–8467, DOI: 10.1523/JNEUROSCI.1132-22.2022 (2022). Publisher: Society for Neuroscience Section: Symposia.

20. Losonczy, A. & Magee, J. C. Integrative properties of radial oblique dendrites in hippocampal CA1 pyramidal neurons. *Neuron* **50**, 291–307, DOI: 10.1016/j.neuron.2006.03.016 (2006).

21. Stimberg, M., Brette, R. & Goodman, D. F. Brian 2, an intuitive and efficient neural simulator. *eLife* **8**, e47314, DOI: 10.7554/eLife.47314 (2019).

22. Caze, R. Demonstration that sublinear dendrites enable linearly non-separable computations, DOI: 10.5281/ZENODO.7875425 (2023). Language: en.

Acknowledgements

B.S.G. would like to acknowledge the support from the Basic Science Program Basic Research Program at the National Research University Higher School of Economics (HSE University). This work was partially supported by the ANR projects "CerebComp" and "InTempCode".

Author contributions statement

R.C., A.T. and D.A.D. conceived the experiments, R.C., D.A.D. and B.S.G. conceived the theory, R.C. carried out the computational research A.T. conducted the experiments, R.C. and A.T. analysed the results, D.A.D, B.S.G and R.C. wrote the manuscript. All authors reviewed the manuscript.