Exoribonuclease RNase R protects Antarctic *Pseudomonas syringae* Lz4W from DNA damage and oxidative stress

Running Title: Role of RNase R in protecting against DNA damage and oxidative stress.

Pragya Mittal[1, 2,*], Rashmi Sipani[1, 3], Apuratha Pandiyan[1, 4] Shaheen Sulthana[1, 6], Anurag K Sinha[1, 5], Ashaq Hussain[1], Malay K Ray[1] and Theetha L Pavankumar[1, 6, *].

Affiliations:

[1] Centre for Cellular and Molecular Biology (CCMB), Council of Scientific and Industrial Research (CSIR), Uppal Road, Hyderabad-500007, India.

Present addresses

[2] MRC Human Genetics Unit, Institute of Genetics and Cancer, University of Edinburgh, EH4 2XU Edinburgh, United Kingdom.

[3] PopVax Pvt. Ltd., IV Floor, Atal Incubation Centre-CCMB Annex-II, Medical Biotechnology Complex, Hyderabad, India.

[4] Department of Biological Sciences, Indian Institute of Science Education and Research (IISER) Mohali, SAS Nagar, Punjab, India.

[5] National Food Institute, Technical University of Denmark, Kgs. Lyngby, Denmark.

[6] Department of Microbiology and Molecular Genetics, and Department of Molecular and Cellular Biology, University of California, Davis, CA 95616, USA.

* Address correspondence to Theetha L Pavankumar, pavan@ucdavis.edu or to Pragya Mittal, pragya.mittal@ed.ac.uk. The authors have declared no conflict of interest.

Abstract

RNase R is a highly processive, 3' - 5' exoribonuclease involved in RNA degradation,

maturation, and processing in bacteria. In *Pseudomonas syringae* Lz4W, RNase R interacts

with the RNase E to form the RNA degradosome complex and is essential for growth at low

temperature. RNase R is also implicated in general stress response in many bacteria. We show

here that *rnr* gene (encoding RNase R) deletion mutant of *P. syringae* is highly sensitive to

various DNA damaging agents and oxidative stress. RNase R is a multidomain protein

comprised of CSD, RNB and S1 domains. We investigated the role of each domain of RNase R

and its catalytic activity in nucleic acid damage and oxidative stress response. Our results

revealed that RNB domain alone without its exoribonuclease activity is sufficient to protect

against DNA damage and oxidative stress. We also show that association of RNase R with

degradosome complex is not required for this function. Finally, based on homology modelling of

RNB domain of RNase R of *P. syringae*, we propose that the presence of a tri-helix wedge

domain may be involved in protection against DNA damage and oxidative stress in *P. syringae*

through a putative helicase activity. Our study for the first time has discovered a hitherto

unknown role of RNase R in protecting *P. syringae* Lz4W against DNA damage and oxidative

stress.

Importance

Bacterial exoribonucleases play a crucial role in RNA maturation, degradation, quality

control and its turnover. In this study, we have uncovered a previously unknown role of 3'- 5'

exoribonuclease RNase R of *P. syringae* Lz4W in DNA damage and oxidative stress response.

Here, we show that neither the catalytic exoribonuclease function of RNase R, nor its

association with the RNA degradosome is essential for this function. Nevertheless, a putative

helicase function of RNase R appears to have an important role in protecting against DNA

damage and oxidative stress. Interestingly, in *P. syringae* Lz4W, the hydrolytic RNase R

exhibits physiological roles similar to phosphorolytic 3'- 5' exoribonuclease PNPase of *E. coli*.

Our data suggests that during the course of evolution, mesophilic *E. coli* and psychrotrophic *P.*

syringae have apparently swapped these exoribonucleases to adapt to their respective

environmental growth conditions.

Key words: *Pseudomonas syringae*, RNase R, DNA damage, Oxidative stress, RNA

degradation, and RNA unwinding.

Introduction

Exoribonucleases play an important role in determining the levels of RNA in the cells (1). Some of the exoribonucleases also associate with degradosome, a multi-protein complex involved in degradation and processing of cellular RNA (2, 3). The RNR superfamily of enzymes that includes proteins like RNase II, RNase R, and the eukaryotic Rrp44/Dis3 are present in all domains of life (4). RNase R, encoded by *rnr* gene, is a 3′ to 5′ highly processive exoribonuclease involved in degradation, maturation and processing of RNA in bacteria (5-8). RNase R plays an important role in RNA quality control, such as degradation of defective tRNA and removal of aberrant fragments of the 16S and 23S rRNAs (7, 9). RNase R is capable of processing dsRNA without the help of a helicase, hence implicated in removal of mRNAs with stable stem loops, including repetitive extragenic palindromic (REPs) elements (6, 10) It is also involved in processing of tmRNA (transfer-messenger RNA) and is required for the process of trans-translation (11). Structurally, RNase R is a multidomain protein consists of RNA binding domains cold-shock domains (CSD) and S1, and the catalytic RNB domain (12). The catalytic RNB domain contains four highly conserved Aspartate residues that help to coordinate Mg^{2+} ions in the catalytic center to facilitate ribonucleolytic activity (12-14).

In *Pseudomonas syringae* Lz4W, RNase R associates with endoribonuclease RNase E, along with DEAD box helicase RhlE to form the degradosome complex (15). The *rnr* gene *P. syringae* Lz4W and *P. putida* is implicated in survival at low temperature (7, 16-18). The *P. syringae* Lz4W cells lacking RNase R accumulate unprocessed 3' ends of 16S rRNAs in polysomes, making the translation machinery inefficient at low temperature, causing cold sensitivity (7). RNase R is also involved in general cellular stress responses. In *E. coli*, RNase R levels increase under cold shock, stationary phase, and heat shock (19, 20).

In a quest to understand the role of RNase R of *P. syringae* Lz4W (RNase R[Ps]) in general cellular stress response, we subjected the *rnr* mutant to various types of stresses and found that *rnr* mutant cells are sensitive to oxidative stress and DNA damaging agents. We show that this role of RNase R is independent of its association with the degradosome complex in *P. syringae* Lz4W. Using a series of domain deletion constructs and mutating catalytically important residues, we discovered that the catalytic RNB domain alone without its exoribonuclease function is sufficient to protect *rnr* mutant of *P. syringae* from DNA damage and oxidative stress. Our data further indicates that a putative helicase function of RNB domain is likely to play a role in protecting *P. syringae* Lz4W cells from DNA damage and oxidative stress.

Results

P. syringae rnr mutant cells are sensitive to oxidative stress and DNA damaging agents

In an attempt to understand the role of RNase R in general stress response in *P. syringae*, the *rnr* mutant was subjected to various conditions that include low and high temperatures (4 and 37 ⁰C), osmotic stress (1M NaCl), envelope stress (10% SDS, 100 mg/ml Lysozyme with 0.5 mM EDTA), antibiotics treatment (100 μg/ml Rifampicin and Spectinomycin), oxidative (0.25 mM Hydrogen peroxide, 15 μM Paraquat) and DNA damage stress (24 J/m^2.s ultraviolet radiation, 0.25 mM Cisplatin, 1 μg/ml Mitomycin C and 2.5 mM Hydroxyurea). As shown in Table 1, we observed that the sensitivity of *rnr* mutant against osmotic, envelope and antibiotic stresses is comparable to that of wild-type (*wt*) cells. We confirmed that the *rnr* mutant cells are cold sensitive, as shown earlier (7). Nevertheless, we found that *rnr* mutant is also sensitive to high temperature stress (Table 1).

Interestingly, we discovered that the *rnr* mutant cells are highly sensitive to the oxidative stress induced by H_2O_2 and Paraquat; and to DNA damage induced by ultraviolet (UV) radiation, Mitomycin C (MMC) and Hydroxyurea (HU) (Table 1). H_2O_2 and Paraquat generate reactive oxygen species (ROS), such as hydroxyl and superoxide radicals that can easily react with biological macromolecules including DNA, RNA, proteins, and lipids (21). We quantified the effect of these agents on the growth and viability of *rnr* mutant in comparison to the *wt* cells. We assessed the colony-forming ability of cells upon treatment with various doses of H_2O_2, Paraquat, UV radiation, MMC and HU. The *rnr* mutant displayed more than 100-fold higher sensitivity compared to *wt* cells to all the agents tested, suggesting a cryptic role of RNase R in protecting *P. syringae* cells from DNA damage and oxidative stress (Figure 1A). Spot assays on ABM plates containing the indicated amounts of H_2O_2, Paraquat, MMC, HU and UV irradiation, further confirmed that *rnr* mutant of *P. syringae* is highly sensitive to all the agents (Figure 1C). In contrast, we observed that the mesophilic *E. coli* cells lacking RNase R (Δrnr) are refractory to oxidative stress and DNA damaging agents (Figure S1) suggesting that the protective role of RNase R against DNA damage is specific to *P. syringae*.

Furthermore, introducing the *rnr* gene encoding RNase R^{Ps} on a low copy plasmid (pGL10::*rnr*^{Ps}) rescued the growth defects of *rnr* mutant caused by DNA damage/oxidative stress confirming that the functional loss of RNase R causes the sensitivity (Figure 1A & 1C). Interestingly, when RNase R of *E. coli* (pGL*rnr*^{Ec}) is overexpressed in *P. syringae* Lz4W, it also rescued the growth defect of *rnr* mutant of *P. syringae* (Figure 1A & 1C). In *E. coli*, another

exoribonuclease PNPase has been shown to protect cells against oxidative stress and DNA

damage (22-25). We examined if PNPase of *P. syringae* has similar functions and could

complement the loss of *rnr* gene against oxidative and DNA damage stress. However, we

observed that *rnr* mutants over-expressing PNPase could not rescue the sensitivity of *rnr*

mutant to oxidative and DNA damage stress (Figure S2).

RNase R is known to upregulate under different stress conditions in *E. coli* (19, 20),

hence we examined the levels of RNase R in *P. syringae* Lz4W cells by Western blotting with

polyclonal anti-RNase R antibodies (15). The levels of RNase R remained unchanged after

treating with stress causing agents (Figure 1B). As a control, the expression of DNA repair

protein RecA was induced quickly upon treatment with UV irradiation as shown earlier (26). This

is in line with a previous report that RNase R of *P. syringae* Lz4W is not induced under cold

stress and in stationary phase (7). This suggests that unlike *E. coli*, the RNase R^{Ps} is not a

stress inducible protein.

110 RNase R protects *P. syringae* cells against oxidative stress and DNA damage

111 independent of RNA degradosome complex

RNase R associates with endonuclease RNase E and RNA helicase RhlE to form the

RNA degradosome complex in *P. syringae* Lz4W (15). The C-terminal domain of RNase E acts

as a scaffold for the formation of degradosome complex with RNase R and RhlE in *P. syringae*

(15). Thus, we constructed a C-terminal deletion mutant of RNase E (*rne*$^{\Delta595-1074}$) to investigate

the role of RNase R associated degradosome complex in oxidative stress and DNA damage.

The generation of C-terminal deleted *rne* mutant (*rne*$^{\Delta595-1074}$) was carried out as described in

materials and methods. The *rne*$^{\Delta595-1074}$ strain was subjected to oxidative stress and DNA

damage causing agents. Unlike the *rnr* mutant, the *rne*$^{\Delta595-1074}$ strain is not sensitive to the DNA

damage and oxidative stress causing agents and showed resistance similar to *wt* (Figure 2).

The results suggest that the requirement of RNase R in protecting *P. syringae* cells against

oxidative stress and DNA damage is an intrinsic property of RNase R and is independent of its

association with the RNA degradosome complex.

124 The RNB domain of RNase R is sufficient in protecting *P. syringae* Lz4W against

125 oxidative stress and DNA damage

The RNase R^{Ps} is a multidomain protein comprising of N-terminal RNA binding cold

shock domain CSD (1 – 225 aa), a central nuclease domain RNB (226 – 665 aa) and the RNA

binding domain S1 at the C-terminus (666 – 885 aa) (Figure 3A). To understand the role of each

domain in protecting *P. syringae* Lz4W cells against oxidative stress and DNA damage, we

constructed a series of domain deletion mutants of RNase R: i) a mutant lacking CSD domain

(ΔCSD), ii) a mutant lacking S1 domain (ΔS1), and iii) a mutant lacking both CSD and S1

domains (RNB) (Figure 3A). In our viability assays, the *rnr* mutants complemented with the

domain deleted mutants of RNase R showed resistance to the oxidative stress and DNA

damaging agents similar to *wt* cells (Figure 3B & C). This suggests that the RNB domain of

RNase R alone, without the RNA binding CSD and S1 domains is sufficient to protect *P.*

syringae Lz4W against oxidative stress and DNA damage.

137   The exoribonuclease activity of RNase R is not required for protecting *P. syringae* Lz4W

138   against oxidative stress and DNA damage

The catalytic site of RNase R^{Ps} has four conserved aspartate residues in the active site

located at positions 276, 282, 284 and 285 (Figure 4A). D276 and D285 have been proposed to

be important for coordinating with Mg^{2+} ion, D282 for holding the RNA in place while D284 is

known for being the residue that cleaves the phosphodiester bond (27). It was shown in *E. coli*

that D280 (equivalent to the D284 residue of *P. syringae*) is the only crucial residue for the

catalytic activity of RNase R without affecting binding of the RNA (14). This residue is also

shown to play a crucial role in the activity of RNase II of *E. coli* and RNase R of *Legionella*

pneumophila (28). Hence, we mutated the equivalent Asp284 in RNase R to alanine,

overexpressed and purified the mutant protein (RNaseR^{D284A}) as described under materials and

methods. SDS-PAGE profile of RNaseR^{D284A} is shown in Fig 4B(i). RNase R^{D284A} show double

protein bands on a SDS-polyacrylamide gel. The lower band is a product of C-terminally

degraded RNase R protein as observed earlier (10). We couldn't separate these two bands due

to marginal difference in their molecular weights. We analyzed the exoribonucleolytic activity of

RNaseR^{D284A} protein on Poly(A) and *malE-malF* substrates and compared with the wild-type.

Wild-type RNase R degraded both poly(A) and *malE-malF* substrates within 10 mins. However,

despite using high protein concentration of RNaseR^{D284A} mutant (0.5 μM vs 15 μM) it failed to

degrade both substrates even after prolonged incubation (Fig 4B (ii & iii)). This shows that the

D284A mutation leads to complete loss of exoribonuclease activity of RNase R^{Ps}.

In our viability assays, all four single amino acid mutants protected Δ*rnr* mutant cells

from oxidative and DNA damage stress, similar to *wt* cells (Figure 5). However, the D282A,

D284A and D285A mutants showed marginal sensitivity to stress induced by hydroxyurea (HU)

suggesting that the functional role of these residues in the activity of RNase R have a plausible

role in protecting particularly against HU induced stress. In general, the data suggest that the

role of RNase R in protecting cells against oxidative stress and DNA damage is independent of

exoribonuclease activity of RNase R.

To further substantiate our finding, we also combined the following mutations on full

length RNase R: D284A + D285A; D276A + D284A; D276A + D284A + D285A. We examined

their role in the function of RNase R in protecting against stress (Supplementary Figure 3). All

the double and triple point mutants also protected Δ*rnr* cells against oxidative stress and DNA

damage similar to *wt* cells. This further confirms that the exoribonuclease activity of RNase R is

not essential for the protection against oxidative stress and DNA damage.

The RNB domain alone with D284A mutation can protect *P. syringae* Lz4W cells against

oxidative stress and DNA damage.

RNB is the main catalytic domain responsible for the exoribonuclease activity of RNase

R. As mentioned above the four conserved Aspartate residues important for the catalytic activity

of RNase R are located in the motif-1 of the RNB domain (Figure 4A(ii)). To further corroborate

our finding that the exoribonuclease function of RNase R^{Ps} is not required for protection against

DNA damage and oxidative stress; we mutated the D284 and D284+D285 residues on catalytic

RNB domain alone to alanine. RNB domain with D284A and D284A+D285A mutations was

expressed in Δ*rnr* strain. The viability assays for the RNB domain with mutations show that the

mutants moderately protect cells against oxidative stress and DNA damage. However, the point

mutations slightly affected the ability of RNB domain to protect cells against Micromycin C,

Paraquat and Hydroxyurea (Figure 5A).

In *E. coli*, RNB domain is also suggested to play an important role in RNA unwinding. It

has been recently suggested that a triple helix wedge region in the RNB domain of *E. coli*

RNase R is responsible for unwinding the duplex-RNA substrates (29). Our homology modelling

of RNase R^{Ps} based on the crystal structure of *E. coli* RNase R, also shows the presence of a

tri-helix wedge region in the RNB domain of RNase R^{Ps} (**Figure 5B**). Although we haven't

examined the role of tri-helix wedge of RNase R^{Ps} in RNA unwinding function, it is possible that

it may exhibit helicase activity similar to the tri-helix wedge region in the RNB domain of *E. coli*

RNase R.

Discussion

Ribonucleases are important modulators involved in various cellular processes by virtue of their RNA processing and degradation activity in maintaining the dynamic transcript levels in bacterial cells. Exoribonucleases like RNase R, PNPase and RNaseJ1 are multifunctional enzymes and are implicated in various biological processes such as stress tolerance, protection from DNA damage and oxidative stress apart from their role in RNA metabolism (19, 20, 22, 24, 30, 31). RNase R is important for the RNA quality control and implicated in various stress conditions in bacteria (5, 20). In this study, we have uncovered a novel role of RNase R in protection of *P. syringae* Lz4W from oxidative stress and DNA damaging agents such as H_2O_2, Paraquat, MMC, Hydroxyurea and UV radiation.

PNPase is a phosphorolytic 3'- 5' exoribonuclease an important component of *E. coli* RNA degradosome complex. However, in *P. syringae*, instead of PNPase, the hydrolytic 3'-5' exoribonuclease RNase R interacts with the RNA degradosome complex (15). Interestingly, *pnp* mutant of *E. coli* and the *rnr* mutant of *P. syringae* exhibit similar phenotypic behaviors. The *pnp* mutant of *E. coli* and the *rnr* mutant of *P. syringae* are both sensitive to cold and both enzymes are shown to be involved in maturation of 3' end of 16S rRNA (32-35). RNase R of *Pseudomonas putida* is also implicated in survival at low temperature (7, 16-18). PNPase is also implicated in protecting *E. coli* cells against oxidative damage (31) and in DNA repair process by virtue of its 3'→5' DNase activity in *B. subtilis* and *E. coli* (22-25).

We show here that the RNase R is involved in oxidative damage and DNA repair process in *P. syringae* Lz4W. Unlike *P. syringae* RNase R, the *rnr* mutant of *E. coli* has no discernable phenotype and is refractory to these stress conditions. This implies that the protective role of RNase R is either specific to *P. syringae* Lz4W or possibly specific to *Pseudomonas* species. It is intriguing that although RNase R of *E. coli* is not involved in DNA damage and oxidative stress response in *E. coli*, yet it protects *P. syringae* Lz4W from DNA damage and oxidative stress. It is possible that the PNPase[Ec] and RNase R[Ps] have evolutionarily swapped their functions with respect to their roles in the cellular milieu of these two bacteria. Although, we don't know why RNase R has been chosen over PNPase (unlike *E. coli)* to protect *P. syringae* Lz4W from oxidative stress and DNA damage, we speculate that the highly processive, hydrolytic RNase R is selected over the less processive, phosphorolytic PNPase to conserve the energy at low temperature.

RNA degradosome is a multi-enzyme complex that significantly contributes to the

steady-state profiles of RNA transcripts in bacteria (36). Our laboratory has previously reported

a novel degradosome complex comprising of the RNase E, the exoribonuclease RNase R and a

DEAD-box RNA helicase RhlE in *P. syringae* Lz4W (15). By deleting of C-terminal domain of

RNase E that is responsible for the assembly of degradosome complex, we showed that the

association of RNase R with the degradosome complex is not needed to protect against

oxidative and DNA damage stress, thus suggesting that the DNA damage resistance phenotype

is an intrinsic function of RNase R.

We further investigated the possible role of each domain of multi-domain RNase R in

protecting *P. syringae* cells from oxidative and DNA damage stress. Interestingly, we found the

catalytic RNB domain of RNase R^{Ps} alone is sufficient for protecting *P. syringae* cells from

oxidative and DNA damage stress, while the CSD and S1 domains are dispensable. In *E. coli*,

RNB domain alone degrades structured RNA (12). Therefore, we envisaged that the

degradation of structured RNAs either by full-length RNase R^{Ps} or by the catalytic RNB domain

itself could play a protective role against oxidative and DNA damage stress in *P. syringae*

Lz4W. We replaced the highly conserved catalytically important aspartate residues of RNase

R^{Ps} with alanine. These mutants showed no apparent sensitivity to oxidative stress and DNA

damaging agents, indicating the RNA degradation function of RNase R is not required. Further,

the double- and triple-point mutations of the aspartate residues in combinations also resulted in

similar results, confirming that the ribonuclease function of RNase R^{Ps} is not essential. Thus, the

ability of RNase R to degrade highly stabilized RNA secondary structures does not have a role

in protecting *P. syrin*gae cells against DNA damage and oxidative stress. Intriguingly, our data

shows that the catalytically inactive RNase R^{Ps} and RNB domain itself can protect cells from

oxidative and DNA damage stress, implying the importance of RNB domain alone with no

exoribonuclease activity in this scenario.

If the association of RNase R with degradosome complex and its exoribonuclease

activity is not essential, then how does RNase R protect *P. syringae* cells from DNA damage

and oxidative stress? How does a catalytically inactive RNB domain itself without CSD and S1

domains could rescue *P. syringae* cells from oxidative and DNA damage stress? However, our

data suggests that the RNB domain without catalytic function has a cryptic functional role which

helps in protecting Lz4W cells from these stress conditions.

RNase R is known to process structured dsRNA without the help of a helicase and is

implicated in removal of mRNAs with stable stem loops (6). It was earlier shown that cold

sensitivity of *csdA* mutant of *E. coli* can be complemented by overexpressing RNase R, possibly

through its helicase function (37). Importantly, the RNB domain of RNase R is also shown to

possess helicase activity in *E. coli* (32, 38, 39). A recent study showed the presence of a winged

tri-helix domain in RNB domain of *E. coli* that is responsible for the helicase function (29). This

prompted us to investigate the presence of a similar tri- helix wedged domain in RNase R of *P.*

syringae. Our homology modelling of *P. syringae* RNase R based on the RNase R of *E. coli*,

shows a similar tri-helix wedge in RNB domain. Hence, we hypothesize that RNase R could

participate in protection of *P. syringae* Lz4W against oxidative and DNA damage stress via its

helicase activity. This is further supported by a recent observation in *P. syringae* Lz4W that the

DEAD box helicases CsdA and SrmB are involved in protection against DNA damage and

oxidative stress (40).

This study uncovered a hitherto unknown role of RNase R in the physiology of cold

adapted bacterium *P. syringae* Lz4W. Here, we report a novel role of the RNase R in protecting

P. syringae Lz4W against DNA damage and oxidative stress. Our study indicate that the

protective role of RNase R is apparently specific to *P. syringae* Lz4W. Our structural homology

modelling provides an insight into the presence of a tri-helix wedge region in the RNB domain of

RNase R. which might play a role in protecting against oxidative and DNA damage stress by

virtue of its putative helicase activity. Although our data biochemically doesn't not establish the

helicase activity of RNB domain of RNase R, the genetic evidence provided here is

comprehendible. In future it would be of relevance to elucidate the mechanism by which RNase

R protects *P. syringae* Lz4W cells from DNA damage and oxidative stress.

Materials and Methods

Bacterial strains, plasmids and growth conditions

Bacterial strains and plasmids used in this study are described in Table 2. *Pseudomonas*

syrinage Lz4W cells were grown in Antarctic Bacterial medium (ABM: 5g liter^{-1} peptone and 2g

liter^{-1} yeast extract) at both 22ºC and 4ºC. *E. coli* strains were routinely grown in Luria-Bertani

medium at 37ºC. The growth media, when required, were supplemented with Antibiotics as

follows: 100 µg ml^{-1} Ampicillin, 50 µg ml^{-1} Kanamycin, 20µg ml^{-1} tetracycline, and 400 µg ml^{-1}

rifampicin. For complementation studies, plasmids were mobilized into *P. syringae* strain by

conjugation using *E. coli* S17-1 helper strain. For growth analysis, cultures were grown in ABM

overnight and were inoculated in a fresh media in 1:100 dilutions and OD_{600} (O.D at 600 nm) was measured at various time intervals as indicated.

Spot assays and quantitation of sensitivity to DNA damaging agents

Cells were grown overnight at 22 ºC in AB media containing appropriate antibiotics and inoculated into fresh AB media with 1:100 dilutions. To measure the sensitivity to NaCl (1M), SDS (10% w/v), Lysozyme (100 mg/ml) + EDTA (500 mM), Ampicillin (100 µg), Rifampicin (100 µg), Spectinomycin (100 µg), Paraquat (20 µM), Mitomycin C (1000 ng), H_2O_2 (0.2 mM), Hydroxyurea (2.5 mM), Cisplatin (0.2 mM); up to 5-fold serial dilutions of exponentially grown cells at 22 ºC were made and spotted on ABM agar plates containing stress inducing agents as indicated. Plates were incubated at 22 ºC for 48 h. For UV sensitivity, up to 5-fold serially diluted exponentially grown cultures were spotted on ABM plates and exposed to UV radiations at indicated doses at the rate of 3 J/m^2/sec and plates were incubated in dark at 22 ºC for 48 h.

For quantitative analysis, exponentially growing cells were incubated for 30 mins with oxidative stress/DNA damaging agents at indicated concentrations, washed, then serially diluted, and plated on ABM-Agar plates or LB agar plates (for *E. coli* strains). Plates were incubated at 22 ºC for 48 h and the colony forming units (cfu) were scored. The percentage of cell survival was calculated by considering the cfu from untreated cells as 100%.

Reagents and general DNA recombinant methods

Routine molecular biology techniques like Genomic DNA isolation, restriction enzyme digestion, PCR, ligation, transformation, conjugation, Southern hybridization, Western analysis and RNA isolation were performed as described earlier (41). All restriction enzymes, T4 DNA ligase, T4 Polynucleotide kinase were purchased from New England Biolabs. Oligos used in the study were from Bioserve India (Hyderabad, India). PCR was carried out using proof reading Pfx DNA polymerase from Invitrogen (San Deigo, CA, USA). DNA sequencing was performed using ABI Prism dye terminator cycle sequencing method (Perkin-Elmer) and analyzed on an automated DNA sequencer (ABI model 3700; Applied Biosystems). Antibodies were purchased from Santa Cruz Biotechnology.

For Southern hybridization, genomic DNA was isolated from WT and *ΔC-rne* mutant cells, digested with PstI enzyme. The products of restriction digest were separated on a 1% w/v Agarose gel and transferred onto Hybond N+ membrane (Amersham Biosciences). For the probe, PCR amplicon of full-length *rne* gene was labeled with [α-^{32}P] dATP using random primer

316 labeling kit (Jonaki, BARC, India). Radioactive signals were detected and quantified by

317 phosphorimager (Fuji FLA 3000).

Construction of mutants of RNase R and RNB by site directed mutagenesis and genetic

complementation studies

The rnr^{Ec}, rnr^{Ps} genes and the domain deletion mutants of rnr^{Ps} were amplified from the

genomic DNA of Lz4W using the primers listed in Table 3. All constructs contain 6x-Histidine tag

at the N-terminal end of the protein. For D276A, D282A, D284A and D285A point-mutations in

RNase R^{Ps}, site-directed mutagenesis of pGL*rnr* (7) was carried out using self-complementary

primers harboring indicated mutations (listed in Table 3). Subsequently, the double D284A +

D285A, D276A + D284 and triple D276A+D284A+D285A mutations were generated by using

respective complementary primers using pGLrnr plasmid with preexisting mutations. Similarly, the

D284A and D284A+D285A mutations were introduced into pGLRNB vector. Mutations and

integrity of full-length genes were confirmed by DNA sequencing.

Purification of RNase R D284A **mutant and biochemical assays**

The wild-type RNase R and D284A mutant derivatives were purified by expressing these

proteins in *rnr* null mutant of *P. syringae* Lz4W as escribed earlier (10). RNase R activity was

assayed *in vitro* by its ability to degrade ssRNA substrate Poly(A) and structured RNA substrate

malE-malF as described earlier (10). Briefly, Poly(A) was labelled at 5'-end using [γ^{32}P] and the

malE-malF substrate was internally labelled using (α^{32}P). Assays were carried out in 20-µl

reaction mixtures containing 25 mM Tris-HCl (pH 7.0), 100 mM KCl, 0.25 mM MgCl$_2$, 5 mM

DTT. 10 pmol of labelled substrate was treated with 0.5 µM RNase R^{WT} and 15 µM RNase

R^{D284A} at the indicated time-points at room temperature, and stopped by adding 10mM EDTA,

0.2% SDS and 1 mg/ml of proteinase K. RNA degradation products were resolved on 8%

denaturing 8 M urea PAGE and degradation products were analysed by Phosphoimager (Fuji).

Western analysis

For examining the induction of RNase R, the exponentially grown wild-type *P. syringae*

cells were incubated with DNA damaging agents. 1 ml aliquot was collected before treating cells

with DNA damaging agents. 5 ml of cells were treated separately with Paraquat (15 µM), MMC

(1000 ng), UV (18 J/m^2), and incubated at 22ºC. 1 ml of aliquot was collected at 30, 60 and 120

min intervals, Cells were further lysed by sonication in presence of Protease inhibitor cocktail

(Roche, USA). Protein concentration from cell extracts was measured using Bio-rad Protein

Assay kit, also by the absorbance at 280 nm (A_{280}) using the Nanodrop spectrophotometer.

Equal amount of protein was loaded and resolved on a 10 % w/v SDS-PAGE. The resolved

proteins were later electro-blotted onto Hybond C-Extra membrane (Amersham Biosciences)

and immunodetection was carried with polyclonal rabbit Anti-RNase R antibodies as described

earlier (15). For detecting the levels of RecA expression, cells were treated with UV (18 J/m^2)

and incubated at 22 °C in dark. Culture aliquots were collected after 30 minutes of treatment

and immuno-detected by polyclonal anti-RecA rabbit antibodies as described earlier (26).

Generation of C-terminal deletion of *rne* **(RNase E gene)**

For generating C-terminal deletion of *rne* gene, a 784 bp fragment from N-terminal of *rne*

gene was first amplified using primer set Rne_Nterm_FP_BamHI and Rne_Nterm_RP_HindIII

having the restriction enzyme sites for BamHI and HindIII respectively **(Table 2).** A second

fragment from C-terminal of *rne* gene was PCR amplified using the primer set

Rne_Cterm_FP_HindIII and Rne_Nterm_RP_SalI having the sites for restriction enzymes

HindIII and SalI (Table 2). Both fragments were initially digested with HindIII and ligated. The

ligated product was further digested with BamHI and SalI and cloned into pBlueScript-KS

vector. The resultant plasmid pBS(N+C)*rne* was linearized by HindIII, and a tetracycline

resistance gene was cloned to obtain the pBS(N+C)*rne*-tet plasmid. The tetracycline resistance

gene (*tet*) cassette (~2.4 KB) was obtained from the pMOS^{tet} (42) as a HindIII fragment. Plasmid

construct was confirmed by sequencing and PCR analysis. Plasmid pBS(N+C)*rne*-tet plasmid

was alkaline denatured as described earlier (42). Alkali denatured plasmid DNA was

electroporated into the wild-type *P. syringae* cells. Recombinants were selected on ABM plates

containing tetracycline and the knock-out was confirmed by Southern and Western analysis.

Modeling of RNase R^{Ps} protein

The amino acid sequence of Lz4W RNase R was taken from NCBI (GenBank:

AUB77091.1). A model of RNase R^{Ps} was constructed using the SWISS-MODEL (43). The

closest template identified was PDB: 5xgu.1, which refers to the RNase R structure from

Escherichia coli. The RMSD between these two structures was then calculated using the

Matchmaker tool of Chimera (44). The RMSD value was calculated between 613 pruned atom

pairs is 0.248 angstroms; (across all 625 pairs: 0.930). The two structures were superimposed

as shown in the figure 5B.

Statistical analysis

Data was calculated and analyzed by using Prism 5 (GraphPad Software Inc, USA).

Acknowledgements

380   We thank Dr. Manjula Reddy for gifting us WT and *rnr* mutant cells of *E. coli*. This study
381   was funded by Council for Scientific and Industrial Research, New Delhi, India.

References

1.   Zuo Y, Deutscher MP. 2001. Exoribonuclease superfamilies: structural analysis and phylogenetic distribution. *Nucleic Acids Res* 29:1017-26.
2.   Carpousis AJ, Van Houwe G, Ehretsmann C, Krisch HM. 1994. Copurification of *E. coli* RNAase E and PNPase: evidence for a specific association between two enzymes important in RNA processing and degradation. *Cell* 76:889-900.
3.   Miczak A, Kaberdin VR, Wei CL, Lin-Chao S. 1996. Proteins associated with RNase E in a multicomponent ribonucleolytic complex. *Proc Natl Acad Sci U S A* 93:3865-9.
4.   Matos RG, Barria C, Moreira RN, Barahona S, Domingues S, Arraiano CM. 2014. The importance of proteins of the RNase II/RNB-family in pathogenic bacteria. *Front Cell Infect Microbiol* 4:68.
5.   Cheng ZF, Deutscher MP. 2003. Quality control of ribosomal RNA mediated by polynucleotide phosphorylase and RNase R. *Proc Natl Acad Sci U S A* 100:6388-93.
6.   Cheng ZF, Deutscher MP. 2005. An important role for RNase R in mRNA decay. *Mol Cell* 17:313-8.
7.   Purusharth RI, Madhuri B, Ray MK. 2007. Exoribonuclease R in *Pseudomonas syringae* is essential for growth at low temperature and plays a novel role in the 3' end processing of 16 and 5 S ribosomal RNA. *J Biol Chem* 282:16267-77.
8.   Zuo Y, Vincent HA, Zhang J, Wang Y, Deutscher MP, Malhotra A. 2006. Structural basis for processivity and single-strand specificity of RNase II. *Mol Cell* 24:149-56.
9.   Arraiano CM, Andrade JM, Domingues S, Guinote IB, Malecki M, Matos RG, Moreira RN, Pobre V, Reis FP, Saramago M, Silva IJ, Viegas SC. 2010. The critical role of RNA processing and degradation in the control of gene expression. *FEMS Microbiol Rev* 34:883-923.
10.   Sulthana S, Rajyaguru PI, Mittal P, Ray MK. 2011. rnr gene from the antarctic bacterium *Pseudomonas syringae* Lz4W, encoding a psychrophilic RNase R. *Appl Environ Microbiol* 77:7896-904.
11.   Richards J, Mehta P, Karzai AW. 2006. RNase R degrades non-stop mRNAs selectively in an SmpB-tmRNA-dependent manner. *Mol Microbiol* 62:1700-12.
12.   Vincent HA, Deutscher MP. 2009. The roles of individual domains of RNase R in substrate binding and exoribonuclease activity. The nuclease domain is sufficient for digestion of structured RNA. *J Biol Chem* 284:486-94.
13.   Barbas A, Matos RG, Amblar M, Lopez-Vinas E, Gomez-Puertas P, Arraiano CM. 2008. New insights into the mechanism of RNA degradation by ribonuclease II: identification of the residue responsible for setting the RNase II end product. *J Biol Chem* 283:13070-6.
14.   Matos RG, Barbas A, Arraiano CM. 2009. RNase R mutants elucidate the catalysis of structured RNA: RNA-binding domains select the RNAs targeted for degradation. *Biochem J* 423:291-301.
15.   Purusharth RI, Klein F, Sulthana S, Jager S, Jagannadham MV, Evguenieva-Hackenberg E, Ray MK, Klug G. 2005. Exoribonuclease R interacts with

endoribonuclease E and an RNA helicase in the psychrotrophic bacterium *Pseudomonas syringae* Lz4W. *J Biol Chem* 280:14572-8.

16. Fonseca P, Moreno R, Rojo F. 2008. Genomic analysis of the role of RNase R in the turnover of *Pseudomonas putida* mRNAs. *J Bacteriol* 190:6258-63.

17. Pavankumar TL, Mittal P, Hallsworth JE. 2020. Molecular insights into the ecology of a psychrotolerant *Pseudomonas syringae*. *Environ Microbiol* doi:10.1111/1462-2920.15304.

18. Reva ON, Weinel C, Weinel M, Bohm K, Stjepandic D, Hoheisel JD, Tummler B. 2006. Functional genomics of stress response in *Pseudomonas putida* KT2440. *J Bacteriol* 188:4079-92.

19. Cairrao F, Cruz A, Mori H, Arraiano CM. 2003. Cold shock induction of RNase R and its role in the maturation of the quality control mediator SsrA/tmRNA. *Mol Microbiol* 50:1349-60.

20. Chen C, Deutscher MP. 2005. Elevation of RNase R in response to multiple stress conditions. *J Biol Chem* 280:34393-6.

21. Schieber M, Chandel NS. 2014. ROS function in redox signaling and oxidative stress. *Curr Biol* 24:R453-62.

22. Cardenas PP, Carrasco B, Sanchez H, Deikus G, Bechhofer DH, Alonso JC. 2009. *Bacillus subtilis* polynucleotide phosphorylase 3'-to-5' DNase activity is involved in DNA repair. *Nucleic Acids Res* 37:4157-69.

23. Cardenas PP, Carzaniga T, Zangrossi S, Briani F, Garcia-Tirado E, Deho G, Alonso JC. 2011. Polynucleotide phosphorylase exonuclease and polymerase activities on single-stranded DNA ends are modulated by RecN, SsbA and RecA proteins. *Nucleic Acids Res* 39:9250-61.

24. Carzaniga T, Sbarufatti G, Briani F, Deho G. 2017. Polynucleotide phosphorylase is implicated in homologous recombination and DNA repair in *Escherichia coli*. *BMC Microbiol* 17:81.

25. Rath D, Mangoli SH, Pagedar AR, Jawali N. 2012. Involvement of pnp in survival of UV radiation in *Escherichia coli* K-12. *Microbiology* (Reading) 158:1196-1205.

26. Sinha AK, Pavankumar TL, Kamisetty S, Mittal P, Ray MK. 2013. Replication arrest is a major threat to growth at low temperature in Antarctic *Pseudomonas syringae* Lz4W. *Mol Microbiol* 89:792-810.

27. Frazao C, McVey CE, Amblar M, Barbas A, Vonrhein C, Arraiano CM, Carrondo MA. 2006. Unravelling the dynamics of RNA degradation by ribonuclease II and its RNA-bound complex. *Nature* 443:110-4.

28. Charpentier X, Faucher SP, Kalachikov S, Shuman HA. 2008. Loss of RNase R induces competence development in *Legionella pneumophila*. *J Bacteriol* 190:8126-36.

29. Chu LY, Hsieh TJ, Golzarroshan B, Chen YP, Agrawal S, Yuan HS. 2017. Structural insights into RNA unwinding and degradation by RNase R. *Nucleic Acids Res* 45:12015-12024.

30. Han R, Jiang J, Fang J, Contreras LM. 2022. PNPase and RhlB Interact and Reduce the Cellular Availability of Oxidized RNA in *Deinococcus radiodurans*. *Microbiol Spectr* 10:e0214022.

31. Wu J, Jiang Z, Liu M, Gong X, Wu S, Burns CM, Li Z. 2009. Polynucleotide phosphorylase protects *Escherichia coli* against oxidative stress. *Biochemistry* 48:2012-20.

32. Awano N, Inouye M, Phadtare S. 2008. RNase activity of polynucleotide phosphorylase is critical at low temperature in *Escherichia coli* and is complemented by RNase II. *J Bacteriol* 190:5924-33.

471   33.   Chen C, Deutscher MP. 2010. RNase R is a highly unstable protein regulated by growth
472   phase and stress. *RNA* 16:667-72.
473   34.   Phadtare S. 2012. *Escherichia coli* cold-shock gene profiles in response to over-
474   expression/deletion of CsdA, RNase R and PNPase and relevance to low-temperature
475   RNA metabolism. *Genes Cells* 17:850-74.
476   35.   Sulthana S, Deutscher MP. 2013. Multiple exoribonucleases catalyze maturation of the
477   3' terminus of 16S ribosomal RNA (rRNA). *J Biol Chem* 288:12574-9.
478   36.   Carpousis AJ. 2007. The RNA degradosome of Escherichia coli: an mRNA-degrading
479   machine assembled on RNase E. *Annu Rev Microbiol* 61:71-87.
480   37.   Awano N, Xu C, Ke H, Inoue K, Inouye M, Phadtare S. 2007. Complementation analysis
481   of the cold-sensitive phenotype of the *Escherichia coli* csdA deletion strain. *J Bacteriol*
482   189:5808-15.
483   38.   Awano N, Rajagopal V, Arbing M, Patel S, Hunt J, Inouye M, Phadtare S. 2010.
484   *Escherichia coli* RNase R has dual activities, helicase and RNase. *J Bacteriol* 192:1344-
485   52.
486   39.   Hossain ST, Deutscher MP. 2016. Helicase Activity Plays a Crucial Role for RNase R
487   Function *in Vivo* and for RNA Metabolism. *J Biol Chem* 291:9438-43.
488   40.   Hussain A, Ray MK. 2022. DEAD box RNA helicases protect Antarctic *Pseudomonas*
489   *syringae* Lz4W against oxidative stress. *Infect Genet Evol* 106:105382.
490   41.   Sambrook J, E. Fritsch, and T. Maniatis. . 1989. *Molecular cloning: a laboratory manual.*,
491   vol I - III. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
492   42.   Pavankumar TL, Sinha AK, Ray MK. 2010. All three subunits of RecBCD enzyme are
493   essential for DNA repair and low-temperature growth in the Antarctic *Pseudomonas*
494   *syringae* Lz4W. *PLoS One* 5:e9412.
495   43.   Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de
496   Beer TAP, Rempfer C, Bordoli L, Lepore R, Schwede T. 2018. SWISS-MODEL:
497   homology modelling of protein structures and complexes. *Nucleic Acids Res* 46:W296-
498   W303.
499   44.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE.
500   2004. UCSF Chimera--a visualization system for exploratory research and analysis. *J*
501   *Comput Chem* 25:1605-12.
502   45.   Shivaji S, Rao NS, Saisree L, Sheth V, Reddy GS, Bhargava PM. 1989. Isolation and
503   identification of *Pseudomonas* spp. from Schirmacher Oasis, Antarctica. *Appl Environ*
504   *Microbiol* 55:767-70.
505   46.   Simon RP, U.; Pühler, A. . 1983. A Broad Host Range Mobilization System for *In Vivo*
506   Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. *Nature*
507   *Biotechnology* 1:784-791.
508   47.   Bidle KA, Bartlett DH. 1999. RecD function is required for high-pressure growth of a
509   deep-sea bacterium. *J Bacteriol* 181:2330-7.

 Figure legends

Figure 1 *S*ensitivity of *P. syringae* of Δ*rnr* and complemented strains to DNA damage and

oxidative stress causing agents. A, Percentage cell survival of WT, Δ*rnr*-pGL10, Δ*rnr*-

pGL*rnr*^{Ps} and Δ*rnr*- pGL*rnr*^{Ec} upon treating with DNA damage and oxidative stress causing

agents at indicated concentrations. Data are from the three independent experiments with mean

± SE. **B,** Expression levels of RNase R of *P. syringae* upon treating with DNA damage and

oxidative stress causing agents, determined by Western analysis using polyclonal RNase R

specific antibodies. The bottom panel shown the expression level of RecA protein upon UV

irradiation as a positive control. **C,** Qualitative assessment of cell viability by spot assays. WT,

Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*^{Ps} and Δ*rnr*- pGL*rnr*^{Ec} treated with DNA damage and oxidative stress

causing agents at indicated concentrations.

Figure 2 *S*ensitivity of C-terminal domain deletion mutant of *rne* gene (*rne* $^{Δ595 - 1074}$)

encoding RNase E compared to Δ*rnr* P. syringae cells. Qualitative assessment of cell

viability by spot assays. WT, Δ*rnr* and Δ*rne* $^{Δ595 − 1074}$ were exposed to DNA damage and

oxidative stress causing agents at indicated concentrations.

Figure 3 *S*ensitivity of Δ*rnr P. syringae* strain complemented with domain/s deleted

mutants of RNase R. A, Schematic representation of structural domains of *P. syringae* RNase

R. The domain/s deleted RNase R constructs; cold shock domain (CSD, 1 – 225 aa), catalytic

RNB domain (226 – 665 aa), and S1 domain (666 – 885 aa) are shown. **B**, Percentage of cell

survival of WT, Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*-*ΔCSD*, Δ*rnr*- pGL*rnr*-*ΔS1* and Δ*rnr*- pGLRNB*(ΔCSD*

+ΔS1) upon treating with DNA damage and oxidative stress causing agents at indicated

concentrations. Data are from the three independent experiments with mean ± SE. **C**,

Qualitative assessment of cell viability by spot assays. WT, Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*-*ΔCSD*,

Δ*rnr*- pGL*rnr*-*ΔS1* and Δ*rnr*- pGLRNB*(ΔCSD +ΔS1)* were treated with DNA damage and

oxidative stress causing agents at indicated concentrations.

Figure 4 *S*ensitivity of Δ*rnr P. syringae* strain complemented with catalytic site mutants

of full-length RNase R to DNA damage and oxidative stress. A, (i) The multiple sequence

alignment of the catalytic region of *P. syringae* RNase R with RNase R of closely related

bacteria including *E. coli*. Boxed and highlighted regions indicate the highly conserved

catalytically important aspartate residues in motif-I of catalytic RNB domain of RNase R. **(ii)**

Schematic representation of structural domains of *P. syringae* RNase Rand the motif-I of RNB

with catalytically important residues are shown. **B.** Purification of RNase R^{D284A} and its

exoribonuclease activity on single-stranded and structured RNA substrates. (i), SDS-PAGE

analysis of purified RNase R^{D284A}, '*' indicate the degraded products of purified RNase R^{D284A}

protein. (ii), shows the time-dependent exoribonuclease activity of wild-type RNase R^{WT} and

RNase R^{D284A} on single stranded Poly(A) and (iii), *malE-malF* structured RNA substrate at

indicated time-points. The end products of size 4-5 mer from degradation of Poly(A), and NMPs

(nucleotide monophosphate) from *malE-malF* substrates are indicated. **C**, Qualitative

assessment of cell viability by spot assays. The WT, Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*-D276A, *rnr*-

pGL*rnr*-D282A, *rnr*- pGL*rnr*-D284A and *rnr*- pGL*rnr*-D285A were exposed to indicated

concentrations of DNA damage and oxidative stress causing agents.

Figure 5 *Sensitivity of Δrnr P. syringae* **strain complemented with catalytic site mutants**

of RNB domain of RNase R. A, Qualitative assessment of cell viability by spot assays. WT,

Δ*rnr*-pGL10, Δ*rnr*- pGLRNB, Δ*rnr*- pGLRNB-D284A and Δ*rnr*- pGLRNB-D284+D285A were

treated with DNA damage and oxidative stress causing agents at indicated concentrations. **B,**

Structural model of *P. syringae* RNase R constructed by superimposing on the structure of *E.*

coli RNase R (PDB:5xgu.1) using SWISS-MODEL. The tri-helix wedge region of RNB domain of

P. syringae RNase R is shown in black color.

Figure S1 *Sensitivity of Δrnr E. coli* **strain to DNA damage and oxidative stress causing**

agents. Qualitative assessment of cell viability by spot assays of WT, Δ*rnr* and Δ*recB* mutant of

E. coli. RecB mutant was used as a positive control against DNA damaging agents. Cells were

exposed to DNA damage and oxidative stress causing agents at indicated concentrations.

Figure S2 *Sensitivity of Δrnr P. syringae* **strain complemented with PNPase of** *P.*

syringae. Qualitative assessment of cell viability by spot assays. WT, Δ*rnr*-pGL10 and Δ*rnr*-

pGL*pnp* were exposed to DNA damage and oxidative stress causing agents at indicated

concentrations.

Figure S3 *Sensitivity of Δrnr P. syringae* **strain complemented with double- and triple-**

mutants of catalytic site residues of RNase R. Qualitative assessment of cell viability by spot

assays. WT, Δ*rnr*-pGL10, *rnr*- pGL*rnr*-(D276A+D284A), *rnr*- pGL*rnr*-(D276A+D285A) and *rnr*-

pGL*rnr*-(D276A+D284A+D285A) treated with DNA damage and oxidative stress causing agents

at indicated concentrations.

Figure 1 *Sensitivity of P. syringae* of Δ*rnr* and complemented strains to DNA damage and oxidative stress causing agents. A, Percentage cell survival of WT, Δ*rnr*-pGL10, Δ*rnr*-pGL*rnr*^{Ps} and Δ*rnr*- pGL*rnr*^{Ec} upon treating with DNA damage and oxidative stress causing agents at indicated concentrations. Data are from the three independent experiments with mean ± SE. **B,** Expression levels of RNase R of *P. syringae* upon treating with DNA damage and oxidative stress causing agents, determined by Western analysis using polyclonal RNase R specific antibodies. The bottom panel shown the expression level of RecA protein upon UV irradiation as a positive control. **C,** Qualitative assessment of cell viability by spot assays. WT, Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*^{Ps} and Δ*rnr*- pGL*rnr*^{Ec} treated with DNA damage and oxidative stress causing agents at indicated concentrations.

Figure 2 *S*ensitivity of C-terminal domain deletion mutant of *rne* gene (*rne* *$^{\Delta 595\text{-}1074}$*)
encoding RNase E compared to Δ*rnr P. syringae* cells. Qualitative assessment of cell
viability by spot assays. WT, Δ*rnr* and Δ*rne* $^{\Delta 595 - 1074}$ were exposed to DNA damage and
oxidative stress causing agents at indicated concentrations.

Figure 3 *Sensitivity of Δrnr P. syringae* **strain complemented with domain/s deleted mutants of RNase R. A**, Schematic representation of structural domains of *P. syringae* RNase R. The domain/s deleted RNase R constructs; cold shock domain (CSD, 1 – 225 aa), catalytic RNB domain (226 – 665 aa), and S1 domain (666 – 885 aa) are shown. **B**, Percentage of cell survival of WT, Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*-ΔCSD, Δ*rnr*- pGL*rnr*-ΔS1 and Δ*rnr*-pGLRNB*(ΔCSD +ΔS1)* upon treating with DNA damage and oxidative stress causing agents at indicated concentrations. Data are from the three independent experiments with mean ± SE. **C**, Qualitative assessment of cell viability by spot assays. WT, Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*-ΔCSD, Δ*rnr*-pGL*rnr*-ΔS1 and Δ*rnr*-pGLRNB*(ΔCSD +ΔS1)* were treated with DNA damage and oxidative stress causing agents at indicated concentrations.

FIGURE 3

Figure 4 *Sensitivity of Δrnr P. syringae* **strain complemented with catalytic site mutants of full-length RNase R to DNA damage and oxidative stress.** **A, (i)** The multiple sequence alignment of the catalytic region of *P. syringae* RNase R with RNase R of closely related bacteria including *E. coli*. Boxed and highlighted regions indicate the highly conserved catalytically important aspartate residues in motif-I of catalytic RNB domain of RNase R. **(ii)** Schematic representation of structural domains of *P. syringae* RNase R and the motif-I of RNB with catalytically important residues are shown. **B.** Purification of RNase R^{D284A} and its exoribonuclease activity on single-stranded and structured RNA substrates. (i), SDS-PAGE analysis of purified RNase R^{D284A}, '*' indicate the degraded products of purified RNase R^{D284A} protein. (ii), shows the time-dependent exoribonuclease activity of wild-type RNase R^{WT} and RNase R^{D284A} on single stranded Poly(A) and (iii), *malE-malF* structured RNA substrate at indicated time-points. The end products of size 4-5 mer from degradation of Poly(A), and NMPs (nucleotide monophosphate) from *malE-malF* substrates are indicated. **C**, Qualitative assessment of cell viability by spot assays. The WT, Δ*rnr*-pGL10, Δ*rnr*- pGL*rnr*-D276A, *rnr*- pGL*rnr*-D282A, *rnr*-pGL*rnr*-D284A and *rnr*- pGL*rnr*-D285A were exposed to indicated concentrations of DNA damage and oxidative stress causing agents.

FIGURE 4

A

FIGURE 5

Figure 5 *Sensitivity of Δrnr P. syringae* **strain complemented with catalytic site mutants of RNB domain of RNase R. A,** Qualitative assessment of cell viability by spot assays. WT, Δ*rnr*-pGL10, Δ*rnr*- pGLRNB, Δ*rnr*- pGLRNB-D284A and Δ*rnr*- pGLRNB-D284+D285A were treated with DNA damage and oxidative stress causing agents at indicated concentrations. **B,** Structural model of *P. syringae* RNase R constructed by superimposing on the structure of *E. coli* RNase R (PDB:5xgu.1) using SWISS-MODEL. The tri-helix wedge region of RNB domain of *P. syringae* RNase R is shown in black color.