Catalytic activity of the Bin3/MEPCE methyltransferase domain is dispensable for 7SK snRNP

function in *Drosophila melanogaster*

Ryan J Palumbo, Steven D Hanes

Department of Biochemistry & Molecular Biology, SUNY Upstate Medical University

750 East Adams Street, 4283 Weiskotten Hall, Syracuse, New York, 13210

PalumboR@upstate.edu, HanesS@upstate.edu

Unexpected *in vivo* results with Bin3

Keywords: Bin3, Bmc1, Bcdin3, MEPCE, Mepcea, 7SK, P-TEFb, RNA Polymerase II, Promoter-Proximal

Pausing, Transcription Elongation

ABSTRACT

Methylphosphate Capping Enzyme (MEPCE) monomethylates the gamma phosphate at the 5'

end of the 7SK noncoding RNA, a modification thought to protect 7SK from degradation. 7SK serves as

a scaffold for assembly of a snRNP complex that inhibits transcription by sequestering the positive

elongation factor P-TEFb. While much is known about the biochemical activity of MEPCE *in vitro*, little

is known about its functions *in vivo*, or what roles— if any—there are for regions outside the conserved

methyltransferase domain. Here, we investigated the role of Bin3, the *Drosophila* ortholog of MEPCE,

and its conserved functional domains in *Drosophila* development. We found that *bin3* mutant females

had strongly reduced rates of egg-laying, which was rescued by genetic reduction of P-TEFb activity,

suggesting that Bin3 promotes fecundity by repressing P-TEFb. *bin3* mutants also exhibited

neuromuscular defects, analogous to a patient with *MEPCE* haploinsufficiency. These defects were also

rescued by genetic reduction of P-TEFb activity, suggesting that Bin3 and MEPCE have conserved roles

in promoting neuromuscular function by repressing P-TEFb. Unexpectedly, we found that a Bin3

catalytic mutant (Bin3^{Y795A}) could still bind and stabilize 7SK and rescue all *bin3* mutant phenotypes,

indicating that Bin3 catalytic activity is dispensable for 7SK stability and snRNP function *in vivo*. Finally,

we identified a metazoan-specific motif (MSM) outside of the methyltransferase domain and generated

mutant flies lacking this motif (Bin3$^{\Delta MSM}$). Bin3$^{\Delta MSM}$ mutant flies exhibited some—but not all—*bin3*

mutant phenotypes, suggesting that the MSM is required for a 7SK-independent, tissue-specific

function of Bin3.

INTRODUCTION

Regulation of eukaryotic RNA polymerase II (RNAPII) occurs at multiple stages of the

transcription cycle, including RNAPII recruitment to promoters by enhancer binding proteins, the

establishment of the pre-initiation complex, and promoter release. In higher eukaryotes an additional

mechanism, promoter-proximal pausing, has been shown to be critical for RNAPII regulation in

development and in response to signaling events (Margaritis and Holstege 2008; Gilmour 2009; Chiba

et al. 2010; Levine 2011; Li and Gilmour 2011; Nechaev and Adelman 2011; Adelman and Lis 2012; Zhou

et al. 2012; Gaertner and Zeitlinger 2014; Jonkers and Lis 2015; Liu *et al.* 2015; Mayer *et al.* 2017; Chen *et*

al. 2018; Core and Adelman 2019; Dollinger and Gilmour 2021; Gonzalez *et al.* 2021; Abuhashem *et al.*

2022). Pausing occurs when RNAPII stalls after transcribing approximately 30-60 nucleotides, and is

induced by the binding of negative elongation factors (Yamaguchi *et al.* 1999; Wang *et al.* 2010) and by

hypo-phosphorylation of serine 2 (Ser2) in the heptad repeat (YSPTSPS) of the carboxy-terminal

domain of the large subunit of RNAPII (reviewed in Brookes and Pombo 2009). Promoter proximal

pausing is relieved by the kinase activity of Positive Transcription Elongation Factor b (P-TEFb; Marshall

and Price 1995), which is a heterodimer composed of Cyclin-dependent kinase 9 (Cdk9; Marshall *et al.*

1996) and the regulatory cyclin, Cyclin T (CycT; Peng *et al.* 1998). P-TEFb phosphorylates negative

elongation factors (Wada *et al.* 1998; Cheng and Price 2007) and Ser2 (Zhou *et al.* 2000; Ramanathan *et*

al. 2001; Shim *et al.* 2002; Ni *et al.* 2004), which allows RNAPII to enter productive elongation (Ni *et al.*

2008).

To maintain a balance between transcription elongation and pausing, P-TEFb activity is highly

regulated; without this regulation, P-TEFb becomes "hyper-activated" and induces aberrant

transcription elongation (Schneeberger *et al.* 2019). Key to this regulation is the 7SK snRNP, which

sequesters (and thereby represses) P-TEFb (Nguyen *et al.* 2001; Yang *et al.* 2001). The "core" 7SK

snRNP is composed of the 7SK noncoding RNA, which acts as a scaffold for the proteins

Methylphosphate Capping Enzyme (MEPCE) and La-Related Protein 7 (LARP7) (Krueger *et al.* 2008;

Barboric *et al.* 2009; Xue *et al.* 2010; Muniz *et al.* 2013; Brogie and Price 2017; Yang *et al.* 2022). MEPCE

and LARP7 protect the 5' and 3' ends of 7SK from exoribonucleolytic degradation, respectively

(Jeronimo *et al.* 2007; Markert *et al.* 2008; Barboric *et al.* 2009; Xue *et al.* 2010; Muniz *et al.* 2013). After

formation of the core snRNP, 7SK RNA is reversibly bound by a homodimer of Hexamethylene Bis-

Acetamide Inducible Protein (HEXIM) (Blazek *et al.* 2005; Li *et al.* 2005). Binding of HEXIM to 7SK

induces a conformational change that allows HEXIM to bind to P-TEFb (Michels *et al.* 2004),

functionally sequestering and inactivating P-TEFb (Yik *et al.* 2004). Upon cellular stress and other

cellular signals (reviewed in Liu *et al.* 2015), binding of accessory proteins to 7SK (Herreweghe *et al.*

2007; Barrandon *et al.* 2007; Krueger *et al.* 2008; Lemieux *et al.* 2015; Bugai *et al.* 2019) and a

concomitant conformational change in the secondary structure of 7SK RNA (Krueger *et al.* 2010; Yang

et al. 2022) releases P-TEFb from the repressive 7SK snRNP. P-TEFb is then able to relieve pausing to

enable transcription elongation.

MEPCE is the long sought-after RNA methyltransferase that uses the methyl donor S-adenosyl-

L-methionine (SAM) to add a monomethyl cap to the gamma phosphate on the 5' end of 7SK (Gupta *et*

al. 1990a; Shumyatsky *et al.* 1990; Shimba and Reddy 1994), an RNA polymerase III transcript (Zieve

and Penman 1976; Murphy *et al.* 1986, 1987, 1989; Shumyatsky *et al.* 1990). MEPCE is human ortholog

of Bin3, a *Drosophila* protein that is the founding member of the Bin3/MEPCE/Bmc1 family of RNA

methyltransferases (Zhu and Hanes 2000; Jeronimo *et al.* 2007; Barboric *et al.* 2009; Cosgrove *et al.*

2012; Páez-Moscoso *et al.* 2022; Porat *et al.* 2022). We originally discovered Bin3 as an interacting

partner of the Bicoid homeodomain protein (Zhu and Hanes 2000). Bicoid acts as both an activator of

zygotic gene transcription, and a translational repressor of maternal *caudal* mRNA during

embryogenesis; both functions are critical for anterior-posterior patterning (Dubnau and Struhl 1996;

Rivera-Pomar *et al.* 1996; Chan and Struhl 1997; Niessing *et al.* 1999, 2000, 2002; Cho *et al.* 2005). We

had hypothesized that Bin3 was an arginine methyltransferase that methylated Bicoid to alter its

function, transitioning Bicoid from being a transcriptional activator to a translational repressor, or vice

versa. However, identification and characterization of MEPCE in human cells (originally called BCDIN3)

demonstrated that it is actually an RNA methyltransferase that targets 7SK to form a snRNP that

represses transcription (Jeronimo *et al.* 2007). We later showed that early in *Drosophila* embryogenesis,

Bin3 stabilizes 7SK to form a novel snRNP containing Bin3 and Bicoid (and potentially other

translational factors) that represses translation of maternal *caudal* mRNA at the anterior of the embryo

(Singh *et al.* 2011). It is important to note that Bicoid is an insect-specific protein, and therefore, the

translational repression function of Bin3 is also likely to be insect-specific. The canonical role of

Bin3/MEPCE in regulating transcription elongation, and the importance of this regulation to normal

development, has not been extensively studied in *Drosophila* or in other model organisms (see below).

Much of what we know about the 7SK RNA binding and capping functions of MEPCE comes

from *in vitro* or cell culture-based studies focusing on the methyltransferase activity of MEPCE and the

stepwise assembly of the 7SK snRNP. The monomethyl cap is believed to protect the 5' end of 7SK

from exoribonucleolytic degradation (Shumyatsky *et al.* 1993). However, MEPCE is peculiar in that

unlike most enzymes, it remains bound to the product of the reaction it catalyzes, calling into question

the essentiality of methyl capping. MEPCE that is constitutively bound to capped 7SK facilitates the

stable binding of LARP7 (Xue *et al.* 2010) to form the core 7SK snRNP (Muniz *et al.* 2013). Several

residues in the active site have been identified as being essential for binding to SAM and/or 7SK (Xue *et*

al. 2010; Shelton *et al.* 2018), and for methyltransferase activity (Yang *et al.* 2019). Pre-association of

MEPCE with 7SK was found to enhance the recruitment of LARP7 to 7SK and to MEPCE itself (Xue *et al.*

2010). However, whether methyltransferase activity is required *in vivo* for 7SK binding and stability, and

7SK snRNP function, has never explicitly been tested. Moreover, these studies have focused primarily

on the function of the methyltransferase domain. During evolution, MEPCE orthologs have acquired

extensive protein sequence flanking the methyltransferase domain, the conservation and function(s) of

which have not been explored.

Underscoring the need to understand the function of MEPCE *in vivo*, there is a

neurodevelopmental disorder that is caused by heterozygosity for a nonsense mutation in *MEPCE* that

results in P-TEFb hyper-activation (Schneeberger *et al.* 2019). However, our ability to study the *in vivo*

function of MEPCE and its orthologs is limited by the fact that MEPCE function appears to be essential

in vertebrate model organisms. For example, transfecting HeLa cells with siRNAs to destabilize 7SK

(which phenocopies the destabilization of 7SK in the absence of MEPCE function) results in apoptosis

(Haaland *et al.* 2005); treating zebrafish embryos with morpholino oligonucleotides targeting *mepcea*

(the zebrafish ortholog of *MEPCE*; originally named *bcdin3*) is embryonic lethal (Barboric *et al.* 2009),

precluding the ability to study Mepcea function in adult zebrafish; and deleting 7SK in mice (which also

phenocopies the destabilization of 7SK in the absence of MEPCE function) is lethal (Xu *et al.* 2022),

precluding the ability to study MEPCE function in adult mice. However, deleting *bin3* (the *Drosophila*

ortholog of *MEPCE*) does not affect the viability of adult *Drosophila* (Singh *et al.* 2011). Therefore,

Drosophila Bin3 provides a model to study the *in vivo* functions of MEPCE in metazoan development.

Here, we investigated the role of Bin3 in *Drosophila* development. We found that Bin3 is

required for fecundity and has a conserved role in promoting neuromuscular function, by repressing P-

TEFb. Unexpectedly, we found that Bin3 catalytic activity is dispensable for 7SK stability and snRNP

function *in vivo*, challenging the paradigm that methyl capping is essential for 7SK stability. Finally, we

identified a metazoan-specific motif (MSM) outside of the methyltransferase domain that confers a

7SK-independent, tissue-specific function to Bin3. These studies illustrate the importance of

establishing a model system in which to study this important and conserved enzyme.

MATERIALS AND METHODS

Fly stocks and husbandry

Fly stocks from the Bloomington *Drosophila* Stock Center (BDSC) used in this study are listed in

Table S1. Fly stocks we constructed for this study are listed in **Table S2**. The genotypes of maternal and

paternal flies, and the genotypes of their progeny used for experiments, and the figures in which those

experiments appear, are listed in **Table S3**. Fly were reared at 25ºC and 60-80% humidity on a 12-hour

light/dark cycle, on either Nutri-Fly MF food (Genesee Scientific) with ~0.064 M propionic acid (Sigma

Aldrich, P5561; or Apex Bioresearch Products, 20-271), or food made based on the Nutri-Fly MF

formula: 25 g/L Inactive Dry Yeast (Genesee Scientific), 89.5 g/L Dry Molasses (Genesee Scientific), 57

g/L Fly Stuff Yellow Cornmeal (Genesee Scientific), 5.84 g/L Nutri-Fly *Drosophila* Agar, Gelidium

(Genesee Scientific), ~0.064 M propionic acid (Apex Bioresearch Products).

bin3 **mutants**

We previously described two excision alleles that remove some (*bin3*$^{4-7}$) or all (*bin3*$^{2-7}$) of the *bin3*

coding exons (Parks *et al.* 2004; Singh *et al.* 2011). Both alleles were made in the same background, and

when placed in *trans* to produce *bin3*$^{4-7}$/*bin3*$^{2-7}$ mutants, incurred background effects likely due to

homozygosity for recessive second-site mutations. To produce *bin3* mutants that avoid these

confounding effects, we crossed flies heterozygous for *bin3*$^{2-7}$ (**Fig. 1a**) with flies heterozygous for a

deficiency uncovering *bin3* (*Df*, **Fig. 1a**), which is in a different genetic background than *bin3*$^{2-7}$ (Ryder *et*

al. 2004).

Generation of transgenic *bin3*⁺ rescue flies

We previously showed that Bin3 is required for embryonic viability by demonstrating a *bin3* cDNA under

control of the maternal *Hsp83* promoter rescued the embryonic lethality of eggs laid by *bin3* mutant

females (Singh *et al.* 2011). To examine the role of Bin3 more broadly, we needed a rescue approach

that would faithfully express *bin3* under control of its own regulatory sequences in all *bin3*-expressing

tissues, not just in the ovary. To do this, we used BAC clone CH322-154G8 (BACPAC Genomics) that

comprises a 22.065 kb fragment of chromosome 2R, containing the *bin3* P1 promoter (Singh *et al.*

2011), and 3 of the 4 predicted mRNA isoforms of *bin3* (*bin3-RA*, *bin3-RC*, and *bin3-RD*). This BAC was

inserted into the *attP* landing site on chromosome 3R in BDSC stock #9744 (*y^1 w^{1118}; PBac{y$^+$-attP-*

9A}VK00027) by BestGene, Inc.

Generation of the pUAS-DSCP-XbaI-attB vector via NEBuilder HiFi DNA Assembly

High-copy plasmids pNP (Qiao *et al.* 2018; Wang *et al.* 2019) and pUASP-attB (Drosophila Genomics

Resource Center Stock 1358; https://dgrc.bio.indiana.edu//stock/1358; RRID:DGRC_1358; Takeo *et al.*

2012) were purified from bacterial cells grown to confluence at 37ºC in LB + 100 µg/ml ampicillin using

the Monarch Plasmid Miniprep Kit T1010L from New England BioLabs, Inc. (NEB). To create pUAS-

DSCP-XbaI-attB (the vector into which *bin3* and 3xHA sequences would be cloned, see below), the

10XUAS-DSCP promoter was amplified from plasmid pNP with Q5 High-Fidelity DNA Polymerase

(NEB) using primers UP0394 and UP0395, and inserted into pUASP-attB that had been digested with

*Xho*I (NEB) and *Pst*I (NEB), via NEBuilder HiFi DNA Assembly (NEB.). NEBuilder HiFi DNA Assembly

destroyed the *Xho*I site, and partially eliminated one of the two GAGA enhancers in pUASp-attB .

Ovarian RNA Purification and Reverse Transcription for NEBuilder HiFi DNA Assembly

one to three-day old *w^{1118}* (BDSC #5905) females were incubated with males in yeasted vials overnight.

Five pairs of ovaries were dissected from two to four-day old females in ice-cold 1X PBS, and flash-

frozen in liquid nitrogen. Total RNA was extracted using the Monarch Total RNA Miniprep Kit (NEB),

according to the manufacturer's instructions for RNA purification from "Tissues". Total RNA (1 µg) was

reverse transcribed using ProtoScript II First Strand cDNA Synthesis Kit (NEB, E6560S) with primer

UP0118, which anneals to the very end of *bin3* coding sequence (CDS). See **Table S4** for primer

sequence.

Purification of BAC clone CH322-154G8 for NEBuilder HiFi DNA Assembly

EPI300 bacterial cells (Lucigen) harboring the BAC clone CH322-154G8 (BACPAC Genomics) were

grown to confluence at 37ºC in LB + 12.5 µg/ml chloramphenicol, and plasmid replication was induced

by diluting 1:10 in LB + 12.5 µl/ml chloramphenicol and 1X CopyControl Induction Solution (Lucigen).

The BAC was purified purified using the Monarch Plasmid Miniprep Kit (NEB).

Generation of *UAS-DSCP-3xHA-bin3* transgenic flies via NEBuilder HiFi DNA Assembly

To generate plasmid *pUAS-DSCP-3xHA-bin3-attB*, which expresses Bin3 C-terminally tagged with 3xHA,

under UAS control: Fragment #1, comprising the *bin3-RA* 5' -UTR to the end of the *bin3* CDS, was

amplified from *bin3* cDNA using Q5 High-Fidelity DNA Polymerase, and primers UP0423 and UP0414.

The 5' end of Fragment #1 is homologous to sequences immediately upstream of the *Xba*I site in

pUAS-DSCP-XbaI-attB; the 3' end of Fragment #1 is homologous to 5' end of a synthetic dsDNA

fragment comprising Gly-Gly-Gly-Ser-3xHA (G$_4$S-3xHA; Integrated DNA Technologies) codon-

optimized for *Drosophila* with the IDT Codon Optimization Tool (Integrated DNA Technologies).

Fragment #2, comprising the entire *bin3-RA* 3' -UTR, was amplified from the BAC clone CH322-154G8

using Q5 High-Fidelity DNA Polymerase, and primers UP0415 and UP0424. The 5' end of Fragment #2

is homologous to the 3' end of the G$_4$S-3xHA fragment; the 3' end of Fragment #2 is homologous to

sequences immediately downstream of the *Xba*I site in pUAS-DSCP-XbaI-attB. Fragment #1, G$_4$S-

3xHA, and Fragment #2 were assembled in this order into the *Xba*I site of pUAS-DSCP-XbaI-attB via

NEBuilder HiFi DNA Assembly. The *Xba*I site was destroyed.

To generate plasmid *pUAS-DSCP-3xHA-bin3*^{Y795A}*-attB*, which expresses catalytically-dead Bin3^{Y795A}, C-

terminally tagged with 3xHA ,under UAS control: Fragment #3, comprising the *bin3-RA* 5' -UTR to 11 nt

downstream of the Y795 codon, was amplified from *bin3* cDNA using Q5 High-Fidelity DNA

Polymerase, and primer UP0423 and the mutagenic primer UP0419, which introduces a Y795A point

mutation. The 5' end of Fragment #3 is homologous to sequences immediately upstream of the *Xba*I

site in pUAS-DSCP-XbaI-attB; the 3' end of Fragment #3 is homologous to the 5' end of Fragment #4.

Fragment #4, comprising 11 nt upstream of the Y795 codon to the end of the *bin3-RA* CDS was

amplified from cDNA using Q5 High-Fidelity DNA Polymerase, and mutagenic primer UP0420, which

introduces a Y795A point mutation, and primer UP0414. The 5' end of Fragment #4 is homologous to

the 3' end of Fragment #3; the 3' end of Fragment #4 is homologous to 5' end of the G$_4$S-3xHA

fragment. Fragment #3, Fragment #4, G$_4$S-3xHA, and Fragment #2 (see above) were assembled in this

order into the *Xba*I site of pUAS-DSCP-XbaI-attB via NEBuilder HiFi DNA Assembly. The *Xba*I site was

destroyed.

To generate plasmid *pUAS-DSCP-3xHA-bin3*$^{\Delta MSM}$*-attB*, which expresses Bin3$^{\Delta MSM}$, C-terminally tagged

with 3xHA, under UAS control: Fragment #5, comprising the *bin3-RA* 5' -UTR to 12 nt downstream of

the L326 codon, was amplified from *bin3* cDNA using Q5 High-Fidelity DNA Polymerase, and primer

UP0423 and the mutagenic primer UP0417, which eliminates codons 311-326. The 5' end of Fragment

#5 is homologous to sequences immediately upstream of the *Xba*I site in pUAS-DSCP-XbaI-attB; the 3'

end of Fragment #5 is homologous to the 5' end of Fragment #6. Fragment #6, comprising 12 nt

upstream of the F311 codon to the end of the *bin3-RA* CDS was amplified from cDNA using Q5 High-

Fidelity DNA Polymerase and mutagenic primer UP0418, which eliminates codons 311-326, and primer

UP0414. The 5' end of Fragment #6 is homologous to the 3' end of Fragment #5; the 3' end of

Fragment #6 is homologous to 5' end of the G_4S-3xHA fragment. Fragment #5, Fragment #6, G_4S-

3xHA, and Fragment #2 (see above) were assembled in this order into the *Xba*I site of pUAS-DSCP-

XbaI-attB via NEBuilder HiFi DNA Assembly. The *Xba*I site was destroyed.

See **Table S4** for primer sequences and the sequence of the G_4S-3xHA synthetic dsDNA fragment used

to make these 3 constructs. Sequencing of each construct revealed that all UTRs corresponded to the

bin3-RA mRNA isoform of *bin3*. Interestingly, each construct had an in-frame insertion of an Ala residue

between L573 and V574, when mapped to the reference sequence of *bin3* (NCBI Reference Sequence

NM_165468.2). These 3 constructs were recombined into the *attP* landing site on chromosome 3L in

BDSC stock #8622 ($y^1 w^{67c23}$; $P\{y^{+t7.7}=CaryP\}attP2$) by BestGene, Inc.

Expression of *UAS-DSCP-3xHA-bin3* transgenes using the Trojan GAL4 approach

We used the UAS/GAL4 system (Brand and Perrimon 1993) and the Trojan GAL4 approach (Diao and

White 2012; Diao *et al.* 2015; Lee *et al.* 2018) to induce the expression of *UAS-DSCP-3xHA-bin3* (hereon

referred to as *UAS-bin3*) transgenes. Intronic Trojan GAL4 insertions produce a null allele of a gene

while also expressing GAL4 under the regulatory sequences of that gene. A Trojan GAL4 insertion into

one of the *bin3* introns (*bin3$^{MI08045-TG4.0}$*, hereon referred to as *bin3$^{TG4.0}$*) in *trans* to a deficiency (*Df*)

uncovering *bin3* produces *bin3*-null (*bin3$^{TG4.0}$/Df*) flies that express GAL4 in the spatiotemporal pattern of

endogenous *bin3*. (**Fig. 4a**). When transgenes comprising *UAS-bin3* cDNAs (**Fig. 5d-f**) are incorporated

into *bin3$^{TG4.0}$/Df* flies, GAL4 induces expression of Bin3 proteins in the pattern of endogenous *bin3* in an

otherwise *bin3*-null background.

Total RNA purification from ovaries and RT-qPCR

Five one to three-day old females were incubated with males in yeasted vials overnight. Five pairs of

ovaries were dissected from two to four-day old females in ice-cold 1X PBS, and flash-frozen in liquid

nitrogen. Total RNA was extracted using the Monarch Total RNA Miniprep Kit (NEB), according to the

manufacturer's instructions for RNA purification from "Tissues". RT-qPCR was performed on 10 ng total

RNA using the Luna Universal One-Step RT-qPCR Kit (NEB, E3005L). *bin3*, 7SK, U6, and *rp49* mRNAs

were reverse transcribed and amplified with primers UP0493 + UP0118, UP0146 + UP0147, OW1123 +

OW1124, and UP0113 + UP0114, respectively (see **Table S4** for primer sequences). Reactions were

performed using a CFX Opus 384 Real-Time PCR System (Bio-Rad).

Protein extraction and Western Analysis

Ten one to three-day old females were incubated with males in yeasted vials overnight. Ten pairs of

ovaries were dissected from two to four-day old females in ice-cold 1X PBS, and flash-frozen in liquid

nitrogen. Protein was exacted as previously described {Prudêncio.2015}. Extracts were clarified through

0.22 μM Spin-X columns (Costar), and mixed 1:1 with 2X Sample Loading Buffer (Bio-Rad) + 5% β-

mercaptoethanol, and boiled for 5 minutes. Extracts were spun down, and 15 μl was loaded onto 4-20%

TGX gels (Bio-Rad), and ran at 100 V for 10 minutes, then 300 V for 20 minutes. Protein was transferred

to PVDF-LF membranes (Bio-Rad) using a Trans-Blot Turbo Transfer System (Bio-Rad) using the High

MW preset (1.3 A, 25 V, 10 minutes). Membranes were blocked in EveryBlot Blocking Buffer (Bio-Rad)

for 5 minutes, and then incubated with primary antibodies diluted in EveryBlot Blocking Buffer for 1

hour at room temperature. Membranes were washed with TBST (1X Tris-Buffered Saline, 0.2% Tween-

20) for 5 minutes 3 times, and then incubated with secondary antibodies diluted in EveryBlot Blocking

Buffer for 1 hour at room temperature. Membranes were with TBST for 5 minutes 3 times, dehydrated

with 100% methanol, and imaged using a ChemiDoc MP Imaging System with Image Lab Touch

software (Bio-Rad). Primary antibodies used were 1 μg/ml rabbit anti-HA (abcam, ab9110), and 1 μg/ml

mouse anti-beta Actin (abcam, ab170325), and 5 µg/ml IgG purified from sheep anti-CycT antiserum

(Nguyen *et al.* 2012). using Pierce Protein G Magnetic Beads according to the manufacturer's

instructions. Secondary antibodies were from Jackson ImmunoResearch, Inc., and used at 1 µg/ml:

Alexa Fluor 488 AffiniPure Donkey Anti-Rabbit IgG (711-545-152), Alexa Fluor 488 AffiniPure Donkey

Anti-Sheep IgG (713-545-147), Alexa Fluor 594 AffiniPure Donkey Anti-Mouse IgG (715-585-150), and

Alexa Fluor 647 AffiniPure Donkey Anti-Mouse IgG (715-605-150).

Egg-Laying Assay

Ten one to three-day old virgin females were incubated with wild-type males *en masse* in yeasted vials

overnight. Ten two to four-day old females were split individually into fresh vials with two Oregon-R

males, and incubated for 24 hours. After 24 hours, eggs were counted to determine the number of eggs

laid/female/day. Experiments were performed in biological triplicate.

Immunofluorescence staining of ovaries

Immunofluorescence staining of ovaries was performed as described (Maimon and Gilboa 2011). Ten 1-

4 day old females were incubated with males in yeasted vials overnight, and then 10 pairs of ovaries

were dissected from 2-5 day old females in ice-cold 1X PBS. The following steps were performed at RT

unless otherwise indicated. Ovaries were fixed in 1 ml of Image-iT Fixative Solution (Thermo Scientific,

R37814) + 0.3% Triton X-100 for 20 minutes. Ovaries were washed and permeabilized with 1 ml of 1% PT

Buffer (1X PBS, 1% Triton X-100) for 5 minutes, 10 minutes, and 45 minutes. Ovaries were blocked with

1 ml PBT (1X PBS, 1% BSA, 0.3% Triton X-100) for 1 hour, and then incubated in 250 µl PBT and primary

antibodies overnight at 4°C. Primary antibodies used were 1 µg/ml rabbit anti-HA (abcam ab9110) and 5

µg/ml mouse anti-Vasa (Developmental Studies Hybridoma Bank, Vasa 46F11). Ovaries were washed

with 1 ml 0.3% PT Buffer (1X PBS, 0.3% Triton X-100) 3 times for 30 minutes each, blocked with 1 ml

PBT-NDS (1X PBS, 1% BSA, 0.3% Triton X-100, 5% normal donkey serum) for 1 hour, and incubated in

250 µl PBT-NDS containing 10 µg/ml Hoechst 33342 (Thermo Scientific, H3570) and 2 µg/ml secondary

antibodies for 2 hour in the dark. Secondary antibodies were from Jackson ImmunoResearch, Inc.:

Alexa Fluor 488 AffiniPure Donkey Anti-Rabbit IgG (711-545-152) and Alexa Fluor 647 AffiniPure Donkey

Anti-Mouse IgM (715-605-020). Ovaries were washed with 1 ml 0.3% PT Buffer in the dark 3 times for 30

minutes each, then mounted in 50 µl VECTASHIELD PLUS Antifade Mounting Medium (Vector Labs,

Inc., H-1900-10).

Single-molecule fluorescence *in situ* hybridization (smFISH) in ovaries

smFISH on adult ovaries was performed essentially as described (Raj and Tyagi 2010; Abbaszadeh and

Gavis 2016). Ten 1-4 day old females were incubated with males in yeasted vials overnight, and then 10

pairs of ovaries were dissected from 2-5 day old females in ice-cold 1X PBS. The following steps were

performed at RT unless otherwise indicated. Ovaries were fixed in 1 ml of Image-iT Fixative Solution

(Thermo Scientific, R37814) for 30 minutes, then washed with PBST (1X PBS, 0.1% Tween-20) 3 times

for 5 minutes each. Ovaries were permeabilized by incubating in 7:3 PBST:methanol for 5 minutes, 1:1

PBST:methanol for 5 minutes, 3:7 PBST:methanol for 5 minutes, 100% methanol for 10 minutes, 3:7

PBST:methanol for 5 minutes, 1:1 PBST:methanol for 5 minutes, and 7:3 PBST:methanol for 5 minutes.

Ovaries were then washed with PBST 4 times for 5 minutes each, followed by pre-hybridized in Wash

Buffer A (20% Stellaris RNA FISH Wash Buffer A (Biosearch Technologies Cat# SMF-WA1-60) and 10%

deionized formamide) for 5 minutes. Ovaries were incubated in Stellaris RNA FISH Hybridization Buffer

(Biosearch Technologies Cat# SMF-HB1-10) with 10% deionized formamide and 125 µM Quasar 640-

labeled oligonucleotide probes, overnight at 37ºC in the dark. Oligonucleotide probes against 7SK were

designed using the LGC Biosearch Technologies' Stellaris RNA FISH Probe Designer

(https://www.biosearchtech.com/support/tools/design-software/stellaris-probe-designer). Ovaries

were washed in the dark with Wash Buffer A for 30 minutes at 37ºC, and then with Wash Buffer A with

10 µg/ml Hoechst 33342 (Thermo Scientific, H3570) for 30 minutes at 37ºC, and finally with Stellaris RNA

FISH Wash Buffer B (Biosearch Technologies Cat# SMF-WB1-20) for 5 minutes at RT. Ovaries were

mounted in 50 µl VECTASHIELD PLUS Antifade Mounting Medium (Vector Labs, Inc., H-1900-10).

Ovariole Counting Assay

One to three-day old females were incubated with sibling males in yeast vials overnight. Two to four-

day females were anesthetized, transferred to 1.7 ml tubes, flash-frozen in liquid nitrogen, and stored at

-80ºC. Frozen females were thawed on ice or at RT and dissected in a 1:1 mixture of methanol and 1X

PBS. 50% methanol dehydrates and extracts lipids from the gonadal sheath surrounding the ovary and

from the clear portions of the ovarioles (turning them opaque), making ovarioles easy to separate and

count. The mean number of ovarioles per ovary, as well as the absolute value of the difference in

ovariole number between each ovary per female were calculated (Lobell *et al.* 2017). Experiments were

performed in biological duplicate.

Climbing Assay

Ten 2-5 day old females were transferred to 15 ml tubes, and allowed to acclimate for 3'. Flies were

tapped to the bottom of the tube, and climbing was recorded for 12 seconds. Flies were then allowed to

reacclimatize for another 3 minutes before performing the subsequent trial. 10 trials were performed

per genotype to determine the percent of flies climbing 8 cm in 12 seconds per trial. All genotypes for a

particular experiment were tested at the same time. Experiments were performed in biological

triplicate.

Modeling of Bin3 protein sequence onto MEPCE

Bin3 was modeled onto 309 amino acids of the MEPCE active site in complex with *S*-adenosyl-*L*-

homocysteine (SAH) and stem loop 1 of 7SK (PDB 6dcb; Yang *et al.* 2019) using hhpred (Söding 2005;

Hildebrand *et al.* 2009; Meier and Söding 2015; Zimmermann *et al.* 2018; Gabler *et al.* 2020) and

MODELLER (Šali *et al.* 1995; Zimmermann *et al.* 2018; Gabler *et al.* 2020). Alignment of the Bin3 model

to 6dcb, RMSD calculation, and modeling of the Y421A and Y795A mutations was performed using

PyMol (version 2.5.4).

Bin3 Disorder Prediction

Disorder in Bin3 was predicted by IUPred3, using the IUPred3 short disorder analysis type (Mészáros *et*

al. 2018; Erdős and Dosztányi 2020; Erdős *et al.* 2021).

RNA Immunoprecipitation (RIP)

Twenty-five one to three-day old females with or without transgenes expressing *bin3-3xHA* cDNAs

were incubated with males in yeasted vials overnight. Twenty-five pairs of ovaries were dissected from

two to four-day old females in ice-cold, 1X PBS, and fixed in 1 ml 1.8% formaldehyde (a mixture of 55%

1X PBS and 45% Image-iT Fixative Solution, Thermo Scientific, R37814) for 15 minutes at RT on a

nutating rotor. Fixation was terminated by addition of 163 µl 2.5 M glycine to a final concentration of

~350 mM, and incubation for 5 minutes at RT on a nutating rotor. Ovaries were washed briefly with with

1X PBS 3 TIMES, and then flash-frozen in liquid nitrogen. Ovaries were homogenized by hand with a

blue pestle on ice in 50 µl ice-cold RIPA Lysis and Extraction Buffer (Thermo Scientific, 89900)

supplemented with 1 mM DTT, 3X Halt Protease and Phosphatase Inhibitor Cocktail (Thermo Scientific,

78440), and 1 U/µl SUPERase•In RNase Inhibitor (Invitrogen, AM2696). 200 µl of additional buffer was

added to the homogenate, and samples were incubated on ice for 15 minutes to allow for complete

nuclear lysis. Homogenate was pelleted by centrifugation at top speed for 1 minute. 2.5 µl of

homogenate (1/100th homogenate volume) was removed and kept on ice in a total of 300 µl 1X Monarch

DNA/RNA Protection Reagent (NEB, T2011L) as the input sample. The remaining homogenate was

used for immunoprecipitation. 25 µl (1/10th homogenate volume) of Pierce Anti-HA Magnetic Beads

(Thermo Scientific, 88836) were washed with 175 µl 1X TBS-T (Pierce 20X TBS Tween 20 Buffer,

Thermo Scientific, 28360), and then with 1 ml 1X TBS-T. Homogenate was added to the washed beads,

and immunoprecipitation was performed for 4 hours at 4°C with end-over-end rotation. Beads were

washed briefly with 1 ml 1X TBS-T 3 times. Bin3-3xHA complexes were purified by two sequential

competitive elutions with 75 µl of Pierce HA Peptide (Thermo Scientific, 26184) diluted to 2 mg/ml in

Nuclease-Free Water (NEB, B1500L), at 37°C for 15 minutes each. The eluates were pooled, and the 150

µl total eluate was added to 150 µl 2X Monarch DNA/RNA Protection Reagent. All samples (inputs and

eluates) were incubated at 65-70°C for 30 minutes to reverse formaldehyde crosslinks. Total RNA was

extracted using the Monarch Total RNA Miniprep Kit (NEB), according to the manufacturer's

instructions for RNA purification from "Tissues". Total RNA was eluted in 100 µl Nuclease-Free Water.

2.9 µl of each sample was used in 10-µl reactions to reverse transcribe and amplify 7SK and U6 snRNAs

with primers UP0146 + UP0147 and OW1123 + OW1124, respectively (see **Table S4** for primer

sequences). Reactions were performed using the Luna Universal One-Step RT-qPCR Kit (NEB, E3005L)

and a CFX Opus 384 Real-Time PCR System (Bio-Rad).

RESULTS

Bin3 is required for female fecundity

Reproductive success is influenced by fertility (offspring viability) and fecundity (the rate of

offspring production). Previously, we showed that Bin3 plays a role in fertility, promoting embryonic

viability by a mechanism of *caudal* translation repression (Singh *et al.* 2011). During that and other

studies, we noticed that *bin3* mutants lay fewer eggs than wild type flies. Therefore, we asked if Bin3

also plays a role in fecundity by promoting a normal egg-laying rate. To do this, we compared the egg-

laying rates of flies that were either wild type for *bin3* (wt), heterozygous for *bin3* (*bin3²⁻⁷/+*), or deleted

entirely for *bin3* (*bin3²⁻⁷* in *trans* to a deficiency that removes the *bin3* locus; *bin3²⁻⁷/Df*). The genomic

regions deleted by *bin3²⁻⁷* and *Df* are depicted in **Fig. 1a**. As expected, *bin3* mRNA levels are reduced by

half in *bin3²⁻⁷/+* heterozygotes and are undetectable in *bin3²⁻⁷/Df* mutants (**Fig. 1b**). Interestingly, we

found that *bin3²⁻⁷/Df* females laid eggs at a significantly reduced rate compared to wt (P = .0001) and

bin3²⁻⁷/+ (P < .0001) females (**Fig. 2a**). The egg-laying rate defect was specific to the loss of *bin3*, as a

single copy of a genomic fragment containing the *bin3* ORF (*bin3⁺*, **Fig. 1a**) restored *bin3* mRNA

accumulation by half (**Fig. 1b**) and rescued the egg-laying defect of *bin3²⁻⁷/Df* females (P < .0001, **Fig.**

2a). Rescue was comparable to that of *bin3²⁻⁷/+* controls. These results demonstrate that Bin3 is

required for egg-laying.

Bin3 promotes female fecundity by repressing P-TEFb

We found that 7SK RNA levels were strongly reduced in *bin3²⁻⁷/Df* females compared to wt (P <

.0001, **Fig. 1c**), as we have previously shown (Singh *et al.* 2011). The canonical function of Bin3/MEPCE

is to stabilize 7SK RNA (Jeronimo *et al.* 2007; Singh *et al.* 2011). 7SK RNA in turn acts as a scaffold for

proteins that form a snRNP that sequesters and represses P-TEFb (Krueger *et al.* 2008; Barboric *et al.*

2009; Xue *et al.* 2010; Muniz *et al.* 2013; Brogie and Price 2017; Yang *et al.* 2022). Therefore,

destabilization of 7SK in $bin3^{2-7}/Df$ females should lead to deregulation (activation) of P-TEFb, which

could be responsible for the egg-laying defect. If this is true, then we expect that reducing the activity

of P-TEFb in $bin3^{2-7}/Df$ females would rescue the egg-laying defect. To genetically reduce the activity of

P-TEFb, we introduced a mutant allele of *CycT* ($CycT^{jlB2}$) into $bin3^{2-7}/Df$ flies, and confirmed that this

mutation reduces the protein levels of the largest of the two CycT isoforms (**Fig. 1e**). Indeed, we found

that reducing the amount of CycT in $bin3^{2-7}/Df$ females rescued the egg-laying defect (P < .0001, **Fig.**

2a), while having no effect on 7SK RNA levels (**Fig. 1c**). These results suggest that Bin3 stabilizes 7SK

RNA to form the 7SK snRNP, which then sequesters and represses P-TEFb to promote a normal egg-

laying rate.

Bin3 and 7SK are expressed ubiquitously in the ovary

To gain insight into the role of Bin3 in egg-laying, we determined where Bin3 (and its substrate

7SK) is expressed and localized in ovaries, which are composed of 16-20 individual ovarioles that can be

thought of as assembly lines of germline development (**Fig. 2b**). We used the UAS/GAL4 (Brand and

Perrimon 1993) and Trojan GAL4 systems (see Materials and Methods; Diao and White 2012; Diao *et al.*

2015; Lee *et al.* 2018), where a Trojan GAL4 insertion into the *bin3* ORF was used to drive the expression

of a 3xHA-epitope-tagged *bin3* transgene in the spatiotemporal pattern of endogenous *bin3* ($bin3^{TG4.0}$ >

UAS-3xHA-bin3). Bin3-3xHA was detected by immunofluorescence staining of ovaries. We found that

Bin3 is expressed in and localized to the nucleic of all ovarian cells (**Fig. 2, c, c'**). Bin3 was found to be

expressed in the germline cells of the germarium (located at the anterior of each ovariole), and in the 15

nurse cells that supply the oocyte with mRNAs and proteins. Bin3 was also found to be expressed in the

somatic cells surrounding the germline cells and the oocyte, which communicate with germline cells to

promote egg chamber and oocyte maturation. Control flies lacking the *UAS-3xHA-bin3* transgene

($bin3^{TG4.0}$/+) yielded no detectable HA staining (**Fig. 2, d, d'**).

To identify where in the ovary 7SK RNA is expressed and localized, we performed single-

molecule fluorescence *in situ* hybridization (Raj and Tyagi 2010; Abbaszadeh and Gavis 2016) on ovaries

from wt and *bin3²⁻⁷/Df* females. We found that in wt ovarioles, 7SK follows an identical expression

pattern to that of Bin3 (compare **Fig. 2, e, e'** with **Fig. 2, c, c'**). As expected, 7SK staining was

undetectable in *bin3²⁻⁷/Df* ovarioles (**Fig. 2, f, f'**), which do not accumulate 7SK RNA (**Fig. 1c**). Taken

together, Bin3 and 7SK are both expressed in all cells of the ovarioles and localize to nuclei—

commensurate with their role in transcription elongation regulation.

Bin3 is not required to establish the proper number of ovarioles

The total number of ovarioles per ovary directly correlates with the number of eggs insects can

lay (Wayne *et al.* 1997; Wayne and Mackay 1998), and is established by proper proliferation and

migration of the somatic terminal filament cells (TFCs; Sahut-Barnola *et al.* 1995, 1996). Given that we

found *bin3²⁻⁷/Df* mutant females had an egg-laying defect, and that Bin3 is expressed in the somatic

cells of the ovarioles, we hypothesized that Bin3 promotes egg-laying by functioning in the TFCs to

establish ovarioles. To test this hypothesis, we used the UAS/GAL4 system (Brand and Perrimon 1993)

with two different *bab1-GAL4* drivers (Cabrera *et al.* 2002) to induce expression of the *UAS-3xHA-bin3*

transgene specifically in the somatic TFCs of *bin3²⁻⁷/Df* females to determine whether this rescued the

egg-laying rate. Unfortunately, we were unable to recover *bin3²⁻⁷/Df; bab1-GAL4/UAS-bin3* flies using

either driver. Both *bab1-GAL4* drivers are expressed in several cell types in addition to the TFCs (Cabrera

et al. 2002), and therefore it is likely that ectopic expression of *bin3* in all *bab1*-expressing cells is lethal.

As an alternative approach, we compared ovariole numbers in *bin3²⁻⁷/Df* females with those of

wt and *bin3²⁻⁷/+* controls. To do this, we dissected ovaries and determined (i) the average number of

ovarioles per ovary, and (ii) the difference in the number of ovarioles within a pair of ovaries (i.e.,

"ovariole asymmetry"; Lobell *et al.* 2017). Unexpectedly, we did not find a significant difference in

either the mean number of ovarioles per ovary (**Fig. 2g**) or in ovariole asymmetry (**Fig. 2h**) in $bin3^{2-7}/Df$

females compared to either wt or $bin3^{2-7}/+$ control females.

Together, these results suggest that Bin3 is not required to establish ovarioles, and therefore

promotes egg-laying by some other mechanism. These results also suggest that Bin3 does not have a

function in TFCs relevant to the establishment of ovarioles. However, these results do not rule out some

other function of Bin3 in TFCs (*e.g.*, cell-cell communication with germline cells) that might promote

egg-laying. The lethality of inducing *bin3* expression in TFCs using the *bab1-GAL4* drivers precludes the

ability to test this hypothesis in any straightforward way.

Expression of Bin3 in the germline is not sufficient to promote fecundity

Bin3 is also expressed in germline cells, therefore, it remained possible that Bin3 functions in

the germline to promote a normal egg-laying rate. To test this hypothesis, we used a *nanos-*

GAL4::VP16 driver (Doren *et al.* 1998) to induce expression of the *UAS-3xHA-bin3* transgene specifically

in the germline cells of $bin3^{2-7}/Df$ females, and asked whether this improved the egg-laying rate (**Fig.**

2g). Results showed that germline expression of *bin3* did not rescue the egg-laying rate defect of $bin3^{2-7}$

$^{7}/Df$; *nanos-GAL4::VP16/UAS-bin3* females compared to $bin3^{2-7}/Df$; *nanos-GAL4::VP16/+* mutant females

(P = .9942, **Fig. 2i**), suggesting that Bin3 does not function in germline cells to promote a normal egg-

laying rate.

Bin3 has a conserved role in neuromuscular function

A nonsense mutation in the *MEPCE* gene of a patient with impaired neuromuscular ability was

shown to reduce MEPCE protein levels, causing P-TEFb hyper-activation and aberrant RNAPII

transcription elongation (Schneeberger *et al.* 2019). If a conserved function of Bin3/MEPCE is to repress

P-TEFb to promote normal neurodevelopment, then it would be expected that *bin3* mutant flies would

also have neuromuscular defects, and that these defects should be rescued by reducing CycT levels

(lowering P-TEFb activity).

Two ways in which neuromuscular function can be assessed in flies is by examining climbing

(Manjila and Hasan 2018) and resting wing posture (Baehrecke 1997). Normally, flies climb to the top of

an enclosure after being knocked to the bottom, and fold their wings one-over-the-other when resting.

We found that a significant percentage of *bin3²⁻⁷/Df* flies had a climbing defect compared to *bin3²⁻⁷/+*

control flies (P = .0003, **Fig. 3a**). Additionally, whereas control *bin3²⁻⁷/+* flies properly folded their wings

(**Fig. 3b**), *bin3²⁻⁷/Df* mutant flies had a *held-out wings* phenotype (**Fig. 3c**). Both defects were specific to

loss of *bin3*, as the *bin3⁺* genomic fragment rescued the climbing (*P* = .0004, **Fig. 3a**) and resting wing

posture defects of *bin3²⁻⁷/Df* flies (compare **Fig. 3d** with **Fig. 3c**). Importantly, both climbing (*P* = .0008,

Fig. 3a) and wing posture (compare **Fig. 3e** with **Fig. 3c**) defects of *bin3²⁻⁷/Df* flies were also rescued by

reducing CycT levels. These results suggest that Bin3 and MEPCE share a conserved role in

neuromuscular function from *Drosophila* to humans, most likely by repressing P-TEFb and preventing

aberrant transcription elongation.

Bin3 catalytic activity is dispensable for 7SK stability and snRNP function *in vivo*

The importance of MEPCE catalytic residues in 7SK binding and capping have been studied

mostly *in vitro* (Xue *et al.* 2010; Shelton *et al.* 2018; Yang *et al.* 2019). To determine the importance of

Bin3 catalytic activity *in vivo,* we sought to generate flies expressing a catalytically-dead Bin3 protein.

We first analyzed the crystal structure of the methyltransferase domain of human MEPCE (PDB 6dcb,

Yang *et al.* 2019). As shown in **Fig. 4a**, Tyrosine 421 (Y421) is juxtaposed between the 5' gamma

phosphate of 7SK and S-adenosyl-*L*-homocysteine (SAH; used in place of the S-adenosyl-*L*-methionine

(SAM) methyl donor for the purposes of crystallization). Y421 hydrogen bonds to the 7SK gamma

phosphate. Mutation of this tyrosine to alanine (Y421A) completely abolished MEPCE catalytic activity

in vitro (Yang *et al.* 2019). We modeled Bin3 onto the crystal structure (**Fig. 4a**) and found that Y795 in

Bin3 (red) aligned well with Y421 in MEPCE (black). We created models (**Fig. 4b**) representing the

Y421A mutation in MEPCE (black) and the concomitant Y795A mutation in Bin3 (red), which revealed a

complete loss of coordination between 7SK, SAH (i.e., SAM), and MEPCE/Bin3.

Given the *in vitro* importance of the Y421 residue in MEPCE for catalytic activity, and that Y795

in Bin3 aligns with Y421, we generated flies expressing either wt Bin3 (Bin3^{wt}) or Bin3^{Y795A} and tested

whether Bin3^{Y795A} was able to (i) bind and stabilize 7SK, and (ii) rescue *bin3* mutant phenotypes. To do

this, we again used the Trojan GAL4 approach. Importantly, not only do Trojan GAL4 insertions express

GAL4 in the pattern of the gene they are inserted into, but they simultaneously generate a null allele of

the inserted gene. Indeed, *bin3* Trojan GAL4 insertion flies (*bin3$^{TG4.0}$/Df*) (**Fig. 4d**) had nearly

undetectable levels of both *bin3* mRNA (*P* < .0001, **Fig. 4e**) and 7SK RNA (*P* < .0001 **Fig. 4f**) compared to

wt, confirming that *bin3$^{TG4.0}$* is a null allele of *bin3*. Additionally, we found that GAL4 expressed from

bin3$^{TG4.0}$ induced both *UAS-bin3^{wt}* (**Fig. h**) and *UAS-bin3^{Y795A}* (**Fig. 4i**) transgenes similarly at the mRNA

(*P* = .2188, **Fig. 4k**) and protein (*P* = .9836, **Fig. 4l**) levels. Thus, these flies could be used to study the

importance of Bin3 catalytic activity *in vivo*.

Bin3 catalytic activity is expected to be required for capping and stabilization of 7SK *in vivo*;

therefore, in Bin3^{Y795A} mutant flies, 7SK should be destabilized. Remarkably, there was no significant

difference in 7SK levels between Bin3^{Y795A} flies (*bin3$^{TG4.0}$/Df; UAS-bin3^{Y795A}*/+) and Bin3^{wt} flies (*bin3$^{TG4.0}$/Df;*

UAS-bin3^{wt}/+; *P* = .9996, **Fig. 5a**), suggesting that Bin3 catalytic activity is not required for the stability

of 7SK *in vivo*. How then might Bin3 stabilize 7SK in the absence of catalytic activity? As suggested by

prior studies (Xue *et al.* 2010), Bin3 might form a stable complex with 7SK independent of its catalytic

function, and this may protect it from degradation. Indeed, we find that Bin3^{Y795A} immunoprecipitated

as much 7SK RNA as did Bin3^{wt} (*P* = .5350, **Fig. 5b**), suggesting that the Y795A mutation does not

diminish the binding of Bin3 to 7SK, and that binding alone is sufficient for stabilization *in vivo*.

Given that Bin3^{Y795A} could bind and stabilize 7SK, we expected that Bin3^{Y795A} would rescue the

fecundity, climbing, and wing posture defects of *bin3* mutant flies. First, we showed that *bin3$^{TG4.0}$/Df*

flies recapitulate the fecundity (compare **Fig. 6a** to **Fig. 2a**), climbing (compare **Fig. 6b** with **Fig. 3a**),

and wing posture (compare **Fig. 6d** with **Fig. 3c**) defects of *bin3$^{2-7}$/Df* flies. As expected, Bin3^{Y795A}

expression rescued the fecundity (*P* = .0008, **Fig. 6a**), climbing (*P* < .0001, **Fig. 6b**), and wing posture

(compare **Fig. 6f** to **Fig. 6d**) defects of *bin3$^{TG4.0}$/Df* flies, similar to Bin3^{wt} (*P* > .9999 for both fecundity

and climbing). Therefore, despite the importance of catalytic activity to 7SK capping *in vitro*, Bin3

catalytic activity and therefore the methylphosphate cap do not seem to be required *in vivo* for 7SK

stability, nor is it likely to affect 7SK snRNP assembly and function (**Fig. 6i**).

A metazoan-specific motif in Bin3 is required to promote specific neuromuscular functions

Much attention has been given to the methyltransferase domain of Bin3 orthologs; however,

metazoans have acquired over evolutionary time additional protein sequences outside of the

methyltransferase domain (Cosgrove *et al.* 2012). To determine whether there is conserved homology

between Bin3 orthologs in these regions, we aligned several Bin3 orthologs that have been previously

characterized, including *Drosophila* Bin3 (Zhu and Hanes 2000; Singh *et al.* 2011; Cosgrove *et al.* 2012),

human MEPCE (Xue *et al.* 2010; Muniz *et al.* 2013; Shelton *et al.* 2018; Schneeberger *et al.* 2019; Yang *et*

al. 2019; Lee *et al.* 2020; Zhang *et al.* 2021), zebrafish Bcdin3 (Barboric *et al.* 2009), and *S. pombe* Bmc1

(Páez-Moscoso *et al.* 2022; Porat *et al.* 2022). We discovered a 16-amino acid motif that is almost

entirely conserved in metazoans (flies, zebrafish, and humans), but is absent in the fission yeast *S.*

pombe (**Fig. 4c**). Therefore, we have named this motif the <u>M</u>etazoan-<u>S</u>pecific <u>M</u>otif (MSM). The N-

terminal portion of this motif is predicted to be more structured (IUPred3 score 0.2733 to 0.3764), while

the C-terminal portion is predicted to be less structured (IUPred3 score 0.4062 to 0.5887). Given the

strong conservation of the MSM, we hypothesized that the MSM must contribute to some important

function of Bin3 and its metazoan orthologs. To test this hypothesis, we used the Trojan GAL4 system

to generate flies expressing either Bin3^{wt} or Bin3 deleted of the entire 16-aa MSM (Bin3$^{\Delta MSM}$) and asked

whether Bin3$^{\Delta MSM}$ could (i) bind and stabilize 7SK, and (ii) rescue *bin3* mutant phenotypes.

554   We found that GAL4 expressed from *bin3*$^{TG4.0}$ induced both the *UAS-bin3*^{wt} (**Fig. 4h**) and *UAS-*

bin3$^{\Delta MSM}$ (**Fig. 4j**) transgenes to express similar levels of mRNA (*P* = .3840, **Fig. 4k**) and protein (*P* =

.7994, **Fig. 4l**). Given that the MSM is outside of the Bin3 methyltransferase domain, we did not expect

that the Bin3 MSM would be required for 7SK stability or binding. Accordingly, there was no significant

difference in 7SK levels (*P* = .3846, **Fig. 5a**) between Bin3$^{\Delta MSM}$ flies (*bin3*$^{TG4.0}$*/Df; UAS-bin3*$^{\Delta MSM}$*/+*) and

Bin3^{wt} flies (*bin3*$^{TG4.0}$*/Df; UAS-bin3*^{wt}*/+*). Also, Bin3$^{\Delta MSM}$ immunoprecipitated as much 7SK (*P* = .9938, **Fig.**

5b) as did Bin3^{wt}.

561   Given that 7SK was bound and stabilized by Bin3$^{\Delta MSM}$, we expected that Bin3$^{\Delta MSM}$ would rescue

the fecundity, climbing, and wing posture defects of *bin3*$^{TG4.0}$*/Df* flies. Indeed, Bin3$^{\Delta MSM}$ expression

rescued the fecundity (*P* = .0061, **Fig. 6a**) and climbing (*P* < .0001) defects of *bin3*$^{TG4.0}$*/Df* flies, similar to

Bin3^{wt} (*P* = .6681, **Fig. 6a**; *P* = .0685, **Fig. 6b**). However, Bin3$^{\Delta MSM}$ did not rescue the *held-out wings*

phenotype of *bin3*$^{TG4.0}$*/Df* flies (compare **Fig. 6g** with **Fig. 6d**), suggesting that while the MSM is

dispensable for Bin3 function in the contexts of egg-laying and climbing, the Bin3 MSM plays a critical

role in normal resting wing posture, independent of 7SK (**Fig. 6i**). Interestingly, reducing CycT dosage in

Bin3$^{\Delta MSM}$ flies (*bin3*$^{TG4.0}$*/Df; UAS-bin3*$^{\Delta MSM}$*/CycT*^{j11B2}) rescued the *held-out wings* defect (compare **Fig. 6h**

with **Fig. 6g**). We think that because Bin3$^{\Delta MSM}$ binds and stabilizes 7SK (and as such, forms the 7SK

snRNP to repress P-TEFb; **Fig. 5b**) it is unlikely that the MSM directly contributes to P-TEFb repression;

instead, genetically reducing P-TEFb activity probably had a non-specific rescuing effect by globally

tapering transcription elongation.

Bin3 binds to the U6 spliceosomal RNA, but is not required for its stability

MEPCE in humans (Jeronimo *et al.* 2007; Muniz *et al.* 2013) and Bmc1 in *S. pombe* (Páez-

Moscoso *et al.* 2022; Porat *et al.* 2022) bind to the U6 spliceosomal snRNA, but are not required for U6

stability. We asked whether Bin3 in *Drosophila* also binds to U6 and whether Bin3 is also dispensable for

U6 stability.

We found that Bin3^{wt} (P = .0113), Bin3^{Y795A} (P = .0033), and Bin3$^{\Delta MSM}$ (P = .0336) all bind to U6

snRNA, compared to un-tagged control. We also found that there is no significant difference in the

amount of U6 bound by Bin3^{Y795A} (P = .7671) or Bin3$^{\Delta MSM}$ (P = .8486) compared to Bin3^{wt} (**Fig. 5d**). These

results show for the first time that Bin3 in *Drosophila* binds to U6, and that neither catalytic activity nor

the MSM are required for binding. It is not surprising that Bin3$^{\Delta MSM}$ binds to U6, as Bmc1 lacks this

region (Cosgrove *et al.* 2012), yet it binds to U6 (Páez-Moscoso *et al.* 2022; Porat *et al.* 2022). And, like

MEPCE and Bmc1, we found that Bin3 is not required for U6 stability. U6 levels (**Fig. 5c**) were not

different between flies expressing Bin3^{wt} and Bin3^{Y795A} (P = .8794) or Bin3$^{\Delta MSM}$ (P = .7529), and were not

significantly affected in *bin3$^{2-7}$/Df* flies (P = .9314, **Fig. 1d**). U6 levels were reduced by approximately half

in *bin3$^{TG4.0}$/Df* flies (P = .0471, **Fig. 4g**); however, these data were just below the threshold for

significance (α = 0.05). Taken together, Bmc1, Bin3, and MEPCE have a conserved function in binding to

the U6 snRNA, but are not required for its stability.

DISCUSSION

In this study, we showed that Bin3, the *Drosophila* ortholog of the 7SK methylphosphate

capping enzyme, MEPCE, has a conserved role in promoting normal metazoan development by

regulating the P-TEFb positive elongation factor (**Figs. 7a, 8a**). We further investigated Bin3 by

determining the *in vivo* requirement for two different regions of the Bin3 protein: the methyltransferase

domain, and a previously undiscovered motif that is specific to metazoan orthologs of Bin3. To our

surprise, we found that the catalytic activity of the methyltransferase domain appears to be

dispensable for the function of Bin3 as a constituent of the repressive 7SK snRNP (**Fig. 7b**). By contrast,

the metazoan-specific motif in Bin3 is required for a tissue-specific function, one that is 7SK-

independent (**Figs. 7c, 8b**). We also provide the first evidence in *Drosophila* for binding of Bin3 to the U6

spliceosomal snRNA. As in other model organisms (Jeronimo *et al.* 2007; Muniz *et al.* 2013; Páez-

Moscoso *et al.* 2022; Porat *et al.* 2022), we found that Bin3 was not required for U6 stability.

Bin3 represses P-TEFb to promote normal fly physiology

The canonical function of MEPCE is to *repress* gene expression by binding to and stabilizing 7SK

snRNA, which acts as a scaffold for proteins that sequester P-TEFb (Peterlin *et al.* 2011; Cosgrove *et al.*

2012). This inhibits P-TEFb, which normally phosphorylates RNAPII to promote transcription

elongation. Consequently, loss of MEPCE function destabilizes 7SK, dissociates the repressive snRNP,

releasing P-TEFb (Fujinaga *et al.* 2023). Hyper-activated P-TEFb induces aberrant gene expression by

ectopic phosphorylation of RNAPII and other factors (He *et al.* 2008; Schneeberger *et al.* 2019). A non-

canonical function of MEPCE is to *activate* gene expression by binding to the tail of histone H4 and

recruiting free P-TEFb to chromatin. This non-canonical function was found be required for the

induction of tumorigenic genes in a breast cancer cell line (Shelton *et al.* 2018). Here, we show that Bin3

contributes to normal fly physiology through a mechanism most consistent with its canonical function

in repression of P-TEFb. We found that the egg-laying, climbing, and wing posture defects of *bin3*

mutants were rescued by genetically reducing the activity of P-TEFb with a mutation in *CycT* (which is

needed for P-TEFb kinase activity). Thus, in fly development, Bin3 appears to repress P-TEFb (and gene

expression) rather than to activate P-TEFb (**Figs. 7a, 8a**).

Bin3 is required for reproductive success

Previously, we found that Bin3 plays a role in fertility by promoting embryonic viability (Singh *et*

al. 2011). Here, we show that Bin3 is required for fecundity by supporting a normal egg-laying rate (**Fig.**

7a). Thus, Bin3 has a role in two different aspects of reproductive success: fertility (offspring viability),

and fecundity (the rate of offspring production).

Although Bin3 and its substrate 7SK are expressed in all germline and somatic cells of the ovary,

germline expression of Bin3 was dispensable for a normal egg-laying rate. A germline function for Bin3

might occur later in development, consistent with our previous finding that Bin3 appears to regulate the

translation of *grk* in late-stage oocytes (Singh *et al.* 2011), which is important for embryonic patterning.

A previous report demonstrated that both wild-caught flies harboring indels in *bin3*, and

transgenic flies harboring transposon insertions into *bin3*, have reduced numbers of ovarioles (Lobell *et*

al. 2017). However, we found that our *bin3²⁻⁷/Df* mutant females did not exhibit reduced ovariole

numbers compared to controls. We suggest that the difference between our findings and those of

Lobell *et al*. (2017) might be attributable to differences in genetic backgrounds. Nevertheless, the wild-

type number of ovarioles established in our *bin3* mutants is not sufficient to promote a normal egg-

laying rate. Egg-laying is in part a neuromuscular process, as muscle contractions of the ovary and

uterus promote successful oviposition (i.e., egg-laying; Deady and Sun 2015; Irizarry and Stathopoulos

2015; Liao and Nässel 2020; Garrett *et al.* 2023). It is possible, therefore, that defects in neuromuscular

function that we observe in *bin3* mutants (see below) might also explain why *bin3* mutant females lay

fewer eggs, *e.g.*, due to a reduction in the frequency or strength of these contractions.

Conserved role of Bin3 and MEPCE in metazoan neurodevelopment

642   A patient with a neurodevelopmental disorder characterized by mobility and musculature

deficiencies was shown to carry a heterozygous nonsense mutation in *MEPCE* (Schneeberger *et al.*

2019). Similarly, we found that *bin3* mutant flies exhibited difficulties in climbing and an inability to

properly fold their wings, phenotypes that are characteristic of neuromuscular defects (Shukla *et al.*

2013; Kairamkonda and Nongthomba 2014; Sanhueza *et al.* 2014; Manjila and Hasan 2018). Many

mutations in *Drosophila* can affect wing posture (see below), including those in the highly-conserved

JAK/STAT pathway (Harrison *et al.* 1998; Callus and Mathey-Prevot 2002; Ayala-Camargo *et al.* 2013;

Hatini *et al.* 2013; Johnstone *et al.* 2013). Therefore, the conserved roles of Bin3 and MEPCE in

metazoan neurodevelopment may occur through the JAK/STAT pathway. While the exact mechanisms

remain to be determined, the phenotypes we observe in *bin3* mutants (egg-laying, climbing, wing

posture) can be attributable to neuromuscular defects, suggesting a strong conservation in the

developmental function of Bin3 and MePCE in *Drosophila* and human development.

654   Bin3/MEPCE methylates the gamma phosphate on the 5′ end of the 7SK snRNA, while another

class of RNA methyltransferases methylates the N^6 position of internal adenosines (N^6-

methyladenosine, m^6A) on "RRACH" motifs (Wang *et al.* 2018) in a variety of RNAs (Lence *et al.* 2018).

Interestingly, both human 7SK (Warda *et al.* 2017; Leger *et al.* 2021; Perez-Pepe *et al.* 2023) and mouse

7SK (Xu *et al.* 2022) contain the m^6A modification. In HeLa cells, methylation of 7SK by the METLL3

m^6A "writer" protein in response to EGF signaling allows hnRNPs to associate with 7SK; this causes

HEXIM to dissociate from the 7SK snRNP and release P-TEFb, which activates transcriptional

elongation (Perez-Pepe *et al.* 2023). In mice, methylation of 7SK at adenine 281 by METLL3 is required

for the binding of m^{6}A "reader" proteins to establish proper 7SK secondary structure (Xu *et al.* 2022).

Drosophila 7SK contains a single RRACH motif at position 397-401; however, it is not known whether

the adenine within this motif is methylated, or if m^{6}A at that position would serve functions similar to

those in HeLa cells or mice. Intriguingly, flies mutant for m^{6}A writers (and m^{6}A readers) exhibit the

same neuromuscular defects in both climbing and resting wing posture (Lence *et al.* 2016; Kan *et al.*

2017) as flies mutant for *bin3* (this study). Therefore, it is possible that *Drosophila* 7SK might also

contain the m^{6}A modification (at A399), and that m^{6}A is important for 7SK stability and function in

neurodevelopmental processes in *Drosophila*.

670   Although Bin3 and the m^{6}A pathway might function synergistically to promote 7SK stability and

function, they seem to act antagonistically on transcription elongation: Bin3 is required to *prevent*

RNAPII pause-release, whereas the m^{6}A pathway has been shown in *Drosophila* to *promote* RNAPII

pause-release (Akhtar *et al.* 2021). Clearly, there is a complex balance between 7SK stability (via Bin3

binding and possible m^{6}A methylation) as well as 7SK functions that sequester P-TEFb (Bin3-mediated

function) or release P-TEFb (m^{6}A modification function) that is governed by the activities of diverse

classes of RNA methyltransferases, which in *Drosophila* contribute to neuromuscular function.

Moreover, it appears that m6A modifications in 7SK might serve species-specific roles in 7SK biology.

678   Neuromuscular disorders arise from developmental defects in motoneurons and/or the muscles

that they control. We attempted to identify the cells and tissues that require Bin3 for neuromuscular

function by performing an RNAi screen using over 90 GAL4 drivers active in specific neuron and muscle

cell types that drive the expression of two different short hairpin RNAs (shRNAs) targeting *bin3* coding

sequence (Ni *et al.* 2011). However, we did not identify any conditions under which *bin3* RNAi

knockdown recapitulated the neuromuscular phenotypes of *bin3* mutant flies. The failure of our screen

might be due to technical reasons such as weak GAL4 drivers and/or insufficient levels of RNAi

knockdown, or they could be due to Bin3 being required in more than one tissue simultaneously for

neuromuscular activity. Indeed, knockdown of *bin3* induced by the Trojan-GAL4 *bin3* allele (which also

functions as a null allele) in the spatiotemporal expression pattern of *bin3* recapitulated the *held-out*

wings phenotype of *bin3* mutant flies (data not shown), suggesting that Bin3 might be required in

multiple cell types simultaneously, at least for normal resting wing posture.

Bin3 catalytic activity is dispensable for 7SK stability and snRNP function *in vivo*

MEPCE modifies the 5′ end of 7SK RNA by catalyzing the addition of a monomethyl cap to the

gamma phosphate (Gupta *et al.* 1990b; Shumyatsky *et al.* 1990; Jeronimo *et al.* 2007; Xue *et al.* 2010;

Yang *et al.* 2019). It has long been thought that the monomethyl cap is essential for 7SK stability. For

example, *in vitro* transcribed capped 7SK was more stable than uncapped 7SK when injected into

Xenopus oocytes (Shumyatsky *et al.* 1993). However, whether 7SK requires a monomethyl cap for

stability *in vivo* has never been adequately addressed. We and others have shown that Bin3 and its

orthologs are essential for 7SK stability, as deletion of *bin3* (Singh *et al.* 2011), siRNA-mediated

knockdown of *MEPCE* in human cells (Jeronimo *et al.* 2007; Xue *et al.* 2010), and morpholino-mediated

disruption of *mepcea* splicing in zebrafish embryos (Barboric *et al.* 2009) reduces 7SK levels.

Importantly, these approaches do not explicitly test the *in vivo* requirement for methyltransferase

activity, which requires the use of catalytic mutants.

Catalytic mutants of MEPCE have been characterized *in vitro*. A triple mutation in the active site

(V447A, L448A, and D449A) abolished methyltransferase activity, likely by preventing the interaction

between MEPCE and the methyl donor *S*-adenosyl- -methionine (SAM). However, this mutant is also

defective for binding to 7SK and therefore it is not a catalytic mutant *per se* (Xue *et al.* 2010). A double

mutant in the active site (G451A and G455A) also abolished MEPCE methyltransferase activity, again,

presumably by preventing the interaction between MEPCE and SAM; however, this mutant does bind

to 7SK (Shelton *et al.* 2018). Finally, a single amino acid mutation in the active site (Y421A) reduced

methyltransferase activity to 0.6% that of the wt enzyme. This is likely due to loss of hydrogen bonds

between Y421 and both an adjacent residue (R425) and the 7SK gamma phosphate, which is the

recipient of the methyl group from SAM (Yang *et al.* 2019). Importantly, Y421A is not expected to

disrupt binding to either the methyl donor co-factor or the RNA substrate, but rather to disrupt the

chemistry of methyl transfer to the gamma phosphate (Yang *et al.* 2022). Therefore, given the precise

and specific nature of this active site mutation, our preferred choice for *in vivo* study of the importance

of methyl capping was a Y795A substitution in Bin3 (orthologous to Y421A in MEPCE).

Unexpectedly, we found that the Bin3[Y795A] mutant did not exhibit any of the physiological

defects exhibited by *bin3* deletion mutants, suggesting that catalytic activity is not required for Bin3

function *in vivo* (**Fig. 8a**). This result could be explained, in part, by our finding that Bin3[Y795A] still bound

and stabilized 7SK no differently than Bin3[wt], despite its expected lack of methyltransferase activity

(**Fig. 7b**). Thus, flies expressing the Bin3[Y795A] allele retain the ability to assemble the 7SK snRNP to

repress P-TEFb and promote normal development. Our findings suggest that *in vivo* capping of

endogenous 7SK is not required for its stability *per se*, so long as Bin3 is able to bind to 7SK.

That the Bin3[Y795A] catalytic mutant protein was still able to bind to 7SK is not entirely surprising.

Y421 in MEPCE (analogous to Y795 in Bin3) is located within a disordered region that becomes ordered

upon binding to 7SK, forming an alpha helix (Yang *et al.* 2022); this disorder-to-order transition is

required for MEPCE methyltransferase activity. When LARP7 binds to 7SK and then associates with

MEPCE, the alpha helix containing Y421 becomes disordered again (Yang *et al.* 2022); this order-to-

disorder transition induced by LARP7 binding inhibits MEPCE methyltransferase activity (Xue *et al.*

2010; Brogie and Price 2017; Yang *et al.* 2022). Therefore, MEPCE bound by LARP7 effectively

phenocopies MEPCE with a Y421A mutation, which is catalytically-dead (Yang *et al.* 2019). Thus, it

appears that Y421 in MEPCE and Y795 in Bin3 are not essential for 7SK binding, but instead are

essential only for methyl capping.

Our finding that the Bin3[Y795A] catalytic mutant protein not only bound 7SK, but stabilized 7SK

despite the expected lack of catalytic activity, suggests that RNA binding contributes more to 7SK

stability than does methyl capping, challenging the assumption that the gamma monomethyl cap is

essential for 7SK stability. This is underscored by the fact that U6 snRNA, which binds to and is

presumed to be capped by Bin3 and its orthologs, is stable in the absence of these proteins (Jeronimo *et*

al. 2007; Muniz *et al.* 2013; Páez-Moscoso *et al.* 2022; Porat *et al.* 2022).

Bin3 and its orthologs are peculiar enzymes, in that they remain associated with the products of

the reaction they catalyze (Xue *et al.* 2010; Yang *et al.* 2019, 2022). MEPCE has a higher affinity both for

capped 7SK and SAH (the products) than uncapped 7SK and SAM (the reactants) (Yang *et al.* 2019).

Therefore, the gamma monomethyl cap might be required not for 7SK stability *per se*, but to enhance

the binding of MEPCE to the 5′ end of 7SK to keep the enzyme bound after catalyzing the

methyltransferase reaction. Constitutive binding of MEPCE to 7SK would serve the dual purposes of

physically protecting 7SK from inorganic phosphatases (for which the gamma monomethyl cap is a

substrate; Shumyatsky *et al.* 1990), and aiding in the recruitment of LARP7, which binds weakly to 7SK

in the absence of MEPCE (Xue *et al.* 2010), but enhances the affinity of MEPCE for 7SK after binding

(Muniz *et al.* 2013; Yang *et al.* 2022). However, our finding that the Bin3[Y795A] catalytic mutant bound as

much 7SK as Bin3[wt] despite the lack of a gamma monomethyl cap is not consistent with such a model

(at least not *in vivo*). Instead, our results support a model in which 7SK RNA secondary structure—and

not the gamma monomethyl cap—is required for Bin3 and its orthologs to bind to 7SK. Indeed, MEPCE

bound equally well to transfected 7SK harboring either the normal gamma monomethyl cap or a tri-

methyl guanosine (TMG) cap (Muniz *et al.* 2013), and Bmc1 (the *S. pombe* ortholog of Bin3) binds to

both gamma monomethyl-capped U6 RNA and TMG-capped telomerase RNA (Páez-Moscoso *et al.*

2022; Porat *et al.* 2022). These findings suggest that a gamma monomethyl cap structure is not

specifically required for constitutive or high-affinity binding by MEPCE and its orthologs. Alternatively,

7SK secondary structure features likely mediate initial association with MEPCE (i.e., prior to the

association of LARP7, which enhances the affinity of MEPCE for 7SK). When MEPCE initially binds to

7SK in the "nascent" conformation (adopted during 7SK synthesis), there is extensive hydrogen

bonding between the methyltransferase domain and the 5′ hairpin of 7SK (Yang *et al.* 2019). Binding of

MEPCE to 7SK requires the first two (Singh *et al.* 1990) to three (Muniz *et al.* 2013) base pairs of the 5′

hairpin, and a portion of the single-stranded DNA 3′ to the hairpin (Shumyatsky *et al.* 1994; Muniz *et al.*

2013; Yang *et al.* 2019). Subsequently, LARP7 binds to the complex of MEPCE and 7SK, and the

conformation of 7SK changes from a "nascent" conformation to a "closed" conformation (adopted after

formation of the core snRNP of MEPCE, 7SK, and LARP7). In this conformation, MEPCE forms

additional hydrogen bonds to 7SK, and thereby has higher affinity for the "closed" conformation of 7SK

than for the "nascent" conformation of 7SK (Yang *et al.* 2022). Therefore, both 7SK secondary structure

and association with LARP7 are likely the major contributors to the constitutive binding of MEPCE to

7SK.

One possible circumstance under which the gamma monomethyl cap on 7SK might be essential

for stability is when the 7SK snRNP is actively repressing P-TEFb. For HEXIM to bind to 7SK and

sequester P-TEFb, 7SK must undergo a conformational change from the "closed" conformation (i.e.,

the "core" snRNP conformation) to the "linear" conformation (i.e., the "repressive" snRNP

conformation). In this conformation, the 5′ end of 7SK is located outside of the MEPCE active site

(Yang *et al.* 2022), and therefore, in the context of the repressive 7SK snRNP, the 5′ end of 7SK might

be solvent-exposed, which would necessitate a gamma monomethyl cap to protect 7SK from

exoribonucleolytic degradation. However, these findings are based on crystal structures using a

recombinant MEPCE comprising only the methyltransferase domain. Our finding that 7SK is not only

bound but is also stabilized by Bin3^{Y795A} catalytic mutant protein (just like Bin3^{wt} protein) leaves open

the possibility that there are regions outside of the methyltransferase domain in full-length

Bin3/MEPCE that protects the 5′ end of 7SK, even when located outside of the active site. Moreover,

the finding that the 5′ end of 7SK might not be bound by MEPCE when in the "linear" conformation

further supports a model in which the gamma monomethyl cap is not required for constitutive binding

of MEPCE to 7SK.

If the monomethyl cap is not required for either 7SK stability or for Bin3 binding, then what is

the purpose of the methyltransferase domain in Bin3 and its orthologs? One possibility is that the

methyltransferase domain is a vestige of an ancestral ortholog of Bin3 that did not remain

constitutively bound to its RNA substrates, and therefore, the monomethyl cap would have been

essential to protect these RNAs from 5′-3′ exoribonucleolytic degradation. The acquisition of

additional protein binding constituents during evolution of a primordial 7SK snRNP complex might have

necessitated that Bin3 remain associated with 7SK (after catalyzing the methyltransferase reaction) to

facilitate the stable interaction of these additional components (like Larp7). This process would have

rendered the monomethyl capping reaction dispensable for 7SK stability, as we observe. A second

possibility, supported by the strong conservation of the methyltransferase domain over evolutionary

time, argues that catalytic activity has an important function that has yet to be identified, perhaps on

substrates other than 7SK.

A metazoan-specific motif in Bin3 facilitates a 7SK-independent, tissue-specific function in

neurodevelopment

S. pombe Bmc1, the most ancient ortholog of Bin3 studied to date, consists only of the

conserved methyltransferase domain (Cosgrove *et al.* 2012; Páez-Moscoso *et al.* 2022; Porat *et al.*

2022). Over the course of evolution, Bin3 orthologs gained additional protein sequence flanking the

methyltransferase domain (Cosgrove *et al.* 2012). We found a 16-amino acid motif N-terminal to the

methyltransferase domain that is highly conserved in metazoans, and is mostly unstructured. Given its

location outside of the methyltransferase domain, we were not surprised to find that this metazoan-

specific motif (MSM) was not required for Bin3 to bind or stabilize 7SK *in vivo*, and consequently, was

dispensable for fecundity and climbing ability. What we did not expect, however, was that flies

expressing Bin3 deleted of the MSM (Bin3$^{\Delta MSM}$) were unable to properly fold their wings, phenocopying

bin3 null mutants (**Fig. 7c**). These findings suggest that the Bin3 MSM plays a tissue-specific role in gene

regulation independent of 7SK (**Fig. 8b**). There is likely one or more MSM-interacting proteins that help

Bin3 regulate gene expression. These proteins might be specifically expressed in the tissues involved in

wing posture (*e.g.*, the direct and indirect flight muscles), or they might be ubiquitously expressed, but

the interaction with the Bin3 MSM is essential for Bin3 function only in tissues contributing to normal

resting wing posture. Moreover, this protein might be incorporated into the core snRNP of 7SK, Bin3,

and Larp7 to generate a novel 7SK snRNP, or the association with Bin3 MSM might be independent of

the 7SK snRNP altogether.

A 7SK-independent function of Bin3 is not without precedent. It was previously shown that

fibroblasts from a patient with a nonsense mutation in *MEPCE* (*MEPCE* haploinsufficient cells) and cells

deleted of *LARP7* (*LARP7* KO cells) both had reduced levels of 7SK, as expected, but that genes up-

regulated in *MEPCE* haploinsufficient cells were not affected in *LARP7* KO cells. Moreover, exogenous

MEPCE rescued the up-regulation phenotype of *MEPCE* haploinsufficient cells, despite only leading to a

small increase in 7SK levels. Based on these findings, Schneeberger *et al*. (2019) hypothesized that

MEPCE might have a 7SK-independent role in regulating transcription, potentially by interacting with

as-yet unknown proteins. Our finding that Bin3$^{\Delta MSM}$ binds and stabilizes 7SK but is not sufficient to

rescue all *bin3* mutant phenotypes supports this hypothesis and suggests that the as-yet unknown

proteins that interact with Bin3/MEPCE to regulate transcription does so through the MSM,

independent of the 7SK snRNP.

Conclusions

In summary, we have carried out the first detailed *in vivo* study of the highly-conserved

Bin3/MEPCE RNA methyltransferase in a metazoan host. The surprising results demonstrate the value

of rigorously testing models for enzyme function in whole organisms. Our study establishes *Drosophila*

as a genetically tractable developmental model for further dissection of Bin3/MEPCE and 7SK non-

coding RNA pathways, and how their disruption may be associated with human disease.

DATA AVAILABILITY

Plasmids and *Drosophila* stocks are available upon request. All reagent inquiries should be directed to

RJP. All data required to confirm the conclusions made in this article are presented in the text, figures,

and tables.

ACKNOWLEDGEMENTS

We are grateful to the Bloomington *Drosophila* Stock Center (NIH Grant P40OD018537) for fly stocks,

the *Drosophila* Genomics Resource Center (NIH Grant 2P40OD010949) for plasmid pUASP-attB; Jian-

Quan Ni for plasmid pNP; BestGene, Inc. for generating transgenic flies; and David Price for anti-CycT

antiserum. We are also grateful to Scott Neal, Mark Bayfield, and Jennifer Porat for helpful discussions.

FUNDING

This work and RJP was supported by grants to S. Hanes (NSF MCB-1515076 and NIH R-01 GM123985-

01) and B. Knutson (NIGMS R01-GM141033).

CONFLICTS OF INTEREST

None declared.

 FIGURE LEGENDS

Figure 1. Molecular characterization of *bin3* mutant and rescue flies. (a) Graphic depicting the ~26 kb

bin3 locus on chromosome 2R (cytogenetic bands 42A13-42A14). Two promoters, P1 and P2 (grey),

drive expression of *bin3* (Zhu and Hanes 2000; Singh *et al.* 2011). The *bin3* ORF (salmon) comprises 6

coding exons. *bin3$^\Delta$* refers to the *bin3$^{2-7}$* excision allele created by FLP/FRT-mediated recombination

between transposon insertions *PBac{RB}bin3^{e02231}* and *P{XP}d02161* (Parks *et al.* 2004; Singh *et al.* 2011).

Df refers to deficiency *Df(2R)ED1612*, which deletes ~829 kb of chromosome 2R, including promoter P1

and all *bin3* coding sequence (Ryder *et al.* 2004). *bin3$^\Delta$* and *Df* in *trans* produce *bin3* mutant flies. *bin3$^+$*

(sky blue) is a transgenic insertion of a BAC comprising ~20 kb of chromosome 2R, including promoter

P1 and all *bin3* coding sequence. In *bin3$^{2-7}$/Df* flies, this insertion rescues *bin3* mutant phenotypes. **(b-d)**

Bar graphs showing the average amount of *bin3* mRNA (b), 7SK snRNA (c), and U6 snRNA (d) in total

RNA extracted from ovaries of females of the indicated genotypes, for three biological replicates. Note

that *bin3* mRNA was below the level of detection in *bin3$^{2-7}$/Df* and *bin3$^{2-7}$/Df; CycT^{j11B2}/+* ovaries for 2 of

the three biological replicates. RNA levels were normalized to *rp49* mRNA and made relative to wt

using CFX Maestro 2.0 (version 5.2.008.0222, Bio-Rad). Error bars represent the standard error of the

mean. Statistical significance was calculated by unpaired t-test using CFX Maestro 2.0 (version

5.2.008.0222, Bio-Rad). *P* values of <.05 (*), <0.001 (***), <.0001 (****), and not significant (ns) are

indicated. **(e)** Western blot for CycT (Nguyen *et al.* 2012) and Actin in extracts prepared from ovaries of

females of the indicated genotypes. The *P{lacW}j11B2^{j11B2}* insertion (Spradling *et al.* 1999) into one of the

two different *CycT* 5' -UTRs reduces expression of the large isoform of CycT.

Figure 2. Bin3 is ubiquitously expressed in the ovary and is required for egg-laying, but is not

required to establish ovarioles and does not function in germline to promote egg-laying. (a) Scatter

plot showing the egg-laying rates of females of the indicated genotypes for each of three biological

replicates. Error bars represent the standard deviation. Statistical significance compared to *bin3²⁻⁷/Df*

was calculated by nested one-way ANOVA with Dunnett's Correction for multiple comparisons, using

Prism (version 9.4.1, GraphPad). *P* values of <.001 (***) and <.0001 (****) are indicated. **(b)** Cartoon

depicting a single ovariole, of which 16-20 make up a single adult ovary. Germline cells are indicated in

pink, nurse cell nuclei (arrow) are indicated in purple, oocytes are indicated in blue, and somatic cells

surrounding the germline and oocytes are indicated in purple. **(c to d')** Immunofluorescence staining of

adult ovaries for 3xHA-Bin3 (green), Vasa (a cytoplasmic marker for germline cells; blue), and Hoechst

staining of DNA (red). Scale bars = 100 μm. A Trojan-GAL4 phase 0 insertion in *bin3* (*bin3^{TG4.0}*) was used

to drive expression of a *UAS-3xHA-bin3* transgene in the pattern of *bin3* expression (*bin3^{TG4.0} > UAS-bin3-*

3xHA). 3xHA-Bin3 localizes to the nuclei of all germline and somatic cells (C-C'). No HA staining was

detected in *bin3^{TG4.0}* control ovaries (D-D'), demonstrating the specificity of HA staining in *bin3^{TG4.0} >*

UAS-3xHA-bin3 ovaries. **(e-f')** Single-molecule fluorescence *in situ* hybridization (smFISH) of adult

ovaries for 7SK (green) and Hoechst staining of DNA (purple). Scale bars = 100 μm. 7SK localizes to the

nuclei of all germline and somatic cells in wt ovaries (E-E'). 7SK staining is absent in *bin3²⁻⁷/Df* ovaries

(F-F'), which do not accumulate 7SK (see Fig. 1c). **(g)** Scatter plot showing the mean number of

ovarioles per ovary per female of the indicated genotypes for each of two biological replicates. Error

bars represent the standard deviation. Statistical significance compared to *bin3²⁻⁷/Df* was calculated by

nested one-way ANOVA with Dunnett's Correction for multiple comparisons, using Prism (version

9.4.1, GraphPad). *P* values of not significant (ns) are indicated. **(h)** Scatter plot showing the absolute

value of the difference between ovariole number within a pair of ovaries per female of the indicated

genotypes for each of two biological replicates. Error bars represent the standard deviation. Statistical

significance compared to *bin3²⁻⁷/Df* was calculated by nested one-way ANOVA with Dunnett's

Correction for multiple comparisons, using Prism (version 9.4.1, GraphPad). *P* values of not significant

(ns) are indicated. **(i)** Scatter plot showing the egg-laying rates of females of the indicated genotypes

for each of three biological replicates. Error bars represent the standard deviation. Statistical

significance compared to *bin3²⁻⁷/Df; nanos-GAL4::VP16/+* was calculated by nested one-way ANOVA

with Dunnett's Correction for multiple comparisons, using Prism (version 9.4.1, GraphPad). *P* values of

<.05 (*), <.001 (***) and not significant (ns) are indicated.

Figure 3. Bin3 is required for neuromuscular function. (a) Scatter plot showing the percent of flies of

the indicated genotypes climbing 8 cm in 12" in each of 10 trials, for each of three biological replicates.

Error bars represent the standard deviation. Statistical significance compared to *bin3²⁻⁷/Df* was

calculated by nested one-way ANOVA with Dunnett's Correction for multiple comparisons, using Prism

(version 9.4.1, GraphPad). *P* value of <.001 (***) is indicated. **(b-e)** Images of females of the indicated

genotypes, exhibiting either a normal resting wing posture or a *held-out wings* phenotype. **(f)** Scatter

plot showing the percent of flies of the indicated genotypes climbing 8 cm in 12" in each of 4 trials, for

each of three biological replicates. Error bars represent the standard deviation. Statistical significance

was calculated by nested t-test, using Prism (version 9.4.1, GraphPad). *P* value of not significant (ns) is

indicated. **(g-h)** Images of females of the indicated genotypes, exhibiting either a normal resting wing

posture or a *held-out wings* phenotype.

Figure 4. Molecular characterization of the *bin3^{TG4.0}* Trojan GAL4 allele and Bin3 protein mutants.

(a) Bin3 modeled onto the crystal structure of 309 amino acids of the MEPCE active site (Yang *et al.*

2019). Y421 in MEPCE (black) is juxtaposed between the 7SK 5' gamma phosphate (orange and

salmon) and *S*-adenosyl-*L*-homocysteine (SAH, blue and gold), and is essential for MEPCE catalytic

activity *in vitro* (Yang *et al.* 2019). Y795 in Bin3 (red) aligns with Y421 in MEPCE (RMSD = 0.000). **(b)**

Same as in (a), but with the Y421A (MEPCE, black) and Y795A (Bin3, red) tyrosine-to-alanine mutations

introduced. Y421A completely eliminated MEPCE catalytic activity *in vitro* (Yang *et al.* 2019). **(c)** Partial

alignment of Bin3 orthologs from *D. melanogaster*, *H. sapiens* (MEPCE), *D. rerio* (BCDIN3), and *S. pombe*

(Bmc1). Conserved residues are in boldface. **(d)** Graphic depicting the *bin3* locus, as in **Fig. 1a**. *bin3*$^{TG4.0}$

refers to a transgenic insertion of a Trojan GAL4 phase 0 (TG4.0) construct into the second intron in the

bin3 ORF (Lee *et al.* 2018). A splice acceptor (SA) in the insertion causes the TG4.0 cassette to be spliced

into the *bin3* mRNA; transcription is terminated at the end of the cassette by a synthetic poly(A)

sequence. When the chimeric *bin3*$^{TG4.0}$ mRNA is translated, T2A, a viral peptide sequence, disrupts

translation and causes the release of the nonfunctional Bin3:T2A chimeric peptide from the ribosome;

subsequently, the ribosome resumes translation of full-length GAL4 protein. *Df* refers to deficiency

Df(2R)BSC313, which deletes ~625 kb of chromosome 2R, including promoters P2 and P1, and all *bin3*

coding sequence (Cook *et al.* 2012). *bin3*$^{TG4.0}$ and *Df* in *trans* produce *bin3* mutant flies that express *GAL4*

in the pattern of *bin3* expression. GAL4 then binds to UAS and drives expression of *UAS-bin3* transgenes

(see **Fig. 4h-j**). **(e-g)** Bar graphs showing the average amount of *bin3* mRNA (e), 7SK snRNA (f), and U6

snRNA (g) in total RNA extracted from ovaries of females of the indicated genotypes, for three

biological replicates. RNA levels were normalized to *rp49* mRNA and made relative to wt using CFX

Maestro 2.0 (version 5.2.008.0222, Bio-Rad). Error bars represent the standard error of the mean.

Statistical significance was calculated by unpaired t-test using CFX Maestro 2.0 (version 5.2.008.0222,

Bio-Rad). *P* values of <.05 (*), <0.01 (**), <.0001 (****), and not significant (ns) are indicated. **(h-j)**

Graphic depicting *UAS-DSCP-3xHA-bin3* transgenes. UAS-DSCP (grey) is a GAL4-responsive promoter

that is active in both germline and somatic cells (Pfeiffer *et al.* 2008). *bin3* cDNAs encode Bin3 proteins

that are either wt (h; sky blue), catalytically-dead (I; Y795A, pink), or deleted of the MSM (j; ΔMSM,

maroon). Each construct is tagged at the C-terminus with 3xHA that is codon-optimized for *Drosophila*

(green). When these transgenes are independently crossed into the *bin3*$^{TG4.0}$/*Df* background (see **Fig.**

4d), flies will express only 3xHA-Bin3, 3xHA-Bin3^{Y795A}, or 3xHA-Bin3^{ΔMSM} proteins, respectively. **(k)** Bar

graph showing the average amount of *bin3* mRNA in total RNA extracted from ovaries of females of the

indicated genotypes, for three biological replicates. RNA levels were normalized to *rp49* mRNA and

made relative to *bin3*$^{TG4.0}$*/Df; UAS-bin3/*+ using CFX Maestro 2.0 (version 5.2.008.0222, Bio-Rad). Error

bars represent the standard error of the mean. Statistical significance was calculated by unpaired t-test

using CFX Maestro 2.0 software (version 5.2.008.0222, Bio-Rad). *P* value of not significant (ns) is

indicated. All three *UAS-bin3* transgenes are expressed to similar levels. **(i)** Representative Western blot

for 3xHA-Bin3 proteins and Actin in extracts prepared from ovaries of females of the indicated

genotypes. The relative amounts of 3xHA-Bin3^{Y795A} and 3xHA-Bin3$^{\Delta MSM}$ relative to 3xHA-Bin3 (wt) ± the

standard deviation from two biological replicates are indicated below. Statistical significance was

calculated by ordinary one-way ANOVA with Dunnett's Correction for multiple comparisons with a

single pooled variance, using Prism (version 9.4.1, GraphPad). *P* values of <0.05 (*) and not significant

(ns) are indicated. All three transgenes produce similar amounts of 3xHA-Bin3 proteins.

Figure 5. Neither catalytic activity nor the metazoan-specific motif are required for 7SK or U6

binding and stability. (a and **c)** Bar graphs showing the average amount of 7SK (a) and U6 (c) RNAs in

total RNA extracted from ovaries of females of the indicated genotypes, for three biological replicates.

RNA levels were normalized to *rp49* mRNA and made relative to *bin3*$^{TG4.0}$*/Df; UAS-bin3/*+ using CFX

Maestro 2.0 (version 5.2.008.0222, Bio-Rad). Error bars represent the standard error of the mean.

Statistical significance was calculated by unpaired t-test using CFX Maestro 2.0 (version 5.2.008.0222,

Bio-Rad). *P* value of not significant (ns) is indicated. **(b** and **d)** Bar graphs showing the average amount

of 7SK (b) and U6 (d) RNA bound by 3xHA-Bin3 proteins immunoprecipitated with anti-HA magnetic

beads from extracts prepared from ovaries of females of the indicated genotypes, for three biological

replicates. The ratio of immunoprecipitated RNA to input RNA, from flies expressing 3xHA-Bin3

proteins relative to flies not expressing 3xHA-Bin3 proteins (*bin3*$^{TG4.0}$/+), was calculated using CFX

Maestro 2.0 (version 5.2.008.0222, Bio-Rad). Error bars represent the standard error of the mean.

Statistical significance compared to *bin3*[TG4.0]/+ (black asterisks) or *bin3*[TG4.0]/*Df; UAS-bin3*/+ (sky blue

asterisks) was calculated by one-way ANOVA with Benjamini-Hochberg correction for false discovery

rate, using CFX Maestro 2.0 (version 5.2.008.0222, Bio-Rad). *P* values of <0.05 (*), <.01 (**), <.0001

(****), and not significant (ns) are indicated.

Figure 6. Phenotypic characterization of Bin3[Y795A] **and Bin3**[ΔMSM] **mutant proteins *in vivo*. (a)** Scatter

plot showing egg-laying rates of females of the indicated genotypes for each of three biological

replicates. Error bars represent the standard deviation. Statistical significance compared to *bin3*[TG4.0]/*Df*

(salmon asterisks) or *bin3*[TG4.0]/*Df; UAS-bin3*/+ (sky blue asterisks) was calculated by nested one-way

ANOVA with Dunnett's Correction for multiple comparisons, using Prism (version 9.4.1, GraphPad). *P*

values of <.01 (**), <0.001 (***), <.0001 (****), and not significant (ns) are indicated. **(b)** Scatter plot

showing the percent of flies of the indicated genotypes climbing 8 cm in 12" in each of 10 trials, for each

of three biological replicates. Error bars represent the standard deviation. Statistical significance

compared to *bin3*[TG4.0]/*Df* (salmon asterisks) or *bin3*[TG4.0]/*Df; UAS-bin3*/+ (sky blue asterisks) was calculated

by nested one-way ANOVA with Dunnett's Correction for multiple comparisons, using Prism (version

9.4.1, GraphPad). *P* values of <.0001 (****) and not significant (ns) are indicated. **(c-h)** Images of

females of the indicated genotypes, exhibiting either a normal resting wing posture or a *held-out wings*

phenotype.

Figure 7. Summary of Bin3[wt]**, Bin3**[Y795A]**, and Bin3**[ΔMSM] **protein functions. (a)** Bin3[wt] binds, caps, and

stabilizes 7SK, which allows for P-TEFb to be sequestered and repressed, which prevents RNAPII pause-

release and promotes egg-laying, climbing, and normal resting wing posture. **(b)** Bin3[Y795A] is able to

bind and stabilize 7SK despite being catalytically-dead; consequently, Bin3[Y795A] also promotes egg-

laying, climbing, and normal resting wing posture. **(c)** Bin3[ΔMSM] binds, caps, and stabilizes 7SK, and

represses P-TEFb; this is sufficient to promote normal egg-laying and climbing, but is not sufficient to

promote normal wing posture. This suggests the MSM confers a 7SK-independent, tissue-specific

function to Bin3.

1000

Figure 8. Model for the role of Bin3 in *Drosophila* development. (a) Bin3 binds, caps, and stabilizes

7SK; however, methyltransferase activity is not required for 7SK stability *in vivo*. Stabilized 7SK

assembles a snRNP that represses P-TEFb, which prevents RNAPII transcription elongation. **(b)** The

Bin3 MSM confers a 7SK-independent, tissue-specific function to Bin3. The MSM might mediate an

interaction with a tissue-specific factor, or this interaction might only be important in specific tissues.

Bin3 and this factor then repress transcription elongation by a mechanism that is likely downstream of

P-TEFb.

SUPPLEMENTAL TABLES

Table S1. Public fly stocks used in this study. This table lists the publicly available fly stocks used in

this study, including their genotypes as they appear in the paper, their detailed genotypes, the stock

center from which they were procured, and the stock number.

1012

1013 **Table S2. Fly stocks produced for this study.** This table lists the fly stocks produced for this study,

1014 including their genotypes as they appear in the paper, their detailed genotypes, and a detailed

1015 description of how each fly stock was produced.

1016

Table S3. Flies used for experiments. This table lists the flies used in each figure, including their

genotypes as they appear in the paper, their detailed genotypes, the genotypes of the female and male

flies crossed to produce flies used for experiments, and the figure(s) in which each fly is used.

REFERENCES

Abbaszadeh E. K., and E. R. Gavis, 2016 Fixed and live visualization of RNAs in Drosophila oocytes and

embryos. Methods 98: 34–41. https://doi.org/10.1016/j.ymeth.2016.01.018

Abuhashem A., V. Garg, and A.-K. Hadjantonakis, 2022 RNA polymerase II pausing in development:

orchestrating transcription. Open Biol 12: 210220. https://doi.org/10.1098/rsob.210220

Adelman K., and J. T. Lis, 2012 Promoter-proximal pausing of RNA polymerase II: emerging roles in

metazoans. Nat Rev Genet 13: 720–731. https://doi.org/10.1038/nrg3293

Akhtar J., Y. Renaud, S. Albrecht, Y. Ghavi-Helm, J.-Y. Roignant, *et al.*, 2021 m6A RNA methylation

regulates promoter- proximal pausing of RNA polymerase II. Mol Cell 81: 3356-3367.e6.

https://doi.org/10.1016/j.molcel.2021.06.023

Ayala-Camargo A., A. M. Anderson, M. Amoyel, A. B. Rodrigues, M. S. Flaherty, *et al.*, 2013 JAK/STAT

signaling is required for hinge growth and patterning in the Drosophila wing disc. Dev Biol 382: 413--

426. https://doi.org/10.1016/j.ydbio.2013.08.016

Baehrecke E. H., 1997 who encodes a KH RNA binding protein that functions in muscle development.

Dev Camb Engl 124: 1323–32.

Barboric M., T. Lenasi, H. Chen, E. B. Johansen, S. Guo, *et al.*, 2009 7SK snRNP/P-TEFb couples

transcription elongation with alternative splicing and is essential for vertebrate development. Proc

National Acad Sci 106: 7798–7803. https://doi.org/10.1073/pnas.0903188106

Barrandon C., F. Bonnet, V. T. Nguyen, V. Labas, and O. Bensaude, 2007 The Transcription-Dependent

Dissociation of P-TEFb-HEXIM1-7SK RNA Relies upon Formation of hnRNP-7SK RNA Complexes.

Mol Cell Biol 27: 6996–7006. https://doi.org/10.1128/mcb.00975-07

Blazek D., M. Barboric, J. Kohoutek, I. Oven, and B. M. Peterlin, 2005 Oligomerization of HEXIM1 via

7SK snRNA and coiled-coil region directs the inhibition of P-TEFb. Nucleic Acids Res 33: 7000–7010.

https://doi.org/10.1093/nar/gki997

Brand A. H., and N. Perrimon, 1993 Targeted gene expression as a means of altering cell fates and

generating dominant phenotypes. Development 118: 401–415.

https://doi.org/10.1242/dev.118.2.401

Brogie J. E., and D. H. Price, 2017 Reconstitution of a functional 7SK snRNP. Nucleic Acids Res 45: 6864–

6880. https://doi.org/10.1093/nar/gkx262

Brookes E., and A. Pombo, 2009 Modifications of RNA polymerase II are pivotal in regulating gene

expression states. Embo Rep 10: 1213–1219. https://doi.org/10.1038/embor.2009.221

Bugai A., A. J. C. Quaresma, C. C. Friedel, T. Lenasi, R. Düster, *et al.*, 2019 P-TEFb Activation by RBM7

Shapes a Pro-survival Transcriptional Response to Genotoxic Stress. Mol Cell 74: 254-267.e10.

https://doi.org/10.1016/j.molcel.2019.01.033

Cabrera G. R., D. Godt, P. Fang, J. Couderc, and F. A. Laski, 2002 Expression pattern of Gal4 enhancer

trap insertions into the bric à brac locus generated by P element replacement. Genesis 34: 62–65.

https://doi.org/10.1002/gene.10115

Callus B. A., and B. Mathey-Prevot, 2002 SOCS36E, a novel Drosophila SOCS protein, suppresses

JAK/STAT and EGF-R signalling in the imaginal wing disc. Oncogene 21: 4812--4821.

https://doi.org/10.1038/sj.onc.1205618

Chan S.-K., and G. Struhl, 1997 Sequence-specific RNA binding by Bicoid. Nature 388: 634–634.

https://doi.org/10.1038/41692

Chen F. X., E. R. Smith, and A. Shilatifard, 2018 Born to run: control of transcription elongation by RNA

polymerase II. Nat Rev Mol Cell Bio 19: 464–478. https://doi.org/10.1038/s41580-018-0010-5

Cheng B., and D. H. Price, 2007 Properties of RNA Polymerase II Elongation Complexes Before and

After the P-TEFb-mediated Transition into Productive Elongation*. J Biol Chem 282: 21901–21912.

https://doi.org/10.1074/jbc.m702936200

Chiba K., J. Yamamoto, Y. Yamaguchi, and H. Handa, 2010 Promoter-proximal pausing and its release:

Molecular mechanisms and physiological functions. Exp Cell Res 316: 2723–2730.

https://doi.org/10.1016/j.yexcr.2010.05.036

Cho P. F., F. Poulin, Y. A. Cho-Park, I. B. Cho-Park, J. D. Chicoine, *et al.*, 2005 A New Paradigm for

Translational Control: Inhibition via 5 -3 mRNA Tethering by Bicoid and the eIF4E Cognate 4EHP.

Cell 121: 411–423. https://doi.org/10.1016/j.cell.2005.02.024

Cook R. K., S. J. Christensen, J. A. Deal, R. A. Coburn, M. E. Deal, *et al.*, 2012 The generation of

chromosomal deletions to provide extensive coverage and subdivision of the Drosophila

melanogaster genome. Genome Biol 13: R21. https://doi.org/10.1186/gb-2012-13-3-r21

Core L., and K. Adelman, 2019 Promoter-proximal pausing of RNA polymerase II: a nexus of gene

regulation. Gene Dev 33: 960–982. https://doi.org/10.1101/gad.325142.119

Cosgrove M. S., Y. Ding, W. A. Rennie, M. J. Lane, and S. D. Hanes, 2012 The Bin3 RNA

methyltransferase targets 7SK RNA to control transcription and translation. Wiley Interdiscip Rev

Rna 3: 633–647. https://doi.org/10.1002/wrna.1123

Deady L. D., and J. Sun, 2015 A Follicle Rupture Assay Reveals an Essential Role for Follicular

Adrenergic Signaling in Drosophila Ovulation. Plos Genet 11: e1005604.

https://doi.org/10.1371/journal.pgen.1005604

Diao F., and B. H. White, 2012 A Novel Approach for Directing Transgene Expression in Drosophila :

T2A-Gal4 In-Frame Fusion. Genetics 190: 1139–1144. https://doi.org/10.1534/genetics.111.136291

Diao F., H. Ironfield, H. Luan, F. Diao, W. C. Shropshire, *et al.*, 2015 Plug-and-Play Genetic Access to

Drosophila Cell Types using Exchangeable Exon Cassettes. Cell Reports 10: 1410–1421.

https://doi.org/10.1016/j.celrep.2015.01.059

Dollinger R., and D. S. Gilmour, 2021 Regulation of Promoter Proximal Pausing of RNA Polymerase II in

Metazoans. J Mol Biol 433: 166897. https://doi.org/10.1016/j.jmb.2021.166897

Doren M. V., A. L. Williamson, and R. Lehmann, 1998 Regulation of zygotic gene expression in

Drosophila primordial germ cells. Curr Biol 8: 243–246. https://doi.org/10.1016/s0960-

9822(98)70091-0

Dubnau J., and G. Struhl, 1996 RNA recognition and translational regulation by a homeodomain

protein. Nature 379: 694–699. https://doi.org/10.1038/379694a0

Erdős G., and Z. Dosztányi, 2020 Analyzing Protein Disorder with IUPred2A. Curr Protoc Bioinform 70:

e99. https://doi.org/10.1002/cpbi.99

Erdős G., M. Pajkos, and Z. Dosztányi, 2021 IUPred3: prediction of protein disorder enhanced with

unambiguous experimental annotation and visualization of evolutionary conservation. Nucleic Acids

Res 49: gkab408-. https://doi.org/10.1093/nar/gkab408

Fujinaga K., F. Huang, and B. M. Peterlin, 2023 P-TEFb: The master regulator of transcription

elongation. Mol Cell 83: 393–403. https://doi.org/10.1016/j.molcel.2022.12.006

Gabler F., S. Nam, S. Till, M. Mirdita, M. Steinegger, *et al.*, 2020 Protein Sequence Analysis Using the

MPI Bioinformatics Toolkit. Curr Protoc Bioinform 72: e108. https://doi.org/10.1002/cpbi.108

Gaertner B., and J. Zeitlinger, 2014 RNA polymerase II pausing during development. Development 141:

1179–1183. https://doi.org/10.1242/dev.088492

Garrett E. C., A. M. Bielawski, E. Ruchti, L. M. Sherer, I. Waghmare, *et al.*, 2023 The matricellular protein

Drosophila CCN is required for synaptic transmission and female fertility. Genetics.

https://doi.org/10.1093/genetics/iyac190

Gilmour D. S., 2009 Promoter proximal pausing on genes in metazoans. Chromosoma 118: 1–10.

https://doi.org/10.1007/s00412-008-0182-4

Gonzalez M. N., D. Blears, and J. Q. Svejstrup, 2021 Causes and consequences of RNA polymerase II

stalling during transcript elongation. Nat Rev Mol Cell Bio 22: 3–21. https://doi.org/10.1038/s41580-

020-00308-8

Gupta S., R. K. Busch, R. Singh, and R. Reddy, 1990a Characterization of U6 small nuclear RNA cap-

specific antibodies. Identification of gamma-monomethyl-GTP cap structure in 7SK and several

other human small RNAs. J Biol Chem 265: 19137–19142. https://doi.org/10.1016/s0021-

9258(17)30635-x

Gupta S., R. K. Busch, R. Singh, and R. Reddy, 1990b Characterization of U6 small nuclear RNA cap-

specific antibodies. Identification of gamma-monomethyl-GTP cap structure in 7SK and several

other human small RNAs. J Biological Chem 265: 19137–42.

Haaland R. E., C. H. Herrmann, and A. P. Rice, 2005 siRNA depletion of 7SK snRNA induces apoptosis

but does not affect expression of the HIV-1 LTR or P-TEFb-dependent cellular genes. J Cell Physiol

205: 463–470. https://doi.org/10.1002/jcp.20528

Harrison D. A., P. E. McCoon, R. Binari, M. Gilman, and N. Perrimon, 1998 Drosophila unpaired encodes

a secreted protein that activates the JAK signaling pathway. Gene Dev 12: 3252--3263.

https://doi.org/10.1101/gad.12.20.3252

Hatini V., E. Kula-Eversole, D. Nusinow, and S. J. D. Signore, 2013 Essential roles for stat92E in

expanding and patterning the proximodistal axis of the Drosophila wing imaginal disc. Dev Biol 378:

38--50. https://doi.org/10.1016/j.ydbio.2013.02.016

He N., N. S. Jahchan, E. Hong, Q. Li, M. A. Bayfield, *et al.*, 2008 A La-Related Protein Modulates 7SK

snRNP Integrity to Suppress P-TEFb-Dependent Transcriptional Elongation and Tumorigenesis. Mol

Cell 29: 588–599. https://doi.org/10.1016/j.molcel.2008.01.003

Herreweghe E. V., S. Egloff, I. Goiffon, B. E. Jády, C. Froment, *et al.*, 2007 Dynamic remodelling of

human 7SK snRNP controls the nuclear level of active P-TEFb. Embo J 26: 3570–3580.

https://doi.org/10.1038/sj.emboj.7601783

Hildebrand A., M. Remmert, A. Biegert, and J. Söding, 2009 Fast and accurate automatic structure

prediction with HHpred. Proteins Struct Funct Bioinform 77: 128–132.

https://doi.org/10.1002/prot.22499

Irizarry J., and A. Stathopoulos, 2015 FGF signaling supports Drosophila fertility by regulating

development of ovarian muscle tissues. Dev Biol 404: 1–13.

https://doi.org/10.1016/j.ydbio.2015.04.023

Jeronimo C., D. Forget, A. Bouchard, Q. Li, G. Chua, *et al.*, 2007 Systematic Analysis of the Protein

Interaction Network for the Human Transcription Machinery Reveals the Identity of the 7SK Capping

Enzyme. Mol Cell 27: 262–274. https://doi.org/10.1016/j.molcel.2007.06.027

Johnstone K., R. E. Wells, D. Strutt, and M. P. Zeidler, 2013 Localised JAK/STAT Pathway Activation Is

Required for Drosophila Wing Hinge Development. Plos One 8.

https://doi.org/10.1371/journal.pone.0065076

Jonkers I., and J. T. Lis, 2015 Getting up to speed with transcription elongation by RNA polymerase II.

Nat Rev Mol Cell Bio 16: 167–177. https://doi.org/10.1038/nrm3953

Kairamkonda S., and U. Nongthomba, 2014 Beadex Function in the Motor Neurons Is Essential for

Female Reproduction in Drosophila melanogaster. Plos One 9: e113003.

https://doi.org/10.1371/journal.pone.0113003

Kan L., A. V. Grozhik, J. Vedanayagam, D. P. Patil, N. Pang, *et al.*, 2017 The m6A pathway facilitates sex

determination in Drosophila. Nat Commun 8: 1--16. https://doi.org/10.1038/ncomms15737

Krueger B. J., C. Jeronimo, B. B. Roy, A. Bouchard, C. Barrandon, *et al.*, 2008 LARP7 is a stable

component of the 7SK snRNP while P-TEFb, HEXIM1 and hnRNP A1 are reversibly associated.

Nucleic Acids Res 36: 2219–2229. https://doi.org/10.1093/nar/gkn061

Krueger B. J., K. Varzavand, J. J. Cooper, and D. H. Price, 2010 The Mechanism of Release of P-TEFb

and HEXIM1 from the 7SK snRNP by Viral and Cellular Activators Includes a Conformational Change

in 7SK. Plos One 5: e12335. https://doi.org/10.1371/journal.pone.0012335

Lee P.-T., J. Zirin, O. Kanca, W.-W. Lin, K. L. Schulze, *et al.*, 2018 A gene-specific T2A-GAL4 library for

Drosophila. Elife 7: e35574. https://doi.org/10.7554/elife.35574

Lee S., H. Liu, R. Hill, C. Chen, X. Hong, *et al.*, 2020 JMJD6 cleaves MePCE to release positive

transcription elongation factor b (P-TEFb) in higher eukaryotes. Elife 9: e53930.

https://doi.org/10.7554/elife.53930

Leger A., P. P. Amaral, L. Pandolfini, C. Capitanchik, F. Capraro, *et al.*, 2021 RNA modifications

detection by comparative Nanopore direct RNA sequencing. Nat Commun 12: 7198.

https://doi.org/10.1038/s41467-021-27393-3

Lemieux B., M. Blanchette, A. Monette, A. J. Mouland, R. J. Wellinger, *et al.*, 2015 A Function for the

hnRNP A1/A2 Proteins in Transcription Elongation. Plos One 10: e0126654.

https://doi.org/10.1371/journal.pone.0126654

Lence T., J. Akhtar, M. Bayer, K. Schmid, L. Spindler, *et al.*, 2016 m6A modulates neuronal functions

and sex determination in Drosophila. Nature 540: 242–247. https://doi.org/10.1038/nature20568

Lence T., C. Paolantoni, L. Worpenberg, and J.-Y. Roignant, 2018 Mechanistic insights into m6A RNA

enzymes. Biochimica Et Biophysica Acta Gene Regul Mech.

https://doi.org/10.1016/j.bbagrm.2018.10.014

Levine M., 2011 Paused RNA Polymerase II as a Developmental Checkpoint. Cell 145: 502–511.

https://doi.org/10.1016/j.cell.2011.04.021

Li Q., J. P. Price, S. A. Byers, D. Cheng, J. Peng, *et al.*, 2005 Analysis of the Large Inactive P-TEFb

Complex Indicates That It Contains One 7SK Molecule, a Dimer of HEXIM1 or HEXIM2, and Two P-

TEFb Molecules Containing Cdk9 Phosphorylated at Threonine 186*. J Biol Chem 280: 28819–

28826. https://doi.org/10.1074/jbc.m502712200

Li J., and D. S. Gilmour, 2011 Promoter proximal pausing and the control of gene expression. Curr Opin

Genet Dev 21: 231–235. https://doi.org/10.1016/j.gde.2011.01.010

Liao S., and D. R. Nässel, 2020 Drosophila Insulin-Like Peptide 8 (DILP8) in Ovarian Follicle Cells

Regulates Ovulation and Metabolism. Front Endocrinol 11: 461.

https://doi.org/10.3389/fendo.2020.00461

Liu X., W. L. Kraus, and X. Bai, 2015 Ready, pause, go: regulation of RNA polymerase II pausing and

release by cellular signaling pathways. Trends Biochem Sci 40: 516–525.

https://doi.org/10.1016/j.tibs.2015.07.003

Lobell A. S., R. R. Kaspari, Y. L. S. Negron, and S. T. Harbison, 2017 The Genetic Architecture of

Ovariole Number in Drosophila melanogaster: Genes with Major, Quantitative, and Pleiotropic

Effects. G3 Genes Genomes Genetics 7: 2391–2403. https://doi.org/10.1534/g3.117.042390

Maimon I., and L. Gilboa, 2011 Dissection and Staining of Drosophila Larval Ovaries. J Vis

Exp. https://doi.org/10.3791/2537

Manjila S., and G. Hasan, 2018 Flight and Climbing Assay for Assessing Motor Functions in Drosophila.

Bio-protocol 8. https://doi.org/10.21769/bioprotoc.2742

Margaritis T., and F. C. P. Holstege, 2008 Poised RNA Polymerase II Gives Pause for Thought. Cell 133:

581–584. https://doi.org/10.1016/j.cell.2008.04.027

Markert A., M. Grimm, J. Martinez, J. Wiesner, A. Meyerhans, *et al.*, 2008 The La-related protein LARP7

is a component of the 7SK ribonucleoprotein and affects transcription of cellular and viral

polymerase II genes. Embo Rep 9: 569–575. https://doi.org/10.1038/embor.2008.72

Marshall N. F., and D. H. Price, 1995 Purification of P-TEFb, a Transcription Factor Required for the

Transition into Productive Elongation (∗). J Biol Chem 270: 12335–12338.

https://doi.org/10.1074/jbc.270.21.12335

Marshall N. F., J. Peng, Z. Xie, and D. H. Price, 1996 Control of RNA Polymerase II Elongation Potential

by a Novel Carboxyl-terminal Domain Kinase*. J Biol Chem 271: 27176–27183.

https://doi.org/10.1074/jbc.271.43.27176

Mayer A., H. M. Landry, and L. S. Churchman, 2017 Pause & go: from the discovery of RNA polymerase

pausing to its functional implications. Curr Opin Cell Biol 46: 72–80.

https://doi.org/10.1016/j.ceb.2017.03.002

Meier A., and J. Söding, 2015 Automatic Prediction of Protein 3D Structures by Probabilistic Multi-

template Homology Modeling. Plos Comput Biol 11: e1004343.

https://doi.org/10.1371/journal.pcbi.1004343

Mészáros B., G. Erdős, and Z. Dosztányi, 2018 IUPred2A: context-dependent prediction of protein

disorder as a function of redox state and protein binding. Nucleic Acids Res 46: gky384-.

https://doi.org/10.1093/nar/gky384

Michels A. A., A. Fraldi, Q. Li, T. E. Adamson, F. Bonnet, *et al.*, 2004 Binding of the 7SK snRNA turns the

HEXIM1 protein into a P-TEFb (CDK9/cyclin T) inhibitor. Embo J 23: 2608–2619.

https://doi.org/10.1038/sj.emboj.7600275

Muniz L., S. Egloff, and T. Kiss, 2013 RNA elements directing in vivo assembly of the 7SK/MePCE/Larp7

transcriptional regulatory snRNP. Nucleic Acids Res 41: 4686–4698.

https://doi.org/10.1093/nar/gkt159

Murphy S., M. Tripodi, and M. Melli, 1986 A sequence upstream from the coding region is required for

the transcription of the 7SK RNA genes. Nucleic Acids Res 14: 9243–9260.

https://doi.org/10.1093/nar/14.23.9243

Murphy S., C. D. Liegro, and M. Melli, 1987 The in vitro transcription of the 7SK RNA gene by RNA

polymerase III is dependent only on the presence of an upstream promoter. Cell 51: 81–87.

https://doi.org/10.1016/0092-8674(87)90012-2

Murphy S., A. Pierani, C. Scheidereit, M. Melli, and R. G. Roeder, 1989 Purified octamer binding

transcription factors stimulate RNA polymerase III-mediated transcription of the 7SK RNA gene. Cell

59: 1071–1080. https://doi.org/10.1016/0092-8674(89)90763-0

Nechaev S., and K. Adelman, 2011 Pol II waiting in the starting gates: Regulating the transition from

transcription initiation into productive elongation. Biochimica Et Biophysica Acta Bba - Gene Regul

Mech 1809: 34–45. https://doi.org/10.1016/j.bbagrm.2010.11.001

Nguyen V. T., T. Kiss, A. A. Michels, and O. Bensaude, 2001 7SK small nuclear RNA binds to and inhibits

the activity of CDK9/cyclin T complexes. Nature 414: 322–325. https://doi.org/10.1038/35104581

Nguyen D., B. J. Krueger, S. C. Sedore, J. E. Brogie, J. T. Rogers, *et al.*, 2012 The Drosophila 7SK snRNP

and the essential role of dHEXIM in development. Nucleic Acids Res 40: 5283–5297.

https://doi.org/10.1093/nar/gks191

Ni Z., B. E. Schwartz, J. Werner, J.-R. Suarez, and J. T. Lis, 2004 Coordination of Transcription, RNA

Processing, and Surveillance by P-TEFb Kinase on Heat Shock Genes. Mol Cell 13: 55–65.

https://doi.org/10.1016/s1097-2765(03)00526-4

Ni Z., A. Saunders, N. J. Fuda, J. Yao, J.-R. Suarez, *et al.*, 2008 P-TEFb Is Critical for the Maturation of

RNA Polymerase II into Productive Elongation In Vivo. Mol Cell Biol 28: 1161–1170.

https://doi.org/10.1128/mcb.01859-07

Ni J. Q., R. Zhou, B. Czech, L. P. Liu, L. Holderbaum, *et al.*, 2011 A genome-scale shRNA resource for

transgenic RNAi in Drosophila. Nat Methods 8: 405--407. https://doi.org/10.1038/nmeth.1592

Niessing D., N. Dostatni, H. Jäckle, and R. Rivera-Pomar, 1999 Sequence interval within the PEST motif

of Bicoid is important for translational repression of caudal mRNA in the anterior region of the

Drosophila embryo. Embo J 18: 1966–1973. https://doi.org/10.1093/emboj/18.7.1966

Niessing D., W. Driever, F. Sprenger, H. Taubert, H. Jäckle, *et al.*, 2000 Homeodomain Position 54

Specifies Transcriptional versus Translational Control by Bicoid. Mol Cell 5: 395–401.

https://doi.org/10.1016/s1097-2765(00)80434-7

Niessing D., S. Blanke, and H. Jäckle, 2002 Bicoid associates with the 5 -cap-bound complex of caudal

mRNA and represses translation. Gene Dev 16: 2576–2582. https://doi.org/10.1101/gad.240002

Páez-Moscoso D. J., D. V. Ho, L. Pan, K. Hildebrand, K. L. Jensen, *et al.*, 2022 A putative cap binding

protein and the methyl phosphate capping enzyme Bin3/MePCE function in telomerase biogenesis.

Nat Commun 13: 1067. https://doi.org/10.1038/s41467-022-28545-9

Parks A. L., K. R. Cook, M. Belvin, N. A. Dompe, R. Fawcett, *et al.*, 2004 Systematic generation of high-

resolution deletion coverage of the Drosophila melanogaster genome. Nat Genet 36: 288--292.

https://doi.org/10.1038/ng1312

Peng J., N. F. Marshall, and D. H. Price, 1998 Identification of a Cyclin Subunit Required for the Function

ofDrosophila P-TEFb*. J Biol Chem 273: 13855–13860. https://doi.org/10.1074/jbc.273.22.13855

Perez-Pepe M., A. W. Desotell, H. Li, W. Li, B. Han, *et al.*, 2023 7SK methylation by METTL3 promotes

transcriptional activity. Sci Adv 9. https://doi.org/10.1126/sciadv.ade7500

Peterlin B. M., J. E. Brogie, and D. H. Price, 2011 7SK snRNA: a noncoding RNA that plays a major role in

regulating eukaryotic transcription. Wiley Interdiscip Rev Rna 3: 92–103.

https://doi.org/10.1002/wrna.106

Pfeiffer B. D., A. Jenett, A. S. Hammonds, T.-T. B. Ngo, S. Misra, *et al.*, 2008 Tools for neuroanatomy

and neurogenetics in Drosophila. Proc National Acad Sci 105: 9715–9720.

https://doi.org/10.1073/pnas.0803697105

Porat J., M. E. Baidouri, J. Grigull, J.-M. Deragon, and M. A. Bayfield, 2022 The methyl phosphate

capping enzyme Bmc1/Bin3 is a stable component of the fission yeast telomerase holoenzyme. Nat

Commun 13: 1277. https://doi.org/10.1038/s41467-022-28985-3

Qiao H. H., F. Wang, R. G. Xu, J. Sun, R. Zhu, *et al.*, 2018 An efficient and multiple target transgenic

RNAi technique with low toxicity in Drosophila. Nat Commun 9. https://doi.org/10.1038/s41467-018-

06537-y

Raj A., and S. Tyagi, 2010 Single Molecule Tools: Fluorescence Based Approaches, Part A. Methods

Enzymol 472: 365–386. https://doi.org/10.1016/s0076-6879(10)72004-8

Ramanathan Y., S. M. Rajpara, S. M. Reza, E. Lees, S. Shuman, *et al.*, 2001 Three RNA Polymerase II

Carboxyl-terminal Domain Kinases Display Distinct Substrate Preferences*. J Biol Chem 276: 10913–

10920. https://doi.org/10.1074/jbc.m010975200

Rivera-Pomar R., D. Niessing, U. Schmidt-Ott, W. J. Gehring, and H. Jäckle, 1996 RNA binding and

translational suppression by bicoid. Nature 379: 746–9. https://doi.org/10.1038/379746a0

Ryder E., F. Blows, M. Ashburner, R. Bautista-Llacer, D. Coulson, *et al.*, 2004 The DrosDel collection: A

set of P-element insertions for generating custom chromosomal aberrations in Drosophila

melanogaster. Genetics 167: 797--813. https://doi.org/10.1534/genetics.104.026658

Sahut-Barnola I., D. Godt, F. A. Laski, and J.-L. Couderc, 1995 Drosophila Ovary Morphogenesis:

Analysis of Terminal Filament Formation and Identification of a Gene Required for This Process. Dev

Biol 170: 127–135. https://doi.org/10.1006/dbio.1995.1201

Sahut-Barnola I., B. Dastugue, and J.-L. Couderc, 1996 Terminal filament cell organization in the larval

ovary of Drosophila melanogaster: ultrastructural observations and pattern of divisions. Roux's

Archives Dev Biology 205: 356–363. https://doi.org/10.1007/bf00377215

Šali A., L. Potterton, F. Yuan, H. van Vlijmen, and M. Karplus, 1995 Evaluation of comparative protein

modeling by MODELLER. Proteins Struct Funct Bioinform 23: 318–326.

https://doi.org/10.1002/prot.340230306

Sanhueza M., L. Zechini, T. Gillespie, and G. Pennetta, 2014 Gain-of-function mutations in the ALS8

causative gene VAPB have detrimental effects on neurons and muscles. Biol Open 3: 59--71.

https://doi.org/10.1242/bio.20137070

Schneeberger P. E., T. Bierhals, A. Neu, M. Hempel, and K. Kutsche, 2019 de novo MEPCE nonsense

variant associated with a neurodevelopmental disorder causes disintegration of 7SK snRNP and

enhanced RNA polymerase II activation. Sci Rep-uk 9: 12516. https://doi.org/10.1038/s41598-019-

49032-0

Shelton S. B., N. M. Shah, N. S. Abell, S. K. Devanathan, M. Mercado, *et al.*, 2018 Crosstalk between the

RNA Methylation and Histone-Binding Activities of MePCE Regulates P-TEFb Activation on

Chromatin. Cell Reports 22: 1374–1383. https://doi.org/10.1016/j.celrep.2018.01.028

Shim E. Y., A. K. Walker, Y. Shi, and T. K. Blackwell, 2002 CDK-9/cyclin T (P-TEFb) is required in two

postinitiation pathways for transcription in the C. elegans embryo. Gene Dev 16: 2135–2146.

https://doi.org/10.1101/gad.999002

Shimba S., and R. Reddy, 1994 Purification of human U6 small nuclear RNA capping enzyme. Evidence

for a common capping enzyme for gamma-monomethyl-capped small RNAs. J Biological Chem 269:

12419–23.

Shukla V., F. Habib, A. Kulkarni, and G. S. Ratnaparkhi, 2013 Gene Duplication, Lineage-Specific

Expansion, and Subfunctionalization in the MADF-BESS Family Patterns the Drosophila Wing

Hinge. Genetics 196: 481–496. https://doi.org/10.1534/genetics.113.160531

Shumyatsky G. P., S. V. Tillib, and D. A. Kramerov, 1990 B2 RNA and 7SK RNA, RNA polymerase III

transcripts, have a cap-like structure at their 5 end. Nucleic Acids Res 18: 6347–6351.

https://doi.org/10.1093/nar/18.21.6347

Shumyatsky G., D. Wright, and R. Reddy, 1993 Methylphosphate cap structure increases the stability of

7SK, B2 and U6 small RNAs in Xenopus oocytes. Nucleic Acids Res 21: 4756–4761.

https://doi.org/10.1093/nar/21.20.4756

Shumyatsky G., S. Shimba, and R. Reddy, 1994 Capping signals correspond to the 5' end in four

eukaryotic small RNAs containing gamma-monomethylphosphate cap structure. Gene Expression

4: 29--41.

Singh R., S. Gupta, and R. Reddy, 1990 Capping of Mammalian U6 Small Nuclear RNA In Vitro Is

Directed by a Conserved Stem-Loop and AUAUAC Sequence: Conversion of a Noncapped RNA into

a Capped RNA. Mol Cell Biol 10: 939–946. https://doi.org/10.1128/mcb.10.3.939-946.1990

Singh N., H. Morlock, and S. D. Hanes, 2011 The Bin3 RNA methyltransferase is required for repression

of caudal translation in the Drosophila embryo. Dev Biol 352: 104–115.

https://doi.org/10.1016/j.ydbio.2011.01.017

Söding J., 2005 Protein homology detection by HMM–HMM comparison. Bioinformatics 21: 951–960.

https://doi.org/10.1093/bioinformatics/bti125

Spradling A. C., D. Stern, A. Beaton, E. J. Rhem, T. Laverty, *et al.*, 1999 The Berkeley Drosophila

Genome Project Gene Disruption Project: Single P-Element Insertions Mutating 25% of Vital

Drosophila Genes. Genetics 153: 135–177. https://doi.org/10.1093/genetics/153.1.135

Takeo S., S. K. Swanson, K. Nandanan, Y. Nakai, T. Aigaki, *et al.*, 2012 Shaggy/glycogen synthase

kinase 3β and phosphorylation of Sarah/regulator of calcineurin are essential for completion of

Drosophila female meiosis. Proc National Acad Sci 109: 6382–6389.

https://doi.org/10.1073/pnas.1120367109

Wada T., T. Takagi, Y. Yamaguchi, D. Watanabe, and H. Handa, 1998 Evidence that P-TEFb alleviates

the negative effect of DSIF on RNA polymerase II-dependent transcription in vitro. Embo J 17: 7395–

7403. https://doi.org/10.1093/emboj/17.24.7395

Wang X., S. Hang, L. Prazak, and J. P. Gergen, 2010 NELF Potentiates Gene Transcription in the

Drosophila Embryo. Plos One 5: e11498. https://doi.org/10.1371/journal.pone.0011498

Wang S., P. Chai, R. Jia, and R. Jia, 2018 Novel insights on m6A RNA methylation in tumorigenesis: a

double-edged sword. Mol Cancer 17: 101. https://doi.org/10.1186/s12943-018-0847-4

Wang F., H.-H. Qiao, R.-G. Xu, J. Sun, R. Zhu, *et al.*, 2019 pNP Transgenic RNAi System Manual in

Drosophila. Bio-protocol 9: 1--12. https://doi.org/10.21769/bioprotoc.3158

Warda A. S., J. Kretschmer, P. Hackert, C. Lenz, H. Urlaub, *et al.*, 2017 Human METTL16 is a N6-

methyladenosine (m6A) methyltransferase that targets pre-mRNAs and various non-coding RNAs.

Embo Rep 18: 2004–2014. https://doi.org/10.15252/embr.201744940

Wayne M. L., J. B. Hackett, and T. F. C. Mackay, 1997 QUANTITATIVE GENETICS OF OVARIOLE

NUMBER IN DROSOPHILA MELANOGASTER. I. SEGREGATING VARIATION AND FITNESS.

Evolution 51: 1156–1163. https://doi.org/10.1111/j.1558-5646.1997.tb03963.x

Wayne M. L., and T. F. C. Mackay, 1998 Quantitative Genetics of Ovariole Number in Drosophila

melanogaster. II. Mutational Variation and Genotype-Environment Interaction. Genetics 148: 201–

210. https://doi.org/10.1093/genetics/148.1.201

Xu X., L. Ma, X. Zhang, S. Guo, W. Guo, *et al.*, 2022 A positive feedback circuit between RN7SK snRNA

and m6A readers is essential for tumorigenesis. Mol Ther J Am Soc Gene Ther.

https://doi.org/10.1016/j.ymthe.2022.12.013

Xue Y., Z. Yang, R. Chen, and Q. Zhou, 2010 A capping-independent function of MePCE in stabilizing

7SK snRNA and facilitating the assembly of 7SK snRNP. Nucleic Acids Res 38: 360–369.

https://doi.org/10.1093/nar/gkp977

Yamaguchi Y., T. Takagi, T. Wada, K. Yano, A. Furuya, *et al.*, 1999 NELF, a Multisubunit Complex

Containing RD, Cooperates with DSIF to Repress RNA Polymerase II Elongation. Cell 97: 41–51.

https://doi.org/10.1016/s0092-8674(00)80713-8

Yang Z., Q. Zhu, K. Luo, and Q. Zhou, 2001 The 7SK small nuclear RNA inhibits the CDK9/cyclin T1

kinase to control transcription. Nature 414: 317–322. https://doi.org/10.1038/35104575

Yang Y., C. D. Eichhorn, Y. Wang, D. Cascio, and J. Feigon, 2019 Structural basis of 7SK RNA 5´-γ-

phosphate methylation and retention by MePCE. Nat Chem Biol 15: 132–140.

https://doi.org/10.1038/s41589-018-0188-z

Yang Y., S. Liu, S. Egloff, C. D. Eichhorn, T. Hadjian, *et al.*, 2022 Structural basis of RNA conformational

switching in the transcriptional regulator 7SK RNP. Mol Cell 82: 1724-1736.e7.

https://doi.org/10.1016/j.molcel.2022.03.001

Yik J. H. N., R. Chen, A. C. Pezda, C. S. Samford, and Q. Zhou, 2004 A Human Immunodeficiency Virus

Type 1 Tat-Like Arginine-Rich RNA-Binding Domain Is Essential for HEXIM1 To Inhibit RNA

Polymerase II Transcription through 7SK snRNA-Mediated Inactivation of P-TEFb. Mol Cell Biol 24:

5094–5105. https://doi.org/10.1128/mcb.24.12.5094-5105.2004

Zhang B., Z. Liu, J. Liu, K. Cao, W. Shan, *et al.*, 2021 Long non-coding RNA ST8SIA6-AS1 promotes the

migration and invasion of hypoxia-treated hepatocellular carcinoma cells through the miR-

338/MEPCE axis. Oncol Rep 45: 73–82. https://doi.org/10.3892/or.2020.7864

Zhou M., M. A. Halanski, M. F. Radonovich, F. Kashanchi, J. Peng, *et al.*, 2000 Tat Modifies the Activity

of CDK9 To Phosphorylate Serine 5 of the RNA Polymerase II Carboxyl-Terminal Domain during

Human Immunodeficiency Virus Type 1 Transcription. Mol Cell Biol 20: 5077–5086.

https://doi.org/10.1128/mcb.20.14.5077-5086.2000

Zhou Q., T. Li, and D. H. Price, 2012 RNA Polymerase II Elongation Control. Biochemistry-us 81: 119–

143. https://doi.org/10.1146/annurev-biochem-052610-095910

Zhu W., and S. D. Hanes, 2000 Identification of Drosophila Bicoid-interacting proteins using a custom

two-hybrid selection. Gene 245: 329–339. https://doi.org/10.1016/s0378-1119(00)00048-2

Zieve G., and S. Penman, 1976 Small RNA species of the HeLa cell: Metabolism and subcellular

localization. Cell 8: 19–31. https://doi.org/10.1016/0092-8674(76)90181-1

Zimmermann L., A. Stephens, S.-Z. Nam, D. Rau, J. Kübler, *et al.*, 2018 A Completely Reimplemented

MPI Bioinformatics Toolkit with a New HHpred Server at its Core. J Mol Biol 430: 2237–2243.

https://doi.org/10.1016/j.jmb.2017.12.007

Fig. 2

(a)

Egg-Laying Assay

Eggs/female/day

wt — ***
$bin3^{2-7}/+$ — ****
$bin3^{2-7}/Df$
$bin3^{2-7}/Df; bin3^+/+$ — ****
$bin3^{2-7}/Df; CycT^{j11B2}/+$ — ****

(b)

Germline cells
Somatic cells
Oocyte

Egg chambers

Germarium

Nurse cells

(c) 3xHA-Bin3 DNA Vasa
$bin3^{TG4.0} > UAS-3xHA-bin3$

(c′) 3xHA-Bin3
$bin3^{TG4.0} > UAS-bin3-3xHA$

(d) 3xHA-Bin3 DNA Vasa
$bin3^{TG4.0}/+$

(d′) 3xHA-Bin3
$bin3^{TG4.0}/+$

(e) 7SK DNA
wt

(e′) 7SK
wt

(f) 7SK DNA
$bin3^{2-7}/Df$

(f′) 7SK
$bin3^{2-7}/Df$

(g) Ovarioles Per Ovary

ns   ns   ns

wt $bin3^{\Delta}/+$ $bin3^{\Delta}/Df$

(h) Ovariole Asymmetry

ns   ns   ns

wt $bin3^{\Delta}/+$ $bin3^{\Delta}/Df$

(i)

Egg-laying Assay with Germline-Specific Bin3 expression

Eggs/female/day

wt — *
$bin3^{2-7}/+; nos-GAL4::VP16/+$ — ***
$bin3^{2-7}/Df; nos-GAL4::VP16/+$
$bin3^{2-7}/Df; nos-GAL4::VP16/UAS-bin3$ — ns

Fig. 3

(a) **Climbing Assay**

Percent of flies climbing 8 cm in 12" (y-axis, 0 to 150)

*** *bin3²⁻⁷/+*

bin3²⁻⁷/Df

*** *bin3²⁻⁷/Df; bin3⁺/+*

*** *bin3²⁻⁷/Df; CycTʲ¹¹ᴮ²/+*

(b) *bin3²⁻⁷/+*

(c) *bin3²⁻⁷/Df*

(d) *bin3²⁻⁷/Df; bin3⁺/+*

(e) *bin3²⁻⁷/Df; CycTʲ¹¹ᴮ²/+*

Fig. 4

(a) Catalytic Domain (wild-type)
Y421 (MePCE)
Y795 (Bin3)
SAH
7SK γ-Pi

(b) Catalytic Domain (Y to A mutation)
Y421A (MePCE)
Y795A (Bin3)
SAH
7SK γ-Pi

(c)
Metazoan-Specific Motif (MSM)

Bin3 (*D. melanogaster*) 301-RKEVI**VP**P**TKFLLGGNI**S**DPLNLNSL**QN**E**NTSNASS-336
MePCE (*H. sapiens*)   175-SDCDS**VLP**SN**FLLGGNI**F**DPLNLNSL**LD**E**EVSRTLN-210
BCDIN3 (*D. rerio*)   115-SESES**VLP**TN**FLLGGNI**F**DPLNLNSL**LD**E**EVNKALN-150
Bmc1 (*S. pombe*)   ------------------------------------

(d) *bin3* locus on chromosome 2R
P2   P1   *bin3*
Inserted in *bin3^TG4.0*: SA | T2A | GAL4 | poly(A)
Deleted in *Df*
~30.3 kb

(e) *bin3* mRNA Levels — mRNA Level (relative to wt)
ns, ****

(f) 7SK RNA Levels — RNA Level (relative to wt)
, **

(g) U6 RNA Levels — RNA Level (relative to wt)
ns, *
wt; *bin3^TG4.0*/+; *bin3^TG4.0*/*Df*

(h) *UAS-bin3^wt*: UAS — DSCP — *bin3^wt* cDNA — 3xHA
(i) *UAS-bin3^Y795A*: UAS — DSCP — *bin3^Y795A* cDNA — 3xHA
(j) *UAS-bin3^ΔMSM*: UAS — DSCP — *bin3^ΔMSM* cDNA — 3xHA

(k) *3xHA-bin3* mRNA Levels — mRNA Level (relative to *UAS-bin3*)
ns, ns

bin3^TG4.0/*Df; UAS-bin3^wt*/+
bin3^TG4.0/*Df; UAS-bin3^Y795A*/+
bin3^TG4.0/*Df; UAS-bin3^ΔMSM*/+

(l) 3xHA-Bin3 Protein Levels

bin3^TG4.0	+	+	+	+
UAS-3xHA-bin3^wt	−	+	−	−
UAS-3xHA-bin3^Y795A	−	−	+	−
UAS-3xHA-bin3^ΔMSM	−	−	−	+

250 kDa
150 kDa — 3xHA
50 kDa
37 kDa — Actin

0.0 ± 0.0 (*)   1.0 ± 0.0   1.1 ± 0.071 (ns)   0.85 ± 0.35 (ns)

(a)

Egg-Laying Assay with *UAS-bin3* Transgenes

(b)

Climbing Assay with *UAS-bin3* Transgenes

Fig. 7

		Bin3 binds 7SK	Bin3 caps 7SK	Bin3 stabilizes 7SK	7SK snRNP assembles and represses P-TEFb	**Repressed P-TEFb promotes RNAPII pausing to support…**		
						Fecundity	Climbing	Wing posture
(a)	Bin3[wt]	✓	✓	✓	✓	✓	✓	✓
(b)	Bin3[Y795A]	✓	✗	✓	✓	✓	✓	✓
(c)	Bin3[MSM]	✓	✓	✓	✓	✓	✓	✗

Fig. 8

(a) <u>7SK-dependent mechanism</u>

Bin3

Methyltransferase
Activity

*Not required for
7SK stability or
function*

Hexim

Bin3

Larp7

Cdk9

CycT

Kinase
Activity

RNAPII
transcription
elongation
repressed

(b) <u>Tissue-specific, 7SK-independent mechanism</u>

?

MSM

Bin3

?

MSM

Bin3

RNAPII
transcription
elongation
repressed