In vitro modelling of anterior primitive streak patterning with human pluripotent stem cells identifies the path to notochord progenitors

M. Robles-Garcia, C. Thimonier, K. Angoura, E. Ozga, H. MacPherson, G. Blin[*]

Institute for Regeneration and Repair, Institute for Stem Cell Research, School of Biological Sciences, The University of Edinburgh, 5 Little France Drive, Edinburgh BioQuarter, Edinburgh EH16 4UU, UK.

* Author for correspondence (guillaume.blin@ed.ac.uk)

Abstract

Notochord progenitors (NotoPs) represent a scarce yet crucial embryonic cell population, playing important roles in embryo patterning and eventually giving rise to the cells that form and maintain intervertebral discs. The mechanisms regulating NotoPs emergence are unclear. This knowledge gap persists due to the inherent complexity of cell fate patterning during gastrulation, particularly within the anterior primitive streak (APS), where NotoPs first arise alongside other important progenitors including neuro-mesodermal and endodermal progenitors.

To gain insights into this process, we use micropatterning together with FGF and the WNT pathway activator CHIR9901, to guide the development of human embryonic stem cells into reproducible patterns of APS cell fates. We show that small variations in CHIR9901 dosage dictate the downstream dynamics of endogenous TGFbeta signalling which in turn controls cell fate decisions. We show that sustained NODAL signalling induces endoderm while NODAL inhibition is needed for NMP specification. Furthermore, we unveil a crosstalk between TGFbeta and WNT signaling pathways, wherein TGFbeta inhibition enhances WNT activity. Finally, we demonstrate that the timely inhibition of TGFbeta signalling is imperative for the emergence of NotoPs.

Our work elucidates the signalling regimes underpinning NotoPs emergence and provides novel insights into the regulatory mechanisms controlling the balance of APS cell fates during gastrulation.

Introduction

In vertebrate embryos, the tissues of the posterior axis, including the spinal cord, the cartilage, bones and muscles of the spine, as well as the gut, are all laid down progressively in an anterior-to-posterior direction. This evolutionary-conserved process, termed axial elongation (reviewed in (Henrique et al., 2015; Neijts et al., 2014; Wymeersch et al., 2021), is governed by lineage-restricted progenitors emerging during gastrulation in the anterior portion of the primitive streak (APS) (Fig1A).

Among these, notochord progenitors (NotoPs) give rise to the notochord, a cylindrical structure that forms along the ventral midline in chordate embryos (Balmer et al., 2016; Stemple, 2005). Work in animal models has shown that the notochord ensures essential functions during development, both as a source of localised signalling molecules necessary for the proper patterning of adjacent tissues (Streit and Stern, 1999) and as a mechanical structure contributing to the straightness of the rostro-caudal axis as the embryo elongates (Bagnat and Gray, 2020; McLaren and Steventon, 2021). Eventually, the cells that form the notochord contribute to the nucleus pulposus (Choi et al., 2008; McCann et al., 2012), the central region of intervertebral discs responsible for the homeostasis of the surrounding fibro-cartilagenous tissue (reviewed in (Wise et al., 2020)).

Previous work in mouse embryos has shown that the specification of the notochord requires the cooperation of WNT and NODAL signalling (Dunn et al., 2004; Lickert et al., 2002; Merrill et al., 2004; Vincent et al., 2003; Yamamoto et al., 2001). However, the exact signalling requirements for the specification of NotoPs has remained elusive. Indeed, in spite of previous efforts (Colombier et al., 2020; Diaz-Hernandez et al., 2020; Winzi et al., 2011; Zhang et al., 2020), the derivation of NotoPs from pluripotent stem cells remains inefficient unless transcription factors regulating the notochord fate are overexpressed and even these conditions lead to mixed population of cells including alternative cell fates from the mesoderm and endoderm lineage (Schifferl et al., 2023; Warin et al., 2024). Furthermore, apart from one exception (Xu et al., 2021), 3D organoids that mimic axial elongation also lack notochord in spite of the fact that WNT and NODAL signalling are both active in these in vitro models (Beccari et al., 2018; Cermola et al., 2021; Moris et al., 2020; Turner et al., 2017) suggesting there may be additional unknown cues needed for notochord specification.

This gap in understanding stems from the relative rarity of NotoPs in the embryo and from the complexity and rapidity of the cell fate decisions during gastrulation. In mouse embryos, NotoPs emerge in close proximity to both the definitive endoderm (Beddington and Robertson, 1989; Scheibner et al., 2021), and neuromesodermal progenitors (NMPs), the bi-fated cells that form the somitic mesoderm and the spinal cord (Cambray and Wilson, 2002; Cambray and Wilson, 2007; Forlani et al., 2003; Koch et al., 2017; Tzouanacou et al., 2009). All three populations, endoderm, NotoPs and NMPs, have been reported to emerge at around mid-gastrulation (Ang and Rossant, 1994; Cambray and Wilson, 2002; Pour et al., 2022; Scheibner et al., 2021; Sulik et al., 1994; Yamanaka et al., 2007), eventually forming a progenitor zone organised along the antero-posterior axis of the embryo. On the anterior side of the primitive streak, endodermal progenitors undergo partial epithelial to mesenchymal transition and intercalate into the underlying visceral endoderm to form the gut endoderm (Kwon et al., 2008; Scheibner et al., 2021; Viotti et al., 2014). NotoPs establish

the ventral epithelial layer of the node, demarcating the rostral and caudal regions of the embryo (Bakker et al., 2016; Kinder et al., 2001). NMPs populate the epithelial space directly adjacent and posterior to the node and together with NotoPs form the progenitor growth zone that fuels axial elongation (Fig 1A) (Abdelkhalek et al., 2004; Wymeersch et al., 2019). How the patterning and balanced proportion of these populations is established is not understood.

Here, we set out to identify the signalling requirements distinguishing NotoPs from other APS cell fates in a human context. To tackle this challenge and circumvent the technical and ethical limitations inherent to research on rare embryonic cell populations, we use micropatterning to guide the development of human embryonic stem cells (hESC) into reproducible patterns of APS cell fates. We start with an NMP-inducing medium containing FGF and CHIR99021 (CHIR), a potent activator of the WNT pathway (Cohen and Goedert, 2004). First, we show that in contrast to cells grown as a 2D monolayer culture, cells grown on micropatterns preferentially differentiate towards mes-endodermal lineages. We show that partial inhibition of endogenous NODAL signalling prevents endoderm specification and restores the emergence of NMPs but fails to induce NotoPs. Then we uncover that hESC grown on micropatterns respond non-monotonically to increasing doses of CHIR with distinct downstream temporal dynamics of NODAL signalling. We also identify a crosstalk between WNT and NODAL wherein NODAL inhibition potentiates WNT signalling activity. By inhibiting NODAL and BMP signalling 24h post CHIR induction when the cells are still uncommitted to endoderm, we are able to efficiently redirect the cells from the endoderm to the NotoP fate.

Overall, our study uncovers signalling cross-talks and dynamics that correlate with key lineage restrictions prior to axial elongation and identifies the path to the notochord lineage.

Results

hESC colony confinement directs patterning of mesendoderm-biased cell fates.

In order to study the mechanisms underlying NotoPs specification in a human context, we set out to establish an *in vitro* system that would mimic aspects of the formation of the axial progenitor zone using hESC. NotoPs emerge in close proximity to NMPs within a small, confined region of the embryo (Cambray and Wilson, 2007). Furthermore, scRNAseq analysis have shown that *in vitro* derived NMPs may contain a rare population of cells with a node-like signature (Edri et al., 2019). Therefore, we hypothesised that combining an NMP derivation medium (Frith and Tsakiridis, 2019; Gouti et al., 2014) with geometrical confinement (Blin, 2021) would elicit the self-organisation of hESC colonies into distinct domains of APS cell fates including NotoPs (Fig 1B)

Figure 1 Geometrical confinement elicits radial patterning of endoderm and mesoderm biased cell fates *A Diagrammatic representation of the putative organisation and hierarchies of cell fates around the node in a human embryo at the end of gastrulation (left). Extra-embryonic structures are omitted. Relevant cell lineages are indicated together with the associated cell fate markers used in this study (Right). B Illustration of the protocol used in C to E (Top) and confocal max projections showing unconstrained and micropatterned cultures (Bottom). The yellow outline indicate the colony shown at higher resolution in C. Scale bar: 200µm. C-E Confocal max projections of micropatterned colonies stained at 48h, scale bar: 100µm. The corresponding Z-projection is shown below each image with yellow arrowheads pointing at SOX17+ or SOX2- cells lining the central domain at the bottom of the colony. Radial profiles of cell type abundance are shown on the right. c: number of analysed colonies, n: number of analysed nuclei, shaded area: 95% confidence interval. F Confocal images of colonies stained at day 5 of differentiation. Wide fields are shown on the left and a representative colony on the right. Images are max projections except for the selected colony stained with HAND1, ColV and Fibronectin where a z-slice was selected to better show the localisation of the positive signal. Scale bar: 100µm. APS: anterior primitive streak, Epi: Epiblast, NotoP: Notochord Progenitors, NMPs:*

Neuromesodermal progenitors. NE: Neurectoderm, PXM: paraxial mesoderm, LPM: Lateral plate
mesoderm, Ant.E: Anterior endoderm.

To test this idea, we stained the cells 48h post CHIR and FGF induction for the NMP markers TBXT
and SOX2 and included the endodermal marker SOX17 (Kanai-Azuma et al., 2002; Viotti et al., 2014)
as a proxy for the emergence of additional APS cell fates that do not normally appear in NMP
differentiation monolayers (Frith and Tsakiridis, 2019) (Fig 1B). In line with previous reports (Frith et
al., 2018; Gouti et al., 2014), cells cultured in conventional 2D dishes co-expressed TBXT and SOX2
(Frith et al., 2018; Gouti et al., 2014) and remained negative for SOX17 (Fig 1B). In sharp contrast,
cells grown on micropatterns consistently formed radially organised domains of cell fate markers, with
a TBXT/SOX2 domain located between a SOX2-only domain in the centre and a SOX17 domain at
the periphery (Fig 1B and 1C). We confirmed that this phenomenon was reproducible across several
human pluripotent cell lines (Sup Fig 1).
We next tested whether different colony sizes may change the proportion of cell fates on micropatterns
(Sup Fig 2A). (Kwon et al., 2008; Scheibner et al., 2021; Viotti et al., 2014)Using our established
quantitative immunofluorescence pipeline (Blin et al., 2019; Wisniewski et al., 2019), we measured the
proportion of each individual population (Sup Fig 2B) and plotted these proportions as a function of the
radial distance from the colony edge (Sup Fig 2C). We found that all three domains, i.e SOX17,
TBXT/SOX2 and SOX2-only domains, remained located at a consistent distance from the colony edge
across all colony diameters except for colonies below 320μm where this rule did not apply as strictly.
Incidently, the central domain of SOX2 expressing cells increased in size proportionally with colony
diameter effectively increasing the percentage of SOX2-only cells. These observations may suggest
that the mechanisms driving the radial cell fate organisation in this context is boundary-driven as
reported in other micropatterned colony systems (Etoc et al., 2016; Martyn et al., 2019; Warmflash et
al., 2014). For the subsequent experiments, we decided to use 500μm colonies because this diameter
offered a good compromise for analysis and imaging with a clear radial fate marker distribution (Fig 1
C).

We next asked about the developmental state of the cells forming the central SOX2 domain. In mouse
embryos, SOX2 is initially co-expressed with OCT4 in the pluripotent epiblast and remains expressed
in the developing neurectoderm while OCT4 becomes progressively lost as the cells exit pluripotency
(Avilion et al., 2003; Osorno et al., 2012). In our colonies, OCT4 was still expressed in the central
SOX2 domain at this stage indicating that the cells had not yet exited pluripotency (Fig 1D).

Next, we tested for the presence of NotoPs. We first looked for the co-expression of FOXA2 and TBXT

which are both essential for the development of the notochord (Ang and Rossant, 1994; Lolas et al.,

2014; Tamplin et al., 2011; Yamanaka et al., 2007). Cells expressing FOXA2 could be found all around

the colony spanning a domain of approximately 100μm from the edge (Fig 1E). This domain could be

divided into a SOX17+ outter domain indicative of the endodermal fate and a FOXA2/TBXT inner

domain co-localising with the SOX2/TBXT domain shown in Fig 1C. Since these markers are also

transiently co-expressed in the nascent mesendoderm during gastrulation (Burtscher and Lickert,

2009), we performed FISH against the NotoP-specific marker NOTO (Abdelkhalek et al., 2004;

Plouhinec et al., 2004). We did not observe any positive cells for this marker suggesting that the

TBXT/FOXA2+ cells in these colonies likely represent an early mesendoderm population at 48h.

To further characterise lineages emerging in micropatterned colonies, we repeated the same protocol

as in Fig 1B and cultured the cells for an additional 3 days in unsupplemented N2B27 (Fig 1F). Over

time, the colonies established a 3-dimensional structure composed of a cavity-comprising SOX2-

positive core surrounded by a mass of SOX17 endodermal cells and clusters of HAND1+ cells. The

existence of HAND1+ cells in these colonies indicate that some of the cells differentiated either to

lateral plate mesoderm or extraembryonic mesoderm (Pham et al., 2022). On the other hand, we were

unable to find evidence of notochord-like cells co-expressing TBXT, FOXA2 and SOX9 which is

normally found in the notochord (Bagheri-Fam et al., 2006), confirming that the TBXT/FOXA2+

population identified at 48h failed to engage in the notochord lineage. We were also unable to find

TBXT/SOX2 double positive cells in these structures suggesting that NMPs were also absent.

The 3-dimensional multi-tissue architecture observed here was remarkably consistent across colonies

(Fig 1F). Interestingly, colonies adopted a 3-dimensional organisation as early as 48h with the

formation of a dome-like structure where SOX2+ cells were elevated in comparison to other cell types

(Fig 1C, D, E and Sup Fig2A). In particular, we consistently observed that SOX17+ cells, while

abundant at the periphery, also formed a sparse epithelial layer lining the bottom of the colony (arrow

heads in Fig 1C, E and Sup Fig2A), perhaps reflecting the behaviour of the nascent endoderm in vivo

which undergoes partial EMT as it segregates from the mesoderm to form the gut endoderm

epithelium during gastrulation (Kwon et al., 2008; Scheibner et al., 2021; Viotti et al., 2014). Staining

for Collagen V and Fibronectin at day 5 (Fig 1F), two extracellular matrix proteins expressed in the

endoderm and mesoderm in human embryos (Zhao et al., 2024), revealed that the cells secreted their

own extra-cellular matrix (ECM). Both proteins formed a basal membrane surrounding the SOX2+

domain while Fibronectin also formed a complex network inside the endodermal domain. These

observations indicate that the cells organised their own ECM which likely contributed to maintaining

the separation between cell fate domains and the overall morphogenetic process observed in the colonies.

Altogether, these initial experiments allowed us to establish an *in vitro* system where hESC organise into stereotypic patterns of cell fates including endoderm and mesoderm and undergo reproducible complex morphogenesis over time. While this system provides a good starting point, we were unable to detect NotoPs in these conditions, raising the question of which additional signals might be required to generate this cell type. Furthermore, NMPs did not persist on micropatterns despite the use of an NMP inducing medium suggesting that confinement potentiates endogenous cues that deflect the cells from axial cell fates towards alternative lineages.

2- Dynamics of NODAL and WNT signalling correlate with the loss of axial cell fates and the emergence of definitive endoderm and lateral plate mesoderm

To gain insights into the mechanisms driving cell fate diversification and the loss of axial lineages on micropatterns, we performed time course experiments and bulk RNA Nanostring analysis. We used a panel of probes consisting of the 780 genes included in the standard human embryonic stem cell gene panel together with 30 additional custom probes (listed in Sup Table 1). This panel covered a wide array of genes involved in differentiation, metabolism, signalling pathways and the cell cycle. In order to analyse our dataset, we used the Bioconductor package moanin (Varoquaux and Purdom, 2020) which allowed us to group individual genes based on their temporal profile (see Methods). We identified 7 clusters that are shown in Fig 2A. Cluster 1 and 2 identify genes that are expressed at the start and then progressively downregulated. As expected, these include pluripotency markers such as MYC, OCT4, DPPA4, DNMT3B and ZFP42. We found NANOG to be highest at around 36h and then lost rapidly consistently with the fact that NANOG is re-expressed in the posterior epiblast at the onset of gastrulation in mouse embryos (C et al., 2018; Hart et al., 2004; Osorno et al., 2012). We also found SOX2 in cluster 2 as a gene that is rapidly downregulated and starts to re-emerge at around 72h most likely as a result of its expression in the central domain undergoing neural differentiation (Fig 1H).

Fig 2 Dynamics of NODAL, WNT and BMP signalling correlate with the loss of axial cell fates and the emergence of definitive endoderm and lateral plate mesoderm - A *Nanostring time course analysis. Cells were treated as in Fig. 1B. For each cluster, a model spline of the cluster is shown at the top of the gene expression heatmap. Genes are ordered from top to bottom in decreasing order of maximum differential expression. Colours represent the mRNA count normalised on a per gene basis. **B** Time course analysis via quantitative immunofluorescence of 500µm colonies. Left: montage of confocal max projections, Middle: stacked bar plot showing the relative proportions of individual cell populations over time, Right: Radial profiles of the percentage of cells expressing the markers indicated at the top of the plots. **C** Widefield images of 500µm colonies stained via branched DNA FISH at selected time points.*

Cluster 4 and 5 identified genes that become expressed as early as 12h post-induction. Cluster 4 regrouped genes with a sustained expression after 36h while the genes in cluster 5 peaked at 36h and

decreased thereafter. These two clusters revealed a clear APS signature with the expression of genes
associated with the APS in mice including MIXL1 (Hart et al., 2002), LHX1 (Costello et al., 2015),
EOMES (Costello et al., 2011), GSC, CDX2, FOXA2 (Burtscher and Lickert, 2009), KLF5 (Aksoy et
al., 2014) as well as the NMP-associated gene NKX1-2 (Albors et al., 2018). Encouragingly we also
found a peak of expression of SLIT2 and CAV1 at 36h, recently reported as markers of human
embryonic notochord (Paillat et al., 2023; Warin et al., 2024). However, NMPs and NotoPs associated
genes decreased over time indicating that axial progenitors failed to emerge. We confirmed this using
quantitative immunofluorescence and observed that while the majority of the cells expressed
TBXT/SOX2 at 36h initiating from the periphery at 24h, these markers were progressively lost in
favour of SOX17 (Fig 2B and Sup Fig 3). Similarly, TBXT/FOXA2 co-expressing cells increased in
proportion until 36h and were replaced by FOXA2 only cells at the periphery suggesting that some of
these cells were on their way to form endoderm (Fig 2B and Sup Fig 3).

In fact, many of the genes found in cluster 4 and 5 of our Nanostring dataset are known regulators of
the endodermal lineage. For example, both MIXL1 and KLF5 are required for specification of the
definitive endoderm in the mouse (Aksoy et al., 2014; Hart et al., 2002; Moore-Scott et al., 2007) and
LHX1, while expressed in the node, works together with OTX2 (found expressed transiently in cluster
3) to define anterior endoderm (Costello et al., 2015). Furthermore, looking at cluster 6 and 7, where
genes become upregulated from 48h onwards, we could confirm the emergence of additional
endodermal markers such as FOXA1 (Ang and Rossant, 1994), PDGFRA and GATA4 as well as
several ECM components, some of which likely secreted by the endoderm including FN1, COL4A2,
COL5A1, COL5A2, FBN2 and FLNC.

Endoderm was not the only lineage emerging in our colonies as we could observe clear evidence of
mesoderm specification. GATA6 (Morrisey et al., 1996; Zhao et al., 2005) and EOMES (Costello et al.,
2011), found in cluster 4, are both involved in the specification of the endoderm and the cardiac
mesoderm lineage in the streak, and cluster 6 and 7 regrouped many genes associated with the
lateral plate mesoderm including GATA3, HAND1, ISL1, TBX3 (Washkowitz et al., 2012) and MESP1
alongside the EMT markers SLUG, SNAIL and MSX2.

Overall, these data confirm our previous observations and show that cells on micropatterns follow a
coherent developmental program. However, while the cells initially follow the route towards axial cell
fates (i.e express APS markers), the cells eventually differentiate towards alternative lineages
including endoderm and lateral plate mesoderm.

We thus turned our focus towards endogenous signalling pathways that may explain the observed
endoderm and mesoderm differentiation. NODAL is a known driver of mesodermal and endodermal

specification (Robertson, 2014) and its expression is positively regulated by the canonical WNT
pathway (Ben-Haim et al., 2006; Norris et al., 2002); a pathway that we stimulate with CHIR in our
cultures. Our Nanostring data showed upregulation and sustained expression of the WNT target gene
LEF1 (Cadigan and Waterman, 2012) as early as 12h while NODAL expression peaked at 24h before
decreasing progressively (FIG 2A). This peak of NODAL was followed by a peak of CER1 expression
at 36h in line with previous report showing that NODAL induces its own inhibitor (Colombier et al.,
2020; Funa et al., 2015). FISH confirmed the temporal expression profile of NODAL and CER1 (Fig2
C) and showed that NODAL expression was widespread across the colony at 24h alongside its
antagonist LEFTY2. Interestingly, CER1 (Belo et al., 1997; Perea-Gomez et al., 2002) and the WNT
antagonist DKK1, both expressed in the hypoblast of primate embryos (Bergmann et al., 2022) were
expressed in a domain that overlapped spatially and temporally with the SOX17 domain (Fig2 A and
B) indicating that at least a fraction of the endoderm emerging in micropattern colonies may adopt an
hypoblast identity.

Altogether, our data revealed the implementation of a regulatory network of signalling molecules in
micropatterns involving WNT, NODAL and their respective inhibitors.

3- Nodal inhibition rescues NMP emergence

Our results raise the possibility that endogenous NODAL signalling is responsible for the loss of axial
cell fates and the emergence of definitive endoderm and lateral plate mesoderm on micropatterns. To
confirm this hypothesis, we next decided to titrate NODAL activity using increasing doses of the
NODAL receptors inhibitor SB431542 (SB) (Inman et al., 2002) (Fig 3A). We observed that even low
amounts of SB was sufficient to strongly reduce the proportion of SOX17 and FOXA2 positive cells
confirming that endogenous NODAL signalling activity is indeed responsible for the emergence of
endoderm in this system (Fig 3A-1µM). Further increasing the dose of SB progressively expanded the
central domain of SOX2 single positive cells restricting TBXT expression to the periphery. We also
noted that higher doses of SB, depleted the FOXA2/TBXT population while maintaining a high
percentage of SOX2/TBXT population at the periphery (Fig3 A 3µM and 10µM). We hypothesised that
this SOX2/TBXT positive and FOXA2 negative cell population may be engaging in the NMP cell fate
and could drive elongation.

Figure 3 Nodal inhibition rescues NMP specification and enables tissue elongation. *A SB dose response in 500µm colonies fixed at 48h. (i) Max projections of confocal z-stacks (top) and z-projection (bottom). (ii) Stacked bar plots showing the relative proportions of individual cell populations. (iii) Heatmaps showing the radial profile of average nuclear marker intensity scaled*

To test this idea, we cultured the cells for 3 days in CHIR/FGF with or without 3µM or 10µM SB and for an additional 3 days without CHIR/FGF while maintaining SB concentration (Fig 3B and C). Since previous work with 3D models of axial elongation showed that the starting number of cells is critical to the competence of cell aggregates to elongate (Beccari et al., 2018; Martins et al., 2020; Moris et al., 2020; Olmsted and Paluh, 2021; Sanaki-Matsumiya et al., 2022; Turner et al., 2017; Veenvliet et al., 2020), we tested two different colony diameters (500µm and 300µm). As expected, we did not observe elongation in the control with either of these diameters but we did see elongation in SB containing conditions. To our surprise, we observed that competence to elongate was a function of both colony diameter and SB concentration, with elongation occurring consistently across colonies only with 3µM SB on 500µM micropatterns and with 10µM SB on 300µm micropatterns (Fig 3B and C). We next checked whether SOX2/TBXT double positive putative NMPs were maintained in these structures (Fig 3C). Indeed, we observed that elongation correlated with the conditions favouring the highest abundance of SOX2/TBXT cells which were located towards the bottom of the dish while elongated structures comprised a cavity and were exclusively composed of SOX2 positive cell.

Together, these results demonstrate that suppression of NODAL signalling is necessary for the specification of NMPs and that adequate NODAL suppression and colony size combinatorially dictate uniaxial growth on micropatterns.

4- Small variations in CHIR dosage leads to radically distinct cell fate patterning outcomes

None of the tested conditions so far allowed us to find NOTO+ cells using FISH staining at 48h (not shown) raising the question of what other cues would be needed for NotoPs emergence. To gain further insights into the mechanisms of cell fate patterning on micropatterns upon CHIR exposure, and perhaps identify clues about the missing conditions for NotoPs emergence, we decided to test a range of CHIR concentrations and monitor markers of cell fates 48h post-induction via quantitative

immunofluorescence (FIG 4A,B) and Nanostring analysis (Fig 4C). In the absence of CHIR, all the
cells remained OCT4/SOX2 double positive and negative for TBXT, SOX17 and FOXA2 (FIG 4A, B
and Sup Fig 4 A, B) showing that exogenous stimulation of the WNT pathway is required to initiate
differentiation. More importantly, we found a clear dose-dependent effect of CHIR on the levels and
spatial distributions of cell fate markers: The percentage of SOX2+ cells was negatively correlated to
CHIR concentration while the percentage of TBXT+ cells increased proportionally, in line with the fact
that TBXT is a known direct target of the WNT/β-catenin pathway (Arnold et al., 2000). On the other
hand, SOX17 and FOXA2 did not correlate linearly with CHIR levels as these markers emerged at
intermediate CHIR concentrations and were rare or absent at higher concentrations (Fig 4A and Sup
Fig4 B). Instead, high doses of CHIR induced the expression of the lateral plate marker HAND1 (Sup
Fig 4C). Our Nanostring analysis supported these results with a peak of endodermal markers at 1 and
2µM CHIR while colonies treated with the highest doses of CHIR expressed the LPM markers HAND1,
TBX3 and the posterior marker CDX2 as well as HOX genes. These data suggest that endodermal
cells are replaced by more posterior mesodermal fates at high CHIR concentration.

Interestingly, when considering the spatial distribution of these markers, we observed a clear inward
shift of the different cell populations when CHIR was increased (FIG 3A iii): TBXT was expressed at
the periphery with 1µM CHIR and progressively expressed throughout the colony with increasing
CHIR concentrations. In parallel, SOX2 was downregulated at the periphery with 2µM CHIR and
higher concentrations were required to repress SOX2 in the centre. As a result, TBXT/SOX2 double
positive cells were found at the periphery at 1µM CHIR but were progressively shifted and restricted to
the centre, forming a smaller fraction of the total population with increasing CHIR concentration where
TBXT single positive cells became predominant. Finally, SOX17 appeared restricted at the edge at
1µM CHIR, but spanned a 150 µm domain at 2µM, while at 3µM, the few SOX17+ cells present in the
colony were located at around 100µm from the border (FIG 4A iii – yellow line). These observations
further support the idea that CHIR-induced patterning is initiated from the periphery of the colony.

Together, our data show a complex, non-linear dose-dependent action of CHIR on cell fates, with
intermediate doses of CHIR inducing endodermal cell fates predominantly, and higher doses inducing
more posterior (FOXA2 negative) mesodermal fates.

Fig 4 Cell fate patterning does not correlate linearly with CHIR dosage - A *CHIR dose response in 500µm colonies fixed at 48h. (i) Images show the max projections of confocal z-stacks and a z-projection underneath. (ii) Stacked bar plot showing the relative proportions of individual cell populations for each CHIR concentration. (iii) Radial profiles of the percentage of each population. c: number of analysed colonies, n: number of analysed nuclei, shaded area: 95% confidence interval. **B** Heatmaps showing the radial profile of average nuclear marker intensity scaled between 0 and 1 on a per-marker basis. **C** Nanostring analysis of colonies grown in FGF and increasing doses of CHIR harvested at 48h. The colour scale corresponds to the log2 ratio of mRNA count in the sample relative to hESC.*

5- CHIR dosage correlates with distinct downstream NODAL signalling dynamics

Our previous results raise the possibility of a non-linear effect of CHIR concentration on downstream signalling activity which would in turn define cell fate outcomes on micropatterns. We first checked the spatio-temporal dynamics of the canonical WNT pathway using the target gene LEF1 as a proxy (Fig 5A and Bi). We found that LEF1 expression increased monotonically over time at a rate proportional to CHIR concentration and we did not see any clear evidence of LEF1 spatial patterning along the colony radius. These data suggested that all the cells were all equally competent to respond to CHIR regardless of their position in the colony and that they did so proportionally to the dose of CHIR they received, suggesting that additional signals drive radial patterns of cell fates.

Since we and others have found that CHIR induces NODAL signalling on micropatterns (Fig 2 - (Martyn et al., 2018; Massey et al., 2019), we next investigated the spatio-temporal dynamics of NODAL signalling across CHIR concentrations (Fig 5B-D). During the first 24h, the rate of increase in NODAL transcripts correlated with the dose of CHIR. However, while NODAL kept increasing after 24h with 1 and 2µM CHIR, NODAL expression rapidly decreased with 3µM CHIR and dropped even faster with 4µM (Fig 5A and B(ii)). Nuclear SMAD2/3 levels (the downstream signalling effector of the pathway) were consistent with the temporal profiles of NODAL expression. Indeed, nuclear SMAD2/3 first increased proportionally to CHIR concentrations at early time points and then dropped earlier at 3 and 4µM CHIR than at 1 and 2µM CHIR (Fig 5D and B(iii)) confirming the non-linear dependence of NODAL signalling to CHIR dosage.

Interestingly, when looking closely at the spatial pattern of NODAL signalling in the colonies, we observed that NODAL expression was evenly distributed during the first 12h at all CHIR concentrations (Fig 5C), consistently with the notion that all the cells responded to CHIR by activating NODAL expression. However, a radial gradient of NODAL expression starting from the periphery became apparent at 24h in colonies treated with 1 and 2µM CHIR. The NODAL expression domain expanded slightly at 1µM CHIR correlating with the positioning of a few cells with nuclear SMAD2/3 at 36h and 48h (Fig 5D). With 2µM CHIR, NODAL expression expanded to the centre as soon as 36h and became even stronger in the centre at 48h consistently with the presence of cells with nuclear SMAD2/3 across the entire colony at 36h. These results show that a radial gradient of NODAL expression (Fig 5C) precedes the emergence of endoderm at the periphery of colonies treated with 1µM and 2µM CHIR (Fig 2B and 4A). Such a wave of NODAL expression was not clearly apparent at 3µM and 4µM CHIR, perhaps because radial expression occurred too rapidly between 12h and 24h for our experiment to capture this process.

Fig 5 Small variations in CHIR dosage correlate with distinct downstream dynamics of NODAL signalling. *A Cartoons describing the position of image crops shown in B and D-G relative to their respective colony (Left) and the organisation of the bins shown in heatmaps (Middle). Colour sales used in the figure are provided on the right. **B-G** Time course analysis of WNT and NODAL signalling in 500μm hESC colonies across CHIR concentrations. **B and E** show immunofluorescence analysis of*

nuclear LEF1 and nuclear SMAD2/3 levels respectively. Images show selected z-slice across a
*confocal z-stack. **D, F and G** show FISH analysis of NODAL, LEFTY2 and CER1 transcript levels.*
Images are 2D widefield images. For B, and D-G, the radial profiles of signal intensities over time are
*provided below images. **C** shows the temporal profiles or gene expression averaged across entire*
colonies. Lines indicate the mean average expression across colonies the shaded area indicate the
95% confidence interval. The figure is representative of two independent experiments.

Interestingly, nuclear SMAD2/3 did not strictly follow NODAL expression. Furthermore, NODAL expression dynamics (FIG5 Bi) indicated the existence of a negative feedback that occurs faster with higher doses of CHIR. Inhibitors of the pathway include LEFTY2 and CER1 (Aykul et al., 2015). LEFTY2 expression followed closely the spatio-temporal dynamic of NODAL expression (FIG5 E and B (iv)) while CER1 harboured a similar pattern of expression albeit in a temporally delayed manner (FIG5 E and B (v)) supporting the idea that LEFTY2 and CER1 may indeed contribute to the temporal profile of NODAL expression and nuclear SMAD2/3. However, the fact that there exists a decoupling between the level of these inhibitors and the downregulation kinetic of NODAL (with 2µM CHIR for example) implies that other mechanisms are also at play and it will be interesting to elucidate this in the future.

All Together, our results demonstrate that even very small variations in CHIR concentration induce distinct NODAL signalling dynamics which correlate with cell fate outcomes. Importantly, a slow and sustained increase in NODAL expression together with a low expression of LEF1 was found in colonies which formed endoderm. On the other hand, a sharp transient expression of NODAL together with a strong increase in LEF1 was found in colonies where mesoderm was the most abundant. These experiments allowed us to gain insights into the mechanisms driving fate patterning in micropatterned colonies and begin to characterise a system that will be useful for more detailed mechanistic studies in the future.

6- Abrupt Nodal and BMP inhibition is required for the spontaneous emergence of NotoPs

We next turned to the question of how to further modify signalling in order to achieve NotoPs differentiation. Specification of NotoPs requires the cooperation of WNT and NODAL signals in the mouse embryo (Lickert et al., 2002; Vincent et al., 2003; Yamamoto et al., 2001). However, our previous results show that varying WNT activity alone is not sufficient to elicit NotoPs despite the consequences of WNT activity on downstream NODAL signalling. Therefore, we then asked whether

endogenous BMP signalling may be preventing NotoPs emergence in our colonies. BMP signalling
has been shown to inhibit specification of the notochord (Yasuo and Lemaire, 2001) and our
Nanostring data showed that BMP2 and BMP4 are indeed expressed in our system (Fig 2 and Fig
4C), most likely downstream of NODAL signalling (Chhabra et al., 2019; Repina et al., 2023). We
confirmed these results via FISH against BMP2 across CHIR concentrations at 48h (Sup Fig 5). Given
these considerations, we hypothesised that the NODAL and BMP signalling dynamics established
spontaneously within our colonies is inadequate for NotoPs emergence and that a tight exogenous
control of TGFβ signalling together with sustained WNT activity is instead necessary.

To test this ideas, we inhibited NODAL and BMP signalling using small molecule inhibitors added
either throughout differentiation or at 24h when a peak of NODAL expression was observed and when
putative precursors of NotoPs may be present (TBXT/FOXA2 double positive cells were already
present and SOX17+ cells were still absent at 24h). To monitor NotoPs emergence, we used FISH
against the NotoP marker NOTO and against CHRD, a BMP inhibitor expressed in the APS and
rapidly restricted to the node in mice (Bachiller et al., 2000).

As expected, we observed high NODAL expression and an absence of NOTO signal in the control (Fig
6A and B). Interestingly, we also found a broad domain of CHRD transcripts across the colony
probably reflecting cells transiting through an early APS mesendoderm state. Treatment with 0.1μM of
the BMP inhibitor LDN added at 24h had no effect on NOTO but led to an increase in CHRD and
NODAL expression indicating that BMP signalling may be one of the factors negatively feeding back
onto NODAL expression. More importantly, NODAL inhibition from 0h abolished NODAL and resulted
in the presence of rare NOTO/CHRD double positive cells at the periphery of the colony, indicating
that some NotoPs can be specified in this condition. Even more importantly, NODAL inhibition from
24h onwards induced a large domain of strong NOTO and CHRD co-expression localised within a
150μm distance from the colony edge. Addition of LDN at 24h further potentiated this effect
confirming that BMP signalling inhibits NotoPs specification. NotoPs emergence was reproducible
across all colonies within the experiment (Fig 7B) as well as with other cell lines (Sup Fig 6) confirming
the robustness of these results. Furthermore, manipulating colony shape to maximise the proportion of
cells experiencing edge effects such as lines of 200μm width enabled us to generate colonies with
covered almost entirely of NOTO+ cells (Fig 7C)

Fig 6 Timely TGFbeta inhibition is required for the spontaneous emergence of NotoPs on
micropatterns. *A* *FISH staining for NODAL, CHRD and NOTO in 500µm micropattern colonies*

*induced with 2µM CHIR and 20ng/ml FGF2 and fixed at 48h post induction. Timed treatment with the Nodal or BMP inhibitors are indicated. **B** Shows box plots of average signal intensities in each individual colony (left) and heatmaps of binned signal intensities along the radial distance from the edge. Asterisks in box plots indicate the p-values of unpaired t-tests (** p<0.05; *** p<0.001). C Representative widefield images of micropatterned colonies stained via FISH at 48h with SL added at 24h. Yellow arrows indicate a 100µm domain from the edge. D Representative confocal z-stack max projections of colonies stained 2 days post-induction with CHIR and FGF and with or without SB and LDN (SL) treatment added at 0h or 24h. **E** Representative confocal z-stack max projections of colonies grown for 3 days in 2µM CHIR and 20ng/ml FGF containing medium and an additional 3 days in unsupplemented N2B27 with or without SL treatment added at 0h or 24h. **F** Confocal image showing a sagital view of a whole-mount immuno-stained early bud mouse embryo. Notice the SOX9 expression in the crown cells of the node and the nascent notochord. Scale bar: 100µm.*

Staining for SOX17, SOX2, TBXT and FOXA2 in these conditions at 48h (Fig 7D) showed that endoderm (FOXA2/SOX17 double positive cells) was eliminated from colonies treated with SB/LDN added at 24h and that instead the periphery of the colonies contained a large amount of TBXT+/ FOXA2+ and SOX17- cells. This result support the notion that FOXA2+/TBXT+ cells formed at 24h may be competent to form either endoderm or NotoPs depending on whether these cells experience sustained NODAL signalling or an abrupt inhibition respectively.

Finally, to determine whether the NOTO positive cells found at 48h were able to form notochord, we cultured the cells for three days with CHIR and FGF and then three more days in N2B27 alone with or without SB/LDN added at 0h or 24h (Fig 7E). We stained the cells for FOXA2, TBXT and SOX9 which are co-expressed in the node and the nascent notochord in early bud mouse embryos (Fig 7F) as well as SOX17 to distinguish cells that may have formed endoderm. While SOX9 and TBXT were absent from FOXA2/SOX17 co-expressing cells in the control colonies, we found a significant number of FOXA2, TBXT and SOX9 co-expressing cells when TGFbeta inhibitors were added in the medium. This number was further increased when SB and LDN were added at 24h.

All together our data provide evidence that delayed TGFbeta inhibition in micropattern colonies efficiently produce notochord-competent NotoPs.

7- TGFbeta inhibition potentiates WNT signalling response

Finally, we set out to understand the changes in signalling downstream of TGFbeta inhibition which

might explain the specification of NotoPs. We used Nanostring to find differentially expressed genes in

48h colonies with or without the NODAL inhibitor SB added at 24h (Sup Fig 7).

Figure 7 TGFbeta inhibition potentiates WNT signalling responsiveness in a domain that

overlaps with NotoPs emergence. *Time course analysis of LEF1 expression in 500µm hESC*

colonies. The line plot on the left shows the temporal profiles or average nuclear LEF1 intensity within

50µM of the colonies edges. The shaded area indicate the 95% confidence interval. Images on the

right show representative image crops as indicated in Fig 5A. The radial profiles of signal intensities

over time are provided as heatmaps below images.

Interestingly, we observed the upregulation of several NOTCH signalling related genes when NODAL

was inhibited at 24h as well as genes indicating a more anterior identity such as GBX2 and HOXB2.

Importantly, we also found a strong increase in the WNT target gene LEF1. To further elucidate how

WNT signalling is impacted by TGFbeta inhibition and how this may be associated with NotoPs

emergence, we performed a time course analysis of LEF1 expression in colonies treated with FGF

and 2µM CHIR with or without SB and LDN added at 0h or 24h (Fig 7). We also included a 4µM CHIR

condition to get a comparative measure of LEF1 levels in these colonies. Strikingly, when SB and LDN

were added at 0h to colonies treated with 2µM CHIR, LEF1 intensity increased at the same rate as

what was observed for 4µM CHIR treated colonies. When SB and LDN were added at 24h to 2µM

CHIR treated colonies, LEF1 expression sharply increased to reach the levels observed in colonies

treated with 4µM CHIR. These results are in line with our Nanostring data and demonstrate that

TGFbeta inhibition strongly potentiates the cells responsiveness to canonical WNT signalling.

Importantly, LEF1 intensity was the highest at the periphery of the colonies treated with SB and LDN

added at 24h. This domain overlapped with the domain of NOTO+ cells suggesting that high WNT
activity drives NotoPs emergence. Given that 4μM CHIR treatment generates a high level of WNT
activity and a transient peak of NODAL expression (Fig 5) we wondered whether NotoPs could be
obtained by simply inhibiting BMP signalling in 4μM colonies (Sup Fig 8). This, however did not result
in any NOTO positive staining and neither did the addition of SB and LDN at 24h in these colonies.
These results indicate that a tight control of the temporal profiles of WNT and TGFbeta signalling is
needed for NotoPs emergence.
561
562 Together our data clarify how WNT and TGFbeta signalling cooperate in order to specify the
563 notochordal lineage. While WNT and TGFbeta signalling are initially necessary to induce an early APS
564 cell state, a timely and abrupt inhibition of TGFbeta signalling is necessary to maximise WNT
565 signalling response and define NotoPs.
566
567 **Discussion**
568
The series of lineage restrictions that occur in the APS during gastrulation remain challenging to
investigate *in vivo* and this is especially true in a human context. Here, using spatial confinement with
micropatterns, we were able to direct the development of hESC into reproducible radial patterns of all
the APS cell fates including NotoPs (Colombier et al., 2020). This system enabled us to gain insights
into 1) the mechanisms that regulate temporal patterns of signalling cues and 2) to delineate the
signalling sequences distinguishing APS cell fates from one another (Figure 8).
575

576
577 **Figure 8 Graphical summary of cell fate outcomes and associated signalling dynamics. *A***
578 *Diagram summarising the spatial organisation of cell fates at 48h on micropatterns when treated with*
579 *20ng/ml FGF and with 1μM CHIR (i), 2μM CHIR (ii, iv, v), 3μM CHIR (iii) and SB and LDN added at*

*24h (iv) or 0h (v). THE WNT and NODAL signalling profiles associated with these patterns of differentiation is indicated as a line plot underneath each colony. **B** shows the putative hierarchy of cell fate lineages and the signalling conditions defining lineage restrictions.*

Our first experiment showed that cells on micropatterns treated with 2µM CHIR and 20ng/ml FGF2 predominantly underwent endodermal differentiation on micropatterns (Fig 1B). This was at first surprising to us given that the same concentrations of CHIR and FGF in 2D monolayer cultures normally lead to homogenous NMP differentiation (Fig 1B). This also raised the question of what makes the cells on micropatterns adopt a distinct fate than in monolayer cultures.

Previous work using micropatterns and BMP or WNT ligands as the triggers of differentiation have shown that cells located at the colony boundary respond more efficiently to signalling molecules than cells in the centre due to a loss of epithelial integrity on the colony edges (Etoc et al., 2016; Legier et al., 2023; Martyn et al., 2018; Martyn et al., 2019; Warmflash et al., 2014). Our data indicate that a similar boundary-driven mechanism is likely taking place here as well given that the size of the peripheral differentiation domain remained constant with increasing colony sizes (Sup Fig 2). In our case however, the cells were treated with CHIR, a cell-permeable molecule that bypasses ligand/receptor interactions and is therefore unlikely to be affected by a loss of epithelial integrity. Our LEF1 staining showed that all the cells were indeed equally responsive to CHIR (Figure 5A). Nevertheless differential responsiveness to ligands may matter when secondary signals downstream of CHIR are secreted. Our data support this idea as NODAL expression was initiated throughout the colony at early time points (as a result of equal responsiveness to CHIR) before a wave of NODAL expression travelling from the periphery towards the centre became apparent (Figure 5D). Such a wave, which was also described in other systems (Heemskerk et al., 2019; Liu et al., 2022; Martyn et al., 2019) may be the result of the positive self-regulation of NODAL (Liu et al., 2022) together with a high responsiveness to NODAL at the periphery. As NODAL signalling is a notorious driver of the endodermal lineage (Nowotschin et al., 2019) and NODAL inhibition prevents endoderm emergence (Fig 3), we conclude that confinement potentiates NODAL signalling downstream of CHIR which in turn drives endodermal differentiation on micropatterns.

Our data are in agreement with and complementary to a recent preprint showing that treatment of micropatterned colonies with WNT3a and FGF8 ligands generates a similar spatio-temporal pattern of NODAL expression (Fig5J of (Ortiz-Salazar et al., 2024)). Both systems ultimately lead to the formation of reproducible and self-organised 3-dimensional structures composed of definitive endoderm with a core of epiblast cells as shown in Fig 1F. These micropattern systems therefore

provide a powerful platform - complementary to other 3D human in vitro models (Moris et al., 2020;
Vianello and Lutolf, 2020) - to further understand endoderm specification and morphogenesis in a
human context.

Importantly, we found that endoderm only emerges on micropatterns in a very small CHIR
concentration range around 2μM. Our CHIR dose-response experiment (Fig 4) was particularly
revealing and helped us to delineate the signalling sequences that segregate individual APS cell fates
from one another (summarised in Fig 8). We show that even very small variations in CHIR
concentrations radically change the proportion of endodermal and mesodermal cell fates (Fig 4 and
Fig 8) and that this effect was mediated by a non linear relationship between CHIR concentration and
the downstream dynamics of NODAL expression (Fig5C-E). 1 and 2μM CHIR induced a progressive
and sustained increase in NODAL production favouring the endodermal lineage while higher CHIR
concentrations resulted in a sharp increase followed by a rapid reduction of NODAL expression which
correlated with an abundance of mesoderm (Fig 4, 5 and 8). It is useful to compare these findings with
the results from Ortiz-salazar et al. These authors reported that a 10-fold increase in WNT3a ligand
concentration did not alter endoderm emergence while CHIR (10μM) induced mesoderm
differentiation. Overall, our data are in agreement and complementary and may identify the range of
CHIR concentrations that relate to the physiological, ligand-based activation of the pathway.

Our findings also suggest the existence of a negative feedback on NODAL expression when NODAL
reaches high levels of expression (Fig 5B). Secreted antagonists of the pathway, LEFTY2 and CER1
may contribute to the downregulation of NODAL at high CHIR concentrations but are unlikely to be the
sole negative regulators of NODAL given the decoupling between the level of expression of these
antagonists and the kinetic of NODAL downregulation (Fig 5). Furthermore, we found heterogeneous
levels of nuclear SMAD2/3 in individual colonies with domains of NODAL expression that did not
overlap necessarily with nuclear SMAD2/3 (Fig 5D and E). One possible reason include an
antagonism between NODAL and BMP. Indeed, BMP has been shown to antagonise NODAL in other
contexts (see for example (Pereira et al., 2012)) and we found here that higher CHIR dosages
correlated with higher levels of BMP2 transcripts (Sup Fig 5). It will be interesting to use our
micropattern system to elucidate this question in the future.

Most importantly, our data sheds some light on how distinct signalling regimes are established and
associated with individual cell fates (Fig 8B):

1) We have found that NMP specification requires an environment where NODAL activity is
maintained at a minimum (Fig 3). This idea is in agreement with a recent report showing that the
maintenance of NMPs *in vitro* requires NODAL inhibition (Kelle et al., 2024). This notion is also
compatible with work in mouse embryos showing that NMPs appear later than endoderm, at a time
when NODAL signalling activity starts to decrease (Lawson et al., 1991). Furthermore, neighbouring
NotoPs - as a source of NODAL inhibitors - may protect NMPs from advert differentiation during axial
elongation. Indeed a release of NODAL inhibition might explain why NotoPs ablation results in early
termination of axial elongation (Abdelkhalek et al., 2004; McLaren and Steventon, 2021; Saunders et
al., 2024; Wymeersch et al., 2019).

2) Our observations also show that a proportion of the cells differentiate to lateral plate mesoderm (Fig
1F), most likely under the influence of endogenous BMP signalling (Heemskerk et al., 2019). This is
perhaps not surprising as fate mapping experiments have shown that cardiac mesoderm arise directly
adjacent and posterior to the definitive endoderm (Lawson et al., 1991; Tam et al., 1997) and lineage
tracing using FOXA2-cre mouse lines have demonstrated that a proportion of FOXA2 expressing cells
in the streak are fated to form cardiac ventricles (Bardot et al., 2017). It will be interesting in the future
to confirm the lineage hierarchy and the identity of the HAND1+ cells emerging in our system.

3) Finally, our manipulations of NODAL and BMP signalling has enabled us to identify the signalling
requirements for the specification of the notochord lineage (Fig 6A and 8B) and unveiled a signalling
cross-talk where NODAL signalling dampens WNT activity (Fig 7). We propose a model where initial
WNT and NODAL signalling combinatorially specify epiblast cells to an early multipotent APS state
(co-expressing TBXT and FOXA2) which can commit to either endoderm or notochord. While
prolonged NODAL exposure keeps WNT activity low and drives endoderm differentiation, a sharp
inhibition of NODAL signalling potentiates WNT activity and redirects the cells to the NotoP fate (Fig
8B). Our results and model are consistent with observations in the Xenopus where a sudden drop of
p-Smad2 correlates with the emergence of the notochord (Schohl and Fagotto, 2002) but it will be
important to test our model in an *in vivo* setting and determine mechanisms that are evolutionary
conserved or that differ in the human.

Reassuringly, our data are reproducible across multiple cell lines (Sup Fig 6) and consistent with a
recent study reporting a similar requirement for delayed TGFbeta signalling inhibition for the derivation
of NotoPs on micropatterns (Rito et al., 2023). These authors also report a cross-talk between hippo
signalling and FGF activity at the colony boundary overlapping with the domain of NotoPs emergence.

Our data showing that TGFbeta inhibition potentiates WNT activity in this domain nicely complement
these findings. It will be important in the future to elucidate whether and how the specific mechanical
and biochemical environment defined at the colony boundaries *in vitro* are defined inside the embryo.

Conclusion

NotoPs are regarded as a promising cell type for drug discovery or cell therapy and much remains to
be learned about the healthy and pathological development of the notochord. Encouragingly, recent
evidence suggest that NotoPs persist as a small transcriptionally stable population throughout axial
elongation (Wymeersch et al., 2019) making it likely that NotoPs may be expandable in culture if the
correct conditions can be identified. Our work together with the recent work from Rito and colleagues
(Rito et al., 2023) provides insights that will inform the development of reliable NotoPs derivation
protocols. It will be interesting to test if a fully functional node can be reproduced *in vitro* and
determine what other signalling cues define the proportions and maturation of individual node sub-
populations in the future.

Materials and Methods

Cell culture

Experiments were conducted with the MasterShef7 hESC line obtained from the University of
Sheffield, the RC17 hESC line from the University of Edinburgh (De Sousa et al) and the NAS2 hIPSC
line from the Kunath lab (Devine et al). Approval for the use of hESC was granted by the steering
committee for the UK Stem Cell bank and for the Use of Stem Cell Lines. Cell lines were propagated
at 37°C and 5% CO_2 in mTSER Plus medium (100-0276, Stemcell Technologiess) on Geltrex
(A1413302, Life Technologies) coated 6-well plates (3516, Corning Incorporated). Wells were coated
for 30 minutes at 37°C using a 100µg/mL Geltrex solution diluted in Magnesium and Calcium
containing DPBS (14080-048, GIBCO). Passaging was performed every 2 to 3 days using Accutase
(00-4555-56, ThermoFisher Scientific). The cells were Mycoplasma tested prior to running
experiments. Only cells with a passage number below 30 were used for all experiments.

Micropattern fabrication

Micropatterning was done in Ibidi 8-well µSlides (IB-80826, Ibidi) with a PRIMO bioengineering
plateform (Alveole) mounted on a Nikon Ti2 Widefield microscope using a protocol adapted from
Alveole: The surface of ibidi slides was first passivated as follows: slides were plasma treated (Harrick
Plasma Cleaner) for 1 min 30 seconds at high intensity in a 200 mTorr vacuum and then incubated at

room temperature for 1h with 200µL/well of 200 µg/mL Poly-D-Lysine (A-003-E, Merck Millipore)
diluted in 0.1M HEPES pH 8.4 (H3784-25G, Sigma-Aldrich). The wells were washed twice with ddH$_2$O
and once with 0.1M HEPES pH 8.4 and then incubated for 1h in the dark with 125µL/well of 80mg/mL
mPEG-SVA (MPEG-SVA-5000, Laysan Bio) freshly dissolved in 0.1M HEPES pH 8.4. Slides were
then washed profusely with ddH$_2$O, air dried and stored at 4°C until further processing (1 week
maximum). The PRIMO insolation step was next performed less than one day prior to plating the cells:
passivated wells were covered with 8µL PLPP gel (Cairn Research, 1µL PLPP gel/well diluted with
7µL 70% Ethanol) in the dark and left to dry for ~30min at room temperature. Slides were then
insolated with PRIMO through a 20X lens with a dose of 50mJ/cm^2. All micropattern shapes were
designed in Inkscape and converted to binary tiff files in ImageJ prior to loading in the Alveole
Leonardo software. After insolation, PLPP gel was removed with 3 ddH2O washes and the slides were
air dried and stored at 4°C until use.
730

Culture on micropatterns

Micropatterned Ibidi slides were first rehydrated for 5 min in Magnesium and Calcium containing
DPBS (14080-048, GIBCO; thereafter DPBS++). Matrix coating was then performed by incubating the
wells at room temperature for 30 min on a rocker with a mixture of 40µg/mL rhVitronectin-N (A14700,
ThermoFisher) and 10µg/mL rhLaminin521 (A29249, GIBCO) diluted in DPBS++. Wells were washed
3 times with DPBS++ and left in the last wash whilst preparing the cells for seeding to ensure that the
wells were not left to dry. For seeding, 80% confluent cells were dissociated to single cells with
Accutase, and resuspended in seeding medium composed of mTESR Plus supplemented with 10µM
Y-27632 (1254, Tocris Bio-Techne) and 1:100 Penicillin/Streptomycin (10,000U/mL pen, 10,000mg/mL
strep; 15140-122, Invitrogen). Cells were plated onto micropatterns at a density of 150 000 cells/well
in 250µL of seeding medium. The cells were left to adhere for 3h at 37°C. After attachment, the excess
of cells was removed by gentle pipetting and applying fresh seeding medium. The cells were left to
settle and cover the patterns overnight until induction of differentiation the next morning. Cells were
washed once in N2B27 to remove traces of growth factors present in seeding medium. Differentiation
was then induced using N2B27 medium supplemented with Penicillin/Streptomycin (1:100), CHIR
99021 at concentrations indicated in figures legends (4423/10, Tocris Bio-Techne), 20ng/ml human
bFGF (PHG6015, ThermoFisher Scientific). SB 431542 (1614, Tocris Bio-Techne) was used at 10µM
unless specified otherwise and LDN 193189 (72147, StemCell Technologies) at 0.1µM.
749

Immunofluorescence

Mouse embryos were staged according to (Downs and Davies, 1993) and stained as described
previously in (Wong, 2021). Cells grown on micropatterns were fixed with 4% PFA (CHE2036,
Scientific Laboratory Supplies), washed 3 times with a solution of PBS and 0.1% Triton X-100

(A16046, Alfa Aesar; hereafter PBST) and left overnight at 4°C in blocking solution containing 3%
Donkey Serum (D9663, Merck) and 0.03% Sodium Azide (40-2000-01, Severn Biotech Ltd) in PBST.
All primary antibodies were incubated overnight at 4°C and secondary antibodies at room temperature
for 3h. All washing steps were performed in PBST. Some co-staining required the use of primary
antibodies raised in the same species. In these cases, the staining was performed with either pre-
conjugated antibodies only, or sequentially using a non-conjugated antibody first, its corresponding
secondary antibody next, followed by a blocking step using species-specific serum (3% Goat Serum
(G9023-10ML, Sigma) or 3% Rabbit Serum (R9133, Merck)) and finally applying the conjugated
antibody. Slide were washed 3 times in PBST and kept sealed at 4°C until imaging.
763
764
765

Primary Antibody	Cat. no, supplier	Dilution
SOX2	14-9811-82, ThermoFisher Scientific	1:200
TBX6	AF4744, R&D Systems	1:400
PAX6	561462, BD Bioscience	1:300
TBXT	NL2085R Conjugated, R&D Systems	1:300
SOX17	AF1924, R&D Systems	1:200
CDX2	Ab76541, Abcam	1:200
OCT4	AP3724A, Abcepta	1:200
FOXA2	8186, Cell Signalling	1:400
SMAD2/3	610842, BD Bioscience	1:100
SOX9	Ab185230 Conjugated, Abcam	1:250
CDH1	SIGMA, MABT26	1:100
CDH2	BD Bioscience, 610182	1:200

766
767

Secondary Antibody	Cat. no, supplier	Dilution
Donkey-Anti-Goat-405	Abcam, ab175665	1:1000

Donkey-Anti-Mouse-405	Abcam, ab175658	1:1000
Donkey-Anti-Rat-488	Abcam, ab150153	1:1000
Donkey-Anti-Mouse-488	Abcam, ab150105	1:1000
Donkey-Anti-Rabbit-A568	Abcam, ab175470	1:1000
Donkey-Anti-Mouse-568	Abcam, ab175472	1:1000
Donkey-Anti-Rabbit-647	Abcam, ab150075	1:1000
Donkey-Anti-Rat-647	Abcam, ab150155	1:1000
Donkey-Anti-Mouse-647	Abcam, ab150107	1:1000

Fluorescent In Situ Hybridization

Branched DNA FISH was performed using the viewRNA Cell Plus assay from ThermoFisher Scientific (88-19000-99) according to the manufacturer's protocol. FISH probes used in this study are listed in the table below. Slides were imaged on the same day with a Nikon TiE widefield inverted microscope and a dry 20X lens.

FISH Probe-Channel	Cat.no
NOTO-488	VA4-3101015-VC
CHRD-546	VA1-3005411-VC
NODAL-647	VA6-3167894-VC
LEFTY2-546	VA1-3004742-VC
DKK1-488	VA4-3082344-VC
CER1-488 BMP2-647	VA4-3089614-VC VA6-3168104-VC

Immunofluorescence Imaging and Image analysis

Embryos were imaged in PBST using a Leica TCS SP8 Confocal and a 25X water immersion lens. All images were annotated and contrast-adjusted using FIJI (Schindelin et al., 2012). Micropatterned colonies were imaged with an Opera Phenix Plus (Perkin Elmer). Ibidi slides contained around 64 colonies per well, out of which ~20 colonies were selected for analysis in each experiment. To ensure

an unbiased sampling of the colonies, the slides were first fully scanned with a 10x lens to generate
overview images of the LMBR (nuclear envelope marker) signal only. These images were then
processed with an automated pipeline in the harmony software (Perkin Elmer). This pipeline rejected
colonies with an unexpected area or roundness and then randomly sampled 20 colonies from the pool
of valid colonies. Sampled colonies were next imaged with a 20X lens to generate 3D multichannel z-
stacks with voxel size of 0.59 x 0.59 x 1µm. Opera images were then exported as Tiff files for further
analysis. Nuclear segmentation was performed on the LMBR signal as described previously (Blin et
al., 2019). Raw images and nuclear masks were imported into Pickcells
(https://pickcellslab.frama.io/docs/) to compute nuclear features including 3D spatial coordinates and
average intensities in all fluorescence channels. The tsv file created in PickCells was then analysed in
python. Our Jupyter Notebooks and data files are available in our Gitlab repository
(https://framagit.org/pickcellslab/data/2023_haxioms).

FISH Imaging and Image analysis

Micropatterened colonies were imaged on an inverted widefield microscope (Nikon Eclipse Ti) with a
long distance 10x lens to image entire wells (8mm x 8mm). Images were were then segmented using
CellPose 2.0 (Stringer et al., 2021) to identify individual colonies. We used the livecell model as a
starting point and manually adjusted the model according to the CellPose tutorial using at least 5
different images to ensure accurate segmentation across our conditions. Images were scaled down for
the segmentation process to save on computational resources and scaled back up to their original
sizes using ImageJ prior to loading into PickCells for further analysis. PickCells was used to compute
average intensities in each channel within each colony, as well as background intensity in the
immediate vicinity of the colony (bounding box intensity – colony intensity) and radial intensities as
described in Fig 5. Data were next analysed in python. The colonies were selected based on their
compactness and surface area, to avoid including abnormally sized and shaped colonies (e.g. partially
broken colonies). The background intensity was subtracted from the average and radial colony
intensities and plotted in python.

Nanostring analysis

RNA samples were prepared using an Absolutely RNA microprep kit (cat.no 400805, Agilent
technologies) and Nanostring profiling was performed using nCounter technology as per the
manufacturer's instructions. We used a panel of probes consisting of the 780 genes included in the
standard human embryonic stem cell gene panel together with 30 additional custom probes (all genes
are listed in Sup Table 1). Normalisation of raw data was accomplished in the Nanostring dedicated

nCounter software. Next, raw counts were imported in R (R Core Team, 2013) and analysed with the

Bioconductor package moanin (Varoquaux and Purdom, 2020). We first applied an initial cut-off to

filter out all the genes where the max count was below 100. The data was then log2 transformed. We

kept only the top 50% most variable genes based on the median absolute deviation (mad) metric over

time. A spline was then fitted onto each individual gene profile and we grouped genes into 7 clusters

using kmeans clustering on the parameters of the fitted splines to obtain the heatmaps shown in Fig

2A. R scripts and data are available in our Gitlab repository

(https://framagit.org/pickcellslab/data/2023_haxioms).

Mouse husbandry

Mouse work was carried out under the UK Home Office project license PPL PEEC9E359, approved by

the University of Edinburgh Animal Welfare and Ethical Review Panel and in compliance with the

Animals (Scientific Procedures) Act 1986. Mice used were of C57Bl6/J strain background. The mice

were kept under a 12 h light/dark cycle, and the embryo age was denoted day 0 on the midpoint of the

dark cycle the day the plug was found.

Funding:

This work was funded by a BBSRC project grant to GB ref:BB/W002310/1, a BBSRC Alert equipment

grant to GB ref: BB/T017961/1 and a WT ISSF3 award to GB ref: IS3-R1.16 19/20. MRG is funded by

a BBSRC EASTBIO studentship. EO was funded by a BSDB Gurdon studentship.

Acknowledgements:

We thank Tilo Kunath for the provision of RC17 and NAS2 cell lines. We also thank Sally Lowell,

Anestis Tsakiridis and Anne Camus for their critical review of the manuscript. We also thank Val Wilson

for helping us with mouse embryos. We thank Justyna Cholewa Waclaw and Matthieu Vermeren for

their help with imaging and Alison Munro for running the Nanostring samples.

References

Abdelkhalek, H. B., Beckers, A., Schuster-Gossler, K., Pavlova, M. N., Burkhardt, H., Lickert, H., Rossant, J., Reinhardt, R., Schalkwyk, L. C., Müller, I., et al. (2004). The mouse homeobox gene Not is required for caudal notochord development and affected by the truncate mutation. *Genes Dev.* **18**, 1725–1736.

Aksoy, I., Giudice, V., Delahaye, E., Wianny, F., Aubry, M., Mure, M., Chen, J., Jauch, R., Bogu, G. K., Nolden, T., et al. (2014). Klf4 and Klf5 differentially inhibit mesoderm and endoderm differentiation in embryonic stem cells. *Nat Commun* **5**, 3719.

Albors, A. R., Halley, P. A. and Storey, K. G. (2018). Lineage tracing of axial progenitors using Nkx1-2CreERT2 mice defines their trunk and tail contributions. *Development* **145**,.

Ang, S. L. and Rossant, J. (1994). HNF-3 beta is essential for node and notochord formation in mouse development. *Cell* **78**, 561–574.

Arnold, S. J., Stappert, J., Bauer, A., Kispert, A., Herrmann, B. G. and Kemler, R. (2000). Brachyury is a target gene of the Wnt/beta-catenin signaling pathway. *Mech. Dev.* **91**, 249–258.

Avilion, A. A., Nicolis, S. K., Pevny, L. H., Perez, L., Vivian, N. and Lovell-Badge, R. (2003). Multipotent cell lineages in early mouse development depend on SOX2 function. *Genes Dev* **17**, 126–140.

Aykul, S., Ni, W., Mutatu, W. and Martinez-Hackert, E. (2015). Human Cerberus Prevents Nodal-Receptor Binding, Inhibits Nodal Signaling, and Suppresses Nodal-Mediated Phenotypes. *PLoS One* **10**, e0114954.

Bachiller, D., Klingensmith, J., Kemp, C., Belo, J. A., Anderson, R. M., May, S. R., McMahon, J. A., McMahon, A. P., Harland, R. M., Rossant, J., et al. (2000). The organizer factors Chordin and Noggin are required for mouse forebrain development. *Nature* **403**, 658–661.

Bagheri-Fam, S., Barrionuevo, F., Dohrmann, U., Günther, T., Schüle, R., Kemler, R., Mallo, M., Kanzler, B. and Scherer, G. (2006). Long-range upstream and downstream enhancers control distinct subsets of the complex spatiotemporal Sox9 expression pattern. *Developmental Biology* **291**, 382–397.

Bagnat, M. and Gray, R. S. (2020). Development of a straight vertebrate body axis. *Development* **147**,.

Bakker, B. S. de, Jong, K. H. de, Hagoort, J., Bree, K. de, Besselink, C. T., Kanter, F. E. C. de, Veldhuis, T., Bais, B., Schildmeijer, R., Ruijter, J. M., et al. (2016). An interactive three-dimensional digital atlas and quantitative database of human development. *Science* **354**, aag0053.

Balmer, S., Nowotschin, S. and Hadjantonakis, A.-K. (2016). Notochord morphogenesis in mice: Current understanding & open questions. *Developmental Dynamics* **245**, 547–557.

Bardot, E., Calderon, D., Santoriello, F., Han, S., Cheung, K., Jadhav, B., Burtscher, I., Artap, S., Jain, R., Epstein, J., et al. (2017). Foxa2 identifies a cardiac progenitor population with ventricular differentiation potential. *Nat Commun* **8**, 14428.

Beccari, L., Moris, N., Girgin, M., Turner, D. A., Baillie-Johnson, P., Cossy, A.-C., Lutolf, M. P., Duboule, D. and Arias, A. M. (2018). Multi-axial self-organization properties of mouse embryonic stem cells into gastruloids. *Nature* **562**, 272–276.

Beddington, R. S. and Robertson, E. J. (1989). An assessment of the developmental potential of embryonic stem cells in the midgestation mouse embryo. *Development* **105**, 733–737.

Belo, J. A., Bouwmeester, T., Leyns, L., Kertesz, N., Gallo, M., Follettie, M. and De Robertis, E. M. (1997). Cerberus-like is a secreted factor with neutralizing activity expressed in the anterior primitive endoderm of the mouse gastrula. *Mech. Dev.* **68**, 45–57.

Ben-Haim, N., Lu, C., Guzman-Ayala, M., Pescatore, L., Mesnard, D., Bischofberger, M., Naef, F., Robertson, E. J. and Constam, D. B. (2006). The nodal precursor acting via activin receptors induces mesoderm by maintaining a source of its convertases and BMP4. *Dev. Cell* **11**, 313–323.

Bergmann, S., Penfold, C. A., Slatery, E., Siriwardena, D., Drummer, C., Clark, S., Strawbridge, S. E., Kishimoto, K., Vickers, A., Tewary, M., et al. (2022). Spatial profiling of early primate gastrulation in utero. *Nature* **609**, 136–143.

Blin, G. (2021). Quantitative developmental biology in vitro using micropatterning. *Development* **148**,.

Blin, G., Sadurska, D., Migueles, R. P., Chen, N., Watson, J. A. and Lowell, S. (2019). Nessys: A new set of tools for the automated detection of nuclei within intact tissues and dense 3D cultures. *PLOS Biology* **17**, e3000388.

Burtscher, I. and Lickert, H. (2009). Foxa2 regulates polarity and epithelialization in the endoderm germ layer of the mouse embryo. *Development* **136**, 1029–1038.

C, M., G, C., Ka, J., Ac, H., Gg, S. and J, N. (2018). Oct4 regulates the embryonic axis and coordinates exit from pluripotency and germ layer specification in the mouse embryo. *Development (Cambridge, England)* **145**,.

Cadigan, K. M. and Waterman, M. L. (2012). TCF/LEFs and Wnt Signaling in the Nucleus. *Cold Spring Harb Perspect Biol* **4**, a007906.

Cambray, N. and Wilson, V. (2002). Axial progenitors with extensive potency are localised to the mouse chordoneural hinge. *Development* **129**, 4855–4866.

Cambray, N. and Wilson, V. (2007). Two distinct sources for a population of maturing axial progenitors. *Development* **134**, 2829–2840.

Cermola, F., D'Aniello, C., Tatè, R., Cesare, D. D., Martinez-Arias, A., Minchiotti, G. and Patriarca, E. J. (2021). Gastruloid Development Competence Discriminates Different States of Pluripotency. *Stem Cell Reports* **16**, 354–369.

Chhabra, S., Liu, L., Goh, R., Kong, X. and Warmflash, A. (2019). Dissecting the dynamics of signaling events in the BMP, WNT, and NODAL cascade during self-organized fate patterning in human gastruloids. *PLOS Biology* **17**, e3000498.

Choi, K.-S., Cohn, M. J. and Harfe, B. D. (2008). Identification of nucleus pulposus precursor cells and notochordal remnants in the mouse: Implications for disk degeneration and chordoma formation. *Developmental Dynamics* **237**, 3953–3958.

Cohen, P. and Goedert, M. (2004). GSK3 inhibitors: development and therapeutic potential. *Nat Rev Drug Discov* **3**, 479–487.

Colombier, P., Halgand, B., Chédeville, C., Chariau, C., François-Campion, V., Kilens, S., Vedrenne, N., Clouet, J., David, L., Guicheux, J., et al. (2020). NOTO Transcription Factor Directs Human Induced Pluripotent Stem Cell-Derived Mesendoderm Progenitors to a Notochordal Fate. *Cells* **9**, 509.

Costello, I., Pimeisl, I.-M., Dräger, S., Bikoff, E. K., Robertson, E. J. and Arnold, S. J. (2011). The T-box transcription factor Eomesodermin acts upstream of Mesp1 to specify cardiac mesoderm during mouse gastrulation. *Nat Cell Biol* **13**, 1084–1091.

Costello, I., Nowotschin, S., Sun, X., Mould, A. W., Hadjantonakis, A.-K., Bikoff, E. K. and Robertson, E. J. (2015). Lhx1 functions together with Otx2, Foxa2, and Ldb1 to govern anterior mesendoderm, node, and midline development. *Genes Dev* **29**, 2108–2122.

Diaz-Hernandez, M. E., Khan, N. M., Trochez, C. M., Yoon, T., Maye, P., Presciutti, S. M., Gibson, G. and Drissi, H. (2020). Derivation of notochordal cells from human embryonic stem cells reveals unique regulatory networks by single cell-transcriptomics. *Journal of Cellular Physiology* **235**, 5241–5255.

Dunn, N. R., Vincent, S. D., Oxburgh, L., Robertson, E. J. and Bikoff, E. K. (2004). Combinatorial activities of Smad2 and Smad3 regulate mesoderm formation and patterning in the mouse embryo. *Development* **131**, 1717–1728.

Edri, S., Hayward, P., Jawaid, W. and Arias, A. M. (2019). Neuro-mesodermal progenitors (NMPs): a comparative study between pluripotent stem cells and embryo-derived populations. *Development* **146**, dev180190.

Etoc, F., Metzger, J., Ruzo, A., Kirst, C., Yoney, A., Ozair, M. Z., Brivanlou, A. H. and Siggia, E. D. (2016). A Balance between Secreted Inhibitors and Edge Sensing Controls Gastruloid Self-Organization. *Developmental Cell* **39**, 302–315.

Forlani, S., Lawson, K. A. and Deschamps, J. (2003). Acquisition of Hox codes during gastrulation and axial elongation in the mouse embryo. *Development* **130**, 3807–3819.

Frith, T. J. R. and Tsakiridis, A. (2019). Efficient Generation of Trunk Neural Crest and Sympathetic Neurons from Human Pluripotent Stem Cells Via a Neuromesodermal Axial Progenitor Intermediate. *Current Protocols in Stem Cell Biology* **49**, e81.

Frith, T. J., Granata, I., Wind, M., Stout, E., Thompson, O., Neumann, K., Stavish, D., Heath, P. R., Ortmann, D., Hackland, J. O., et al. (2018). Human axial progenitors generate trunk neural crest cells in vitro. *eLife* **7**, e35786.

Funa, N. S., Schachter, K. A., Lerdrup, M., Ekberg, J., Hess, K., Dietrich, N., Honoré, C., Hansen, K. and Semb, H. (2015). β-Catenin Regulates Primitive Streak Induction through Collaborative Interactions with SMAD2/SMAD3 and OCT4. *Cell Stem Cell* **16**, 639–652.

Gouti, M., Tsakiridis, A., Wymeersch, F. J., Huang, Y., Kleinjung, J., Wilson, V. and Briscoe, J. (2014). In Vitro Generation of Neuromesodermal Progenitors Reveals Distinct Roles for Wnt Signalling in the Specification of Spinal Cord and Paraxial Mesoderm Identity. *PLOS Biology* **12**, e1001937.

Hart, A. H., Hartley, L., Sourris, K., Stadler, E. S., Li, R., Stanley, E. G., Tam, P. P. L., Elefanty, A. G. and Robb, L. (2002). Mixl1 is required for axial mesendoderm morphogenesis and patterning in the murine embryo. *Development* **129**, 3597–3608.

Hart, A. H., Hartley, L., Ibrahim, M. and Robb, L. (2004). Identification, cloning and expression analysis of the pluripotency promoting Nanog genes in mouse and human. *Developmental Dynamics* **230**, 187–198.

Heemskerk, I., Burt, K., Miller, M., Chhabra, S., Guerra, M. C., Liu, L. and Warmflash, A. (2019). Rapid changes in morphogen concentration control self-organized patterning in human embryonic stem cells. *eLife* **8**, e40526.

Henrique, D., Abranches, E., Verrier, L. and Storey, K. G. (2015). Neuromesodermal progenitors and the making of the spinal cord. *Development* **142**, 2864–2875.

Kanai-Azuma, M., Kanai, Y., Gad, J. M., Tajima, Y., Taya, C., Kurohmaru, M., Sanai, Y., Yonekawa, H., Yazaki, K., Tam, P. P. L., et al. (2002). Depletion of definitive gut endoderm in Sox17-null mutant mice. *Development* **129**, 2367–2379.

Kelle, D., Ugur, E., Rusha, E., Shaposhnikov, D., Livigni, A., Horschitz, S., Davoudi, M., Blutke, A., Bushe, J., Sterr, M., et al. (2024). Capture of Human Neuromesodermal and Posterior Neural Tube Axial Stem Cells. 2024.03.26.586760.

Kinder, S. J., Tsang, T. E., Wakamiya, M., Sasaki, H., Behringer, R. R., Nagy, A. and Tam, P. P. L. (2001). The organizer of the mouse gastrula is composed of a dynamic population of progenitor cells for the axial mesoderm. *Development* **128**, 3623–3634.

Koch, F., Scholze, M., Wittler, L., Schifferl, D., Sudheer, S., Grote, P., Timmermann, B., Macura, K. and Herrmann, B. G. (2017). Antagonistic Activities of Sox2 and Brachyury Control the Fate Choice of Neuro-Mesodermal Progenitors. *Developmental Cell* **42**, 514-526.e7.

Kwon, G. S., Viotti, M. and Hadjantonakis, A.-K. (2008). The Endoderm of the Mouse Embryo Arises by Dynamic Widespread Intercalation of Embryonic and Extraembryonic Lineages. *Developmental Cell* **15**, 509–520.

Lawson, K. A., Meneses, J. J. and Pedersen, R. A. (1991). Clonal analysis of epiblast fate during germ layer formation in the mouse embryo. *Development* **113**, 891–911.

Legier, T., Rattier, D., Llewellyn, J., Vannier, T., Sorre, B., Maina, F. and Dono, R. (2023). Epithelial disruption drives mesendoderm differentiation in human pluripotent stem cells by enabling TGF-β protein sensing. *Nat Commun* **14**, 349.

Lickert, H., Kutsch, S., Kanzler, B., Tamai, Y., Taketo, M. M. and Kemler, R. (2002). Formation of multiple hearts in mice following deletion of beta-catenin in the embryonic endoderm. *Dev Cell* **3**, 171–181.

Liu, L., Nemashkalo, A., Rezende, L., Jung, J. Y., Chhabra, S., Guerra, M. C., Heemskerk, I. and Warmflash, A. (2022). Nodal is a short-range morphogen with activity that spreads through a relay mechanism in human gastruloids. *Nat Commun* **13**, 497.

Lolas, M., Valenzuela, P. D. T., Tjian, R. and Liu, Z. (2014). Charting Brachyury-mediated developmental pathways during early mouse embryogenesis. *PNAS* **111**, 4478–4483.

Martins, J.-M. F., Fischer, C., Urzi, A., Vidal, R., Kunz, S., Ruffault, P.-L., Kabuss, L., Hube, I., Gazzerro, E., Birchmeier, C., et al. (2020). Self-Organizing 3D Human Trunk Neuromuscular Organoids. *Cell Stem Cell* **26**, 172-186.e6.

Martyn, I., Kanno, T. Y., Ruzo, A., Siggia, E. D. and Brivanlou, A. H. (2018). Self-organization of a human organizer by combined Wnt and Nodal signalling. *Nature* **558**, 132–135.

Martyn, I., Brivanlou, A. H. and Siggia, E. D. (2019). A wave of WNT signaling balanced by secreted inhibitors controls primitive streak formation in micropattern colonies of human embryonic stem cells. *Development* **146**,.

Massey, J., Liu, Y., Alvarenga, O., Saez, T., Schmerer, M. and Warmflash, A. (2019). Synergy with TGFβ ligands switches WNT pathway dynamics from transient to sustained during human pluripotent cell differentiation. *PNAS* **116**, 4989–4998.

McCann, M. R., Tamplin, O. J., Rossant, J. and Séguin, C. A. (2012). Tracing notochord-derived cells using a Noto-cre mouse: implications for intervertebral disc development. *Dis Model Mech* **5**, 73–82.

McLaren, S. B. P. and Steventon, B. J. (2021). Anterior expansion and posterior addition to the notochord mechanically coordinate zebrafish embryo axis elongation. *Development* **148**, dev199459.

Merrill, B. J., Pasolli, H. A., Polak, L., Rendl, M., García-García, M. J., Anderson, K. V. and Fuchs, E. (2004). Tcf3: a transcriptional regulator of axis induction in the early embryo. *Development* **131**, 263–274.

Moore-Scott, B. A., Opoka, R., Lin, S.-C. J., Kordich, J. J. and Wells, J. M. (2007). Identification of molecular markers that are expressed in discrete anterior–posterior domains of the endoderm from the gastrula stage to mid-gestation. *Developmental Dynamics* **236**, 1997–2003.

Moris, N., Anlas, K., van den Brink, S. C., Alemany, A., Schröder, J., Ghimire, S., Balayo, T., van Oudenaarden, A. and Martinez Arias, A. (2020). An in vitro model of early anteroposterior organization during human development. *Nature* **582**, 410–415.

Morrisey, E. E., Ip, H. S., Lu, M. M. and Parmacek, M. S. (1996). GATA-6: a zinc finger transcription factor that is expressed in multiple cell lineages derived from lateral mesoderm. *Dev Biol* **177**, 309–322.

Neijts, R., Simmini, S., Giuliani, F., van Rooijen, C. and Deschamps, J. (2014). Region-specific regulation of posterior axial elongation during vertebrate embryogenesis. *Developmental Dynamics* **243**, 88–98.

Norris, D. P., Brennan, J., Bikoff, E. K. and Robertson, E. J. (2002). The Foxh1-dependent autoregulatory enhancer controls the level of Nodal signals in the mouse embryo. *Development* **129**, 3455–3468.

Nowotschin, S., Hadjantonakis, A.-K. and Campbell, K. (2019). The endoderm: a divergent cell lineage with many commonalities. *Development* **146**,.

Olmsted, Z. T. and Paluh, J. L. (2021). Co-development of central and peripheral neurons with trunk mesendoderm in human elongating multi-lineage organized gastruloids. *Nat Commun* **12**, 3020.

Ortiz-Salazar, M. A., Camacho-Aguilar, E. and Warmflash, A. (2024). Endogenous Nodal switches Wnt interpretation from posteriorization to germ layer differentiation in geometrically constrained human pluripotent cells. 2024.03.13.584912.

Osorno, R., Tsakiridis, A., Wong, F., Cambray, N., Economou, C., Wilkie, R., Blin, G., Scotting, P. J., Chambers, I. and Wilson, V. (2012). The developmental dismantling of pluripotency is reversed by ectopic Oct4 expression. *Development* **139**, 2288–2298.

Paillat, L., Coutant, K., Dutilleul, M., Le Lay, S. and Camus, A. (2023). Three-dimensional culture model to study the biology of vacuolated notochordal cells from mouse nucleus pulposus explants. *Eur Cell Mater* **45**, 72–87.

Perea-Gomez, A., Vella, F. D. J., Shawlot, W., Oulad-Abdelghani, M., Chazaud, C., Meno, C., Pfister, V., Chen, L., Robertson, E., Hamada, H., et al. (2002). Nodal antagonists in the anterior visceral endoderm prevent the formation of multiple primitive streaks. *Dev. Cell* **3**, 745–756.

Pereira, P. N. G., Dobreva, M. P., Maas, E., Cornelis, F. M., Moya, I. M., Umans, L., Verfaillie, C. M., Camus, A., de Sousa Lopes, S. M. C., Huylebroeck, D., et al. (2012). Antagonism of Nodal signaling by BMP/Smad5 prevents ectopic primitive streak formation in the mouse amnion. *Development* **139**, 3343–3354.

Pham, T. X. A., Panda, A., Kagawa, H., To, S. K., Ertekin, C., Georgolopoulos, G., van Knippenberg, S. S. F. A., Allsop, R. N., Bruneau, A., Chui, J. S.-H., et al. (2022). Modeling human extraembryonic mesoderm cells using naive pluripotent stem cells. *Cell Stem Cell* **29**, 1346-1365.e10.

Plouhinec, J.-L., Granier, C., Le Mentec, C., Lawson, K. A., Sabéran-Djoneidi, D., Aghion, J., Shi, D. L., Collignon, J. and Mazan, S. (2004). Identification of the mammalian Not gene via a phylogenomic approach. *Gene Expression Patterns* **5**, 11–22.

Pour, M., Kumar, A. S., Farag, N., Bolondi, A., Kretzmer, H., Walther, M., Wittler, L., Meissner, A. and Nachman, I. (2022). Emergence and patterning dynamics of mouse-definitive endoderm. *iScience* **25**, 103556.

R Core Team (2013). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.

Repina, N. A., Johnson, H. J., Bao, X., Zimmermann, J. A., Joy, D. A., Bi, S. Z., Kane, R. S. and Schaffer, D. V. (2023). Optogenetic control of Wnt signaling models cell-intrinsic embryogenic patterning using 2D human pluripotent stem cell culture. *Development* **150**, dev201386.

Rito, T., Libby, A. R. G., Demuth, M. and Briscoe, J. (2023). Notochord and axial progenitor generation by timely BMP and NODAL inhibition during vertebrate trunk formation. 2023.02.27.530267.

Robertson, E. J. (2014). Dose-dependent Nodal/Smad signals pattern the early mouse embryo. *Seminars in Cell & Developmental Biology* **32**, 73–79.

Sanaki-Matsumiya, M., Matsuda, M., Gritti, N., Nakaki, F., Sharpe, J., Trivedi, V. and Ebisuya, M. (2022). Periodic formation of epithelial somites from human pluripotent stem cells. *Nat Commun* **13**, 2325.

Saunders, D., Camacho, C. and Steventon, B. (2024). Spinal Cord Elongation Enables Proportional Regulation of the Zebrafish Posterior Body. 2024.04.02.587732.

Scheibner, K., Schirge, S., Burtscher, I., Büttner, M., Sterr, M., Yang, D., Böttcher, A., Ansarullah, Irmler, M., Beckers, J., et al. (2021). Epithelial cell plasticity drives endoderm formation during gastrulation. *Nat Cell Biol* **23**, 692–703.

Schifferl, D., Scholze-Wittler, M., Luque, A. V., Pustet, M., Wittler, L., Veenvliet, J. V., Koch, F. and Herrmann, B. G. (2023). Genome-wide identification of notochord enhancers comprising the regulatory landscape of the Brachyury (T) locus in mouse. *Development* dev.202111.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**, 676–682.

Schohl, A. and Fagotto, F. (2002). Beta-catenin, MAPK and Smad signaling during early Xenopus development. *Development* **129**, 37–52.

Stemple, D. L. (2005). Structure and function of the notochord: an essential organ for chordate development. *Development* **132**, 2503–2512.

Streit, A. and Stern, C. D. (1999). Mesoderm patterning and somite formation during node regression: differential effects of chordin and noggin. *Mech Dev* **85**, 85–96.

Stringer, C., Wang, T., Michaelos, M. and Pachitariu, M. (2021). Cellpose: a generalist algorithm for cellular segmentation. *Nat Methods* **18**, 100–106.

Sulik, K., Dehart, D. B., Inagaki, T., Carson, J. L., Vrablic, T., Gesteland, K. and Schoenwolf, G. C. (1994). Morphogenesis of the murine node and notochordal plate. *Developmental Dynamics* **201**, 260–278.

Tam, P. P., Parameswaran, M., Kinder, S. J. and Weinberger, R. P. (1997). The allocation of epiblast cells to the embryonic heart and other mesodermal lineages: the role of ingression and tissue movement during gastrulation. *Development* **124**, 1631–1642.

Tamplin, O. J., Cox, B. J. and Rossant, J. (2011). Integrated microarray and ChIP analysis identifies multiple Foxa2 dependent target genes in the notochord. *Dev Biol* **360**, 415–425.

Turner, D. A., Girgin, M., Alonso-Crisostomo, L., Trivedi, V., Baillie-Johnson, P., Glodowski, C. R., Hayward, P. C., Collignon, J., Gustavsen, C., Serup, P., et al. (2017). Anteroposterior polarity and elongation in the absence of extra-embryonic tissues and of spatially localised signalling in gastruloids: mammalian embryonic organoids. *Development* **144**, 3894–3906.

Tzouanacou, E., Wegener, A., Wymeersch, F. J., Wilson, V. and Nicolas, J.-F. (2009). Redefining the Progression of Lineage Segregations during Mammalian Embryogenesis by Clonal Analysis. *Developmental Cell* **17**, 365–376.

Varoquaux, N. and Purdom, E. (2020). A pipeline to analyse time-course gene expression data.

Veenvliet, J. V., Bolondi, A., Kretzmer, H., Haut, L., Scholze-Wittler, M., Schifferl, D., Koch, F., Guignard, L., Kumar, A. S., Pustet, M., et al. (2020). Mouse embryonic stem cells self-organize into trunk-like structures with neural tube and somites. *Science* **370**, eaba4937.

Vianello, S. and Lutolf, M. P. (2020). In vitro endoderm emergence and self-organisation in the absence of extraembryonic tissues and embryonic architecture. *bioRxiv* 2020.06.07.138883.

Vincent, S. D., Dunn, N. R., Hayashi, S., Norris, D. P. and Robertson, E. J. (2003). Cell fate decisions within the mouse organizer are governed by graded Nodal signals. *Genes Dev.* **17**, 1646–1662.

Viotti, M., Nowotschin, S. and Hadjantonakis, A.-K. (2014). SOX17 links gut endoderm morphogenesis and germ layer segregation. *Nature Cell Biology* **16**, 1146–1156.

Warin, J., Vedrenne, N., Tam, V., Zhu, M., Yin, D., Lin, X., Guidoux-D'halluin, B., Humeau, A., Roseiro, L., Paillat, L., et al. (2024). In vitro and in vivo models define a molecular signature reference for human embryonic notochordal cells. *iScience* **27**, 109018.

Warmflash, A., Sorre, B., Etoc, F., Siggia, E. D. and Brivanlou, A. H. (2014). A method to recapitulate early embryonic spatial patterning in human embryonic stem cells. *Nat Meth* **11**, 847–854.

Washkowitz, A. J., Gavrilov, S., Begum, S. and Papaioannou, V. E. (2012). Diverse functional networks of Tbx3 in development and disease. *Wiley Interdiscip Rev Syst Biol Med* **4**, 273–283.

Winzi, M. K., Hyttel, P., Dale, J. K. and Serup, P. (2011). Isolation and characterization of node/notochord-like cells from mouse embryonic stem cells. *Stem Cells and Development* **20**, 1817–1828.

Wise, C. A., Sepich, D., Ushiki, A., Khanshour, A. M., Kidane, Y. H., Makki, N., Gurnett, C. A., Gray, R. S., Rios, J. J., Ahituv, N., et al. (2020). The cartilage matrisome in adolescent idiopathic scoliosis. *Bone Research* **8**, 1–13.

Wisniewski, D., Lowell, S. and Blin, G. (2019). Mapping the Emergent Spatial Organization of Mammalian Cells using Micropatterns and Quantitative Imaging. *J Vis Exp*.

Wymeersch, F. J., Skylaki, S., Huang, Y., Watson, J. A., Economou, C., Marek-Johnston, C., Tomlinson, S. R. and Wilson, V. (2019). Transcriptionally dynamic progenitor populations organised around a stable niche drive axial patterning. *Development* **146**,.

Wymeersch, F. J., Wilson, V. and Tsakiridis, A. (2021). Understanding axial progenitor biology in vivo and in vitro. *Development* **148**,.

Xu, P.-F., Borges, R. M., Fillatre, J., de Oliveira-Melo, M., Cheng, T., Thisse, B. and Thisse, C. (2021). Construction of a mammalian embryo model from stem cells organized by a morphogen signalling centre. *Nat Commun* **12**, 3277.

Yamamoto, M., Meno, C., Sakai, Y., Shiratori, H., Mochida, K., Ikawa, Y., Saijoh, Y. and Hamada, H. (2001). The transcription factor FoxH1 (FAST) mediates Nodal signaling during anterior-posterior patterning and node formation in the mouse. *Genes Dev* **15**, 1242–1256.

Yamanaka, Y., Tamplin, O. J., Beckers, A., Gossler, A. and Rossant, J. (2007). Live Imaging and Genetic Analysis of Mouse Notochord Formation Reveals Regional Morphogenetic Mechanisms. *Developmental Cell* **13**, 884–896.

Yasuo, H. and Lemaire, P. (2001). Role of Goosecoid, Xnot and Wnt antagonists in the maintenance of the notochord genetic programme in Xenopus gastrulae. *Development* **128**, 3783–3793.

Zhang, Y., Zhang, Z., Chen, P., Ma, C. Y., Li, C., Au, T. Y. K., Tam, V., Peng, Y., Wu, R., Cheung, K. M. C., et al. (2020). Directed Differentiation of Notochord-like and Nucleus Pulposus-like Cells Using Human Pluripotent Stem Cells. *Cell Reports* **30**, 2791-2806.e5.

Zhao, R., Watt, A. J., Li, J., Luebke-Wheeler, J., Morrisey, E. E. and Duncan, S. A. (2005). GATA6 Is Essential for Embryonic Development of the Liver but Dispensable for Early Heart Formation. *Mol Cell Biol* **25**, 2622–2631.

Zhao, C., Reyes, A. P., Schell, J. P., Weltner, J., Ortega, N. M., Zheng, Y., Björklund, Å. K., Baqué-Vidal, L., Sokka, J., Torokovic, R., et al. (2024). A Comprehensive Human Embryogenesis Reference Tool using Single-Cell RNA-Sequencing Data. 2021.05.07.442980.

848