Title: Investigating the impact of paternal aging on murine sperm miRNA profiles and their potential link to autism spectrum disorder

Kazusa Miyahara[+], Misako Tatehana[+], Takako Kikkawa, and Noriko Osumi*

Department of Developmental Neuroscience, Center for Advanced Research and Translational Medicine (ART), Tohoku University Graduate School of Medicine, Sendai, Japan.

[+]These authors contributed equally to this work

*Correspondence to: Noriko Osumi, PhD
Department of Developmental Neuroscience, Center for Advanced Research and Translational Medicine (ART), Tohoku University School of Medicine, Sendai, Japan.
TEL: 022-717-8201; FAX: 022-717-8205
E-mail: osumi@med.tohoku.ac.jp

Abstract

Paternal aging has consistently been linked to an increased risk of neurodevelopmental
disorders, including autism spectrum disorder (ASD), in offspring. Recent evidence has
highlighted the involvement of epigenetic factors. In this study, we aimed to investigate
age-related alterations in microRNA (miRNA) profiles of mouse sperm and analyze
target genes regulated by differentially expressed miRNAs. Microarray analyses were
conducted on sperm samples from mice at different ages: 3 months (3M), over 12M,
and beyond 20M. We identified 26 miRNAs with differential expression between the
3M and 20M, 34 miRNAs between the 12M and 20M, and 2 miRNAs between the 3M
and 12M mice. The target genes regulated by these miRNAs were significantly
associated with apoptosis/ferroptosis pathways and the nervous system. The findings
suggest that aging alters the sperm miRNA profiles and that the target genes regulated
by these differentially expressed miRNAs are associated with apoptosis and ASD,
implying a potential link between paternal aging and an increased risk of
neurodevelopmental disorders. The observed age-related changes in sperm miRNA
profiles have the potential to impact sperm quality and subsequently affect offspring
development.

Introduction

With the increasing trend of delayed parenthood, there has been a burgeoning focus on understanding the intricate relationship between aging and germ-line cells. While the effect of maternal aging on offspring, such as an elevated risk of miscarriage and Down syndrome, have garnered significant attention, the impact of male aging has received comparatively less recognition. However, recent research has revealed the advancing paternal age not only influences sperm morphology and motility, but also has implications for DNA integrity and epigenetic factors[1-4]. These alterations in sperm composition due to paternal aging have been implicated in perturbations during the developmental process of offspring, potentially contributing to spectrum of health issues[5].

In recent decades, investigations into the origins of health and disease in future generations have gained paramount significance. The concept of "Developmental Origin of Health and Disease (DOHaD)" has conventionally emphasized the impact of environmental factors on fetal development within the maternal uterus. However, there is now an emerging recognition that paternal factors also play a pivotal role in the risk of diseases manifestation in children, giving rise to the concept of the "Paternal Origin of Health and Disease (POHaD)"[6,7]. For example, a study has suggested more than 800 differential DNA methylation regions between fathers who have children with autism spectrum disorder (ASD) and those who do not have, suggesting that epigenetic alterations of sperm are key for occasion of ASD[8]. Paternal aging, in particular, has been linked to not only to adverse incidents during pregnancy, such as miscarriage[9] and stillbirth[10], but also postnatal health issues, including low birth weight[11], congenital heart defects[12], and malignant tumors[13]. Although the concept of POHaD is a relatively nascent, delving into this perspective holds the potential to provide insights into diseases that remain incompletely understood.

The concept of POHaD has emerged as a compelling framework, offering substantial evidence linking paternal factors to a range of neuropsychiatric disorders in offspring. Notably, advanced paternal age has been consistently demonstrated an increased risk of schizophrenia[14], bipolar disorder[15], attention-deficit/hyperactivity disorder (ADHD)[16], neurocognitive dysfunction[17], ASD[18] in subsequent generations. The global surge in ASD[19] has been influenced by a complex interplay of social and biological factors, encompassing updated diagnostic criteria, heightened social awareness, maternal drug exposure, and parental aging[18,20-22]. Remarkably, a meta-analysis of six million subjects across five countries has revealed that paternal aging exerts a more pronounced impact

on ASD risk compared to maternal aging[18]. Consequently, further exploration on the effects of POHaD may hold significant potential in providing invaluable insights into understanding pathophysiology of ASD.

The precise mechanism underlying the impact of paternal aging on the health of the next generation remains incompletely understood. *De novo* mutations, which are mutations that are not present in eighter parent but are observed in the child, have been associated with paternal aging and are believed to contribute to the increased disease risk in offspring. Specifically, *de novo* mutations of paternal origin have been found to accumulate with advancing paternal age, particularly in individuals diagnosed with ASD and schizophrenia[24,25]. Nevertheless, it is important to note that *de novo* mutations derived from aged fathers account for only a fraction, approximately 10-20%, of the overall risk of psychiatric disorders in children[26,27]. This suggests the involvement of additional factors that necessitate careful consideration in understanding the complete picture.

Epigenetic modifications in sperm have emerged as another plausible mechanism underlying the effects of paternal aging effects on offspring[28]. We have previously revealed age-related modifications in DNA methylation in mouse sperm[4] and changes in histone modifications within the testis[29], suggesting their potential influence on offspring development. Micro RNAs (miRNAs) are another crucial epigenetic factor involved in gene expression regulation by inhibiting translation. These short non-coding RNA molecules (21-25 nucleotides) form complexes known as RNA-induced silencing complexes with their mRNA target sequences, ultimately suppressing translation[30]. The biomedical research field actively investigates miRNAs due to their potential as biomarkers Notably, aberrant miRNA profiles have been identified in individuals with ASD, with altered miRNA expression observed in various samples such as the brain, blood, serum, and saliva[31-33]. Similar findings have been reported in animal models of ASD[34]. However, the the precise mechanisms by which these changes in miRNA expression contribute to the pathogenesis of ASD remain unclear.

In a previous study on human sperm samples, age-related alterations in miRNA expression have been reported, with certain miRNAs exhibiting increased levels while others show decreased levels[2]. Additionally, alterations in mouse sperm miRNA expression, induced by factors such as diet and exercise, can impact the insulin sensitivity of female offspring[35]. Nevertheless, the comprehensive changes in overall miRNA profiles and the precise role of the target genes regulated by these altered miRNAs remain elusive. Therefore, the primary objective of this study is to

meticulously examine the expression profiles of miRNAs in sperm obtained from aged mice in comparison to young counterparts. Furthermore, we aim to unravel the potential functions of differentially expressed miRNA's target genes, shedding light on their roles in the context of paternal aging and potential implications for offspring mental health.

Materials and methods

Animals

C57BL/6J mice at 3 months old were purchased from a breeder (Charles River

Laboratories, Japan) and were raised to 12 or 20 months old at the Institute for Animal

Experimentation at the Tohoku University Graduate School of Medicine; they were then

used for miRNA and mRNA analysis. The mice were then sampled for microarray

analyses at 3 months old (3M), > 12M or > 20M (Fig. 1). All animals were housed in

standard cages in a temperature- and humidity-controlled room with a 12-hour

light/dark cycle (light on at 8 am), and they had free access to standard food and water.

All experimental procedures were approved by the Ethics Committee for Animal

Experiments of the Tohoku University Graduate School of Medicine (#2017- MED210,

#2019- MED280-03, 2020BeLMO-005-01 and 2020BeA-015-01), and the animals were

treated according to the National Institutes of Health guidance for the care and use of

laboratory animals.

Sperm collection for miRNA and mRNA analyses

For microarray analyses, three mice for each age were used. Mice were anesthetized

with isoflurane (MSD Animal Health) and perfused with PBS (pH 7.4) to remove blood.

After perfusion, the cauda epididymis was removed from fat and other connective

tissues to collect sperm by retrograde perfusion[36-38]. Sperm were centrifuged on a 27%

Percoll solution (2005-001 or 2001-017, NK system). The pellet was resuspended in

PBS and recentrifuged ($400 \times g$ for 2 min) to remove excess Percoll. The purity of the

pellet was confirmed by microscopy. Samples were then subjected to extracting miRNA

and mRNA.

RNA extraction and microarray analysis

Sperm samples were subjected to extract total RNA including miRNA or mRNA by

using miRNeasy Mini Kit (217004, QIAGEN). We confirmed whether total RNA

including miRNA was sufficiently collected by Agilent Small RNA Reagent kit

(5067-1549) with 2100 Bioanalyzer Instrument (G2939BA). RNA samples were

subjected to microarray analyses. Clariom™ D Assay for the mouse (902513,

ThermoFisher Scientific) and GeneChip™ miRNA 4.0 Array (902412, ThermoFisher

Scientific) were used for mRNA and miRNA analyses, respectively. Transcript

expression values were normalized using Signal Space Transformation (SST-RMA)

method. Raw intensity data (i.e., CEL files) was analyzed by using Transcriptome

Analysis Console (TAC) Software version 4.0.2 (ThermoFisher Scientific). We

identified differentially expressed miRNAs based on > 2-fold up- or down-regulation

along with a *P*-value < 0.05. Microarray data have been deposited into the Gene Expression Omnibus database under accession number GSE232620 and are available at the following URL: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE232620. Pathway analyses were conducted using the Reactome pathway analysis package in R[39]. Networks were constructed using Cytoscape[40], which visualized relationships between enriched pathways and genes. Pearson`s correlation test was utilized to analyze the correlation between miRNAs (i.e., *mmu-let-7b-5p*, *mmu-miR-10a-5p* and *mmu-miR-24-3p*) and their target genes related with apoptosis (i.e., *Ppp3r1* and *Bcl2l11*). We also compared among groups the expression levels of the five miRNAs that have previously been reported as candidate miRNAs (i.e., miR-10a-5p, miR-146a-5p, miR-34c-5p, miR-143-3p, and miR-146b-5p) transmitted from sperm to fertilized egg[41]. In these analyses, *P* < 0.05 was considered significant. We used Bonferroni's multiple comparison test to verify the significance of the microarray results.

Analyses exploring possible function of target genes of altered miRNAs
To investigate possible functions of the target genes of differentially expressed miRNAs in relation to ASD, we utilized the autism-related gene database SFARI[42]. We constructed a network of miRNA-target gene interactions by using Cytoscape. Additionally, we also conducted Pearson`s correlation test between the expression levels of miRNAs (*mmu-miR-466j*, *mmu-miR-24-3p*, *mmu-miR-690*, and *mmu-let-7b-5p*) and their target SFARI genes (*Oxtr*, *Gabrb2*, *Ctnnb1*, and *Grik2*).

Results

Age-related changes in expression of miRNAs in the mouse sperm

We first analyzed age-related changes in sperm miRNA expression patterns. 298, 273,

and 364 miRNAs were expressed in sperm samples of 3 M, 12 M, and 20 M,

respectively (Fig. 2A). There were 221 miRNAs commonly expressed in 3 M and 12 M,

and 248 miRNAs commonly expressed in 3 M and 20 M. In 12 M and 20 M, 229

miRNAs were commonly expressed, and 210 miRNAs were expressed in all three

groups (Fig. 2B).

The miRNAs were then compared between the two groups, respectively. We considered

the miRNAs that showed > 2-fold up- or down-regulation along with $P < 0.05$ in

expression as significantly different. Two miRNAs were upregulated in the sperm

sample collected from 12M compared with the sperm of 3M, 21 miRNAs were

upregulated, and 5 miRNAs were downregulated in 3M vs 20M samples, and 26 and 8

miRNAs were up- or down-regulated, respectively, in 12M vs 20M samples (Fig. 2B-E).

The differentially expressed miRNAs at each comparison are summarized in

Supplementary Table S1.

No overlap was observed between the miRNAs that exhibited alterations in the 3M vs

12M comparison and those in the 3M vs 20M comparison (Fig. 2F). In the 3M vs 12M

comparison, two miRNAs that displayed changes in the 3M vs 12M comparison were

also found to be altered (4.5% of the total altered miRNAs). Furthermore, out of the 26

miRNAs that showed alteration in the 3M vs 20M comparison, 16 of them were

similarly altered in the 12M vs 20M comparison (36% of the total altered miRNAs).

Notably, this subset of miRNAs included *let-7b-5p*, *miR-10a-5p*, and *miR-24-3p,* which

target apoptosis-related genes (Fig. 2F; Supplementary Table S2). These results indicate

a distinct age-associated shift in the sperm miRNA profiles, particularly when

comparing 3M vs 20M and 12M vs 20M age groups.

Targets of age-related miRNAs in the mouse sperm

Next, miRNAs significantly altered in the comparison between groups were listed and

their target genes (Supplementary Table S2) were used for enrichment analyses by using

an R package for Reactome Pathway Analysis[39]. Although, the 3M vs 12M comparison

did not have any pathways that were significantly different (Fig. 3A), comparison

between 3M and 20M samples yielded differential pathways including "Dimerization of

procaspase-8", "CASP8 activity is inhibited", "Regulation by c-FLIP", "Apoptosis",

"Caspase activation via Death Receptors in the presence of ligand", and "Caspase

activation via extrinsic apoptotic signaling pathway" (Fig. 3B). Most of these pathways are thus related to the regulation of apoptosis.

In our comparison between 12M and 20M, we also found several pathways that showed significant differences. These included the same apoptosis-related pathways such as "Caspase activation via Death Receptors", "Dimerization of procaspase-8" and "Apoptosis", as well as other pathways such as "Signaling by Receptor Tyrosine Kinases", "Transmission across Chemical Synapses", and "NCAM signaling for neurite out-growth" and "Neuronal System" (Fig. 3C). Those pathways related to the nervus system were not significantly enriched in comparison between 3M vs 20M (Fig. 3B). Furthermore, four genes related to ferroptosis, another type of cell death, including *Acsl4*, *Hmox1*, *Slc11a2*, and *Slc3a2,* were identified as target genes of altered miRNAs between 3M vs 20M. Among them, *Acsl4* and *Hmox1* were also target genes in comparison between 12M vs 20M. Therefore, our findings suggest that aging may be linked with cell death during spermatogenesis.

In summary, we found more differentially expressed miRNAs in comparison of 12M vs 20M sperm samples, which yielded more target genes in unique pathways. The significant pathways in comparison between 3M vs 20M samples overlapped with those that were significant between 12M vs. 20M (Fig. 3D). These results indicate that the target genes of miRNAs that change with aging were significantly enriched in pathways associated with cell death and the nervous system.

Changes in mRNA expression of age-associated miRNA target genes

We measured expression levels of target genes using a commercial microarray kit and the top 10 enriched pathways and genes were visualized using Cytoscape[40] (Fig. 4). Among these pathways, we focused on the apoptosis-related pathways (such as "Caspase activation via Death Receptors", "Dimerization of procaspase-8" and "Apoptosis") and investigated whether the expression levels of the target genes regulated by altered miRNAs were affected by aging (Supplementary Table S3). The network representations in Fig. 4 revealed that the expression levels of most genes, including *Casp9*, *Tnfsf10*, and *Fadd*, exhibited minimal changes. However, the expression levels of *Ppp3r1,* regulated by *mmu-let-7b-5p,* and *Bcl2l11,* regulated by *mmu-miR-10a-5p* and *mmu-miR-24-3p,* increased with sperm aging. Notably, *Ppp3r1* showed a significant increase in 20M mice compared to 3M mice (fold change = 1.5, *P* = 0.029). To validate the association between changes in gene expression, we conducted Pearson's correlation tests. The results demonstrated a significant positive correlation between the expression levels of *Ppp3r1* and *mmu-let-7b-5p* (Fig. 5A: *R* = 0.840, *P* =

0.005). Although *Bcl2l11* also exhibited a positive trend with *mmu-miR-10a-5p* and *mmu-miR-24-3p,* the correlation did not reach statistical significance (Fig. 5B and 4C; *mmu-miR-10a-5p*: $R = 0.646$, $P = 0.060$; *mmu-miR-24-3p*; $R = 0.613$, $P = 0.079$).

Target genes of altered miRNAs related with ASD

Since paternal aging is a risk for ASD[18,22], we analyzed whether the target genes of the age-related miRNAs might have some relation with ASD. We searched the autism-related gene database SFARI[42], and identified an overlap of 28 and 38 SFARI genes with the target genes of differentially-expressed miRNAs between 3M vs 20M and 12M vs 20M, respectively (Supplementary Table S4). No SFARI genes were identified as miRNA targets in comparisons with 3M and 12M. The relationship between miRNAs and SFARI genes is visualized in Fig. 6 as a network. We focused on four SFARI genes (i.e., *Oxtr*, *Gabrb2*, *Ctnnb1*, and *Grik2*), whose expression levels differed significantly between the groups, and confirmed correlations of the expression levels and their regulatory miRNAs. These included significant negative correlations between *mmu-miR-466j* vs *Oxtr* (Fig. 7A; $R = -0.701$, $p= 0.035$) and *mmu-miR-24-3p* vs *Gabrb2* (Fig. 7B; $R = -0.707$, $p=0.033$), while there was a tendency of positive correlation between *mmu-miR-690* vs *Ctnnb1* (Fig. 7C; $R = 0.615$, $p=0.078$). A negative correlation between *mmu-let-7b-5p* vs *Grik2* was not significant (Fig. 7D; $R = -0.246$, $P = 0.523$).

Changes in expression of miRNAs that may possibly be transmitted from sperm to fertilized eggs

Finally, we focused on candidate miRNAs that can be carried from sperm to the fertilized egg, based on a previous mouse research comparing miRNA profiles of sperm, oocytes and fertilized eggs[41]. We examined, using the microarray, expression of the miRNAs, *miR-10a-5p*, *miR-146a-5p*, *miR-34c-5p*, *miR-143-3p*, *miR-146b-5p*. Our results revealed significant upregulation of *miR-10a-5p* and *miR-146a-5p* at 20M in the mouse (Fig. 7E-I). This finding highlights the potential significance of *miR-10a-5p* and *miR-146a-5p* as candidate sperm miRNAs transmitted to the zygote and their possible implications for subsequent offspring development.

Discussion

The present study aimed to investigate the effect of paternal aging on miRNA profiles in mouse sperm. We identified differentially expressed miRNAs with aging, which were significantly enriched in pathways related to apoptosis and the nervous system. We observed that the expression levels of *Ppp3r1* and *Bcl2l11*, two target genes of these miRNAs, showed trends of changes with aging and correlations with their respective regulatory miRNAs. Moreover, we found upregulation of *miR-10a-5p* and *miR-146a-5p* at 20M, suggesting the possibility of these miRNAs being transmitted to the zygote and having impact on offspring's development. Our study provided evidence that age-related changes in miRNAs might have an impact on the regulation of miRNAs in sperm, which may subsequently affect sperm quality and health of offspring.

Changes in expression of miRNAs with aging: implications for cell death or the nervous system

Our analyses of miRNAs in sperm samples obtained from mice at different ages (3M, 12 M, and 20M) revealed distinct patterns of miRNA expression. Notably, when comparing the miRNA profiles between the youngest (3M) and the oldest (20M) groups, we observed significant associations with pathways related to apoptosis (Fig. 3B). This finding aligns with previous studies demonstrating an increase in apoptosis and DNA damage in human and mouse sperm with advanced age[43,44].

Through our analyses, we identified miRNA target genes that both promote (e.g., *Bcl2l11*, *Casp9*, and *Ppp3r1*) and inhibit (e.g., *Tnfsf10*, *Fadd*, and *Fas*) apoptosis. The expression levels of these genes play critical role in regulating apoptosis. For instance, decreased expression of *Bcl2l11* and *Casp9* has been associated with suppression of sperm apoptosis[45]. Furthermore, exposure of recombinant tumor necrosis factor-related apoptosis-inducing ligand (coded by *Tnfsf10*) can modulate sperm apoptosis[46]. Our findings revealed variations in the expression levels of certain genes, including *Bcl2l11* and *Ppp3r1,* across different age groups, suggesting that changes in miRNA profile may contribute to the aberrant expression of these target genes.

Given that 20M represents a highly advanced age, it is important to consider that the aging process is accompanied by multitude of factors, including increased oxidative stress, lipid peroxidation, iron accumulation, and chronic inflammation. These factors may collectively contribute to heightened aging-related stress, which may not only promote apoptosis and ferroptosis but also increase the occurrence of necrosis. Our analysis revealed the presence of pathways related to apoptosis and ferroptosis in the

comparison between 12M and 20M groups (Fig. 3C), further supporting the notion that changes in miRNA profile of sperm resulting from paternal aging can influence different forms of cell death.

Our investigation uncovered several intriguing aging-related pathways, such as "Signaling by Receptor Tyrosine Kinases", "Transmission across Chemical Synapses", "Neuronal System" and "NCAM signaling for neurite out-growth". The consistency of these findings is particularly noteworthy, considering the shared expression of many genes in both the brain and testis[47]. One example is genes related with calcium signaling, a crucial process in both neuron and sperm. Calcium channels play a central role in neurotransmission in the brain[48] and are essential for sperm motility and capacitation[49]. In our study, we identified target genes involved in calcium signaling, including *Camkk1* and *Cacna2d2*. *Cacna2d2* codes for the alpha-2/delta subunit of the voltage-dependent calcium channel complex, while *Camkk1* codes for the Ca^{2+}/calmodulin-dependent protein kinase α, both of which are essential for proper calcium signaling in sperm[50,51]. Consequently, aberrant expression of these genes due to miRNAs may disrupt sperm motility and capacitation[49], both of which are also associated with paternal aging[1,52]. Thus, our findings suggest that changes in signaling pathways related to calcium signaling and other pathways identified in sperm from aged mice could contribute to the observed decline of sperm quality in aged males.

Changes in expression levels of genes caused by altered miRNAs with aging

Our study provides compelling evidence that altered miRNAs have an impact on the expression levels of specific genes in the mouse sperm. Particularly noteworthy were the age-related changes in the expression levels of the two apoptosis-related genes, *Ppp3r1* and *Bcl2l11*. These changes demonstrated a proportional association with the expression levels of their regulatory miRNAs, namely *mmu-let-7b-5p*, *mmu-miR-10a-5p*, and *mmu-miR-24-3p*, as quantitatively determined by microarray analyses. Previous research has indicated that the upregulation of these three miRNAs is associated with apoptosis[53-55]. For instance, Huang, Y. et al. has demonstrated that *let-7b-5p* promotes apoptosis, and that knockdown of *let-7b-5p* suppresses apoptosis in cultured human cells[53]. These findings lend further support to our hypothesis that these three miRNAs may indeed contribute to sperm apoptosis.

Paternal aging and ASD risk of offspring

We further investigated the impact of altered miRNAs on autism-related SFARI genes through microarray analyses. Although changes in the expression levels of SFARI genes were generally limited, intriguing associations were found between the expression

levels of *Oxtr* and *Gabrb2* and their regulatory miRNAs. *Oxtr* encodes an oxytocin receptor, a crucial component in regulation of social behavior[56]. A meta-analysis has previously identified four polymorphisms in the human *OXTR* gene (rs7632287, rs237887, rs2268491, and rs2254298) that are linked to ASD[57]. Similarly, *Gabrb22* encodes the gamma-aminobutyric acid (GABA) type A receptor subunit beta2, an important component of the inhibitory system in the brain[58]. Disruptions of GABA system have been implicated in ASD risk[59]. Polymorphisms on *GABRB2*, such as rs2617503 and rs12187676, have been implicated in ASD risk[60]. Although the roles of *Oxtr* and *Gabrb2* in germ-line cells are not well understood, their crucial functions in brain development are known unknown. Aberrant expressions of these genes due to miRNA changes in sperm may provide partial insights into the relationship between paternal aging and the increased risk of ASD in offspring.

Limitation

While our study provides valuable insights into the role of miRNAs in paternal aging and their potential association with ASD risk, several limitations should be acknowledged. First, to fully understand the physiological function of these miRNAs in fertilized eggs and their relationship to ASD, additional investigations involving techniques such as miRNA injection or knockdown in fertilized mouse eggs are necessary. Despite the relative low abundance of miRNAs in sperm compared to the fertilized egg, they still hold promise as potential biomarkers for predicting the physical conditions of offspring. In addition, it is important to acknowledge that our study focused on miRNAs and their target genes, but other regulatory mechanisms and factors may also contribute to the development of ASD. Exploring additional epigenetic modifications and factors that influence gene expression could provide a more comprehensive understanding of the paternal contribution to ASD.

Conclusion

In this study, we aimed to investigate epigenetic changes in male germline cells induced by aging and their potential impact on the development of the next generation. Our findings shed light on the alterations in the miRNA profile of sperm as a result of aging, which can subsequently lead to changes in gene expression levels. These changes have the potentials to contribute to a wide range of symptoms associated with ASD and other transgenerational epigenetic changes. Future research is needed to elucidate the precise mechanism by which the sperm epigenome influences the disease risk in the offspring. Such investigations will not only deepen our understanding of the function of the sperm

epigenome functions but also contribute to the development of novel therapeutic and preventive strategies for the diseases in the next generation. In addition to the well-established importance of the maternal environment in the DOHaD concept, our study highlights the growing relevance of investigating the effects of POHaD. By exploring both maternal and paternal contributions, we can gain comprehensive insights into the complex interplay between genetics, epigenetics, and environmental factors in shaping the health outcome of future generations.

References

1.   Gunes, S., Hekim, G. N., Arslan, M. A. & Asci, R. Effects of aging on the male reproductive system. *J. Assist. Reprod. Genet.* **33**, 441-454, doi:10.1007/s10815-016-0663-y (2016).

2.   Paoli, D. *et al.* Cytological and molecular aspects of the ageing sperm. *Hum. Reprod.* **34**, 218-227, doi:10.1093/humrep/dey357 (2019).

3.   Moskovtsev, S. I., Willis, J. & Mullen, J. B. Age-related decline in sperm deoxyribonucleic acid integrity in patients evaluated for male infertility. *Fertil. Steril.* **85**, 496-499, doi:10.1016/j.fertnstert.2005.05.075 (2006).

4.   Yoshizaki, K. *et al.* Paternal age affects offspring via an epigenetic mechanism involving REST/NRSF. *EMBO Rep* **22**, e51524, doi:10.15252/embr.202051524 (2021).

5.   Sharma, R. *et al.* Effects of increased paternal age on sperm quality, reproductive outcome and associated epigenetic risks to offspring. *Reprod. Biol. Endocrinol.* **13**, 35, doi:10.1186/s12958-015-0028-x (2015).

6.   Soubry, A. POHaD: why we should study future fathers. *Environ Epigenet* **4**, dvy007, doi:10.1093/eep/dvy007 (2018).

7.   Soubry, A. Epigenetics as a Driver of Developmental Origins of Health and Disease: Did We Forget the Fathers? *Bioessays* **40**, doi:10.1002/bies.201700113 (2018).

8.   Garrido, N. *et al.* Sperm DNA methylation epimutation biomarker for paternal offspring autism susceptibility. *Clin. Epigenetics* **13**, 6, doi:10.1186/s13148-020-00995-2 (2021).

9.   du Fosse, N. A., van der Hoorn, M. P., van Lith, J. M. M., le Cessie, S. & Lashley, E. Advanced paternal age is associated with an increased risk of spontaneous miscarriage: a systematic review and meta-analysis. *Hum. Reprod. Update* **26**, 650-669, doi:10.1093/humupd/dmaa010 (2020).

10.   Urhoj, S. K., Andersen, P. K., Mortensen, L. H., Davey Smith, G. & Nybo Andersen, A. M. Advanced paternal age and stillbirth rate: a nationwide register-based cohort study of 944,031 pregnancies in Denmark. *Eur. J. Epidemiol.* **32**, 227-234, doi:10.1007/s10654-017-0237-z (2017).

11.   Mao, Y. *et al.* Association Between Paternal Age and Birth Weight in Preterm and Full-Term Birth: A Retrospective Study. *Front. Endocrinol. (Lausanne)* **12**, 706369, doi:10.3389/fendo.2021.706369 (2021).

12.   Malik, S. *et al.* Association between congenital heart defects and small for gestational age. *Pediatrics* **119**, e976-982, doi:10.1542/peds.2006-2742 (2007).

13.   Urhoj, S. K., Jespersen, L. N., Nissen, M., Mortensen, L. H. & Nybo Andersen, A. M. Advanced paternal age and mortality of offspring under 5 years of age: a

register-based cohort study. *Hum. Reprod.* **29**, 343-350, doi:10.1093/humrep/det399 (2014).

D Malaspina, S. H., S Fennig, D Heiman, D Nahon, D Feldman, E S Susser. Advancing paternal age and the risk of schizophrenia. *Arch. Gen. Psychiatry* **58**, 361-367, doi:10.1001/archpsyc.58.4.361 (2001).

Emma M Frans, S. S., Abraham Reichenberg, Paul Lichtenstein, Niklas Långström, Christina M Hultman. Advancing Paternal Age and Bipolar Disorder. *Arch. Gen. Psychiatry* **65**, 1034-1040, doi:10.1001/archpsyc.65.9.1034. (2008).

D'Onofrio, B. M. *et al.* Paternal age at childbearing and offspring psychiatric and academic morbidity. *JAMA Psychiatry* **71**, 432-438, doi:10.1001/jamapsychiatry.2013.4525 (2014).

Saha, S. *et al.* Advanced paternal age is associated with impaired neurocognitive outcomes during infancy and childhood. *PLoS Med.* **6**, e40, doi:10.1371/journal.pmed.1000040 (2009).

Sandin, S. *et al.* Autism risk associated with parental age and with increasing difference in age between the parents. *Mol. Psychiatry* **21**, 693-700, doi:10.1038/mp.2015.70 (2016).

Jon Baio, L. W., Deborah L Christensen, Matthew J Maenner, Julie Daniels, Zachary Warren, Margaret Kurzius-Spencer, Walter Zahorodny, Cordelia Robinson Rosenberg, Tiffany White, Maureen S Durkin, Pamela Imm, Loizos Nikolaou, Marshalyn Yeargin-Allsopp, Li-Ching Lee, Rebecca Harrington, Maya Lopez, Robert T Fitzgerald, Amy Hewitt, Sydney Pettygrove, John N Constantino, Alison Vehorn, Josephine Shenouda, Jennifer Hall-Lande, Kim Van Naarden Braun, Nicole F Dowling. Prevalence of Autism Spectrum Disorder Among Children Aged 8 Years – Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2014. *MMWR Surveill. Summ.* **67**, 1-23, doi:10.15585/mmwr.ss6706a1. (2018).

Abraham Reichenberg, R. G., Mark Weiser, Michealine Bresnahan, Jeremy Silverman, Susan Harlap, Jonathan Rabinowitz, Cory Shulman, Dolores Malaspina, Gad Lubin, Haim Y Knobler, Michael Davidson, Ezra Susser. Advancing Paternal Age and Autism. *Arch. Gen. Psychiatry* **63**, 1026-1032, doi:10.1001/archpsyc.63.9.1026. (2006).

Weintraub, K. The prevalence puzzle: Autism counts. *Nature* **479**, 22-24, doi:10.1038/479022a. (2011).

Sato, A., Kotajima-Murakami, H., Tanaka, M., Katoh, Y. & Ikeda, K. Influence of Prenatal Drug Exposure, Maternal Inflammation, and Parental Aging on the Development of Autism Spectrum Disorder. *Front Psychiatry* **13**, 821455, doi:10.3389/fpsyt.2022.821455 (2022).

Sanders, S. J. *et al.* De novo mutations revealed by whole-exome sequencing are strongly associated with autism. *Nature* **485**, 237-241, doi:10.1038/nature10945 (2012).

Kong, A. *et al.* Rate of de novo mutations and the importance of father's age to disease risk. *Nature* **488**, 471-475, doi:10.1038/nature11396 (2012).

O'Roak, B. J. *et al.* Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations. *Nature* **485**, 246-250, doi:10.1038/nature10989 (2012).

Gratten, J. *et al.* Risk of psychiatric illness from advanced paternal age is not predominantly from de novo mutations. *Nat. Genet.* **48**, 718-724, doi:10.1038/ng.3577 (2016).

Taylor, J. L. *et al.* Paternal-age-related de novo mutations and risk for five disorders. *Nat Commun* **10**, 3043, doi:10.1038/s41467-019-11039-6 (2019).

Schagdarsurengin, U., Paradowska, A., & Steger, K. Analysing the sperm epigenome: roles in early embryogenesis and assisted reproduction. *Nature reviews. Urology* **9**, 609-619, doi:10.1038/nrurol.2012.183 (2012).

Tatehana, M., Kimura, R., Mochizuki, K., Inada, H. & Osumi, N. Comprehensive histochemical profiles of histone modification in male germline cells during meiosis and spermiogenesis: Comparison of young and aged testes in mice. *PLoS One* **15**, e0230930, doi:10.1371/journal.pone.0230930 (2020).

Krol, J., Loedige, I. & Filipowicz, W. The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet* **11**, 597-610, doi:10.1038/nrg2843 (2010).

Hicks, S. D. & Middleton, F. A. A Comparative Review of microRNA Expression Patterns in Autism Spectrum Disorder. *Front Psychiatry* **7**, 176, doi:10.3389/fpsyt.2016.00176 (2016).

Sehovic, E. *et al.* Identification of developmental disorders including autism spectrum disorder using salivary miRNAs in children from Bosnia and Herzegovina. *PLoS One* **15**, e0232351, doi:10.1371/journal.pone.0232351 (2020).

Hicks, S. D., Carpenter, R. L., Wagner, K. E., Pauley, R., Barros, M., Tierney-Aves, C., Barns, S., Greene, C. D., & Middleton, F. A. Saliva MicroRNA Differentiates Children With Autism From Peers With Typical and Atypical Development. *J. Am. Acad. Child Adolesc. Psychiatry* **59**, 296-308, doi:10.1016/j.jaac.2019.03.017. (2020).

Urdinguio, R. G. *et al.* Disrupted microRNA expression caused by Mecp2 loss in a mouse model of Rett syndrome. *Epigenetics* **5**, 656-663, doi:10.4161/epi.5.7.13055 (2010).

McPherson, N. O., Owens, J. A., Fullston, T. & Lane, M. Preconception diet or

exercise intervention in obese fathers normalizes sperm microRNA profile and metabolic syndrome in female offspring. *Am. J. Physiol. Endocrinol. Metab.* **308**, E805-821, doi:10.1152/ajpendo.00013.2015 (2015).

Nixon, B. *et al.* The microRNA signature of mouse spermatozoa is substantially modified during epididymal maturation. *Biol. Reprod.* **93**, 91, doi:10.1095/biolreprod.115.132209 (2015).

Nixon, B. *et al.* The identification of mouse sperm-surface-associated proteins and characterization of their ability to act as decapacitation factors. *Biol. Reprod.* **74**, 275-287, doi:10.1095/biolreprod.105.044644 (2006).

Baker, M. A., Hetherington, L., Weinberg, A. & Velkov, T. Phosphopeptide analysis of rodent epididymal spermatozoa. *J Vis Exp*, doi:10.3791/51546 (2014).

Yu, G. & He, Q. Y. ReactomePA: an R/Bioconductor package for reactome pathway analysis and visualization. *Mol. Biosyst.* **12**, 477-479, doi:10.1039/c5mb00663e (2016).

Shannon, P. T., Grimes, M., Kutlu, B., Bot, J. J., & Galas, D. J. RCytoscape: tools for exploratory network analysis. *BMC Bioinformatics* **14**, doi:10.1186/1471-2105-14-217 (2013).

Yang, Q., Lin, J., Liu, M., Li, R., Tian, B., Zhang, X., Xu, B., Liu, M., Zhang, X., Li, Y., Shi, H., & Wu, L. Highly sensitive sequencing reveals dynamic modifications and activities of small RNAs in mouse oocytes and early embryos. *Science advances* **2**, doi:10.1126/sciadv.1501482 (2016).

Banerjee-Basu, S., & Packer, A. SFARI Gene: an evolving database for the autism research community. *Dis. Model. Mech.* **3**, 133-135, doi:10.1242/dmm.005439 (2010).

Siddighi, S., Chan, C. A., Patton, W. C., Jacobson, J. D. & Chan, P. J. Male age and sperm necrosis in assisted reproductive technologies. *Urol. Int.* **79**, 231-234, doi:10.1159/000107955 (2007).

Stobezki, R., Titus, S., Halicka, D., Darzynkiewicz, Z. & Oktay, K. Declining BRCA-Mediated DNA Repair in Sperm Aging and its Prevention by Sphingosine-1-Phosphate. *Reprod. Sci.* **27**, 940-953, doi:10.1007/s43032-019-00098-1 (2020).

Ding, Y. *et al.* MicroRNA-222 Transferred From Semen Extracellular Vesicles Inhibits Sperm Apoptosis by Targeting BCL2L11. *Front Cell Dev Biol* **9**, 736864, doi:10.3389/fcell.2021.736864 (2021).

Zauli, G. *et al.* Soluble TRAIL is present at high concentrations in seminal plasma and promotes spermatozoa survival. *Reproduction* **148**, 191-198, doi:10.1530/REP-14-0144 (2014).

Matos, B., Publicover, S. J., Castro, L. F. C., Esteves, P. J. & Fardilha, M. Brain and testis: more alike than previously thought? *Open Biol* **11**, 200322, doi:10.1098/rsob.200322 (2021).

Brini, M., Cali, T., Ottolini, D. & Carafoli, E. Neuronal calcium signaling: function and dysfunction. *Cell. Mol. Life Sci.* **71**, 2787-2814, doi:10.1007/s00018-013-1550-7 (2014).

Darszon, A., Nishigaki, T., Beltran, C. & Trevino, C. L. Calcium channels in the development, maturation, and function of spermatozoa. *Physiol. Rev.* **91**, 1305-1355, doi:10.1152/physrev.00028.2010 (2011).

Jose, O. *et al.* Recombinant human ZP3-induced sperm acrosome reaction: evidence for the involvement of T- and L-type voltage-gated calcium channels. *Biochem. Biophys. Res. Commun.* **395**, 530-534, doi:10.1016/j.bbrc.2010.04.059 (2010).

Nguyen, T. M., Combarnous, Y., Praud, C., Duittoz, A. & Blesbois, E. Ca2+/Calmodulin-Dependent Protein Kinase Kinases (CaMKKs) Effects on AMP-Activated Protein Kinase (AMPK) Regulation of Chicken Sperm Functions. *PLoS One* **11**, e0147559, doi:10.1371/journal.pone.0147559 (2016).

de La Rochebrochard, E. & Thonneau, P. Paternal age >or=40 years: an important risk factor for infertility. *Am. J. Obstet. Gynecol.* **189**, 901-905, doi:10.1067/s0002-9378(03)00753-1 (2003).

Huang, Y., Liu, Y., Huang, J., Gao, L., Wu, Z., Wang, L., & Fan, L. Let-7b-5p promotes cell apoptosis in Parkinson's disease by targeting HMGA2. *Mol. Med. Report.* **24**, 820, doi:doi.org/10.3892/mmr.2021.12461 (2021).

Liu, J., Chen, Z., Cui, Y., Wei, H., Zhu, Z., Mao, F., Wang, Y., & Liu, Y. Berberine promotes XIAP-mediated cells apoptosis by upregulation of miR-24-3p in acute lymphoblastic leukemia. *Aging* **12**, 3298-3311, doi:doi.org/10.18632/aging.102813 (2020).

Ma, Y. *et al.* miR-10a-5p Promotes Chondrocyte Apoptosis in Osteoarthritis by Targeting HOXA1. *Mol Ther Nucleic Acids* **14**, 398-409, doi:10.1016/j.omtn.2018.12.012 (2019).

Gimpl, G., & Fahrenholz, F. The oxytocin receptor system: structure, function, and regulation. *Physiol. Rev.* **81**, 629-683, doi:10.1152/physrev.2001.81.2.629 (2001).

LoParo, D. & Waldman, I. D. The oxytocin receptor gene (OXTR) is associated with autism spectrum disorder: a meta-analysis. *Mol. Psychiatry* **20**, 640-646, doi:10.1038/mp.2014.77 (2015).

Petroff, O. A. GABA and glutamate in the human brain. *Neuroscientist* **8**, 562-573,

doi:10.1177/1073858402238515 (2002).

59   Hussman, J. P. Suppressed GABAergic inhibition as a common factor in suspected etiologies of autism. *J. Autism Dev. Disord.* **31**, 247-248, doi:10.1023/a:1010715619091. (2001).

60   Ma, D. Q., Whitehead, P. L., Menold, M. M., Martin, E. R., Ashley-Koch, A. E., Mei, H., Ritchie, M. D., Delong, G. R., Abramson, R. K., Wright, H. H., Cuccaro, M. L., Hussman, J. P., Gilbert, J. R., & Pericak-Vance, M. A. Identification of significant association and gene-gene interaction of GABA receptor subunit genes in autism. *Am. J. Hum. Genet.* **77**, 377-388, doi:10.1086/433195 (2005).

Acknowledgements

We would like to express our sincere gratitude to Prof. Eileen McLaughlin at University of Wollongong for the online discussions and many helpful suggestions. We would also like to express our deep gratitude to Prof. Takashi Shinohara at Kyoto University Graduate School of Medicine and Dr. Hiroko Hamada at Tohoku University Graduate School of Medicine for their various advice on experiments and teaching experimental skills. We would also like to express our deepest gratitude to Prof. Yasuhisa Matsui, Prof. Takahiro Arima, Mr. Jasper Germeraad at Tohoku University Graduate School of Medicine for their critical reading and valuable feedback, as well as our excellent technical staff, Ms. Sayaka Makino, for taking care of the experimental mice and preparing the experimental environment.

Data availability statement

Microarray data were deposited into the Gene Expression Omnibus database under accession number GSE232620 and are available at the following URL: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE232620.

Authors` contribution

MT, and NO designed the study. MT, KM, and TK performed the experiments. MT and KM undertook the bioinformatic and statistical analyses. KM, MT and NO wrote the manuscript. All authors contributed and have approved the final manuscript.

Funding information

This work was supported by a grant from The Canon Foundation.

Conflict of interest

The authors declare no conflict of interest.

Figure legends

Figure 1. Experimental diagram for the strategy of analyzing miRNAs. Sperm samples were collected from the cauda epididymis of three C57BL/6J mice at 3M, > 12M, and > 20M via retrograde perfusion. Total RNA, including miRNAs, was extracted from the collected samples and subjected to microarray analyses. Further details can be found in the text.

Figure 2. Analyses of differentially expressed miRNAs with aging. (**A**) Venn diagram showing the number of miRNAs expressed in sperm samples from mice aged 3M, 12M and 20M, respectively. A total of 447 miRNAs were expressed. White letters indicate the numbers of common miRNAs and their percentages against the total numbers. (**B**) Differentially expressed miRNAs were defined as those showing > 2-fold up- or down-regulation and $P < 0.05$. Red bars indicate upregulated miRNAs and green bars indicate downregulated miRNAs. (**C-E**) Volcano plots showing differentially expressed miRNAs in sperm samples from mice aged 3M vs 12M (**C**), 3M vs 20M (**D**), and 12M vs 20M (**E**). Differentially expressed miRNAs were defined as those showing > 2-fold up- or down-regulation and $P < 0.05$. Red dots indicate upregulated miRNAs and green dots indicate downregulated miRNAs. (**F**) Venn diagram showing differentially expressed miRNAs in sperm samples comparing 3M vs 12M, 3M vs 20M, and 12M vs 20M. A total of 46 differentially expressed miRNAs were identified. White letters indicate the number of common miRNAs and their percentage against the total number.

Figure 3. Pathway analyses of miRNA target genes. Pathway analyses of miRNA target genes were performed on miRNA target genes significantly altered the sperm samples of 3M vs 12M, 3M vs 20M, and 12M vs 20M, respectively using the Reactome pathway analysis package in R. The number of genes associated with each pathway is shown on the horizontal axis in (**A-C**). The gene ratio in (**D**) indicates the percentage of genes associated with each pathway among the genes listed as targets of the miRNAs.

Figure 4. Network of biological pathways and genes. Network of significantly enriched pathways and target genes belonged to those pathways when comparing 3M vs 20M (**A**) and 12M vs 20M　(**B**). Node color indicates the fold change of the gene expression. Node size indicates the significance of the differential expression; larger sizes indicate smaller *p* values.

Figure 5. Correlation analysis between apoptosis-related genes, *Ppp3r1* and *Bcl2l11*, and their regulator miRNAs. Scatter plot shows the correlation between *Ppp3r1* and *mmu-let-7b-5p* (**A**), *Bcl2l11* and *mmu-miR-10a-5p* (**B**), and *Bcl2l11* and *mmu-miR-24-3p* (**C**).

Figure 6. Network of miRNAs and SFARI genes. Network of significantly altered miRNAs and their target autism-related SFARI genes when comparing 3M vs 20M (**A**) and 12M vs 20M (**B**). Node color indicates the fold change of the miRNA and target gene expression. Node size indicates the significance of the differential expression; larger sizes indicate smaller *p* values.

Figure 7. Expression levels of specific miRNAs. (**A-D**) Correlation analyses between SFARI genes whose expression levels were varied between groups and their regulatory miRNAs. (**A**) *mmu-miR-466j* vs *Oxtr*, (**B**) *mmu-miR-24-3p* vs *Gabrb2*, (**C**) *mmu-miR-690* vs *Ctnnb1*, (**D**) *mmu-let-7b-5p* vs *Grik2*. (**E-I**) Expression scores in this microarray analysis of miRNAs that are previously reported to be inherited from sperm to fertilized egg [41]. (**E**) *miR-10a-5p*, (**F**) *miR-146a-5p*, (**G**) *miR-34c-5p*, (**H**) *miR-143-3p*, and (**I**) *miR-146b-5p*. Bars show the mean of the signals of the three samples in each group. Bonferroni's multiple comparison test was used to test for significant differences. *$P < 0.05$, ** $P < 0.01$.

Figure 1

Figure 2

Figure 3

(A)

Enrichment Analysis of target genes of
differentially expressed miRNAs between **3M vs 12M**

(B)

Enrichment Analysis of target genes of
differentially expressed miRNAs between **3M vs 20M**

(C)

Enrichment Analysis of target genes of
differentially expressed miRNAs between **12M vs 20M**

(D)

Figure 4

(A)

(B)

Fold change

-5.0   1.0   5.0

Figure 5

(A)

(B)

(C)

Figure 6

(A)

(B)

Figure 7