Genetic diversity, determinants, and dissemination of *Burkholderia pseudomallei*

lineages implicated in melioidosis in northeast Thailand

Rathanin Seng[1], Chalita Chomkatekaew[2,3], Sarunporn Tandhavanant[1], Natnaree Saiprom[1],

Rungnapa Phunpang[1], Janjira Thaipadungpanit[4], Elizabeth M Batty[2,5], Nicholas PJ Day[2,5],

Wasun Chantratita[6], T. Eoin West[1,7,8], Nicholas R Thomson[9], Julian Parkhill[3], Claire

Chewapreecha[2,4,9,10]*, Narisara Chantratita[1,2]*

[1] Department of Microbiology and Immunology, Faculty of Tropical Medicine, Mahidol University,

Bangkok, Thailand.

[2] Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol

University, Bangkok, Thailand.

[3]Department of Veterinary Medicine, University of Cambridge, UK

[4]Department of Clinical Tropical Medicine, Faculty of Tropical Medicine, Mahidol University,

Bangkok, Thailand.

[5]Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of

Oxford, Oxford, UK.

[6]Center for Medical Genomics, Faculty of Medicine Ramathibodi Hospital, Mahidol University,

Bangkok, Thailand

[7]Division of Pulmonary, Critical Care & Sleep Medicine, Department of Medicine, University of

Washington, Seattle, Washington, USA.

[8] Department of Global Health, University of Washington, Seattle, Washington, USA.

[9]Parasites and Microbes, Wellcome Sanger Institute, Cambridge, UK.

[10] Previous Affiliations: Bioinformatics and Systems Biology Program, School of Bioresource

and Technology, King Mongkut University of Technology Thonburi, Bangkok, Thailand.

*Contribute equally

For sequencing, and data analysis - correspond to Claire Chewapreecha

(claire@tropmedres.ac)

For sample collection, sampling processing and metadata - correspond to Narisara Chantratita

(narisara@tropmedres.ac)

Abstract

Melioidosis is an often-fatal neglected tropical disease caused by an environmental bacterium *Burkholderia pseudomallei*. However, our understanding of the disease-causing bacterial lineages, their dissemination, and adaptive mechanisms remains limited. To address this, we conducted a comprehensive genomic analysis of 1,391 *B. pseudomallei* isolates collected from nine hospitals in northeast Thailand between 2015 and 2018, and contemporaneous isolates from neighbouring countries, representing the most densely sampled collection to date. Our study identified three dominant lineages with unique gene sets enhancing bacterial fitness, indicating lineage-specific adaptation strategies. Crucially, recombination was found to drive lineage-specific gene flow. Transcriptome analyses of representative clinical isolates from each dominant lineage revealed heightened expression of lineage-specific genes in environmental versus infection conditions, notably under nutrient depletion, highlighting environmental persistence as a key factor in the success of dominant lineages. The study also revealed the role of environmental factors – slope of terrain, altitude, direction of rivers, and the northeast monsoons - in shaping *B. pseudomallei* geographical dispersal. Collectively, our findings highlight persistence in the environment as a pivotal element facilitating *B. pseudomallei* spread, and as a prelude to exposure and infection, thereby providing useful insights for informing melioidosis prevention and control strategies.

Introduction

Melioidosis, a severe infectious disease, affects an estimated 165,000 cases globally each year, of which 89,000 are fatal[1]. The disease is caused by *Burkholderia pseudomallei*, a Gram-negative bacillus found in soil and contaminated water across tropical and sub-tropical regions. Historically, limited access to microbiology laboratories for culture-confirmed diagnosis led to underreporting, particularly in lower- and middle-income countries[2]. However, improved infrastructure and awareness have led to increases in reported cases across South Asia, Southeast Asia, East Asia[3–8] and Australia[9,10] In Southeast Asia, the disease incidence is often linked to agriculture practice, particularly during the rainy seasons when rice paddy fields are flooded for planting. The flooded terrain enables the bacterium in the soil to surface, potentially exposing farmers to *B. pseudomallei* and subsequently leading to melioidosis[11]. Additionally, many cases of melioidosis have been associated with severe weather events[12–14]. While climate likely influences human encounters with *B. pseudomallei*, further investigation is needed to fully understand the mechanisms linking environmental factors to melioidosis epidemiology.

Understanding the population structure, dissemination and adaptation of *B. pseudomallei* in

these climatically-challenged endemic regions requires a large-scale, geographically and

chronologically densely-sampled, genetic dataset. Previous studies, albeit limited in sample

size, have demonstrated that *B. pseudomallei* dissemination is driven by both anthropogenic

and environmental factors[15–18]. Streams[19,20], monsoons, typhoons and cyclones[13,14,21,22] were

identified as significant contributors to bacterial dissemination, highlighting the importance of

bacterial persistence across a range of environmental conditions. *B. pseudomallei* exhibits

remarkable survival capabilities across diverse environments, spanning from wet to dry,

nutrient-depleted soil[23–28] thereby enabling the bacterium to thrive in various ecological niches.

Previous studies have noted the temporal and geographical co-existence of multiple *B.*

pseudomallei lineages[15,16,29]. However, little is known about their distinct genetic content and

adaptive strategies. Identification of lineage-specific genes associated with bacterial persistence

and disease escalation will be essential to develop disease control strategies.

In this study, we conducted a population genomics analysis using combined *B. pseudomallei*

isolates from melioidosis patients across nine provinces in the northeast Thailand including

Buriram, Khon Kaen, Mahasarakam, Mukdahan, Nakhon Phanom, Roi-Et, Sisaket, Surin and

Udon Thani[8]; totaling 1,265 isolates collected from July 2015 to December 2018. Additionally,

we incorporated contemporary environmental and clinical collections from Thailand and

neighbouring countries[15,29–34], consisting of 15 clinical isolates and 111 environmental isolates

(**Figure 1a**, **Supplementary data 1**). Our comprehensive analysis, including a total of 1,391

isolates, revealed the population structure, dissemination patterns, and genetic diversity of this

bacterium. We identified genetic determinants associated with dominant lineages and

investigated their biological functions and expression conditions in three representative isolates,

each representing a dominant lineage. This provides insights into the strategies employed by *B.*

pseudomallei lineages for successful persistence in the environment, ultimately leading to

human exposure and infection.

Results

Population structure analysis revealed the successful *B. pseudomallei* lineages and

mixture of clinical and environmental isolates

To define the population structure of clinical and environmental *B. pseudomallei* in a

hyperendemic area of northeast Thailand and neighbouring regions (n = 1,391), we performed

four independent approaches. PopPUNK[35] analysis was performed on genome assemblies

(**Supplementary data 2**). Additionally, we constructed three maximum-likelihood (ML)
phylogenies[36], each based on different sets of single nucleotide polymorphisms (SNPs): core
genomes (n= 77,156 SNPs), core gene multilocus typing[37] (cgMLST, n = 46,945 SNPs), and
seven-gene multilocus typing genes[38] (MLST, n = 31 SNPs). These approaches facilitated the
grouping of isolates with close genetic similarity into distinct lineages. Notably, both PopPUNK
analysis and core genome SNPs phylogeny yielded consistent results (**Supplementary Figure**
1) clustering the population into three dominant lineages (**Figure 1b**). The average pairwise
core SNP distance within each dominant lineage was 549, 351, and 517 SNPs for lineage 1, 2,
and 3, respectively, in contrast to the pairwise core SNP distance of 1,087 SNPs within the total
population (**Figure 1c**). This lower pairwise core SNP distance across lineages confirmed the
genetic relatedness as defined by PopPUNK and core genome SNP phylogenetic analysis.
While cgMLST displayed conservation for two out of three dominant lineages, MLST exhibited
inconsistencies across dominant lineages with lower phylogenetic resolution and poorer
bootstrap support compared to other methodologies (**Supplementary Figure 1**). Consequently,
we relied on the population delineated by PopPUNK and core genome SNP phylogeny for
subsequent investigations.
115
The three predominant lineages (denoted as lineage 1 to 3) comprised 312, 297, 125 isolates,
respectively. They accounted for 52.8% of the studied population and persisted throughout the
sampling period. Interestingly, each lineage peaked during the rainy season, correlating
agricultural practices at the onset of rainfall with increased environmental exposure and
subsequent melioidosis infections (**Figure 1d**). Despite the small sample size of environmental
isolates, we observed a clustering of these isolates with clinical isolates within each dominant
lineage, indicating their core genetic similarities and shared origin. The ratio of environmental to
clinical isolates varied across lineages (Chi-square test with Monte Carlo resampling p-value
5.00×10^{-4}, **Supplementary Figure 2**). Due to the substantially lower number of environmental
isolates used and incomplete geographical distribution matching between clinical and
environmental isolates, caution is warranted in interpreting these results. Nevertheless, our
findings highlight a mixing of environmental and clinical samples, suggesting that clinical
isolates could serve as a surrogate for tracking the dissemination of an environmental
bacterium, especially in the absence of equally comprehensive environmental samples.
130
Genetic evidence identifies patterns of *B. pseudomallei* dissemination in Northeast
Thailand

We examined the dissemination patterns of *B. pseudomallei* in northeast Thailand. Except for
Mahasarakam where samples were limited, all three dominant lineages were present across the
rest of eight studied provinces (**Figure 1a**). This prompted us to focus the analysis on these
dominant lineages to identify consistent geographical distributions underlying their spread in the
region. We generated lineage-specific phylogenies to improve genetic resolution for
transmission analysis. Additionally, we reconstructed ancestral histories of provincial origins,
quantified the number of inter-provincial transmissions to examine transmission patterns, and
estimated the time of the most recent common ancestor of each dominant lineage and its sub-
lineages (**Supplementary Figures 3 – 5**, see Methods). This allowed us to link the emergence
of these lineages to historical events that might have affected their transmission dynamics.
While transmission signals could reflect distinct dissemination patterns in each dominant lineage
or its sub-lineages, we also considered the possibility of shared factors leading to a uniform
geographical distribution. Notably, we observed consistent dissemination patterns in 14 out of
28 provincial pairs across the three dominant lineages (**Figure 2a, Supplementary Figure 6**).
Eight out of these 14 provincial pairs potentially correlated with the slope of terrain altitude
between provinces or the natural flow of rivers in the region (**Figure 2b**). These pairs included
"Udon Thani-to-Khon Kaen", "Udon Thani-to-Buriram", "Udon Thani-to-Mukdahan", "Udon
Thani-to-Surin", "Khon Kaen-to-Buriram", "Nakhon Phanom-to-Mukdahan", "Roi-Et-to-Surin" and
"Surin-to-Sisaket". Northeast Thailand is described as a saucer-shaped plateau, with elevations
ranging from over 200 meters above sea level in the northwestern corner (parts of Udon-Thani,
and Khon-Kaen) to less than 100 meters in the southeast (parts of Mukdahan and Sisaket), and
gradually descending toward the Mekong River in the east[39,40]. The Mun River originates from
elevated hills in central Thailand, streaming eastward through Buriram, Surin, Sisaket before
merging with the Mekong River. Similarly, the Chi River, a tributary to the Mun, also originates
from the central Thailand mountains. The Chi flows eastward through Khon Kaen,
Mahasarakam, Roi-Et and converges with the Mun in Sisaket[40].

Additionally, another set of three out of 14 conserved patterns coincided with the wind direction
of the northeastern monsoon during the dry season ("Nakhon Phanom-to-Buriram", "Mukdahan-
to-Buriram", and "Mukdahan-to-Surin"). Thailand experiences two predominant monsoon
seasons: the southwest monsoon from May to October and the northeast monsoon from
November to April. The southwest monsoon brings heavy rainfall from the southwest to
northeast, often marking the start of the agriculture season (**Figure 1c**)[41,42]. Conversely, the
northeast monsoon brings dry winds from the northeast to southwest. Our observation implies

the potential for dry winds to transport aerosolised soil contaminated with *B. pseudomallei*
westward. Despite that previous air sampling during Thailand's rainy season did not detect *B.*
pseudomallei, exploring the impact of dry winds during the northeastern monsoon is essential.
This consideration becomes even more pertinent given reports of the long-range transport of
particles and small organic matters, such as PM2.5 and PM10, via the northeast monsoon
elsewhere in Southeast Asia[43,44]. Furthermore, we successfully estimated the time of most
recent of ancestry for a sub-lineage 1.3, a descendant of lineage 1. Our finding revealed that
this sub-lineage emerged around 2011 (95% HPD of 2000-2014, **Supplementary Figure 3**).
The age of this sub-lineage implies that older lineages, such as its parent lineage 1 likely
experienced multiple monsoon seasons, which possibly resulted in the observed patterns. Apart
from the terrain slope, inland rivers, canal systems[45] and regular monsoons[41,42]; various factors
likely contributed to shaping *B. pseudomallei* dissemination in northeast Thailand.
Anthropogenic activities, such as human migration between the provinces, may also contribute
to the observed pattern; however without access to comprehensive human movement data, this
aspect remains challenging to investigate.

Genetic markers potentially contributing to the emergence of successful *B. pseudomallei*
lineages

The co-existence of multiple *B. pseudomallei* lineages within the same geographical areas and
timeframe implies the presence of diverse adaptive strategies which enable them to thrive in a
shared ecological niche. While some smaller lineages may be sporadically detected, the
persistence of the three dominant lineages throughout the sampling period supports their fitness
and successful adaptive strategies in this niche. We next sought to identify genes that were
present in isolates that form each dominant lineage, or its sub-lineage; but absent in non-
dominant lineages (see Methods). Out of total 15,237 genes in the pan-genome outlined from
this population (see Methods), 5,577 genes were conserved across the entire population while
9,660 genes were variably present (accessory genes). Dominant lineage-specific genes were
defined as accessory genes present in ≥95% of isolates within any of the dominant lineages or
their sub-lineages, but present in ≤15% of isolates outside these lineages. Among these, 247
genes were identified as lineage-specific with their specificity to each dominant lineage and sub-
lineage tabulated in **Supplementary data 3**. The majority of dominant lineage-specific genes
were poorly characterised and annotated as hypothetical proteins (**Figure 3**). To gain insights
into the potential functions, we annotated them using Gene Ontology (GO terms)[46] which
classify them by Biological process, Molecular function, and Cellular component (Figure 3b, see

Methods). Of the 247 dominant lineage genes, GO terms could be assigned to 27 genes for
Biological Process, 68 for Molecular Function, and 12 for Cellular component. For genes that
could be assigned GO terms, functions involved in "DNA integration", "DNA recombination", and
"DNA methylation" might indicate their potential roles in horizontal gene acquisition and
protection against incoming foreign DNA through site-specific DNA methylation. Furthermore,
GO terms associated with "DNA binding" and "Regulation of DNA-templated transcription" may
suggest lineage-specific regulation of the expression of these genes.

Lineage-specific genes were selectively expressed
To delve deeper into the functionality of dominant lineage-specific genes, we explored their
expression patterns across both environmental and infection conditions. We selected
representative strains - "K96243" (lineage 1 – sub-lineage 1.1), "UKMD286" (lineage 2 – sub-
lineage 2.1), and "UKMH10" (lineage 3 – sub-lineage 3.2) - all are clinical isolates with pre-
existing gene expression profiles under infection and environmental conditions[47–49]. For
environmental conditions, K96243 was exposed to water[47], while UKMD286 and UKMH10 were
cultivated in a soil extract medium[48,49] to mimic *B. pseudomallei* in the environment. For
infection conditions, K96243 and UKMD286 were used in murine challenges[47,48] and isolated
from mice organs, while UKMH10 was subjected to human plasma[49] to simulate host infection.
This approach facilitated the comparison of differentially expressed lineage-specific genes
between environmental and infection conditions. Although each representative strain carried a
complete set of lineage-specific genes for their respective sub-lineages (K96243 with 47 genes,
UKMD286 with 27 genes, and UKMH10 with 14 genes), their collective representation
accounted for 69 out of 247 total lineage-specific genes (27.9%) due to observed genetic
diversity within the dominant lineage. Notably, 11 out of 47 lineage-specific genes in K96243
and 6 out of 27 lineage-specific genes in UKMD286 were up-regulated in the environmental
conditions (**Figure 4a to 4c, Supplementary data 4**). Notably, none of the lineage-specific
genes showed up-regulation during the infection condition. The remaining lineage-specific
genes did not exhibit preferential expression in either environmental or infection conditions. The
elevated expression level of lineage-specific genes in the environmental condition was
unexpected considering that all representative strains were clinical isolates. This observation
potentially suggests that dominant lineage-specific genes may play more substantial roles in
bacterial environmental survival than in host pathogenicity.

To thrive in its environmental habitat, *B. pseudomallei* must cope with ranges of physical,
chemical and biological stresses such as desiccation, temperature fluctuations, osmotic
changes, oxidative stress, UV exposure, nutrient scarcity, changes in pH, exposure to heavy
metals, competition from antibiotics released by other microbes, and predation by eukaryotes.
We leveraged the extensive condition-wide transcriptome data spanning 62 conditions available
for K96243[47], a representative strain from lineage 1, to compare the expression patterns of
lineage-1-specific genes with the rest of genes in the K96243 genome (**Figure 4d**). Our analysis
revealed that lineage-1-specific genes within K96243 exhibited a higher level of gene
expression when the bacterial cell experienced nutrient deprivation compared to other genes in
K96243 genome (Two-sided Fisher's exact test p-value = 9.23×10^{-5}). This finding implies that
lineage 1 might possess an adaptive strategy to persist in nutrient-depleted soil, which is not
uncommon in melioidosis endemic areas[27,28,50], before being acquired by a human host and
subsequently causing the disease.
245

Example of lineage-specific genes
The majority of lineage-specific genes were located within genomic islands (GI)[30,51,52]. These
regions are characterised by anomalies in %G+C content or dinucleotide frequency signatures,
or the presence of genes associated with mobile genetic elements such as insertion sequence
(IS) elements and bacteriophages. Notably, we observed that a cluster of genes specific to
lineage 1 (*BPSS2060* to *BPSS2072*) formed a mosaic structure within a putative metabolic
island known as GI 16[30]. Although several variations of GI 16 have been reported (GI16, GI16.1,
GI16.2, GI16a, GI16b, and GI16b.1)[51], it typically spans 60 kb (*BPSS2051* to *BPSS2090*) and
carries several known virulence determinants and genes that enhance metabolic versatility.
While certain virulence factors, such as the filamentous haemagglutinin (*BPSS2053*) required
for host cell adhesion and its processing protein were conserved across multiple lineages
observed in our study, genes encoding functions that potentially expand the metabolic repertoire
were specific to dominant lineages. For example, the mosaic structure of GI 16 (*BPSS2060* to
BPSS2072), specific to lineage 1 (**Supplementary data 3**), contains genes involved in
alternative nutrient catabolism and anabolism (*BPSS2060*, *BPSS2065*, *BPSS2067*, *BPSS2068*,
and *BPSS2072*), transcriptional regulation (B*PSS2061*), and substrate transport (*BPSS2064*,
BPSS2071). Out of 11 lineage-specific genes located in the mosaic structure of GI 16, eight
were found to be upregulated during the early phase of nutrient starvation while remaining silent
during infections. This finding reflects the functional division of GI16 where its lineage-specific
mosaic structure contains genes that contribute to metabolic versatility, while its core structure

encodes virulent determinants associated with disease implications. It is important to note that
the structure of GI 16 may vary across different regions due to the plasticity of genomic islands
and changes in selection pressures. While this observation is significant for a dominant lineage
in northeast Thailand, it may not be generalisable to other geographical locations.

Lineage-specific genes were introduced by homologous recombination
Homologous recombination has been shown to play a significant role in facilitating the
acquisition and loss of genes, and the generation of mosaic structures within the GI of *B.*
pseudomallei[53]. To better understand its association with lineage-specific genes, we identified
recombination events and quantified the rates of recombination in the dominant lineages. The
ratio of polymorphisms introduced through recombination compared to those introduced by
mutation (*r/m*) was 3.7, 4.6 and 2.2 for lineages 1, 2, and 3 respectively (**Table 1)**. A very high
proportion of genes underwent recombination at least once: 99.5% of genes in lineage 1, 99.9%
in lineage 2, and 96.6% in lineage 3. Furthermore, every lineage-specific gene within each
dominant lineage underwent recombination (**Supplementary Figure 8**). The bacterial restriction
modification (RM) systems prevent the invasion of foreign DNA and restrict gene flow between
B. pseudomallei lineages[31]. Notably, components of this system including a type I restriction
system and modification methylase were among dominant lineage-specific genes (*BPSL0947-*
BPSL0948 in lineage 1, and their homologues in lineage 2 and 3). They may act as a barrier for
homologous recombination and potentially modulate lineage-specific genetic diversity. This
highlights the intricate interplay between recombination, lineage-specific genes, and the RM
system in shaping the genetic landscape of *B. pseudomallei* in northeast Thailand.

Discussion
Our analysis of *B. pseudomallei* population genomics enhances our understanding of the
evolution and adaptive strategies employed by the dominant lineages in the melioidosis
hyperendemic region of northeast Thailand and neighbouring countries. Through an
unprecedentedly dense sampling effort between 2015 and 2018, we were able to determine the
co-existence of three dominant lineages, characterise their dissemination patterns, and identify
lineage-specific genes possibly contributing to their success during the studied period. By
analysing transcriptome data from representative strains of each dominant lineage, we gained
further insights into their adaptive strategies, particularly emphasising bacterial persistence in
the environment as crucial for subsequent host acquisition and infection. Nevertheless, our
study has a few limitations.

Due to limited environmental surveillance and the scarcity of environmental isolates in
Southeast Asia, our study primarily relied on clinical isolates. Nonetheless, the co-occurrence of
clinical and environmental isolates within the same lineage, coupled with the direct acquisition of
clinical isolates from the environment, suggests that our findings may hold broader implications
for environmental isolates. Furthermore, the lineage classification report in our study is subject
to potential alterations over time with the introduction of new data. As *B. pseudomallei*
continuously adapts to environmental pressures, the composition of lineages 1, 2, and 3, along
with their respective lineage-specific genes may shift. New lineages with superior fitness or
selective advantages, of different adaptive strategies could potentially outcompete the existing
dominant lineages. Consequently, this may lead to emergence of new lineage classifications in
the future. Continued surveillance efforts including both clinical and environmental samples will
be essential. This ongoing monitoring will be pivotal in identifying alterations within the bacterial
population, uncovering new adaptive strategies, and evaluating their impact on disease
dynamics over time.

Our understanding of the significance of bacterial persistence as a strategy for successful
lineages stems from the observed up-regulation of dominant lineage-specific genes in
environmental conditions, along with the increased expression of lineage-1-specific genes
during nutrient deprivation. However, it is essential to recognise the genetic diversity within each
dominant lineage. The representative strains used in our study carried a subset of the lineage-
specific genes corresponding to their lineage. As a result, lineage-specific genes absent in
these strains, often annotated as hypothetical proteins, remained unexplored in our analysis.
Moreover, our detailed characterisation of lineage-1-specific genes using a comprehensive
transcriptome dataset of 62 distinct conditions might not capture the full spectrum of conditions
encountered by *B. pseudomallei* in its natural habitat. There is a possibility that other adaptive
strategies were overlooked in this study, indicating scope for future exploration. Despite these
shortcomings, our dataset and analysis currently represent one of the most comprehensive
efforts to date. Future research will prioritise the generation of a more extensive condition-wide
transcriptome, covering a broader range of conditions and incorporating strains with diverse
genetic variations to identify other adaptive strategies employed by *B. pseudomallei*.

Our findings underscore the significance of environmental persistence in driving the success of
dominant lineages 1 and 2, notably highlighting lineage-1-specific genes in mediating bacterial

survival under nutrient depletion. It remains uncertain whether lineage 3 adopts a similar
strategy. Nevertheless, our results align with previous soil sampling studies, which consistently
observed a higher prevalence of *B. pseudomallei* in nutrient-depleted compared to nutrient-rich
soil[28,50]. Additionally, a molecular evolutionary study also supports the species' long-term
adaptation to survive nutrient scarcity[27]. The northeast region of Thailand, where our samples
were primarily collected, has inherently low-fertility soil. This is exacerbated by intensified
agriculture, monoculture, excessive synthetic fertilizer use, and poor land management,
resulting in depleted soil nutrients and organic matter[54]. This presents a challenging
environment for *B. pseudomallei* to thrive, thereby potentially selecting for successful lineages
with persistent traits as observed in our study.

Our analyses also highlight the role of various factors such as differences in terrain altitude[55,56],
river flow dynamics[40], and the northeast monsoon[41] contribute to shaping the dissemination
patterns of *B. pseudomallei*. These drivers of dissemination are influenced by both natural and
human activities. For instance, strong winds can carry dried soil particles[57], potentially
containing *B. pseudomallei* over distances. Climate change-induced alterations in vegetation
cover might expose soil to rainfall and winds[58], impacting the bacterial spread. Additionally,
deforestation can disrupt natural barriers like trees and shrubs, accelerating water runoff[57,58] and
potentially facilitating the wide-ranging dissemination of *B. pseudomallei* during flood.
Considering these dynamics, the strategy of bacterial persistence likely plays a pivotal role in its
widespread dissemination within the region, thereby influencing disease prevalence. Therefore,
an effective disease control strategy should integrate both environmental and clinical public
health measures to effectively mitigate the impact of melioidosis.

Materials and Methods

Data collection and bacterial isolates

The *B. pseudomallei* isolates in our study included a cohort[8] from northeast Thailand gathered between July 2015 to December 2018 consisting of 1,265 clinical isolates. We also incorporated a contemporaneous dataset from Thailand and neighbouring regions[15,29–34], comprising 15 clinical and 111 environmental isolates sourced from previous publications. In total, 1,391 *B. pseudomlalei* genomes were used in this study. Their metadata and accession numbers were documented in **Supplementary data 1**.

The northeast Thailand collection was collected from patients participated in our longitudinal cohort study[7]. The patients were from nine provinces including Udon Thani (n = 230), Mukdahan (n = 198), Roi Et (n = 195), Surin (n = 170), Nakhon Phanom (n = 135), Buriram (n = 123), Sisaket (n = 107), Khon Kaen (n = 96) and Maha Sarakham (n = 11), who were admitted to nine hospitals included in our cohort. The ethical approval for the cohort study was obtained from the Ethics Committee of the Faculty of Tropical Medicine, Mahidol University (MUTM 2015-002-001 and MUTM 2021-055-01). The isolates were obtained from various clinical samples, including blood (71.8%), pus (12.5%), sputum (9.9%), body fluid (3.5%), urine (1.9%) and tissue (0.4%) (**Supplementary data 1**). As latent infection accounts for < 5% of the cases, the majority of clinical cases likely directly acquired from the environment[8]. The numbers of enrolled cases were lower at the beginning of the study due to the delayed sample collection in some study sites, resulting in inconsistent number of bacterial isolates used across different sites. When applicable, a permutation test was performed to ensure that an unequal number of isolates did not impact the temporal or spatial analysis.

Culture confirmation of *B. pseudomallei*, DNA extraction and whole genome sequencing

All 1,265 *B. pseudomallei* samples from the northeast Thailand collection were cultured on Ashdown, selective agar plates and confirmed the species using latex agglutination test and matrix-laser absorption ionisation mass spectrometry (MALDI-TOF MS). A single colony from Ashdown agar plate was subjected to culture in Luria-Bertani (LB) broth and subsequently used for DNA extraction. Genomic DNA was extracted using QIAamp DNA Mini Kit (Qiagen, Germany). All genomic DNA were processed for the 150-base-read library preparation and sequencing using Illumina HiSeq2000 system with 100-cycle paired-end runs at Wellcome Sanger Institute, Cambridge UK. An average of 71X read depth was achieved. To control the potential contamination in each sample with other closely related species, we assigned

391   taxonomic identity using Kraken[59] v.1.1.1. We then estimated the genome completeness and
392   species confirmation using CheckM[60] v.1.2.2 and FastANI[61] v.1.31, respectively. The quality
393   control data of the 1,265 genomes were listed in **Supplementary data 2**.
394

Genome assembly and mapping from short read data
Short reads were *de novo* assembled using Velvet v.1.2.10[62] followed by optimisation as
previously described[29], with their quality scores reported in **Supplementary data 2**. Short reads
were also mapped against several reference genomes, including a strain K96243[30] (accession
numbers BX571965 and BX571966) to determine the whole population structure, and lineage-
specific references to improve the resolution for lineage-specific analyses. We selected
K96243[30] genome as a population-wide reference due to its origin in northeast Thailand,
aligning with the geographical focus of our study. Additionally, its well-characterised and
complete genome further streamlined subsequent analyses. For all mapping, variants were
called using Snippy v.4.6.0 (https://github.com/tseemann/snippy). To avoid mapping errors and
false SNPs, we filtered out SNPs covered by less than 10 reads and found in a frequency of
less than 0.9.
407

Defining whole population structure
PopPUNK clustering
PopPUNK v.2.6.0[35] was run on 1,391 assembled genomes. To define the core and accessory
distance between each pair of isolates, the assemblies were hashed at different k-mers. The
population model was fit using command line "poppunk-runner.py --fit-model --distance
<database.dists> --output <database> --full-db --ref-db <database> --min-kmer 15 –max-kmer
31 --max-a-dist 0.53 --K 4 –k-step 2" with the result density of 0.028, transitivity of 0.992, and
network score of 0.8961.
416

Maximum likelihood phylogenies from core genome SNP, cgMLST and MLST
An alignment of full genome was created by mapping whole genome sequences of each *B.*
pseudomallei against a complete genome of K96243[30] strain. From this alignment, 4,221
cgMLST loci based on a scheme described in[37] were extracted and concatenated to form
cgMLST alignment. Additionally, seven MLST loci, as per scheme described in [38] were
extracted from the same alignment and concatenated to create the MLST alignment. Core
genome SNP alignment was identified from a full genome alignment using snp-sites[63] v.2.5.1,
with genomic islands[51] masked. Separate maximum likelihood phylogenies were constructed for

core genome SNP alignment, cgMLST alignment, and MLST alignment using IQ-TREE[36]
v.2.0.3. Standard model selection in IQ-TREE determined the best-fit model as
TVM+F+ASC+R6 for all three phylogenies. To access the robustness of the phylogenetic trees,
a 1,000 bootstrap support was performed for each tree.
429
Comparison of population structure outlined phylogeny constructed from core genome
SNPs, cgMLST, MLST and PopPUNK
To test for consistency between phylogenetic trees constructed from core genome SNPs,
cgMLST and MLST alignment, we use the R package treespace[64] v. 1.1.4.3 to explore the tree
tip distributions. We compared pairwise tree distances within the first 100 bootstraps within each
alignment category (indicative of bootstrap support strength), and across trees generated from
different alignment categories (indicating proximity between tree categories). The tree pairwise
distances were computed, and principal components (PCs) were derived with eigenvalues
calculated for different PCs. The similarity among phylogenies from each alignment category
was assessed using two PCs dimensions, which jointly accounted for >90% of variability in
pairwise distance (**Supplementary Figure 1a**). The scatter plot of PCs revealed a close
clustering of bootstrap trees from core genome SNP and cgMLST, while the bootstrap trees
from MLST alignment showed greater dispersion, highlighting less consistency in the trees
generated by the MLST approach. We further compared the consistency between the median
phylogenetic tree of each alignment category and PopPUNK classification was visually
compared using iTOL[65] (**Supplementary Figure 1b**).
446
Specific lineage analysis
To investigate the dissemination and genetic diversity within each dominant lineage, we
conducted individual genome alignments, recombination removal, and maximum likelihood
phylogeny. To enhance the sensitivity of variant, we selected closely related genomes as
references for each lineage. Specifically, the complete genome *B. pseudomallei* strain K96243
(accession numbers BX571965 and BX571966) served as the mapping reference for lineage 1,
while new reference genomes were created for lineages 2 and 3.
454
For lineage 2, we chose a representative isolate 27035_8#57 and subjected it to long-read
sequencing on a local MinION sequencer following the manufacturer's standard protocol
(Oxford Nanopore Technologies, Oxford, United Kingdom). A complete hybrid assembly of the
long-read and short-read sequence data of this strain was performed using Unicycler[66] v.0.8.4.

The resulting hybrid assembly of the 27035_8#57 genome was employed as the mapping
reference for this lineage.

In the absence of representative complete genomes for lineage 3, we selected the best quality
de novo assembly of isolate 27035_8#119 and orientated its contigs according to strain K96243
using ABACAS[67] v.1.3.1. This genome was used as a mapping reference for lineage 3.

For lineage-specific mapping, Snippy v.4.6.0 as employed as in the whole population analysis.
All genome alignments were subjected to Gubbins[68] v.3.1.3, a recombination identification tool,
to detect and remove recombination fragments. This process determined the genetic diversity
introduced by horizontally acquired elements and vertically inherited SNPs, thereby producing
recombination-free SNP alignments for phylogenetic reconstruction. Maximum-likelihood
phylogenies were constructed using recombination-free SNP alignment of each dominant
lineage using IQ-TREE[36] v.2.0.3 with TVM+F+ASC+R6 and 1,000 replicates of bootstrap
support. The overall proportion of nodes with ≥80% bootstrap support of lineage-specific
phylogenies reached 83.5%.

Dating the timeline for lineages and sub-lineages
To enable dating analysis, we further divided each lineage into sub-lineages using R package
rhierbaps[69] v.1.1.4. We inferred evolutionary timeline and estimated the age of each lineage and
sub-lineage based on the isolate's collection date. A Bayesian molecular dating provided in R
package BactDating[70] v.1.1.1. was employed to assess the temporal signals by examining a
positive correlation between the isolate's collection date and the root-to-tip distance.
Recombination removed phylogenies were used in this analysis. A date-randomisation test,
consisting of 100 permutations, was performed to assess the robustness of the temporal signal
compared to noise.

Notably, the temporal signals were discernable at the sub-lineage level rather than the broader
lineage level. Among the 10 sub-lineages, only one sub-lineage (lineage 1.3) exhibited a
positive correlation in their clock signals. Given the limited sample size of lineage 1.3, we
employed a strict clock model to prevent parameter over-fitting. We ran three independent
Markov chain Monte Carlo (MCMC) chains, each spanning at least 100 million iterations, and
sampled every 10,000 steps. The prior mutation rate derived from Pearson and colleagues was
used. Visual inspection of the trace from each MCMC chain confirmed signal convergence, with

effective sampling size values > 200 for key parameters. Visualisation of results were performed

using the R package ggtree[71] v.3.10.0 to generate credibility time-calibrated phylogeny for each

sub-lineage (**Supplementary Figures 3**).

Ancestral state reconstruction analysis

Ancestral trait reconstruction was conducted to discern the dissemination patterns of *B.*

pseudomallei among provinces in northeast Thailand, focusing on dominant lineages. Due to

varying number of isolates among provinces, the analysis excluded Mahasarakham, which had

a limited dataset (n = 11), resulting in the analysis of eight provinces: Buriram, Khon Kaen,

Mukdahan, Nakhon Phanom, Roi Et, Sisaket, Surin, and Udon Thani. This approach yielded 28

potential province-to-province transmission combinations. To mitigate sampling biases, we sub-

sampled the phylogeny of each dominant lineage to have an equal number of isolates per

province (n = 15 isolates) and permuted 1,000 times. Using the stochastic character mapping

function (*make.simmap*) from the R package phytools[72] v.1.9.16, we conducted 100 simulations

(nsim = 100) to reconstruct the provincial origins at each node in the sub-sampled phylogeny

(1,000 phylogenies per lineage). This allowed us to quantify transition events (Markov jumps)

between province pairs and determine the cumulative branch length associated with each

province (Markov rewards). A Mann-Whitney U test, with Bonferroni correction for multiple

comparisons was applied to compare the transition event counts among provinces

(**Supplementary Figure 6**).

Pan-genome analysis

All the study genomes were annotated using Prokka[73] v.1.14.5, and further used in the pan-

genome analysis. Each genome has a median of 5,845 coding sequences (CDS) predicted onto

each genome with a range of 5,642 to 6,142 CDS per genome. Panaroo[74] v.1.3.3 was

employed to estimate the pan-genome with a sensitive option and a cut-off sequence identity of

92% derived from previous study[15]. The number of estimated genes falls within a comparable

range to previous studies from a single population[29,53].

Identification of dominant lineage-specific genes

We determined lineage-specific genes by assessing their prevalence within dominant lineage or

any of their sub-lineages, requiring a high occurrence (95%) within these specific groups while

maintaining a low presence in non-dominant lineages. To achieve this, three thresholds were

employed: strict (95 % occurrence in dominants vs 5% occurrence in non-dominants),

intermediate (95% in dominants vs 10% in non-dominants), and relaxed (95% in dominants vs

15% in non-dominants). Based on visual examination of gene distribution patterns (**Figure 3a,**

Supplementary Figure 7), the relaxed threshold was used to maximise the number of genes

included in subsequent analysis.

Identification of Gene Ontology (GO: terms)

Amino acid sequences of lineage-specific genes were submitted to InterPro database[46]

(https://www.ebi.ac.uk/interpro/) which characterised the function of lineage-specific genes

based on biological processes, molecular functions, and cellular compartments **(Figure 3;**

Supplementary data 3).

Transcriptomic analysis of dominant lineage-specific genes

Lineage 1 transcriptome analysis

The analysis focused on an expression profile of a strain K96243, which serves as a

representative of lineage 1. Data was sourced from microarray experiment generated by Ooi *et*

al.[47] and accessed through the Gene Expression Omnibus (GEO) under accession number

GSE43205.

To understand the difference between environmental and infection conditions, we compared the

expression profile of K96243 being exposed to water and K96243 recovered from infected mice.

To simulate environmental conditions, K96243 was cultured to log phase in LB medium,

subsequently washed with sterile deionising water, and suspended in water. To emulate

infection conditions, BALB mice were infected with 1000 CFU of *B. pseudomallei*, and bacteria

were harvested from the lungs three days post-infection. Two replicates were performed for

each condition. We retrieved microarray data from GEO using the R package GEOquery[75]

v.2.58.0, and differential gene expression analysis was performed using the R package limma[76]

v.3.58.1.

We used binary expression patterns reported in Ooi *et al.*[47] to compare the expression profiles

of lineage-specific genes against the remaining genes in K96243 across 62 conditions. This

enabled the comparison of the count of expressed genes within the lineage-specific category

against the remainder of the genes for each condition using Fisher's exact test, with multiple

testing adjustments via Benjamini-Hochberg corrections.

Lineage 2 transcriptome analysis

This analysis focused on the expression profile of a strain UKMD286, representative of lineage

2. RNAseq data was obtained from an experiment conducted by Ghazali *et al.*[48] and accessed

through the European Nucleotide Archive (ENA) (E-MTAB-11200).

To simulate environmental conditions, UKMD286 was cultured in BHIB medium overnight,

resuspended, and inoculated into soil extract medium. For infection condition, BALB mice were

infected with UKMD286, and the bacteria were harvested from spleens five days post infection.

Each experiment was conducted with two replicates. FastQC v.0.11.9 and FastXtool v.0.0.14

were used to pre-processed sequenced reads. Raw reads were aligned to UKDM286 genome

using Hisat2[77] v. 2.2.1 with differential gene expression performed using the R package

DESeq2[78] v.1.40.2.

Lineage 3 transcriptome analysis

We used a strain UKMH10 to represent the expression profile of lineage 3. Data was originated

from an RNAseq experiment conducted by Kong *et al.*[49] and was accessed through the

European Nucleotide Archive (ENA) (PRJEB53338)

To replicate environmental conditions, UKMH10 was cultured in LB medium overnight and sub-

cultured into soil extract medium. To simulate infection conditions, UKMH10 was cultured in LB

medium overnight and inoculated into human plasma, then incubated at 37 C to mimic the

human body temperature. Bacterial cells were harvested once the absorbance reading of the

bacterial cultures at 600 nm (OD600) reached 0.5. Each experiment was performed with two

replicates. FastQC v.0.11.9 and FastXtool v.0.0.14 were used to pre-processed sequenced

reads. Raw reads were aligned to UKMH10 genome using Hisat2[77] v.2.2.1 with differential gene

expression performed using the R package DESeq2[78] v.1.40.2.

Acknowledgements

We would like to thank physician and microbiology staff at the Udon Thani Hospital, Khon Kaen

Hospital, Srinakarin Hospital, Nakhon Phanom Hospital, Mukdahan Hospital, Roi Et Hospital,

Surin Hospital, Sisaket Hospital and Buriram Hospital for reporting the case and providing the

facilities for bacterial collection. We sincerely appreciate the transcriptomic data of B.

pseudomallei UKMD286 and UKMH10 provided by Professor Sheila Nathan and colleagues.

We are grateful to Dr. Soe Htet Aung for supporting geographical map plot. This work was

funded by Mahidol University (MU-KMUTT Biomedical engineering and Biomaterials

Consortium) and Royal Golden Jubilee Ph.D. Programme (RGJ-ASEAN) (http://rgj.trf.or.th)

through RS and NC. CCho was funded by Wellcome International Master Fellowship

(221418/Z/20/Z). CChe was funded by Wellcome International Intermediate Fellowship

(216457/Z/19/Z) and Sanger International Fellowship. NC and TEW were supported by the

National Institute of Allergy and Infectious Diseases of the National Institutes of Health

(NIAID/NIH) (https://www.nih.gov) under Award Number U01AI115520. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health. This research was funded in part, by the Wellcome Trust [220211]. For the

purpose of Open Access, the author has applied a CC BY public copyright license to any Author

Accepted Manuscript version arising from this submission. The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data availability

The genome sequence data presented in this study can be found in online repositories. The

ENA under study accession number PRJEB25606 and PRJEB35787. The accession numbers

for individual genomes, and their annotated assemblies are listed in **Supplementary data 1**.

Code availability

The analyses used public software including Kraken v.1.1.1, CheckM v.1.2.2, FastANI v.1.31,

FastQC v.0.11.9, FastXtool v.0.0.14, Unicycler v.0.8.4, ABACAS v.1.3.1, Velvet v.1.2.10,

Prokka v.1.13.4, Panaroo v.1.2.9, Snippy v.4.6.0, PopPUNK v.2.6.0, IQ-TREE v.2.0.3, Gubbins

v.3.1.3, Hisat2 v.2.2.1 and R packages: treespace v.1.1.4.3, BactDating v.1.1.1, ggtree v.3.10.0,

phytools v.1.9.16, GEOquery v.2.58.0, limma v.3.58.1 and DESeq2 v.1.40.2

Ethics and Inclusion statements

The research is led by local researchers and include local contributions throughout all research

process including the study design, study implementation, data ownership, intellectual

properties and authorship for publication.

Author contributions

RS, CChe and NC conceived, designed the study and write the original draft. NC and CChe

administrated and supervised the project. NC acquired funding to collect isolates, while CChe

acquired funding for sequencing and downstream analysis. NC, RS, TEW, and NS collected

and identified bacterial isolates. NC, RS and RP collected clinical data. CChe, NRT, NC, RS,

JT, EB and WC performed whole-genome sequencing. NPJD and NC contributed reagents.

NRT and JP contributed software tools. RS, CCho, and CChe performed bioinformatics

analyses. CChe, NC, NRT and JP interpret the analyses. CChe wrote the revised draft. All

authors read and approved the manuscript.

Table 1 Recombination in dominant lineages.

Lineage	Percent of CDS impacted by recombination at least once (recombinant CDS/total CDSs in the reference genome)	Percent of dominant lineage-specific genes underwent recombination at least once (recombinant gene/lineage-specific genes identified in the reference genome)	Average *r/m* (number of SNPs introduced by recombination/ SNPs introduced by substitutions)		
			Internal nodes (95% CI)	Terminal nodes (95% CI)	Average (95% CI)
1	99.5% (5981/6010)	100% (47/47)	2.8 (2.3-3.3)	4.6 (4.0-5.3)	3.7 (3.3-4.1)
2	99.9% (5963/5971)	100% (65/65)	5.0 (3.9-6.0)	4.3 (3.6-4.9)	4.6 (4.0-5.2)
3	96.6% (5338/5523)	100% (20/20)	1.4 (1.1-1.8)	2.9 (2.2-3.7)	2.2 (1.8-2.6)

 Figure and table legend

Figure 1 Distribution of *B. pseudomallei* genomes used in this study (a) Geographical
representation of the countries and provinces sampled for the 1,391 *B. pseudomallei* genomes
used in this study. Pie-chart summarises the proportion of dominant lineage 1, 2, and 3
presented at each location with the chart size proportional to the number of the samples
collected (b) An unrooted phylogenetic tree colour-coded by dominant lineages (c) Histogram
depicting the distribution of clinical *B. pseudomallei* isolates from the northeast Thailand cohort
throughout 2015-2018 sampling period. The shaded blue area represents the period of rainy
seasons. (d) Boxplots summarising the pairwised core genome SNP distances among isolates
in this study, shown in a logarithmic scale. The distribution is depicted for the entire population
and each dominant lineage.

Figure 2 Dissemination patterns in northeast Thailand. (a) Province-to-province transmission
patterns influenced by northeast Thailand geographical landscape. Nodes present provinces,
denoted by abbreviation and ordered by altitude: U-Udon Thani, K-Khon Kaen, B-Buriram, R -
Roi Et, N-Nakhon Phanom, Su-Surin, M-Mukdahan, and Si-Sisaket. Rivers are depicted in blue
with major rivers including the Great Mekong River, the Chi River, and the Mun River and their
flow direction annotated. (b) Average altitude of provinces in meters above sea level. Error bars
present 95% confidence interval. The northwest provinces exhibit higher altitudes, gradually
declining towards the southeast. For (a) and (b) solid arrows illustrating transmission
directionality explained by altitude differences. Dotted arrows represent transmission
directionality influenced by northeast monsoon winds. Grey arrows signify patterns with unclear
explanation.

Figure 3 Dominant lineage-specific genes and their Gene Ontology (GO terms). (a) The
heatmap represents lineage-specific genes (right) detected in each isolate, aligned with the
phylogeny (left). Lineage-specific genes shared across multiple dominant lineages are
highlighted in yellow. Lineage-specific genes from lineage 1, 2, 3 are coloured in green, red, and
purple, respectively. Additionally, the colour stripes provide information on the lineage and sub-
lineage membership (b) Bar plots displays the frequency of GO annotations of lineage-specific
genes in each dominant lineage categorised by biological process, molecular function, and
cellular compartment. The pie-charts summarise the proportion of lineage-specific genes with
assigned GO terms (black).

Figure 4 Transcriptome analysis of representative strains: K96243 (lineage 1), UKMD286
(lineage 2) and UKMH10 (lineage 3) (a to c) Volcano plots demonstrate differential gene
expression (DGE) between environmental and infection conditions. Vertical dotted lines
represent the statistical cut-off at log two-fold change, while horizontal dotted lines display the
statistical cut-off at the adjusted p-value of 0.05 on a negative log scale. Each dot represents a
gene, with lineage-1, lineage-2, and lineage-3-specific coloured in green, red, and purple,
respectively. (d) Binary expression profile of lineage-1-specific genes across different
conditions. A star denotes significant differences in the gene expression profile of lineage-1-
specific genes compared the remaining genes of strain K96243.

Supplementary Figure 1 Comparison of approaches used in in outlining the population
structure (a) A scatter plot displays the first two principal components (PCs) derived from
pairwise distances between trees. Each dot represents a bootstrap tree and is colour-coded by
the method used to generate the tree: core genome SNP (blue), cgMLST (red) and MLST
(green). (b) A scree plot summarises eigenvalues computed for each PCs. (c to d) The median
phylogenetic trees constructed from core genome SNP, cgMLST, and MLST and their
consistency with PopPUNK clustering method.

Supplementary Figure 2 Distribution of environmental and clinical isolates by each lineage.
Barplots highlight the co-detection of environmental (green) and clinical isolates (red) across
dominant lineages 1, 2, and 3.

Supplementary Figure 3 Lineage 1 specific analysis (a) A recombination removed lineage 1
phylogeny with colour stripes displaying its sub-lineage structure, year of collection, and
sampling province (left to right). (b) A map of northeast Thailand showing the distribution of
each isolate and the region's river system. (c) Time-calibrated phylogeny of sub-lineage 1.3 with
blue error bars indicating 95% highest posterior density interval, with the estimated mutational
rate consistent with previous study.

Supplementary Figure 4 Lineage 2 specific analysis (a) A recombination removed lineage 2
phylogeny with colour stripes highlighting sub-lineage structure, year of collection, and sampling
province (left to right). (b) A map of northeast Thailand with the region's river system. Dots
present the distribution of individual samples.

Supplementary Figure 5 Lineage 3 specific analysis (a) A recombination removed lineage 3
phylogeny with colour stripes highlighting sub-lineage structure, year of collection, and sampling

province (left to right). (b) A map of northeast Thailand showing the distribution of each isolate
and the region's river system.

Supplementary Figure 6 Transmission patterns and evolutionary time spent at each province.
(a to c) Proportion of transition events (Markov jumps) among provincial pairs for lineage 1, 2,
and 3 respectively. The pairs were denoted as province1 – province 2, with transitions from
province 1 to province 2 shown in red, and transitions from province 2 to province 1 in green.
The Man-Whitney U test was conducted for each pair to assess differences in transition
frequency by direction, with Bonferroni correction applied for multiple tests. (d to f) Total branch
length from provincial trait reconstruction (Markov rewards) for lineage 1, 2, and 3, respectively.

Supplementary Figure 7 Selection criteria for lineage-specific genes. Scatter plots show the
frequency distribution of lineage-specific (red) against other genes (black), based on their
distribution within the dominant lineages and their sub-lineages (horizontal axis) compared to
their distribution in non-dominant lineages.

Supplementary Figure 8 Recombination patterns detected in lineage 1, 2, and 3. From left to
right: the recombination-removed phylogeny of each lineage, a stripe representing the sampling
year and sub-lineage classification, and heatmaps displaying recombination patterns identified
in chromosome 1 and 2. The top orange lines mark the genome coordinates. For each lineage,
their respective lineage-specific genes are highlighted in blue at the top of the panel. Each
heatmap represents recombination blocks aligned with the phylogeny. Recombination events
occurring at the internal nodes are coloured in red, while those occurring at the external
branches are coloured in blue. The recombination hotspot is plotted at the bottom of each
heatmap.

Supplementary data 1 Epidemiological data, isolate and accession codes for both short reads
and annotated assembly used in this study deposited in the European Nucleotide Archive (ENA)
(n = 1,391) (provided as a separate excel file).
712
713   **Supplementary data 2** Quality control information of newly sequenced *B. pseudomallei* from
714   northeast Thailand collection (n = 1,265) (provided as a separate excel file).
715
716   **Supplementary data 3** List of dominant lineage-specific genes and their Gene Ontology (GO)
717   terms (provided as a separate excel file).

Supplementary data 4 List of dominant lineage-specific genes and their expression profile under infectious and environmental conditions generated by Ooi *et al.* 2013, Ghazali *et al.* 2023 and Kong *et al.* 2023 (provided as a separate excel file).

References

1. Limmathurotsakul, D. *et al.* Predicted global distribution of Burkholderia pseudomallei and burden of melioidosis. *Nat Microbiol* **1**, 15008 (2016).

2. Birnie, E. *et al.* Global burden of melioidosis in 2015: a systematic review and data synthesis. *The Lancet Infectious Diseases* **19**, 892–902 (2019).

3. Mohapatra, P. R. & Mishra, B. Burden of melioidosis in India and South Asia: Challenges and ways forward. *The Lancet Regional Health - Southeast Asia* **2**, 100004 (2022).

4. Zheng, X., Xia, Q., Xia, L. & Li, W. Endemic Melioidosis in Southern China: Past and Present. *TropicalMed* **4**, 39 (2019).

5. Tran, Q. T. L. *et al.* Child Melioidosis Deaths Caused by *Burkholderia pseudomallei* – Contaminated Borehole Water, Vietnam, 2019. *Emerg. Infect. Dis.* **28**, 1689–1693 (2022).

6. Win, M. M. *et al.* Enhanced melioidosis surveillance in patients attending four tertiary hospitals in Yangon, Myanmar. *Epidemiol. Infect.* **149**, e154 (2021).

7. Thabit, A. M. *et al.* Etiologies of tropical acute febrile illness in West Pahang, Malaysia: A prospective observational study. *Asian Pac J Trop Med* **13**, 115 (2020).

8. Chantratita, N. *et al.* Characteristics and one year outcomes of melioidosis patients in Northeastern Thailand: A prospective, multicenter cohort study. *The Lancet Regional Health - Southeast Asia* **9**, 100118 (2023).

9. Hodgetts, K. *et al.* Melioidosis in the remote Katherine region of northern Australia. *PLoS Negl Trop Dis* **16**, e0010486 (2022).

10. Gassiep1, I., Ganeshalingam, V., Chatfield, M. D., Harris, P. N. A. & Norton, R. E. The epidemiology of melioidosis in Townsville, Australia. *Transactions of The Royal Society of Tropical Medicine and Hygiene* **116**, 328–335 (2022).

11. Hinjoy, S. *et al.* Melioidosis in Thailand: Present and Future. *TropicalMed* **3**, 38 (2018).

12. Bulterys, P. L. *et al.* Climatic drivers of melioidosis in Laos and Cambodia: a 16-year case series analysis. *Lancet Planet Health* **2**, e334–e343 (2018).

13. Chai, L. Y. A. & Fisher, D. Earth, wind, rain, and melioidosis. *The Lancet Planetary Health* **2**, e329–e330 (2018).

14. Liu, X. *et al.* Association of Melioidosis Incidence with Rainfall and Humidity, Singapore, 2003–2012. *Emerg. Infect. Dis.* **21**, 159–162 (2015).

15.   Chewapreecha, C. *et al.* Global and regional dissemination and evolution of
Burkholderia pseudomallei. *Nat Microbiol* **2**, 16263 (2017).

16.   Zheng, H. *et al.* Genetic diversity and transmission patterns of Burkholderia
pseudomallei on Hainan island, China, revealed by a population genomics analysis. *Microbial*
Genomics **7**, (2021).

17.   Sarovich, D. S. *et al.* Phylogenomic Analysis Reveals an Asian Origin for African
Burkholderia pseudomallei and Further Supports Melioidosis Endemicity in Africa. *mSphere*
1, e00089-15 (2016).

18.   Price, E. P., Currie, B. J. & Sarovich, D. S. Genomic Insights Into the Melioidosis
Pathogen, Burkholderia pseudomallei. *Curr Trop Med Rep* **4**, 95–102 (2017).

19.   Mayo, M. *et al.* A cluster of melioidosis cases from an endemic region is clonal and is
linked to the water supply using molecular typing of Burkholderia pseudomallei isolates. *The*
American Journal of Tropical Medicine and Hygiene **65**, 177–179 (2001).

20.   Draper, A. D. K. *et al.* Association of the Melioidosis Agent *Burkholderia pseudomallei*
with Water Parameters in Rural Water Supplies in Northern Australia. *Appl Environ Microbiol*
76, 5305–5307 (2010).

21.   Chen, Y.-L. *et al.* The Concentrations of Ambient Burkholderia Pseudomallei during
Typhoon Season in Endemic Area of Melioidosis in Taiwan. *PLoS Negl Trop Dis* **8**, e2877
(2014).

22.   Chen, P.-S. *et al.* Airborne Transmission of Melioidosis to Humans from Environmental
Aerosols Contaminated with B. pseudomallei. *PLoS Negl Trop Dis* **9**, e0003834 (2015).

23.   Limmathurotsakul, D. *et al.* Melioidosis Caused by *Burkholderia pseudomallei* in Drinking
Water, Thailand, 2012. *Emerg. Infect. Dis.* **20**, 265–268 (2014).

24.   Yip, T.-W. *et al.* Endemic Melioidosis in Residents of Desert Region after Atypically
Intense Rainfall in Central Australia, 2011. *Emerg. Infect. Dis.* **21**, 1038–1040 (2015).

25.   Baker, A. L., Ezzahir, J., Gardiner, C., Shipton, W. & Warner, J. M. Environmental
Attributes Influencing the Distribution of Burkholderia pseudomallei in Northern Australia.
PLoS ONE **10**, e0138953 (2015).

26.   Pumpuang, A. *et al.* Survival of Burkholderia pseudomallei in distilled water for 16 years.
Transactions of the Royal Society of Tropical Medicine and Hygiene **105**, 598–600 (2011).

27.   Chewapreecha, C. *et al.* Co-evolutionary Signals Identify Burkholderia pseudomallei
Survival Strategies in a Hostile Environment. *Mol Biol Evol* **39**, msab306 (2022).

28.   Hantrakun, V. *et al.* Soil Nutrient Depletion Is Associated with the Presence of
Burkholderia pseudomallei. *Appl. Environ. Microbiol.* **82**, 7086–7092 (2016).

29.   Chewapreecha, C. *et al.* Genetic variation associated with infection and the environment
in the accidental pathogen Burkholderia pseudomallei. *Commun Biol* **2**, 428 (2019).

30.   Holden, M. T. G. *et al.* Genomic plasticity of the causative agent of melioidosis,
Burkholderia pseudomallei. *Proceedings of the National Academy of Sciences* **101**, 14240–
14245 (2004).

31.   Nandi, T. *et al. Burkholderia pseudomallei* sequencing identifies genomic clades with
distinct recombination, accessory, and epigenetic profiles. *Genome Res.* **25**, 129–141 (2015).

32.   Liechti, N. *et al.* Whole-Genome Assemblies of 16 Burkholderia pseudomallei Isolates
from Rivers in Laos. *Microbiol Resour Announc* **10**, e01226-20 (2021).

33.   Rachlin, A. *et al.* Whole-genome sequencing of Burkholderia pseudomallei from an
urban melioidosis hot spot reveals a fine-scale population structure and localised spatial
clustering in the environment. *Sci Rep* **10**, 5443 (2020).

34.   Webb, J. R. *et al.* Myanmar Burkholderia pseudomallei strains are genetically diverse
and originate from Asia with phylogenetic evidence of reintroductions from neighbouring
countries. *Sci Rep* **10**, 16260 (2020).

35.   Lees, J. A. *et al.* Fast and flexible bacterial genomic epidemiology with PopPUNK.
Genome Res. **29**, 304–316 (2019).

36.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Molecular Biology and Evolution* **37**, 1530–1534 (2020).

37.   Lichtenegger, S. *et al.* Development and Validation of a Burkholderia pseudomallei Core
Genome Multilocus Sequence Typing Scheme To Facilitate Molecular Surveillance. *J Clin*
Microbiol **59**, e00093-21 (2021).

38.   Godoy, D. *et al.* Multilocus Sequence Typing and Evolutionary Relationships among the
Causative Agents of Melioidosis and Glanders, *Burkholderia pseudomallei* and *Burkholderia*
mallei. *J Clin Microbiol* **41**, 2068–2079 (2003).

39.   Lofjle, E. & Kubiniok, J. Landform Development and Bioturbation of the Khorat Plateau,
Northeast Thailand. *Nat Hist Bull Siam Soc* 199–216 (1996).

40.   NASA. Thailand Elevation Map.
https://www.floodmap.net/elevation/CountryElevationMap/?ct=TH (2020).

41.   Thai Meteorological Department. The Climate of Thailand. (2016).

42.   Arpornrat, T., Ratjiranukool, S., Ratjiranukool, P. & Sasaki, H. Evaluation of Southwest
Monsoon Change over Thailand by High-resolution Regional Climate Model under High RCP
Emission Scenario. *J. Phys.: Conf. Ser.* **1144**, 012112 (2018).

43.   Hien, P. D., Bac, V. T. & Thinh, N. T. H. PMF receptor modelling of fine and coarse

PM10 in air masses governing monsoon conditions in Hanoi, northern Vietnam. *Atmospheric*

Environment **38**, 189–201 (2004).

44.   Hien, T. T. *et al.* Current Status of Fine Particulate Matter (PM2.5) in Vietnam's Most

Populous City, Ho Chi Minh City. *Aerosol Air Qual. Res.* **19**, 2239–2251 (2019).

45.   Floch, P. & Molle, F. Marshalling water resources: A chronology of irrigation

development in the Chi-Mun River basin Northeast Thailand. (2007).

46.   The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing

strong. *Nucleic Acids Research* **47**, D330–D338 (2019).

47.   Ooi, W. F. *et al.* The Condition-Dependent Transcriptional Landscape of Burkholderia

pseudomallei. *PLoS Genet* **9**, e1003795 (2013).

48.   Ghazali, A.-K. *et al.* Transitioning from Soil to Host: Comparative Transcriptome Analysis

Reveals the Burkholderia pseudomallei Response to Different Niches. *Microbiol Spectr* **11**,

e03835-22 (2023).

49.   Kong, C. *et al.* Transcriptional landscape of Burkholderia pseudomallei cultured under

environmental and clinical conditions. *Microbial Genomics* **9**, (2023).

50.   Manivanh, L. *et al.* Burkholderia pseudomallei in a lowland rice paddy: seasonal

changes and influence of soil depth and physico-chemical properties. *Sci Rep* **7**, 3031

(2017).

51.   Tuanyok, A. *et al.* Genomic islands from five strains of Burkholderia pseudomallei. *BMC*

Genomics **9**, 566 (2008).

52.   Bertelli, C. *et al.* Enabling genomic island prediction and comparison in multiple

genomes to investigate bacterial evolution and outbreaks. *Microbial Genomics* **8**, (2022).

53.   Spring-Pearson, S. M. *et al.* Pangenome Analysis of Burkholderia pseudomallei:

Genome Evolution Preserves Gene Order despite High Recombination Rates. *PLoS ONE*

10, e0140274 (2015).

54.   *White gold: the commercialisation of rice farming in the Lower Mekong Basin*. (Palgrave

Macmillan, 2020).

55.   Keyes, C. F. *Isan: Regionalism in Northeastern Thailand*. (Southeast Asia Program,

Department of Asian Studies, Cornell University, 1967).

56.   Charusiri, P., Daorerk, V., Archibald, D., Hisada, K. & Ampaiwan, T. Geotectonic

Evolution of Thailand: A new synthesis. *Journal of the Geological Society of Thailand* 1–20

(2002).

57. Borrelli, P. *et al.* An assessment of the global impact of 21st century land use change on

soil erosion. *Nat Commun* **8**, 2013 (2017).

58. Singh, B. K. *et al.* Climate change impacts on plant pathogens, food security and paths

forward. *Nat Rev Microbiol* **21**, 640–656 (2023).

59. Wood, D. E. & Salzberg, S. L. Kraken: ultrafast metagenomic sequence classification

using exact alignments. *Genome Biol* **15**, R46 (2014).

60. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Res.* **25**, 1043–1055 (2015).

61. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat*

Commun **9**, 5114 (2018).

62. Zerbino, D. R. & Birney, E. Velvet: Algorithms for de novo short read assembly using de

Bruijn graphs. *Genome Res.* **18**, 821–829 (2008).

63. Page, A. J. *et al.* SNP-sites: rapid efficient extraction of SNPs from multi-FASTA

alignments. *Microbial Genomics* **2**, (2016).

64. Jombart, T., Kendall, M., Almagro-Garcia, J. & Colijn, C. TREESPACE: Statistical

exploration of landscapes of phylogenetic trees. *Molecular Ecology Resources* **17**, 1385–

1392 (2017).

65. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic

tree display and annotation. *Nucleic Acids Research* **49**, W293–W296 (2021).

66. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial

genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**, e1005595

(2017).

67. Assefa, S., Keane, T. M., Otto, T. D., Newbold, C. & Berriman, M. ABACAS: algorithm-

based automatic contiguation of assembled sequences. *Bioinformatics* **25**, 1968–1969

(2009).

68. Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of recombinant

bacterial whole genome sequences using Gubbins. *Nucleic Acids Research* **43**, e15–e15

(2015).

69. Tonkin-Hill, G., Lees, J. A., Bentley, S. D., Frost, S. D. W. & Corander, J. Fast

hierarchical Bayesian analysis of population structure. *Nucleic Acids Research* **47**, 5539–

5549 (2019).

70.   Didelot, X., Croucher, N. J., Bentley, S. D., Harris, S. R. & Wilson, D. J. Bayesian
inference of ancestral dates on bacterial phylogenetic trees. *Nucleic Acids Research* **46**,
e134–e134 (2018).
71.   Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. GGTREE⬚: an R package for
visualization and annotation of phylogenetic trees with their covariates and other associated
data. *Methods Ecol Evol* **8**, 28–36 (2017).
72.   Revell, L. J. phytools: an R package for phylogenetic comparative biology (and other
things). *Methods Ecol Evol* **3**, 217–223 (2012).
73.   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–
2069 (2014).
74.   Tonkin-Hill, G. *et al.* Producing polished prokaryotic pangenomes with the Panaroo
pipeline. *Genome Biol* **21**, 180 (2020).
75.   Davis, S. & Meltzer, P. S. GEOquery: a bridge between the Gene Expression Omnibus
(GEO) and BioConductor. *Bioinformatics* **23**, 1846–1847 (2007).
76.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Research* **43**, e47–e47 (2015).
77.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915
(2019).
78.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
906

Figure 1

Figure 2

a

Wind direction of Northeast monsoon

Direction of the Great Mekong River flow

Direction of Mun and Chi rivers flow

U	Udon Thani
K	Khon Kaen
B	Buriram
R	Roi-Et
N	Nakhon Phanom
Su	Surin
M	Mukdahan
Si	Sisaket

Patterns associated with altitude variances and natural flow of rivers

Patterns associated with wind direction of seasonal monsoon

Unexplained patterns

b

Udon Thani
Khon Kaen
Buriram
Roi-Et
Nakhon Phanom
Surin
Mukdahan
Sisaket

Height above sea level (m)

Figure 3

Figure 4

a Lineage 1: K96243

Lineage 1 specific genes
Other genes

environment   infection

−log10(p-val)

log2FC

b Lineage 2: UKMD286

Lineage 2 specific genes
Other genes

environment   infection

−log10(p-val)

log2FC

c Lineage 3: UKMH10

Lineage 3 specific genes
Other genes

environment   infection

−log10(p-val)

log2FC

d Binary expression patterns of Lineage 1 - specific genes mapped to K96243 genome across different conditions

Up-regulation

Down-regulation

** p-val = 0.00765