Vision rivals audition in alerting humans for fast action

Niklas Dietze[1] & Christian H. Poth[2]

[1] https://orcid.org/0000-0001-7103-8717

[2] https://orcid.org/0000-0003-1621-4911

Neuro-Cognitive Psychology and Center for Cognitive Interaction Technology, Bielefeld University, Bielefeld, Germany

Corresponding author:

Niklas Dietze

Bielefeld University

Department of Psychology

P.O. box 10 01 31

D-33501 Bielefeld

niklas.dietze@uni-bielefeld.de

Statements and Declarations

Funding: This research was funded by the Deutsche Forschungsgemeinschaft (DFG; grant number 429119715 to CHP).

Conflicts of interest: The authors declare there are no conflicts of interest.

Ethics approval: The study was in accordance with the ethical guidelines of the German Psychological Association (DGP) and approved by Bielefeld University's ethics committee.

Consent to participate: All participants gave written informed consent before participation.

Availability of data and materials: All data, analysis code, and experiment code have been made publicly available on the Open Science Framework (https://osf.io/7g4d5/). The design, sampling and analysis plan were preregistered (https://osf.io/fud4k).

Author Contributions: Conceptualisation**:** CHP; Methodology: ND, CHP; Software: ND, CHP; Formal analysis: ND, CHP; Investigation: ND, Data Curation: ND, CHP; Writing – Original Draft: ND, CHP; Writing – Review & Editing: CHP, ND; Visualisation: ND; Supervision: CHP; Funding Acquisition: CHP

Acknowledgements: We acknowledge support for the publication costs by the Open Access Publication Fund of Bielefeld University and the DFG.

Abstract

Successful behaviour requires that humans act promptly upon the ubiquitous rapid changes in the environment. Prompt actions are supported by phasic alertness: the increased readiness for perception and action elicited by warning stimuli (*alerting cues*). Audition is assumed to induce phasic alertness for action faster and more strongly than other senses. Here, we show that vision can be equally effective as audition. We investigated the temporal evolution and the effectiveness of visual and auditory alerting for action in a speeded choice task, while controlling for basic sensitivity differences between the modalities that are unrelated to action control (by matching auditory and visual stimuli according to reaction times in a prior simple detection task). Results revealed that alerting sped up responses, but this happened equally fast and equally strong for visual and auditory alerting cues. Thus, these findings argue that vision rivals audition in phasic alerting for prompt actions, and suggest that the underlying mechanisms work across both modalities.

Keywords: cognitive control, action, perception, phasic alertness, arousal, choice reaction, temporal expectation

1. Introduction

Humans must rapidly perceive and act upon sudden changes in the environment. For instance, drivers have to slam on the brake when a pedestrian or deer suddenly cuts the road. In such situations, split seconds of reaction time can be vital for survival. To handle time-sensitive situations, the human brain is equipped with powerful alerting mechanisms that support fast action upon stimuli in the environment (Hackley, 2009; Petersen & Posner, 2012). *Alerting effects* are the behavioural signature of these mechanisms: Responses in visual detection and discrimination tasks are faster (Dietze & Poth, 2022; Fan et al., 2002; Hackley, 2009; Poth, 2020), more accurate (Matthias et al., 2010; Petersen et al., 2017; Wiegand et al., 2017), or faster at the cost of errors (Han & Proctor, 2022; McCormick et al., 2019; Posner et al., 1973) when target stimuli are preceded by warning stimuli (alerting cues). Alerting effects are assumed to arise because alerting cues induce phasic alertness, a temporary state of arousal that heightens the overall readiness for perception and action (Posner & Petersen, 1990; Sturm & Willmes, 2001). Arguing for such a global, unspecific modulation of perception and action, the performance benefits occur even though alerting cues provide no information about what response would be correct for the impending target. Thus, rather than promoting a specific modulation through low-level influences on the detection of target stimuli, alerting seems to also support the higher-level processes, such as decision-making for selecting among response alternatives (Fan et al., 2002; Hackley, 2009; Poth, 2020) and perceptual encoding for object recognition (Haupt et al., 2019; Petersen et al., 2017; Wiegand et al., 2017). In line with these global benefits for cognitive processing, alerting can temporarily restore spatially impaired visual perception in neurological patients (Robertson et al., 1998) and alerting trainings can accelerate visual perception in aging (Penning et al., 2021). Alongside clinical applications, the beneficial effects of alerting cues on action in safety-

critical situations have long been recognised and used to guide the design of warning

signals (Wogalter & Mayhorn, 2005).

Audition elicits the fastest and strongest alerting effects in humans (Bertelson &

Tisseyr, 1969; Davis & Green, 1969; Harvey, 1980) and non-human primates (Chapman et

al., 1986; Spidalieri et al., 1983). It has been proposed that auditory stimuli are more

arousing than stimuli in other modalities such as vision (Nissen, 1977; Ulrich & Mattes,

1996). Across visual tasks, reactions have been found sped up, when auditory alerting

cues preceded visual targets shortly (< 150 ms; Posner et al., 1976), but the effect

dissipated for longer cue-target onset asynchronies (CTOAs). The effect seems to be

phasic and automatic, as it arises quickly but cannot be retained to prolong the readiness

for perception and action (Posner et al., 1976). In contrast, this pattern was not observed

for visual alerting cues, which has been taken as evidence that the visual modality cannot

deliver the fast, short-lasting, and automatic alerting effect. At least for longer CTOAs (≥

150 ms), it was found that auditory alerting cues and visual alerting cues can induce

similar reaction time benefits when controlled for location cueing effects caused by

different target and warning signal locations (Rodway, 2005). This shows that the task

paradigm and temporal dynamics play a crucial role in alerting effects.

Audition has higher temporal resolution than vision (Artieda & Pastor, 1996), so that

auditory alerting cues should affect performance across CTOAs in a temporally more

precise way. Across modalities it has been shown, that reaction times generally decrease

with increasing CTOAs (Näätänen, 1971). This has been interpreted as evidence that

participants' temporal expectation for the target increases over time, because all used

CTOAs were equally likely, so that with every passing CTOA, targets became more and

more likely to appear (Niemi & Näätänen, 1981). Temporal expectation contributes to

enhanced response readiness (Nobre & van Ede, 2017), and it can support action in

addition or interaction to automatic alerting (Lu et al., 2014). The classic studies that

compared vision and audition in alerting used fixed or uniformly distributed CTOAs

(Bertelson & Tisseyr, 1969; Davis & Green, 1969). Thus, differences in the alerting

capability could stem from interactions between their intrinsic temporal dynamics and

mechanisms for temporal expectation. Therefore, temporal expectation has to be taken

into account when visual and auditory alerting are compared in their effectiveness. It is

assumed that this can be achieved by drawing CTOAs from non-aging probability

distributions, in which the probability that a target appears now given it has not appeared

yet is constant over time (Näätänen, 1971). Such a manipulation has been shown to keep

the temporal expectation of the targets constant, and thus reduces its confound with

response readiness (Petersen et al., 2017; Weinbach & Henik, 2012). Crucially, it is

unknown how differences in auditory and visual alerting effects unfold over longer time-

periods using non-aging probability distributions.

The advantage of audition over vision does not necessarily stem from privileged

access of audition to mechanisms preparing for perception and action, but could rather be

a by-product of audition's low-level characteristics. That is, reaction times have been found

to decline with increasing psychophysical intensity of alerting cues (Adams & Behar, 1966)

and auditory processing is generally faster than visual processing (Grondin, 2010; Hove et

al., 2013), as transduction times are higher in the auditory than the visual system

(Recanzone, 2009; Torre et al., 1995). All this hints at that audition dominates vision in

alerting for action due to differences in their low-level characteristics. Arguing against this

hypothesis, however, are findings tying audition more closely to mechanisms for action

control. Specifically, auditory stimuli seem to result in greater activations in the motor

cortex compared with visual stimuli (Fujioka et al., 2012; Grahn & Rowe, 2009). This larger

neuronal activity could ultimately translate to faster responses, meaning that audition

indeed has a privileged access to alerting for action. For the discussed reasons, the

question if audition indeed dominates vision in phasic alertness is still wide open. The

privilege of audition for alerting could be inherited from audition's higher low-level

sensitivity and processing speed.

Besides the behavioural effects, differences in alerting can be investigated through

physiological markers: Alertness is assumed to be regulated by neuromodulation of

cortical processing from the locus coeruleus-norepinephrine system (Aston-Jones &

Cohen, 2005; Petersen & Posner, 2012; Posner & Petersen, 1990). Bursts of

norepinephrine impact the arousal level and thereby support a wide range of cognitive

functions. This neuronal activation can be indirectly assessed with changes in pupil size

(Nieuwenhuis et al., 2011; Reimer et al., 2016). The size of the pupil increases with the

intensity of an alerting cue (Petersen et al., 2017), in line with the idea that it reflected

intensity-dependent arousal. Thus, assuming that this pupil dilation was a measure for

arousal (e.g., Aston-Jones & Cohen, 2005; Joshi & Gold, 2020; Poth, 2021), the

advantages of audition for eliciting phasic alertness due to increased arousal should

become manifest in stronger pupil dilations for auditory alerting cues than for cues in other

modalities. In addition to pupil responses, saccade latencies have also been linked to the

current state of alertness (Bocca & Denise, 2006). It has been found that saccadic

latencies are facilitated when preceded by an alerting cue (Ross & Ross, 1980). For these

reasons, not only reaction times but also physiological arousal measures should be

included in comparisons between alerting modalities.

Therefore, to settle this issue, we investigated the effectiveness and temporal evolution

of visual and auditory alerting in a choice reaction task, while equating for the simple

detection times between the modalities. First, participants performed a simple detection

task to identify the intensities of auditory and visual alerting cues that yielded to similar

reaction times (Experiment 1). Following the matching procedure, participants performed a

speeded choice with the individually determined intensities of visual and auditory alerting

cues (Experiment 2). However, unlike matching procedures focusing on perceived

stimulus intensities (cf. Stevens & Marks, 1965), we explicitly did not equalise the stimulus intensities at a perceptual level but regarding speeded responses. Using reaction times, we controlled for the speed advantage of audition with respect to earlier processing that is required for a simple reaction without a choice. In contrast to simple reactions, choice reactions measure additional processes of action control, such as response selection and error avoidance (Stuss et al., 2005). To control for temporal expectation, CTOAs across trials were drawn from a non-aging probability distribution, following the gold standard outlined above (Weinbach & Henik, 2012). Surprisingly, however, we still found traces of temporal expectation in our results, casting doubt on a clear dissociation of temporal expectation and alerting. Our findings show that with matched alerting cues based on the known reaction times in a simple detection task, visual alerting is just as effective as auditory alerting, both in terms of behavioural performance as well as physiological arousal responses (pupil dilation and latencies of spontaneous saccades). As such, the findings show that vision can rival audition when the stimulus intensities are matched based on the simple detection times of visual and auditory alerting cues. Phasic alertness seems to arise equally from different senses, which may hint at modality-general alerting mechanisms.

2. Method

2.1. Participants

Seven participants from the university including one of the authors (participant ND) took part in the preregistered experiments (https://osf.io/fud4k). They were between 21 and 41 years old (*median* = 27 years), six were female and one was male. Due to the psychophysical nature of the present investigation and the large number of trials (7000 trials in Experiment 1, 52500 trials in Experiment 2), we limited the sample size to seven participants (one more participant than initially preregistered because one of the authors

participated in the study) which should suffice to establish the existence of effects in the

population as long as all participants show qualitatively similar results (Anderson &

Vingrys, 2001; Smith & Little, 2018). All participants reported normal or corrected-to-

normal vision, normal hearing, and gave written informed consent before participation. The

study was in accordance with the ethical guidelines of the German Psychological

Association (DGP) and approved by university's ethics committee.

2.2. Apparatus and stimuli

Both experiments took place in the same dark room with the only light sources being the

display monitor and the operator monitor located behind the participant. Participants were

seated in front of the preheated display monitor (Poth & Horstmann, 2017) at a viewing

distance of 71 cm with their head on a chin-and-forehead rest. Eye-movements were

recorded with an Eyelink 1000 (SR Research, Ottawa, ON, Canada) at a sampling rate of

1000 Hz. The CRT monitor (G90fB, ViewSonic, Brea, CA, USA) ran at a refresh rate of 85

Hz and a resolution of 1024 x 768. Stimuli were controlled using MATLAB R2014b (The

MathWorks, Natick, MA, USA) and the Psychophysics toolbox (Brainard, 1997; Kleiner et

al., 2007; Pelli, 1997). Responses were collected with a standard external computer

mouse placed in front of the participants. Auditory stimuli were presented using

loudspeakers placed 45 cm to the left and right of screen centre (Philips Multimedia

Speaker System A 1.2 Fun Power/MMS 101, Philips, Amsterdam). All visual stimuli were

white figures (102.6 cd/m^2) presented on a black background (0.03 cd/m^2). A filled circle

with a diameter of 0.28° of visual angle was used as a fixation point for both experiments.

In Experiment 1, the target stimulus was either a larger frame with a diameter of 5° of

visual angle presented at the centre of the screen or a sine tone with a frequency of 700 or

900 Hz (both frequencies occurring equally often and in random order across trials) to

avoid habituation effects. In Experiment 2, the same stimuli were used as alerting stimuli.

The target stimulus was a filled square of 0.5° of visual angle which appeared either 6.5°

to the left or to the right of screen centre. To check the precise timing of the custom

experimental program, we measured the auditory and visual stimuli externally (cf. Poth et

al., 2018) with a microphone capsule and a BPW-34 photodiode, sampled at 2.5 kHz using

a TDS 2022 B oscilloscope (Tektronix, Beaverton, OR, USA). Fifteen runs, in which an

auditory alerting stimulus or a visual alerting stimulus preceding a visual target stimulus

with CTOAs from 153 ms to 529 ms were measured. We found that the CTOAs for the

visual stimuli were on average 0.2 ms shorter than programmed. For the runs with both

modalities, the CTOAs were on average 1.5 ms shorter.

2.3. Procedure of Experiment 1

The first experiment served as a baseline measurement to match visual and auditory cues

based on the trimmed mean reaction times. At the beginning of the experiment and after

250 trials or 20 broken fixations, a 9-point calibration of the eye-tracker was conducted. In

this experiment, 10 logarithmically spaced stimulus intensities for both modalities were

presented each 50 times randomly throughout the session after a random fixation period of

750 ms to 1250 ms drawn from a uniform distribution for a duration of 50 ms. Participants

had to respond as quickly as possible to the stimulus onset. An additional 1500 ms for

pupillometry followed after response collection. Visual intensity varied from 5.79 cd/m² to

102.6 cd/m² and auditory intensity varied from 36.6 db(A) to 64.2 db(A). Each participant

conducted one session (1000 trials for each participant, 7 x 1000 = 7000 trials for the

whole sample). On average, the time to complete a session was about 65 min.

2.4. Procedure of Experiment 2

In Experiment 2, the matched stimuli based on the trimmed mean reaction times in the

prior simple detection task for each participant were used to measure the alerting effects

for both modalities. The same calibration conventions as in Experiment 1 were applied to Experiment 2. Trials started with the presentation of the fixation point. After a random fixation period of 750 ms to 1250 ms drawn from a uniform distribution, either an auditory cue (1/3 of trials), a visual cue (1/3 of trials) or no cue (1/3 of trials) was presented for 50 ms. The CTOAs were drawn from a non-aging probability distribution in steps of 47.04 ms from 153 ms to 3965 ms with a probability of 0.1. After the waiting period, the target stimulus was presented for 200 ms. If participants moved their eyes outside of the fixation window of 2.5° before target onset, the trial was randomly repeated during the remaining sequence. Participants were asked to respond as quickly as possible to the location of the (6.5° to the left or to the right of screen centre) target by pressing the corresponding mouse button (Fig. 1). Again, an additional 1500 ms for pupillometry followed after response collection. Each participant conducted 10 sessions (10 x 750 = 7500 trials for each participant, 7 x 10 x 750 = 52500 trials for the whole sample). However, technical problems prevented recording of 268 trials resulting to a final sample of 52232 trials. Controlling for fatigue, no more than two sessions were completed on the same day. Between two sessions a maximum of seven days elapsed. All sessions were completed on average within a period of six weeks (*min* = 14 days, *max* = 94 days). The time to complete a session was about 60 min. Before and after each session, participants filled out a short questionnaire on subjective relaxation as part of a larger assessment across studies (cf. Steghaus & Poth, 2022).

2.5. Behavioural analyses

Statistical analyses were performed in R (4.0.5, R Core Team, 2021). Following the preregistered protocol, reaction times for each experimental condition were compared using a Bayesian multilevel model created in the Stan computational framework (https://mc-stan.org) accessed with the R-package brms (2.16.3, Bürkner, 2017). We used

a Bayesian model instead of a frequentist approach to quantify evidence in favour of the

null hypothesis. In addition to several other advantages, Bayesian models are more robust

with small sample sizes (Stegmueller, 2013). We analysed reaction times as a function of

alerting condition and CTOA with a random intercept per participant, a random slope per

alerting condition and a random slope per CTOA. The model fit was assessed with the

expected log predictive density based on approximate leave-one-out cross-validations. For

both fixed effects and the random effects of the final multilevel model, we specified

conservative priors. The model was run with 4 MCMC chains and 4000 iterations for each

chain. Complementary, pairwise comparisons between the alerting effects (mean

difference between no cue trials and alert trials) across all CTOAs and participants were

tested using paired *t*-tests with Cohen's d_z effect size (Cohen, 1988) and Bayesian *t*-tests

using standard settings of the R-package BayesFactor (0.9.12-42; Morey & Rouder, 2021),

whose Bayes factors quantify evidence in favour of the null hypothesis (BF_{01}) or evidence

in favour of the alternative hypothesis (BF_{10}). All data processing and visualisations were

made using the tidyverse libraries (Wickham et al., 2019) and data.table (Dowle et al.,

2023). Error trials (1.5 %), trials with CTOAs greater than 1518 ms (3.9 %) and trials for

which the reaction time was greater than 2.5 SD (2.3 %) were removed from the analyses.

Table 1 Matched stimulus intensities for each participant based on the mean reaction times.

Participant	Auditory intensity Decibel (A)	RT (ms) Mean (SD)	Visual intensity Candela (cd/m²)	RT (ms) Mean (SD)
1	36.6	206 (31)	8.11	208 (33)
2	52.3	272 (77)	55.56	265 (51)
3	36.6	280 (90)	11.27	273 (88)
4	36.6	259 (73)	15.60	271 (74)
5	36.6	340 (71)	11.27	351 (72)
6	36.6	261 (80)	11.27	266 (74)
7	64.2	307 (15)	102.6	300 (84)

2.7. *Pupil analyses*

The average pupil size after the pupil normalised from the pupillary light reflex was

computed and compared between conditions. Baseline pupil diameter on each trial was

defined as the mean pupil diameter in the interval 100 ms prior to cue onset. We used the

subtraction method to compute baseline corrected pupil responses for each participant.

Mean alerting induced pupil dilations (mean difference between no cue trials and alert

trials) for each participant were compared using paired *t*-tests with Cohen's d_z effect size

and Bayesian *t*-tests. The same exclusion criteria as the behavioural analyses were

applied to the pupil data.

1.1. *Saccade analyses*

Mean saccade latencies of the first saccade relative to target onset were extracted and

compared between alerting conditions (mean difference between no cue trials and alert

trials). We also used paired *t*-tests with Cohen's d_z effect size and Bayesian *t*-tests. On top

Figure 2 Manual responses of Experiment 2. Participants' mean reaction times fitted
with a polynomial regression in the three experimental conditions. Confidence bands
depict the 95 % confidence interval.

of the previous exclusion criteria, trials were not used if the latency of the first saccade was

less than 0 ms or greater than 1000 ms (15.8 %) relative to target onset.

3. Results

1.1. Behavioural results

The matched intensities of Experiment 1 are presented in Table 1 and the mean reaction

time results of Experiment 2 are visualised in Figure 2. In Experiment 2, reaction times

were generally shorter when preceded by a visual alerting , $t(209) = 18.490$, $p < .001$, $d_z =$

1.276, $BF_{10} = 2.9$ x 10^{42}, and an auditory alerting cue, $t(209) = 17.434$, $p < .001$, $d_z =$

1.203, $BF_{10} = 1.8$ x 10^{39}, compared to the no cue condition (cf. Fan et al., 2002; Hackley,

2009; Poth, 2020). Differences in reaction times between trials with an alerting cue and

trials without an alerting cue diminished with increasing CTOAs. In the early CTOAs, the

phasic alerting effect was about 30 ms. In later CTOAs, it decreased up to a point where

differences between conditions were no longer present. This was particularly due to the

change in the no cue condition. Reaction times decreased with longer CTOAs which is

surprising considering the use of non-aging probability distributions to control for temporal

expectation. Thus, it seems that traces of temporal expectation contributed to the present

results. We also found that visual alerting was not inferior to auditory alerting. Based on

the observations of the posterior distribution, auditory alerting was with 98.5 % confidence

not better than visual alerting. This means that the hypothesis being tested (auditory

alerting > visual alerting) was in 98.5 % of the posterior samples more likely than the

alternative hypothesis (auditory alerting ≤ visual alerting). The posterior mean of the

difference was -3.3 ms with a 95 % credible interval from -6.2 ms to -0.4 ms. On average

participants responses were 24.2 ms (SD = 19.0 ms) faster following the visual alerting

cue and 20.9 ms (SD = 17.4 ms) faster following the auditory alerting cue (Fig. 3). In

summary, near 100 % of the tested cases revealed that auditory alerting was not superior

to visual alerting. This was supported by the pairwise comparisons, $t(209)$ = -4.319, p =

1.000, d_z = -0.298, BF_{01} = 73217, showing that reaction times following an auditory alerting

cue were not shorter than reaction times following a visual alerting cue. Importantly, these

findings cannot be explained by a speed-accuracy trade-off as the overall error rate was

low (1.5 %) and differences between the cue conditions only very minor (no cue: 0.4 %,

visual: 0.6 %, auditory cue: 0.5 %). Pairwise comparisons showed no differences between

the no cue and visual alerting cue, $t(209)$ = -0.292, p = .771, d_z = 0.020, BF_{01} = 12.429, no

cue and auditory alerting cue, $t(209)$ = -0.476, p = .635, d_z = 0.033, BF_{01} = 11.593, and

visual alerting cue and auditory alerting cue, $t(209)$ = 0.215, p = .830, d_z = 0.015, BF_{01} =

12.671. Note, that these findings did not change with the exclusion of participant ND. The

same analyses without participant ND are reported in the supplementary material.

1.2. *Pupillary results*

Pupil responses were observed in all three cue conditions (Fig. 4 a). As such, part of the

responses must be attributed to target onset and detection. Nevertheless, differences

between conditions provide an estimate about the arousal induced by the alerting signals

(Petersen et al., 2017). Directional pairwise comparisons between the pupil alerting

effects, $t(209) = 0.381$, $p = .352$, $d_z = 0.026$, $BF_{01} = 0.545$, revealed indecisive evidence on

the hypothesis that auditory alerting is superior to visual alerting. However, pairwise

comparisons testing the non-directional hypothesis, $t(209) = 0.381$, $p = .704$, $d_z = 0.026$,

$BF_{01} = 12.068$, showed that the pupil dilations were not different between modalities. On

average pupil responses from baseline for the interval from 1500 ms to 1600 ms relative to

cue onset were .053 mm *(SD = .123 mm)* following the visual alerting cue and .052 mm

(SD = .115 mm) following the auditory alerting cue.

1.1. *Saccadic latency results*

As an additional eye-movement marker for phasic alertness, we investigated the latencies

of the oculomotor response. Similar to the manual reaction times, saccadic latencies after

Figure 4 Pupil responses and saccadic latencies of Experiment 2. **a** Grand averages of pupil dilation for the three experimental conditions relative to cue onset. Shaded area shows the analysed time period after the pupil recovered from the pupillary light reflex. Confidence bands depict the 95 % confidence interval. **b** Saccadic latencies fitted with a polynomial regression in the three experimental conditions. Confidence bands depict the

target onset were shortened when preceded by a visual alerting cue, $t(209) = 6.501$, $p <$

.001, $d_z = 0.449$, $BF_{10} = 1.4 \times 10^7$, and an auditory alerting cue, $t(209) = 7.467$, $p < .001$, d_z

$= 0.515$, $BF_{10} = 3.1 \times 10^9$, compared to the no cue condition (Fig. 4 b) (cf. Ross & Ross,

1980). Again, using directional pairwise comparisons between the saccadic alerting

effects, $t(209) = 0.124$, $p = 0.451$, $d_z = 0.009$, $BF_{01} = 0.822$, revealed indecisive evidence

on the hypothesis that auditory alerting leads to faster saccade latencies than visual

alerting. But, pairwise comparisons testing the non-directional hypothesis, $t(209) = 0.124$,

$p = 0.902$, $d_z = 0.009$, $BF_{01} = 12.865$, showed that the saccadic alerting effects are not

different between modalities. On average saccadic latencies were 18.7 ms ($SD = 41.8$ ms)

faster following a visual alerting cue and 19.1 ms ($SD = 37.0$ ms) faster following an

auditory alerting cue compared to the condition without a preceding cue. Overall, the

saccade responses occurred on average 205 ms after the motoric response (Fig. 5 a),

which suggests that the motoric response delayed the temporal execution of an eye-

movement (Reimer et al., 2020). Note, however, most of the spontaneous saccades

Figure 5 Inter-response intervals and saccadic landing positions of Experiment 2. **a** Frequency distribution of the inter-response interval, the differences between reaction times and saccade latencies. Negative inter-response intervals indicate that the manual responses were executed prior to the saccades. **b** Density plot of average eye position after the first saccade in degrees of visual angle. Bright yellow squares show the locations with the highest density of saccade landing positions. Red X's depict the two possible

landed close to screen centre (Fig. 5 b). Only 1.6 % of the saccades were directed towards

the target locations (no cue: 0.5 %, visual: 0.5 %, auditory cue: 0.6 %). Most importantly,

across all saccades no differences in the horizontal landing positions were found between

the no cue and visual alerting cue, $t(209) = -1.657$, $p = .099$, $d_z = -0.114$, $BF_{01} = 3.375$, no

cue and auditory alerting cue, $t(209) = -1.947$, $p = .053$, $d_z = -0.134$, $BF_{01} = 2.032$, and

visual alerting cue and auditory alerting cue, $t(209) = 0.021$, $p = .984$, $d_z = 0.001$, $BF_{01} =$

12.960. Thus, although participants' saccades were facilitated, we did not find that visual

or auditory alerting promoted orienting towards the target.

4. Discussion

We discovered that in contrast to common views, visual and auditory alerting supports fast

action equally well. We employed a novel procedure for matching intensities of alerting

cues according to reaction times in a prior simple detection task to control for low-level

differences between audition and vision. For alerting cues with matched intensities,

auditory and visual alerting both elicited faster responses, and this was the case across

CTOAs up to around 1000 ms. This phasic alerting effect was consistent for both

modalities across all participants. Crucially, these findings contradict classic studies that

have reported large auditory advantages of 31 ms with a foreperiod of 30 ms (Bertelson &

Tisseyr, 1969) and 36 ms with a foreperiod of 50 ms (Davis & Green, 1969). In the present

study, we used a wide range of CTOAs including very short foreperiods drawn from a non-

aging probability distribution. Nevertheless, auditory alerting was never significantly better

than visual alerting. In fact, participants responses were on average even slightly faster

(for 3.4 ms) with a preceding visual cue. Thus, this indicates that the previously reported

advantages of audition for alerting (Bertelson & Tisseyr, 1969; Davis & Green, 1969;

Harvey, 1980) could be reflected by the higher sensitivity and processing speed of

audition. Once controlled for these low-level characteristics of audition, vision rivals audition in alerting humans for perception and action.

Increasing the level of alertness is also expressed in physiological arousal responses. For instance, the pupil dilates once participants are alerted (Aston-Jones & Cohen, 2005; Petersen et al., 2017), and the magnitude of dilation seems positively correlated with intensity levels of the alerting cue (Petersen et al., 2017). Thus, for the present study, differences in pupil responses would highlight any modality specific influences. However, we observed similar pupil responses between the auditory and visual alerting modality. Although at the individual level, pupil responses varied due to differences in intensity levels, dilations in the no cue condition were always systematically weaker than the alerting conditions. In addition, we found a similar relationship in saccadic latencies. Saccadic latencies were equally facilitated when preceded by auditory and visual alerting cues. In contrast to manual reaction times, spontaneous saccades are more automatic and less prone to confounding factors such as the speed-accuracy trade-off and response criterion (Ross & Ross, 1980). Therefore, these patterns of results offer converging evidence that there is no auditory dominance in phasic alertness.

In the present study, we focused on visual targets for two main reasons. First, studies that have reported an auditory dominance found the greatest effects on behaviour with auditory stimuli preceding visual targets (Bertelson & Tisseyr, 1969; Davis & Green, 1969; Harvey, 1980). Second, the visual domain is most relevant for action control (Posner et al., 1976). Previous accounts have postulated that cross-modal auditory advantages have been caused by different locations for visual alerting cues and visual targets (Rodway, 2005; Turatto et al., 2002). In classic experiments, visual cues typically appear at different locations than the visual target while auditory alerting cues only convey little spatial information (Bertelson & Tisseyr, 1969; Davis & Green, 1969; Harvey, 1980; Posner et al., 1976). Hence, it is easier to shift attention from an auditory cue to a visual

target than from a visual cue to a visual target (Rodway, 2005). This is in line with the

hypothesis that visual input typically engages more attention than any other modality

(Posner et al., 1976). Since attention is highly focused to the visually cued location, it

makes it more difficult to shift attention away to the following visual target (Posner et al.,

1976). One way to minimise this location cueing effect, would be to use cues that do not

convey any spatial properties (Wright & Richard, 2003). For our present study we used a

relatively large white frame presented around screen centre which should not have caused

any location specific sensory processing (Matthias et al., 2010). Even when the present

manipulation did not fully control for such an influence, our results argue against a cost

induced by an additional attentional shift caused by the visual warning signal. That is, we

did not find any auditory advantage in the early CTOAs. However, the present data cannot

rule out any differences in CTOAs below 153 ms.

We found that the magnitude of the phasic alerting effect decreased with

longer CTOAs. In particular, it was observed that the mean reaction times in the condition

without a cue converged in an exponential fashion with the other two alerting conditions at

around 500 ms. This contradicts with findings suggesting no changes in phasic alertness

with CTOAs up to 1000 ms (Lu et al., 2014). Moreover, it has been reported when using

non-aging probability distributions that reaction times slightly increase as a function of

CTOA (Lu et al., 2014; Saban et al., 2019). Typically, faster reaction times with increased

CTOAs are only observed when using aging or accelerated-aging probability distributions

(Fuentes & Campoy, 2008; Lu et al., 2014). This is because the probability for the target

appearance becomes greater over time, which in turn leads to a build-up in temporal

expectation (Coull, 2009). The present pattern of results was quite similar to previous

studies that did not use non-aging probability distributions, and thus questions the

procedure to control for temporal expectation. Here, we used a fine-grained temporal

resolution with time-steps of 47 ms and many CTOAs per bin up to 1518 ms which draws

a clear picture of the temporal evolution of reaction times. This suggests that response

readiness cannot be completely dissociated from temporal expectancy when using non-

aging probability distributions as both processes might be triggered by the alerting signal.

In fact, more recently it was proposed that that the hazard rate, which is kept constant

using non-aging probability distributions, may not be suitable to account for temporal

expectancy (see also Grabenhorst et al., 2019).

The present study focused on the most common warning modalities: vision and

audition. It would be interesting to find out whether the same findings on alerting extend to

other modalities such as touch and temperature. Future studies using the same paradigm

would benefit from smaller intensity increments of the alerting cues, so that the detection

times could be equated more precisely. In addition, a monitor with ultra-high temporal

resolution (Poth et al., 2018) could potentially unravel minor differences in the time-course

between the modalities.

In conclusion, we found that alerting works equally well with visual and auditory

alerting cues as long as low-level differences are controlled for. This delivers first time

evidence that phasic alertness exerts its effects in a modality-general way because the

underlying mechanisms do not grant privileged access to a particular sense.

References

Adams, C., & Behar, I. (1966). Stimulus change properties of the RT ready signal.

Psychonomic Science, 6, 389–390. https://doi.org/10.3758/BF03330951

Anderson, A. J., & Vingrys, A. J. (2001). Small samples: Does size matter? *Investigative*

Opthalmology & Visual Science, 47(7), 1411–1413.

Artieda, J., & Pastor, M. A. (1996). Neurophysiological mechanisms of temporal

perception. In *Time, Internal Clocks and Movement* (115. Aufl., S. 1–25).

Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus—Norepinephrine function: Adaptive gain and optimal performance. *Annual Review of Neuroscience*, *28*(1), 403–450. https://doi.org/10.1146/annurev.neuro.28.061604.135709

Bertelson, P., & Tisseyr, F. (1969). The time-course of preparation: Confirmatory results with visual and auditory warning signals. *Acta Psychologica*, 145–154. https://doi.org/10.1016/0001-6918(69)90047-X

Bocca, M.-L., & Denise, P. (2006). Total sleep deprivation effect on disengagement of spatial attention as assessed by saccadic eye movements. *Clinical Neurophysiology*, *117*(4), 894–899. https://doi.org/10.1016/j.clinph.2006.01.003

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Bürkner, P.-C. (2017). brms: An R package for bayesian multilevel models using Stan. *Journal of Statistical Software*, *80*(1). https://doi.org/10.18637/jss.v080.i01

Chapman, C. E., Spidalieri, G., & Lamarre, Y. (1986). Activity of dentate neurons during arm movements triggered by visual, auditory, and somesthetic stimuli in the monkey. *Journal of Neurophysiology*, *55*(2), 203–226. https://doi.org/10.1152/jn.1986.55.2.203

Cohen, J. (1988). *Statistical power analysis for the behavioral sciences.* (2nd ed.). Lawrence Erlbaum Associates.

Coull, J. T. (2009). Neural substrates of mounting temporal expectation. *PLoS Biology*, *7*(8), e1000166. https://doi.org/10.1371/journal.pbio.1000166

Davis, R., & Green, F. A. (1969). Intersensory differences in the effect of warning signals on reaction time. *Acta Psychologica*, *30*, 155–167. https://doi.org/10.1016/0001-6918(69)90048-1

Dietze, N., & Poth, C. H. (2022). Phasic alertness is unaffected by the attentional set for orienting. *Journal of Cognition*, *5*(1), 46. https://doi.org/10.5334/joc.242

Dowle, M., Srinivasan, A., Gorecki, J., Chirico, M., Stetsenko, P., Short, T., Lianoglou, S.,

Antonyan, E., Bonsch, M., Parsonage, H., Ritchie, S., Ren, K., Tan, X., Saporta, R.,

Seiskari, O., Dong, X., Lang, M., Iwasaki, W., Wenchel, S., … Schwen, B. (2023).

data.table: Extension of „data.frame". https://cran.r-

project.org/web/packages/data.table/index.html

Fan, J., McCandliss, B. D., Sommer, T., Raz, A., & Posner, M. I. (2002). Testing the

efficiency and independence of attentional networks. *Journal of Cognitive*

Neuroscience, *14*(3), 340–347. https://doi.org/10.1162/089892902317361886

Fuentes, L. J., & Campoy, G. (2008). The time course of alerting effect over orienting in

the attention network test. *Experimental Brain Research*, *185*(4), 667–672.

https://doi.org/10.1007/s00221-007-1193-8

Fujioka, T., Trainor, L. J., Large, E. W., & Ross, B. (2012). Internalized timing of

isochronous sounds is represented in neuromagnetic beta oscillations. *Journal of*

Neuroscience, *32*(5), 1791–1802. https://doi.org/10.1523/JNEUROSCI.4107-

11.2012

Grabenhorst, M., Michalareas, G., Maloney, L. T., & Poeppel, D. (2019). The anticipation

of events in time. *Nature Communications*, *10*(1), 5802.

https://doi.org/10.1038/s41467-019-13849-0

Grahn, J. A., & Rowe, J. B. (2009). Feeling the beat: Premotor and striatal interactions in

musicians and nonmusicians during beat perception. *Journal of Neuroscience*,

29(23), 7540–7548. https://doi.org/10.1523/JNEUROSCI.2018-08.2009

Grondin, S. (2010). Timing and time perception: A review of recent behavioral and

neuroscience findings and theoretical directions. *Attention, Perception, &*

Psychophysics, *72*(3), 561–582. https://doi.org/10.3758/APP.72.3.561

Hackley, S. A. (2009). The speeding of voluntary reaction by a warning signal. *Psychophysiology*, *46*(2), 225–233. https://doi.org/10.1111/j.1469-8986.2008.00716.x

Han, T., & Proctor, R. W. (2022). Effects of a neutral warning signal on spatial two-choice reactions. *Quarterly Journal of Experimental Psychology*, *75*(4), 754–764. https://doi.org/10.1177/17470218211037604

Harvey, N. (1980). Non-informative effects of stimuli functioning as cues. *Quarterly Journal of Experimental Psychology*, *32*(3), 413–425. https://doi.org/10.1080/14640748008401835

Haupt, M., Ruiz-Rizzo, A. L., Sorg, C., & Finke, K. (2019). Phasic alerting effects on visual processing speed are associated with intrinsic functional connectivity in the cingulo-opercular network. *NeuroImage*, *196*, 216–226. https://doi.org/10.1016/j.neuroimage.2019.04.019

Hove, M. J., Fairhurst, M. T., Kotz, S. A., & Keller, P. E. (2013). Synchronizing with auditory and visual rhythms: An fMRI assessment of modality differences and modality appropriateness. *NeuroImage*, *67*, 313–321. https://doi.org/10.1016/j.neuroimage.2012.11.032

Joshi, S., & Gold, J. I. (2020). Pupil size as a window on neural substrates of cognition. *Trends in Cognitive Sciences*, *24*(6), 466–480. https://doi.org/10.1016/j.tics.2020.03.005

Kleiner, M., Brainard, D. H., Pelli, D., Ingling, A., Murray, R., & Broussard, C. (2007). What's new in psychtoolbox-3. *Perception*, *36*(14), 1–16.

Lu, S., Wang, W., & Cai, Y. (2014). Temporal expectancy modulates phasic alerting in both detection and discrimination tasks. *Psychonomic Bulletin & Review*, *22*(1), 235–241. https://doi.org/10.3758/s13423-014-0664-8

Matthias, E., Bublak, P., Müller, H. J., Schneider, W. X., Krummenacher, J., & Finke, K. (2010). The influence of alertness on spatial and nonspatial components of visual attention. *Journal of Experimental Psychology: Human Perception and Performance*, *36*(1), 38–56. https://doi.org/10.1037/a0017602

McCormick, C. R., Redden, R. S., Hurst, A. J., & Klein, R. M. (2019). On the selection of endogenous and exogenous signals. *Royal Society Open Science*, *6*(11), 190134. https://doi.org/10.1098/rsos.190134

Morey, R. D., & Rouder, J. N. (2021). *BayesFactor: Computation of Bayes Factors for common designs.* https://cran.r-project.org/web/packages/BayesFactor/index.html

Näätänen, R. (1971). Non-aging fore-periods and simple reaction time. *Acta Psychologica*, *35*(4), 316–327. https://doi.org/10.1016/0001-6918(71)90040-0

Niemi, P., & Näätänen, R. (1981). Foreperiod and simple reaction time. *Psychological Bulletin*, *89*(1), 30. https://doi.org/10.1037/0033-2909.89.1.133

Nieuwenhuis, S., De Geus, E. J., & Aston-Jones, G. (2011). The anatomical and functional relationship between the P3 and autonomic components of the orienting response: P3 and orienting response. *Psychophysiology*, *48*(2), 162–175. https://doi.org/10.1111/j.1469-8986.2010.01057.x

Nissen, M. J. (1977). Stimulus intensity and information processing. *Perception & Psychophysics*, *22*(4), 338–352. https://doi.org/10.3758/BF03199699

Nobre, A. C., & van Ede, F. (2017). Anticipated moments: Temporal structure in attention. *Nature Reviews Neuroscience*, *19*(1), 34–48. https://doi.org/10.1038/nrn.2017.141

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spatial Vision*, *10*(4), 437–442.

Penning, M. D., Ruiz-Rizzo, A. L., Redel, P., Müller, H. J., Salminen, T., Strobach, T., Behrens, S., Schubert, T., Sorg, C., & Finke, K. (2021). Alertness training increases

visual processing speed in healthy older adults. *Psychological Science*, *32*(3), 340–

353. https://doi.org/10.1177/0956797620965520

Petersen, A., Petersen, A. H., Bundesen, C., Vangkilde, S., & Habekost, T. (2017). The

effect of phasic auditory alerting on visual perception. *Cognition*, *165*, 73–81.

https://doi.org/10.1016/j.cognition.2017.04.004

Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years

after.   *Annual   Review   of   Neuroscience*,   *35*(1),   73–89.

https://doi.org/10.1146/annurev-neuro-062111-150525

Posner, M. I., Klein, R., Summers, J., & Buggie, S. (1973). On the selection of signals.

Memory & Cognition, *1*(1), 2–12. http://dx.doi.org/10.3758/BF03198062

Posner, M. I., Nissen, M. J., & Klein, R. M. (1976). Visual dominance: An information-

processing account of its origins and significance. *Psychological Review*, *83*(2),

157–171. https://doi.org/10.1037/0033-295X.83.2.157

Posner, M. I., & Petersen, S. E. (1990). The attention system of the human brain. *Annual*

Review   of   Neuroscience,   *13*(1),   25–42.

https://doi.org/10.1146/annurev.ne.13.030190.000325

Poth, C. H. (2020). Phasic alertness reverses the beneficial effects of accessory stimuli on

choice reaction. *Attention, Perception, & Psychophysics*, *82*(3), 1196–1204.

https://doi.org/10.3758/s13414-019-01825-1

Poth, C. H. (2021). Urgency forces stimulus-driven action by overcoming cognitive control.

ELife, *10*, e73682. https://doi.org/10.7554/eLife.73682

Poth, C. H., Foerster, R. M., Behler, C., Schwanecke, U., Schneider, W. X., & Botsch, M.

(2018). Ultrahigh temporal resolution of visual presentation using gaming monitors

and   G-Sync.   *Behavior   Research   Methods*,   *50*(1),   26–38.

https://doi.org/10.3758/s13428-017-1003-6

Poth, C. H., & Horstmann, G. (2017). Assessing the monitor warm-up time required before

a psychological experiment can begin. *The Quantitative Methods for Psychology*,

13(3), 166–173. https://doi.org/10.20982/tqmp.13.3.p166

R Core Team. (2021). *R: A language and environment for statistical computing.* R

Foundation for Statistical Computing. https://www.R-project.org/

Recanzone, G. H. (2009). Interactions of auditory and visual stimuli in space and time.

Hearing Research, *258*(1–2), 89–99. https://doi.org/10.1016/j.heares.2009.04.009

Reimer, C. B., Tudge, L., & Schubert, T. (2020). Saccadic eye movements in dual tasking:

No impairment of spatial planning, but delayed execution of saccades. *Quarterly*

Journal   of   Experimental   Psychology,   *74*(2),   326–343.

https://doi.org/10.1177/1747021820960693

Reimer, J., McGinley, M. J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D. A., & Tolias,

A. S. (2016). Pupil fluctuations track rapid changes in adrenergic and cholinergic

activity   in   cortex.   *Nature   Communications*,   *7*(1),   13289.

https://doi.org/10.1038/ncomms13289

Robertson, I. H., Mattingley, J. B., Rorden, C., & Driver, J. (1998). Phasic alerting of

neglect patients overcomes their spatial deficit in visual awareness. *Nature*,

395(6698), 169–172. https://doi.org/10.1038/25993

Rodway, P. (2005). The modality shift effect and the effectiveness of warning signals in

different   modalities.   *Acta   Psychologica*,   *120*(2),   199–226.

https://doi.org/10.1016/j.actpsy.2005.05.002

Ross, L. E., & Ross, S. M. (1980). Saccade latency and warning signals: Stimulus onset,

offset, and change as warning events. *Perception & Psychophysics*, *27*(3), 251–

257. https://doi.org/10.3758/BF03204262

Saban, W., Weinbach, N., & Gabay, S. (2019). Monocular channels have a functional role

in phasic alertness and temporal expectancy. *Attention, Perception, &*

Psychophysics, *81*(3), 752–763. https://doi.org/10.3758/s13414-018-01653-9

Smith, P. L., & Little, D. R. (2018). Small is beautiful: In defense of the small-N design.

Psychonomic Bulletin & Review, *25*(6), 2083–2101.

Spidalieri, G., Busby, L., & Lamarre, Y. (1983). Fast ballistic arm movements triggered by

visual, auditory, and somesthetic stimuli in the monkey. II. Effects of unilateral

dentate lesion on discharge of precentral cortical neurons and reaction time.

Journal of Neurophysiology, *50*(6), 1359–1379.

https://doi.org/10.1152/jn.1983.50.6.1359

Steghaus, S., & Poth, C. H. (2022). Assessing momentary relaxation using the Relaxation

State Questionnaire (RSQ). *Scientific Reports*, *12*(1), 16341.

https://doi.org/10.1038/s41598-022-20524-w

Stegmueller, D. (2013). How many countries for multilevel modeling? A comparison of

frequentist and Bayesian approaches. *American Journal of Political Science*, *57*(3),

748–761. https://doi.org/10.1111/ajps.12001

Stevens, J. C., & Marks, L. E. (1965). Cross-modality matching of brightness and

loudness. *Proceedings of the National Academy of Sciences*, *54*, 407–411.

Sturm, W., & Willmes, K. (2001). On the functional neuroanatomy of intrinsic and phasic

alertness. *NeuroImage*, *14*(1), S76–S84. https://doi.org/10.1006/nimg.2001.0839

Stuss, D. T., Alexander, M. P., Shallice, T., Picton, T. W., Binns, M. A., Macdonald, R.,

Borowiec, A., & Katz, D. I. (2005). Multiple frontal systems controlling response

speed. *Neuropsychologia*, *43*(3), 396–417.

https://doi.org/10.1016/j.neuropsychologia.2004.06.010

Torre, V., Ashmore, J., Lamb, T., & Menini, A. (1995). Transduction and adaptation in

sensory receptor cells. *The Journal of Neuroscience*, *15*(12), 7757–7768.

https://doi.org/10.1523/JNEUROSCI.15-12-07757.1995

Turatto, M., Benso, F., Galfano, G., & Umiltà, C. (2002). Nonspatial attentional shifts

between audition and vision. *Journal of Experimental Psychology: Human*

Perception and Performance, *28*(3), 628–639. https://doi.org/10.1037/0096-

1523.28.3.628

Ulrich, R., & Mattes, S. (1996). Does immediate arousal enhance response force in simple

reaction time? *The Quarterly Journal of Experimental Psychology Section A*, *49*(4),

972–990. https://doi.org/10.1080/713755672

Weinbach, N., & Henik, A. (2012). Temporal orienting and alerting – the same or different?

Frontiers in Psychology, *3*(26). https://doi.org/10.3389/fpsyg.2012.00236

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund,

G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S.,

Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., … Yutani, H. (2019).

Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686.

https://doi.org/10.21105/joss.01686

Wiegand, I., Petersen, A., Finke, K., Bundesen, C., Lansner, J., & Habekost, T. (2017).

Behavioral and brain measures of phasic alerting effects on visual attention.

Frontiers in Human Neuroscience, *11*(176).

https://doi.org/10.3389/fnhum.2017.00176

Wogalter, M., & Mayhorn, C. (2005). Providing cognitive support with technology-based

warning systems. *Ergonomics*, *48*(5), 522–533.

https://doi.org/10.1080/00140130400029258

Wright, R. D., & Richard, C. M. (2003). Sensory mediation of stimulus-driven attentional

capture in multiple-cue displays. *Perception & Psychophysics*, *65*(6), 925–938.

https://doi.org/10.3758/BF03194824

Supplemental material

Using the sample without participant ND (n = 6), we also found that visual alerting was not

inferior to auditory alerting. Auditory alerting was with 98.2 % confidence not better than

visual alerting. This means that the hypothesis being tested (auditory alerting > visual

alerting) was in 98.2 % of the posterior samples more likely than the alternative hypothesis

(auditory alerting ≤ visual alerting). The posterior mean of the difference was -3.6 ms with

a 95 % credible interval from -7.1 ms to -0.3 ms. On average participants responses were

26.6 ms (SD = 19.1 ms) faster following the visual alerting cue and 23.0 ms (SD = 17.7

ms) faster following the auditory alerting cue (Fig. S1). This was supported by the pairwise

comparisons, $t(179)$ = -4.151, p = 1.000, d_z = -0.309, BF_{01} = 549, showing that reaction

times following an auditory alerting cue were not shorter than reaction times following a

visual alerting cue.

Figure S1 Alerting effects of Experiment 2 without participant
ND. Based on the multilevel model, solid red points show the
fitted values for the aggregated alerting effects of each
participant and CTOA. Error bars depict the 95 % credible