Deferoxamine as adjunct therapeutics for tuberculosis.

Sandeep R. Kaushik[1], Nidhi Yadav[1], Ashish Gupta[1], Sukanya Sahu[1], Nikhil Bhalla[1], Poonam

Dagar[1], Amit Kumar Mohapatra[1], Adyasha Sarangi[1], Taruna Sharma[2], Bichitra Biswal[3], Amit

Kumar Pandey[2], Ranjan Kumar Nanda[1,*]

[1]Translational Health Group, International Centre for Genetic Engineering and

Biotechnology, New Delhi-110067, India.

[2]Mycobacterial Pathogenesis Laboratory, Centre for Tuberculosis Research, Translational

Health Science and Technology Institute, Faridabad, Haryana-121004, India.

[3]Structural and Functional Biology Laboratory, National Institute of Immunology,

New Delhi-110067, India.

Running title: Deferoxamine for tuberculosis treatment.

*Corresponding author

Ranjan Kumar Nanda (PhD)

Translational Health Group, International Centre for Genetic Engineering and Technology,

New Delhi-110067, India

Telephone: 91-11-26741358; E mail: ranjan@icgeb.res.in

Abstract:

Dysregulated iron metabolism is reported in tuberculosis patients; therefore, it represents an

opportunity for developing host-directed therapeutics. This study monitored the

antimycobacterial properties of an iron chelator, i.e., Deferoxamine (DFO/D), and its impact

on the transcript and metabolite levels of *Mycobacterium tuberculosis* (Mtb) *in vitro*. For *in*

vivo validation, a group of mice received ferric carboxymaltose to create an iron overload

condition, and controls were aerosol-infected with Mtb H37Rv. Mtb-infected mice received

isoniazid (INH/H) and rifampicin (RIF/R) in combination with or without DFO before tissue-

specific CFU assay, liver metabolite screening and iron quantification using mass

spectrometry. DFO showed antimycobacterial properties comparable to INH *in vitro*. DFO

treatment deregulated (log2DFO/control>±1.0) Mtb transcript (n=137) levels, the majority of

which encode for iron-containing proteins and proteins involved in stress response. DFO

treatment up-regulated Rv3622c (PE32), Rv2353c (PPE39) and Rv3022A (PE29) genes and

conditional knocking down of ABC transporter like *irtA* by anhydrotetracycline (Atc)

inducible CRISPR interference (CRISPRi) approach compromised Mtb growth showing their

potential involvement in iron metabolism. Global Mtb metabolite analysis using GC-MS

identified a set of 5 deregulated metabolites indicating a perturbed pentose phosphate

pathway and inositol phosphate metabolism in the host upon DFO treatment. Iron-overloaded

mice exhibited significantly higher tissue mycobacterial burden at two weeks post-infection,

and the efficacy of INH and RIF were compromised, corroborating with previous reports.

Iron chelation by DFO or combined with/adjunct to RIF and INH significantly reduced the

lung/tissue mycobacterial burden at four weeks post-treatment, specifically in the first (~0.5

log) and second weeks (~0.5 log) of treatment. The intracellular pro-inflammatory cytokine

levels in the lung CD4+ T cells of INH and RIF-treated groups with or without DFO were

similar, suggesting DFO has a direct role in Mtb survival and metabolism rather than improved infection, and the efficacy of INH and RIF were compromised, corroborating with previous reports. DFO adjunct to RIF and INH treatment significantly altered liver arginine biosynthesis, which directly neutralizes ammonia and is immune-supportive. Conventionally, DFO is used for treating acute iron toxicity that is common in thalassemic patients, and this study demonstrates DFO has potential as adjunct therapeutics for tuberculosis.

Keywords: Tuberculosis, Iron chelation, Deferoxamine, Host-directed therapeutics, Metabolomics, Iron overloading, Transcriptomics.

Introduction: *Mycobacterium tuberculosis* (Mtb), the causative agent of Tuberculosis (TB), primarily infects the lower respiratory tract. After reaching the lungs, through the lymphatic and immune cell system, it disseminates to other body parts, causing Miliary and extra-pulmonary TB in different organs like the urinogenital system, brain, spinal cord, and liver.[1] The duration of treatment modalities for TB patients infected with drug-sensitive Mtb strains is 6–9 months and may prolong up to 24 months or more if drug-resistant Mtb strains cause the infection.[2] Shortening TB treatment duration by adopting additional host-directed therapeutics might help in increasing adherence and may also reduce the relapse rate, minimize Mtb drug resistance development, and may help in achieving the eradication of TB.[3]

Iron is an essential micronutrient for the host and Mtb's survival. In the host, iron is absorbed from the food materials through the intestine and is involved in erythropoiesis, respiration and other essential functions. Meanwhile, Mtb, with the help of siderophores like mycobactin and carboxymycobactin, chelates the host Fe^{3+} ion and intracellularly stores it in Fe^{2+} form with the help of iron storage proteins.[4,5] Iron supplementation in TB patients is reported to worsen clinical outcomes. In contrast, iron chelation in *in vitro* Mtb cultures prevents its growth and significantly reduces Mtb load in *in vivo* settings.[6,7,8,9,10] Infecting mice or macrophages with Mtb strain deficient in synthesis or uptake of siderophores leads to an inefficient establishment of active infection.[4] Mtb experiences iron deprivation in the endosome after being engulfed by interstitial macrophages. This compartment of interstitial macrophages is not permissive for Mtb growth. However, in alveolar macrophages, Mtb seldom experiences iron deprivation, and therefore, iron is readily available for Mtb inside its endosome.[11] Iron deprivation in Mtb compromises functions of enzymes that need iron as

cofactors as those involved in the TCA cycle and oxidative phosphorylation; therefore, its

deprivation is known to cause metabolic reprogramming in the host.[12]

The actinomycetes *Streptomyces pilosus* produces deferoxamine (DFO), that acts as a

siderophore, having strong affinity towards iron (Fe^{3+}), and forms a stable ferrioxamine

complex. DFO is hydrophilic and is used for the treatment of iron-toxicity which is

associated with various conditions including hemochromatosis. Excess iron *in vitro* increases

the viability and proliferation of Mtb, whereas chelation using DFO affects both these

activities, whereas iron chelation by silybin only impacts Mtb proliferation.[13] Chelating host

iron by DFO may also deprive Mtb from iron, which is essential for its survival and

establishment of the active TB disease.

In this study, we aimed to elucidate the anti-Mtb potential of DFO by monitoring its

influence on Mtb survival, levels of transcripts involved in iron metabolism, impact on tissue

metabolism and whether at early treatment time points it accelerates Mtb clearance adjunct to

the existing anti-TB drugs in C57BL/6 mice. Our findings indicate that iron deprivation

caused by DFO can augment earlier Mtb clearance from the host.

Results

Deferoxamine affects mycobacterial growth *in vitro*: Following the existing literature, the

DFO dose was fixed and used in all our experiments.[14,15] The growth curve of Mtb showed a

time-dependent increase in controls, whereas replicates treated with DFO at 0.1 mg/mL or

INH at 0.1 μg/mL alone or combined compromised Mtb growth (Figure 1A and B). The

growth kinetics of Mtb in the presence of DFO in 7H9 media were comparable to INH-

treated groups (Figure 1B). Furthermore, the combined treatment of DFO and INH also

compromised Mtb growth. On day five, post-inoculation, a significant reduction in Mtb

survival was observed in the presence of DFO, INH as well as their combination (Figure 1C).

However, as expected, DFO alone or in combination with INH significantly reduced

intracellular iron, zinc and copper levels in Mtb (Figure 1D, Supplementary Figure S1A and

B). From this we concluded that DFO may have a bactericidal effect, as Mtb relies on these

divalent ions for various processes like two-metal ion catalysis that are essential for its

survival and growth.

Because Gallium nitrate is reported to have antimycobacterial properties, we used it as an

additional control at 40 μg/mL concentration in subsequent experiments.[21] Electron

micrographs of Mtb treated with DFO and INH-RIF, harvested on day five post-treatment,

showed changes in the cell wall compared to the control and Gallium nitrate (GN) treated

Mtb (Figure 1E and F, Supplementary Figure S2). The kill kinetics of Mtb in the presence of

DFO were comparable to INH at day four and in combination (DFO+GN+INH) had

significantly lower CFU count at day eight post-incubation, whereas GN in combination with

DFO showed comparable CFU at both day 4 and 8 (Figure 1G). GN exhibited a less

pronounced anti-Mtb potential than DFO and INH, so it was excluded from further

experiments. These results revealed that DFO restricts iron availability, reduces CFU count,

has anti-Mtb potential, and causes morphological changes in Mtb.

DFO limits iron, leading to deregulated iron metabolism in Mtb: To understand the direct

impact of DFO on the Mtb transcriptome machinery, RNA-seq analysis of Mtb treated with

DFO and controls was performed. Quality control of RNA-seq data showed ~18.2±1.6

million high-quality reads based on a Phred-score $\geq$ 30 with negligible adapter

contamination. Upon alignment of the reads with the reference Mtb genome, we observed

alignment rates of 74.3±4.1 % and 97.1±0.8 % of DFO-treated and control Mtb cultures,

respectively. Out of a set of 3,931 identified Mtb genes, 3,826 showed a counts/million

(CPM) value $\geq$ 1 in all replicates of each condition and 137 genes were deregulated

(log$_2$DFO/control>$\pm$1.0, adjusted p-value $\leq$ 0.05) (Figure 2A, B, and C). The top 10 down

regulated genes included Rv1593c, Rv1596, Rv0263c, Rv1595, Rv0757, Rv1594, Rv0264c,

Rv0262c, Rv1130, and Rv1131. The top 10 up regulated genes included Rv0758, Rv2353c,

Rv0094c, Rv2812, Rv1587c, Rv2813, Rv1553, Rv1290A, Rv1917c, and Rv2370c (Figure

2C). Principal component analysis (PCA) of all the identified transcripts and unsupervised

hierarchical clustering of the deregulated gene sets showed two distinct clusters of the DFO-

treated and control samples (Figure 2D, Supplementary Figure S3A). In this study, Mtb

treated with DFO showed up-regulation of Rv1094, Rv1594 and the down-regulation of

Rv0327, Rv1553, Rv2276, Rv3251c, Rv3260c, and Rv3406 supporting earlier reports

(Figure 2E).[16,17]

Iron availability impacts the levels of transporters like Rv1348 (*irtA*), Rv1349 (*irtB*) and

mycobactin utilization protein Rv2895c (*viuB*). Since *irtAB* functions as an ABC transporter

complex involved in iron scavenging, it's essential role in iron transport was verified by

creating a conditional knockdown strain utilizing an anhydrotetracycline (Atc) inducible

CRISPR interference (CRISPRi) approach. A scrambled sgRNA (single guide RNA) was

used as a control to assess the gene-specific effect. The suppression of both the genes (*irtA*

and *irtB*) in the presence of Atc in the CRISPR knockdown strain of *IrtA* was observed

(Supplementary Figure S4A). Interestingly, the Atc-induced *irtA*-KD Mtb did not grow in

7H9 media, confirming its essentiality for its growth and multiplication (Supplementary

Figure S4B). DFO-treated Mtb had significantly higher Rv1348 levels with insignificant

changes in Rv1349 and Rv2895c levels (Figure 2F). DFO treatment for five days had a

negligible impact on Rv2027c (*dosT*), Rv3132c (*dosS*), and Rv3133 (*dosR*) genes, which are

known players in the induction of dormancy and persistor Mtb formation (Figure 2G).[18] DAVID gene ontology clustering revealed >1.2–3.0-fold enrichment of transcripts encoding proteins having ATPase activity, indicating DFO's impact on Mtb energy metabolism (Figure 2H). DFO also impacted the TCA cycle, oxidative phosphorylation, secondary metabolite synthesis, butanoate metabolism, fatty acid biosynthesis, and host-immune response (Figure 2H and I, Supplementary Figure S3B and C). DFO treatment led to upregulation of biological processes such as cellular response to starvation, chemical homeostasis, iron starvation, anaerobic respiration, and siderophore biosynthesis in Mtb (Figure 2I, shown in red in Supplementary Figure S3B). RNA modification, Heme metabolism, Nicotinamide, Cobalamin, Tetrapyrrole, Porphyrin, Arginine, Glutamine, Vitamins, Alpha-amino acid, Aromatic amino acid, branched-chain amino acid, aspartate biosynthesis/metabolism were downregulated upon DFO treatment (Figure 2I). Interestingly, downregulation of RNA modification, macromolecule methylation, fatty acid metabolism, dicarboxylic acid metabolic process and aromatic compound biosynthesis in DFO-treated Mtb were observed (shown in green in Supplementary Figure S3C). RNA-seq analysis of DFO-treated Mtb suggested that it has a direct impact on Mtb transcriptome and, as a result, may also alter the associated biological functions. A complete list of DEGs is included in Supplementary Table S1.

Earlier reports elegantly explained the role of various Mtb genes in iron metabolism and how iron uptake is a highly regulated process that can be disrupted for therapeutic intervention.[19] Out of the reported genes, we observed upregulation of Rv3622c (PE32), Rv2353c (PPE39) and Rv3022A (PE29), and the rest showed marginal changes which could be due to experimental differences (Supplementary Table S2).

DFO treatment alters Mtb metabolome: Further, the impact of DFO on Mtb metabolome was monitored using GC-MS analysis. PCA plot revealed that metabolites of DFO treated

and control H37Rv Mtb clustered separately (Figure 3A and B). DFO-treated H37Rv Mtb

showed significantly higher abundance (log2FC >± 1.0, -log10 p-value ≥ 1.3) of five

metabolites (Citric acid, Ribofuranose, Erythrotetrofuranose, Myoinositol and Propanoic

acid) compared to control (Figure 3C). Pathway impact analysis revealed that streptomycin

biosynthesis, inositol phosphate metabolism and pentose phosphate pathway were

significantly overrepresented in DFO-treated Mtb (Figure 3D). These findings indicate that

DFO impacts the Mtb metabolome and causes detrimental effects on Mtb through iron

deprivation. However, the target of DFO in Mtb is unknown, as with many antibiotics,

including pyrazinamide. Next, we aimed to monitor whether DFO can limit intracellular Mtb

growth and duplication.

DFO shows anti-mycobacterial activity in Mtb-infected J774A.1 cell culture: The

confocal images of the Mtb-infected J774A.1 cells that received DFO treatment or control

were analysed using NIS-Element software. The DFO-pretreated J774A.1 cells showed

decreased intracellular mycobacterial burden in a dose-dependent manner (Supplementary

Figure S4C). Interestingly, we also observed hypertrophy, marked by increased cell size in

DFO-treated cells compared to control.

Iron supplementation leads to liver iron loading, higher tissue mycobacterial load and

compromises the efficacy of INH and RIF: To validate the hypothesis that iron

supplementation is not beneficial in TB patients, we created an iron overload mice model and

infected it with a low aerosol dose of Mtb**.** The method adopted for iron overloading in

C57BL/6 mice is presented in Supplementary Figure S5A. Prussian blue staining of the liver

samples of iron-overloaded groups also showed higher iron accumulation and distribution

(Supplementary Figure S5B, S7A and B). Iron overloading had a negligible effect on the

mice's body weight during and after the course (Supplementary Figure S5C). However, the

iron-overloaded mice group showed significantly higher liver iron levels than the control

group, and the INH-RIF treatment had a marginal reduction (Supplementary Figure S5D,

S6A and B). Prussian blue staining and ICP-MS analysis of the liver of iron-overloaded mice

showed increased iron levels till 75 days post-treatment (Supplementary Figure S6C, D and

S7D, E). The tissue (lungs, spleen, and liver) mycobacterial burden was significantly higher

in the iron-overloaded mice group at 15 days post-infection compared to the controls

(Supplementary Figure S6E, F and G). The whole blood and thigh muscle iron levels of iron-

overloaded mice did not show significant variations (Data not shown). As expected, the INH-

RIF treatment cleared the tissue mycobacterial burden in control groups; however, the

mycobacterial clearance in the lungs and spleen of iron overloaded mice groups was delayed

and took longer time (>30 days of treatment) to bring it to lower than the limit of detection.

Interestingly, the liver mycobacterial burden of the iron-overloaded group was similar to the

control groups at 45- and 90-days post-infection. This demonstrated that higher host iron

levels compromise the anti-mycobacterial efficacy of first-line anti-TB drugs.

Iron chelation by DFO significantly reduces the tissue mycobacterial burden in

C57BL/6 mice: Mtb infected C57BL/6 mice treated with DFO alone showed significant

reduction in the liver iron levels as seen by Prussian blue staining and ICP-MS (Figure 4A, B,

C and Supplementary Figure S7C, G). The decrease in iron levels was followed by a

significant reduction in lung Mtb load within 30 days post-treatment and continued up to 75

days post-treatment compared to the infected control (90 dpi, Figure 4C and D). No change in

the spleen mycobacterial burden was observed (Figure 4E). However, the liver of DFO-

treated mice showed a significantly lower mycobacterial load (Figure 4F). This suggested

that DFO alone has a direct effect that indirectly affects Mtb clearance in the lung tissues, so

we studied the impact of DFO in combination with the anti-TB drugs in vivo in Mtb-infected

mice.

DFO enhances lung bacterial clearance, increases recruitment of CD4$^+$ T cells and

macrophages impacting liver metabolite profile with the existing anti-TB drugs:

Significantly lower liver iron levels were observed in RIF-INH treatment with or without

DFO (Figure 5A, B, C and Supplementary Figure S7F, H). DFO adjunct to anti-tuberculosis

drugs (InRHD) showed no difference in the lung bacterial burden at 30 days post-treatment

(Figure 5D). Interestingly, Mtb-infected C57BL/6 mice treated with DFO adjunct to INH-RIF

(grouped as InRHD) showed a significant reduction in lung mycobacterial burden at early

time points, i.e., one- and two-weeks post-treatment, compared to the group receiving only

INH-RIF (grouped as InRH) treatment (Figure 6A, B, and F). We monitored the immune cell

distribution in an independent experiment and observed that InRHD group had a significantly

higher frequency of CD4$^+$ T cells and F4/80+ macrophages at one-week post-treatment

compared to the InRH group and In and InRH groups, respectively (Figure 6B and C). The

InRHD group had a significantly lower frequency of CD4$^+$ T cells and F4/80+ macrophages

at two-week post-treatment compared to the Mtb infected (In) group and group receiving TB

drugs (InRH), respectively (Figure 6A and G).

Liver global metabolome analysis of InRHD vs InRH group completing the first week of

treatment, using GC-MS, showed 20 deregulated (log2 Fold change $>\pm1.0$; p-value< 0.05)

metabolites (Figure 6D). DFO treatment impacted the liver alanine, glutamic acid, lactic acid,

and malonic acid levels (Figure 6D). Pathway enrichment analysis showed an

overrepresentation of alanine, aspartate and glutamate, pyruvate metabolism and arginine

biosynthesis in the treated group (Figure 6E). After one-week of INH-RIF treatment with or

without DFO (both InRH and InRHD), liver pantenoic acid, lactic acid and alloxanoic acid

levels were deregulated compared to the Mtb-infected control (In) (Supplementary Figure S8A and B). Similarly, seven deregulated metabolites were identified in the liver of the DFO-treated group, completing two weeks of treatment (Figure 6H). The liver of the InRHD group had significantly higher levels of proteogenic amino acids (valine and isoleucine) and non-proteinogenic amino acids (ornithine and aminoadipic acid) compared to the InRH group (Figure 6H). Arginine biosynthesis was overrepresented in the livers of mice receiving DFO as an adjunct to RIF and INH at 1 and 2 w.p.t. (Figure 6E and I). Long-chain fatty acids like oleic acid, arachidonic acid and octadecanoic acid were deregulated in the liver of the drug-treated mice (InRH, InRHD) as compared to the infected control (In) in the second week of treatment (Supplementary Figure S8C and D). The drug metabolism pathway for INH metabolism was enriched in the liver of mice groups receiving treatment (Supplementary Figure S9A, B and D). The arachidonic acid pathway was over-represented in the first week of treated groups (Supplementary Figure S9B). In contrast, biosynthesis of unsaturated fatty acids (linoleic and alpha-linoleic acid) was enriched in treated groups compared to the Mtb infected controls at first- and second-week post-treatment (Supplementary Figure S9C and D). These findings suggested that DFO accelerated Mtb clearance in C57BL/6 mice and impacted the liver metabolite levels significantly differently in adjunct to the INH and RIF than when treated with anti-TB drugs alone.

DFO treatment adjunct to INH and RIF minimally effects cytokine production in the lung CD4+ T-cells: Increased CD4+ T cells and their proinflammatory cytokine releasing levels significantly improve lung Mtb clearance. Therefore, the effect of DFO on the frequency of CD4+ T cells and their pro-inflammatory cytokine production capacity after stimulation with PMA and ionomycin was monitored. The frequency of lung $CD3^+CD4^+$ cells in Mtb-infected C57BL/6 mice receiving INH-RIF with or without DFO at two weeks

post-treatment was similar (Supplementary Figure S10A and B). The intracellular pro-

inflammatory cytokines (IFN-γ, TNF-α and IL-17A) levels in the lung $CD4^+$ T-cells of INH

and RIF-treated groups with or without DFO were found to be similar (Supplementary Figure

S10C). These findings suggested that DFO does not directly impact the CD4+ T cell

frequency and pro-inflammatory cytokine production efficiency of $CD4^+$ T cells, indicating

that DFO does not hamper the effect of the immune system against active Mtb infection.

Discussion: Numerous biological processes, including cell wall synthesis, ATP synthesis,

DNA coiling, transcription, and translation in Mtb, have been instrumental in developing

anti-TB drugs.[19] However, limited studies have attempted to understand how limiting iron in

the host affects Mtb growth and influences the effectiveness of anti-TB agents. Additionally,

the effect of iron limitation on the survivability of Mtb has received less attention.[18,19] In this

study, we attempted to generate evidence of the pivotal role played by the host iron in Mtb

survival and how manipulating iron availability through chemical agents, such as iron

chelators, may impact Mtb clearance both *in vitro* and *in vivo* (Figure 7).

Mtb cultured in 7H9 media with DFO exhibited notable bactericidal activity akin to the

effectiveness of INH (Figure 1). Gallium nitrate is a ferric mimetic known to disrupt iron

metabolism in Mycobacteria, and it decreases Mtb proliferation in infected

macrophages.[18,19,20] We observed that GN was less effective in inhibiting Mtb growth than

DFO and INH *in vitro*. The observation of our *in vitro* study aligns with previous reports

corroborating the anti-mycobacterial role of DFO.[9] Iron deprivation targets mycobacterial

membrane integrity by enhancing membrane permeability and disrupting membrane

homeostasis reported in *M. smegmatis*.[25] Electron micrograph images of H37Rv Mtb treated

with DFO and INH showed cell wall damage, and a decreased envelope thickness compared

to the controls (Figure 1F and Supplementary Figure S2). DFO also impacts Mtb's

intracellular copper and zinc levels (Supplementary Figures S1A and B). Transition metals

like copper and zinc in low concentrations are essential for Mtb viability but, at high

concentrations, can be toxic. Many enzymes in Mtb require these metal ions for their activity,

like cytochrome C oxidase and superoxide dismutase, which require copper, copper, and

zinc, respectively. So, the effect of DFO seems to be mainly due to iron deprivation and

partly because of copper and zinc chelation as well.

Recent studies on iron deprivation in Mtb showed the up regulation of specific iron-

containing proteins that aid in iron storage, genes involved in the uptake of iron and down

regulation of the TCA cycle, and anabolic pathways for amino acid, vitamins, heme, and

secondary metabolite biosynthesis.[16,17] We also observed perturbed transcript levels of genes

encoding proteins containing iron-sulfur clusters or iron alone. It seems that upon iron

chelation by DFO, Mtb tries to scavenge the residual iron from surroundings by up regulating

the genes involved in iron transport (Rv1348), iron-containing proteins (Rv1094, Rv1594)

and storage (Rv0327c, Rv1553, Rv2276, Rv3251c, Rv3260c and Rv3406). RNA-seq analysis

revealed that the gene signatures indicative of starvation and abiotic stress are affected, and

DFO has a direct and detrimental effect on Mtb. Mtb is widely known to form a persister

population when it encounters biotic and abiotic stress within the host or *in vitro*. However,

the dos regulon encoded by Rv2027c(*dosS*), Rv3132c(*dosT*), and Rv3133c(*dosR*) did not

show changes on day 5 of DFO treatment.18 Rv1348(irtA) seems critical for Mtb survival, as

the CRISPRi knockdown assay revealed. The results suggested that DFO containing regimen

may not lead to the generation of viable but non culturable bacteria (data not shown).

Mtb H37Rv global metabolite analysis revealed that DFO-induced metabolome changes

persist even in the absence of DFO as myoinositol levels were significantly upregulated in

DFO-pretreated Mtb as compared to Mtb cultured in rich media (7H9 supplemented with

10% OADC) (Figure 3C). Mycobacteria use myoinositol to synthesize phosphatidylinositol

and mycothiol, an essential phospholipid required for cell wall synthesis and maintaining

redox homeostasis in Mtb, respectively.[26] Global metabolite analysis also indicated that DFO

treatment affects the pentose phosphate pathway in Mtb. These results aligned with our

transcriptomics analysis, which indicated that DFO treatment perturbs metabolic pathways in

Mtb directly.

Several reports have shown that iron supplementation in mice, whether administered orally or

peritoneally, supports the growth of non-tuberculous mycobacteria (NTM); however, the

studies on Mtb are limited.[27] Iron supplementation in mice significantly increased the Mtb

burden in spleen and lung tissues aligning with our findings.[28] Our study delved into the

impact of host iron status on Mtb survival and growth *in vivo*, revealing that iron overloading

achieved via intravenous administration of ferric carboxymaltose promoted higher

mycobacterial load in the lung, liver, and spleen several weeks post supplementation

(Supplementary Figure S8). However, this effect waned within a few months after

discontinuing the iron overloading.

Limiting iron through DFO administration to Mtb-infected mice significantly reduced liver

iron levels and mycobacterial burden compared to the infected control group (Figure 4C and

4F). Furthermore, liver iron levels in the iron-overloaded C57BL/6 mice significantly

declined upon DFO treatment. We also assessed the effect of DFO when used as an adjunct

to anti-TB drugs like INH and RIF. We observed substantially lower lung bacterial burden

between animals treated with INH and RIF with and without DFO. It is critical to highlight

that DFO administration, in conjunction with the existing TB drugs, reduced tissue Mtb load

significantly at early time points. This suggests that DFO, as an adjunct to existing TB drugs,

holds promise for potentially shortening the duration of anti-TB treatment by various

mechanisms (Figure 7).

As the liver plays a critical role in regulating iron homeostasis by detecting even minor

fluctuations in the systemic iron requirements and accumulating excess iron.[29] Various

regulatory mechanisms within the liver govern the expression of the iron regulatory genes,

storage capacity, and iron mobilization. Dysregulation of these functions can lead to iron

imbalance, a primary contributor to iron-related disorders. Iron supplementation in C57BL/6

mice had no discernible effect on their weight, corroborating earlier reports in Sprague-

Dawley rat pups.[30] To assess the effect of iron chelation on the liver, we observed lower liver

mycobacterial burden in the DFO-treated groups. We also monitored the liver metabolome of

Mtb-infected mice receiving INH and RIF treatment with or without DFO (Figure 6D and H).

It is well known that anti-TB drugs cause drug-induced liver injury (DILI) in model

organisms and humans. Global liver metabolite analysis showed an enriched arginine

biosynthesis pathway in the DFO-treated group, which could be contributing to liver

detoxification.[31] Arginine, an essential antioxidant for the liver, is reported to neutralize

ammonia, detoxifying the liver. Arginine supplementation helps in liver regeneration, and the

production of antimicrobial compounds like nitric oxide (NO) in macrophages and as an

adjunct to DOTS treatment, is beneficial to TB patients.[31,32] Arginine, alanine, and valine

abundance were higher in the liver of DFO-treated mice with potential immune-supportive

functions.[33] Alanine is converted to pyruvate by the alanine aminotransferase (ALT) enzyme

in the liver, which contributes to glucose production by gluconeogenesis and enhances

immune cell function against pathogens.[34] DFO treatment enriched linoleic acid, alpha-

linolenic acid metabolism, and arachidonic acid metabolism compared to Mtb-infected

controls at the first and second week of treatment. Linoleic acid is also a precursor for

arachidonic acid, which gives rise to an essential class of inflammatory mediators called

eicosanoids to elicit an immune response against pathogens. DFO has been reported to

increase glycolytic metabolism in Mtb-infected human monocyte-derived macrophages

(MDM) while concurrently enhancing the early innate immune response by increasing the

expression of IL-1β and TNF-α both at transcript and protein levels.[35] Interestingly, we did

not observe any significant changes in the intracellular cytokine levels (IFN-γ, IL-17A and

TNF-α) in CD4$^+$ T-cells harvested from the lungs of Mtb-infected C57BL/6 mice receiving

DFO in combination with the first-line anti-TB drugs which may be a cell type-specific

activity suggesting the direct role of DFO on macrophage cytokine production with minimal

effect on cytokine production the CD4$^+$ T-lymphocytes (Supplementary Figure S10C).[36] But

we did observe increased recruitment of CD4$^+$ T cells in the lungs of the InRHD group at the

first week of treatment but observed reduced frequency of these cells at the second week of

treatment. It is likely that the lungs of mice receiving DFO adjunct to INH and RIF had lower

CD4$^+$ T cell and macrophage frequency because of effective bacterial clearance observed two

weeks post-treatment. Importantly, DFO was observed to reversibly inhibit DNA synthesis in

human B and T lymphocytes with limited impact on RNA and protein synthesis.[36] In the

presence of equimolar FeCl3, the effects of DFO in these cells were also reversed.[36] We

hypothesized that the combined effects of DFO with first-line anti-TB drugs could yield

enhanced bactericidal effects and potentially reduce TB treatment duration. A recent report

demonstrated that DFO treatment in BCG-infected primary hMDMs increased bedaquiline

bactericidal activity, indicating its additive effects with the second-line anti-TB drugs.[9]

Further research is required to assess the efficacy of DFO in combination with these drugs

against drug-resistant Mtb isolates. DFO can be introduced in the host by the subcutaneous,

intramuscular, intravenous or oral route. Although iron chelation by DFO is the gold

standard, it can also cause side effects like hypotension, fever, dysuria, leg cramps, rashes,

pruritus, and free radical-mediated alveolar damage, leading to pulmonary toxicity, which

intermittent infusions could avoid.

Targeting other critical trace elements like copper, zinc, and magnesium, which are essential

for Mtb survival and growth, could also be a promising avenue for adjunct TB treatment

strategies.[37,38,39] Other iron chelators like Desferri-Exochelin may have a similar impact and

need additional investigation.[40] Additionally, exploring the role of gene polymorphisms in

NRAMP1, the HFE gene associated with hemochromatosis, HP and α2 macroglobulin genes,

which contribute to TB susceptibility, may facilitate the identification of the populations that

could benefit the most from iron chelation therapy.[10,41] Currently, there is no effective TB

vaccine available for adults, and drug-resistant TB cases are on rise, it is crucial now than

ever to invest resources in repurposing of FDA-approved or novel drugs for TB treatment.

Several FDA-approved drugs have emerged as potential host-directed therapeutic candidates

for combination therapy with the existing TB drugs. DFO, an FDA approved drug with pro-

inflammatory properties for treating conditions like hemochromatosis in humans, which has

shown the potential to bolster host immunity in the present study, leading to improved Mtb

clearance and, therefore, has the potential to serve as an agent for host-directed adjunct

therapy.[42] Our findings on the earlier clearance of Mtb in DFO in combination to RIF and

INH offers a new insights into manipulating host iron metabolism to shorten the treatment

duration (Figure 7). Host-directed therapies involving iron chelation merit further validation

to determine their impact on limiting Mtb drug resistance development, if any. Particular

attention should be given to the potential toxicity of DFO treatment, and introducing an

equivalent amount of $FeCl_3$ in cellular systems negates its function.[30] It is worth exploring

whether DFO, as an adjunct, could also reduce the dosage of existing TB drugs, which

presents an exciting avenue for future research. New therapeutic interventions targeting host

or pathogen iron metabolism, such as iron chelation, are poised to contribute significantly to

the fight against TB.

Experimental procedures

Ethical statement: The experiments performed in this study were approved by the Institute

Biosafety   Committee   and   the   Institute   Animal   Ethics   Committee

(ICGEB/IAEC/07032020/TH-11) of the International Center for Genetic Engineering and

Biotechnology, New Delhi.

Mycobacterial H37Rv culture: *Mycobacterium tuberculosis* H37Rv strain (OD = 0.02)

were inoculated in 7H9 media (BD Difco Middlebrook) supplemented with glycerol (0.2%),

tween 80 (0.05%) and OADC (Oleate, albumin dextrose and catalase, 10%) in the absence or

presence of individual drugs or in combination in the tuberculosis aerosol challenge facility

(TACF, BSL-III) at ICGEB New Delhi. Mtb were grown at $37^{o}C$ in a shaker incubator set at

180 rpm. Individual drug solutions were prepared at their minimum inhibitory concentrations

(MICs) for DFO (0.1 mg/ml) and isoniazid (INH, 0.1 µg/ml) using milliQ water. Mtb growth

was monitored by measuring absorbance at 600 nm at an interval of 24 hours up to 5 days

post inoculation using a spectrophotometer to monitor the drugs' bactericidal activity. On the

fifth day, the harvested control and drug treated Mtb were inoculated on 7H11 agar plates

supplemented with OADC (10%) and BD MGIT™ - PANTA™ antibiotic Mixture for colony

forming unit (CFU) assay. On the fifth day, a fraction of the Mtb culture (0.8 OD, 133 M)

were centrifuged at 3,500 g for 10 minutes at room temperature and elemental analysis was

performed in the pellet and the cultured media as a separate fraction by using inductively

coupled plasma mass spectrometry (ICP-MS) analysis (Figure 1A).

Transmission electron microscopy of Mtb: Mtb cultures (0.1 OD) were inoculated with

DFO, INH, and Gallium nitrate (GN, 4µg/ml) at MIC and control, incubated at 180 rpm at

37^{o}C. Mtb cultures were harvested 5 days post-treatment and centrifuged at 3,500 g for 5

minutes, followed by washing with phosphate buffer saline (PBS) by centrifuging at 3,500 g

for 5 minutes. The pellet containing Mtb was fixed with 2% glutaraldehyde for an hour. The

fixed bacterial pellet was washed twice with PBS and then passed five times through a 26 G

needle to make a single-cell suspension resuspended in 100 µl ddH$_2$O. 10 µl suspension was

loaded onto a carbon grid and stained with 5% uranyl acetate. After air drying the grid, it was

loaded in the Transmission Electron Microscope (Tecnai G2 spirit, HT120kV) and the

micrographs were captured (Figure 1E).

Kill kinetics of Mtb in the presence of DFO, GN, INH and in combination: Log phase

(OD= 0.2) Mtb H37Rv culture were inoculated in 7H9 media (2 ml) supplemented with

OADC (10 %), glycerol (0.2 %) and tween 80 (0.05 %) in presence of $10 \times$ MIC of DFO (1

mg/ml), GN (40 µg/ml), INH (1 µg/ml) and in combination (DIG: DFO +INH + GN) and

were incubated at 37^{o}C at 180 rpm. Aliquots (100 µl) of each culture condition were

harvested at days 0, 4 and 8 for CFU assay by plating diluted samples on 7H11 agar plates

supplemented with OADC (10 %) and BD MGIT™ - PANTA™ and incubated at 37^{o}C.

Mycobacterial colonies were enumerated after 3 weeks post-plating.

Transcriptome analysis of DFO treated Mtb: Mtb H37Rv cultures (0.1 OD/ml) treated

with DFO (0.1 mg/ml) and control were harvested on day 5 by centrifuging at 3,500 g for 10

minutes at room temperature. The pellet after washing with Tris-EDTA (1 M) to remove

extra media and resuspended in Trizol reagent (1 ml, Invitrogen Inc., USA). The samples

were subjected to bead beating (4 cycles, 45 sec each) after adding zirconia beads (~200 µl,

0.1 mm) at 6.5 m/s speed with incubation of 30 sec on ice after each cycle using Biospec

Mini-Beadbeater-16. Chloroform (0.2 ml) was added to the samples and after briefly shaking

the tubes, the samples were incubated on ice (for 5 min) and centrifuged at 13,400 g, 4℃ for

20 min. The upper aqueous layer was transferred to a fresh tube and RNA was precipitated by

adding isopropyl alcohol (0.5 ml). After incubating the samples at -80ºC for 20 min, were

centrifuged at 13,400 g for 10 min at 4ºC. The pellet was washed with 75% ethanol and

resuspended in nuclease-free water. RNA was quantified using a Qubit BR RNA estimation

kit (Invitrogen Inc., USA) as per the manufacturer's protocol and RNA integrity was assessed

using Agilent TapeStation (Agilent Inc., USA). Samples (0.5 µg each) above a RIN value of

7 were sent for RNA sequencing to NGB Diagnostics Pvt. Ltd. Briefly, library preparation

was performed using NEBNext Ultra RNA library prep kit (NEB, USA). The uniquely

indexed libraries were sequenced on the Illumina NovaSeq 6000 platform with 150×2 paired-

end chemistry (Figure 2A).

The Next Generation Sequencing (NGS) data in fastq format was analyzed using FastQC

(v0.12.1) to determine the quality and adapter contamination. Adapter contamination was

addressed using bbduk.sh (from BBmap toolkit v39.06) with options (ktrim=l k=23 mink=11

hdist=1 tpe tbo). Hisat2 (v2.2.1) was used for the alignment of reads to the Mtb genome

(Refseq Accession: NC_000962.3) which was subsequently sorted using Samtools (v1.13).

The sorted alignment files in BAM format were used as input for counting reads using

featureCounts (v2.0.3). The read count matrix was formatted so that it can be used as input

for iGEAK! and iDEP.96. EdgeR was used to identify the differentially expressed genes with

a minimum CPM value of 1, a $\log_2$DFO/control>1.0 and adjusted p-value $\leq$ 0.05 calculated

using Benjamin-Hochberg method. To identify the deregulated pathways, the same raw read

matrix was analyzed using the pre-ranked FGSEA option in iDEP.96.

CRISPRi knockdown strain preparartion: The pLJR965 CRISPRi plasmid (kind gift by

Dr. Sarah M. Fortune) backbone was digested with BsmBI. Subsequently, sgRNAs designed

to target the *irtA* ORF was cloned using previously established methods.[40] Confirmation of

successful cloning was obtained through partial sequencing. The resultant plasmid containing

the *irtA*-targeting sgRNA was then introduced into Mtb via electroporation. Bacterial cultures

were allowed to recover in 7H9 medium for 24 hours, after which were plated onto 7H11

agar supplemented with kanamycin (50 µg/ml) for the selection of transformants. The

inducible system was triggered by anhydrotetracycline (Atc, 50 ng/ml) supplementation on

every third day.

Global metabolite analysis of DFO-treated Mtb H37Rv: Mtb H37Rv cultures (normalized

with bacterial number) treated with DFO (0.1 mg/ml) and control were harvested on day 5 by

centrifuging at 3,500 g for 10 minutes at room temperature (Figure 3A). Mtb pellet harvested

from DFO-treated and control condition was incubated in drug-free 7H9 media with OADC

(10%) for 6 days and then equal number of bacteria were harvested for metabolomics

analysis. The pellet was resuspended in 80% methanol along with ribitol (2 µl, 0.5 mg/ml) as

a spike in standard and transferred to bead beating tubes containing 0.1 mm zirconia beads.

The samples were subjected to bead beating using Biospec Mini-Beadbeater-16 at 3650

oscillations/minute, 30 seconds on/off cycles six times with incubation on ice during the off

cycle. After bead beating, the samples were incubated on ice for 30 minutes followed by

centrifugation at 10,000 g for 10 minutes at 4°C to collect the supernatant (800 µl). The

extracted metabolites were filtered using a 0.2 µm, nylon membrane filter (#726-2520) and

stored at -20°C until further processing. Equal volume of the extracted metabolites from all

the samples were pooled to prepare two QC samples.

For derivatization, samples were dried at 40°C using vacuum concentrator (Labconco

Centrivap, USA) and incubated with Methoxamine hydrochloride (MeOX-HCl, 2%, 40 µl) at

60°C for 2 hours at 900 rpm in a thermomixer (Eppendorf, USA). After adding N-methyl-N-

(trimethylsilyl) trifluoroacetamide (MSTFA, 70 µl), it was incubated at 60°C for 30 minutes

at 900 rpm in a thermomixer. After incubation, the samples were centrifuged at 10,000 g for

10 minutes at 25°C, the supernatant was transferred to GC vial inserts for GC-MS analysis

using a 7890 Gas Chromatography (Agilent Technologies, Santa Clara, CA) coupled to a

Pegasus 4D GC × GC time-of-flight mass spectrometer (LECO Corporation). The samples

were randomized using Research Randomizer (www.randomizer.org) and processed together

with one QC sample ran before and after the sequence. The derivatized samples (1 μL) were

injected to an HP-5ms GC column (30 meters length, 250 μm internal diameter) in splitless

mode using Helium as a carrier gas at a constant flow rate (1 ml/min) and secondary column

was Rxi-17 of 1.5 m length and 250 μm diameter. Electron ionization (EI) mode was fixed at

−70 eV to scan ions of 33 to 600 m/z range at acquisition rate of 20 spectra/second. The ion

source temperature was set at 220°C. GC method parameters used for acquisition were: 50°C

for 1 min, ramp of 8.5°C/min to 200°C, ramp of 6°C/ min to 280°C, hold for 5 minutes.

Secondary oven temperature offset was set at +5°C relative to GC oven temperature.

Similarly, the modulator temperature offset was set at +15°C relative to the secondary oven

temperature. Transfer line temperature was set at 225°C. A solvent delay of 600 seconds was

used, and all data acquisition was completed within 24 hours of derivatization. A commercial

standard mix of amino acids were derivatized and run using the same GC-MS method to

confirm the identity of the important metabolic features. All GC-MS raw data files of the

study groups were aligned using the "Statistical Compare" feature of ChromaTOF (4.50.8.0,

Leco, CA, USA). Peaks with a width of 1.3 seconds and the signal to noise ratio (S/N)

threshold of ≥50 were selected. Putative identity of the peaks was assigned using a NIST

(National Institute of Standards and Technology, USA) library (version 11.0; 243,893

spectra) search qualifying a similarity match of >600 at a mass threshold of 100. The .csv

file, generated from the above analysis, was manually curated to check the peak identity.

Analytes present in more than 50% of the samples of a study group were selected for

statistical analysis. Ribitol peak area from the sample was used for normalizing the liver

metabolite data. Fold change analysis and p-value were calculated using MetaboAnalyst

software (version 6.0). Metabolic features that showed $\log_2$ fold change $\geq \pm 1.0$ with p-value <

0.05 between the study groups were selected as deregulated molecules. Pathway enrichment

analysis for deregulated molecules was performed using Pathway analysis plugin in the

MetaboAnalyst software.

J774A.1 infection with H37Rv and DFO treatment: The J774A.1 cell was seeded on

sterile round cover glass in a 6-well plate. The cells were pretreated with different DFO

concentrations (0, 0.25, 2.5 and 25 mM) for 24 hours before introducing Mtb H37Rv at 1:10

multiple of infection (MOI). After washing, the cells were stained with DAPI (1 μg/ml,

Invitrogen #D1306), MitoTracker™ Red CMXRos (200 nM, Invitrogen #M7512) and fixed

at different time points and finally slides were prepared for confocal microscopy by using the

prolong gold antifade mountant (Invitrogen, Cat# P36934). These fixed cells were used for

fluorescence microscopy using a laser scanning confocal microscope (Nikon A1R, USA), and

images were analyzed using NIS-Element software.

Iron supplementation in C57BL/6 mice and Mtb aerosol infection: C57BL/6 mice (8-10

weeks old, male) from the host institute (ICGEB) animal house were transferred to the

tuberculosis aerosol challenge BSL-III facility (TACF) for adaptation for at least one week

prior to the infection experiment. The animals were kept in a specific pathogen-free

environment, 12-hours daylight conditions, and food and water were provided *ad-libitum*. A

set of the study mice received ferric-carboxymaltose (Ferinject, LUPIN ltd, India, 0.5 mg

iron/mice in 100 μl saline) through intravenous injection, while the control mice received

saline (100 μl). These iron-supplemented mice received a total of four doses of ferric-

carboxymaltose injections on every fourth day. After a week of rest, the iron-supplemented

and control mice were aerosol infected with a low dose (100-120 CFU) of Mtb H37Rv strain

using Madison chamber present in TACF and sacrificed at 15-, 45- and 90-days post-

infection (dpi) to monitor the tissue bacterial load. A subgroup of Mtb infected C57BL/6

mice received either INH-RIF (InIR, 0.1 g/l each, changed alternate day) in drinking water

ad-libitum or DFO (InD, 150 mg/kg body weight of mice, intraperitoneal injection) or both

(InIRD) starting 15 dpi for 75 days (i.e., 90 dpi). DFO dose was introduced intraperitoneally

every alternate day till treatment completion. At the end of the study, mice were

anaesthetized using 5% Isoflurane, and organs were harvested from the humanely euthanized

mice. Harvested tissues (lungs, spleen, and liver) were processed using a homogenizer in

sterile PBS (1 ml) and an aliquot (100 µl) of it was inoculated on 7H11 agar plates

supplemented with OADC (10%) and BD MGIT™ - PANTA™ antibiotic mixture at

appropriate dilution in sterile PBS and the colonies were counted post 3 weeks of incubation

at $37^{o}C$ in the incubator. To a part of the liver tissue, chilled methanol was added and stored

at $-80^{o}C$ for elemental analysis using ICP-MS and metabolite profiling using gas

chromatography-mass spectrometry (GC-MS).

Tissue histology analysis: A part of the harvested tissues was stored in 10% formaldehyde

for histology analysis. Briefly, processed tissues were paraffin-embedded and 5 µm thick

sections were used for slide preparation at an accredited pathology lab. Prussian blue staining

was used for monitoring the tissue iron distribution. Histology slides were scanned in Zeiss

microscope and analyzed using Zen 3.1 software (Supplementary Figure S7).

Elemental analysis using ICP-MS: Equal number of bacteria (0.8 OD) were pelleted and to

the pellet, HNO_3 (70%, 150 µl) was added and transferred to sample vials (MG5, Anton Paar,

USA) and then H_2O_2 (30%, 50 µl) was added. Similarly, to 1 ml culture filtrates, HNO_3

(70%, 500 µl) and H_2O_2 (30%, 150 µl) were added. The sealed sample vials were subjected

to microwave (Anton Paar, USA) digestion using a ramp of 250W for 15 min with a hold

time of 5 min at 250W. Similarly, liver , thigh muscle (50 mg each) tissues or equal volumes

of whole blood (50 µl) were taken in a digestion vial (64MG5 vials) and after adding HNO_3

(400 μl) and H_2O_2 (100 μl) were incubated for 20 min at room temperature, and then the vials

were sealed and subjected to microwave digestion. The initial power was set at a ramp of 10

min and 150W with a max temperature of 140°C, followed by a 15 min hold. Then, a second

ramp of 15 min from 150-250W was used, and a final hold was used to bring the temperature

to 55°C at the highest fan speed. After diluting the samples with trace metal-free water, iron

(57Fe) levels were monitored in the digested samples using ICP-MS (iCAPTM TQ ICP-MS,

Thermo Scientific, USA). Thermo Scientific Qtegra[TM] Intelligent Scientific Data Solution

(ISDS) software was used to operate and control the instrument. Briefly, the torch was

warmed up for 30 min in single-quad Kinetic energy discrimination (SQ-KED) mode with

helium to remove the polyatomic nuclei interference and then autotuned in normal mode

followed by advanced KED mode. Then, the multi-element standard (#92091, Sigma Aldrich,

USA) of different concentrations was run at the same settings to prepare the standard plot. A

sample blank was run before each sample, and a quality control (QC) of known

concentrations of standards was run after every 15–20 samples to monitor and maintain

uniform signal intensity throughout the run. Data analysis was done using Qtegra software

(Thermo Scientific, USA).

Flow sorting of CD4$^+$ T cells and macrophages from the lungs: We performed an

independent experiment to monitor the immune cell distribution in the lungs of Mtb infected

C57BL/6 mice. The right lobes of the lungs were harvested and incubated with collagenase (2

mg/ml) and DNase (1 mg/ml) in RPMI-1640 (2 ml) at 37°C for 30 min. The enzymatically

treated tissues were passed through a strainer (70 μm) before centrifuging at 400 g for 10 min

to collect the pellet. Ammonium chloride potassium (ACK) lysis buffer was added to the

pellet, incubated for 2 min at room temperature and quenched by adding RPMI-1640. The

cells were centrifuged and resuspended in RPMI-1640 and counted using a hemocytometer.

10 million cells from each sample were incubated with an antibody cocktail of LIVE/DEAD,

CD45, CD3, CD4 and F4/80 surface markers (Live dead fixable stain #L34975, CD4-SB600 #63-0042-82 from Invitrogen and CD3-Alexa Fluor 700 #152316, CD45-PerCP/Cy5.5 #103132 and F4/80-PE #123110 from BioLegend) for 40 mins on ice. After staining, these cells were fixed using paraformaldehyde (4%) washed with FACS buffer and finally resuspended in the FACS buffer. Immune cell sorting was done using BD FACS Aria III Fusion (BD Biosciences, USA) present in TACF, and the data was analyzed using FCSExpress (DeNovo software, Version 6.0).

Liver metabolite profiling using GC-TOF MS: For metabolites isolation, liver tissues (100 mg) were finely chopped and transferred to bead-beating tubes (2 ml) with zirconium beads (2 mm, 250 mg). Chilled methanol (80%, 1 ml) was added to the tubes along with ribitol (5 μl, 2 mg/ml) as a spike in standard. The tissues were subjected to bead beating using Biospec Mini-Beadbeater-16 at 3650 oscillations/minute, 30 seconds on/off cycles six times with incubation on ice during the off cycle. After bead beating, the samples were incubated on ice for 30 minutes followed by centrifugation at 10,000 g for 10 minutes at 4°C to collect the supernatant (800 μl). The extracted metabolites were filtered (0.2 μm, nylon membrane filter #726-2520) and stored at -20°C until further processing. Equal volume (120 μl) of the extracted metabolites from all the samples from a particular time point were pooled to prepare a QC sample. For liver metabolite derivatization, data acquisition and analysis using GC-MS, the same procedure as followed for global Mtb H37Rv metabolite analysis was adopted.

Intracellular cytokine staining and flow cytometry: The right lobes of the lungs were harvested and incubated with collagenase (2 mg/ml) and DNase (1 mg/ml) in RPMI-1640 (2 ml) at 37°C for 30 min. The enzymatically treated tissues were passed through a strainer (70 μm) before centrifuging at 400 g for 10 min to collect the pellet. ACK lysis buffer was added to the pellet, incubated for 2 min at room temperature and quenched by adding RPMI-1640.

The cells were centrifuged and resuspended in RPMI-1640, then counted before seeding in a 12-well plate and incubated overnight with phorbol 12-myristate 13-acetate (5 ng, PMA) and ionomycin (0.75 µg). After washing, these cells were incubated with brefeldin (5 µg) and monensin (1.3 µg) for 3 hrs. The cells were washed with FACS buffer (1% FBS in PBS) and incubated with an antibody cocktail of LIVE/DEAD, CD3 and CD4 surface markers (Live dead fixable stain #L34975, CD3-FITC #11-0032-82 from Invitrogen, USA and CD4-BV785 #100552 from BioLegend) for 40 mins on ice. After staining, these cells were fixed using paraformaldehyde (4%) and permeabilized using Permeabilization buffer (#00-8333-56, Invitrogen, USA). These cells were incubated with an antibody cocktail of intracellular cytokines: TNF-α, IL-17A and IFN-γ (TNF-α-APC #17-7321-82, IL-17A-efluor450 #48-7177-82, IFN-γ-PerCP Cy5.5 #45-7311-82 from Invitrogen, USA). Flow cytometry data were acquired using BD LSR Fortessa X-20 (Supplementary Figure S10A) and analyzed using FCSExpress (DeNovo software, Version 6.0).

Statistical analysis: All data are presented as mean ± standard deviation (SD) values. Statistical analyses were performed using Origin 2020, GraphPad Prism (Version 8.4.2) and Microsoft Excel using an unpaired two-tailed t-test to identify group-specific variations. Differences were considered statistically significant with $p < 0.05$ at 95% confidence.

Data availability: All the related data is presented in the manuscript and the transcriptome data generated from DFO-treated Mtb, and control could be accessed from NCBI-Gene Expression Omnibus (GEO) using accession ID: GSE264613.

Supporting information: This article contains supporting information.

Acknowledgements: We acknowledge the Department of Biotechnology (DBT), Government of India, for supporting activities through research grants to RKN and supporting the Tuberculosis Aerosol Challenge Facility at ICGEB and ICGEB New Delhi for providing access to the ICP-MS facility and core support to RKN. Sandeep R. Kaushik

received a Senior Research Fellowship, and Nidhi Yadav and Ashish Gupta received a Junior

Research Fellowship from the DBT. Sukanya Sahu received a Senior Research Fellowship,

and Poonam Dagar received a Junior Research Fellowship from the Council of Scientific and

Industrial Research (CSIR) in New Delhi. Nikhil Bhalla is supported by the DBT-funded

project (Grant ID: National Network Project of National Institute of Immunology, New Delhi

-[40267]). We thank Ms Mothe Sravya for helping with manuscript preparation.

Author contributions: SRK, NY, and RKN conceptualized and designed experiments; SRK,

SS, AKM, N and AG performed the animal experiments; SRK, SS, AKM, NY, AG, PD and

AS carried out *in vitro,* and the laboratory profiling experiments; NB analyzed the Mtb RNA

sequencing data; SRK and NY analyzed the data and prepared the figures; TS and AKP

generated the CRISPRi knockdown of the Mtb strain; BB shared resources and reagents to

carry out the project; RKN generated funds for this work; SRK and RKN wrote the first draft

of the manuscript and revised it, incorporating the comments of all co-authors.

Conflict of interest: The authors declare that they have no conflicts of interest with the

contents of this article.

References :

1. https://www.cdc.gov/tb/topic/basics/default.htm (accessed on 31/05/2023)

2. Revised National Tuberculosis Control Programme

(RNTCP) https://tbcindia.gov.in/WriteReadData/NSP Draft 20.02.2017 1.pdf (accessed on

31/05/2023)

3. Hawn TR, Matheson AI, Maley SN, Vandal O. Host-directed therapeutics for tuberculosis:

can we harness the host? Microbiol Mol Biol Rev. 2013 Dec;77(4):608-27.

4. Sritharan M. Iron homeostasis in *Mycobacterium tuberculosis*: mechanistic insights into

siderophore-mediated iron Uptake. J Bacteriol. 2016;198(18):2399-09.

5. G. Rodriguez GM, Sharma N, Biswas A, Sharma N. The iron response of *Mycobacterium*

tuberculosis and its implications for tuberculosis pathogenesis and novel therapeutics. Front

Cell Infect Microbiol. 2022 May;12:876667.

6. Boelaert JR, Vandecasteele SJ, Appelberg R, Gordeuk VR. The effect of the host's iron

status on tuberculosis. J Infect Dis. 2007 Jun 15;195(12):1745-53.

7. Phelan JJ, Basdeo SA, Tazoll SC, McGivern S, Saborido JR, Keane J. Modulating iron for

metabolic support of TB Host Defense. Front Immunol. 2018 Oct 15;9:2296.

8. Lounis N, Maslo C, Boelaert JR, Bonnafous P, Trnffot-Pernot C, Baohong J, Grosset J.

Impact of iron loading and iron chelation on murine tuberculosis. Clin Microbiol Infect.

199;5(11):687-692.

9. Cahill C, O'Connell F, Gogan KM, Cox DJ, Basdeo SA, O'Sullivan J, Gordon SV, Keane

J, Phelan JJ. The iron chelator desferrioxamine increases the efficacy of bedaquiline in

primary human macrophages infected with BCG. Int J Mol Sci. 2021 Mar 13;22(6):2938.

10. Schaible UE, Collins HL, Priem F, Kaufmann SH. Correction of the iron overload defect

in beta-2-microglobulin knockout mice by lactoferrin abolishes their increased susceptibility

to tuberculosis. J Exp Med. 2002 Dec 2;196(11):1507-13.

11. Pisu D, Huang L, Grenier JK, Russell DG. Dual RNA-Seq of Mtb-Infected Macrophages

In Vivo Reveals Ontologically Distinct Host-Pathogen Interactions. Cell Rep. 2020

Jan;30(2):335-350.e4.

12. Pal R, Hameed S, Fatima Z. Iron deprivation affects drug susceptibilities of Mycobacteria

targeting membrane integrity. J Pathog. 2015;938523.

13. Cronjé L, Edmondson N, Eisenach KD, Bornman L. Iron and iron chelating agents

modulate *Mycobacterium tuberculosis* growth and monocyte-macrophage viability and

effector functions. FEMS Immunol Med Microbiol. 2005;45(2):103-12.

14. Rock, J.M., Hopkins, F.F., Chavez, A., Diallo, M., Chase, M.R., Gerrick, E.R., Pritchard,

J.R., Church, G.M., Rubin, E.J., Sassetti, C.M. and Schnappinger, D. Programmable

transcriptional repression in mycobacteria using an orthogonal CRISPR interference

platform. Nature Microbiology 2017 Feb 6, 2(4): 1-9.

15. Vilchèze C, Hartman T, Weinrick B, Jacobs Jr WR. Mycobacterium tuberculosis is

extraordinarily sensitive to killing by a vitamin C-induced Fenton reaction. Nature

communications. 2013 May 21;4(1):1881.

16. Kurthkoti K, Amin H, Marakalala MJ, Ghanny S, Subbian S, Sakatos A, Livny J, Fortune

SM, Berney M, Rodriguez GM. The capacity of *Mycobacterium tuberculosis* to survive iron

starvation might enable it to persist in iron-deprived microenvironments of human

granulomas. mBio. 2017 Aug 15;8(4):e01092-17.

17. Alebouyeh S, Cárdenas-Pestana JA, Vazquez L, Prados-Rosales R, Del Portillo P, Sanz J,

Menéndez MC, García MJ. Iron deprivation enhances transcriptional responses to in vitro

growth arrest of *Mycobacterium tuberculosis*. Front Microbiol. 2022 Oct 4;13:956602.

18. Sivaramakrishnan S, de Montellano PR. The DosS-DosT/DosR Mycobacterial sensor

system. Biosensors (Basel). 2013;3(3):259-282.

19. Horsburgh CR, Barry E, Lange C. Treatment of Tuberculosis. N Engl J Med 2015;

373:2149-2160

20. Theriault ME, Pisu D, Wilburn KM, Lê-Bury G, MacNamara CW, Michael Petrassi H,

Love M, Rock JM, VanderVen BC, Russell DG. Iron limitation in M. tuberculosis has broad

impact on central carbon metabolism. Commun Biol. 2022 Jul 9;5(1):685.

21. Rodriguez GM. Control of iron metabolism in *Mycobacterium tuberculosis*. Trends

Microbiol. 2006 Jul;14(7):320-7.

22. Olakanmi O, Britigan BE, Schlesinger LS. Gallium disrupts iron metabolism of

mycobacteria residing within human macrophages. Infect Immun. 2000;68(10):5619-27.

23. D. Choi SR, Switzer B, Britigan BE, Narayanasamy P. Gallium porphyrin and gallium

nitrate synergistically inhibit mycobacterial species by targeting different aspects of

iron/heme metabolism. ACS Infect Dis. 2020;6(10):2582-91.

24. Olakanmi O, Kesavalu B, Pasula R, Abdalla MY, Schlesinger LS, Britigan BE. Gallium

nitrate is efficacious in murine models of tuberculosis and inhibits key bacterial Fe-dependent

enzymes. Antimicrob Agents Chemother. 2013 Dec;57(12):6074-80.

25. Pal, R., Hameed, S., & Fatima, Z. (2015). Iron deprivation affects drug susceptibilities of

mycobacteria targeting membrane integrity. J pathog., 2015.

26. Movahedzadeh F, Wheeler PR, Dinadayala P, Av-Gay Y, Parish T, Daffé M, Stoker NG.

Inositol monophosphate phosphatase genes of Mycobacterium tuberculosis. BMC Microbiol.

2010 Feb 18;10:50.

27. Kochan I. The role of iron in bacterial infections, with special consideration of host-

tubercle bacillus interaction. Curr Top Microbiol Immunol. 1973;60:1-30.

28. Agoro R, Mura C. Iron supplementation therapy, a friend and foe of Mycobacterial

infections? Pharmaceuticals (Basel). 2019 May 17;12(2):75.

29. Anderson ER, Shah YM. Iron homeostasis in the liver. Compr Physiol. 2013

Jan;3(1):315-30.

30. McMillen S, Lönnerdal B. Postnatal iron supplementation with ferrous sulfate vs. ferrous

is-glycinate chelate: Effects on iron metabolism, growth, and central nervous system

development in sprague dawley rat pups. Nutrients. 2021 Apr 22;13(5):1406.

31. Lee DY, Kim EH. Therapeutic effects of amino acids in liver diseases: current studies

and future perspectives. Journal of Cancer Prevention. 2019 Jun;24(2):72.

32. Jeong I, Park JS, Cho YJ, Yoon HI, Song J, Lee CT, Lee JH. Drug-induced

hepatotoxicity of anti-tuberculosis drugs and their serum levels. Journal of Korean medical

science. 2015 Feb 1;30(2):167-72.

33. I. Schön T, Elias D, Moges F, Melese E, Tessema T, Stendahl O, Britton S, Sundqvist T.

Arginine as an adjuvant to chemotherapy improves clinical outcome in active tuberculosis.

Eur Respir J. 2003 Mar;21(3):483-8.

34. Qian K, Zhong S, Xie K, Yu D, Yang R, Gong DW. Hepatic ALT isoenzymes are

elevated in gluconeogenic conditions including diabetes and suppressed by insulin at the

protein level. Diabetes/metabolism research and reviews. 2015 Sep;31(6):562-71.

35. Phelan JJ, McQuaid K, Kenny C, Gogan KM, Cox DJ, Basdeo SA, O'Leary S, Tazoll SC,

Ó Maoldomhnaigh C, O'Sullivan MP, O'Neill LA, O'Sullivan MJ, Keane J. Desferrioxamine

supports metabolic function in primary human macrophages infected with *Mycobacterium*

tuberculosis. Front Immunol. 2020 May 13;11:836.

36. Lederman HM, Cohen A, Lee JW, Freedman MH, Gelfand EW. Deferoxamine: a

reversible S-phase inhibitor of human lymphocyte proliferation. Blood. 1984 Sep;64(3):748-

53.

37. Mehdiratta K, Singh S, Sharma S, Bhosale RS, Choudhury R, Masal DP, Manocha A,

Dhamale BD, Khan N, Asokachandran V, Sharma P, Ikeh M, Brown AC, Parish T, Ojha AK,

Michael JS, Faruq M, Medigeshi GR, Mohanty D, Reddy DS, Natarajan VT, Kamat SS,

Gokhale RS. Kupyaphores are zinc homeostatic metallophores required for colonization of

Mycobacterium tuberculosis. Proc Natl Acad Sci U S A. 2022 Feb 22;119(8):e2110293119.

38. Piddington DL, Kashkouli A, Buchmeier NA. Growth of *Mycobacterium tuberculosis* in

a defined medium is very restricted by acid pH and Mg(2+) levels. Infect Immun. 2000

Aug;68(8):4518-22.

39. Dow A, Sule P, O'Donnell TJ, Burger A, Mattila JT, Antonio B, Vergara K, Marcantonio

E, Adams LG, James N, Williams PG, Cirillo JD, Prisic S. Zinc limitation triggers

anticipatory adaptations in *Mycobacterium tuberculosis*. PLoS Pathog. 2021 May

14;17(5):e1009570.

40. Hodges YK, Weinberger HD, Stephens J, Horwitz MA, Horwitz LD. Desferri-Exochelin,

a lipid-soluble, hexadentate iron chelator, effectively removes tissue iron. Translational

Research. 2006 Aug 1;148(2):63-71.

41. Li X, Yang Y, Zhou F, Zhang Y, Lu H, Jin Q, Gao L. SLC11A1 (NRAMP1)

polymorphisms and tuberculosis susceptibility: updated systematic review and meta-analysis.

PLoS One. 2011 Jan 25;6(1):e15831.

42. Parker JB, Griffin MF, Downer MA, Akras D, Berry CE, Cotterell AC, Gurtner GC,

Longaker MT, Wan DC. Chelating the valley of death: Deferoxamine's path from bench to

wound clinic. Front Med (Lausanne). 2023 Feb 15;10:1015711.

 Figure legends

Figure 1: Deferoxamine (DFO) chelates iron and shows mycobactericidal effect in the

in-vitro **condition**. **A.** Schematic presentation of the method used for *in vitro* Mtb culture

with DFO and isoniazid (INH) for growth kinetics and ICP-MS. **B.** Growth kinetics of

Mycobacterium tuberculosis H37Rv strain in the absence and presence of different drugs

(DFO, INH, DFO+INH) and control. **C.** Survived Mycobacteria cultures on day five post-

drug-treated conditions and control as observed in CFU assay, showing the significant

bactericidal effect of the drugs alone or in combination. **D.** Intracellular elemental levels in

drug-treated and controlled mycobacterial culture at 5th-day post-treatment quantified using

ICP-MS. **E.** Schematic presentation of the methodology used for TEM sample preparation. **F.**

Transmission electron micrographs of untreated, DFO, INH, and Gallium nitrate (GN) treated

Mtb, respectively, harvested at the 5th day post-drug treatment. **G.** Kill curve of Mtb with

DFO, GN, INH and combination at 10x MIC of each drug. **H.** CFU plot for day four post-

inoculation**.** CFU plot for day eight post-inoculation. CF: culture filtrate, CFU: colony

forming units, and dashed line in the plots represent the limit of detection (LOD). Each time

point had five biological replicates per group. Results are determined using unpaired, non-

parametric two-tailed t-tests. ns: not significant. ns: not significant; *: $p<0.05$; **: $p<0.01$;

***: $p<0.001$; ****: $p<0.005$ at 95% confidence interval.

Figure 2: Transcriptome analysis of DFO-treated Mtb showed deregulated iron

metabolism and several other pathways. A. Schematic presentation of the method used for

RNA extraction from DFO-treated Mtb H37Rv and control for RNAseq analysis. **B.**

Identified transcripts with CPM $\geq$ 1, from the RNA-seq analysis and the deregulated

($\log_2$DFO/control $\geq \pm$ 1.0, $p < 0.05$) ones. **C.** Volcano plot showing the deregulated genes in

DFO-treated Mtb compared to control. **D.** Principal component analysis (PCA) of the RNA-

seq data (transcripts: 3826) of DFO treated Mtb and control **E.** Heatmap showing transcript

levels of iron deprivation marker genes of DFO treated Mtb and control. **F.** Transcript levels

of known iron transporters (Rv1348, Rv1349) and Mycobactin protein Utilization ViuB

(Rv2895c) in DFO-treated Mtb and control. **G.** Transcript levels of dormancy-inducing genes

dosT (Rv2027c), *dosS* (Rv3132) and *dosR* (Rv3133) in DFO-treated Mtb and controls. **H.**

Gene Ontology analysis of the deregulated transcripts using the DAVID tool in DFO-treated

Mtb. **I.** Deregulated biological processes, as identified from the pre-ranked FGSEA (iDEP.96

tool) in DFO-treated Mtb compared to control.

Figure 3: DFO alters Mtb global metabolite profile in-vitro. A. Schematic presentation of

the method used for metabolite profiling from DFO-treated Mtb H37Rv and control using

GC-MS. **B.** Principal component analysis showing DFO-treated Mtb H37Rv clustering

separately from the control group. **C.** Deregulated metabolites observed between DFO-

treated and control Mtb. **D.** Pathway analysis showing overrepresented metabolic pathways in

DFO-treated Mtb.

Figure 4: DFO aids in Mycobacterial clearance in C57BL/6 mice. A. Schematic

presentation of the method used to infect C57BL/6 mice with *Mycobacterium tuberculosis*

H37Rv strain and treatment using DFO. **B.** Prussian Blue stained liver histology for DFO-

treated mice showing decreased liver iron accumulation indicated by black arrows. **C.** Liver

iron levels were estimated using ICP-MS in the infected control compared with Mtb-infected

mice receiving DFO treatment. Mycobacterial load in the lungs **(D)**, spleen **(E)** and liver **(F)**

of the control and DFO treated *Mycobacteria tuberculosis* H37Rv infected C57BL/6 mice.

Each time point had five biological replicates per group. The dashed line in the plots

represents the LOD. i.p.: intra-peritoneal injection; dpi: days post infection; CFU: colony

forming units. Results are determined using unpaired, non-parametric two-tailed t-tests. ns:

not significant. *: p<0.05; **: p<0.01; ***: p<0.005; p-values at 95% confidence interval.

Figure 5: DFO adjunct to anti-tuberculosis drugs (INH/H and RIF/D) minimally affects lung mycobacterial burden at later timepoint. **A.** Schematic presentation of the method used for Mycobacterial infection and treatment using INH/H and RIF/R with or without DFO/D. **B.** Prussian blue staining of the liver tissue of RHD-treated mice (InRHD) shows decreased iron accumulation indicated by black arrows compared to IR-treated and infected control mice. **C** Liver iron levels estimated using ICP-MS in the infected controls and infected mice receiving INH and RIF with DFO. **D.** Lung mycobacterial load of the *Mycobacteria tuberculosis* H37Rv infected C57BL/6 mice receiving INH and RIF with or without DFO. Each time point had five biological replicates per group. The dashed line in the plots represents the LOD. i.p.: intra-peritoneal injection; dpi: days post-infection; wpt: weeks post-treatment, CFU: colony forming units. Results are determined using unpaired, non-parametric two-tailed t-tests. ns: not significant. **: p<0.01, ns: not significant; up: upregulated, down: downregulated, p-values at 95% confidence interval.

Figure 6: **DFO adjunct to anti-tuberculosis drugs (INH/H and RIF/D) accelerated early mycobacterial clearance in C57BL/6 mice and impacts host liver metabolome A.** Schematic presentation of the method used for Mycobacterial infection, treatment using INH and RIF with or without DFO and monitoring host tissues at early time points **B.** Lung mycobacterial load one wpt. **C.** Frequency of CD4$^+$ T cells and F4/80+ macrophages in the lungs of *Mycobacteria tuberculosis* H37Rv infected C57BL/6 mice receiving INH and RIF with or without DFO at the first week of treatment. **D.** Liver metabolome of Mycobacterial infected mice treated with anti-tuberculosis drugs with DFO vs without DFO at the first week of treatment. **E.** Scatter dot plot showing pathway enrichment analysis of deregulated

metabolites from **D** plot. **F.** Lung mycobacterial load two wpt. **G.** Frequency of CD4$^+$ T cells

and F4/80+ macrophages in lungs of *Mycobacteria tuberculosis* H37Rv infected C57BL/6

mice receiving INH and RIF with or without DFO at the second week of treatment. **H.** Liver

metabolome of Mycobacterial infected mice treated with anti-tuberculosis drugs with DFO

vs without DFO at the second week of treatment. **I.** Scatter dot plot showing pathway

enrichment analysis of deregulated metabolites from **H** plot. Each time point had five

biological replicates per group except InRH group for immune cell distribution at 2 wpt. The

dashed line in the plots represents the LOD. i.p.: intra-peritoneal injection; dpi: days post-

infection; wpt: weeks post-treatment, CFU: colony forming units. Results are determined

using unpaired, non-parametric two-tailed t-tests. ns: not significant. *: $p<0.05$; **: $p<0.01$;

****: $p<0.0001$, ns: not significant; up: upregulated, down: downregulated, p-values at 95%

confidence interval.

Figure 7: Schematic showing the effect of iron overloading and iron chelation on

Mycobacterial growth, and tissues of Mtb infected mice. Mtb: Mycobacterium

tuberculosis, INH: Isoniazid, RIF: Rifampicin, IDO: Indoleamine 2, 3-dioxygenase, TNF:

Tumour necrosis factor, IFN: Interferon, IL: Interleukin.

Supplementary Table and Figure legends

Supplementary Figure S1: A and B. Intracellular copper and zinc levels in drug-treated and control Mycobacterial culture at 5th-day post-treatment quantified using ICP-MS. Results are determined using unpaired, non-parametric two-tailed t-tests. n = 5 per group, ns: not significant; *: $p<0.05$, **: $p<0.01$; ****: $p<0.0001$ at 95% confidence interval.

Supplementary Figure S2: Transmission electron micrographs of untreated Mtb **(A)**, DFO treated **(B)**, INH treated **(C)**, GN treated **(D)** harvested at 5th day post-drug treatment. Images with dashed boundaries are referred in the main figures.

Supplementary Figure S3: DFO alters iron metabolism and various pathways of Mtb H37Rv. A. Heatmap showing the deregulated (up in red and down in green) genes in the DFO (D1, D2, D3) vs control Mtb (C1, C2) as evident from the Z-score values. **B.** Pathway analysis using Preranked FGSEA in IDEP.96 tool. Red indicates upregulated and green shows down-regulated pathways in Mtb. The intensity of the colour and diameter of the node corresponds to the number of genes impacted in a particular pathway. **C.** Iron-sulfur cluster and iron-binding proteins impacted upon DFO treatment. Green shows presence and Black shows absence of the Iron-sulfur/Iron-binding nature of the protein.

Supplementary Figure S4: irtA knock down in Mtb inhibits its growth and DFO exposure reduces Mtb growth in J774A1 cells. A. Relative suppression of *irtA* and *irtB* expression levels in *irtAKD* and cntrl sgRNA strains (mean ± SEM, n = 3 biological replicates). **B.** Relative suppression of *irtA* and *irtB* expression levels in *irtAKD* and cntrl sgRNA strains **C.** Fluorescent confocal microcopy images acquired at 60X magnification showing the effect of DFO pretreatment on Mtb H37Rv infected J774A.1 murine cells. Blue color represents nuclei stained with DAPI and red color indicates mitochondria stained with

MitoTracker red. Results are determined using unpaired, non-parametric two-tailed t-tests

**P $\leq$ 0.005 and ***P $\leq$ 0.0005, ****P 852 $\leq$ 0.00005. Data represents mean $\pm$ SEM

(standard error mean). (KD=knock down, cntrl=control)

Supplementary Figure S5: A. Schematic representation of the method used to create iron

loading conditions B. Prussian blue staining of the liver tissue at 40X magnification

indicating high iron accumulation shown by black arrows in the FcHealthy C57BL/6 mice as

compared to healthy controls C. Body weight of the C57BL/6 mice before, during and after

iron loading for FcHealthy in comparison to healthy controls D. Liver iron levels, estimated

using ICP-MS at different time points during iron loading as compared to healthy controls.

Scale bar for histology images: 50 μm, Fc: ferric carboxymaltose supplemented group.

Results are determined using unpaired, non-parametric two-tailed t-tests where ns: not

significant; **: $p<0.01$; ***: $p<0.001$ at 95% confidence interval.

Supplementary Figure S6: A. Schematic presentation of the method used to create iron

loading condition, Mycobacterial infection and treatment (with isoniazid: INH and

rifampicin: RIF) experiment. B. Liver iron levels, as estimated using ICP-MS, in iron loaded

and control mice groups. C. Prussian Staining of liver tissues at 40X magnification indicates

high iron accumulation shown by black arrows in iron-loaded mice and did not change

significantly upon drug treatment (D). Mycobacterial load in the lungs (E), spleen (F) and

liver (G) of the Mycobacteria tuberculosis H37Rv infected C57BL/6 mice. Dashed lines in

the CFU plots represent the limit of detection. n = 5/group/time points; i.v.: intra-venous

injection; dpi: days post-infection; CFU: colony forming units; scale bar for histology

images: 50 μm. Results are determined using unpaired, non-parametric two-tailed t-tests

where ns: not significant, p values at 95% confidence interval.

Supplementary Figure S7: Histology images of liver tissue stained with Prussian blue

staining showing iron accumulation in H (A), FcH (B), In (C), FcIn (D), InRH (E), FcInRH

(F) as compared to minimum iron in InD (G) and InRHD (H) groups at 4X, 10X, 40X

magnification respectively. Scale bar: 50μm, In: Mtb infected C57BL/6 mice; FcIn: Mtb

infected Fe supplemented C57BL/6 mice; InD: Deferoxamine (DFO) treated Mtb infected

mice; InRH: Isoniazid (INH) and Rifampicin (RIF) treated Mtb infected C57BL/6 mice;

FcInRH: INH and RIF treated Mtb infected iron supplemented C57BL/6 mice; InRHD: Mtb

infected C57BL/6 mice receiving both DFO, INH and RIF; Healthy: Control mice (H);

FcHealthy: Control mice with iron supplementation (FcH).

Supplementary Figure S8: Liver metabolome changes upon treatment with INH and RIF

with or without DFO at 1st and 2nd weeks. InRH vs In **(A)** and InRHD vs In **(B)** InRH vs In

(A) and InRHD vs In. ns: non-significant, up: upregulated, down: down regulated, p value at

95 % confidence interval, In: H37Rv infected control, InRH: H37Rv infected and INH+ RIF

treated, InRHD: H37Rv infected and INH+RIF +DFO treated, wpt: weeks post treatment, p

values at 95 % confidence interval.

Supplementary Figure S9: Pathway enrichment analysis results for treated groups in

comparison to infected controls at one and two wpt. wpt: weeks post treatment, p values at

95% confidence interval.

Supplementary Figure S10: A. Schematic presentation of the method used for Mtb

infection, treatment and intracellular cytokine staining of the lung cells **B.** CD3+ and CD4+ T

cell frequency in the infected (In) vs INH+ RIF treated with or without DFO **C.** Percentage of

CD4+ T-cells that are IFN-γ, IL-17A and TNF-α positive in the infected (In) vs INH+ RIF

treated with or without DFO. n = 5/group/time points for In, InRH group and n = 4 for

InRHD group; Results are determined using unpaired, non-parametric two-tailed t-tests. ns:

not significant.

Supplementary Table S1: The list of DEG in DFO-treated Mtb H37Rv compared to

control as determined using edgeR in iDEP.96 platform. The DEGs having FC>2

(log2FC>1), at BH adjusted P-value < 0.05 were considered significant.

Supplementary Table S2: Expression levels of genes involved in iron metabolism in

Mycobacterium tuberculosis. Normalized expression matrix was used for paired, 2-tailed

student's t-test for determination of P-values. The gene set was borrowed from a review by

Rodriguez et. al. (Tends in microbiology, 14(7), 2006).

Figure 1

Figure 2

Figure 3

Figure 4

A

B 40X

C

D

E

F

Figure 5

A

B

C

D

Figure 6

Figure 7

Graphical abstract