Single-cell transcriptomic and TCR analysis of human Cytomegalovirus (hCMV)-specific

memory T cells reveals effector and pre-effectors of CD8+- and CD4+-cytotoxic T cells

Running title: Single-cell RNA-Seq analysis of hCMV-specific T cells

Raunak Kar[1], Somdeb Chattopadhyay[1], Anjali Sharma[2,#], Kirti Sharma[1,#], Shreya Sinha[1],

Gopalakrishnan Aneeshkumar Arimbasseri[1,*], Veena S. Patil[1,*,$]

[1]National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi

[2]Vardhman Mahavir Medical College and Safdarjung Hospital, New Delhi.

[#]Equal contribution

[*]Correspondence can be addressed to GAA (aneesh@nii.ac.in) or VSP (veena@nii.ac.in)

[$]Lead contact

Key Words: hCMV, antigen-specific T cells, CD4-CTLs, CD8-CTLs, T_{REGs}, single-cell RNA-

Seq, single-cell TCR-Seq, TCR

Abstract

Background

Latent human Cytomegalovirus (hCMV) infection can pose a serious threat of reactivation and disease occurrence in immune-compromised individuals. Though, T cells are at the core of the protective immune response to hCMV infection, a detailed characterization of different T cell subsets involved in hCMV immunity is lacking.

Results

Here, in an unbiased manner, we characterized over 8000 hCMV-reactive peripheral memory T cells isolated from seropositive human donors, at a single-cell resolution by analyzing their single-cell transcriptomes paired with the T cell antigen receptor (TCR) repertoires. The hCMV-reactive T cells were highly heterogeneous and consisted of different developmental and functional memory T cell subsets such as, long-term memory precursors and effectors, T helper-17, T regulatory cells (T_{REGs}) and cytotoxic T lymphocytes (CTLs) of both CD4 and CD8 origin. The hCMV-specific T_{REGs}, in addition to being enriched for molecules known for their suppressive functions, showed enrichment for the interferon response signature gene sets. The hCMV-specific CTLs were of two types, the pre-effector and effector-like. The co-clustering of hCMV-specific CD4-CTLs and CD8-CTLs in both pre-effector as well as effector clusters suggest shared transcriptomic signatures between them. The huge TCR clonal expansion of cytotoxic clusters suggest a dominant role in protective immune response to CMV.

Conclusions

The study uncovers the heterogeneity in the hCMV-specific memory T cells reveling many functional subsets with potential implications in better understanding of hCMV-specific T cell immunity. The data presented can serve as a knowledge base for designing vaccines and therapeutics.

Background

The acquisition of immunological memory to infections is the hallmark of protective immune response. During this conserved process of T cell immunological memory development, the naive T cells that have not previously encountered antigen, differentiate during the primary infection, into memory T cells that have specialized functions in immune defence to a subsequent infection with the same pathogen. A small number of antigen-specific memory T cells can have an enormous impact in directing or misdirecting an immune response upon secondary infection. However, the rarity of antigen-specific memory T cells have limited the detailed characterization of these cells until very recently. The utilization of single-cell genomics to dissect the heterogeneity and understand th global gene expression patterns of these rare antigen-specific memory T cells has advanced our understanding of various disease-specific T cell memory repertoire in an unbiased manner [1-7].

CD4$^+$ memory T cells are classified based on their functional properties and in addition more recently also based on their transcriptomes, into multiple subsets such as but not limited to, T helper (T$_H$)1, T$_H$2, T$_H$17, T$_H$1/17, T follicular helper (T$_{FH}$), and T regulatory (T$_{REG}$) cells [8-12]. For example, T$_{FH}$ cells are the specialized providers of help to B cells, and T$_{REG}$s, the regulators of the immune response with their immunosuppressive properties [11, 13, 14]. Furthermore, each of those subtypes can exhibit great functional diversity and plasticity [12]. The peripheral CD4$^+$ as well as CD8$^+$ memory T cells can also be classified based on their developmental stages as long-lived precursor memory and short-lived effector memory T cells such as, stem cell memory T cells (T$_{SCM}$), central memory T cells (T$_{CM}$), and effector memory T cells (T$_{EM}$), effector memory T cells expressing CD45RA (T$_{EMRA}$) [15]. The CD8$^+$ T cells are known for their cytotoxic functions, and the CD4$^+$ T cells, predominantly, for their helper function where they provide help to other immune cell types including the CD8$^+$ T cells in resolving the infection. However, others and we have shown that CD4$^+$ T cells can also function

as cytotoxic T cells and are enriched in the CD4-T_{EMRA} compartment and show marked T cell

antigen receptor (TCR) clonal expansion similar to the CD8-T_{EMRA} cells in many acute and

chronic viral infections including the cytomegalovirus (CMV) [1, 3, 4, 6, 16-22].

Following initial infection, the human CMV (hCMV) establishes persistence or latency

in immune-competent individuals, although intermittent viral shedding can occur [23]. In the

absence of an effective immune response especially in immune-compromised individuals, the

virus can reactivate and cause the disease. Hence, reactivation of hCMV poses a higher risk for

individuals undergoing transplant or chemotherapy, immune-compromised Human

Immunodeficiency Virus (HIV) infected individuals, and new-borns of hCMV seropositive

mothers [23]. Both CD4$^+$ and CD8$^+$ T cells have been shown to play important roles in the

immune response to CMV [24, 25]. While hCMV-specific CD8$^+$ T cells have extensively been

studied, studies on CD4$^+$ T cells are sparse [2]. Considering that the emergence of the hCMV-

specific CD4$^+$ T cells precedes that of CD8$^+$ T cells, they were mostly thought to perform

supportive roles by enhancing the CD8$^+$ T cell responses to the hCMV infection. However,

subsequent studies have demonstrated additional roles of CD4+ T cells, other than the

supportive one towards CD8$^+$ T cell, in CMV immunity (reviewed in [2]). The poorer or lack

of hCMV-specific CD4$^+$ T cell responses have been associated with persistent viral shedding

in urine and saliva as well as severe disease outcomes [24-27]. In individuals undergoing stem

cell or organ transplantation, hCMV-specific CD4$^+$ and CD8$^+$ T cells have been shown to

abrogate reactivation [25-29]. Further, during the primary infection, the role of CD4$^+$ T cells

has been established in resolution of the disease symptoms [30, 31]. The identification of

hCMV-specific T cells has classically relied on the measurement of intracellular cytokine

production, predominantly interferon gamma (IFNγ) and to a lesser extent interleukin 10

(IL10), in response to stimulation with peptides derived from various hCMV ORFs (Open

Reading Frame) or total viral lysate [2, 6, 16, 32-35]. Several studies, in healthy seropositive

individuals, have identified a higher percentage of hCMV-specific T cells, reaching almost as high as ~5-10% of total CD4$^+$ and CD8$^+$ T cells for epitopes derived from pp65 and IE-1 proteins of hCMV, with majority of them showing terminally differentiated effector memory phenotype [16, 33, 34]. In addition to producing IFNγ, the hCMV-reactive CD4$^+$ T cells have been shown to make cytolytic molecules such as Granzymes, Perforin, and CD107a (degranulation marker), and showed efficient lysis of hCMV infected dendritic cells and controlled virus dissemination *in vitro*, suggesting prominent role of cytotoxic CD4$^+$ T cells in hCMV immunity [16-20]. CD4$^+$ T cells expressing CD25 and FOXP3 (T$_{REGs}$), producing IL10, an immunosuppressive cytokine, have also been observed and are implicated in immune suppression in CMV infection and reactivation [36-39]. However, a detailed characterization of the spectrum of hCMV-specific CD4$^+$ memory T cells with respect to their heterogeneity and molecular signatures have been lacking. Hence, the identification and characterization of the overall hCMV-specific memory T cells with respect to their molecular signatures, in CMV seropositive individuals who are otherwise healthy, will enable the identification of correlates of protective immune response to hCMV.

Hence, in this study, to isolate the spectrum of hCMV-specific T cells, independent of their cytokine profile and Human Leukocyte Antigen (HLA) restriction or not limited to a single peptide (peptide-MHC multimers), we used ARTE (Antigen-Reactive T cell Enrichment) assay [40, 41]. The ARTE assay utilizes the surface expression of co-stimulatory molecules (CD154 and CD137) upon TCR engagement with cognate peptide-MHC (Major Histocompatibility Complex) complex [40, 41]. To dissect the heterogeneity across T cell memory subsets and TCR clonal diversity of hCMV-specific memory T cells, we isolated and performed single-cell transcriptomics analysis of over 8000 hCMV-specific memory T cells, isolated based on the surface expression of CD154$^+$ and/or CD137$^+$ across 10 seropositive donors, after stimulation with pool of hCMV peptides derived from T cell immuno-dominant

epitopes of pp65 and IE-1 proteins. Unbiased clustering based on single-cell transcriptomes identified nine distinct clusters of various functional subsets including, T_{REGs}, T_H17, T_H1 and $T_H1/17$, and cytotoxic CD4$^+$ T cells (CD4-CTLs). The hCMV-antigen specific T_{REGs} were enriched for molecules linked to their suppressive function and interferon response genes ($T_{REG-IFN}$). The transcriptomics analysis combined with TCR repertoire analysis revealed that the cytotoxic cells are predominantly of 2 types; pre-effectors and effectors and are highly clonally expanded. Interestingly, we observed that the transcriptomes of pre-effectors and effectors of CD4-CTLs and CD8-CTLs were largely indistinguishable. Together, in an unbiased approach, our study has identified many previously known as well as novel subsets of hCMV-specific memory T cells and their unique transcriptomic signatures, which potentially have implications in understanding the reactivation of hCMV in immune-compromised individuals. This knowledge base can be an important resource for designing vaccines and therapeutics.

Results

Single-cell transcriptomic analysis of CMV-specific memory T cells

To identify and understand the heterogeneity and clonal diversity of the spectrum of hCMV-specific T cells across different memory subsets in an unbiased way, we performed single-cell transcriptomic (scRNA-Seq) and single-cell TCR (T Cell antigen Receptor) repertoire analysis (scTCR-Seq) of >8000 hCMV-specific memory T cells (Fig. 1A and B). To isolate the spectrum of hCMV-specific T cells independent of their cytokine profile and HLA restriction, we utilized the widely used ARTE (Antigen-Reactive T cell Enrichment) assay where the surface expression of co-stimulatory markers such as CD154 and CD137 is analysed upon the engagement of T cell antigen receptor (TCR) with cognate pMHC (peptide Major Histocompatibility Complex) complex [1, 3, 7, 40-42]. For single-cell omics experiments, we

isolated T cells using florescence activated cell sorting (FACS) based on the surface expression

of CD154 and CD137 upon stimulation of PBMCs (Peripheral Blood Mononuclear Cells) from

10 hCMV-seropositive and high peptide responder individuals with peptide pools derived from

immunodominant pp65 and IE-1 (Fig. 1A and Additional file 1: Fig. S1A-C). To compare the

hCMV-specific memory T cells across known memory subsets, the peptide responding T cells,

from different $CD4^+$ memory T cell compartments (naïve T cells (T_N; $CD45RA^+CCR7^+$),

central memory T cells (T_{CM}; $CD45RA^-CCR7^+$), effector memory T cells (T_{EM}; $CD45RA^-$

$CCR7^-$), and effector memory T cells expressing CD45RA (T_{EMRA}; $CD45RA^+CCR7^-$) [15]),

were FACS sorted separately and pooled before performing single-cell omics experiments (Fig.

1A and Additional file 1: Fig. S1B and C). Further, considering the enrichment of CD4-CTLs

(cytotoxic T lymphocytes) in the CD4-T_{EMRA} subset, to parallelly compare the hCMV-specific

CD4-CTLs and CD8-CTLs we also included hCMV-reactive T cells from CD8-CTL

compartments ($CD8^+CCR7^-$ (CD8-T_{EM} and -T_{EMRA})) in the single-cell omics experiments (Fig.

1A and Additional file 1: Fig. S1B and C) [4]. An unbiased clustering of 6899 hCMV-specific

memory T cells (out of 8062 cells) that passed our quality control parameters (QC; please see

methods section), identified 9 distinct clusters based on transcriptomes (Fig. 1B), and the

clustering was not influenced by donors or the experimental batch (Additional file 1: Fig. S1D

and E; Additional file 2: Table S1). Different developmental T cell memory subsets were

distinguished based on the surface expression of oligo-tagged antibodies (antibody derived

tags; adt) against CD45RA, CCR7, CD95 and IL7Rα as well as *CCR7*, *CD27*, *CD28* transcripts

(Fig. 1C and D; Additional file 2: Table S2) [4, 15]. Transcripts for *CD4* were distributed across

all clusters while *CD8B* transcripts were majorly found in clusters 2 and 4 (Fig. 1E). Cells in

cluster 2 and 4 were mostly T_{EMRA} cells of both CD4 and CD8 origin as they expressed

CD45RA (adt CD45RA) and lacked the expression of CCR7 protein (adt CCR7), as well as

transcripts for *CCR7*, *CD28* and *CD27* (Fig. 1C and D) [4, 15]. T_N cells were very few and

were mostly found in cluster 6 (CD45RA$^+$CCR7$^+$IL7Rα$^+$) (Fig. 1C and D). T$_{CM}$ and T$_{EM}$ cells

were distributed across multiple clusters (Clusters 0, 1, 3, 5, 6, 7) (Fig. 1C and D). These

observations were independently confirmed by the enrichment of subset-specific gene

signatures from T$_{CM}$, T$_{EM}$, and T$_{EMRA}$, as well as common gene signatures from T$_{CM}$-T$_{EM}$, and

T$_{EM}$-T$_{EMRA}$ [4] (Fig. 1F; Additional file 1: Fig. S1F; Additional file 2: Table S3). Both CD4-

and CD8-T$_{EMRA}$ cells (clusters 2 and 4) majorly showed higher expression of *CD137* while

other T$_{CM}$ and T$_{EM}$ clusters (0, 3, 6) showed higher expression of *CD154* (Fig. 1G).

Interestingly many of the T cell activation markers showed dynamic expression amongst the

clusters; *CD154* and *CD137* showed opposing expression and similar observations were made

for *CD69* and *IL2RA* (Fig. 1D and G). Further, co-expression analysis of various T cell

activation markers in combination, confirmed the variability in expression (Fig. 1G). Most

strikingly, the *CD137*^{hi}*CD154*^{lo} cells showed low *CD69* expression, while the *CD154* and

CD137 co-expressing cells expressed similar amounts of *TNFRSF4* transcripts (encoding

OX40) (Fig. 1G). Together, these results show that our data well represents various memory

subsets that show differential expression of T cell activation markers.

hCMV-specific memory T cells are heterogeneous and can be classified into multiple

functional T cell subsets

The differential expression analysis of transcripts amongst the nine clusters of hCMV-specific

T cells, identified cluster specific expression of transcripts, revealing many functional T cell

subsets (Fig. 2A and B; Additional file 2: Table S4). Multiple clusters showed enrichment for

long-term memory markers such as *AQP3*, *CCR7*, *TCF7*, *IL7R*, *LTB*, *JUNB, SELL* (Fig. 2A

and B) [3, 4]. Cluster 1 showed enrichment for *FOXP3*, *BATF*, *TIGIT* and *TNFRSF1B*

transcripts that are known to be expressed in T regulatory cells (T$_{REG}$) (Fig. 2B). Interestingly,

cluster 1 also showed enrichment of transcripts encoding interferon (IFN) response signature

genes (*ISG20*, *ISG15*, *IFIT3*, *OAS1*, *MX1*) (Fig. 2B). In an independent analysis, cluster 1 cells

showed a higher enrichment signature score for overall T_{REG} and IFN response signature gene sets (Fig. 2C; Additional file 2: Table S3), suggesting that cells in cluster 1 are T_{REG} cells and are possibly responding to IFN signalling ($T_{REG-IFN}$). Actively cycling cells were found in cluster 3, one of the clusters enriched for T_{CM} signatures (Fig. 2B and C; Additional file 2: Table S3) [43]. Cells in clusters 2 and 4, enriched for T_{EMRA}-specific gene sets, expressed transcripts encoding cytolytic molecules such as *GZMB*, *PRF1* and *NKG7* and showed enrichment for cytotoxicity gene signatures (Fig. 1F, Fig. 2B and C; Additional file 1: Fig. S1F; Additional file 2: Tables S3 and S4) [4], suggesting that clusters 2 and 4 are cytotoxic clusters. The presence of mixture of cells expressing *CD4* and *CD8* transcripts in both the cytotoxic clusters 2 and 4 suggest possible similarities between CD4-CTLs and CD8-CTLs.

We next interrogated the single-cell transcriptome of hCMV-specific T cells for functional T helper (T_H) cell subtypes such as T_H1, T_H17 and $T_H1/17$, that are known to have role in viral infections [3, 44]. Multiple clusters showed expression of T_H subset-specific transcripts: T_H17 (*CCR4*, *CCR6*, *IL4I1*, *CTSH*), T_H1 (*CXCR3*, *TNF*, *IFNG*, *CSF2*) and $T_H1/17$ (*CXCR3, DPP4*) [9, 10] (Fig. 2D; Additional file 2: Table S4). Interestingly, we found several clusters co-expressing transcripts from different T_H subsets, suggesting each cluster may have cells belonging to more than one T_H subtype. Independently, module enrichment analysis using T_H subset-specific gene sets corroborated these results (Additional file 1: Fig. S2A; Additional file 2: Table S3). The T_H1-specific gene sets were enriched in cluster 2 and 4, the cytotoxic clusters (Additional file 2: Fig. S2A). Considering there are several overlapping genes between T_H1 and cytotoxic gene sets, this observation is in alignment with the known literature [4, 9, 10]. T_H17 signature genes showed highest enrichment in cluster 0, followed by clusters 7, 3 and 1. $T_H1/17$ signatures were mostly enriched in cluster 0 followed by clusters 5 and 7 (Additional file 1: Fig. S2A) [9, 10]. Together these observations indicate that each hCMV-specific T_H subset is at different developmental stages of the memory (long-term memory and

221 effector memory) and clustering was predominantly influenced by the developmental stages of

222 memory T cells

TCR clonally expanded hCMV-specific T cells are majorly observed in cytotoxic clusters.

To understand the TCR clonal diversity amongst the hCMV-specific T cells we analysed the

single-cell TCR-Seq data generated from the same cells. We recovered TCRs from 83.56%

(5765 cells of 6899 cells) of the total QC-ed cells in our data set, of which 21.40% (1234 of

5765 cells) were clonally expanded (Fig. 2E; Additional file 2: Table S5). Several clusters

shared TCR clonotypes, indicating a shared developmental lineage between these clusters

(Additional file 1: Fig. S2B). A maximum of 14 clonotypes were shared between clusters 2 and

4, followed by clusters 0 and 3 sharing 10 clonotypes (Additional file 1: Fig. S2B). Cells in

cluster 2 and 4 showed highest clonal expansion with ~78% and ~76% of the cells being

expanded respectively (Fig. 2E; Additional file 2: Table S5). As high as 76% of the total

expanded cells (938 of 1234) were found in cluster 2 (50%) and cluster 4 (~26%) (Additional

file 2: Table S5). A few top expanded clonotypes were found in as many as 144, 79 and 51

cells from clusters 2 and 4, indicating the preferential expansion of a few specific clonotypes

(Fig. 2F; Additional file 1: Fig. S2C). Interestingly, close to 58% of the expanded cells carried

only 20 clonotypes (Fig. 2F). Both CD4-CTLs and CD8-CTLs (T_{EMRA}) are shown to be

clonally expanded in several diseases including CMV in both humans and animal models [2-

4, 45, 46]. Clonal expansion and enrichment for cytotoxic signature in clusters 2 and 4

emphasizes the role for cytotoxic cells in hCMV-specific protective immune response [16-20].

Single-cell transcriptome analysis of hCMV-specific T_{REG} cells

T_{REG}s are characterised by the higher expression of CD25 (IL2RA) and the transcription factor

FOXP3, but a lower expression of IL7Rα (CD127) in humans [47]. CMV-specific T_{REG}s have

been reported in both mouse models and humans following primary infections and are shown

to produce IL10, a suppressive cytokine [2, 6, 36-39, 48-50]. However, the detailed molecular

signatures of hCMV-specific memory T_{REGs} have not been well described. The transcript

encoding the T_{REG} specific transcription factor FOXP3 was differentially expressed in clusters

1 (85% of cells) and 7 (Fig. 2B, Fig. 3A and B; Additional file 2: Table S4), however though

not differentially expressed, a portion of cells from cluster 5 showed higher *FOXP3* expression

(41% of cells), while various other clusters (clusters 0 and 3) showed lower levels of *FOXP3*

expression (Fig. 3A and B). Considering FOXP3 can be transiently expressed on TCR activated

cells, we examined the co-expression of other T_{REG} related genes and *FOXP3*. The cells co-

expressing *FOXP3 - IL2RA*, and *FOXP3 - IKZF2* were enriched in cluster 1 and portion of

cluster 5, and as expected they showed lower expression of IL7Rα (Fig. 1C and Fig. 3A) [47].

Clusters 1 and 5 were also characterized by the higher expression of other T_{REG}-specific

transcripts such as *TIGIT*, *BATF*, *TNFRSF1B* and the Ikaros Zinc Finger (IkZF) transcription

factor (TF) family members *IKZF1* encoding Ikaros, *IKZF2* encoding Helios, *IKZF3* encoding

Aiolos, *IKZF4* encoding Eos (Fig. 2B and Fig. 3C). The overall T_{REG} gene signature was

enriched in cluster 1 and fraction of cluster 5 (Fig. 2C). Together these results show that T_{REG}

cells are predominantly in cluster 1 and portion of cluster 5 and the cells from clusters 0 and 3

are potentially non-T_{REG} cells, transiently expressing low levels of *FOXP3* in response to TCR

activation. We next interrogated if the hCMV antigen-specific T_{REGs} in our data set were indeed

memory T_{REGs}. Based on the expression of CD45RA, the FOXP3$^+$ T_{REGs} have been classified

as resting or naïve T_{REGs} (FOXP3^{lo}CD45RA$^+$) and effector T_{REGs} (FOXP3^{hi}CD45RA$^-$) [51, 52].

Further the effector T_{REGs} have been classified as central and effector memory cells based on

the expression of CCR7, CD27 and CD28 [53]. The T_{REGs} in our data set were characterized

by the lack of expression of CD45RA, and a moderate expression of CCR7, *CD27* and *CD28*,

indicating that they are mostly central and effector memory T_{REGs} rather than naïve T_{REGs} (Fig.

1C and D).

Although, T$_{REGs}$ expressing FOXP3, are known for their suppressive function, many

reports have shown the presence of T cells expressing FOXP3 transiently, in response to TCR

activation that are non-suppressive in nature. Hence, we assessed if the hCMV-specific T$_{REGs}$

in our data set are expressing markers associated with the suppressive nature of T$_{REGs}$. The

molecular signatures of T$_{REGs}$ found in clusters 1 and 5 indicated their potential suppressive

nature, compared to cells expressing low levels of *FOXP3* observed in clusters 0 and 3 (Fig.

3A). The T$_{REGs}$ in clusters 1 and 5 were $CD137^+CD154^{lo}$, the activation induced characteristic

of T$_{REGs}$ implicative of epigenetically stable antigen-activated T$_{REGs}$ that retain the high

suppressive potential in humans after *in vitro* or *ex vivo* activation (Fig. 1D and G) [54]. The

T$_{REGs}$ in clusters 1 and 5, showed higher expression of *TIGIT*, *TNFRSF18* (GITR) and *CTLA-*

4, genes that are associated with T$_{REGs}$ containing high affinity TCRs, further suggesting their

antigen-specificity (Fig. 2B and Fig. 3C). The T$_{REGs}$ in clusters 1 and 5 also showed higher

expression of *ENTPD1* (CD39), an ectoenzyme ATP apyrase, *CCR4*, *CTLA4*, Helios (*IKZF2*)

and FAS (CD95), molecules that are reported to be associated with effector and suppressive

function (Fig. 1C, Fig. 2D and Fig. 3C) [55-57]. A suppressive function of Helios in T$_{REGs}$ has

been suggested, where Helios$^+$FOXP3$^+$ T$_{REG}$ cells produced lower amounts of T$_H$17-specific

cytokine, IL17, compared to Helios$^-$FOXP3$^+$ T$_{REG}$ cells, hence promoting the formation of

T$_{REGs}$ rather than T$_H$17 subtype [57]. Consistent with these observations, we found that clusters

0 and 3 that expressed *IL17F* and *IL17A*, expressed low levels of *FOXP3*, but lack the

expression of *IKZF2* (Helios) and these cells showed higher expression of *CD154* and lower

expression of *CD137* (Fig. 2B, Fig. 3A andC) [54, 57]. Together these observations suggest

that the low *FOXP3* expressing cells in clusters 0 and 3 are potentially the induced/transient

T$_{REGs}$ that are possibly expressing FOXP3 transiently in response to activation and are non-

suppressive in nature. Conversely, T$_{REG}$ cells in cluster 1 and 5 are *Helios$^+$FOXP3$^+$*, and did

not express *IL17A* and *IL17F*, indicative of suppressive nature (Fig. 2B, Fig. 3A and C). The

lack of expression of exhaustion markers *PDCD1* and *LAG3* further suggests these are not

associated with anergy or exhaustion (Fig. 3C). Together these results provide multiple

evidences to suggest that the T$_{REGs}$ in clusters 1 and 5 are effector memory T$_{REGs}$ and are

potentially suppressive in nature. One of the mechanisms through which T$_{REGs}$ execute their

suppressive function on effector T cells is through cytokines such as IL10, TGFβ and IL35

(reviewed in [58]). The T$_{REGs}$ in cluster 1 lacked or showed low expression of these suppressor

cytokine transcripts *TGFB1*, *IL10* and *IL35*, indicating that cells in cluster 1 may not be actively

suppressing the hCMV-specific effector T cells, at least in the experimental setting used in the

study (Fig. 3C; Additional file 2: Table S4).

We next interrogated if the T$_{REG}$ subsets in our data set, that showed enrichment for

molecules linked to their suppressive function, are not suppressive towards hCMV-specific

effector T cells, then which cell types are they targeting in hCMV-specific immunity. Studies

using mouse models have shown that T$_{REGs}$ can show remarkable functional plasticity and

execute their suppressive function by expressing the TF and the chemokine receptor of the

target T$_H$ subtypes [14, 59-62]. For example, the T$_{REGs}$ suppressive towards T$_H$1 will express

T-bet (encoded by *TBX21*) and CXCR3 (T$_H$1-T$_{REGs}$), while the ones toward T$_H$2 will express

GATA3 and CCR4 (T$_H$2-T$_{REGs}$), albeit at a reduced level compared to the T$_H$ subtype itself,

without necessarily themselves making the T$_H$ subtype-specific cytokines [14, 60-62].

Interestingly the differential gene expression analysis revealed that the T$_{REGs}$ in cluster 1 and

5, are characterized by the higher expression of T$_H$2-specific TF *GATA3* and chemokine

receptor *CCR4* along with *FOXP3* (Fig. 3D-F). GATA3 has been shown to be important for

FOXP3 expression and T$_{REG}$ functions [61]. Hence, we independently examined the overall

T$_H$2 subset-specific gene signatures using module enrichment score and gene set enrichment

analysis (GSEA). We found that both cluster 1 and 5 are enriched for T$_H$2 subset-specific gene

signatures (Fig. 3G and H), suggesting the T$_{REGs}$ in cluster 1 and 5 are most probably T$_H$2-like

T_{REGs}. A recent study that analysed the T cells in pan cancers has also shown the presence of

T_H2- like T_{REGs} that were marked by a higher expression of *GATA3* [63]. Clusters 1 and 5 were

also characterised by the enrichment of IFN response signature genes ($T_{REG-IFN}$) (Fig. 2B and

C). In asthma and house dust mite allergy, both T_H and T_{REG} subsets enriched for IFN response

gene sets showed dampened T_H2 response in healthy individuals with no prior history of allergy

or asthma [5]. Interestingly, in our data set, a joint enrichment analysis using the house dust

mite allergy-specific T_{REG} gene signature and T_H2-specific gene signature showed enrichment

in cluster 1 and 5 (Fig. 3I) [5]. Together these results suggest that T_{REGs} in clusters 1 and 5 are

likely T_H2-T_{REGs} with IFN response signatures that are potentially suppressive towards T_H2.

The T follicular regulatory cells (T_{FRs}) suppress T_{FH} responses and are characterized by the

expression of IL1R2 and CCR8 [3, 64, 65]. The hCMV-reactive T_{REG} cells in clusters 1 and 5

co-expressed *IL1R2* and *CCR8* transcripts along with *FOXP3* transcripts, hence, may represent

T_{FR} cells (Fig. 3D-F) [3, 64, 65]. These interesting observations warrant further investigation

to understand the relevance of T_H2-T_{REGs} and T_{FRs} in CMV immunity, especially when hCMV-

reactive conventional (non-T_{REGs}) T_H2 and T_{FH} cells were absent in our data set.

hCMV-reactive CD4-CTLs and CD8-CTLs show similar transcriptomic profile

Majority of the cells in clusters 2 and 4 and fraction of cells in clusters 0 and 7 were

characterized by the high expression of transcripts encoding cytolytic molecules such as

Granzyme B (*GZMB*), Perforin (*PRF1*), Granulysin (*GNLY*), and transcription factors *HOPX*

and *ZEB2* and showed enrichment for cytotoxicity and T_{EMRA}-specific signature genes (Fig.

1F, Fig. 2B and C, Fig. 4A and B) [3, 4, 16-20]. Interestingly, these cytotoxic clusters could

not be differentiated based on the expression of *CD4* or *CD8A* and *CD8B* transcripts, but by

other cytotoxicity related molecules (Fig. 1E and Fig. 4A-C). The cytolytic granules have been

shown to contain chemokines such as CCL3, CCL4 and CCL5 in addition to the classical

mediators of cytotoxicity (GZMB, PRF1, GNLY), in both viral and bacterial infections,

indicating their role in cytolysis [66, 67]. Cells in cluster 2 showed higher expression of the

transcripts encoding the cytokines such as *TNF* and *IFNG*, chemokines such as *CCL3*, *CCL4*,

CCL4L2, *XCL1*, *XCL2*, while cluster 4 showed higher expression of *CCL5* (Fig. 4B and C).

The chemokine receptor *CCR5*, the receptor for chemokines *CCL3*, *CCL4* and *CCL5*, and

IFNGR1, encoding the receptor for IFNG, was expressed by the cytotoxic cells in cluster 0,

suggesting these cells may be responding to the cytokines and chemokines made by the cells

in cytotoxic clusters 2 and 4 and are potentially the intermediate population (Fig. 4A and B;

Additional file2: Table S4) [68]. This is further supported by the single-cell trajectory analysis,

where on the pseudo time scale cluster 0 appears before clusters 2 and 4 (Fig. 4D). These results

suggest that the cytotoxic cells in cluster 0 are potentially an intermediate population in the

CTL lineage and these cells are mostly T_{EM} cells as they mostly lack the expression of CD45RA

and CCR7 (Fig. 1C).

Compared to cluster 4, cluster 2 cells showed higher expression of *SLAMF7*, *ZEB2*,

GZMB, *GPR18*, *CD72*, *PDCD1* (PD1), *CRTAM* that are known to be expressed in terminally

differentiated effector cells in viral infections and various cancers [3, 69-72] (Fig. 2B, Fig. 3C,

Fig. 4B and D). On the contrary, cluster 4 was characterized by the higher expression of *GNLY*,

EOMES, *ADGRG1* (GPR56), *KLRD1*, *UCP2* and *CRIP1* (Fig. 2B, Fig. 4B and D), that are

mostly not associated with terminally differentiated effector T cells. For example, CRIP1 has

been shown to be anti-apoptotic in cancers and the antigen-specific cells expressing UCP2

survive better in HIV infection through metabolic reprogramming [73, 74], suggesting the

possible role of these genes in providing survival advantage to cells in cluster 4. A coordinated

expression of CCL5 followed by GNLY and PRF1 provided host-defence against

Mycobacterium tuberculosis (Mtb) infection, where the expression of CCL5 attracted the Mtb

infected macrophages that in-turn triggered the expression of GNLY and PRF1 in CD8 T cells

[75]. Further, a delayed onset of cells expressing GNLY has been shown in multiple infections

[76, 77]. These results suggest that the cytotoxic cells could be at different stages of their development or effector function. The differential expression of many of these cytolytic molecules between clusters 2 and 4, imply that the cells in cluster 2 are potentially the terminal effector cells while those in cluster 4 are pre-effector cells. This observation is further strengthened by the TCR analysis (Fig. 2E and F, Fig. 4E and F; Additional file 1: Fig. S2B and C). The 20 most expanded clonotypes were found majorly in cluster 2 and 4 (Fig. 2F; Additional file 1: Fig. S2C). Cluster 2 is relatively more clonally expanded than cluster 4, where over 67% of the cells harbouring the 20 most expanded clonotypes were found in cluster 2 (Fig. 4F). The trajectory analysis also showed that on a pseudo-time scale, cluster 2 and 4 appears to bifurcate from a common origin of cluster 0 (Fig. 4D). Further, the pathway enrichment analysis of genes differentially expressed between clusters 2 and 4 revealed an enrichment for cytokine and chemokine response pathways in cluster 4, while cluster 2 showed enrichment of pathways associated with pathogen induced cytokine storm (Additional file 1: Fig. S3A and B). The effector T cells utilize glycolysis, as the preferred metabolic pathway that play an important role for cytokine secretion and lytic nature of these effector cells [78-80]. The cells in cluster 2 showed enrichment for many metabolic pathways associated with effector cells such as glycolysis and HIF1α signalling as well as pathways related to anti-viral responses (Additional file 1: Fig. S3A). Together, these combined analysis of the transcriptome, TCR clonotype expansion and sharing along with the trajectory analysis provides the evidence to support the idea that cluster 4 cells are pre-effectors while cluster 2 cells are effectors.

Next to assess if the observed differences between the CTL clusters 2 and 4 are influenced by the proportion and the characteristics of CD4-CTLs and CD8-CTLs, we compared the expression of these transcripts between CD4 and CD8 T cells within clusters 2 and 4. The proportions of CD4 and CD8 T cells within clusters 2 and 4 were comparable (Fig.

4G), indicating that the disproportionate distribution of CD4$^+$ and CD8$^+$ T cells is not the cause of the observed differences between clusters 2 and 4. Overall, the transcriptome profile of CD4, CD8 or cells that could not be assigned as CD4 or CD8 T cells (NA) within each cluster did not vary except for *CD8A* and *CD8B* transcripts (Fig. 4H and I). Transcripts such as *PRF1* and *NKG7* that showed comparable expression between cluster 2 and 4, followed the same pattern even when these two clusters were further sub-divided based on the CD4 and CD8 T cell sub-type. The transcripts that showed higher expression in cluster 2 (*XCL1*, *ZEB2*, *CRTAM* and *GZMB*) and cluster 4 (*CCL5*, *GNLY*, *EOMES*), followed similar pattern of expression in CD4 and CD8 T cells within each of these effector and pre-effector clusters (Fig. 4H and I). Both CD4 and CD8 T cells from cluster 4 showed higher expression of IFN response genes *ISG20* and *IFI6* while those in cluster 2 showed higher expression of cytokines *TNF* and *IFNG* (Fig. 4H and I). We observed that the cells in clusters 2 and 4 have distinct gene expression pattern that is not influenced by either CD4 or CD8 T cell proportion (Fig. 4H and I). Overall, our data has revealed that both hCMV-reactive CD4-CTLs and CD8-CTLs share similar transcriptome profile and can be classified as effectors and pre-effectors of CTLs.

Discussion

A significant world population carries the latent hCMV infection, without showing any symptoms or causing the disease. However, latent infection poses a huge risk of reactivation in immune-compromised individuals such as those undergoing transplant or infants born to hCMV seropositive mothers. Although there is very little risk of the disease occurrence in immune-competent individuals, the latent CMV infection puts a huge burden on the immune system, especially for the elderly, where significant proportions of the T cells are found to be hCMV-specific [46]. Identifying and understanding the characteristics of the hCMV-specific T cell immune memory components in individuals carrying the virus in latent form,

representing protective immune response, can be of high value to design therapy or vaccines

to control CMV infection upon reactivation or maternal to fetal transfer.

Our understanding of *ex vivo*, antigen-specific human T cell responses to pathogens

such as hCMV remains limited, due to the rarity of the cells, the complexity of the biology,

and technical challenges of identifying overall antigen-specific CD4$^+$ memory T cells, since

they can be extremely heterogeneous in the expression of activation associated markers and

cytokines. An added layer of complexity comes from HLA-restriction of individuals and the

prior knowledge about the immune-dominant peptide, limiting the use of pMHC multimers.

Most studies have focused on isolating the total CD4$^+$ or CD8$^+$ T cells reactive to hCMV,

without considering the differences in memory proportions that can inadvertently bias the

analysis towards analysing the predominant memory compartment within CD4$^+$ T cells (T$_{CM}$

or T$_{EM}$) and CD8$^+$ T cells (T$_{EMRA}$). Our strategy to isolate antigen-specific T cells based on the

combination of TCR activation dependent markers (CD154 and CD137) across different

memory subsets followed by high resolution single-cell RNA-Seq paired with single-cell TCR

analysis allowed us to interrogate and compare a larger pool of hCMV (pp65 and IE-1)-reactive

cells from different memory subsets across 10 seropositive and peptide-responsive donors. This

unbiased approach enabled us to overlay the heterogeneity in T cell memory subsets classified

based on developmental stages and functional subsets along with their TCR dependent

activation markers. The expression of CD154 and CD137 in different set of cells that further

differentially co-expressed one or more T cell activation associated markers such as CD69,

CD25, OX40, warrants for careful selection of activation markers while analysing antigen-

specific cells *ex vivo*. In this study, we observed that though activated (based on the expression

of *CD154*, *CD137*, *CD69*, *CD25*, *OX40*), most cells, with the exception of effector cells, did

not actively make cytokines and those that made also showed heterogeneity for the cytokine

profile. Many studies previously have also relied on cytokine profiles to analyse *ex vivo* isolated

antigen-specific cells and hence have mostly analysed effector memory T cells.

The single-cell RNA-Seq of a large number of hCMV reactive cells isolated from 10

different donors, identified several interesting functional subsets; T_H17, T_{REGs}, CD4-CTLs and

CD8-CTLs. Of particular interest were CD4- and CD8-CTLs, that clustered together. The two

cytotoxic clusters observed could not be distinguished based on the CD4- and CD8-T cell

specific transcripts *CD4* or *CD8A*/*CD8B*, but on the basis of other cytotoxic molecules that

were linked to different states of effector functions [72]. The cluster that was predominantly

making cytokine and chemokine (*IFNG*, *TNF*, *XCL1*, *XCL2*, *CCL3*, *CCL4*) also expressed

transcription factor (TF) *ZEB2*, the TF known to be associated with late effector function [4,

72]. The cytotoxic cells in both clusters 4 and 0 appeared to respond to the cytokines and

chemokines by expressing their receptors (*IFNGR1*, *CCR5*). However, the cytotoxic cells

across these clusters expressed molecules linked to cytolytic granules (*PRF1*, *GZMB*, *GNLY*),

and transcription factor *HOPX*. The differential expression of many of the molecules linked to

cytotoxicity and effector functions, suggests the effector and pre-effector like subsets. These

subsets, possibly at different developmental states, might exist to ensure a continuity in immune

response during the entire course of the infection.

Another interesting discovery in our data set is the identification of T_{REGs} with

suppressive ability that showed enrichment for the IFN response gene signatures ($T_{REG-IFN}$)

which has been shown to be associated with a dampened T_H2 response in allergy [5]. The

enrichment of T_H2 signatures as well as the expression of *GATA3* and *CCR4*, the T_H2 associated

transcription factor and chemokine receptor by the T_{REGs} further suggested that these could be

suppressive towards the T_H2 subset. Although a very recent study in pan cancer has also

identified this subset, to the best of our knowledge, presence of T_{REGs} enriched for IFN response

genes has not been reported earlier in viral infections in humans [63]. Based on these

observations, we hypothesize that during the initial stages of the infection where the expansion

of effector cells is necessary, T_{REGs} may not be executing their suppressive function towards

these effectors, instead, suppressing other subtypes of no relevance to the disease. However,

further kinetic studies in both animal models and longitudinal human cohorts of different types

of diseases such as viral and bacterial infections, allergy and cancers are necessary to verify or

refute this hypothesis.

Conclusion

In conclusion, using the single-cell multi-omics approach (transcriptome, TCR repertoire and

expression of a few selected proteins), our study has identified an unprecedented amount of

heterogeneity in hCMV-specific CD4$^+$ memory T cell subsets. The study identified novel

subsets such as $T_{REG\text{-}IFN}$ and pre-effectors and effectors of both CD4-CTLs and CD8-CTLs.

The knowledge base can serve as an important resource for designing vaccines and therapeutic

strategies to control CMV infections, especially in immune-compromised individuals and

infants of seropositive mothers.

Materials and Methods

Study subjects

Study subjects were healthy adult donors who donated blood at the Safdarjung Blood Bank

during 2019-20. The donors consented for use of the plasma and buffy coat samples for

research. Donors were HIV-negative and had no history of Hepatitis C infection. The median

age was 30.5 (24-39) and all these donors were male. Approval for the use of this material for

research was obtained from the institutional human ethics committees from both National

Institute of Immunology, New Delhi and Vardhman Mahavir Medical College (VMMC) and

Safdarjung Hospital, New Delhi, India.

PBMC processing

Peripheral Blood Mononuclear Cells (PBMCs) were isolated from buffy coat samples from healthy blood donors obtained from Safdarjung hospital blood bank by density gradient centrifugation using Ficoll-Paque Premium (GE Healthcare Biosciences). PBMCs were cryopreserved in 90% Fetal Bovine Serum (FBS) supplemented with 10% di-methyl sulfoxide (DMSO).

CMV IgG analysis

ELISA (Enzyme Linked Immunosorbent Assay) was performed on the plasma samples from the donors obtained from the blood bank using Human Anti-Cytomegalovirus IgG ELISA kit (abcam, #ab108724) according to the manufacturer's protocol to check for the donors' sero-status. Donors with the absorbance values at least 10% higher than that of the cut-off control supplied in the kit, were considered positive. Based on this criteria we found 124 out of 126 donors in our cohort were hCMV IgG+; 98.4% positivity) (Additional file 1: Fig. S1A).

PBMC stimulation

To analyse or isolate antigen-specific T cells, the ARTE assay was employed with few minor modifications [40, 41]. The hCMV-specific PepTivators (Miltenyi Biotec) derived from pp65 (UL83) and IE-1 (UL123) were used to stimulate PBMCs from CMV-seropositive donors, according to the manufacturer's protocol. Briefly, PBMCs were thawed and treated with 50 U/mL Benzonase (Sigma) in RPMI before resting overnight in serum-free TexMACS medium (Miltenyi Biotec) supplemented with 1% Penicillin/Streptomycin at 37°C in a 96-well plate. Cells were stimulated for 6 or 24hrs at a density of 1 million/100 µL/well of 96 well plate, by the addition of hCMV peptide pools (pp65 and IE-1 PepTivators) at a final concentration of 1 µg/mL in the presence of a blocking CD40 antibody (1 µg/mL; Miltenyi Biotec) and CD28 antibody (1 µg/mL; eBioscience) [1, 40, 41]. Stimulation with αCD3/CD28 DynaBeads Human T cell activator (Invitrogen) (cell to bead ratio of 1:1) served as positive control while with

water (vehicle control) served as unstimulated negative control. The stimulated cells were

washed and stained with cocktail of fluorescent antibodies (Table S2A) and analysed by flow

cytometry or processed for single-cell RNA-sequencing using 10X genomics as follows. The

hCMV-peptide pool stimulated cells from each donor were washed and individually stained

with Cell-hashtag (HTO:Hash Tag Oligo) TotalSeq C antibody (0.5 µg/condition; Biolegend,

Table S2B) as per manufacturer's recommendations for 30 minutes on ice, and washed twice

with MACS buffer (PBS + 2% FBS + 2mM EDTA). HTO stained PBMCs from all 10 donors

were then pooled and stained with oligo-tagged TotalSeq C antibody for cell surface proteins

for T cell memory markers (CD45RA, CCR7, CD95 and IL7Rα) (0.5 µg/condition; Biolegend,

Table S2B) for 30 minutes on ice. Following washing, the pooled cells were further stained

with fluorescence-conjugated surface antibodies to enable FACS sorting of antigen-specific

cells (Table S2A). Cells were directly FACS sorted without an intermediate MACS column

enrichment step using FACSAria Fusion Cell Sorter (Becton Dickinson) in LoBind 1.5 mL

microcentrifuge (Eppendorf) tubes with 1:1 FBS:PBS supplemented with recombinant RNase

inhibitor (1:100, Takara). The live singlet gated CD3+ T cells were further gated as per the

gating strategy shown in additional file 1: Fig. S1B. List of antibodies used are provided in

additional file 2: Table S2. The flowcytometry data was analysed using FlowJo software

(v10.8).

Single-cell RNA-Seq

The sorted antigen-specific cells from various memory compartments were pooled in desired

ratios and used for single-cell RNA-Seq and TCR-Seq assay using 10X Genomics kit, 5′ V(D)J

with gene expression and cell surface protein expression v1.1 as per the manufacturer's

recommendations. Initial amplification of cDNA libraries and final libraries were performed

for 15 cycles and 14 cycles respectively for gene expression library. V(D)J and cell surface

protein libraries were generated corresponding to each 5′ gene expression library using 9 cycles

of amplification. Libraries were sequenced on Illumina NovaSeq6000 sequencing platform for

paired end (PE)150 reads.

Single-cell RNA-Seq analysis

Sequenced reads from the 5′ single-cell gene expression and associated V(D)J library were

mapped to the GRCh38 reference genome using 10X Genomics' cellranger multi (v6.1.1). The

cell surface protein quantification was run as part of the same pipeline. The Seurat package

(v4.3) in R (v4.2.2) was used to perform all further downstream analyses [81].

For downstream analysis, only cells that passed the QCs were retained; cells with UMI

(Unique Molecular Identifier) counts >500, number of genes >350 and percent mitochondrial

genes <15% were considered as good quality cells. Further, cells were assigned as doublets,

singlets or negatives based on the HTO counts using Demux with default parameters in Seurat

package (v4.3) [82]. Singlets were assigned to their respective donor based on HTO count.

Doublets were removed from downstream analysis. After applying all the quality control

parameters, we retained 6899 cells out of 8062 cells for downstream analysis (85.57% cells

passed the quality parameters). Cells from experiment 1 (3 donors) and experiment 2 (10

donors) were combined using the Seurat package after applying the QCs (Table S1).

For each cell, normalized UMI counts for each gene were obtained by dividing the raw

counts by the total counts across all genes. This was multiplied by a scaling factor of 10,000

and log transformed to yield the final normalized values. Prior to clustering, we filtered genes

by selecting for the top 2000 with highest variance across cells, using *FindVariableFeatures*

with *selection.method="vst"*. This data was then centered and scaled using *ScaleData*, and the

top 50 principal components (PCs) were calculated. Top 18 PCs were selected using *ElbowPlot*

and *JackStrawPlot* for downstream processing. This was used to build a shared nearest

neighbour graph using *FindNeighbors*. The cells were clustered using *FindClusters*, with

algorithm=3 and *resolution=0.2*, determined using the *clustree* function. For visualization and

clustering, a 2D UMAP projection of the top 18 PCs was obtained using the *RunUMAP*

function, with *metric="euclidean"* [81].

The cell surface protein tag data was normalized using the *NormalizeData* function in

Seurat with *normalization.method="CLR"*. Briefly, for each cell, the hashtag counts were

divided by the geometric mean of counts of all unique hashtags prior to log transformation.

Differential gene expression analysis

We identified differentially expressed genes using the MAST algorithm integrated within

Seurat's *FindAllMarkers* function using default parameters on $\log_2(\text{CPM}+1)$ counts [81].

Genes in a cluster were reported as differentially expressed (DE) compared to every other

cluster if their $\log_2$ fold change of expression was > 0.25 and BH-adjusted P-value was < 0.05.

Gene module signature scores and Gene Set Enrichment Analysis (GSEA)

Gene module scores were computed using the *AddModuleScore* function in Seurat, with default

parameters [81]. Briefly, for each cell, the module score is defined by the mean of the signature

gene list (test set) after subtracting the mean expression of an aggregate of control gene lists.

Control gene lists (same number as test gene list) are sampled from bins created based on the

level of expression of the signature gene list. GSEA was performed using Qlucore Omics

Explorer 3.8 software package [83]. List of the genes used are provided in supplementary table

S3.

Trajectory inference

Single cell trajectories were computed using the slingshot algorithm within the dynverse/dyno

(v0.1.2) suite of trajectory inference pipelines [84]. Raw and normalized UMI counts from cells

of clusters 0, 2, 3, 4 and 7 were combined using the dynverse's wrap_expressions function.

Cells were assigned groups by cluster and the number of expected start and end states were set

to 1 using *add_prior_information*. Trajectory inference was run using *infer_trajectories* with

default parameters, with method = ti_slingshot(). We did not specify any start or end cells for

the trajectory.

Pathway enrichment analysis

DE genes between clusters 2 and 4 were fed to Ingenuity Pathway Analysis software from

Qiagen. Enrichment pathways were sourced from Ingenuity core enrichment pathways.

TCR analysis

For each cell, all information about clonality, expressed TCR chains, their sequences and

corresponding UMI counts were obtained as part of the standard 10X Genomics' cellranger

multi pipeline (Table S5).

Assigning CD4 and CD8 T cells

For cells in cluster 2 and 4, we assigned cells as $CD4^+$ T cells, or $CD8^+$ T cells, first based on

the *CD4* and *CD8A*/*CD8B* transcript counts. Cells with > 0 counts for *CD4*, were assigned as

$CD4^+$ T cell (333 cells). For $CD8^+$ T cells, we considered the cell as $CD8^+$ T cell (735 cells) if

the sum of all three CD8 transcripts (*CD8A*, *CD8B*, *CD8B2*) was > 0 and *CD4* transcripts were

undetectable. Cells (556) which could not be assigned as either CD4 or CD8 T cell based on

this criterion, were further assigned as $CD4^+$ (274) and $CD8^+$ (12) T cells if they shared the

clonotypes with an already assigned $CD4^+$ or $CD8^+$ T cell. Cells that could not be assigned to

either CD4 or CD8 category based on either of these criteria, were called as NA (not assigned)

cells (294). Based on these criteria, we had a total of 328 $CD4^+$ T cells, 536 $CD8^+$ T cells and

149 NA cells in cluster 2 (total 1013 cells) and 279 $CD4^+$ T cells, 199 $CD8^+$ T cells and 145

NA cells in cluster 4 (total 623 cells).

Statistical tests

All the statistical tests were done using GraphPad Prism v9.0. Mann Whitney u-test was used

to perform pairwise comparison and *P*-value <0.05 was considered significant.

Data and Code availability

Scripts will be made available on GitHub (https://github.com/ImmunogenomicsLab-NII/CMV-HD). Sequencing data for this study is deposited onto the Gene Expression Omnibus with accession number GSE235604.

References

1. Kusnadi A, Ramirez-Suastegui C, Fajardo V, Chee SJ, Meckiff BJ, Simon H, Pelosi E, Seumois G, Ay F, Vijayanand P, Ottensmeier CH: **Severely ill COVID-19 patients display impaired exhaustion features in SARS-CoV-2-reactive CD8(+) T cells.** *Sci Immunol* 2021, **6**.

2. Lim EY, Jackson SE, Wills MR: **The CD4+ T Cell Response to Human Cytomegalovirus in Healthy and Immunocompromised People.** *Front Cell Infect Microbiol* 2020, **10:**202.

3. Meckiff BJ, Ramirez-Suastegui C, Fajardo V, Chee SJ, Kusnadi A, Simon H, Eschweiler S, Grifoni A, Pelosi E, Weiskopf D, et al: **Imbalance of Regulatory and Cytotoxic SARS-CoV-2-Reactive CD4(+) T Cells in COVID-19.** *Cell* 2020, **183:**1340-1353 e1316.

4. Patil VS, Madrigal A, Schmiedel BJ, Clarke J, O'Rourke P, de Silva AD, Harris E, Peters B, Seumois G, Weiskopf D, et al: **Precursors of human CD4(+) cytotoxic T lymphocytes identified by single-cell transcriptome analysis.** *Sci Immunol* 2018, **3**.

5. Seumois G, Ramirez-Suastegui C, Schmiedel BJ, Liang S, Peters B, Sette A, Vijayanand P: **Single-cell transcriptomic analysis of allergen-specific T cells in allergy and asthma.** *Sci Immunol* 2020, **5**.

6. Lyu M, Wang S, Gao K, Wang L, Zhu X, Liu Y, Wang M, Liu X, Li B, Tian L: **Dissecting the Landscape of Activated CMV-Stimulated CD4+ T Cells in**

Humans by Linking Single-Cell RNA-Seq With T-Cell Receptor Sequencing.
Front Immunol 2021, **12:**779961.

7. Lewis SA, Sutherland A, Soldevila F, Westernberg L, Aoki M, Frazier A, Maiche S, Erlewyn-Lajeunesse M, Arshad H, Leonard S, et al: **Identification of cow milk epitopes to characterize and quantify disease-specific T cells in allergic children.** *J Allergy Clin Immunol* 2023.

8. O'Shea JJ, Paul WE: **Mechanisms underlying lineage commitment and plasticity of helper CD4+ T cells.** *Science* 2010, **327:**1098-1102.

9. Schmiedel BJ, Singh D, Madrigal A, Valdovino-Gonzalez AG, White BM, Zapardiel-Gonzalo J, Ha B, Altay G, Greenbaum JA, McVicker G, et al: **Impact of Genetic Polymorphisms on Human Immune Cell Gene Expression.** *Cell* 2018, **175:**1701-1715 e1716.

10. Arlehamn CL, Seumois G, Gerasimova A, Huang C, Fu Z, Yue X, Sette A, Vijayanand P, Peters B: **Transcriptional profile of tuberculosis antigen-specific T cells reveals novel multifunctional features.** *J Immunol* 2014, **193:**2931-2940.

11. Crotty S: **T Follicular Helper Cell Biology: A Decade of Discovery and Diseases.** *Immunity* 2019, **50:**1132-1148.

12. Sallusto F: **Heterogeneity of Human CD4(+) T Cells Against Microbes.** *Annu Rev Immunol* 2016, **34:**317-334.

13. Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M: **Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases.** *J Immunol* 1995, **155:**1151-1164.

14. Scheinecker C, Goschl L, Bonelli M: **Treg cells in health and autoimmune diseases: New insights from single cell analysis.** *J Autoimmun* 2020, **110:**102376.

667   15.   Mahnke YD, Brodie TM, Sallusto F, Roederer M, Lugli E: **The who's who of T-cell**

668   **differentiation: human memory T-cell subsets.** *Eur J Immunol* 2013, **43:**2797-

669   2809.

670   16.   Jackson SE, Sedikides GX, Mason GM, Okecha G, Wills MR: **Human**

671   **Cytomegalovirus (HCMV)-Specific CD4(+) T Cells Are Polyfunctional and Can**

672   **Respond to HCMV-Infected Dendritic Cells In Vitro.** *J Virol* 2017, **91.**

673   17.   van Leeuwen EM, Remmerswaal EB, Heemskerk MH, ten Berge IJ, van Lier RA:

674   **Strong selection of virus-specific cytotoxic CD4+ T-cell clones during primary**

675   **human cytomegalovirus infection.** *Blood* 2006, **108:**3121-3127.

676   18.   van Leeuwen EM, Remmerswaal EB, Vossen MT, Rowshani AT, Wertheim-van

677   Dillen PM, van Lier RA, ten Berge IJ: **Emergence of a CD4+CD28- granzyme B+,**

678   **cytomegalovirus-specific T cell subset after recovery of primary cytomegalovirus**

679   **infection.** *J Immunol* 2004, **173:**1834-1841.

680   19.   Gamadia LE, Rentenaar RJ, van Lier RA, ten Berge IJ: **Properties of CD4(+) T cells**

681   **in human cytomegalovirus infection.** *Hum Immunol* 2004, **65:**486-492.

682   20.   Pachnio A, Ciaurriz M, Begum J, Lal N, Zuo J, Beggs A, Moss P: **Cytomegalovirus**

683   **Infection Leads to Development of High Frequencies of Cytotoxic Virus-Specific**

684   **CD4+ T Cells Targeted to Vascular Endothelium.** *PLoS Pathog* 2016,

685   **12:**e1005832.

686   21.   Cheroutre H, Husain MM: **CD4 CTL: living up to the challenge.** *Semin Immunol*

687   2013, **25:**273-281.

688   22.   Weiskopf D, Bangs DJ, Sidney J, Kolla RV, De Silva AD, de Silva AM, Crotty S,

689   Peters B, Sette A: **Dengue virus infection elicits highly polarized CX3CR1+**

690   **cytotoxic CD4+ T cells associated with protective immunity.** *Proc Natl Acad Sci U*

691   *S A* 2015, **112:**E4256-4263.

23.   Gandhi MK, Khanna R: **Human cytomegalovirus: clinical aspects, immune**

regulation, and emerging treatments. *Lancet Infect Dis* 2004, **4:**725-738.

24.   Tu W, Chen S, Sharp M, Dekker C, Manganello AM, Tongson EC, Maecker HT,

Holmes TH, Wang Z, Kemble G, et al: **Persistent and selective deficiency of CD4+**

T cell immunity to cytomegalovirus in immunocompetent young children. *J*

Immunol 2004, **172:**3260-3267.

25.   Gratama JW, Brooimans RA, van der Holt B, Sintnicolaas K, van Doornum G,

Niesters HG, Lowenberg B, Cornelissen JJ: **Monitoring cytomegalovirus IE-1 and**

pp65-specific CD4+ and CD8+ T-cell responses after allogeneic stem cell

transplantation may identify patients at risk for recurrent CMV reactivations.

Cytometry B Clin Cytom 2008, **74:**211-220.

26.   Einsele H, Roosnek E, Rufer N, Sinzger C, Riegler S, Loffler J, Grigoleit U, Moris A,

Rammensee HG, Kanz L, et al: **Infusion of cytomegalovirus (CMV)-specific T cells**

for the treatment of CMV infection not responding to antiviral chemotherapy.

Blood 2002, **99:**3916-3922.

27.   Sester M, Sester U, Gartner B, Heine G, Girndt M, Mueller-Lantzsch N, Meyerhans

A, Kohler H: **Levels of virus-specific CD4 T cells correlate with cytomegalovirus**

control and predict virus-induced disease after renal transplantation.

Transplantation 2001, **71:**1287-1294.

28.   Peggs KS, Verfuerth S, Pizzey A, Khan N, Guiver M, Moss PA, Mackinnon S:

Adoptive cellular therapy for early cytomegalovirus infection after allogeneic

stem-cell transplantation with virus-specific T-cell lines. *Lancet* 2003, **362:**1375-

1377.

29.   Lilleri D, Fornara C, Chiesa A, Caldera D, Alessandrino EP, Gerna G: **Human**

cytomegalovirus-specific CD4+ and CD8+ T-cell reconstitution in adult

allogeneic hematopoietic stem cell transplant recipients and immune control of viral infection. *Haematologica* 2008, **93:**248-256.

30. Gamadia LE, Remmerswaal EB, Weel JF, Bemelman F, van Lier RA, Ten Berge IJ: **Primary immune responses to human CMV: a critical role for IFN-gamma-producing CD4+ T cells in protection against CMV disease.** *Blood* 2003, **101:**2686-2692.

31. Lilleri D, Fornara C, Revello MG, Gerna G: **Human cytomegalovirus-specific memory CD8+ and CD4+ T cell differentiation after primary infection.** *J Infect Dis* 2008, **198:**536-543.

32. Jackson SE, Mason GM, Wills MR: **Human cytomegalovirus immunity and immune evasion.** *Virus Res* 2011, **157:**151-160.

33. Khan N, Cobbold M, Keenan R, Moss PA: **Comparative analysis of CD8+ T cell responses against human cytomegalovirus proteins pp65 and immediate early 1 shows similarities in precursor frequency, oligoclonality, and phenotype.** *J Infect Dis* 2002, **185:**1025-1034.

34. Sylwester AW, Mitchell BL, Edgar JB, Taormina C, Pelte C, Ruchti F, Sleath PR, Grabstein KH, Hosken NA, Kern F, et al: **Broadly targeted human cytomegalovirus-specific CD4+ and CD8+ T cells dominate the memory compartments of exposed subjects.** *J Exp Med* 2005, **202:**673-685.

35. Jackson SE, Sedikides GX, Okecha G, Wills MR: **Generation, maintenance and tissue distribution of T cell responses to human cytomegalovirus in lytic and latent infection.** *Med Microbiol Immunol* 2019, **208:**375-389.

36. Terrazzini N, Bajwa M, Vita S, Cheek E, Thomas D, Seddiki N, Smith H, Kern F: **A novel cytomegalovirus-induced regulatory-type T-cell subset increases in size**

during older life and links virus-specific immunity to vascular pathology. *J Infect Dis* 2014, **209:**1382-1392.

37. Tovar-Salazar A, Patterson-Bartlett J, Jesser R, Weinberg A: **Regulatory function of cytomegalovirus-specific CD4+CD27-CD28- T cells.** *Virology* 2010, **398:**158-167.

38. Schwele S, Fischer AM, Brestrich G, Wlodarski MW, Wagner L, Schmueck M, Roemhild A, Thomas S, Hammer MH, Babel N, et al: **Cytomegalovirus-specific regulatory and effector T cells share TCR clonality--possible relation to repetitive CMV infections.** *Am J Transplant* 2012, **12:**669-681.

39. Clement M, Marsden M, Stacey MA, Abdul-Karim J, Gimeno Brias S, Costa Bento D, Scurr MJ, Ghazal P, Weaver CT, Carlesso G, et al: **Cytomegalovirus-Specific IL-10-Producing CD4+ T Cells Are Governed by Type-I IFN-Induced IL-27 and Promote Virus Persistence.** *PLoS Pathog* 2016, **12:**e1006050.

40. Bacher P, Scheffold A: **Flow-cytometric analysis of rare antigen-specific T cells.** *Cytometry A* 2013, **83:**692-701.

41. Bacher P, Schink C, Teutschbein J, Kniemeyer O, Assenmacher M, Brakhage AA, Scheffold A: **Antigen-reactive T cell enrichment for direct, high-resolution analysis of the human naive and memory Th cell repertoire.** *J Immunol* 2013, **190:**3967-3976.

42. Poloni C, Schonhofer C, Ivison S, Levings MK, Steiner TS, Cook L: **T-cell activation-induced marker assays in health and disease.** *Immunol Cell Biol* 2023.

43. Best JA, Blair DA, Knell J, Yang E, Mayya V, Doedens A, Dustin ML, Goldrath AW, Immunological Genome Project C: **Transcriptional insights into the CD8(+) T cell response to infection and memory T cell formation.** *Nat Immunol* 2013, **14:**404-412.

765   44.   Bahr F, Wehner R, Platzbecker U, Wermke M, Shayegi N, Middeke JM, Rollig C,

766   Schetelig J, Ehninger G, Schmitz M, et al: **Reconstitution of interleukin-17-**

767   **producing T helper cells after allogeneic hematopoietic cell transplantation.** *Biol*

768   *Blood Marrow Transplant* 2013, **19:**357-365.

769   45.   Schober K, Voit F, Grassmann S, Muller TR, Eggert J, Jarosch S, Weissbrich B,

770   Hoffmann P, Borkner L, Nio E, et al: **Reverse TCR repertoire evolution toward**

771   **dominant low-affinity clones during chronic CMV infection.** *Nat Immunol* 2020,

772   **21:**434-441.

773   46.   Klenerman P, Oxenius A: **T cell responses to cytomegalovirus.** *Nat Rev Immunol*

774   2016, **16:**367-377.

775   47.   Seddiki N, Santner-Nanan B, Martinson J, Zaunders J, Sasson S, Landay A, Solomon

776   M, Selby W, Alexander SI, Nanan R, et al: **Expression of interleukin (IL)-2 and IL-**

777   **7 receptors discriminates between human regulatory and activated T cells.** *J Exp*

778   *Med* 2006, **203:**1693-1700.

779   48.   Tovar-Salazar A, Weinberg A: **Understanding the mechanism of action of**

780   **cytomegalovirus-induced regulatory T cells.** *Virology* 2020, **547:**1-6.

781   49.   Velaga S, Ukena SN, Hopting M, Ivanyi P, Borchers S, Mischak-Weissinger EM,

782   Hamwi I, Buchholz S, Ganser A, Franzke A: **Reconstitution and phenotype of**

783   **Tregs in CMV reactivating patients following allogeneic hematopoietic stem cell**

784   **transplantation.** *Immunol Invest* 2013, **42:**18-35.

785   50.   Almanan M, Raynor J, Sholl A, Wang M, Chougnet C, Cardin RD, Hildeman DA:

786   **Tissue-specific control of latent CMV reactivation by regulatory T cells.** *PLoS*

787   *Pathog* 2017, **13:**e1006507.

788   51.   Miyara M, Yoshioka Y, Kitoh A, Shima T, Wing K, Niwa A, Parizot C, Taflin C,

789   Heike T, Valeyre D, et al: **Functional delineation and differentiation dynamics of**

human CD4+ T cells expressing the FoxP3 transcription factor. *Immunity* 2009,

30:899-911.

52. Valmori D, Raffin C, Raimbaud I, Ayyoub M: **Human RORgammat+ TH17 cells**

preferentially differentiate from naive FOXP3+Treg in the presence of lineage-

specific polarizing factors. *Proc Natl Acad Sci U S A* 2010, **107:**19402-19407.

53. Matsuki F, Saegusa J, Miyamoto Y, Misaki K, Kumagai S, Morinobu A: **CD45RA-**

Foxp3(high) activated/effector regulatory T cells in the CCR7 + CD45RA-CD27

+ CD28+central memory subset are decreased in peripheral blood from patients

with rheumatoid arthritis. *Biochem Biophys Res Commun* 2013, **438:**778-783.

54. Nowak A, Lock D, Bacher P, Hohnstein T, Vogt K, Gottfreund J, Giehr P, Polansky

JK, Sawitzki B, Kaiser A, et al: **CD137+CD154- Expression As a Regulatory T**

Cell (Treg)-Specific Activation Signature for Identification and Sorting of Stable

Human Tregs from In Vitro Expansion Cultures. *Front Immunol* 2018, **9:**199.

55. Jayaratne HE, Wijeratne D, Fernando S, Kamaladasa A, Gomes L, Wijewickrama A,

Ogg GS, Malavige GN: **Regulatory T-cells in acute dengue viral infection.**

Immunology 2018, **154:**89-97.

56. Deaglio S, Dwyer KM, Gao W, Friedman D, Usheva A, Erat A, Chen JF, Enjyoji K,

Linden J, Oukka M, et al: **Adenosine generation catalyzed by CD39 and CD73**

expressed on regulatory T cells mediates immune suppression. *J Exp Med* 2007,

204:1257-1265.

57. Raffin C, Pignon P, Celse C, Debien E, Valmori D, Ayyoub M: **Human memory**

Helios- FOXP3+ regulatory T cells (Tregs) encompass induced Tregs that

express Aiolos and respond to IL-1beta by downregulating their suppressor

functions. *J Immunol* 2013, **191:**4619-4627.

58.   Shevach EM: **Mechanisms of foxp3+ T regulatory cell-mediated suppression.**

Immunity 2009, **30:**636-645.

59.   Sefik E, Geva-Zatorsky N, Oh S, Konnikova L, Zemmour D, McGuire AM, Burzyn

D, Ortiz-Lopez A, Lobera M, Yang J, et al: **MUCOSAL IMMUNOLOGY.**

Individual intestinal symbionts induce a distinct population of RORgamma(+)

regulatory T cells. *Science* 2015, **349:**993-997.

60.   Koch MA, Tucker-Heard G, Perdue NR, Killebrew JR, Urdahl KB, Campbell DJ:

The transcription factor T-bet controls regulatory T cell homeostasis and

function during type 1 inflammation. *Nat Immunol* 2009, **10:**595-602.

61.   Wang Y, Su MA, Wan YY: **An essential role of the transcription factor GATA-3**

for the function of regulatory T cells. *Immunity* 2011, **35:**337-348.

62.   Zheng Y, Chaudhry A, Kas A, deRoos P, Kim JM, Chu TT, Corcoran L, Treuting P,

Klein U, Rudensky AY: **Regulatory T-cell suppressor program co-opts**

transcription factor IRF4 to control T(H)2 responses. *Nature* 2009, **458:**351-356.

63.   Chu Y, Dai E, Li Y, Han G, Pei G, Ingram DR, Thakkar K, Qin JJ, Dang M, Le X, et

al: **Pan-cancer T cell atlas links a cellular stress response state to immunotherapy**

resistance. *Nat Med* 2023.

64.   Eschweiler S, Clarke J, Ramirez-Suastegui C, Panwar B, Madrigal A, Chee SJ,

Karydis I, Woo E, Alzetani A, Elsheikh S, et al: **Intratumoral follicular regulatory**

T cells curtail anti-PD-1 treatment efficacy. *Nat Immunol* 2021, **22:**1052-1063.

65.   Xie MM, Fang S, Chen Q, Liu H, Wan J, Dent AL: **Follicular regulatory T cells**

inhibit the development of granzyme B-expressing follicular helper T cells. *JCI*

Insight 2019, **4**.

66.   Wagner L, Yang OO, Garcia-Zepeda EA, Ge Y, Kalams SA, Walker BD, Pasternack

MS, Luster AD: **Beta-chemokines are released from HIV-1-specific cytolytic T-**

cell granules complexed to proteoglycans. *Nature* 1998, **391:**908-911.

67.   Caccamo N, Milano S, Di Sano C, Cigna D, Ivanyi J, Krensky AM, Dieli F, Salerno

A: **Identification of epitopes of Mycobacterium tuberculosis 16-kDa protein**

recognized by human leukocyte antigen-A*0201 CD8(+) T lymphocytes. *J Infect*

Dis 2002, **186:**991-998.

68.   Eberlein J, Davenport B, Nguyen TT, Victorino F, Jhun K, van der Heide V,

Kuleshov M, Ma'ayan A, Kedl R, Homann D: **Chemokine Signatures of Pathogen-**

Specific T Cells I: Effector T Cells. *J Immunol* 2020, **205:**2169-2187.

69.   Awwad MHS, Mahmoud A, Bruns H, Echchannaoui H, Kriegsmann K, Lutz R, Raab

MS, Bertsch U, Munder M, Jauch A, et al: **Selective elimination of**

immunosuppressive T cells in patients with multiple myeloma. *Leukemia* 2021,

35:2602-2615.

70.   Sumida H, Cyster JG: **G-Protein Coupled Receptor 18 Contributes to**

Establishment of the CD8 Effector T Cell Compartment. *Front Immunol* 2018,

9:660.

71.   Correa-Rocha R, Lopez-Abente J, Gutierrez C, Perez-Fernandez VA, Prieto-Sanchez

A, Moreno-Guillen S, Munoz-Fernandez MA, Pion M: **CD72/CD100 and PD-1/PD-**

L1 markers are increased on T and B cells in HIV-1+ viremic individuals, and

CD72/CD100 axis is correlated with T-cell exhaustion. *PLoS One* 2018,

13:e0203419.

72.   Omilusik KD, Best JA, Yu B, Goossens S, Weidemann A, Nguyen JV, Seuntjens E,

Stryjewska A, Zweier C, Roychoudhuri R, et al: **Transcriptional repressor ZEB2**

promotes terminal differentiation of CD8+ effector and memory T cell

populations during infection. *J Exp Med* 2015, **212:**2027-2039.

73.   Zhang L, Zhou R, Zhang W, Yao X, Li W, Xu L, Sun X, Zhao L: **Cysteine-rich**

intestinal protein 1 suppresses apoptosis and chemosensitivity to 5-fluorouracil

in colorectal cancer through ubiquitin-mediated Fas degradation. *J Exp Clin*

Cancer Res 2019, **38:**120.

74.   Chaudhuri L, Srivastava RK, Kos F, Shrikant PA: **Uncoupling protein 2 regulates**

metabolic reprogramming and fate of antigen-stimulated CD8+ T cells. *Cancer*

Immunol Immunother 2016, **65:**869-874.

75.   Stegelmann F, Bastian M, Swoboda K, Bhat R, Kiessler V, Krensky AM,

Roellinghoff M, Modlin RL, Stenger S: **Coordinate expression of CC chemokine**

ligand 5, granulysin, and perforin in CD8+ T cells provides a host defense

mechanism against Mycobacterium tuberculosis. *J Immunol* 2005, **175:**7474-7483.

76.   Pena SV, Hanson DA, Carr BA, Goralski TJ, Krensky AM: **Processing, subcellular**

localization, and function of 519 (granulysin), a human late T cell activation

molecule with homology to small, lytic, granule proteins. *J Immunol* 1997,

158:2680-2688.

77.   Pena SV, Krensky AM: **Granulysin, a new human cytolytic granule-associated**

protein with possible involvement in cell-mediated cytotoxicity. *Semin Immunol*

1997, **9:**117-125.

78.   Gubser PM, Bantug GR, Razik L, Fischer M, Dimeloe S, Hoenger G, Durovic B,

Jauch A, Hess C: **Rapid effector function of memory CD8+ T cells requires an**

immediate-early glycolytic switch. *Nat Immunol* 2013, **14:**1064-1072.

79.   Rangel Rivera GO, Knochelmann HM, Dwyer CJ, Smith AS, Wyatt MM, Rivera-
Reyes AM, Thaxton JE, Paulos CM: **Fundamentals of T Cell Metabolism and**
Strategies to Enhance Cancer Immunotherapy. *Front Immunol* 2021, **12:**645242.
80.   Cham CM, Gajewski TF: **Glucose availability regulates IFN-gamma production**
and p70S6 kinase activation in CD8+ effector T cells. *J Immunol* 2005, **174:**4670-
4677.
81.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, Lee MJ,
Wilk AJ, Darby C, Zager M, et al: **Integrated analysis of multimodal single-cell**
data. *Cell* 2021, **184:**3573-3587 e3529.
82.   Stoeckius M, Zheng S, Houck-Loomis B, Hao S, Yeung BZ, Mauck WM, 3rd,
Smibert P, Satija R: **Cell Hashing with barcoded antibodies enables multiplexing**
and doublet detection for single cell genomics. *Genome Biol* 2018, **19:**224.
83.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,
Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP: **Gene set enrichment**
analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. *Proc Natl Acad Sci U S A* 2005, **102:**15545-15550.
84.   Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, Purdom E, Dudoit S:
Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics.
BMC Genomics 2018, **19:**477.

Acknowledgements
We thank the members of Immunogenomics laboratory at National Institute of Immunology
(NII) for useful discussions and suggestions.
Funding

This work was supported by the DBT/Wellcome Trust India Alliance Fellowship (grant

number IA/I/18/2/504012) awarded to VSP and DBT-NII core funding to VSP. RK is

supported by DBT-JRF (Department of Biotechnology-Junior Research Fellowship) for his

PhD.

Author information

Authors and Affiliation

National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi, India

Raunak Kar, Somdeb Chattopadhyay, Kirti Sharma, Shreya Sinha, Gopalakrishnan

Aneeshkumar Arimbasseri, Veena S. Patil

Vardhman Mahavir Medical College and Safdarjung Hospital, New Delhi, India

Anjali Sharma

Corresponding authors

Correspondence to Gopalakrishnan Aneeshkumar Arimbasseri (aneesh@nii.ac.in) or Veena S.

Patil (veena@nii.ac.in)

Lead contact: Veena S. Patil (veena@nii.ac.in)

Contributions

RK and VSP conceived, designed, and executed the study. RK, KS and SS performed and

analysed the experiments under the supervision of VSP, Genomics data was analysed by RK

and SC under the supervision of GAA and VSP, AS supervised patient recruitment and sample

collection, RK and VSP wrote the manuscript, which was further edited and approved by all

authors.

Ethics declaration

Ethics approval and consent to participate

Approval for the use of human material for research was obtained from the institutional human

ethics committees from both National Institute of Immunology, New Delhi and Vardhman

Mahavir Medical College (VMMC) and Safdarjung Hospital, New Delhi, India. The donors

consented for use of the plasma and buffy coat samples for research.

Competing interests

The authors declare that they have no competing interests.

Figure Legends

Fig. 1. Single-cell transcriptomic analysis of hCMV-reactive memory T cell subsets. (A)

Schematic showing the study overview of isolating and single-cell RNA-seq of hCMV-specific

T cells using 10X Genomics platform. Histograms (on the right) shows cells isolated using

FACS for single-cell RNA-Seq from indicated T cell memory compartment from 10 donors.

(B) 2D UMAP of single-cell RNA-Seq data from 6899 cells, using 2000 most variable

transcripts, shows 9 distinct clusters of cells of hCMV-specific T cells. (C) 2D UMAP plots

(top) and violin plots (bottom) show expression of the indicated protein (oligo-tagged

antibodies; adt). Colour scale shows expression levels of individual cells (UMAP) or mean

expression in each cluster (violin plot). Only cells (6067) obtained from experiment 2 with 10

donors are shown. (D) Dot plots show the mean expression (colour) and percentage of

expressing cells (size) for selected marker gene transcripts in each cluster. (E) 2D UMAP plots

of single-cell RNA-Seq data show the expression of the *CD4*, *CD8A* and *CD8B* transcripts.

Scale represents expression level of the given transcript. Cells with zero counts for the indicated

transcript are shown in grey colour. (F) 2D UMAP plots show the T_{CM}, T_{EM} and T_{EMRA}

signature score for each cell. (G) 2D UMAP plots (left) of single-cell RNA-Seq data show the

expression of the *CD154* (top) and *CD137* (bottom) transcripts. Scale represents expression

level of the given transcript. Cells with zero counts for the indicated transcripts are shown in

grey colour. Scatter plots (right) show the co-expression matrix for *CD154* (on X-axis), and

CD137 (on Y-axis) with *CD69* or *TNFRSF4* (colour scale: mean expression) transcripts. For

violin plots in (C) and dot plots in (D) cluster with less than 1% of the total cells is not shown

(cluster 8=56 cells). All expression counts for transcripts are $\log_2$ normalized (CPM+1).

Fig. 2. hCMV-reactive T cell subtypes and their TCR clonal diversity. (A) Heatmap shows

the expression of the most significantly enriched transcripts in each cluster. The top 200

transcripts are shown based on adjusted *P* value < 0.05, $\log_2$ fold change > 0.25 and > 10%

difference in the percentage of cells expressing selected transcript between two groups of cells

compared. Each column represents the average expression of all cells for a given cluster.

Colour scale represents z-score for the expression of each transcript across clusters. (B) Dot

plots show the mean expression (colour) and percentage of expressing cells (size) for selected

marker gene transcript in each cluster. (C) 2D UMAP plots show the IFN response, T_{REG},

cytotoxicity and cell cycle signature score for each cell. (D) Violin plots show the normalized

expression level ($\log_2$(CPM+1)) of the indicated differentially expressed transcript. Colour

scale shows percentage of cells expressing the indicated transcript for each cluster. (E) Bar

graph shows the percentage of cells with expanded clonotypes (frequency≥2) across 9 clusters.

(F) Bar graph shows the frequency of 25 most expanded clonotypes across different clusters.

For B and D, clusters with <1% of the total cells are not shown (cluster 8=56 cells).

Fig. 3. Single-cell transcriptomic analysis of hCMV-reactive T_{REGs}. (A) 2D UMAP plots

of single-cell RNA-Seq data show the expression of the *FOXP3*, *IL2RA*, *IKZF2* (left) and

joint expression of *FOXP3* with *IL2RA* and *FOXP3* with *IKZF2* (right 2 plots) transcripts.

Scale represents $\log_2$ normalized (CPM+1) counts for expression values and scaled density

for joint expression plots. Cells with zero counts for the indicated transcripts are shown in

grey colour. (B) Bar graph shows the percentage of cells expressing *FOXP3* transcript

(log$_2$(CPM+1)>0) in clusters 1 and 5. (C) Dot plots show the mean expression (colour) and

percentage of expressing cells (size) for selected marker gene transcripts in indicated clusters.

Clusters with <1% of the total cells are not shown (cluster 8=56 cells). (D) Scatter plots

showing normalized co-expression level (log2(CPM+1)) between *FOXP3* and *GATA3* and

FOXP3 and *IL2RA* transcripts in the indicated clusters. (E) Bar graph shows the percentage

of cells expressing *GATA3* or *IL1R2* transcripts (log$_2$(CPM+1)>0) within *FOXP3* transcript

expressing cells in clusters 1 and 5. (F) 2D UMAP plots show joint expression of *FOXP3* +

GATA3, *FOXP3* + *CCR4* and *FOXP3* + *IL1R2*, *FOXP3* + *CCR8*. (G) Violin plot shows the

module enrichment score for T_H2-specific gene set across 8 clusters. Colour scale indicates

mean module score. Clusters with <1% of the total cells are not shown (cluster 8=56 cells).

(H) Gene set enrichment plot for T_H2 specific gene sets in cluster 1 and cluster 5 versus all

the other cluster. RES: Running Enrichment Score; NES: Normalized Enrichment Score;

FDR: False Discovery Rate. (I) 2D UMAP plot shows joint expression of T_H2-specific gene

set and allergy induced T_{REG}-specific gene set. Scale represents density for joint expression

plot.

Fig. 4. Single-cell transcriptomic analysis of hCMV-reactive CD4-CTLs and CD8-CTLs.

(A) 2D UMAP plots of single-cell RNA-Seq data show the expression of the indicated

transcripts. All expression counts for transcripts are log$_2$ normalized (CPM+1). Cells with zero

counts for the indicated transcripts are shown in grey colour. (B) Dot plots show the mean

expression (colour) and percentage of expressing cells (size) for selected marker gene

transcripts in each cluster. (C) Violin plots showing the normalized expression level

(log$_2$(CPM+1)) of the indicated differentially expressed transcripts across 8 clusters. Colour

scale shows percentage of cells expressing the indicated transcript for a given cluster. (D)

Trajectory analysis using slingshot shows the distribution of cells from indicated clusters (middle) on a pseudo-scale (left), and expression levels (scale) of the *GZMB*, *GNLY*, *CRTAM* and *KLRD1* transcripts. (E) 2D UMAP plot shows the cells with same clonotypes (colours) shared between cluster 3 and, clusters 2 and 4. (F) Pie chart shows the distribution of percentage of cells expressing the top 20 expanded clonotypes in clusters 2 and 4. (G) Stacked bar graph shows the proportion of CD4, CD8 and NA (not assigned) T cells in each indicated cluster (refer to methods section for classification of cells). (H) Heatmap shows the normalized expression of 50 differentially expressed transcripts from each clusters 2 and 4 when compared to the rest of the clusters. (I) Violin plots show the normalized expression level ($\log_2(CPM+1)$) of the indicated differentially expressed transcript across indicated T cell subsets in clusters 2 and 4. Colour scale shows percentage of cells expressing the indicated transcript. For B and C, clusters with <1% of the total cells are not shown (cluster 8=56 cells).

Additional files

Additional file 1. Supplementary figures.

Fig. S1. Single-cell RNA-Seq of hCMV-specific cells. (A) Graph shows the hCMV-specific IgG positivity (normalized absorbance) in 126 human donors. (B) Contour plots showing the gating strategy used to analyse the activated T cells in the indicated subsets upon stimulation of the PBMCs with hCMV peptide pools. (C) Violin plot shows the normalized T cell reactivity to hCMV-specific peptide pools compared to unstimulated control. Donors above the horizontal dotted line for memory compartments were chosen for further single-cell genomics studies (10 out of 16). (C) Stacked bar graph shows the distribution of cells from different donors across 9 clusters. (D) Stacked bar graph (left) and the 2D UMAPs (right) show the distribution of cells across 9 clusters from 2 different experiments/batch. Experiment 1

included 3 donors (D1, D2 and D7) and experiment 2 included all 10 donors. (E) Violin plots show the module enrichment score for T_{CM}, T_{EM}, T_{EMRA}, T_{CM}-T_{EM}, and T_{EM} -T_{EMRA} specific gene sets across 8 clusters. Colour scale indicates average module score. Clusters with <1% of the total cells are not shown (cluster 8=56 cells).

Fig. S2. Functional characterization and TCR analysis of hCMV-specific T cells. (A) Violin plots show the module enrichment score for T_H1, T_H17, and $T_H1/17$ subset-specific gene sets across 8 clusters. Colour scale indicates average module score. Clusters with <1% of the total cells are not shown (cluster 8=56 cells). (B) UpSet plot shows the TCR clonotypes (frequency on y-axis) and sharing of the clonotypes (bottom, connected dots) across the 9 clusters. (C) Alluvial plot shows the clonally expanded cells (1234 cells) and their distribution and sharing across different clusters. Each connecting line represents a single cell.

Fig. S3. Pathways enriched in hCMV-specific cytotoxic clusters. (A) Pathway enrichment analysis using Ingenuity Pathway Analysis (IPA) (Qiagen) shows enrichment of pathways in cluster 2 compared to cluster 4. (B) Pathway enrichment analysis using IPA shows enrichment of pathways in cluster 4 compared to cluster 2.

Additional file 2: Supplementary Tables

Table S1. Details of cells from single-cell genomics experiments.

Table S2: Details of reagents used in the study

Table S3: Gene list used for enrichment analysis in the study

Table S4. Single-cell cluster wise enrichment of transcripts using MAST (log2FC>0.25; adj pval <0.05)

Table S5: TCR clonotype details of hCMV-reactive single-cells

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. S2

A

■ Overlaps with dataset □ No overlap with dataset ■—■ -log(p-value)

B