Inter and Intralaminar Excitation of Parvalbumin Interneurons in Mouse Barrel Cortex

Running title: Laminar Response Differences of Parvalbumin Interneurons

Kate S. Scheuer[1]

Anna M. Jansson[2]

Xinyu Zhao[3]

Meyer B. Jackson[2]

1. Cellular and Molecular Biology Program, 2. Department of Neuroscience, 3. Waisman Center,

University of Wisconsin-Madison, Madison, Wisconsin, 53705

Author contributions: Data collection (KSS), data analysis (KSS, AMJ), manuscript preparation

(KSS, MBJ, XZ), project conception (MBJ, KSS, XZ)

Corresponding author address: Department of Neuroscience, Room 5505 WIMR II, 1111

Highland Ave., Madison, Wisconsin 53705, Phone: 608-262-2938,

ORCID ID 0000-0003-2964-7091

Email: meyer.jackson@wisc.edu

Keywords: parvalbumin interneuron, genetically-encoded voltage sensor, barrel cortex

Conflict of interest statement: The authors declare no competing financial or non-financial

interests.

Acknowledgements: National Institutes of Health Grants NS127219 and NS093866 to MBJ and

NS105200 XZ. Thanks to Dr. Shane McMahon for methodological contributions.

Key points summary

- Excitatory synaptic responses were imaged in parvalbumin (PV) interneurons in slices of mouse barrel cortex using a targeted genetically-encoded voltage sensor. This approach revealed simultaneous voltage changes in approximately 20 neurons pre slice in response to stimulation.

- PV interneurons residing in layer 2/3 had larger amplitudes, longer half-widths, and longer rise-times than PV interneurons residing in layer 4.

- Responses of PV interneurons residing in either layer 2/3 or layer 4 had shorter latencies to stimulation of layer 4 compared to stimulation of layer 2/3.

- Excitatory synaptic transmission to PV interneurons varies with layer of residence and source of excitation.

Abstract

Parvalbumin (PV) interneurons are inhibitory fast-spiking cells with essential roles in directing the flow of information through cortical circuits. These neurons set the balance between excitation and inhibition, control rhythmic activity, and have been linked to disorders including autism spectrum and schizophrenia. PV interneurons differ between cortical layers in their morphology, circuitry, and function, but how their electrophysiological properties vary has received little attention. Here we investigate responses of PV interneurons in different layers of primary somatosensory barrel cortex (BC) to different excitatory inputs. With the genetically-encoded hybrid voltage sensor, hVOS, we recorded voltage changes simultaneously in many L2/3 and L4 PV interneurons to stimulation in either L2/3 or L4. Decay-times were consistent across L2/3 and L4. Amplitude, half-width, and rise-time were greater for PV interneurons

residing in L2/3 compared to L4. Stimulation in L2/3 elicited responses in both L2/3 and L4 with

longer latency compared to stimulation in L4. These differences in latency between layers could

influence their windows for temporal integration. Thus PV interneurons in different cortical

layers of BC show differences in response properties with potential roles in cortical

computations.

Introduction

Parvalbumin (PV) interneurons are inhibitory neurons defined by their expression of the

calcium-binding protein PV (Tremblay *et al.*, 2016). These fast-spiking cells are present in

cortical layers 2-6 and play critical roles in controlling excitation/inhibition balance (Ferguson &

Gao, 2018; Nahar *et al.*, 2021), and in the generation of gamma oscillations, 30-80 Hz brain

waves implicated in many functions including working memory, attention, and perceptual

binding (Tallon-Baudry *et al.*, 1998; Gonzalez-Burgos *et al.*, 2015). PV interneurons and gamma

oscillations have both been linked to a variety of psychiatric conditions including schizophrenia

and autism spectrum disorder (Gonzalez-Burgos *et al.*, 2015; Lauber *et al.*, 2018; Kayarian *et al.*,

2020).

Primary somatosensory barrel cortex (BC) is an attractive place to study PV interneurons

because of its well-defined functions and architecture (Brecht, 2007; Feldmeyer, 2012;

Feldmeyer *et al.*, 2018; Staiger & Petersen, 2021). BC is defined by the presence of barrels,

cytoarchitectural units in L4 which each correspond to a single vibrissa (Woolsey & Van der

Loos, 1970). Distinct molecular, morphological, and electrophysiological cell types form

complex circuits both within and between cortical layers in BC. Two major PV interneuron

morphological subgroups, basket cells and chandelier cells, are distributed differently across

cortical layers in BC. Chandelier cells form axoaxonic contacts and are not present in L4 (Li &

Huntsman, 2014), while basket cells provide perisomatic inhibition and can be found in L2-6

(Naka & Adesnik, 2016; Frandolig *et al.*, 2019; Staiger & Petersen, 2021). These two

morphological subgroups can form distinct interlaminar circuits (Xu & Callaway, 2009). In

addition to these morphological differences, PV interneurons in different layers may have

different roles in functions such as intracortical and thalamic integration (Staiger & Petersen,

2021). Additionally, optogenetically generated gamma oscillations within a given cortical layer

inhibit locally within that layer but facilitate in other layers. Furthermore, peak gamma

oscillation power was higher for L6 compared to L2/3 (Adesnik, 2018). These results raise the

important question of whether the roles of PV interneurons in different layers reflect differences

in their responses to excitatory synaptic inputs. However a simultaneous assessment of voltage

responses of PV neurons in different cortical layers has not been carried out.

Here we use the genetically-encoded hybrid voltage sensor (hVOS) to record excitatory post-

synaptic potentials (EPSPs) optically from L2/3 and L4 PV interneurons in slices of mouse BC

(Chanda *et al.*, 2005; Wang *et al.*, 2010; Bayguinov *et al.*, 2017). We determined PV interneuron

response amplitude, half-width, latency, rise-time, and decay-time elicited by stimulation in L2/3

and L4. Regardless of stimulation layer, L2/3 PV interneuron responses had higher amplitudes,

longer rise-times, and broader half-widths than L4 PV interneurons. Additionally, responses to

stimulation in L2/3 had longer latencies than responses to L4 stimulation, even after accounting

for the effect of conduction distance. By contrast, responses in these layers had similar decay-

times. Thus, hVOS imaging reveals variations in electrophysiological properties of PV

interneurons between cortical layers. These differences have implications for how the cortex

performs computations and integrates inputs between different layers.

Methods

Animals

PV-Cre driver mice (B6.129P2-Pvalb$^{tm1(cre)Arbr}$/J, JAX strain 017320) were crossed with

Ai35-hVOS1.5 Cre-reporter mice (C57BL/6-*Gt(ROSA)26Sor$^{tm1(CAG-hVOS1.5)Mbja}$*/J, JAX strain

031102) to generate animals with hVOS probe targeted to PV interneurons (Bayguinov *et al.*,

2017). Animal procedures were approved by the University of Wisconsin-Madison School of

Medicine and Public Health Animal Care and Use Committee (IACUC protocol: M005952).

hVOS probe

An hVOS probe was used to image voltage changes in PV interneurons. The probe used

here is comprised of cerulean fluorescent protein (CeFP) tethered to the inner leaflet of the cell

membrane with a truncated h-ras motif (Wang *et al.*, 2010). Cells expressing the probe fluoresce,

and this fluorescence is modulated by a Förster resonance energy transfer interaction with

dipicrylamine (DPA), a small, hydrophobic anion which partitions into the cell membrane and

moves when the membrane potential changes. Depolarization drives DPA towards the CeFP and

fluorescence is quenched. Repolarization drives DPA back away from the CeFP so fluorescence

increases. Fluorescence thus reports voltage changes of cells expressing the hVOS probe

(Chanda *et al.*, 2005; Wang *et al.*, 2010). hVOS has sub-millisecond temporal resolution (Chanda

et al., 2005; Bradley *et al.*, 2009) and can be genetically targeted to specific cell types using a

Cre-lox system (Bayguinov *et al.*, 2017). Our crossing of hVOS Cre-reporter animals with PV

Cre-driver animals produces mice previously shown to have 99.2% targeting specificity and

express the hVOS probe in 83% of PV interneurons (Bayguinov *et al.*, 2017).

Slice preparation

Mice 7-8 weeks old were deeply anesthetized with isoflurane and sacrificed with cervical

dislocation (institutional protocol noted above). Brains were rapidly dissected and placed into

ice-cold cutting solution (in mM: 10 glucose, 125 NaCl, 4 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 6

$MgSO_4$, 1 $CaCl_2$) bubbled with a mixture of 95% O_2 / 5% CO_2. After approximately five

minutes, brains were mounted and cut into 300 μm thick coronal slices with a with a Leica

VT1200S vibratome. Slices were placed into a chamber filled with 95% O_2 / 5% CO_2-bubbled

artificial cerebrospinal fluid (ACSF) with the same composition as cutting solution except with

1.3 mM $MgSO_4$ and 2.5 mM $CaCl_2$, and allowed to recover for at least 45 minutes.

Electrophysiology and Imaging

Imaging experiments were performed in 95% O_2 / 5% CO_2-bubbled ACSF containing 4

μM DPA. Slices were placed into a custom recording chamber, and viewed with a BX51

Olympus microscope. Stimulus pulses 200 μA, 180 μsec were applied with a stimulus isolator

(World Precision Instruments, Sarasota, Florida) through fire-polished, ACSF-filled KG-33 glass

electrodes (King Precision Glass, Claremont, California) with tip diameters of about 6-8 μm.

Stimulating electrodes were positioned in L2/3 or L4 of BC using a micromanipulator. Slices

were illuminated with an LED with peak emission at 435 nm (Prizmatix, Holon, Israel) through a

CFP filter cube. PV interneuron responses were acquired with a CCD-SMQ camera (RedShirt

Imaging, Decatur, Georgia) at 2000 Hz with 80x80 spatial resolution. Bandpass filters of 5 and

10 nm centered at 435 nm were added to the excitation pathway when resting light intensities

saturated the CCD-SMQ camera. Gradient contrast and higher resolution fluorescence images

were captured by directing light to a Kiralux CMOS camera (Thorlabs, Newton, New Jersey).

Data acquisition and analysis was performed with custom software (Chang, 2006).

Identifying individual responsive PV interneurons

PV interneurons have extensive axonal and dendritic arbors which allow them to sample

input from many cells, and to provide strong, widespread inhibition (Fukuda & Kosaka, 2003;

Povysheva *et al.*, 2008; Packer & Yuste, 2011; Hu *et al.*, 2014). Cortical PV interneurons contact

about 43-50% or more of pyramidal cells within about < 200 μm (Packer & Yuste, 2011; Inan *et*

al., 2013), and fast-spiking interneuron to excitatory cell connectivity in BC can be as high as

67% for a sub-group of PV interneurons in L4 (Koelbl *et al.*, 2015). Given the dense dendritic

and axonal arbors of PV interneurons, a plasma membrane label such as the hVOS probe

produces broad diffuse fluorescence throughout a slice, obscuring the fluorescence from PV

interneuron cell bodies. This makes it difficult to identify individual PV interneuron cell bodies,

despite clearly identifiable cortical layers in both gradient contrast (Fig. 1A) and fluorescence

(Fig. 1B) images.

To address this problem, we developed a semi-automated method of analyzing imaging

data for objective, reproducible identification of responsive PV interneuron somata. This method

relied on a hybrid approach using both geometric constraints and K-means clustering of signal-

to-noise ratio (SNR) values. SNR was calculated as the peak stimulus-evoked fluorescence

change divided by the baseline root-mean-square fluorescence in a 20-msec pre-stimulus time

window. Pixels with a SNR below the baseline noise were discarded as clearly unresponsive

(gray pixels near top, Fig. 1C). For the geometric constraints, we required regions of interest

(ROIs) representing responsive PV interneurons to be spatially distinct, with no shared faces.

Acceptable ROIs were groups of up to nine contiguous pixels three or fewer pixels across. With

6 μm pixel dimensions this constrains groups to the size of a murine PV interneuron soma, which

is approximately 20 μm in diameter (Wang *et al.*, 2002; Selby *et al.*, 2007; Kooijmans *et al.*,

2020). This criterion excluded some larger groups of pixels as potentially representing more than

one cell even though they had very clear responses with a high SNR. For example, the red,

orange, and yellow regions in the lower left and lower right corners of Fig. 1C and 1E had high

SNR values but formed groups which were much larger than 3 pixels across and clearly

contained several responsive cells. Pixels in these areas were excluded from analysis. Because a

single pixel (6 μm) is too small to be a cell body and could contain several overlapping PV

interneuron dendrites (<0.5-3 μm in rats, smaller in mice (Muller *et al.*, 2005; Judak *et al.*,

2022)) and/or axons (< 1 μm, (Stedehouder *et al.*, 2019)), single isolated pixels were not

considered to be responsive PV interneurons, again despite their high SNR. Responses from

pixels obscured by the stimulating electrode (outlined in black, Fig. 1C-E) were also excluded.

Finally, responses < 45 μm from the tip of the stimulating electrode were assumed to be the

result of direct stimulation and excluded.

To refine and corroborate this procedure, ROIs corresponding to putative responsive PV

interneurons satisfying the geometric constraints were subjected to one-dimensional K-means

clustering of SNR values. One-dimensional K-means clustering was performed on pixels with

SNR above baseline noise (all but the gray pixels in Fig. 1C). Clustering served two main

purposes. First, it divided pixels into groups with similar SNR values. Pixel clusters with higher

average SNR are more likely to contain cell bodies (yellow, Fig. 1D) while those with lower

average SNR are more likely to contain small processes or lack responsive cells (purple, Fig.

1D). We therefore set a response SNR cutoff for K-means clusters to 5 and excluded pixels in

clusters of pixels with average SNR < this cutoff (purple, Fig. 1D), as they likely contained

processes or unresponsive cells. For acceptable clusters (average SNR > 5) we assumed that if

pixels within a group satisfying the geometric requirements have SNR values in the same K-

means cluster, they are likely to represent the same cell body. K-means clustering identified

groups of geometrically associated pixels with similar SNR and assigned them to specific cells.

This method basically compared each pixel to its neighbors and grouped them based on the

likelihood they represent the same cell. In summary, this method was conservative,

implementing multiple exclusion criteria to focus on small groups of pixels with similar SNR

that represent distinct, spatially separated neurons.

Traces of fluorescence versus time from groups of pixels identified in this way were

manually inspected to verify appropriate responses to stimulation (Fig. 1F). We implemented an

additional cutoff based on amplitude, and discarded pixel groups with average $\Delta F/F < 0.1\%$.

Responses with amplitudes more than 3 times the standard deviation above the mean value (>

1.165%) were also excluded. Such instances were very rare (4 PV interneurons total), and

occurred in particularly dark areas or corners of the field of view where resting light was very

low and dividing resulted in implausibly high values. Because our analysis compares PV

interneuron properties based on cortical layer, occasional cells on a border between cortical

layers were also excluded.

Figure 1. Identifying individual responsive PV interneurons. Gradient contrast image taken with the Kiralux camera (A) and fluorescence image taken with the CCD-SMQ camera (B) of a BC slice. Black stars indicate the tip of the stimulating electrode, and dashed lines indicate layer boundaries in A-E. C. SNR heatmap of slice in A and B. Gray pixels near the top have signals below the baseline noise and were excluded from analysis. The electrode is outlined in black on the lower left edge in C-E. D. K-means cluster map. K-means clustering of SNR was performed on pixels with SNR > baseline (colored in C). The data were best fitted with two clusters, with averages of 4.8 and 9.2. The yellow cluster with higher average SNR is likely to contain responsive PV interneurons, while the purple cluster with lower average SNR probably contains processes and unresponsive neurons. E. SNR heatmap overlaid with identified responsive PV interneurons outlined in black or red (color choice based on ease of view and not cell properties). F. Traces of fluorescence versus time for the PV interneurons outlined and numbered in E show clear depolarization in response to stimulation (triangle). G. Expanded portion of a trace (12 msec) shaded in F illustrating response parameters. Amplitude (red) is the maximum change in fluorescence; latency (purple) is the time from stimulation to half-maximal change in fluorescence; half-width (green) is the time between half-maximal change in fluorescence from depolarization to repolarization; rise-time (blue) is the time between half-maximal and maximal change in fluorescence; decay-time (gold) is the time from peak to half-maximal fluorescence.

Despite the conservative nature of this analysis, our procedure identified an average of 21

responsive PV interneurons per slice. Although anatomical estimates of PV interneuron density

vary widely, conservative estimates suggest that our 480x480 μm field of view may contain up to

approximately 75 PV interneurons (Keller *et al.*, 2018). Our numbers are generally well below

this, supporting our procedure as a conservative method of identifying responsive somata. This

method provided a reproducible, objective, and robust procedure to identify individual

responsive PV interneurons. A test of validity is presented in Results (Fig. 2).

Response parameters were extracted from traces of fluorescence versus time for pixel

groups identified as corresponding to responsive PV interneurons (Fig. 1G). Amplitude is the

maximum change in fluorescence. Latency is the time from stimulation to half-maximal change

in fluorescence during depolarization. To account for the effect of distance on latency, we also

divided the latency by distance to the stimulating electrode (distance-normalized latency). Half-

width is the time between half-maximal change in fluorescence during depolarization and

repolarization. Rise-time is the time from half-maximal to maximal fluorescence during

depolarization. Decay-time is the time from maximum to half-maximal fluorescence during

repolarization. All responses were examined visually and 4 were excluded because noise resulted

in anomalous start times with obvious errors in parameters.

Data processing and statistical tests

Fluorescence traces were processed with a nine-point binomial temporal filter and a

spatial filter with $\sigma = 1$. A baseline determined from a polynomial fit was subtracted. Peak

fluorescence change was divided by resting light intensity to give $\Delta F/F$. Our method of

responsive cell identification yielded 1086 PV interneurons from 52 slices from 7 animals (3

female, 4 male). Relationships between distance and half-width or amplitude were evaluated

with Pearson's product-moment correlation tests using individual PV interneurons as the unit of

analysis. For remaining statistical tests, within a given slice layers with fewer than 8 responsive

PV interneurons were excluded, values were averaged for all PV interneurons within each layer,

and this average was used as the unit of analysis.

230   Normality was evaluated using Shapiro-Wilks tests. All parameters were normally

distributed (amplitude: $W = 0.982$, $p = 0.591$; half-width: $W = 0.986$, $p = 0.757$; rise-time: $W =$

0.975, $p = 0.300$) or log-normally distributed (distance-normalized latency: $W = 0.971$, $p =$

0.210; decay-time: $W = 0.986$, $p = 0.750$). Variance between analysis groups was evaluated with

Levene's tests. Variance did not differ significantly for any parameter based on sex (amplitude:

$F(1,53) = 0.903$, $p = 0.346$; half-width: $F(1,53) = 0.078$, $p = 0.782$; distance-normalized latency:

$F(1,53) = 3.808$, $p = 0.056$; rise-time: $F(1,53) = 0.353$, $p = 0.555$; decay-time: $F(1,53) = 0.252$, p

$= 0.618$). The effect of sex was evaluated with t-tests and showed no significant impact

(amplitude: $t(52.857) = 0.488$, $p = 0.628$; half-width: $t(50.648) = -1.018$, $p = 0.314$; distance-

normalized latency: $t(52.438) = -1.252$, $p = 0.216$; rise-time: $t(48.567) = -0.652$, $p = 0.517$;

decay-time: $t(52.005) = -0.893$, $p = 0.376$).

241   Variance did not differ significantly based on PV interneuron layer or stimulation layer

for any parameter (amplitude ($F(3,51) = 0.465$, $p = 0.708$); half-width ($F(3,51) = 0.593$, $p =$

0.623); distance-normalized latency ($F(3,51) = 1.052$, $p = 0.378$); rise-time ($F(3,51) = 0.596$ $p =$

0.321); decay-time ($F(3,51) = 0.223$, $p = 0.880$)). The effects of stimulation layer and/or PV

interneuron residence layer on each parameter were therefore evaluated with ANOVA and post-

hoc Tukey's honestly significant differences tests.

Code availability

R code, Python code, and custom software available on request.

Results

Validation of single cell identification

251   We tested our procedure of cell identification by examining variations with distance. If

pixel groups actually represent multiple neighboring neurons rather than a single neuron, we

would expect response half-width to broaden and response amplitude to decrease with distance

from the stimulating electrode, as the activation of more distant groups should be less

synchronous compared to closer groups. In plots versus distance neither parameter was

significantly correlated with distance (Fig. 2, half-width: R = 0.006, p = 0.854; amplitude: R =

0.042, p = 0.170), indicating that pixel clusters do not contain more than one PV interneuron.

Figure 2. PV interneuron response half-width and amplitude do not vary with distance. Neither half-width (A, R = 0.006, p = 0.854) nor amplitude (B, Pearson's product-moment correlation: R = 0.042, p = 0.170) are significantly correlated with distance from the stimulating electrode. This is consistent with single-cell responses, as half-width would be expected to increase, and amplitude would decrease with distance for a population response. Each point on the scatterplot corresponds to one PV interneuron. Linear regression best fit lines are shown in blue. N = 1086 cells from 52 slices.

259   *PV interneuron responses vary between cortical layers*

260   Stimulation in L2/3 (Fig. 3A-B, left) or L4 (Fig. 3A-B, right) elicited responses across

261   L2/3 through L5 as shown in SNR heatmaps (Fig. 3C). Although PV interneurons in L5

262   responded to stimulation in either L2/3 or L4, less L5 was present in the field of view selected

263   for study so we did not see enough L5 PV interneuron responses to include in the current

264   analysis. L2/3 and L4 PV interneuron response parameters are presented in Table 1. Comparisons

265   are presented in bar graphs below and will be discussed in detail.

	L2/3 Stimulation		L4 Stimulation	
	L2/3 PV interneurons (N = 322, 29 slices)	L4 PV interneurons (N = 259, 29 slices)	L2/3 PV interneurons (N = 331, 23 slices)	L4 PV interneurons (N = 160, 23 slices)
Amplitude (% Δ F/F)	0.567 ± 0.005	0.363 ± 0.004	0.556 ± 0.005	0.422 ± 0.005
Half-width (msec)	5.29 ± 0.067	4.82 ± 0.049	5.45 ± 0.059	4.71 ± 0.050
Raw Latency (msec)	2.79 ± 0.059	3.89 ± 0.053	3.28 ± 0.050	2.38 ± 0.046
Distance-normalized latency (msec/μm)	0.015 ± 0.0002	0.017 ± 0.0002	0.013 ± 0.0002	0.014 ± 0.0002
Rise-Time (msec)	2.32 ± 0.047	2.17 ± 0.039	2.48 ± 0.044	2.02 ± 0.039
Decay-Time (msec)	2.97 ± 0.051	2.64 ± 0.036	2.97 ± 0.043	2.69 ± 0.037

266   Table 1. PV interneuron response parameters based on residence layer and stimulation layer.
267   Values are mean ± SE, N = number of neurons.

Figure 3. PV interneuron responses in BC. Gradient contrast (A) and fluorescence (B) images of two different slices of BC. L2/3 through L5 are visible within the fields of view. The tip of the stimulating electrode (black or white star) is visible in L2/3 (left) or L4 (right) in A-C. Dashed lines separate layers. C. SNR heatmaps for the slices shown in A and B. Warmer colors correspond to higher SNR regions more likely to contain responsive PV interneurons (color scales and ranges – lower right).

Figure 4. Amplitude, rise-time, and half-width of PV interneurons residing in different layers. L2/3 PV interneuron responses (blue) had higher amplitudes (A), longer rise-times (B), and broader half-widths (C) than responses from PV interneurons in L4 (purple). Decay-time (D) did not differ based on PV interneuron residence layer. Stimulation layer did not significantly impact amplitude, half-width, rise-time, or decay-time.

PV interneuron residence layer significantly impacted amplitude (Fig. 4A, $F_{(1,51)}$ =

32.438, $p < 0.001$), rise-time (Fig. 4B, $F_{(1,51)}$ = 4.753, $p = 0.034$), and half-width (Fig. 4C,

$F_{(1,51)}$ = 4.710, $p = 0.035$), but not decay-time (Fig. 4D, $F_{(1,51)}$ = 1.422, $p = 0.239$). Regardless

of stimulation layer, PV interneuron response amplitudes (0.579 ± 0.014, mean ± SE) were 46%

larger in L2/3 than in L4 (0.396 ± 0.018, mean ± SE, p < 0.001). Rise-times for PV interneurons

residing in L2/3 (2.39 ± 0.070 msec) were also longer than those in L4 (2.13 ± 0.047 msec, p =

0.034). L2/3 PV interneuron response half-widths (5.26 ± 0.108 msec) were broader than L4

response half-widths (4.82 ± 0.091 msec, p = 0.035).

Figure 5. PV interneuron response latency depends on stimulation layer. A. Raw latencies for PV interneuron responses to stimulation in L2/3 (orange) or L4 (green). B. Regardless of residence layer, responses elicited by L2/3 stimulation (orange) have longer distance-normalized latencies than those elicited by L4 stimulation (green, t = (50) = 4.244, p < 0.001).

Latency also differed based on cortical layer (Fig. 5). Unlike amplitude, rise-time, and

half-width, however, latency also depended on stimulation layer. To analyze variations in latency

we must take into account the influence of distance and conduction time (Scheuer *et al.*, 2023).

Distances from the electrode tip to PV interneurons responding to L2/3 (180 ± 5.3 μm) and L4

(178 ± 4.3 μm) intralaminar stimulation were similar, as were distances for interlaminar

responses to L2/3 (237 ± 3.4 μm) and L4 (260 ± 4.1 μm) stimulation. However, average

interlaminar distance (248 ± 4.0 μm) was 39% longer than intralaminar distance (179 ± 4.9 μm).

As expected, distance significantly impacted raw latency (F(1,53) = 49.62, p < 0.001). To

account for this effect, we compared response latencies divided by distance from the tip of the

stimulating electrode (distance-normalized latency). Stimulation layer significantly affected

distance-normalized latency ($F_{(1,51)}$ = 16.478, p < 0.001). Regardless of PV interneuron layer,

distance-normalized latencies of responses to stimulation in L2/3 (0.0162 ± 0.0005 msec/μm)

were significantly longer than those to stimulation in L4 (0.0128 ± 0.0004 msec/μm, p < 0.001).

PV interneurons residing in L2/3 responded to interlaminar L4 stimulation (0.013 ± 0.0003

msec/μm) more quickly than intralaminar L2/3 stimulation (0.016 ± 0.0004 msec/μm, t(26.992)

= 3.243, p = 0.003). However, L4 PV interneurons responded to interlaminar L2/3 stimulation

(0.017 ± 0.0005 msec/μm) more slowly than to intralaminar L4 stimulation (0.013 ± 0.0005

msec/μm). Thus, stimulation layer impacts latency in a manner which cannot be attributed to

differences in distance.

Discussion

Here we used the genetically-encoded hybrid voltage sensor hVOS to investigate PV

interneurons in BC. We observed responses of L2/3 and L4 PV interneuron to stimulation of both

these layers. We developed a semi-automatic method of identifying individual responsive PV

interneurons that combines geometric considerations with statistical K-means clustering of SNR.

This method reliably located responsive cells within the high background fluorescence produced

by the extensive arborization of PV interneurons. Using this method, we were able to identify

cell bodies of ~20 responsive PV interneurons per slice, and determine response amplitude, half-

width, latency, rise-time, and decay-time of their responses to synaptic excitation. To test our

method, we plotted half-width and amplitude versus distance from the electrode tip (Fig 2). The

absence of correlations indicates that our pixel clusters contain single cells rather than multiple

cells. The parameters reported in this study represent basic elements of cortical circuitry that may

be useful in the development of accurate computational models designed to recapitulate the roles

of fast-spiking interneurons in BC microcircuit computations (Avermann *et al.*, 2012), and in the

generation of synchronous activity (Di Garbo *et al.*, 2004; Pervouchine *et al.*, 2006).

312   Previous hVOS studies of PV interneuron activity in somatosensory cortex determined

that spike-like responses similar to action potentials had a peak $\Delta F/F$ of 2.4% (Bayguinov *et al.*,

2017), while unitary synaptic responses elicited by an action potential in a single excitatory

neuron ranged from 0.2 – 0.4% (Canales *et al.*, 2022). Assuming ~100 mV action potentials, the

mean amplitude of 0.494% reported here can be estimated as approximately 21 mV. The

responses reported here are 64% larger than unitary excitatory responses and about one fifth the

amplitude of spike-like responses. They are clearly too small to be action potentials, and are

likely to represent EPSPs elicited by an average of about two excitatory neurons. Consistent with

our assessment that these responses are synaptic potentials, our half-widths of ~5.14 msec are 3.6

times broader than half-widths of PV interneuron spikes recorded with hVOS (Bayguinov *et al.*,

2017). Our half-widths fall in the range of other studies of subthreshold, synaptic responses of

PV interneuron in murine cortex of 4.6 to 22.3 msec (Thomson, 1997; Angulo *et al.*, 1999;

Thomson *et al.*, 2002; Beierlein *et al.*, 2003; Holmgren *et al.*, 2003; Ali & Nelson, 2006; West *et*

al., 2006; Ali *et al.*, 2007; Avermann *et al.*, 2012; Zhou & Roper, 2014). Our 2.8 msec half-

decay-time corresponds to an exponential decay-time of 4.1 msec, which is within the range (3.5

to 12 msec) of previously reported values for EPSPs in murine cortical PV interneurons

(Povysheva *et al.*, 2006; Otsuka & Kawaguchi, 2009; Zaitsev & Lewis, 2013; Athilingam *et al.*,

2017). Thus, parameter values reported here fall within the range of previous reports, and reveal

how PV interneuron properties vary depending on location and source of excitation.

Figure 6. Summary of PV interneuron response differences. Amplitude, distance-normalized latency, rise-time, and half-width vary based on cortical layer. PV interneurons (teal circles) residing in L2/3 had higher amplitudes, slower rise-times, and broader half-widths compared to those in L4. Distance-normalized latencies of responses to stimulation of excitatory cells (purple triangles) in L2/3 were longer than those of responses to stimulation in L4.

Response amplitude, distance-normalized latency, rise-time, and half-width differed

based on cortical layer, and Fig. 6 illustrates the key differences. Regardless of stimulation layer,

response amplitude was greater for PV interneurons in L2/3 compared to L4. EPSP amplitude

depends on a wide variety of factors including number of inputs, dendritic location, and ion

channel and receptor makeup (Ali & Nelson, 2006; Bonsi *et al.*, 2007; Otsuka & Kawaguchi,

2009; Zaitsev *et al.*, 2012; Pala & Petersen, 2018; Guo *et al.*, 2020; Das *et al.*, 2021). The

increased EPSP amplitude in L2/3 compared to L4 PV interneurons may therefore reflect some

of these factors. PV interneuron EPSP amplitudes are more than twice as large if the presynaptic

excitatory cell and PV interneuron are reciprocally connected (Zaitsev & Lewis, 2013). Thus,

the larger EPSP amplitudes of L2/3 PV interneurons could be related to higher reciprocal

connectivity. Calcium-permeable AMPA receptors have also been shown to impact fast-spiking

interneuron synaptic responses (Bonsi *et al.*, 2007; Zaitsev *et al.*, 2012), and differences in their

distribution between layers could contribute to the present findings.

PV interneuron residence layer also impacted rise-time and half-width. L2/3 PV

interneuron responses had longer rise-times and broader half-widths than responses of PV

interneurons in L4. Differences in rise-time could reflect dendritic location, presynaptic release

kinetics, or AMPA receptor subunit composition. Compared to other types of interneurons and

pyramidal cells, PV interneurons in CA1 express higher levels of AMPA receptor subunit GluA1,

higher levels of auxiliary proteins regulating AMPA receptors, and especially high levels of

GluA4 (Yamasaki *et al.*, 2016). Knockout of GluA4, but not GluA3, decreases rise-time (Yang *et*

al., 2011). AMPA receptors on PV interneurons can have multiple subunit combinations (Kondo

et al., 1997). Therefore, differences in rise-time based on PV interneuron residence layer might

reflect layer-specific variation in AMPA receptor subunit composition. Longer rise-times for PV

interneurons in L2/3 compared to L4 likely contribute to the broader half-widths of L2/3 PV

interneurons.

In addition to differences in amplitude, rise-time, and half-width, we also observed

differences in latency between cortical layers. However, while the former properties depended on

the layer in which PV interneurons resided, latency differed depending on the location of the

inputs. Accounting for the effect of distance, both L2/3 and L4 PV interneuron responses to L2/3

stimulation had significantly longer latencies compared to responses to L4 stimulation. Because

stimulation layer impacted latencies regardless of the layer in which the PV interneurons resided,

this difference may reflect a property of the excitatory input rather than postsynaptic properties

of PV interneurons. The inputs could have a faster conduction time along their axons or more

direct axonal paths to their targets. It is also possible that the latency differences reflect

differences in the kinetics of neurotransmitter release between axons originating in L2/3 and L4.

366   The shorter latencies of L2/3 PV interneuron responses to interlaminar L4 excitation

compared to the reverse pathway of L4 PV interneuron responses to interlaminar L2/3 excitation

will enable feedforward excitation along the canonical route from L4 to L2/3 PV interneurons to

occur more quickly than L2/3 to L4 feedback. Inhibition plays a key role in coincidence

detection by controlling the temporal integration window (Pouille & Scanziani, 2001), and

because PV interneurons fire rapidly, they are particularly well-suited to tightly constraining

integration within their synaptic targets. Compared to L4, which processes more basic sensory

information such as touch and whisking, L2/3 functions are related to more complex

somatosensory processing such as object localization (O'Connor *et al.*, 2010), stimulus-specific

adaptation (Yarden *et al.*, 2022), texture discrimination (Allitt *et al.*, 2017), and social touch

(Lenschow & Brecht, 2015). The particularly rapid L4 to L2/3 excitation reported here could

narrow the integration window set by L2/3 PV interneurons on their targets, and therefore may

impact these higher-level sensory processes.

379   This work demonstrates the utility of hVOS voltage imaging as a technique to examine

cortical circuitry of many cells of a specific type simultaneously across multiple cortical layers.

This approach can be used to measure response parameters such as amplitude, half-width,

latency, rise-time, and decay-time, which are important for computations. It also provides an

opportunity to compare these response parameters across cortical layers. Here we observed

layer-based differences in amplitude, rise-time, and latency which hold implications for how BC

integrates interlaminar and intralaminar inputs. Future work building on this approach has the

potential to address the circuit functions of PV interneurons as well as other specific cell types

throughout the brain.

388
389   **References**

390   Adesnik H. (2018). Layer-specific excitation/inhibition balances during neuronal
391   synchronization in the visual cortex. *J Physiol-London* **596,** 1639-1657.
392

393   Ali AB, Bannister AP & Thomson AM. (2007). Robust correlations between action potential
394   duration and the properties of synaptic connections in layer 4 interneurones in neocortical
395   slices from juvenile rats and adult rat and cat. *J Physiol* **580,** 149-169.
396

397   Ali AB & Nelson C. (2006). Distinct Ca2+ channels mediate transmitter release at excitatory
398   synapses displaying different dynamic properties in rat neocortex. *Cereb Cortex* **16,** 386-
399   393.
400

401   Allitt BJ, Alwis DS & Rajan R. (2017). Laminar-specific encoding of texture elements in rat
402   barrel cortex. *J Physiol* **595,** 7223-7247.
403

404   Angulo MC, Rossier J & Audinat E. (1999). Postsynaptic Glutamate Receptors and Integrative
405   Properties of Fast-Spiking Interneurons in the Rat Neocortex. *J Neurophysiol* **82**.
406

407   Athilingam JC, Ben-Shalom R, Keeshen CM, Sohal VS & Bender KJ. (2017). Serotonin
408   enhances excitability and gamma frequency temporal integration in mouse prefrontal
409   fast-spiking interneurons. *Elife* **6**.
410

411   Avermann M, Tomm C, Mateo C, Gerstner W & Petersen CC. (2012). Microcircuits of excitatory
412   and inhibitory neurons in layer 2/3 of mouse barrel cortex. *J Neurophysiol* **107,** 3116-
413   3134.
414

415   Bayguinov PO, Ma YH, Gao Y, Zhao XY & Jackson MB. (2017). Imaging Voltage in Genetically
416   Defined Neuronal Subpopulations with a Cre Recombinase-Targeted Hybrid Voltage
417   Sensor. *Journal of Neuroscience* **37,** 9305-9319.
418

419   Beierlein M, Gibson JR & Connors BW. (2003). Two Dynamically Distinct Inhibitory Networks
420   in Layer 4 of the Neocortex. *J Neurophysiol* **90**.
421

422   Bonsi P, Sciamanna G, Mitrano DA, Cuomo D, Bernardi G, Platania P, Smith Y & Pisani A.
423   (2007). Functional and ultrastructural analysis of group I mGluR in striatal fast-spiking
424   interneurons. *Eur J Neurosci* **25,** 1319-1331.
425

Bradley J, Luo R, Otis TS & DiGregorio DA. (2009). Submillisecond Optical Reporting of
Membrane Potential In Situ Using a Neuronal Tracer Dye. *Journal of Neuroscience* **29,**
9197-9209.

Brecht M. (2007). Barrel cortex and whisker-mediated behaviors. *Curr Opin Neurobiol* **17,** 408-
416.

Canales A, Scheuer KS, Zhao X & Jackson MB. (2022). Unitary synaptic responses of
parvalbumin interneurons evoked by excitatory neurons in the mouse barrel cortex. *Cereb*
Cortex.

Chanda B, Blunck R, Faria LC, Schweizer FE, Mody I & Bezanilla F. (2005). A hybrid approach
to measuring electrical activity in genetically specified neurons. *Nat Neurosci* **8,** 1619-
1626.

Chang PY. (2006). Heterogeneous spatial patterns of long-term potentiation in hippocampal
slices. In *Biophysics*, pp. 144. University of Wisconsin-Madison, Madison, WI.

Das A, Zhu B, Xie Y, Zeng L, Pham AT, Neumann JC, Safrina O, Benavides DR, MacGregor
GR, Schutte SS, Hunt RF & O'Dowd DK. (2021). Interneuron Dysfunction in a New
Mouse Model of SCN1A GEFS. *eNeuro* **8**.

Di Garbo A, Panarese A & Chillemi S. (2004). Phase locking states between fast-spiking
interneurons coupled by electrical and chemical synapses. *Neurocomputing* **58-60,** 159-
164.

Feldmeyer D. (2012). Excitatory neuronal connectivity in the barrel cortex. *Front Neuroanat* **6,**
24.

Feldmeyer D, Qi G, Emmenegger V & Staiger JF. (2018). Inhibitory interneurons and their
circuit motifs in the many layers of the barrel cortex. *Neuroscience* **368,** 132-151.

Ferguson BR & Gao WJ. (2018). PV Interneurons: Critical Regulators of E/I Balance for
Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. *Front Neural Circuits*
12, 37.

Frandolig JE, Matney CJ, Lee K, Kim J, Chevee M, Kim SJ, Bickert AA & Brown SP. (2019).
The Synaptic Organization of Layer 6 Circuits Reveals Inhibition as a Major Output of a
Neocortical Sublamina. *Cell Rep* **28,** 3131-3143 e3135.

Fukuda T & Kosaka T. (2003). Ultrastructural study of gap junctions between dendrites of
parvalbumin-containing gabaergic neurons in various neocortical areas of the adult rat.
Neuroscience **120,** 5-20.

Gonzalez-Burgos G, Cho RY & Lewis DA. (2015). Alterations in cortical network oscillations
and parvalbumin neurons in schizophrenia. *Biol Psychiatry* **77,** 1031-1040.

Guo C, Wang C, He T, Yu B, Li M, Zhao C, Yuan Y & Chen H. (2020). The effect of mGlu2/3 receptors on synaptic activities to different types of GABAergic interneurons in the anterior cingulate cortex. *Neuropharmacology* **175,** 108180.

Holmgren C, Harkany T, Svennenfors B & Zilberter Y. (2003). Pyramidal cell communication within local networks in layer 2/3 of rat neocortex. *J Physiol* **551,** 139-153.

Hu H, Gan J & Jonas P. (2014). Interneurons. Fast-spiking, parvalbumin(+) GABAergic interneurons: from cellular design to microcircuit function. *Science* **345,** 1255263.

Inan M, Blazquez-Llorca L, Merchan-Perez A, Anderson SA, DeFelipe J & Yuste R. (2013). Dense and overlapping innervation of pyramidal neurons by chandelier cells. *J Neurosci* **33,** 1907-1914.

Judak L, Chiovini B, Juhasz G, Palfi D, Mezriczky Z, Szadai Z, Katona G, Szmola B, Ocsai K, Martinecz B, Mihaly A, Denes A, Kerekes B, Szepesi A, Szalay G, Ulbert I, Mucsi Z, Roska B & Rozsa B. (2022). Sharp-wave ripple doublets induce complex dendritic spikes in parvalbumin interneurons in vivo. *Nat Commun* **13,** 6715.

Kayarian FB, Jannati A, Rotenberg A & Santarnecchi E. (2020). Targeting Gamma-Related Pathophysiology in Autism Spectrum Disorder Using Transcranial Electrical Stimulation: Opportunities and Challenges. *Autism Res* **13,** 1051-1071.

Keller D, Ero C & Markram H. (2018). Cell Densities in the Mouse Brain: A Systematic Review. *Front Neuroanat* **12,** 83.

Koelbl C, Helmstaedter M, Lubke J & Feldmeyer D. (2015). A barrel-related interneuron in layer 4 of rat somatosensory cortex with a high intrabarrel connectivity. *Cereb Cortex* **25,** 713-725.

Kondo M, Sumino R & Okado H. (1997). Combinations of AMPA Receptor Subunit Expression in Individual
Cortical Neurons Correlate with Expression of Specific Calcium-
Binding Proteins. *J Neurosci* **17**.

Kooijmans RN, Sierhuis W, Self MW & Roelfsema PR. (2020). A Quantitative Comparison of Inhibitory Interneuron Size and Distribution between Mouse and Macaque V1, Using Calcium-Binding Proteins. *Cereb Cortex Commun* **1,** tgaa068.

Lauber E, Filice F & Schwaller B. (2018). Parvalbumin neurons as a hub in autism spectrum disorders. *J Neurosci Res* **96,** 360-361.

Lenschow C & Brecht M. (2015). Barrel cortex membrane potential dynamics in social touch. *Neuron* **85,** 718-725.

Li P & Huntsman MM. (2014). Two functional inhibitory circuits are comprised of a heterogeneous population of fast-spiking cortical interneurons. *Neuroscience* **265,** 60-71.

Muller JF, Mascagni F & McDonald AJ. (2005). Coupled networks of parvalbumin-immunoreactive interneurons in the rat basolateral amygdala. *J Neurosci* **25,** 7366-7376.

Nahar L, Delacroix BM & Nam HW. (2021). The Role of Parvalbumin Interneurons in Neurotransmitter Balance and Neurological Disease. *Front Psychiatry* **12,** 679960.

Naka A & Adesnik H. (2016). Inhibitory Circuits in Cortical Layer 5. *Front Neural Circuits* **10,** 35.

O'Connor DH, Peron SP, Huber D & Svoboda K. (2010). Neural activity in barrel cortex underlying vibrissa-based object localization in mice. *Neuron* **67,** 1048-1061.

Otsuka T & Kawaguchi Y. (2009). Cortical inhibitory cell types differentially form intralaminar and interlaminar subnetworks with excitatory neurons. *J Neurosci* **29,** 10533-10540.

Packer AM & Yuste R. (2011). Dense, unspecific connectivity of neocortical parvalbumin-positive interneurons: a canonical microcircuit for inhibition? *J Neurosci* **31,** 13260-13271.

Pala A & Petersen CC. (2018). State-dependent cell-type-specific membrane potential dynamics and unitary synaptic inputs in awake mice. *Elife* **7**.

Pervouchine DD, Netoff TI, Rotstein HF, White JA, Cunningham MO, Whittington MA & Kopell NJ. (2006). Low-Dimensional Maps Encoding Dynamics in Entorhinal Cortex and Hippocampus. *Neural Comput* **18**.

Pouille F & Scanziani M. (2001). Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. *Science* **293,** 1159-1163.

Povysheva NV, Gonzalez-Burgos G, Zaitsev AV, Kroner S, Barrionuevo G, Lewis DA & Krimer LS. (2006). Properties of excitatory synaptic responses in fast-spiking interneurons and pyramidal cells from monkey and rat prefrontal cortex. *Cereb Cortex* **16,** 541-552.

Povysheva NV, Zaitsev AV, Rotaru DC, Gonzalez-Burgos G, Lewis DA & Krimer LS. (2008). Parvalbumin-positive basket interneurons in monkey and rat prefrontal cortex. *J Neurophysiol* **100,** 2348-2360.

Scheuer KS, Judge J, Zhao X & Jackson MB. (2023). Velocity of Conduction Between Columns and Layers in Barrel Cortex Reported by Parvalbumin Interneurons. *bioRxiv*, 2022.2007.2027.501767.

Selby L, Zhang C & Sun QQ. (2007). Major defects in neocortical GABAergic inhibitory
circuits in mice lacking the fragile X mental retardation protein. *Neurosci Lett* **412,** 227-
232.

Staiger JF & Petersen CCH. (2021). Neuronal Circuits in Barrel Cortex for Whisker Sensory
Perception. *Physiol Rev* **101,** 353-415.

Stedehouder J, Brizee D, Slotman JA, Pascual-Garcia M, Leyrer ML, Bouwen BL, Dirven CM,
Gao Z, Berson DM, Houtsmuller AB & Kushner SA. (2019). Local axonal morphology
guides the topography of interneuron myelination in mouse and human neocortex. *Elife* **8**.

Tallon-Baudry C, Bertrand O, Peronnet F & Pernier J. (1998). Induced Gamma-Band Activity
during the Delay of a Visual Short-Term Memory Task in Humans. *J Neurosci* **18**.

Thomson AM. (1997). Activity-dependent properties of synaptic transmission at two classes of
connections made by rat enocotical pyramidal axons in vitro. *J Physiol* **502**.

Thomson AM, West DC, Wang Y & Bannister AP. (2002). Synaptic connections and small
circuits involving excitatory and inhibitory neurons in layers 2-5 of adult rat and cat
neocortex: triple intracellular recordings and biocytin labelling in vitro. *Cereb Cortex* **12,**
936-953.

Tremblay R, Lee S & Rudy B. (2016). GABAergic Interneurons in the Neocortex: From Cellular
Properties to Circuits. *Neuron* **91,** 260-292.

Wang D, Zhang Z, Chanda B & Jackson MB. (2010). Improved probes for hybrid voltage sensor
imaging. *Biophys J* **99,** 2355-2365.

Wang Y, Gupta A, Toledo-Rodriguez M, Wu CZ & Markram H. (2002). Anatomical,
physiological, molecular and circuit properties of nest basket cells in the developing
somatosensory cortex. *Cereb Cortex* **12,** 395-410.

West DC, Mercer A, Kirchhecker S, Morris OT & Thomson AM. (2006). Layer 6 cortico-
thalamic pyramidal cells preferentially innervate interneurons and generate facilitating
EPSPs. *Cereb Cortex* **16,** 200-211.

Woolsey TA & Van der Loos H. (1970). The structural organization of layer IV in the
somatosensory region (S1) of mouse cerebral cortex. *Brain Res* **17**.

Xu X & Callaway EM. (2009). Laminar specificity of functional input to distinct types of
inhibitory cortical neurons. *J Neurosci* **29,** 70-85.

Yamasaki M, Fukaya M, Yamazaki M, Azechi H, Natsume R, Abe M, Sakimura K & Watanabe
M. (2016). TARP gamma-2 and gamma-8 Differentially Control AMPAR Density Across
Schaffer Collateral/Commissural Synapses in the Hippocampal CA1 Area. *J Neurosci* **36,**
4296-4312.

608
609 Yang YM, Aitoubah J, Lauer AM, Nuriya M, Takamiya K, Jia Z, May BJ, Huganir RL & Wang
610   LY. (2011). GluA4 is indispensable for driving fast neurotransmission across a high-
611   fidelity central synapse. *J Physiol* **589,** 4209-4227.
612
613 Yarden TS, Mizrahi A & Nelken I. (2022). Context-Dependent Inhibitory Control of Stimulus-
614   Specific Adaptation. *J Neurosci* **42,** 4629-4651.
615
616 Zaitsev AV, Kim KK & Magazanik LG. (2012). The role of calcium-permeable AMPA receptors
617   in disynaptic feedforward inhibition in the rat prefrontal cortex. *Biochemistry (Moscow)*
618   *Supplement Series A: Membrane and Cell Biology* **6,** 198-205.
619
620 Zaitsev AV & Lewis DA. (2013). Functional properties and short-term dynamics of
621   unidirectional and reciprocal synaptic connections between layer 2/3 pyramidal cells and
622   fast-spiking interneurons in juvenile rat prefrontal cortex. *Eur J Neurosci* **38,** 2988-2998.
623
624 Zhou FW & Roper SN. (2014). Reduced chemical and electrical connections of fast-spiking
625   interneurons in experimental cortical dysplasia. *J Neurophysiol* **112,** 1277-1290.
626
627