Multimodal analyses reveal genes driving electrophysiological maturation of neurons in the primate prefrontal cortex

Yu Gao[1,9], Qiping Dong[1,9], Kalpana Hanthanan Arachchilage[1,9], Ryan D. Risgaard[1], Moosa Syed[1], Jie Sheng[1], Danielle K. Schmidt[1], Ting Jin[1], Shuang Liu[1], Sara Knaack[1], Dan Doherty[2], Ian Glass[2], Jon E. Levine[3,4], Daifeng Wang[1,5,6]*, Qiang Chang[1,7,8]*, Xinyu Zhao[1,3]*, André M. M. Sousa[1,3]*

[1]Waisman Center, University of Wisconsin-Madison; Madison, WI, USA

[2]Department of Pediatrics, University of Washington School of Medicine; Seattle, WA, USA

[3]Department of Neuroscience, University of Wisconsin-Madison; Madison, WI, USA

[4]Wisconsin National Primate Research Center, University of Wisconsin-Madison; Madison, WI 53705

[5]Department of Biostatistics and Medical Informatics, University of Wisconsin-Madison; Madison, WI, USA

[6]Department of Computer Sciences, University of Wisconsin-Madison; Madison, WI, USA

[7]Department of Medical Genetics, University of Wisconsin-Madison; Madison, WI, USA

[8]Department of Neurology, University of Wisconsin-Madison; Madison, WI, USA

[9] These authors contributed equally to this work.

*Correspondence: André M. M. Sousa (andre.sousa@wisc.edu), Xinyu Zhao (xinyu.zhao@wisc.edu), Qiang Chang (qchang@waisman.wisc.edu), Daifeng Wang (daifeng.wang@wisc.edu)

SUMMARY

The prefrontal cortex (PFC) is critical for myriad high-cognitive functions and is associated with several neuropsychiatric disorders. Here, using Patch-seq and single-nucleus multiomic analyses, we identified genes and regulatory networks governing the maturation of distinct neuronal populations in the PFC of rhesus macaque. We discovered that specific electrophysiological properties exhibited distinct maturational kinetics and identified key genes underlying these properties. We unveiled that RAPGEF4 is important for the maturation of resting membrane potential and inward sodium current in both macaque and human. We demonstrated that knockdown of *CHD8*, a high-confidence autism risk gene, in human and macaque organotypic slices led to impaired maturation, via downregulation of key genes, including *RAPGEF4*. Restoring the expression of *RAPGEF4* rescued the proper electrophysiological maturation of CHD8-deficient neurons. Our study revealed regulators of neuronal maturation during a critical period of PFC development in primates and implicated such regulators in molecular processes underlying autism.

INTRODUCTION

The dorsolateral prefrontal cortex (dlPFC), a neocortical area highly derived in primates[1-3], is required for higher-order brain functions, including executive functions such as working memory, planning, and decision-making[3-5]. These functions are significantly impaired in brain disorders, including autism spectrum disorders (ASD), schizophrenia, obsessive-compulsive disorder, and neurodegenerative disorders[6-8]. Much of our knowledge on PFC development is largely extrapolated from rodent studies. Although rodents possess a PFC that is homologous to a small portion of the primate PFC, they lack the granular dlPFC[1]. Several comparative studies have

identified gene expression signatures that are unique to the PFC of humans and non-human

primates (NHP)[3,9-12]. Therefore, it is essential to understand the mechanisms regulating dlPFC

development in primates.

The cells that compose the primate dlPFC, especially excitatory and inhibitory neurons,

undergo extensive and dynamic maturation throughout midfetal and late-fetal development, during

which critical neurodevelopmental events, such as circuit assembly and electrophysiological

maturation of neurons occur[13]. Midfetal and late-fetal development are a convergent period of

expression for many ASD genes[14,15], yet the functions of these ASD genes during this period of

development remains unclear. Furthermore, although we have a robust knowledge of the molecular

and cellular mechanisms that govern cell fate specification during cortical development[16-22], and

of the transcriptomic and electrophysiological properties of mature neurons in the adult

neocortex[23-27], our understanding of the gene networks that regulate the early stages of neuronal

maturation, especially electrophysiological maturation, remains elusive. Therefore, a multimodal

investigation of primate dlPFC neurons that combines electrophysiological and functional

genomic analyses during critical periods for neuronal maturation is essential to understand the

mechanisms driving neuronal maturation.

Here, to uncover the molecular mechanisms that regulate the maturation of dlPFC neurons,

we first generated single-nucleus multiomic (snMultiomic) data and performed integrated analyses

of the dlPFC from rhesus macaques (*Macaca mulatta)*, ranging from midfetal to late-fetal periods.

We then performed Patch-seq on acute dlPFC slices from 16 macaques, including the same

specimens profiled for snMultiome. This approach allowed us to integrate our electrophysiological

analyses with gene expression profiles to identify the genes that are important for the maturation

of specific electrophysiological features. Furthermore, we have evaluated the function of select

genes – *RAPGEF4* and *CHD8* – and demonstrated that a loss of function of these genes affects the

morphological and electrophysiological maturation of cortical neurons in organotypic slices of

macaque and human. Finally, Patch-seq analysis of brain slices with *CHD8* knockdown identified

RAPGEF4 as a key mediator for CHD8 regulation of electrophysiological maturation of human

cortical neurons.

RESULTS

Cell type-specific transcriptomic changes underlying the maturation of rhesus macaque

dlPFC cells during midfetal and late-fetal development

We performed snMultiomic analyses on 9 rhesus macaque dlPFC samples across eight prenatal

ages: postconceptional day (PCD) 85, PCD95, PCD100, PCD105, PCD110, PCD125, PCD145,

and two PCD155 (**Figures 1A-1B; Table S1**). After implementing stringent quality control

measures, we retained 76,855 nuclei for further analyses (**Figures S1; Tables S1-S2**). After

removing batch effects, we used unsupervised clustering to group nuclei into transcriptomically

defined groups (**Figures S1B-S1C**). Using this strategy, together with cell type-specific marker

genes[16,23,28], we defined 9 major cell classes that included glutamatergic excitatory neurons (ExN),

GABAergic inhibitory neurons (InN), glial progenitor cells (GPC), astrocytes, oligodendrocyte

precursor cells (OPC), oligodendrocytes, microglia, vascular leptomeningeal cells (VLMC), and

endothelial cells (**Figure S2A**). We re-clustered the ExN and InN nuclei to identify their subclasses

based on the layer (L) and projection identity (intratelencephalic [IT], extratelencephalic [ET],

near-projecting [NP], corticothalamic [CT], and L6B) for ExN, and the developmental origin

(medial ganglionic eminence [MGE], caudal ganglionic eminence [CGE], or dorsal lateral

ganglionic eminence [dLGE]) for InN (**Figures 1C and S2B-S2C**). Interestingly, one small

population of IT neurons exhibited molecular features of both ExN L2-3 IT and ExN L3-5 IT,

indicating that this small population of neurons is yet to be fully specified; we classified this

population as ExN L2-3 IT/L3-5 IT. At the lowest hierarchical level of cell type classification, we

transcriptomically defined 85 cell subtypes (**Figure S2E; Table S3**). By integrating our dataset

with two published snRNA-seq datasets that analyzed early fetal[28] and adult[23] macaque dlPFC, we

observed that our data bridges the embryonic/early fetal cell types with the adult cell types,

facilitating the tracing of the developmental origins of distinct adult cell subtypes (**Figures 1E-**

1G).

As expected, we observed an increasing number of glial cells, including astrocytes, OPCs,

and oligodendrocytes throughout late-fetal development (**Figure 1D**), a period characterized by

minimal cortical neurogenesis and robust gliogenesis[29]. Within the excitatory neurons, we

observed that IT neurons were more abundant than any other subclass, as previously described in

the adult dlPFC[23]. The selective expansion of IT neurons in the primate lineage, especially of

upper-layer (L2-3) neurons[23] and their functional relevance in the cortico-cortical circuits that

govern some of the unique cognitive and behavioral abilities of primates[30], led us to focus our

analysis on the maturational profiles of these IT neuronal populations: ExN L2-3 IT and ExN L3-

5 IT (**Figures 2A-2D**). We performed differential gene expression analysis along the

developmental trajectory (i.e., pseudotime) to identify genes that display dynamic, highly variable

changes during neuronal maturation. Among the top 100 genes (**Table S4**), we observed genes

that exhibit cell type-specific correlation with maturation (**Figures 2C-2D**), including *MEIS2*, a

transcriptional regulator that is induced by retinoic acid and is important for the arealization of the

prefrontal cortex[10], in L2-3 IT, and *FOXP2*, a gene encoding a transcription factor that has recently

been shown to have primate-specific expression in L3-5 IT excitatory neurons[23]. Disruptions in

these genes have been linked to neuropsychiatric disorders[31,32].

Together, our analyses uncovered the transcriptional landscape of primate midfetal and

late-fetal PFC development, highlighting differences between distinct excitatory neuronal

populations throughout maturation.

Chromatin accessibility signatures and gene regulatory networks of macaque midfetal and

late-fetal dlPFC cells

The simultaneous profiling of gene expression and chromatin accessibility in the same cells

allows linking putative regulatory elements to genes, thus providing insights into cell-type gene

regulatory mechanisms in the developing primate brain. After filtering out low-quality nuclei and

implementing other stringent quality control steps, snATAC-seq data from 64,941 nuclei were

retained for downstream analyses (**Figure S1D**). We observed that the clustering of cells based on

accessible chromatin was similar to the clustering utilizing gene expression data (**Figures S1E-**

S1H), showing the congruence between the two datasets (**Figure 3A**). However, the snATAC-seq

data revealed more pronounced differences in the maturation states of cells within each cell

subclass, especially for IT excitatory neurons (**Figure S1E**). To gain a better understanding of the

association of regulatory elements and their target genes during neuronal maturation, we estimated

peak-to-gene links, which identify co-activity between genes and their nearby accessible

chromatin regions, for the excitatory and inhibitory neurons (**Figure S3A**), thus providing a list of

potential regulatory elements for each gene (**Table S5**). We found that peak-gene combinations

were hierarchically divided into three distinct categories: a cluster specific to excitatory neurons

during early midfetal development, a cluster specific to excitatory neurons during late-fetal

development, and a cluster specific to inhibitory neurons during late-fetal development. Functional

enrichment analysis of each group revealed that the clusters specific to late-fetal development were

enriched for terms related to calcium signaling and dendritic development in excitatory neurons,

and synaptic processes in inhibitory neurons, whereas the early-midfetal development cluster was

enriched for transcriptional regulation terms (**Figures S3C-S3E; Table S5**).

Next, since gene expression is regulated by gene regulatory networks (GRNs)[33] and

expression of genes associated with ASD converge in excitatory neurons during these

developmental stages in ExN L2-3 IT and ExN L3-5 IT cell subclasses[14,15], we inferred cell-type

gene regulatory networks (GRNs) for these two cell subclasses using transcriptomic and

epigenomic data together with transcription factor (TF) motif binding sites (**Methods**). These cell-

type GRNs link regulatory elements with TF binding sites to target genes. We particularly focused

on analyzing the subnetworks regulating the genes identified in our developmental maturation

trajectory analysis (**Table S4**) and found nine major TFs regulating the driver genes of the two cell

subclasses. Among these nine TFs, we identified MEIS2 as a key regulator for ExN L2-3 IT,

FOXP2 as a hub TF for ExN L3-5 IT, and MEF2C as an important TF for both ExN subclasses

(**Figure 3B**).

Since upper-layer (L2-3) neurons exhibit selective expansion in the primate lineage[23] and

their roles in the cortico-cortical circuits that are important for the unique cognitive and behavioral

abilities of primates[30], we investigated putative regulatory elements that underlie the

transcriptomic differences between ExN L2-3 IT and ExN L3-5 IT. We calculated differential

accessibility and overrepresented motifs between these neuronal populations throughout midfetal

and late-fetal development. Using this strategy, we identified the MEIS2 motif MA0774.1 as one

of the overrepresented motifs in the ExN L2-3 IT subclass (**Table S6**). Furthermore, we found that

ExN L2-3 IT has both higher *MEIS2* gene expression and a higher MEIS2 motif activity score

when compared to ExN L3-5 IT (**Figure 3C**). Because MEIS2 is a transcription factor enriched in

upper-layer neurons in primates[10,23] and is implicated in neuropsychiatric disorders[32], we

calculated MEIS2 transcription factor footprints across the whole genome in the two cell

subclasses and found that, throughout the analyzed developmental periods, ExN L2-3 IT had a

higher footprint enrichment than ExN L3-5 IT, with the highest enrichment in ExN L2-3 IT during

midfetal development (PCD105-110) and the lowest footprint enrichment in ExN L3-5 IT during

late fetal development (PCD155), indicating the selective involvement of MEIS2 in the regulation

of target genes in ExN L2-3 IT (**Figure 3C**). Indeed, quantitative immunofluorescence showed

that there were significantly fewer MEIS2-expressing cells in L3-5, identified by the expression

of *RORB* (**Figure S3B**), compared to L2-3 of dlPFC at PCD155 (**Figure 3D**).

One of the genes potentially regulated by MEIS2 in IT excitatory neurons is the *Nuclear*

factor I/A (*NFIA*), which has been shown to be regulated by MEIS2 in retinal progenitor cells[34]

and is linked to callosal hypoplasia, likely reflecting malformations of upper-layer excitatory

neurons[35]. By analyzing the distribution of accessible chromatin peaks in the vicinity of *NFIA*

across the two neuronal subclasses and developmental periods, we identified 5 active MEIS2

binding sites within 200kb upstream of the *NFIA* transcription start site (**Figure 3E**). We also

found three peak-to-gene links (peaks that show higher co-activity with *NFIA* gene expression),

with one of the links containing two predicted MEIS2 binding motifs (chr1-163553896-

163555161) approximately 181kb upstream of the *NFIA* transcription starting site, suggesting that

it may be a potential regulatory element for *NFIA*. To validate this finding, we conducted a dual

luciferase assay to assess the activity of the genomic sequence containing the two MEIS2 motifs.

Overexpression of MEIS2 resulted in a significant increase in transcription activity driven by the

genomic sequence. Furthermore, when we introduced mutations into two potential MEIS2 binding motifs, one of these mutations significantly reduced the activity of the genomic sequence (**Figure 3F**). Taken together, our analyses have unveiled key regulators, such as MEIS2, and gene regulatory networks governing the maturation of distinct IT neuronal populations during midfetal and late-fetal primate brain development.

Patch-seq of midfetal and late-fetal macaque dlPFC cells identified transcriptomic signatures defining electrophysiological maturation

The midfetal and late-fetal periods are critical stages in cortical development, marked by notable alterations in neuronal morphology, electrophysiology, and gene expression[13]. Although all the analyzed neurons displayed a relatively immature morphology, PCD155 neurons exhibited significantly greater complexity, with greater total dendritic length, increased branching points, and more endings when compared to neurons from PCD100 brains (**Figures S4B-S4F**). To gain a comprehensive understanding of the molecular mechanisms that drive neuronal maturation, we employed Patch-seq, which allows for sequential electrophysiological and transcriptomic analyses of the same cell[36], in 16 rhesus macaques ranging from PCD85 to PCD155 (**Figures 4A and S4; Table S1**). We used our snRNA-seq data (**Figure 2**) as the reference dataset to infer cell types by mapping the 256 cells profiled by Patch-seq onto the reference UMAP (**Figures 4B and S4I; Table S7**). Approximately 72% of all cells were identified as neurons based on their transcriptomic signature, which included 111 IT neurons (**Table S7**). Our evaluation of the electrophysiological features of three major IT subtypes (L2-3, L3-5, and L6) across different ages revealed that these IT neurons exhibited similar electrophysiological features (**Figures S4J-S4N**) during this developmental window. Therefore, we combined all IT neuron types for downstream analysis.

We employed a pseudotime trajectory analysis to infer a maturation trajectory by

integrating differentially expressed genes across age groups in all identified IT neurons (**Methods**).

Our analysis revealed a trajectory characterized by the progressive maturation of IT neurons from

earlier to later developmental periods (**Figure 4C**). We observed that the proportion of IT neurons

exhibiting action potentials (APs) across six age groups (PCD85-90, PCD95-100, PCD105-110,

PCD125, PCD145, and PCD155) gradually increased from 0% in the PCD85-90 group to 95% in

PCD155, providing strong evidence for electrophysiological maturation (**Figure S4O**). We then

assessed neuronal maturation through the lens of the following features that offer unique

perspectives on maturation: membrane capacitance (C_m), input resistance (R_{in}), resting membrane

potential (RMP), and inward sodium current (I_{Na}). Our analysis revealed that, during early midfetal

development (from PCD85 to PCD100), none of the electrophysiological features showed

significant change (**Figure 4**). We observed a significant increase in C_m from PCD100 to PCD105,

after which the values remained stable (**Figure 4**). This increase in C_m potentially reflects the

enlargement of the cell body and more complex cellular structures during the maturation of

neurons. In contrast, there was a gradual decline in the averaged value of R_{in} after PCD125 (**Figure**

4F). This effect may be attributed to an increase in the number of ion channels in the neuronal

membrane. Moreover, the morphological changes that occur in dendrites (**Figures S4B-S4F**) and

spines during maturation may contribute to alterations in signal processing and integration, further

enhancing neuronal function. We also observed a conspicuous pattern of increasingly negative

RMP from PCD100 to PCD155 (**Figure 4G**), which reflects the developmental maturation in the

ion channel composition of these neurons, allowing for improved electrical signal processing and

transmission. During neuronal development, the current density of I_{Na} amplifies, thereby

increasing the capacity of neurons to generate action potentials. Our results indicate a significant

increase in the magnitude of the I_{Na} from PCD105 to PCD155 (**Figure 4H**). This suggests that IT

neurons in the dlPFC acquire the ability to initiate and transmit APs after PCD105. In summary,

our analyses showed a progression in maturation across four distinct electrophysiological

properties. However, it is noteworthy that the temporal dynamics of the four electrophysiological

properties were not identical. While RMP and I_{Na} exhibited a progressive increase from PCD95-

100 to PCD155, C_m and R_{in} displayed drastic changes only at PCD105-110. Our study has

uncovered distinct profiles in the temporal dynamics of electrophysiological maturation, where

each property exhibits a different rate of maturation.

Similar to IT ExNs, InNs exhibited progressively more negative RMP and I_{Na} from

PCD105-110 to PCD155, whereas R_{in} didn't show any significant differences and C_m became

higher throughout development (**Figures S5A-S5E**). The proportion of InNs with APs increased

from 12.5% in PCD85-90 to 100% in PCD125, PCD145, and PCD155, indicating a faster

electrophysiological maturation than ExNs (**Figure S5F**).

We hypothesized that differential expression of distinct genes drives the divergent trends

of the four electrophysiological features across maturation. By implementing a correlation analysis

between the electrophysiological data and the gene expression levels across all IT neurons, we

identified 135, 163, 193, and 142 genes that are associated with the temporal dynamics of

electrophysiological features of C_m, R_{in}, RMP, and I_{Na}, respectively (**Figure S6A; Table S9**).

Importantly, genes that are not expected to exhibit correlation with electrophysiological features,

including the housekeeping genes *GAPDH* and *TBP*, showed no significant correlation with any

of the electrophysiological features (**Figure S6B**). Several genes, including *CAMK2A* and

MICAL2, exhibited high correlation with all four investigated electrophysiological features

(**Figures 4I-4L**). Other genes, such as *RAPGEF4*, exhibited significant correlation with one or two

of the electrophysiological features. To validate our Patch-seq data, we performed in situ

hybridization on dlPFC sections of PCD100 and PCD155 macaques for select genes that

demonstrated significant correlation with all (*CAMK2A*, *MICAL2*) or some (*RAPGEF4*, *NRGN*)

of the electrophysiological features. Consistent with the Patch-seq results, the expression levels of

all tested genes were significantly higher in PCD155 compared to PCD100 (**Figures 4D and S6C**).

Therefore, our Patch-seq analysis of dlPFC neurons across midfetal and late-fetal

development revealed evidence of neuronal maturation across three distinct aspects:

morphological, transcriptomic, and electrophysiological properties, and identified genes

associated with maturation of specific electrophysiological features.

***RAPGEF4* is crucial for the morphological and electrophysiological maturation of excitatory**

neurons

Our Patch-seq analysis demonstrated a marked upregulation of *RAPGEF4* expression in IT

neurons during cortical development (**Figures 4K-4L**), which was also corroborated by our

snRNA-seq data (**Figure 5A**). RAPGEF4 (Rap Guanine Nucleotide Exchange Factor 4), also

known as EPAC2 (exchange protein directly activated by cyclic AMP 2), is a regulator of small

GTPase signaling and a protein kinase A (PKA)-independent cAMP target[37-39]. We found that

RAPGEF4 expression exhibited a high degree of correlation with various electrophysiological

features, with the highest correlation observed with I_{Na} (r = -0.57) and the second highest

correlation with RMP (r = -0.47) (**Figure S6E**). To confirm that RAPGEF4 is required for neuronal

maturation, we infected ex vivo macaque brain slices with lentivirus (LV) expressing mScarlet

and shRNA against *RAPGEF4* (LV-sh*RAPGEF4*) together with LV expressing mNeonGreen and

control shRNA (LV-sh*NC*) (**Figure 5B**). The neurons with *RAPGEF4* knockdown (sh*RAPGEF4*,

mScarlet$^+$) exhibited reduced dendritic complexity, total dendritic length, branching points, and

number of endings compared to controls (mNeonGreen$^+$mScarlet$^-$) (**Figures 5C-5H**). In addition,

LV-sh*RAPGEF4*-infected neurons exhibited smaller I_{Na} (**Figures 5I and 5M**) and less negative

RMP (**Figures 5I and 5L**), without significant changes in C_m and R_{in}, compared to control neurons

(**Figures 5J-5K; Table S10**). These results indicate that a deficiency of *RAPGEF4* in neurons

leads to delayed maturation of I_{Na} and RMP, which is consistent with our finding that *RAPGEF4*

had a strong correlation with I_{Na} and RMP (**Figures 4K-4L and S6E**).

RAPGEF4 is enriched in neurons compared to other cell types[39]. The levels of *RAPGEF4*

increase in the human cortex throughout development, compared to other brain regions or

organs[9,40]. Based on our observations in macaque, we hypothesized that *RAPGEF4* expression is

also associated with the maturation of electrophysiological properties in human. We infected

human midfetal cortical brain slices with LV-sh*RAPGEF4* and performed patch-clamp recording.

As in macaque, we observed reduced I_{Na} and RMP, but not C_m and R_{in}, in LV-sh*RAPGEF4*-

infected compared to LV-sh*NC*-infected neurons (**Figures 5N-5R; Table S10**). To confirm that

the observed effect was not caused by off-target effects of the shRNA, we replicated the

experiment and observed the same results using a different shRNA targeting *RAPGEF4* in human

neurons (**Figures S6H-S6L**). The role of RAPGEF4 in regulating RMP and I_{Na} maturation in both

macaque and human neurons is consistent with the functional enrichment for "regulation of

GTPase activity" in genes driving neuronal maturation (**Figure S6D**). These data validate our

approach to identify genes that drive the maturation of specific electrophysiological features and

reveal the novel role of *RAPGEF4* in morphological and electrophysiological maturation of

neurons in both macaque and human cortical development.

CHD8 knockdown leads to impaired maturation of human cortical excitatory neurons

Recent advances in genomics have revealed important roles of chromatin regulators during

midfetal PFC development and implicated them in neurodevelopmental disorders[41,42]. CHD8

(Chromodomain Helicase DNA-binding protein 8) is a chromatin remodeling protein that plays a

crucial role in regulating gene expression during brain development and has been linked to ASD[42-

44]. To investigate the role of *CHD8* during human midfetal cortical development, a critical

developmental period for ASD[14,15], we knocked down *CHD8* expression in ex vivo human

midfetal brain slices using LV expressing sh*CHD8* and mScarlet together with LV expressing

shNC and mNeonGreen (**Figures 6A and S7B-S7C**). LV-sh*CHD8*-infected neurons displayed

significantly shorter dendrites compared to control neurons (**Figures 6B, 6D-6E and S7D-S7F**),

suggesting that a deficiency in *CHD8* disrupts morphological maturation of cortical neurons during

midfetal development, which is consistent with previous findings in mice[44].

We then performed Patch-seq on 156 cells from slices that contained both *CHD8*-

knockdown and control cells (**Figures 6A and S7G**). To determine the cellular identity of recorded

cells, we compared their transcriptome with the reference snRNA-seq data and assigned each

queried cell a specific cell type (**Figures 6C and S7H; Table S11**). Out of the 156 cells examined,

there were 55 excitatory neurons infected with LV-sh*NC* and 42 infected with LV-sh*CHD8* (**Table

S11**). LV-sh*CHD8*-infected excitatory neurons had less negative RMP and a significant increase

in R_{in} (**Figures 6I and 6H**) but not in C_m and I_{Na} (**Figures 6G, 6J and S7I**), compared to the

controls. To rule out the possibility that the observed effect is due to shRNA off-target effects, we

repeated the experiment with an alternative shRNA in human cultured neurons and observed

similar deficits in both R_{in} and RMP (**Figures S7J-S7N**). Therefore, *CHD8* knockdown produces

a discernible impact on the electrophysiological properties of midfetal human neurons, which is

consistent with its role during early development[43,44].

RAPGEF4 rescues electrophysiological deficits of CHD8-deficient human cortical neurons

To determine which genes might underlie the observed differences in R_{in} and RMP maturation

caused by *CHD8* knockdown, we identified the differentially expressed genes between LV-

sh*CHD8* and LV-sh*NC* and intersected this list with the genes that were associated with the

maturation of these two electrophysiological features (**Table S9**). We found 10 genes associated

with R_{in}, 11 genes associated with RMP, and 20 genes associated with both (**Figure 7A**). These

included genes that are associated with the overall maturation of neurons, such as *NRGN* and

CACNA1I, the R_{in}-associated *RGS6*, a GTPase regulator important for activity-dependent

morphological and electrical maturation[45], and *RAPGEF4*, which we found to significantly impact

the maturation of RMP (**Figures 7B-7E and 5**).

To determine whether reduced RAPGEF4 levels might contribute to the observed reduced

RMP in *CHD8* deficient neurons, we expressed *RAPGEF4* in human cortical neurons that were

knockdown for *CHD8* (sh*CHD8*) (**Figure 7F**). Exogenously expressed *RAPGEF4* restored normal

maturation of both RMP and I_{Na}, with no significant effect on R_{in} or C_m, in *CHD8*-deficient neurons

(**Figures 7G-7L; ; Table S12**). Therefore, *CHD8* deficiency in primate cortical neurons leads to

reduced expression of genes important for electrophysiological maturation, including *RAPGEF4*,

which may contribute to neuronal maturation deficits.

DISCUSSION

In this study, we employed multimodal approaches including snMultiome and Patch-seq to interrogate mechanisms driving neuronal maturation in the primate dlPFC during midfetal and late-fetal development. We identified dynamic gene expression changes across cell types in their maturation trajectories as well as critical developmental periods for the maturation of specific electrophysiological properties in excitatory and inhibitory neurons.

We confirmed that, based on their molecular profiles, excitatory neurons gain their laminar and projection identities earlier than interneuronal final specification. This corroborates previous findings that pointed to the relevance of extrinsic factors to the final specification of interneurons during perinatal development[20,45]. We focused our analyses on the IT excitatory neurons due to their relevance both to primate brain evolution and their association to neuropsychiatric disorders. Our integrative analysis of gene expression and chromatin accessibility data unveiled gene regulatory networks associated with neuron maturation of both excitatory and inhibitory neurons and highlighted MEIS2 as a key transcriptional factor for L2-3 IT neuronal development and maturation.

By integrating electrophysiology, morphology, and functional genomics analyses of primate dlPFC development, we discovered that different electrophysiological properties mature at different rates. While RMP change gradually and linearly during maturation, C_m and R_{in} exhibit a jump at the end of neurogenesis during midfetal developmental, I_{Na} exhibited significant maturation only at the late-fetal period, which coincides with a significant increase in the number of neurons able to fire action potentials. The distinct rate of maturation among electrophysiological properties reveals the intricate and finely tuned regulation of neuronal maturation during midfetal and late-fetal development of primate PFC.

Our Patch-seq approach allowed us to investigate the genes associated with the maturation

of specific electrophysiological features. We have uncovered genes, including kinases, small GTP

signaling molecules, neurotransmitter receptors, ion channels, and transporters that are associated

with maturation of specific electrophysiological features that likely underlie the differences we

observed in their maturation kinetics. One of these genes is *RAPGEF4*, which we have identified

as being significantly correlated with the maturation of resting membrane potentials and inward

sodium currents. RAPGEF4, a guanine exchange factor in small GTPase pathway, is a direct target

of cAMP and has been shown to regulate dendritic spine remodeling in mice and its deficiency in

mice leads to learning and memory deficits[39,46]. However, its role in primate brain development or

human disease is unknown. Dysregulation of the cAMP pathway has been implicated in a wide

range of neurological disorders and a potential treatment to elevate cAMP levels is currently in

clinical trial for fragile X syndrome[47-49]. The small GTPase pathway has been shown to have

critical roles during maturation of human neurons[50]. Our study demonstrates that RAPGEF4 may

be one of the small GTPase regulators driving neuronal maturation during primate brain

development, especially for RMP, an electrophysiological feature regulated by GTPase signaling.

Finally, to demonstrate the potential of our experimental platform in exploring the

developmental origin of human diseases, we showed that knocking down of a high-confidence

ASD gene – *CHD8* – impaired morphological and electrophysiological maturation of excitatory

neurons in human organotypic slices. This impairment was driven by the downregulation of the

expression of several key genes that we identified to be relevant for the electrophysiological

maturation of neurons, including *RAPGEF4 and RGS6*, two regulators of small GTPase signaling

critical for neuronal maturation[50]. By restoring the expression of *RAPGEF4* in CHD8-deficient

neurons we were able to rescue the deficits in electrophysiological maturation of these neurons.

The chromatin regulator CHD8 is expressed at high levels during early-fetal and midfetal

development and the levels of CHD8 do not exhibit significant upregulation during midfetal to

late-fetal neuronal maturation. How CHD8 and other early expressing chromatin regulators

converging at midfetal development contribute to ASD remains unclear. Our finding sheds light

on how a chromatin remodeler may regulate the physiological maturation of neurons, thus

impairing their normal functioning.

Together, our study has revealed key genes and gene networks driving neuronal maturation

during a critical period of prefrontal development in primates and linked the function of a novel

regulator of the maturation of specific electrophysiological features with a high-confidence ASD

risk gene. Our results demonstrate the power of our multimodal approach and its application for

future studies aiming at understanding how genes regulate neuronal maturation and its implication

in brain disorders. Finally, we summarized our results including cell clusters, genes and networks

in an open-access, searchable web app, which will allow researchers to investigate the role of

different genes on neuronal maturation.

ACKNOWLEDGEMENTS
411

We thank Y. Xing, S. Krebsbach, D. Phan, M. Weidenfeller, and H. Thurston for technical

assistance; K. Knobel at the Waisman IDD Model Core for core services; Dr. Heather Simmons,

Director of Pathology Services (WNPRC) for non-human primate tissue acquisition; and the Birth

Defects Research Laboratory (BDRL) at the University of Washington for human tissue

acquisition. This work was supported by the National Institutes of Health (R01MH118827 and

R01NS105200 to X.Z.; R01HD064743 to Q.C; R01NS064025, R01AG067025, and

RF1MH128695 to D.W; 1R01HD106197 and UM1MH130991 to A.M.M.S; P51 OD011106 to

WNPRC; P50HD105353 to Waisman Center; and R24HD000836 to BDRL). Further support was

provided by the DOD IIRA grant (X.Z.), SFARI pilot grant (X.Z., A.M.M.S., Q.C., and D.W.),

Jenni and Kyle Professorship (X.Z.), Brain and Behavior Research Foundation (29721 to

A.M.M.S.), Brain Research Foundation (BRFSG-2023-11 to A.M.M.S.), and the Medical Scientist

Training Program T32 (GM140935 to R.D.R.).

AUTHOR CONTRIBUTIONS

426 Y.G., Q.D., Q.C., X.Z., and A.M.M.S. conceived and designed the study. Y.G., Q.D., R.D.R.,

427 M.S., D.K.S., and A.M.M.S. performed the experiments. K.H.A., J.S., T.J., S.L., S.K., and D.W.

428 analyzed the genomics data. J.E.L., D.D., I.G. provided the analyzed specimens. Y.G., Q.D.,

429 K.H.A., Q.C., X.Z., and A.M.M.S. wrote the manuscript. All authors edited the manuscript.

DECLARATION OF INTERESTS

432 The authors declare no competing interests.

RESOURCE AVAILABILITY

435 The genomic data has been submitted to GEO (GSE235493). The data can be interactively

436 visualized at https://daifengwanglab.shinyapps.io/DevMacaquePFC/. The code and data used for

437 generating figures can be accessed at Zenodo: https://zenodo.org/records/11123264. All other data

438 are available in the main article or supplemental information. Further information and requests for

439 resources should be directed to the corresponding authors.

Figure 1. Single-nucleus transcriptomic atlas of rhesus macaque dlPFC cells across midfetal

and late-fetal development

(A) The frontal cortex of Rhesus macaque (*Macaca mulatta*) brains, aged from postconceptional

day (PCD) 85 to 155, were dissected and sectioned at 300 μm. Acute dorsolateral prefrontal

cortices were analyzed using Patch-seq and single-nucleus multiome (snRNA-seq and snATAC-

seq). Tissue from the same specimens, as well as human midfetal tissue, was also used to culture

organotypic slices for functional analyses of select genes (*RAPGEF4* and *CHD8*). Validation of

key findings was also performed by multiplexed immunostaining and single-molecule fluorescent

in situ hybridization on dlPFC sections from the contralateral hemisphere of the same specimens

(**Table S1**). Finally, we have performed an integrative analysis of the generated multimodal

dataset.

(B-C) UMAP visualization of all the cell classes and subclasses. Cells are colored according to

age in postconceptional days (PCD), from PCD85-PCD155 (B) cell subclass (C).

(D) Bar plot showing the cell composition of the dlPFC throughout midfetal and late-fetal

development.

(E-G) Integrated UMAP visualization depicting cells obtained from both the current study and

published datasets[22,27]. Cells are color-coded based on their respective dataset (current study in

medium blue) (E), age of donor specimen (embryonic/early fetal in blue, current study in orange,

adult in dark green) (F), and cell type (G). ExN, excitatory neuron; InN, inhibitory neurons; Astro,

astrocytes; Oligo, oligodendrocytes; IPC, intermediate precursor cells; enIPC, excitatory neuron

IPC; inIPC inhibitory neuron IPC; oIPC, oligodendrocyte IPC; OPC, oligodendrocyte precursor

cells; GPC, glial precursor cells; Endo, endothelial cells; RG, radial glia cells; oRG, outer radial

glia cells; tRG, truncated radial glia cells; vRG, ventricular radial glia cells; NESC, neuroepithelial

stem cells; VLMC, vascular leptomeningeal cells; AntVen domain cell, antero-ventral domain

cells; NA, unknown cell type.

Figure 2. Maturational trajectories of L2-3 and L3-5 intratelencephalic-projecting neurons.

(A-B) UMAP visualization of the maturation trajectory of ExN L2-3 IT (A), and ExN L3-5 IT (B) cells; the top panel is colored according to postconceptional days, whereas the bottom panel is colored according to inferred pseudotime.

(C-D) Gene expression variation with pseudotime of genes specific to ExN L2-3 IT (C), ExN L3-5 IT (D) (from the top 100 highly variable genes along the developmental trajectory).

Figure 3. Single-nucleus multiome gene expression and chromatin accessibility of rhesus macaque dlPFC cells across mid- to late-fetal development.

(A) Joint (snRNA-seq and snATAC-seq) UMAP visualization of cell subclasses.

(B) Differential gene regulatory networks inferred for ExN L2-3 IT and ExN L3-5 IT cell

subclasses. Hexagonal nodes indicate transcription factors, and the elliptical nodes indicate the

target genes. The edge colors represent the cell subclass specificity of transcription factor-target

gene relationships (black, conserved in both cell subclasses; red, ExN L2-3 IT specific; blue, ExN

L3-5 IT specific).

(C) UMAP visualization of *MEIS2* expression and motif activity scores for MA0774.1 (MEIS2

transcription factor motif) in L2-3 IT and L3-5 IT cells (top panels); MEIS2 transcription factor

footprints for ExN L2-3 IT and ExN L3-5 IT cells across four developmental periods (bottom

panel).

(D) Immunofluorescent detection of MEIS2 in the macaque dlPFC at PCD155, in L2-3 IT and L3-

5 IT cells shows that a significantly higher number of cells express MEIS2 in L2-3 IT than in L3-

5 IT cells (L3-5 was identified using in situ hybridization against *RORB*, as shown in Figure S3E).

Scale bar, 20 μm. (**$p < 0.01$, *t*-test).

(E) Peak-to-gene link plot for *NFIA* together with the gene expression of *MEIS2* and *NFIA* in ExN

L2-3 IT and ExN L3-5 IT across four developmental periods. The links indicate the co-activity

links between *NFIA* gene expression and chromatin accessibility peaks. The highlighted regions

within the chromatin accessibility peaks show the MEIS2 transcription factor motif (MA0774.1)

binding sites.

(F) Dual-Luciferase reporter assay of *NFIA* genomic sequences containing MEIS2 motifs. OE,

overexpression. *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$ (one-way analysis of variance with Tukey

multiple comparisons). Each dot represents one replicate.

Figure 4. Identification of midfetal and late-fetal neuronal maturation-related genes using

Patch-seq.

(A) Schematic representation of the Patch-seq experimental design for analyzing midfetal and late-

fetal rhesus macaque dlPFC cells.

(B) Integration of the 256 cells profiled by Patch-seq onto the snRNA-seq reference UMAP.

(C) Pseudotime trajectory based on the gene expression profiles of patched IT neurons from 16

individual rhesus macaque fetal brains at 9 distinct fetal ages, with representative

electrophysiological sample traces (voltage responses to depolarizing and hyperpolarizing current

injections, and inward sodium currents).

(D) In situ hybridization detection and quantification (right panel) of *CAMK2A*, *MICAL2*, and

RAPGEF4 in PCD100 and PCD155 dlPFC. Scale bar, 20 μm. (****$p < 0.0001$, *t*-test).

(E-L) Plots illustrating the developmental changes in electrophysiological properties, including

membrane capacitance (C_m, E), input resistance (R_{in}, F), resting membrane potential (RMP, G),

and inward sodium current (I_{Na}, H) of IT neurons across six different age groups. Bottom panels

show the expression levels of C_m-correlated genes (I), R_{in}-correlated genes (J), RMP-correlated

genes (K), and I_{Na}-correlated genes (L) in IT neurons across the same age groups. (**$p < 0.01$,

***$p < 0.001$, ****$p < 0.0001$, one-way ANOVA, Tukey's multiple comparisons test).

Figure 5. Knockdown of *RAPGEF4* impairs morphological and electrophysiological maturation of cortical neurons in macaque organotypic dlPFC slices.

(A) Violin plots showing the expression of *RAPGEF4* in IT neurons from PCD85 to PCD155.

(B) Schematic of the experimental setup for investigating the morphological and electrophysiological alterations after *RAPGEF4* knockdown in ex vivo dlPFC slices.

(C) Representative image depicting *RAPGEF4* knockdown (both mScarlet[+][red] and mScarlet[+]/mNeon[+][yellow]) and control (mNeon[+][green only]) neurons in macaque dlPFC slices. Scale bar, 100 μm.

(D) Sample traces depicting the morphology of both *RAPGEF4* knockdown (bottom panel) and

control (top panel) neurons.

(E) Sholl analysis of dendritic complexity in *RAPGEF4* knockdown neurons compared to control

neurons [$F(1,150) = 14.531$, ***$p < 0.001$, multivariate analysis of variance].

(F-H) Knockdown of *RAPGEF4* decreases total dendritic length (F), number of nodes (G), and

number of endings (H) (*t*-test, ***$p < 0.001$, **$p < 0.01$).

(I) Representative traces of inward sodium currents and voltage responses to depolarizing and

hyperpolarizing current injections for macaque *RAPGEF4* knockdown (right panel) and control

(left panel) neurons.

(J-M) Knockdown of *RAPGEF4* in macaque organotypic slices leads to reduced I_{Na} (M) and less

negative RMP (L), without significantly change in C_m (J) and R_{in} (K) (*t*-test, ***$p < 0.001$, *$p <$

0.05). Each dot represents one patched cell. All cells were collected from four individual fetal

brains, with each fetal brain indicated by a different color: black, dark gray, medium gray, and

light gray.

(N) Representative traces of inward sodium currents and voltage responses to depolarizing and

hyperpolarizing current injections for human *RAPGEF4* knockdown (right panel) and control (left

panel) neurons.

(O-R) Knockdown of *RAPGEF4* in human organotypic slices leads to reduced I_{Na} (R) and less

negative RMP (Q), without significantly change in C_m (O) and R_{in} (P) (*$p < 0.05$). Each dot

represents one patched cell. All cells were collected from four individual fetal brains, with each

fetal brain indicated by a different color: black, dark gray, medium gray, and light gray.

Figure 6. Knockdown of *CHD8* impairs maturation of cortical excitatory neurons.

(A) Schematic diagram illustrating the experimental setup used to study the morphological and electrophysiological changes after knockdown of *CHD8* in ex vivo human fetal cortical slices.

(B) Representative image showing *CHD8* knockdown (mScarlet[+][red], mScarlet[+] and mNeon[+][yellow]) and control (mNeon[+][green]) neurons in human cortical slices. Scale bar, 100 μm.

(C) Integration of the 156 cells profiled by Patch-seq onto the snRNA-seq reference UMAP.

(D) Sample traces showing the morphology of both *CHD8* knockdown and control neurons.

(E) *CHD8* knockdown reduces total dendritic length (**p < 0.01, *t*-test).

(F) Representative traces (voltage response to depolarizing [50 pA] and hyperpolarizing [-50 pA] current injections) for control neurons and *CHD8* knockdown.

570 (G-J) *CHD8* knockdown increases R_{in} (H) and RMP (I) but does not significantly alter C_m (G) and

571 I_{Na} (J) (*p < 0.05, **p < 0.01, *t*-test).

Figure 7. Exogenously expressed RAPGEF rescues neuronal maturation deficits resulting

from CHD8 deficiency.

(A) Volcano plot depiction (-$\log_{10}(P_{adj})$) vs $\log_2$(*fold-change*)) of differentially expressed genes

between LV-sh*NC* and LV-sh*CHD8* -infected excitatory neurons. Genes with expression profiles

that significantly correlate with the RMP (green), R_{in} (purple), or both (red) electrophysiology features are labeled.

(B-E) Violin plots showing the expression levels of *NRGN* (B), *CACNA1I* (C), *RAPGEF4* (D), and *RGS6* (E), highlighting the effect of *CHD8* knockdown on these neuronal maturation genes (Figures 4I-4L).

(F) Schematic of lentiviral vectors used for knocking down *CHD8* and overexpression of *RAPGEF4* in human cortical ex vivo organotypic slices.

(G) Representative traces (voltage response to depolarizing [50 pA] and hyperpolarizing [-50 pA] current injections) for neurons with *CHD8* knockdown and neurons with *CHD8* knockdown and *RAPGEF4* overexpression.

(H) Representative traces of inward sodium currents to depolarizing current injections for neurons with *CHD8* knockdown and neurons with *CHD8* knockdown and *RAPGEF4* overexpression.

(I-L) *RAPGEF4* rescues RMP (K) and I_{Na} (L) but does not significantly alter C_m (I) and R_{in} (J) of neurons with *CHD8* knockdown (*$p < 0.05$, **$p < 0.01$, *t*-test).

REFERENCES

1. Preuss, T.M., and Wise, S.P. (2022). Evolution of prefrontal cortex. Neuropsychopharmacology *47*, 3-19. 10.1038/s41386-021-01076-5.
2. Smaers, J.B., Gomez-Robles, A., Parks, A.N., and Sherwood, C.C. (2017). Exceptional Evolutionary Expansion of Prefrontal Cortex in Great Apes and Humans. Curr Biol *27*, 1549. 10.1016/j.cub.2017.05.015.
3. Sousa, A.M.M., Meyer, K.A., Santpere, G., Gulden, F.O., and Sestan, N. (2017). Evolution of the Human Nervous System Function, Structure, and Development. Cell *170*, 226-247. 10.1016/j.cell.2017.06.036.
4. Levy, R., and Goldman-Rakic, P.S. (1999). Association of storage and processing functions in the dorsolateral prefrontal cortex of the nonhuman primate. J Neurosci *19*, 5149-5158. 10.1523/JNEUROSCI.19-12-05149.1999.
5. Kritzer, M.F., and Goldman-Rakic, P.S. (1995). Intrinsic circuit organization of the major layers and sublayers of the dorsolateral prefrontal cortex in the rhesus monkey. J Comp Neurol *359*, 131-143. 10.1002/cne.903590109.

608   6.   Lewis, D.A., and Mirnics, K. (2006). Transcriptome alterations in schizophrenia:
609   disturbing the functional architecture of the dorsolateral prefrontal cortex. Prog Brain Res
610   *158*, 141-152. 10.1016/S0079-6123(06)58007-0.
611   7.   Cools, R., and Arnsten, A.F.T. (2022). Neuromodulation of prefrontal cortex cognitive
612   function in primates: the powerful roles of monoamines and acetylcholine.
613   Neuropsychopharmacology *47*, 309-328. 10.1038/s41386-021-01100-8.
614   8.   Pattabiraman, K., Muchnik, S.K., and Sestan, N. (2020). The evolution of the human brain
615   and disease susceptibility. Curr Opin Genet Dev *65*, 91-97. 10.1016/j.gde.2020.05.004.
616   9.   Zhu, Y., Sousa, A.M.M., Gao, T., Skarica, M., Li, M., Santpere, G., Esteller-Cucala, P.,
617   Juan, D., Ferrandez-Peral, L., Gulden, F.O., et al. (2018). Spatiotemporal transcriptomic
618   divergence across human and macaque brain development. Science *362*.
619   10.1126/science.aat8077.
620   10.   Shibata, M., Pattabiraman, K., Lorente-Galdos, B., Andrijevic, D., Kim, S.K., Kaur, N.,
621   Muchnik, S.K., Xing, X., Santpere, G., Sousa, A.M.M., and Sestan, N. (2021). Regulation
622   of prefrontal patterning and connectivity by retinoic acid. Nature *598*, 483-488.
623   10.1038/s41586-021-03953-x.
624   11.   Shibata, M., Pattabiraman, K., Muchnik, S.K., Kaur, N., Morozov, Y.M., Cheng, X.,
625   Waxman, S.G., and Sestan, N. (2021). Hominini-specific regulation of CBLN2 increases
626   prefrontal spinogenesis. Nature *598*, 489-494. 10.1038/s41586-021-03952-y.
627   12.   Pletikos, M., Sousa, A.M., Sedmak, G., Meyer, K.A., Zhu, Y., Cheng, F., Li, M.,
628   Kawasawa, Y.I., and Sestan, N. (2014). Temporal specification and bilaterality of human
629   neocortical topographic gene expression. Neuron *81*, 321-332.
630   10.1016/j.neuron.2013.11.018.
631   13.   Silbereis, J.C., Pochareddy, S., Zhu, Y., Li, M., and Sestan, N. (2016). The Cellular and
632   Molecular Landscapes of the Developing Human Central Nervous System. Neuron *89*,
633   248-268. 10.1016/j.neuron.2015.12.008.
634   14.   Willsey, A.J., Sanders, S.J., Li, M., Dong, S., Tebbenkamp, A.T., Muhle, R.A., Reilly,
635   S.K., Lin, L., Fertuzinhos, S., Miller, J.A., et al. (2013). Coexpression networks implicate
636   human midfetal deep cortical projection neurons in the pathogenesis of autism. Cell *155*,
637   997-1007. 10.1016/j.cell.2013.10.020.
638   15.   Parikshak, N.N., Luo, R., Zhang, A., Won, H., Lowe, J.K., Chandran, V., Horvath, S., and
639   Geschwind, D.H. (2013). Integrative functional genomic analyses implicate specific
640   molecular pathways and circuits in autism. Cell *155*, 1008-1021.
641   10.1016/j.cell.2013.10.031.
642   16.   Schmitz, M.T., Sandoval, K., Chen, C.P., Mostajo-Radji, M.A., Seeley, W.W.,
643   Nowakowski, T.J., Ye, C.J., Paredes, M.F., and Pollen, A.A. (2022). The development and
644   evolution of inhibitory neurons in primate cerebrum. Nature *603*, 871-877.
645   10.1038/s41586-022-04510-w.
646   17.   Nowakowski, T.J., Bhaduri, A., Pollen, A.A., Alvarado, B., Mostajo-Radji, M.A., Di Lullo,
647   E., Haeussler, M., Sandoval-Espinosa, C., Liu, S.J., Velmeshev, D., et al. (2017).
648   Spatiotemporal gene expression trajectories reveal developmental hierarchies of the human
649   cortex. Science *358*, 1318-1323. 10.1126/science.aap8809.
650   18.   Bhaduri, A., Sandoval-Espinosa, C., Otero-Garcia, M., Oh, I., Yin, R., Eze, U.C.,
651   Nowakowski, T.J., and Kriegstein, A.R. (2021). An atlas of cortical arealization identifies
652   dynamic molecular signatures. Nature *598*, 200-204. 10.1038/s41586-021-03910-8.

653   19.   Johnson, M.B., and Walsh, C.A. (2017). Cerebral cortical neuron diversity and
654   development at single-cell resolution. Curr Opin Neurobiol *42*, 9-16.
655   10.1016/j.conb.2016.11.001.
656   20.   Lim, L., Mi, D., Llorca, A., and Marin, O. (2018). Development and Functional
657   Diversification of Cortical Interneurons. Neuron *100*, 294-313.
658   10.1016/j.neuron.2018.10.009.
659   21.   Bandler, R.C., Mayer, C., and Fishell, G. (2017). Cortical interneuron specification: the
660   juncture of genes, time and geometry. Curr Opin Neurobiol *42*, 17-24.
661   10.1016/j.conb.2016.10.003.
662   22.   Molyneaux, B.J., Arlotta, P., Menezes, J.R.L., and Macklis, J.D. (2007). Neuronal subtype
663   specification in the cerebral cortex. Nat Rev Neurosci *8*, 427-437. Doi 10.1038/Nrn2151.
664   23.   Ma, S., Skarica, M., Li, Q., Xu, C., Risgaard, R.D., Tebbenkamp, A.T.N., Mato-Blanco,
665   X., Kovner, R., Krsnik, Z., de Martin, X., et al. (2022). Molecular and cellular evolution
666   of the primate dorsolateral prefrontal cortex. Science *377*, eabo7257.
667   10.1126/science.abo7257.
668   24.   Berg, J., Sorensen, S.A., Ting, J.T., Miller, J.A., Chartrand, T., Buchin, A., Bakken, T.E.,
669   Budzillo, A., Dee, N., Ding, S.L., et al. (2021). Human neocortical expansion involves
670   glutamatergic neuron diversification. Nature *598*, 151-158. 10.1038/s41586-021-03813-8.
671   25.   Krienen, F.M., Goldman, M., Zhang, Q., R, C.H.D.R., Florio, M., Machold, R., Saunders,
672   A., Levandowski, K., Zaniewski, H., Schuman, B., et al. (2020). Innovations present in the
673   primate interneuron repertoire. Nature *586*, 262-269. 10.1038/s41586-020-2781-z.
674   26.   Beaulieu-Laroche, L., Brown, N.J., Hansen, M., Toloza, E.H.S., Sharma, J., Williams,
675   Z.M., Frosch, M.P., Cosgrove, G.R., Cash, S.S., and Harnett, M.T. (2021). Allometric rules
676   for mammalian cortical layer 5 neuron biophysics. Nature *600*, 274-278. 10.1038/s41586-
677   021-04072-3.
678   27.   Caglayan, E., Ayhan, F., Liu, Y., Vollmer, R.M., Oh, E., Sherwood, C.C., Preuss, T.M.,
679   Yi, S.V., and Konopka, G. (2023). Molecular features driving cellular complexity of human
680   brain evolution. Nature *620*, 145-153. 10.1038/s41586-023-06338-4.
681   28.   Micali, N., Ma, S., Li, M., Kim, S.K., Mato-Blanco, X., Sindhu, S.K., Arellano, J.I., Gao,
682   T., Shibata, M., Gobeske, K.T., et al. (2023). Molecular programs of regional specification
683   and neural stem cell fate progression in macaque telencephalon. Science *382*, eadf3786.
684   10.1126/science.adf3786.
685   29.   Rash, B.G., Duque, A., Morozov, Y.M., Arellano, J.I., Micali, N., and Rakic, P. (2019).
686   Gliogenesis in the outer subventricular zone promotes enlargement and gyrification of the
687   primate cerebrum. Proceedings of the National Academy of Sciences of the United States
688   of America *116*, 7089-7094. 10.1073/pnas.1822169116.
689   30.   Luria, V., Ma, S., Shibata, M., Pattabiraman, K., and Sestan, N. (2023). Molecular and
690   cellular mechanisms of human cortical connectivity. Curr Opin Neurobiol *80*, 102699.
691   10.1016/j.conb.2023.102699.
692   31.   Li, M., Santpere, G., Imamura Kawasawa, Y., Evgrafov, O.V., Gulden, F.O., Pochareddy,
693   S., Sunkin, S.M., Li, Z., Shin, Y., Zhu, Y., et al. (2018). Integrative functional genomic
694   analysis of human brain development and neuropsychiatric risks. Science *362*.
695   10.1126/science.aat7615.
696   32.   Erdogan, F., Ullmann, R., Chen, W., Schubert, M., Adolph, S., Hultschig, C., Kalscheuer,
697   V., Ropers, H.H., Spaich, C., and Tzschach, A. (2007). Characterization of a 5.3 Mb
698   deletion in 15q14 by comparative genomic hybridization using a whole genome "tiling

path" BAC array in a girl with heart defect, cleft palate, and developmental delay. Am J Med Genet A *143A*, 172-178. 10.1002/ajmg.a.31541.

33. Jin, T., Rehani, P., Ying, M., Huang, J., Liu, S., Roussos, P., and Wang, D. (2021). scGRNom: a computational pipeline of integrative multi-omics analyses for predicting cell-type disease genes and regulatory networks. Genome Med *13*, 95. 10.1186/s13073-021-00908-9.

34. Dupacova, N., Antosova, B., Paces, J., and Kozmik, Z. (2021). Meis homeobox genes control progenitor competence in the retina. Proceedings of the National Academy of Sciences of the United States of America *118*. 10.1073/pnas.2013136118.

35. Negishi, Y., Miya, F., Hattori, A., Mizuno, K., Hori, I., Ando, N., Okamoto, N., Kato, M., Tsunoda, T., Yamasaki, M., et al. (2015). Truncating mutation in NFIA causes brain malformation and urinary tract defects. Hum Genome Var *2*, 15007. 10.1038/hgv.2015.7.

36. Cadwell, C.R., Palasantza, A., Jiang, X., Berens, P., Deng, Q., Yilmaz, M., Reimer, J., Shen, S., Bethge, M., Tolias, K.F., et al. (2016). Electrophysiological, transcriptomic and morphologic profiling of single neurons using Patch-seq. Nat Biotechnol *34*, 199-203. 10.1038/nbt.3445.

37. Bos, J.L. (2003). Epac: a new cAMP target and new avenues in cAMP research. Nat Rev Mol Cell Biol *4*, 733-738. 10.1038/nrm1197.

38. de Rooij, J., Zwartkruis, F.J., Verheijen, M.H., Cool, R.H., Nijman, S.M., Wittinghofer, A., and Bos, J.L. (1998). Epac is a Rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP. Nature *396*, 474-477. 10.1038/24884.

39. Woolfrey, K.M., Srivastava, D.P., Photowala, H., Yamashita, M., Barbolina, M.V., Cahill, M.E., Xie, Z., Jones, K.A., Quilliam, L.A., Prakriya, M., and Penzes, P. (2009). Epac2 induces synapse remodeling and depression and its disease-associated forms alter spines. Nature neuroscience *12*, 1275-1284. 10.1038/nn.2386.

40. Consortium, G.T. (2015). Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. Science *348*, 648-660. 10.1126/science.1262110.

41. Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.Y., Peng, M., Collins, R., Grove, J., Klei, L., et al. (2020). Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiology of Autism. Cell *180*, 568-584 e523. 10.1016/j.cell.2019.12.036.

42. Katayama, Y., Nishiyama, M., Shoji, H., Ohkawa, Y., Kawamura, A., Sato, T., Suyama, M., Takumi, T., Miyakawa, T., and Nakayama, K.I. (2016). CHD8 haploinsufficiency results in autistic-like phenotypes in mice. Nature *537*, 675-679. 10.1038/nature19357.

43. Bernier, R., Golzio, C., Xiong, B., Stessman, H.A., Coe, B.P., Penn, O., Witherspoon, K., Gerdts, J., Baker, C., Vulto-van Silfhout, A.T., et al. (2014). Disruptive CHD8 mutations define a subtype of autism early in development. Cell *158*, 263-276. 10.1016/j.cell.2014.06.017.

44. Durak, O., Gao, F., Kaeser-Woo, Y.J., Rueda, R., Martorell, A.J., Nott, A., Liu, C.Y., Watson, L.A., and Tsai, L.H. (2016). Chd8 mediates cortical neurogenesis via transcriptional regulation of cell cycle and Wnt signaling. Nature neuroscience *19*, 1477-1488. 10.1038/nn.4400.

45. Kepecs, A., and Fishell, G. (2014). Interneuron cell types are fit to function. Nature *505*, 318-326. 10.1038/nature12983.

46. Jones, K.A., Sumiya, M., Woolfrey, K.M., Srivastava, D.P., and Penzes, P. (2019). Loss
of EPAC2 alters dendritic spine morphology and inhibitory synapse density. Molecular
and cellular neurosciences *98*, 19-31. 10.1016/j.mcn.2019.05.001.
47. Berry-Kravis, E., Hicar, M., and Ciurlionis, R. (1995). Reduced cyclic AMP production in
fragile X syndrome: cytogenetic and molecular correlations. Pediatr Res *38*, 638-643.
10.1203/00006450-199511000-00002.
48. Berry-Kravis, E.M., Harnett, M.D., Reines, S.A., Reese, M.A., Ethridge, L.E., Outterson,
A.H., Michalak, C., Furman, J., and Gurney, M.E. (2021). Inhibition of phosphodiesterase-
4D in adults with fragile X syndrome: a randomized, placebo-controlled, phase 2 clinical
trial. Nat Med *27*, 862-870. 10.1038/s41591-021-01321-w.
49. Sherwood, C.C., Subiaul, F., and Zawidzki, T.W. (2008). A natural history of the human
mind: tracing evolutionary changes in brain and cognition. Journal of anatomy *212*, 426-
454. 10.1111/j.1469-7580.2008.00868.x.
50. Wang, L., Pang, K., Zhou, L., Cebrian-Silla, A., Gonzalez-Granero, S., Wang, S., Bi, Q.,
White, M.L., Ho, B., Li, J., et al. (2023). A cross-species proteomic map reveals neoteny
of human synapse development. Nature *622*, 112-119. 10.1038/s41586-023-06542-2.
51. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J.,
Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-
cell data. Cell *184*, 3573-3587 e3529. 10.1016/j.cell.2021.04.048.
52. Stuart, T., Srivastava, A., Madad, S., Lareau, C.A., and Satija, R. (2021). Single-cell
chromatin state analysis with Signac. Nat Methods *18*, 1333-1341. 10.1038/s41592-021-
01282-5.
53. Schep, A.N., Wu, B., Buenrostro, J.D., and Greenleaf, W.J. (2017). chromVAR: inferring
transcription-factor-associated accessibility from single-cell epigenomic data. Nat
Methods *14*, 975-978. 10.1038/nmeth.4401.
54. Hahne, F., and Ivanek, R. (2016). Visualizing Genomic Data Using Gviz and
Bioconductor. Methods Mol Biol *1418*, 335-351. 10.1007/978-1-4939-3578-9_16.
55. Fornes, O., Castro-Mondragon, J.A., Khan, A., van der Lee, R., Zhang, X., Richmond,
P.A., Modi, B.P., Correard, S., Gheorghe, M., Baranasic, D., et al. (2020). JASPAR 2020:
update of the open-access database of transcription factor binding profiles. Nucleic Acids
Res *48*, D87-D92. 10.1093/nar/gkz1001.
56. Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos,
S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell transcriptional landscape
of mammalian organogenesis. Nature *566*, 496-502. 10.1038/s41586-019-0969-x.
57. Amezquita, R.A., Lun, A.T.L., Becht, E., Carey, V.J., Carpp, L.N., Geistlinger, L., Marini,
F., Rue-Albrecht, K., Risso, D., Soneson, C., et al. (2020). Orchestrating single-cell
analysis with Bioconductor. Nat Methods *17*, 137-145. 10.1038/s41592-019-0654-x.
58. Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y.,
Brenner, M., Loh, P.R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate
integration of single-cell data with Harmony. Nat Methods *16*, 1289-1296.
10.1038/s41592-019-0619-0.
59. Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. Nat Protoc *4*,
1184-1191. 10.1038/nprot.2009.97.

60.   Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017). Reversed graph embedding resolves complex single-cell trajectories. Nat Methods *14*, 979-982. 10.1038/nmeth.4402.

61.   Lawrence, M., Huber, W., Pages, H., Aboyoun, P., Carlson, M., Gentleman, R., Morgan, M.T., and Carey, V.J. (2013). Software for computing and annotating genomic ranges. PLoS Comput Biol *9*, e1003118. 10.1371/journal.pcbi.1003118.

62.   Huber, W., Carey, V.J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B.S., Bravo, H.C., Davis, S., Gatto, L., Girke, T., et al. (2015). Orchestrating high-throughput genomic analysis with Bioconductor. Nat Methods *12*, 115-121. 10.1038/nmeth.3252.

63.   Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics *32*, 2847-2849. 10.1093/bioinformatics/btw313.

64.   Rainer, J., Gatto, L., and Weichenberger, C.X. (2019). ensembldb: an R package to create and use Ensembl-based annotation resources. Bioinformatics *35*, 3151-3153. 10.1093/bioinformatics/btz031.

65.   Leng, N., Dawson, J.A., Thomson, J.A., Ruotti, V., Rissman, A.I., Smits, B.M., Haag, J.D., Gould, M.N., Stewart, R.M., and Kendziorski, C. (2013). EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq experiments. Bioinformatics *29*, 1035-1043. 10.1093/bioinformatics/btt087.

66.   Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C., and Chanda, S.K. (2019). Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat Commun *10*, 1523. 10.1038/s41467-019-09234-6.

67.   Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-based analysis of ChIP-Seq (MACS). Genome Biol *9*, R137. 10.1186/gb-2008-9-9-r137.

68.   Onorati, M., Li, Z., Liu, F., Sousa, A.M.M., Nakagawa, N., Li, M., Dell'Anno, M.T., Gulden, F.O., Pochareddy, S., Tebbenkamp, A.T.N., et al. (2016). Zika Virus Disrupts Phospho-TBK1 Localization and Mitosis in Human Neuroepithelial Stem Cells and Radial Glia. Cell Rep *16*, 2576-2592. 10.1016/j.celrep.2016.08.038.

69.   Shen, M., Sirois, C.L., Guo, Y., Li, M., Dong, Q., Mendez-Albelo, N.M., Gao, Y., Khullar, S., Kissel, L., Sandoval, S.O., et al. (2023). Species-specific FMRP regulation of RACK1 is critical for prenatal cortical development. Neuron. 10.1016/j.neuron.2023.09.014.

70.   Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888-1902 e1821. 10.1016/j.cell.2019.05.031.

71.   Stuart, T., Srivastava, A., Madad, S., Lareau, C.A., and Satija, R. (2022). Author Correction: Single-cell chromatin state analysis with Signac. Nat Methods *19*, 257. 10.1038/s41592-022-01393-7.

72.   Young, M.D., and Behjati, S. (2020). SoupX removes ambient RNA contamination from droplet-based single-cell RNA sequencing data. Gigascience *9*. 10.1093/gigascience/giaa151.

73.   McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder: Doublet Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst *8*, 329-337 e324. 10.1016/j.cels.2019.03.003.

74.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-
833   cell transcriptomic data across different conditions, technologies, and species. Nat
834   Biotechnol *36*, 411-420. 10.1038/nbt.4096.
835   75.   Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F.,
836   and Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using
837   UMAP. Nat Biotechnol. 10.1038/nbt.4314.
838   76.   Haghverdi, L., Lun, A.T.L., Morgan, M.D., and Marioni, J.C. (2018). Batch effects in
839   single-cell RNA-sequencing data are corrected by matching mutual nearest neighbors. Nat
840   Biotechnol *36*, 421-427. 10.1038/nbt.4091.
841   77.   Dong, Q., Kim, J., Nguyen, L., Bu, Q., and Chang, Q. (2020). An Astrocytic Influence on
842   Impaired Tonic Inhibition in Hippocampal CA1 Pyramidal Neurons in a Mouse Model of
843   Rett Syndrome. J Neurosci *40*, 6250-6261. 10.1523/JNEUROSCI.3042-19.2020.
844   78.   Hagemann-Jensen, M., Ziegenhain, C., Chen, P., Ramskold, D., Hendriks, G.J., Larsson,
845   A.J.M., Faridani, O.R., and Sandberg, R. (2020). Single-cell RNA counting at allele and
846   isoform resolution using Smart-seq3. Nat Biotechnol *38*, 708-714. 10.1038/s41587-020-
847   0497-0.
848   79.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
849   Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
850   Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.
851   80.   Gao, Y., Shen, M., Gonzalez, J.C., Dong, Q., Kannan, S., Hoang, J.T., Eisinger, B.E.,
852   Pandey, J., Javadi, S., Chang, Q., et al. (2020). RGS6 Mediates Effects of Voluntary
853   Running on Adult Hippocampal Neurogenesis. Cell Rep *32*, 107997.
854   10.1016/j.celrep.2020.107997.
855   81.   Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using
856   real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.
857   10.1006/meth.2001.1262.
858   82.   Gao, Y., Su, J., Guo, W., Polich, E.D., Magyar, D.P., Xing, Y., Li, H., Smrt, R.D., Chang,
859   Q., and Zhao, X. (2015). Inhibition of miR-15a Promotes BDNF Expression and Rescues
860   Dendritic Maturation Deficits in MeCP2-Deficient Neurons. Stem Cells *33*, 1618-1629.
861   10.1002/stem.1950.
862   83.   Guo, Y., Shen, M., Dong, Q., Mendez-Albelo, N.M., Huang, S.X., Sirois, C.L., Le, J., Li,
863   M., Jarzembowski, E.D., Schoeller, K.A., et al. (2023). Elevated levels of FMRP-target
864   MAP1B impair human and mouse neuronal development and mouse social behaviors via
865   autophagy pathway. Nat Commun *14*, 3801. 10.1038/s41467-023-39337-0.
866
867