# Near chromosome-level genome assembly of the microsporidium *Hamiltosporidium tvaerminnensis*

Running head: *Hamiltosporidium tvaerminnensis* genome report

Pascal Angst[1*], Jean-François Pombert[2], Dieter Ebert[1,†], and Peter D. Fields[1,†]

[1] Department of Environmental Sciences, Zoology, University of Basel, Basel, Switzerland
[2] Department of Biology, Illinois Institute of Technology, Chicago, Illinois, USA.
[*] Corresponding author
[†] Shared last author

Pascal Angst
Department of Environmental Sciences, Zoology
University of Basel
Vesalgasse 1, 4051 Basel, +4161 207 03 75, pascal.angst@unibas.ch

OrcIDs
Pascal Angst: 0000-0002-8654-2251
Jean-François Pombert: 0000-0002-3449-8375
Dieter Ebert: 0000-0003-2653-3772
Peter D. Fields: 0000-0003-2959-2524

Keywords: fungi, microsporidia, telomere-to-telomere, Iso-seq, alternative polyadenylation, splicing efficiency, *Hamiltosporidium tvaerminnensis*

## Abstract

Microsporidia are intracellular parasitic fungi whose genomes rank among the smallest of all known eukaryotes. A number of outstanding questions remain concerning the evolution of their large-scale variation in genome architecture, responsible for genome size variation of more than an order of magnitude. This genome report presents a first near-chromosomal assembly of a large-genome microsporidium, *Hamiltosporidium tvaerminnensis*. Combined Oxford Nanopore, Pacific Biosciences, and Illumina sequencing led to a genome assembly of 17 contigs, 11 of which represent complete chromosomes. Our assembly is 21.64 Mb in length, has an N50 of 1.44 Mb, and consists of 39.56 % interspersed repeats. We introduce a novel approach in microsporidia, PacBio Iso-Seq, as part of a larger annotation pipeline for obtaining high-quality annotations of 3,573 protein-coding genes. Based on direct evidence from the full-length Iso-Seq transcripts, we present evidence for alternative polyadenylation and variation in splicing efficiency, which are potential regulation mechanisms for gene expression in microsporidia. The generated high-quality genome assembly is a necessary resource for comparative genomics that will help elucidate the evolution of genome architecture in response to intracellular parasitism.

## Significance

Microsporidia are a model for genome evolution in response to intracellular parasitism, but we lack high-quality resources from species with large genomes. We present a near complete assembly of a large-genome microsporidium, *Hamiltosporidium tvaerminnensis*, and obtain high-quality gene annotations through full-length transcripts using Iso-Seq, a

52    novel approach in microsporidia. Our study provides insights into gene regulation and paves

53    the way for comparative genomic analyses aiming to understand the evolution of genome

54    reduction and expansion in these intracellular parasites.

## Introduction

Microsporidia have become a valuable model in evolutionary biology for studying extreme

parasitism (Murareanu et al., 2021; Wadi & Reinke, 2020). Compared to their fungi relatives,

the genomes of microsporidia have been strongly shaped by their specialized life history of

obligate intracellular parasitism (Corradi, 2015). In particular, the genomes of most

microsporidia are very compact and reduced (Murareanu et al., 2021). However, genome

sizes of microsporidia vary by more than an order of magnitude, making them a suitable

model for studying genome evolution in eukaryotes and parasites (Wadi & Reinke, 2020). A

powerful approach for this is comparative genomics (i.e., comparing the presence or

absence of genetic sequences, their relative location, and their abundance among different

species) for which high-quality genome assemblies are a prerequisite (Kille et al., 2022).

Increasing interest in microsporidia has boosted available genome resources, but most

published genomes are drafts (Williams et al., 2022). Following from recent successful

efforts to obtain telomere-to-telomere assemblies in *Encephalitozoon*, a genus of

microsporidia with small genomes (Mascarenhas dos Santos et al., 2023), we focus here on

a large-genome species, *Hamiltosporidium tvaerminnensis*.

First noted by Green (1957) as *Octosporea bayeri*, the reclassified *H. tvaerminnensis* has

become a model in the study of host–parasite interactions with its only host *Daphnia*

*magna*, a planktonic microcrustacean (Altermatt et al., 2007; Lass & Ebert, 2005; Orlansky &

Ben-Ami, 2019; Routtu & Ebert, 2015; Santos & Ebert, 2022; Vizoso & Ebert, 2004). Haag et

al. (2011) provided the cytological and molecular description of *H. tvaerminnensis*, and

Corradi et al. (2009) produced the first draft of its genome. Recently, genetic and genomic

studies have started using *H. tvaerminnensis* to investigate the evolution of variation in

microsporidian genome architecture (Angst et al., 2022, 2023; de Albuquerque et al., 2020;

Haag et al., 2013, 2020). *Hamiltosporidium tvaerminnensis* has a remarkably large genome

compared to other microsporidia, partly due to the proliferation of transposable genomic

elements (de Albuquerque et al., 2020; Parisot et al., 2014). Draft genomes are available for

the nearest known relative, *H. magnivora*, and more distantly related species, *Thelohania*

*contejeani* (Cormier, Wattier, et al., 2021) and *Edhazardia aedis* (Desjardins et al., 2015).

Demands for comparative genomic approaches are increasing, calling for an improved

genome of *H. tvaerminnensis*, a leading model organism among large-genome

microsporidia. A high-quality genome will allow advancements in the study of genome

architecture evolution in microsporidia; for example, through comparison with

chromosomal assemblies of *Encephalitozoon*. We have integrated Pacific Biosciences

(PacBio) Continuous Long Reads (CLR), Oxford Nanopore Technologies (ONT), and Illumina

DNA sequencing for a near-chromosomal assembly, and we employed PacBio Iso-Seq for

high-quality gene annotation. The first use of Iso-Seq for a microsporidium provides insights

into microsporidian alternative polyadenylation (APA) and splicing efficiency as potential

regulation mechanisms of gene expression in reduced genomes.

## Results and Discussion

### Genome Sequencing and Assembly

We generated a genome assembly of *H. tvaerminnensis* based on PacBio CLR and ONT sequencing reads (total/N50: 17.53 Gb/1.54 Kb and 2.75 Gb/32.68 Kb, respectively). In a subsequent step, we polished this assembly using Illumina paired-end (PE) sequencing reads (total: 66 Gb). The final assembly consisted of 21.64 Mb in 17 contigs with an N50 of 1.44 Mb (Table 1, Figure 1). We used BUSCO to assess the biological completeness of our assembly and found a BUSCO score of 94 %, a relatively high value for a large-genome microsporidium (Cormier, Chebbi, et al., 2021). The structural completeness of our assembly was assessed using scripts from (Mascarenhas dos Santos et al., 2023), which allow for the identification of telomeres. Telomeres are identifiable by terminal, short, tandem repeats– $(TTAGGG)_n$ in the case of *H. tvaerminnensis*, as is the case in most fungi (Rahnama et al., 2020). Eleven contigs represented complete chromosomes flanked by telomeres at both ends (Figure 1, Table S1), four contigs had only one telomere, and two had none. Therefore, we estimated the genome contains 13 – 17 chromosomes. Large multicopy repeats arrayed in tandem and scattered throughout the genome prevented us from fully assembling the chromosomes with major repetitive loci abutting contig ends without telomeres (Figure 1). Overall, 39.56 % of the genome consisted of interspersed repeats (retroelements 7.07 %, DNA transposons 7.04 %, other 25.45 %). High numbers of repeats are common in large-genome microsporidia (de Albuquerque et al., 2020), explaining why previous attempts to assemble the genome of *H. tvaerminnensis* have been less successful in both completeness and contiguity (size = 18.34 Mb; number of contigs = 2,915; N50 = 9.58 Kb; BUSCO score =

87.6 %) (Haag et al., 2020) compared to our long-read sequencing-based assembly. The

highest quality genomes within the microsporidia are from the genus *Encephalitozoon*,

which have an order of magnitude smaller genomes than *H. tvaerminnensis* and telomeric

repeats of (TTAGG)$_n$ on both ends of all eleven chromosomes (Mascarenhas dos Santos et

al., 2023).

**Figure 1. Karyoplot of *H. tvaerminnensis*.** Eleven of the 17 contigs have two telomeres, i.e.,

represent full chromosomes. In total, 39.56 % of the genome are interspersed repeats and

low complexity DNA sequences. The inset shows the repeat length distribution on a base-10

logarithmic scale.

**Table 1. Statistics of the assembly and annotation of the *H. tvaerminnensis* genome.**

| Assembly | |
|---|---|
| Total length (Mb) | 21.64 |
| GC content (%) | 26.60 |
| Contig N50 (Mb) | 1.44 |
| Contig number | 17 |
| BUSCO completeness score (%) | 94 |
| **Annotation** | |
| Total length of repeats (Mb) | 8.56 |
| Number of protein-coding genes | 3,573 |
| Mean gene length (bp) | 1,391 |
| Number of predicted introns | 98 |
| Number of Iso-Seq confirmed introns | 17 |
| Number of genes with confirmed introns | 15 |
| Mean intron length (bp) | 28 |

## Iso-Seq for Genome Annotation

RNA-seq data from intracellular parasites are sparse, and even more sparse are full-length transcript sequences. Iso-Seq helped us to predict both gene structures and untranslated regions (UTRs) more accurately than was previously done for microsporidia. Using 8.5 Gb of Iso-Seq reads (so-called circular consensus sequence (CCS) of cDNA), we obtained annotations for 3,573 protein-coding genes with a mean length of 1,391 bp (Table 1). The number of genes in microsporidia ranges from about 2,000 to 4,000 and is positively correlated with genome size (Jespersen et al., 2022). Compared to previous work, we obtained fewer fragmented genes; the score of fragmented BUSCOs in our annotations was 1.8 % (previously 3.8 %; Haag et al., 2020). We also obtained many fewer introns (98) than previously predicted (890), but our number is still higher than estimates for other large-genome microsporidia annotated with the help of (short-read) RNA-seq (Desjardins et al., 2015). After observing the discrepancy between annotations with and without RNA-seq data, we suspected introns may not be efficiently spliced.

Splicing Efficiency Analysis

Species with reduced genomes, sparse introns or no introns, and reduced spliceosomes

have been suggested to exhibit low splicing efficiencies (Schärfen et al., 2022; Whelan et al.,

2019). Microsporidia are such species. For example, in *E. cuniculi*, splicing efficiency for

small introns is only around 20 % (Grisdale et al., 2013). We used direct evidence from full-

length transcripts and found high variation in splicing efficiency in *H. tvaerminnensis* as has

previously been reported for the reduced genomes of the red alga *Cyanidioschyzon merolae*

(Schärfen et al., 2022) and for parasitic fungi (Sieber et al., 2018). We used 17 introns from

15 genes with good representation in the Iso-Seq data (Table S2). These introns had a mean

length of 28 bp. The percentages of spliced transcripts ranged from 7.5 % to 92.3 % (Table

S2) with a mean of 54.53 %. We did not find a statistical correlation between percent spliced

and the length of introns. In conclusion, we found fewer introns in *H. tvaerminnensis* than

previously suggested and intermediate splicing efficiencies, which is in line with theorized

patterns for reduced-genome species. All but two introns identified in *H. tvaerminnensis*

featured the same splice sites as have been found in (non-)parasitic fungi (i.e., 5'GU...AG3')

(Kupfer et al., 2004). The other two introns had splice sites 5'AA...AG3' and 5'AU...GA3'.

Alternative Polyadenylation

The mechanisms of gene regulation in microsporidia are unexplored. APA is a widespread

and conserved regulatory mechanism by which the availability of multiple polyadenylation

sites can lead to variation in 3'-UTR lengths of mRNA isoforms and thus 3'-UTR content,

which strongly influences gene expression (Mitschka & Mayr, 2022). Here, Iso-Seq data

allowed us to investigate the occurrence of mRNA isoforms encoding the same protein but

differing in their 3'-UTRs resulting from APA. By observing multiple 3'-ends for many genes,

we found direct evidence for APA in the full-length transcripts of *H. tvaerminnensis*.

Upstream of the 3'-ends, a motif analysis of the 500 most expressed genes revealed

enrichment of an UAAA tetramer. The UAAA tetramer was found in 95.8 % of the genes and

87.7 % of them were part of an A[A/U]UAAA hexamer on average 20 nucleotides upstream

of the most common 3'-ends. A[A/U]UAAA is a known motif for cleavage of protein-coding

mRNAs in metazoa (Neve et al., 2017) and is shown to be enriched upstream of 3'-ends in

the bee-infecting microsporidium *Nosema ceranae* (Chen et al., 2021). These motifs were

slightly less enriched upstream of the second and third most common 3'-ends (Table S3). No

motif was strongly enriched that did not contain an UAAA tetramer.

## Materials and Methods

### Sample preparation and sequencing

The *H. tvaerminnensis* isolate used in this study was obtained from the *D. magna* genotype

FI-OER-3-3, collected in July 2005 from a rockpool (59°47'18.9"N 23°10'26.9"E) on the Island

Oeren in the Tvaerminne archipelago, Southwestern Finland. The same isolate has been

Illumina sequenced for earlier genome drafts (Corradi et al., 2009; Haag et al., 2020) and a

biogeographic study (Angst et al., 2022). Genomic Illumina reads from the latter were

reused here (NCBI database, SRA accession: SRX13146514, Bioproject ID: PRJNA780787).

We generated long-read sequencing data by applying a protocol described in Fields et al.,

(2015) for obtaining microbiota-free *D. magna* tissue, but retaining microsporidia, from

which we extracted high-molecular-weight DNA using Genomic-tips (QIAGEN). A standard

PacBio genomic DNA library and sequencing on a SMRTcell in CLR mode on a Sequel I have

previously been done at the Quantitative Genomics Facility service platform at the

Department of Biosystem Science and Engineering (D-BSSE, ETH) Basel, Switzerland (NCBI

database, SRA accession: SRR23724335, Bioproject ID: PRJNA778105). Additionally, a ONT

genomic DNA library was prepared with the SQK-LSK110 ligation kit and sequenced on a

MinION device with a Spot-ON Flow Cell (R9.4.1).

To extract full-length transcripts, we used the RNeasy extraction kit (QIAGEN)

supplying Proteinase K for protein digestion. The NEBNext Single Cell/Low Input cDNA

Synthesis and Amplification Module kit (New England Biolabs) was used to generate cDNA

from the extraction. An Iso-Seq cDNA library was generated with the Iso-Seq Express Oligo

Kit and the SMRTbell Express Template Prep Kit 2.0 (PacBio). The Iso-Seq library was

sequenced using a single SMRTcell on a Sequel I at the D-BSSE (Basel, Switzerland) and

processed with the *isoseq3* pipeline (https://github.com/PacificBiosciences/IsoSeq),

resulting in approximately 8.5 Gb of CCS data.

Genome assembly

An assembly based on the PacBio CLR sequencing data, generated with canu v.2.1.1 (Koren

et al., 2017), was extended *in-silico* using a seed-based approach with Consed v.29 (Gordon

et al., 1998) as described in Pombert et al. (2013). Independently, the Nanopore sequencing

reads were basecalled with bonito v.0.5.3 (https://github.com/nanoporetech/bonito) using

the dna_r9.4.1_e8.1_sup@v3.3 model and then assembled with nextDenovo v.2.5.0

(https://github.com/Nextomics/NextDenovo) using an expected genome size of 20 Mb. The

Nanopore nextDenovo assembly was polished with the Illumina reads and the PacBio CLR

reads using nextPolish v.1.4.1 (Hu et al., 2020), Medaka v.1.7.2

(https://github.com/nanoporetech/medaka), and two runs of Pilon v.1.24 (Walker et al.,

2014) and then merged with the extended PacBio CLR assembly using quickmerge v.0.3

(Solares et al., 2018) and manual curation. The absence of contaminants in this final

assembly was ascertained by BLAST homology searches against the NCBI database and its

completeness was assessed with check_for_telomeres.pl v.0.3 (Mascarenhas dos Santos et

al., 2023). Repeats in the assembly were detected with RepeatModeler v.2.0.2 including the

LTR pipeline (Flynn et al., 2020) and ReapeatMasker v.4.1.2 (Smit et al., 2013) and plotted

with karyoploteR v.1.24.0 (Gel & Serra, 2017) in R v.4.2.2 (R Core Team, 2022).

Genome annotation

The *H. tvaerminnensis* genome was annotated by feeding the Iso-Seq transcriptome data as

extrinsic evidence to the fungal annotation pipeline funannotate v.1.8.13 (Palmer & Stajich,

2022). Briefly, as per guideline for gene prediction using long RNA reads

(https://github.com/Gaius-

Augustus/BRAKER/blob/master/docs/long_reads/long_read_protocol.md), we mapped the

whole-transcript reads from the CCS FASTA produced by *isoseq3* to the repeat-masked

genome using Minimap2 v.2.22 (Li, 2018) with the flags '-ax splice –secondary=no -C5'.

Approximately 25.5 % of the reads aligned to the parasite genome. We then collapsed the

total read set into distinct transcripts (while retaining isoforms) using the

*collapse_isoforms_by_sam.py* script provided by the *cDNA_Cupcake*

(https://github.com/Magdoll/cDNA_Cupcake) pipeline and ran GeneMarkS-T v.5.1 (Tang et

al., 2015) to predict protein-coding regions in the transcripts. The resultant FASTA and GFF

files were used for gene prediction and annotation with the wrapper tool funannotate

v.1.8.13 (Palmer & Stajich, 2022), which relies on TANTAN v.26 (Frith, 2011) for repeat-

masking. Funannotate operated AUGUSTUS v.3.4.0 (Stanke et al., 2006, 2008), GeneMark-ES

v.4.62 (Brůna et al., 2020), GlimmerHMM v.3.0.4 (Majoros et al., 2004), SNAP v.2013_11_29

(Korf, 2004), and EVidenceModeler v.1.1.1 (Haas et al., 2008) for ab-initio gene prediction

and consensus gene structure generation. Functional annotation in funannoate relies on the

combined evidence from InterProScan v.5.55_88.0 (Jones et al., 2014), eggNOG-mapper

v.2.1.9 (Cantalapiedra et al., 2021; Huerta-Cepas et al., 2019), Phobius v.1.01 (Käll et al.,

2004), SignalP v.5.0b (Almagro Armenteros et al., 2019), and antiSMASH v.6.0 (Blin et al.,

2021) as well as on comparisons to Pfam (Mistry et al., 2021), UniProt (The UniProt

Consortium, 2022), MEROPS (Rawlings et al., 2014), dbCAN (Yin et al., 2012) and BUSCO

v.5.4.3 (Manni et al., 2021) databases. Also, the completeness was assessed using BUSCO

with its microsporidia_odb10 database (Creation date: 2020-08-05). For proteins without

functional annotation from this pipeline, we lifted-over functional annotations from the

species reference proteome (UniProt database, Proteome ID: UP000292282) using BLAST

v.2.3.0+ (Camacho et al., 2009), AGAT v.1.0.0 (Dainat, 2022), and GFF3sort v.1.0.0 (Zhu et

al., 2017).

Full-length transcript analyses

For the analysis of splicing efficiency and APA, we followed the methodology described in

Schärfen et al. (2022). Briefly, starting from the BAM file, which was provided by the

sequencing facility, we used lima v.2.6.0 (https://github.com/PacificBiosciences/barcoding)

in "-isoseq" mode to remove IsoSeq template-switching oligo sequences and to orient

subreads in the correct 5' to 3' direction. After aligning subreads to our genome using

minimap2 in "splice:hq" mode, we used SAMtools v.1.16.1 (Li et al., 2009) and BEDtools

v.2.30.0 (Quinlan, 2014) to identify intron-spanning reads. The percentage of spliced introns

was calculated by dividing the number of spliced reads by the total number of reads for

each intron times 100. For this analysis, introns were manually selected based on Iso-Seq

coverage level. A potential correlation between the percent of spliced reads and intron

length was tested in R.

For the analysis of APA, briefly, we used SAMtools, BEDtools, and pysam v.0.20.0

(https://github.com/pysam-developers/pysam) to extract each read's 3'-end coordinate

following Schärfen et al. (2022). Then, we drew the distribution of 3'-end coordinates as a

histogram separately for each gene and applied the find_peaks algorithm from scipy.signal

v.1.7.3 (Virtanen et al., 2020) with height equal to four and distance equal to 18. Hence, we

found the number of peaks (=3'-end coordinates) per gene. We then separately extracted

sequences of the 500 most expressed genes up to 40 nucleotides upstream of the highest

peak, the second highest peak, and the third highest peak using seqtk v.1.3-r106

(https://github.com/lh3/seqtk). For each set of sequences, we did a motif analysis using

XSTREME v5.5.0 from the MEME Suite (Grant & Bailey, 2021) with a minimum motif length

of four and determined the distance of the obtained motif from the peak.

## Data Availability Statement

Raw data is deposited at the NCBI SRA database and the assembled genome as well as

predicted set of protein sequences are available at NCBI GenBank (FIOER33 v3; BioProject ID

PRJNA778105) and at https://doi.org.10.6084/m9.figshare.22794887 and will be publicly

available upon acceptance. (Reviewer links: ftp://ftp-

trace.ncbi.nlm.nih.gov/sra/review/SRP426041_20230526_134124_5035f32c1836f93f1e061

a20b37fb2d0; https://figshare.com/s/7ee5e2ab29e3730a47e7)

## Conflict of Interest

296 The authors have no conflicts of interest to declare.

## Author Contributions

298 All authors designed the study. DE collected the sample. PA and PDF performed the

299 molecular work. PA, JFP, and PDF analyzed the data. PA wrote the manuscript. All authors

300 reviewed the manuscript.

## Acknowledgements

302 We thank Jürgen Hottinger, Urs Stiefel, Michelle Krebs, and Alix Thivolle for help in the

303 laboratory. We thank members of the Ebert group for providing feedback on the study and

304 the manuscript.

## Funding

306 This work was supported by the Swiss National Science Foundation (SNSF) (grant numbers

307 310030B_166677 and 310030_188887 to DE).

## References

Almagro Armenteros, J. J., Tsirigos, K. D., Sønderby, C. K., Petersen, T. N., Winther, O.,

Brunak, S., von Heijne, G., & Nielsen, H. (2019). SignalP 5.0 improves signal peptide

predictions using deep neural networks. *Nature Biotechnology*, *37*(4), Article 4.

https://doi.org/10.1038/s41587-019-0036-z

Altermatt, F., Hottinger, J., & Ebert, D. (2007). Parasites promote host gene flow in a

metapopulation. *Evolutionary Ecology*, *21*(4), 561–575.

https://doi.org/10.1007/s10682-006-9136-6

Angst, P., Ebert, D., & Fields, P. D. (2022). Demographic history shapes genomic variation in

an intracellular parasite with a wide geographical distribution. *Molecular Ecology*,

*31*(9), 2528–2544. https://doi.org/10.1111/mec.16419

Angst, P., Ebert, D., & Fields, P. D. (2023). Population genetic analysis of the microsporidium

*Ordospora colligata* reveals the role of natural selection and phylogeography on its

extremely compact and reduced genome. *G3 Genes|Genomes|Genetics*, jkad017.

https://doi.org/10.1093/g3journal/jkad017

Blin, K., Shaw, S., Kloosterman, A. M., Charlop-Powers, Z., van Wezel, G. P., Medema, M. H.,

& Weber, T. (2021). antiSMASH 6.0: Improving cluster detection and comparison

capabilities. *Nucleic Acids Research*, *49*(W1), W29–W35.

https://doi.org/10.1093/nar/gkab335

Brůna, T., Lomsadze, A., & Borodovsky, M. (2020). GeneMark-EP+: Eukaryotic gene

prediction with self-training in the space of genes and proteins. *NAR Genomics and*

*Bioinformatics*, *2*(2), lqaa026. https://doi.org/10.1093/nargab/lqaa026

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T.

L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*, 421.

https://doi.org/10.1186/1471-2105-10-421

Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., & Huerta-Cepas, J. (2021).

eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain

Prediction at the Metagenomic Scale. *Molecular Biology and Evolution*, *38*(12),

5825–5829. https://doi.org/10.1093/molbev/msab293

Chen, H.-Z., Fan, X.-X., Fan, Y.-C., Wang, J., Zhu, Z.-W., Jiang, H.-B., Zhang, W.-D., Long, Q.,

Xiong, C.-L., Zheng, Y.-Z., Fu, Z.-M., Xu, G.-J., Chen, D.-F., & Guo, R. (2021). Analysis of

alternative splicing and alternative polyadenylation of *Nosema ceranae* genes.

*Mycosystema*, *40*(1), 161–173. https://doi.org/10.13346/j.mycosystema.200216

Cormier, A., Chebbi, M. A., Giraud, I., Wattier, R., Teixeira, M., Gilbert, C., Rigaud, T., &

Cordaux, R. (2021). Comparative Genomics of Strictly Vertically Transmitted,

Feminizing Microsporidia Endosymbionts of Amphipod Crustaceans. *Genome Biology*

*and Evolution*, *13*(1), evaa245. https://doi.org/10.1093/gbe/evaa245

Cormier, A., Wattier, R., Giraud, I., Teixeira, M., Grandjean, F., Rigaud, T., & Cordaux, R.

(2021). Draft Genome Sequences of *Thelohania contejeani* and *Cucumispora*

*dikerogammari*, Pathogenic Microsporidia of Freshwater Crustaceans. *Microbiology*

*Resource Announcements*, *10*(2), e01346-20. https://doi.org/10.1128/MRA.01346-20

Corradi, N. (2015). Microsporidia: Eukaryotic Intracellular Parasites Shaped by Gene Loss

and Horizontal Gene Transfers. *Annual Review of Microbiology*, *69*(1), 167–183.

https://doi.org/10.1146/annurev-micro-091014-104136

Corradi, N., Haag, K. L., Pombert, J.-F., Ebert, D., & Keeling, P. J. (2009). Draft genome

sequence of the *Daphnia* pathogen *Octosporea bayeri*: Insights into the gene content

of a large microsporidian genome and a model for host-parasite interactions.

*Genome Biology*, *10*(10), R106. https://doi.org/10.1186/gb-2009-10-10-r106

Dainat, J. (2022). *AGAT: Another Gff Analysis Toolkit to handle annotations in any GTF/GFF*

*format.* Zenodo. https://doi.org/10.5281/zenodo.7255559

de Albuquerque, N. R. M., Ebert, D., & Haag, K. L. (2020). Transposable element abundance

correlates with mode of transmission in microsporidian parasites. *Mobile DNA*, *11*,

19. https://doi.org/10.1186/s13100-020-00218-8

Desjardins, C. A., Sanscrainte, N. D., Goldberg, J. M., Heiman, D., Young, S., Zeng, Q.,

Madhani, H. D., Becnel, J. J., & Cuomo, C. A. (2015). Contrasting host–pathogen

interactions and genome evolution in two generalist and specialist microsporidian

pathogens of mosquitoes. *Nature Communications*, *6*, 7121.

https://doi.org/10.1038/ncomms8121

Fields, P. D., Reisser, C., Dukić, M., Haag, C. R., & Ebert, D. (2015). Genes mirror geography in

*Daphnia magna*. *Molecular Ecology*, *24*(17), 4521–4536.

https://doi.org/10.1111/mec.13324

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F. (2020).

RepeatModeler2 for automated genomic discovery of transposable element families.

*Proceedings of the National Academy of Sciences*, *117*(17), 9451–9457.

https://doi.org/10.1073/pnas.1921046117

Frith, M. C. (2011). A new repeat-masking method enables specific detection of homologous

sequences. *Nucleic Acids Research*, *39*(4), e23. https://doi.org/10.1093/nar/gkq1212

Gel, B., & Serra, E. (2017). karyoploteR: An R/Bioconductor package to plot customizable

genomes displaying arbitrary data. *Bioinformatics*, *33*(19), 3088–3090.

https://doi.org/10.1093/bioinformatics/btx346

Gordon, D., Abajian, C., & Green, P. (1998). Consed: A Graphical Tool for Sequence Finishing.

*Genome Research*, *8*(3), 195–202. https://doi.org/10.1101/gr.8.3.195

Grant, C. E., & Bailey, T. L. (2021). *XSTREME: Comprehensive motif analysis of biological*

*sequence datasets* (p. 2021.09.02.458722). bioRxiv.

https://doi.org/10.1101/2021.09.02.458722

Green, J. (1957). Parasites and epibionts of Cladocera in rock pools of Tvärminne

archipelago. *Archivum Societatis Zoologicae Botanicae Fennicae 'Vanamo*, *12*, 5–12.

Grisdale, C. J., Bowers, L. C., Didier, E. S., & Fast, N. M. (2013). Transcriptome analysis of the

parasite *Encephalitozoon cuniculi*: An in-depth examination of pre-mRNA splicing in a

reduced eukaryote. *BMC Genomics*, *14*(1), 207. https://doi.org/10.1186/1471-2164-

14-207

Haag, K. L., Larsson, J. I. R., Refardt, D., & Ebert, D. (2011). Cytological and molecular

description of *Hamiltosporidium tvaerminnensis* gen. Et sp. Nov., a microsporidian

parasite of *Daphnia magna*, and establishment of *Hamiltosporidium magnivora*

comb. Nov. *Parasitology*, *138*(4), 447–462.

https://doi.org/10.1017/S0031182010001393

Haag, K. L., Pombert, J.-F., Sun, Y., de Albuquerque, N. R. M., Batliner, B., Fields, P., Lopes, T.

F., & Ebert, D. (2020). Microsporidia with Vertical Transmission Were Likely Shaped

by Nonadaptive Processes. *Genome Biology and Evolution*, *12*(1), 3599–3614.

https://doi.org/10.1093/gbe/evz270

Haag, K. L., Traunecker, E., & Ebert, D. (2013). Single-nucleotide polymorphisms of two

closely related microsporidian parasites suggest a clonal population expansion after

the last glaciation. *Molecular Ecology*, *22*(2), 314–326.

https://doi.org/10.1111/mec.12126

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., White, O., Buell, C. R., &

Wortman, J. R. (2008). Automated eukaryotic gene structure annotation using

EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome

Biology*, *9*(1), R7. https://doi.org/10.1186/gb-2008-9-1-r7

Hu, J., Fan, J., Sun, Z., & Liu, S. (2020). NextPolish: A fast and efficient genome polishing tool

for long-read assembly. *Bioinformatics*, *36*(7), 2253–2255.

https://doi.org/10.1093/bioinformatics/btz891

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H.,

Mende, D. R., Letunic, I., Rattei, T., Jensen, L. J., von Mering, C., & Bork, P. (2019).

eggNOG 5.0: A hierarchical, functionally and phylogenetically annotated orthology

resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research*, *47*(D1),

D309–D314. https://doi.org/10.1093/nar/gky1085

Jespersen, N., Monrroy, L., & Barandun, J. (2022). Impact of Genome Reduction in

Microsporidia. In L. M. Weiss & A. W. Reinke (Eds.), *Microsporidia: Current Advances*

*in Biology* (pp. 1–42). Springer International Publishing. https://doi.org/10.1007/978-

3-030-93306-7_1

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,

Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew,

M., Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: Genome-scale protein

function classification. *Bioinformatics*, *30*(9), 1236–1240.

https://doi.org/10.1093/bioinformatics/btu031

Käll, L., Krogh, A., & Sonnhammer, E. L. L. (2004). A Combined Transmembrane Topology

and Signal Peptide Prediction Method. *Journal of Molecular Biology*, *338*(5), 1027–

1036. https://doi.org/10.1016/j.jmb.2004.03.016

Kille, B., Balaji, A., Sedlazeck, F. J., Nute, M., & Treangen, T. J. (2022). Multiple genome

alignment in the telomere-to-telomere assembly era. *Genome Biology*, *23*(1), 182.

https://doi.org/10.1186/s13059-022-02735-6

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. (2017).

Canu: Scalable and accurate long-read assembly via adaptive *k*-mer weighting and

repeat separation. *Genome Research*, *27*(5), 722–736.

https://doi.org/10.1101/gr.215087.116

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics*, *5*(1), 59.

https://doi.org/10.1186/1471-2105-5-59

Kupfer, D. M., Drabenstot, S. D., Buchanan, K. L., Lai, H., Zhu, H., Dyer, D. W., Roe, B. A., &

Murphy, J. W. (2004). Introns and Splicing Elements of Five Diverse Fungi. *Eukaryotic*

*Cell*, *3*(5), 1088–1100. https://doi.org/10.1128/EC.3.5.1088-1100.2004

Lass, S., & Ebert, D. (2005). Apparent seasonality of parasite dynamics: Analysis of cyclic

prevalence patterns. *Proceedings of the Royal Society B: Biological Sciences*,

*273*(1583), 199–206. https://doi.org/10.1098/rspb.2005.3310

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*,

*34*(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., &

Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools.

*Bioinformatics*, *25*(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Majoros, W. H., Pertea, M., & Salzberg, S. L. (2004). TigrScan and GlimmerHMM: Two open

source ab initio eukaryotic gene-finders. *Bioinformatics*, *20*(16), 2878–2879.

https://doi.org/10.1093/bioinformatics/bth315

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO

Update: Novel and Streamlined Workflows along with Broader and Deeper

Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes.

*Molecular Biology and Evolution*, *38*(10), 4647–4654.

https://doi.org/10.1093/molbev/msab199

Mascarenhas dos Santos, A. C., Julian, A. T., Liang, P., Juárez, O., & Pombert, J.-F. (2023).

Telomere-to-Telomere genome assemblies of human-infecting *Encephalitozoon*

species. *BMC Genomics*, *24*(1), 237. https://doi.org/10.1186/s12864-023-09331-3

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L.,

Tosatto, S. C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., & Bateman, A.

(2021). Pfam: The protein families database in 2021. *Nucleic Acids Research*, *49*(D1),

D412–D419. https://doi.org/10.1093/nar/gkaa913

Mitschka, S., & Mayr, C. (2022). Context-specific regulation and function of mRNA

alternative polyadenylation. *Nature Reviews Molecular Cell Biology*, *23*(12), Article

12. https://doi.org/10.1038/s41580-022-00507-5

Murareanu, B. M., Sukhdeo, R., Qu, R., Jiang, J., & Reinke, A. W. (2021). Generation of a

Microsporidia Species Attribute Database and Analysis of the Extensive Ecological

and Phenotypic Diversity of Microsporidia. *MBio*, *12*(3), e01490-21.

https://doi.org/10.1128/mBio.01490-21

Neve, J., Patel, R., Wang, Z., Louey, A., & Furger, A. M. (2017). Cleavage and

polyadenylation: Ending the message expands gene regulation. *RNA Biology*, *14*(7),

865–890. https://doi.org/10.1080/15476286.2017.1306171

Orlansky, S., & Ben-Ami, F. (2019). Genetic resistance and specificity in sister taxa of

*Daphnia*: Insights from the range of host susceptibilities. *Parasites & Vectors*, *12*(1),

545. https://doi.org/10.1186/s13071-019-3795-y

Palmer, J., & Stajich, J. (2022). *nextgenusfs/funannotate: Funannotate* (1.8.11). Zenodo.

https://zenodo.org/record/2604804

Parisot, N., Pelin, A., Gasc, C., Polonais, V., Belkorchia, A., Panek, J., El Alaoui, H., Biron, D. G.,

Brasset, E., Vaury, C., Peyret, P., Corradi, N., Peyretaillade, É., & Lerat, E. (2014).

Microsporidian genomes harbor a diverse array of transposable elements that

demonstrate an ancestry of horizontal exchange with metazoans. *Genome Biology*

*and Evolution*, *6*(9), 2289–2300. https://doi.org/10.1093/gbe/evu178

Pombert, J.-F., Xu, J., Smith, D. R., Heiman, D., Young, S., Cuomo, C. A., Weiss, L. M., &

Keeling, P. J. (2013). Complete Genome Sequences from Three Genetically Distinct

Strains Reveal High Intraspecies Genetic Diversity in the Microsporidian

*Encephalitozoon cuniculi*. *Eukaryotic Cell*, *12*(4), 503–511.

https://doi.org/10.1128/EC.00312-12

Quinlan, A. R. (2014). BEDTools: The Swiss-army tool for genome feature analysis. *Current*

*Protocols in Bioinformatics*, *47*, 11.12.1-11.12.34.

https://doi.org/10.1002/0471250953.bi1112s47

R Core Team. (2022). *R: The R Project for Statistical Computing*. R Foundation for Statistical

Computing. https://www.r-project.org/

Rahnama, M., Novikova, O., Starnes, J. H., Zhang, S., Chen, L., & Farman, M. L. (2020).

Transposon-mediated telomere destabilization: A driver of genome evolution in the

blast fungus. *Nucleic Acids Research*, *48*(13), 7197–7217.

https://doi.org/10.1093/nar/gkaa287

Rawlings, N. D., Waller, M., Barrett, A. J., & Bateman, A. (2014). MEROPS: The database of

proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Research*, *42*(D1),

D503–D509. https://doi.org/10.1093/nar/gkt953

Routtu, J., & Ebert, D. (2015). Genetic architecture of resistance in *Daphnia* hosts against

two species of host-specific parasites. *Heredity*, *114*(2), 241–248.

https://doi.org/10.1038/hdy.2014.97

Santos, J. L., & Ebert, D. (2022). The effects of temperature and host-parasite interactions on

parasite persistence in a planktonic crustacean. *Journal of Freshwater Ecology*, *37*(1),

555–568. https://doi.org/10.1080/02705060.2022.2134219

Schärfen, L., Zigackova, D., Reimer, K. A., Stark, M. R., Slat, V. A., Francoeur, N. J., Wells, M.

L., Zhou, L., Blackshear, P. J., Neugebauer, K. M., & Rader, S. D. (2022). Identification

of Alternative Polyadenylation in *Cyanidioschyzon merolae* Through Long-Read

Sequencing of mRNA. *Frontiers in Genetics*, *12*.

https://www.frontiersin.org/articles/10.3389/fgene.2021.818697

Sieber, P., Voigt, K., Kämmer, P., Brunke, S., Schuster, S., & Linde, J. (2018). Comparative

Study on Alternative Splicing in Human Fungal Pathogens Suggests Its Involvement

During Host Invasion. *Frontiers in Microbiology*, *9*.

https://www.frontiersin.org/articles/10.3389/fmicb.2018.02313

Smit, A., Hubley, R., & Green, P. (2013). *RepeatMasker Open-4.0*.

http://www.repeatmasker.org

Solares, E. A., Chakraborty, M., Miller, D. E., Kalsow, S., Hall, K., Perera, A. G., Emerson, J. J.,

& Hawley, R. S. (2018). Rapid Low-Cost Assembly of the *Drosophila melanogaster*

Reference Genome Using Low-Coverage, Long-Read Sequencing. *G3*

*Genes|Genomes|Genetics*, *8*(10), 3143–3154.

https://doi.org/10.1534/g3.118.200162

Stanke, M., Diekhans, M., Baertsch, R., & Haussler, D. (2008). Using native and syntenically

mapped cDNA alignments to improve de novo gene finding. *Bioinformatics*, *24*(5),

637–644. https://doi.org/10.1093/bioinformatics/btn013

Stanke, M., Schöffmann, O., Morgenstern, B., & Waack, S. (2006). Gene prediction in

eukaryotes with a generalized hidden Markov model that uses hints from external

sources. *BMC Bioinformatics*, *7*(1), 62. https://doi.org/10.1186/1471-2105-7-62

Tang, S., Lomsadze, A., & Borodovsky, M. (2015). Identification of protein coding regions in

RNA transcripts. *Nucleic Acids Research*, *43*(12), e78.

https://doi.org/10.1093/nar/gkv227

The UniProt Consortium. (2022). UniProt: The Universal Protein Knowledgebase in 2023.

*Nucleic Acids Research*, gkac1052. https://doi.org/10.1093/nar/gkac1052

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,

Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M.,

Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E.,

… van Mulbregt, P. (2020). SciPy 1.0: Fundamental algorithms for scientific

computing in Python. *Nature Methods*, *17*(3), Article 3.

https://doi.org/10.1038/s41592-019-0686-2

Vizoso, D. B., & Ebert, D. (2004). Within-host dynamics of a microsporidium with horizontal

and vertical transmission: *Octosporea bayeri* in *Daphnia magna*. *Parasitology*, *128*(Pt

1), 31–38. https://doi.org/10.1017/s0031182003004293

Wadi, L., & Reinke, A. W. (2020). Evolution of microsporidia: An extremely successful group

of eukaryotic intracellular parasites. *PLOS Pathogens*, *16*(2), e1008276.

https://doi.org/10.1371/journal.ppat.1008276

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C. A.,

Zeng, Q., Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: An Integrated Tool for

Comprehensive Microbial Variant Detection and Genome Assembly Improvement.

*PLOS ONE*, *9*(11), e112963. https://doi.org/10.1371/journal.pone.0112963

Whelan, T. A., Lee, N. T., Lee, R. C. H., & Fast, N. M. (2019). Microsporidian Introns Retained

against a Background of Genome Reduction: Characterization of an Unusual Set of

550   Introns. *Genome Biology and Evolution*, *11*(1), 263–269.

551   https://doi.org/10.1093/gbe/evy260

552 Williams, B. A. P., Williams, T. A., & Trew, J. (2022). Comparative Genomics of Microsporidia.

553   In L. M. Weiss & A. W. Reinke (Eds.), *Microsporidia: Current Advances in Biology* (pp.

554   43–69). Springer International Publishing. https://doi.org/10.1007/978-3-030-93306-

555   7_2

556 Yin, Y., Mao, X., Yang, J., Chen, X., Mao, F., & Xu, Y. (2012). dbCAN: A web resource for

557   automated carbohydrate-active enzyme annotation. *Nucleic Acids Research*, *40*(W1),

558   W445–W451. https://doi.org/10.1093/nar/gks479

559 Zhu, T., Liang, C., Meng, Z., Guo, S., & Zhang, R. (2017). GFF3sort: A novel tool to sort GFF3

560   files for tabix indexing. *BMC Bioinformatics*, *18*(1), 482.

561   https://doi.org/10.1186/s12859-017-1930-3