Disordered regions and folded modules in CAF-1 promote histone deposition in *S. pombe*

Fouad Ouasti[1,2,4,8], Maxime Audin[1,2,8], Karine Freon[3,4,5], Jean-Pierre Quivy[6], Mehdi Tachekort[1,2], Elizabeth Cesard[1,2], Aurélien Thureau[7], Virginie Ropars[1,2], Paloma F. Varela[1,2], Gwenaelle Moal[1,2], Ibrahim Soumana Amadou[3,4,5], Aleksandra Uryga[3,4,5], Pierre Legrand[7], Jessica Andreani[1,2], Raphael Guerois[1,2], Geneviève Almouzni[6], Sarah Lambert[3,4,5]*, Francoise Ochsenbein[1,2]*

[1]Institute Joliot, Commissariat à l'énergie Atomique (CEA), Direction de la Recherche Fondamentale (DRF), F91191 Gif-sur-Yvette, France

[2]Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ. Paris-Sud, Université Paris-Saclay, 91198, Gif-sur-Yvette cedex, France

[3]Institut Curie, PSL Research University, UMR3348, 91400 Orsay, France.

[4] Paris-Saclay University, 91400 Orsay, France.

[5]CNRS, UMR3348, 91400 Orsay France. Equipe Labellisée ligue contre le Cancer.

[6] Institut Curie, PSL Research University, CNRS, Sorbonne Université, Nuclear Dynamics Unit, Équipe Labellisée Ligue contre le Cancer, Paris, France.

[7]Synchrotron SOLEIL, HelioBio group, l'Orme des Merisiers, Départementale 128, 91190 Saint-Aubin, France.

[8]These author contributed equally

Correspondence: francoise.ochsenbein@cea.fr   Phone: 33-1-69089679   Fax: 33-1-69084712; mailto:Sarah.lambert@curie.fr

Abstract

Coupling histone deposition with DNA synthesis is important for genome and epigenome integrity in eukaryotes. Here, we reconstituted *Sp*CAF-1, the major histone deposition factor linked to DNA synthesis in fission yeast and characterized its structure using NMR, SAXS and molecular modeling. We reveal the unfolded nature of the acidic domain, and how it folds upon histone binding, while the long KER helix mediates DNA binding and stimulates *Sp*CAF-1 association with the polymerase accessory factor PCNA. By designing specific *Sp*CAF-1 mutants, we demonstrate that both histone and DNA binding are required for its functions *in vitro* and *in vivo*. We show that PCNA binding not only accelerates nucleosome assembly *in vitro* but is also essential for the proper targeting of the complex to the chromatin *in vivo*. Our structural and functional study reveals how the combination of disordered regions and folded modules in CAF-1 enable the dynamics of multiple interactions to promote histone deposition coupled to DNA synthesis.

Introduction

In eukaryotes, genomic DNA is packaged in a dynamic nucleoprotein complex, the chromatin, which protects DNA and regulates its accessibility. The fundamental repeat unit of chromatin, the nucleosome core particle, comprises 146 bp of DNA wrapped around a histone octamer including a tetramer of histone H3−H4 flanked by two dimers of H2A−H2B[1].. Histone chaperones are critical players in ensuring histone traffic and deposition. Without energy consumption, they escort histones, facilitate their transfer and deposition on DNA, and provide links with DNA based-processes such as DNA replication, repair and gene transcription[2]. In line with these key properties, perturbations of histone chaperones are associated with defects in genome and epigenome maintenance and function as found in cancer, aging and viral infections[3-5]. Discovered over thirty years ago[6], conserved in all eukaryotes[7], the histone chaperone Chromatin Assembly Factor 1 (CAF-1) is central and unique in promoting the deposition of replicative histones H3−H4 in a manner coupled to DNA synthesis, i.e. during DNA replication and repair, and is also involved in heterochromatin maintenance (see[2,8] for review). The unique feature of CAF-1 is that it provides a link with DNA synthesis via its association with the trimeric DNA polymerase processivity factor, Proliferation Cell Nuclear Antigen (PCNA), through PCNA Interacting Protein motifs (PIP)[9-13].

CAF-1 comprises three subunits (**Figure 1a**)[6,14,15]. While progress in uncovering its molecular/genetic properties derives from work in *Saccharomyces cerevisiae,* and biochemical work in human cells, there is still a lack of atomic information. In *cerevisiae,* CAF-1 is a hetero-trimer that binds to a single H3−H4 dimer[16,17] and induces a conformational rearrangement promoting interaction with the DNA. Two complexes co-associate to ensure the deposition of H3−H4 tetramers on DNA in the first step for nucleosome assembly (see[18] for review). Two domains of the large subunit Cac1 contribute to DNA binding[16,19], the conserved low complexity region called KER (for Lys, Glu and Arg rich) and the C-terminal Winged Helix Domain (WHD). These features are conserved in human CAF-1[13]. However, we still miss a complete view of the 3D organisation of the CAF-1 complex, and even most critically miss an understanding of whether its constitutive domains function in synergistic or independent manner to facilitate non only histone deposition but also contribute to a proper address in cells. Finally, the degree of conservation of these properties across species needs to be assessed.

To get further insights into CAF-1 function, we isolated the fission yeast complex (Pcf1-Pcf2-Pcf3) and investigated its binding mode with its three main partners, histones H3−H4, DNA and PCNA. Based on these structural insights, we designed targeted mutations to specifically alter Pcf1 interactions with DNA, PCNA and histones H3−H4. To probe their functions, we also analysed phenotypes of the corresponding mutants in fission yeast.

1 Results

2 Global organization of the full-length *Sp*CAF-1 complex

The large subunit of CAF-1, present in all major groups of eukaryotes, exhibits significant
sequence divergence (16% or 21% between *Sp*Pcf1 and *Sc*Cac1 or *Hs*CHAF1A/p150, respectively).
Given this high sequence divergence, conserved biochemical properties between *Sc*CAF-1 and *Sp*CAF-
1 should reveal important functional features. From sequence alignments, the 6 main conserved regions
previously proposed to contribute to the nucleosome assembly activity of CAF-1 can be inferred in
*Sp*Pcf1 sequence, a KER domain, a single PIP motif, an acidic domain (ED domain), the domains
predicted to bind Pcf2 (2BD) and Pcf3 (3BD) and a C-terminal WHD domain (**Figure S1a**). Although
*Sp*Pcf1 is shorter than *Sc*Cac1 and *Hs*CHAF1A/p150 (544 residues instead of 606 and 956, respectively)
its sequence includes a remarkable high abundance of predicted I̲ntrinsically D̲isordered R̲egions (IDRs)
(**Figure 1b**). These IDRs include the predicted histone-binding domain (Pcf1_ED), the PCNA (PIP
motif) and the DNA binding domain (Pcf-1 KER).

We produced and purified the three subunits of *Sp*CAF-1 separately from bacteria and insect
cells (**Figure 1c,** materials & methods). When isolated, both Pcf2 and Pcf3 are monomeric while Pcf1
forms large soluble oligomers (**Figure S1b**). Mixing the subunits by pairs, we observed stable
complexes for Pcf1-Pcf2 and Pcf1-Pcf3 by size exclusion chromatography (SEC). Pcf2 and Pcf3 did not
interact with each other (**Figure S1b, S1c**) suggesting that the large subunit Pcf1 mediates the complex
assembly. We next reconstituted and isolated the recombinant full-length (FL) *Sp*CAF-1 complex by
SEC (**Figure 1c**). An experimental molecular weight of 179 kDa was calculated using Small Angle X-
ray Scattering (SAXS), consistent with a 1:1:1 stoichiometry (**Figure S1d**). These data are in agreement
with a globular complex with a significant flexibility (**Figure S1e**).

To determine the extent of disorder in the large subunit of *Sp*CAF-1, Pcf1 was produced with
uniform ^{15}N (or ^{15}N-^{13}C) labeling. The CAF-1(^{15}N-Pcf1) complex with unlabeled Pcf2 and Pcf3 was
reconstituted and SEC-purified. Given the size of this complex (167 kDa), we expected that only amide
signals from residues in long disordered regions could be observed by Nuclear Magnetic Resonance
(NMR) spectroscopy. The ^{15}N-^{1}H spectrum shows about 140 amide signals, revealing that up to a quarter
of Pcf1 residues are intrinsically disordered in the full *Sp*CAF-1 complex (**Figure 1d**). These residues
are located in four continuous segments of Pcf1 and define intrinsically disordered regions that we
labeled IDR1 to IDR4 (**Figure 1b**). IDR1 corresponds to the ~50 N-terminal residues of the protein,
IDR2 (181-198) is located between the PIP motif and the 3BD region, IDR3 (355-394) overlaps a large
segment of the acidic ED domain and IDR4 (451-470) is located between the 2BD region and the C-
terminal WHD domain. The boundaries of the four IDRs are in agreement with the segments of Pcf1
predicted to harbor disorder with a high probability (**Figure 1b**).

We next build a model of the *Sp*CAF-1 complex using the AlphaFold2 multimer software (AF2)
with one copy of each full-length protein (**Figure 1e**, **Figures S1f-l**). The model is consistent with our

biochemical data showing that Pcf1 mediates the complex assembly. Also, in agreement with their disordered nature, low values around 0.2-0.3 of the local quality of the model as calculated by the Local Distance Difference Test (pLDDT) were obtained in the four IDR segments with a remarkable match for the delimitations of the four IDR segments by pLDDT values and NMR data (**Figure 1b**). Accordingly, these segments are symbolized with a dashed line in **Figure 1e**. In contrast, significantly high pLDDT was obtained for the 3BD, 2BD and WHD domains of Pcf1 and for the two subunits Pcf2 and Pcf3. (**Figure 1b**, **Figures S1f**). These data allowed to identify four independent modules, not predicted to interact with each other. The first module corresponds to the KER domain of Pcf1, predicted to form a long helix ending by the PIP motif. The second module contains the Pcf2 subunit, composed of 7 WD repeats arranged in a circular fold, and a segment of Pcf1 corresponding to the 2BD domain forming three short beta strands and a short helix (**Figure S1g-i**). In the third module, the 3BD domain of Pcf1 composed of 7 helices and 3 beta strands establishes a large interface with Pcf3, composed of 7 circular WD repeats (**Figure S1j-l**). The fourth domain is the WHD domain. We next used these models to fit our SAXS data allowing flexibility between the four modules. The best model fitted the experimental data with a high accuracy and is in agreement with a relatively globular complex. Superimposing the generated models did not define a unique orientation between the four modules, suggesting that the complex has an inter-module flexibility (**Table S1, Figure S1m-n**).

Taken together, our findings indicate that the large subunit Pcf1 mediates the (1:1:1) complex assembly. Pcf1 includes four IDR, and can organize its key regions (KER, PIP, 3BD, ED, 2BD and WHD) allowing them to be exposed and bound by Pcf2 and Pcf3 simultaneously.

Figure 1: The large *SpCAF-1* subunit includes four intrinsically disordered regions (IDR) a Names for the large, the medium and the small subunits of CAF-1 in *H. sapiens*, *S. cerevisiae* and *S. pombe*. **b** Upper panel: The magenta line shows the predicted disorder of Pcf1 (spot disorder software) and the black line the Cα Local Distance Difference Test (pLDDT) calculated for Pcf1 residues by the AlphaFold2 model of the full *SpCAF-1* complex. Lower panel: Cα chemical shift index calculated for the 101 assigned residues. This Cα chemical shift index is consistent with their disordered nature. The four IDR regions are highlighted with pink semi-transparent vertical bars. The predicted domains of Pcf1 are labeled. **c** General strategy for the production of *SpCAF-1*. The lower panel shows the purification SEC profile and the SDS-PAGE purity of the sample. **d** ¹H-¹⁵N SOFAST-HMQC spectrum of the FL *SpCAF-1* complex composed of uniformly labeled ¹⁵N-Pcf1 and unlabeled Pcf2 and Pcf3 (*SpCAF-1*(¹⁵N-Pcf1)). The assigned signals are labeled. **e** AF2 model of the *SpCAF-1* complex. The four IDR segments are shown with a dashed line. The relative orientation of the four modules is arbitrary.

In the FL *Sp*CAF-1 complex, the acidic ED domain is disordered but folds upon histone binding

We next investigated the interaction of *Sp*CAF-1 with histones H3−H4. A stable complex was isolated by SEC at low (150 mM NaCl) and high (1 M NaCl) salt concentrations, confirming that the reconstituted *Sp*CAF-1 complex tightly binds histones (**Figure 2a**). SAXS measurements at low salt allowed to calculate an experimental molecular weight of 193 kDa for this complex, showing that *Sp*CAF-1 binds a dimer of histones H3−H4 (**Figure S1d**). In addition, these data are compatible with a more extended shape compared to *Sp*CAF-1 alone (**Figure S1e**).

Addition of histones to *Sp*CAF-1(^{15}N-Pcf1) led to a drastic decrease in intensity of the NMR signal specifically for residues in the IDR3 segment (**Figure 2b**, **Figure S2a**). To further characterize this domain, we designed a short construct of Pcf1 (325-396), called Pcf1_ED, corresponding to the IDR3 segment extended in its N-terminus with the conserved acidic segment (325-355) (**Figure 2c, Figure S1a**). We confirmed the fully disordered nature of Pcf1_ED by NMR (**Figure S2b-c**). Signals corresponding to residues 355-394 (IDR3) remarkably overlap in the spectra of Pcf1_ED and *Sp*CAF-1(^{15}N-Pcf1) (**Figure S2a**), showing that this segment was fully flexible in *Sp*CAF-1, and did not interact with other regions of the complex. Upon binding of unlabeled histones H3−H4, we observed the vanishing of almost all NMR signals of ^{15}N Pcf1_ED as in the full *Sp*CAF-1 complex (**Figure 2d**). In contrast, a large part of signals (338-396) did not vanish anymore upon addition of a histone complex preformed with two other histone chaperones known to compete with CAF-1 for histone binding[16], Asf1−H3−H4−Mcm2(69-138) (**Figure 2d**). This region of the ED domain, is indeed in direct competition with Asf1 and Mcm2 whose histone binding modes are well established (**Figure 2e**)[20,21]. Fully consistent with this NMR competition experiment, this segment of Pcf1_ED domain was predicted by AlphaFold2 to interact with histones H3−H4 through the same surface as the one bound by Mcm2 (**Figure 2f, Figure S2d**). Two highly conserved positions in Pcf1, L359 and F380, are thus proposed to mediate histone H3−H4 binding in the same region as Mcm2 (**Figure 2f**). We next used these AlphaFold2 models to fit the SAXS curve of the *Sp*CAF-1−H3−H4 complex allowing reorientation of the different modules (**Figure S1o-p**). Remarkably, all generated models show a significant exclusion of the KER domain from the complex, suggesting that the KER domain of *Sp*CAF-1 becomes more accessible upon histone binding.

Figure 2: The acidic ED domain binds histones alone and in the full CAF-1 complex

Figure 2: The acidic ED domain binds histones alone and in the full *Sp*CAF-1 complex a SEC profile and the SDS-PAGE purity of *Sp*CAF1−H3−H4 histones at 150 mM NaCl and 1 M NaCl. **b** Mapping of the interaction between *Sp*CAF-1(^{15}N-Pcf1) and *Sp*H3−H4 histones, using the intensities ratio (I/I0), where I and I0 are the intensity of the signals ^{1}H-^{15}N SOFAST-HMQC spectra before and after addition of histones, respectively. **c** Sequence Logo of the ED domain generated with a large data set of Pcf1 homologues. The position of the two conserved residues Y340 and W348, mutated in ED* are indicated with stars and conserved Pcf1 L359 and F380 residues with five and four branch stars respectively. **d** Mapping of the interaction between Pcf1_ED or Pcf1_ED* with *Sp*H3−H4 histones using the intensities ratio (I/I0) as in b. Histones were added alone or previously complexed with histones chaperones. **e** Cartoon representation of the complex between human histones H3−H4 (light blue and cyan), Asf1 (light grey) and Mcm2 (dark grey) (PDB: 5BNX). **f** AlphaFold2 model of Pcf1 (353-385) (as red cartoon), corresponding to the segment of the ED domain indicated in red, in complex with histones H3−H4 (light blue and cyan surface) superimposed with Mcm2 and Asf1 as in panel e. The two insets represent zoomed views of the sidechains of the conserved Pcf1 L359 and F380 residues (red sticks) binding into H4 and H3 pockets, respectively. The same four and five branch stars are used to label these positions in the logo panel c.

The NMR competition experiment also reveals that an additional region of Pcf1_ED domain (338-351) is involved in the interaction with H3−H4 but is not competing with the Asf1-Mcm2 module. (**Figure 2d**). In order to alter the interaction of the ED domain with histones without modifying its charge and without interfering with Asf1 or Mcm2 binding, we identified from sequence alignments in this segment (**Figure 2c**), two invariant hydrophobic residues, Y340 and W348, that were mutated into alanines (mutant called ED*, **Figure S1a**). As expected, the isolated Pcf1_ED* domain showed almost no histone binding as observed by the intensity of ^{1}H-^{15}N NMR signals (**Figure 2d**). We next monitored the impact of the ED* mutations in the context of the full *Sp*CAF-1 complex. To do so, the mutations Y340A-W348A were introduced in the full length Pcf1, and the complex reconstituted with the uniformly ^{15}N labeled Pcf1(ED*) and unlabeled Pcf2 and Pcf3 (**Figure S2e**). The ^{1}H-^{15}N NMR spectrum of this mutant was similar that of the WT complex, but upon addition of unlabeled histones H3−H4 no major change was observed (**Figure S2a**), which strongly suggest an alteration of the histone binding of this mutant.

In summary, we identified critical amino-acids in the ED domain involved in H3−H4 binding. We also showed that addition of histones leads to a conformational change in the *Sp*CAF-1 complex with less disorder in the ED domain and an increased accessibility of the KER domain.

*Sp*CAF-1 binds dsDNA longer than 40bp

We next analyzed the DNA binding properties of *Sp*CAF-1. Electrophoretic mobility shift assays (EMSA) were performed with a DNA ladder as substrate in order to determine the minimal DNA size for *Sp*CAF-1 binding. The complex *Sp*CAF-1 showed significant binding for DNAs longer than 40bp (**Figure S3a**). EMSAs with a double-stranded 40bp DNA fragment confirmed the homogeneity of the bound complex. When increasing the *Sp*CAF-1 concentration, additional mobility shifts suggest, a cooperative DNA binding (**Figure 3a**). Micro-Scale Thermophoresis (MST) measurements were next performed using an alexa-488 labeled 40bp dsDNA (**Figure 3b, Table 1**). The curves were fitted with a Hill model with a EC50 value of $0.7 \pm 0.1 \mu M$ and a cooperativity (Hill coefficient, h) of 2.7 ± 0.2, in line with a cooperative DNA binging of *Sp*CAF-1.

Table 1: Experimental affinities of different *Sp*CAF-1 constructs with a 40bp dsDNA measured by Microscale thermophoresis (MST).

Construct	EC50 (μM)	Hill coeff., h
Pcf1_KER	1.1 ± 0.2	3.3 ± 0.5
Pcf1_KER*	12.2 ± 0.7	1.5 ± 0.3
Pcf1_KER-PIP	1.9 ± 0.3	5.2 ± 0.9
Pcf1_WHD	Not detected	Not detected
*Sp*CAF-1 WT	0.7 ± 0.1	2.7 ± 0.2
*Sp*CAF-1(ΔWHD)	0.7 ± 0.1	2.3 ± 0.3
*Sp*CAF-1(KER*)	2.8 ± 0.4	1.3 ± 0.3
*Sp*CAF-1(ED*)	1.0 ± 0.1	2.3 ± 0.1
*Sp*CAF-1(PIP*)	0.7 ± 0.1	2.7 ± 0.3

The KER domain is the main DNA binding region of *Sp*CAF-1

The KER and WHD domains of the CAF-1 large subunit were shown to be involved in DNA binding in *Sc*CAF-1 and *Hs*CAF-1[13,16,19,22]. We were thus interested to explore the conservation of these features in Pcf1. We first isolated the KER domain (Pcf1_KER, **Figure S1a**), and an extended fragment we called Pcf1_KER-PIP (**Figure S1a**), which includes the PIP motif (Q_{172}-L-K-L_{175}-N-N-F_{178}-F_{179}). These domains are predicted by AlphaFold2 to form a long helix with partial disorder at both ends and possible extension over the first half of the PIP motif (**Figure 1d, Figure 3c**). Using a combination of circular dichroism (CD) (**Figure S3b-c**), SEC-SAXS (**Figure S1d-e, Figure S3d**) and NMR (**Figure S3e-f**), we confirmed that the isolated KER domain of Pcf1 domain forms a straight monomeric helix, partially continuing over the PIP motif. This long helix exhibits a strong bias in amino acid composition and remarkably, almost all basic residues are positioned on the same side of the helix (**Figure 3d**) providing a suitable interface for DNA binding[13]. We performed EMSA using a DNA ladder as substrate and we found that Pcf1_KER domain binds DNA that are longer than 40bp, as observed with the full *Sp*CAF-1 complex (**Figure S3e**). EMSAs with double strand 40bp DNA fragment showed the presence of multiple bands for Pcf1_KER bound DNA, indicating a possible cooperative DNA binding of this fragment (**Figure 3d**). Affinity measurements by MST led to a EC50 of 1.1 ± 0.2 µM for Pcf1_KER with a cooperativity around 3, consistent with EMSA experiments (**Figure 3d, Table 1**). The DNA binding properties of Pcf1_KER-PIP are comparable to that of Pcf1_KER (**Figure 3e, Table 1, Figure S3f**). The EC50 obtained for the isolated Pcf1_KER are also close that of the full *Sp*CAF-1 complex (0.7 ± 0.1µM) suggesting that the KER domain constitutes the principal DNA binding domain of *Sp*CAF-1.

Based on these results, we designed a mutant called Pcf1_KER* with a charge inversion for five positive residues at the C-terminus of the potential DNA binding face of the KER helix (R147E-K150E-K154E-R161E-K168E) (**Figure 3c**, **Figure S1a**). The CD analysis of Pcf1_KER* shows this mutant is mainly helical (**Figure S3g**). MST quantification confirmed that the mutation of Pcf1_KER* impaired DNA binding by a factor of 10 even-though residual DNA binding remained (**Figure 3e, Figure S3h-j, Table 1**). The KER* mutation was then introduced in the full complex *Sp*CAF-1(KER*) (**Figure S1a, Figure S2e**) and we confirmed by MST and EMSA its lower affinity for dsDNA (**Figure 3b, Figure S3k, Table 1**). Importantly, the NMR signals of all IDR for this mutant with or without histones were close to that of the WT (**Figure S3l-m**) indicating that the KER* mutation did not impair histone binding.

Figure 3: Pcf1_KER is the main DNA binding domain of *Sp*CAF-1: a EMSA with *Sp*CAF-1 and 40 dsDNA (1µM) revealed with SYBR SAFE staining. **b** Microscale thermophoresis (MST) fitted curves of *Sp*CAF-1 WT and mutants with 40bp dsDNA. **c** Upper panel: Modelled structure of the Pcf1_KER-PIP domain (56-185) rainbow coloured according to the pLDDT of each residue. Red corresponds to pLDDT values of 1 and dark blue of 0. Middle panel same model represented with its electrostatic surface. Lower panel: zoom of the C-terminus of the KER domain and the PIP motif. The five mutated residues are labeled and highlighted with spheres. **d** EMSA of Pcf1_KER binding with a 40bp dsDNA (1µM) revealed with SYBR SAFE staining **e** MST fitted curves of Pcf1_KER constructs and mutants with 40bp dsDNA. **f** Overlay of the calculated model of the WHD domain obtained with the CS-rosetta software (light orange) using NMR assignments of the domain (**Figure S3n**) , with AlphaFold2 (gold) and the structure of Cac1 WHD from budding yeast (PDB 5jbm, in grey)[23](Grey). **g** EMSA revealed with SYBR SAFE staining of Pcf1_WHD domain with a 40 dsDNA (1µM).

The C-terminal of Pcf1 folds as a WHD domain but does not bind DNA

We next isolated the Pcf1_WHD domain (**Figure S1a**) and confirmed by NMR and AlphaFold2 that its global fold is similar to *Sc*WHD [19,23] (**Figure 3f, Figure S3n**). Unexpectedly, Pcf1_WHD does not interact with DNA of any size (**Figure 3g, Figure S3n**). The residues involved in DNA binding in *Sc*WHD, K564 and K568, correspond to S514 and G518 in *Sp*WHD, respectively, leading to a different electrostatic surface, probably not favorable for DNA binding (**Figure S3p-q**). To further investigate the role of the WHD domain of *Sp*CAF-1, the WHD domain was deleted in the reconstituted *Sp*CAF-1(ΔWHD) complex (**Figure S1a Figure S2e**) and analyzed by EMSA and NMR. We observed the similar DNA binding property and IDR properties for *Sp*CAF-1(ΔWHD) and the WT complex (**Table 1, Figure 3b, Figure S3k-m**).

Together our results show that the KER domain constitutes the main DNA binding region of *Sp*CAF-1 and that the WHD domain does not contribute to this binding.

Crosstalk between DNA and PCNA binding

The PIP motif of Pcf1 was found crucial for *Sp*CAF-1 interaction with PCNA *in vivo*[12]. Given its proximity with the KER domain, we further investigated potential cross-talks between PCNA and DNA binding. We first measured by isothermal microcalorimetry (ITC) an affinity of 7.1±1.3 μM between *Sp*PCNA and a short PIP motif segment (**Figure S4a**, **Table 2**). This affinity is in the same range (2-fold less affine) as a peptide isolated from the replicative polymerase delta from *S. pombe* Cdc27 (**Figure S4a**, **Table 2**). In agreement with its consensus sequence, the binding mode of Pcf1_PIP motif to *Sp*PCNA is predicted by AlphaFold to be canonical (**Figure S4b**). Consistently, no binding was observed for the Pcf1_PIP* peptide with 4 Alanine mutations, previously designed to disrupt the PIP motif (**Figure S4a**, **Table 2**)[12]. We next measured the affinity of the longer fragment Pcf1_KER-PIP for *Sp*PCNA and observed an affinity gain of a factor 10 (0.7 ± 1.3 μM) (**Figure S4c, Table 2**), revealing interactions between the KER domain and PCNA. ITC also fits a stoichiometry of ~2 Pcf1_KER-PIP per PCNA trimer, suggesting that, in each PCNA trimer, one monomer remains unbound and potentially accessible for binding to other partners. Pcf1_KER-PIP* did not interact with PCNA confirming the importance of the PIP motif for this association (**Figure S4c, Table 2**). The KER* mutation impaired the interaction of Pcf1_KER*-PIP with PCNA of a factor 10 reaching the affinity of the short isolated Pcf1 PIP peptide (**Figure S4c, Table 2**). Collectively these results show that both the PIP motif and the C-terminal part of the KER domain are involved in PCNA binding.

To reveal possible crosstalk between CAF-1 binding to PCNA and DNA, we analysed, in the presence or absence of dsDNA, the binding of the full *Sp*CAF-1 complexes (WT *Sp*CAF-1, *Sp*CAF-1(PIP*), *Sp*CAF-1(ED*), *Sp*CAF-1(KER*) and *Sp*CAF-1(ΔWHD)) with recombinant *Sp*PCNA, using EMSA (**Figure 4a, Figure S4d**). For all combinations tested, we quantified binding by monitoring the

disappearance of free PCNA (**Figure 4b**) and free DNA (**Figure 4c**). In this assay, only 20% of free PCNA intensity was lost by addition of DNA (**Figure 4b**), probably because the PCNA trimer can slide along the linear DNA and dissociates during the migration. In the absence of DNA, we observe a small but significant decrease of free PCNA upon addition of WT *Sp*CAF-1, in agreement with the relatively low binding affinity between Pcf1_KER helix and PCNA (**Table 2**). In contrast, in the presence of dsDNA, addition of an excess of WT *Sp*CAF-1 leads to the complete disappearing of the free PCNA band and to a large shift of the band corresponding to *Sp*CAF-1-DNA, corresponding to a larger complex engaging CAF-1, PCNA and DNA (**Figure 4a**). *Sp*CAF-1(ED)* and *Sp*CAF-1(ΔWHD) show similar binding properties for PCNA and DNA compared to WT CAF-1. In contrast, *Sp*CAF-1(PIP*) binds DNA like the WT, but is strongly impaired for PCNA binding alone and in the presence of DNA, while *Sp*CAF-1(KER*) is impaired for binding both DNA and *Sp*PCNA. In agreement, the large shifted band corresponding to a *Sp*CAF-1−PCNA−DNA complex is not observed for these two mutants (**Figure S4d**).

Altogether, our data show the stabilization of thee CAF-1−PCNA interaction by DNA that requires both the KER domain and the PIP motif but not the ED and WHD domain. Conversely, the capacity of CAF-1 to bind PCNA does not impair its interaction with DNA.

Figure 4: The *Sp*CAF-1(KER*) mutant is affected for PCNA binding. a EMSA showing interactions of purified *Sp*CAF-1 (at the indicated concentrations), with or without recombinant *Sp*PCNA (3µM) in the presence and absence of 40bp dsDNA (1µM), revealed with Coomassie blue (upper panel) and with SYBR SAFE staining (lower panel). **b** Quantification of bound *Sp*PCNA in the EMSA shown in panel **a** and in **Figure S4d** for *Sp*CAF-1 and mutants. Values are indicated in % compared to the free PCNA reference (PCNA alone in line 1 in panel **a**) after addition of *Sp*CAF-1 (WT or mutant) at the indicated concentration and in the presence (filled bars) or absence (dashed bars) of 40bp dsDNA (1µM). **c** Quantification of bound DNA for EMSA shown in panel **a** and in **Figure S4d** for *Sp*CAF-1 and mutants. Bound DNA in % is compared to the free DNA reference (line 5 in panel **a**) after addition of *Sp*CAF-1 (WT or mutant) at the indicated concentration and in the presence (filled bars) or absence (dashed bars) of *Sp*PCNA (3µM). All experiments were done in duplicates. Mean values are indicated and error bars shows their standard deviation.

Table 2: Interactions parameter with *Sp*PCNA measured by isothermal microcalorimetry (ITC): *The stoichiometry (N) is calculate as a molar ratio of monomeric PCNA.

Ligand	Kd (μM)	ΔG (kCal.M^{-1})	N*	ΔH (kCal.M^{-1})	-TΔS (kCal.M^{-1})
Pcf1_PIP	7.1 ± 1.3	-6.9 ± 0.1	0.97 ± 0.08	-2.9 ± 0.2	-0.39 ± 0.3
Pcf1_PIP*	undetectable	ND	ND	ND	ND
Pcf1_KER-PIP	0.7 ± 0.2	-8.2 ± 0.2	0.64 ± 0.04	+2.9 ± 0.6	-11.2 ± 0.8
Pcf1_KER*-PIP	7.1 ± 1.5	-6.9 ± 1.2	0.7 ± 0.2	+1.0 ± 0.5	-7.9 ± 0.7
Pcf1_KER-PIP*	undetectable	ND	ND	ND	ND
Cdc27_PIP	3.5 ± 0.3	-7.3 ± 0.1	0.9 ± 0.1	-4.8 ± 0.02	-2.4 ± 0.1

In vitro histone deposition properties of *Sp*CAF-1 mutants

We next examined the ability of the full *Sp*CAF-1 complex reconstituted with the four Pcf1 mutants (*Sp*CAF-1(PIP*), *Sp*CAF-1(ED*), *Sp*CAF-1(KER*), *Sp*CAF-1(ΔWHD)) to promote nucleosome assembly mediated by CAF-1 in a complex environment closer to physiological conditions. We used Xenopus high speed egg extract (HSE) that are powerful systems competent for chromatin assembly and effective to exploit depletion/complementation assays[24]. We depleted HSE for the endogenous Xenopus CAF-1 largest subunit (xp150) and assessed the capacity of *Sp*CAF-1(PIP*), *Sp*CAF-1(ED*), *Sp*CAF-1(KER*) and *Sp*CAF-1(ΔWHD) to complement these xp150-depleted extracts[16,24] (**Figure S5**). We monitored nucleosome assembly coupled to DNA synthesis using as a template a circular UV-damaged plasmid enabling to analyse by supercoiling assay and nucleotide incorporation simultaneously both repair synthesis and nucleosome formation (**Figure 5**)[10]. We verified that p150-depleted HSE lacked the capacity to promote nucleosome assembly on labeled DNA when compared to mock depleted HSE, and that the recombinant WT *Sp*CAF-1 complex efficiently rescued the loss of xp150 as attested by the detection of supercoiled form I. In contrast, when we complemented the depleted extract with *Sp*CAF-1 mutant complexes *Sp*CAF-1(ED*), *Sp*CAF-1(KER*), *Sp*CAF-1(ΔWHD) we did not detect the supercoiled form I. This indicates that these mutants cannot promote nucleosome assembly (**Figure 5**). When we used the *Sp*CAF-1(PIP*) mutant, we did not detect supercoiling on labeled DNA at 45 minutes, yet at 2 hours supercoiling ultimately reached levels achieved using the WT *Sp*CAF-1 (**Figure 5**, bottom, synthesized DNA). Interestingly both for 45 and 2 hours of assembly *Sp*CAF-1(PIP*) mutant yielded more supercoiling than any of the *Sp*CAF-1(ED*), *Sp*CAF-1(KER*), *Sp*CAF-1(ΔWHD) mutants. Thus, while mutation in the PIP motif of Pcf1 impaired chromatin assembly at a short time, when more time is given, it allows ultimately to catch up with the wild type. In contrast, none of the *Sp*CAF-1(ED*), *Sp*CAF-1(KER*), *Sp*CAF-1(ΔWHD) mutants could catch up, leading to a *Sp*CAF-1 complex deficient for nucleosome assembly even after longer incubation time. Therefore, these data validate the important role of the amino-acids Y340 and W348 within the ED domain in Pcf1 and the importance to preserve the integrity of the KER and WHD domain to ensure a proper *Sp*CAF-1 mediated nucleosome assembly on synthesized DNA.

Together, these results indicate that the PIP domain provides Pcf1 with the ability to accelerate

nucleosome assembly, yet the integrity of the ED, KER and WD domain proved absolutely mandatory

for an efficient *Sp*CAF-1 mediated nucleosome assembly.

Figure 5: Rescue using Y340 and W348 in the ED domain, the intact KER DNA binding domain and the C-
terminal WHD of Pcf1 in *Sp*CAF-1 mediated nucleosome assembly. Supercoiling analysis after 45 (left) and
120 (right) minutes incubation to monitor chromatin assembly in control mock and xenopus p150-depleted HSE.
Total DNA visualized by EtBr staining (top) and synthesized DNA visualized by biotin detection (bottom) are
shown. The xenopus p150-depleted HSE is either mock complemented (-) or complemented using *Sp*CAF-1
complex composed of wild type Pcf1(WT) or mutants Pcf1(PIP*), Pcf1_ED*, Pcf1(KER*), or Pcf1(ΔWHD) as
indicated. T: pBS plasmid incubated without extract run in parallel serves as a migration control to locate
supercoiled DNA. The position of relaxed (II) and supercoiled (I) DNA are indicated.

Association of *Sp*CAF-1 with histones impacts PCNA interaction *in vivo*

We next investigated the consequences of the four Pcf1 mutations previously characterized *in*

vitro, on *Sp*CAF-1 function *in vivo* by introducing the respective mutations at the endogenous *pcf1* gene.

Both *WT* and mutants were FLAG tagged in their N-terminal part. Immuno-blot of total cell extract with

anti-flag antibody showed that all mutated forms of Pcf1 were expressed to the same level than *WT* Pcf1

(**Figure S6a**).

We first tested PCNA–Pcf1 interaction by co-immunoprecipitation of FLAG-Pcf1 and found

that Pcf1(ΔWHD) showed a similar PCNA interaction than *WT* Pcf1 (**Figure 6a-b** and Figure **S6b**). No

interactions were detected with Pcf1(PIP*) and Pcf1(KER*), in line with the requirement of the KER

and PIP domains for PCNA binding (**Figure 4, Table 2**). Surprisingly, we found that Pcf1(ED*) binds

eight times more to PCNA than the *WT* Pcf1 (**Figure 6a-b**) although the corresponding *Sp*CAF-1(ED*)

bound PCNA with or without DNA *in vitro*, similarly to *WT* (**Figure 4**, **Figure S4d**). This suggest an

interplay *in vivo* between the binding of CAF-1 to PCNA and its capacity to bind histones.

Figure 6: Association of CAF-1 with histone is coupled to PCNA interaction *in vivo*. a Anti-FLAG Pulldown to address PCNA−CAF-1 interaction *in vivo* in indicated strains. **b** Quantification of bound PCNA from (**a**). **c** Example of Pcf2-GFP foci in living cells in indicated strains, according to cell morphology. **d** Quantification of cells showing Pcf2-GFP foci, according to cell morphology in indicated strains. Values are means of at least 3 independent experiments ± standard error. At least 1000 nuclei were analysed per strain. P values are indicated with stars and were calculated using the student test.

To probe this further, we analyzed *Sp*CAF-1 foci that were reported to colocalize with PCNA during DNA replication[12]. Since previously reported GFP-tagged forms of Pcf1 are not fully functional, we made use of cells expressing Pcf2-GFP, a functional tagged form[25]. As expected, Pcf2-GFP formed discrete foci during the bulk of S-phase (septated cells) until late S/early G2 phase (late cytokinesis cells) but not during G2 phase (mono-nucleated cells) in a Pcf1-dependent manner (**Figure 6c-d**). The *pcf1(ΔWHD)* mutation behaved like the *WT* in this assay. In contrast, S-phase Pcf2 foci were undetectable when Pcf1−PCNA interaction is impaired (in *pcf1(KER*)* and *pcf1(PIP*))*. Interestingly, Pcf2-GFP foci were more frequent in all cell cycle phases in *pcf1(ED*)* mutated cells compared to *WT*. Simultaneous acquisition of GFP fluorescence in living *WT* and mutated *pcf1* cells revealed that Pcf2-GFP foci were more abundant and brighter in *pcf1(ED*)* cells compared to *WT* (**Figure S6c**), suggesting a higher concentration of CAF-1 within replication factories. In conclusion, the ability of CAF-1 to localize to replication factories correlates with its association with PCNA *in vivo*, possibly modulated by the histone binding.

The WHD domain specifies CAF-1 function in distinct cellular processes.

In *S. pombe*, CAF-1 is involved in the replication-coupled maintenance of heterochromatin[15]. We employed a strain in which *ura4+* is inserted at the peri-centromeric heterochromatin of the chromosome I (**Figure 7a**, top panel). The expression of *ura4* is repressed by the surrounding heterochromatin resulting in a poor growth on uracil-depleted media and resistance to 5-fluoro-orotic acid (5FOA) (**Figure 7a**, bottom panel). As previously reported, the deletion of *pcf1* resulted in a better cell growth on uracil-depleted media compared to *WT* cells, showing that the heterochromatin is not properly maintained, leading to the derepression of *ura4+*. All mutants, excepted *pcf1(ΔWHD)*, exhibited defects in *ura4* silencing, similar to the one observed in the null mutant. This shows that the inability to interact with histone, PCNA and DNA results in a complete lack of CAF-1 function in maintaining heterochromatin. Interestingly, the WHD domain, while required for chromatin assembly *in vitro* (**Figure 5**), is dispensable for the maintenance of heterochromatin. We thus investigated further the role of this domain.

We analyzed the accumulation of Rad52-GFP foci as a readout of global accumulation of DNA damage (**Figure 7b**). The deletion of *pcf1* led to a modest but significant increase in the frequency of cells showing Rad52-GFP foci. A similar effect was observed in *pcf1(ED*)* mutated cells, while the presence of a CAF-1 complex unable to interact with PCNA resulted in a greater increase (in *pcf1(PIP*)* and *pcf1(KER*)* mutants). In contrast, no significant increase was observed in *pcf1(ΔWHD)* cells. Thus, both CAF-1 interaction with histone and PCNA prevent the accumulation of DNA damage, but histone deposition is not absolutely required.

The deletion of *pcf1* is synthetic lethal with the deletion of *hip1*, the gene encoding one subunit of the fission yeast HIRA complex[25], indicating that in the absence of replication-coupled histone deposition by CAF-1, cell viability relies on H3−H4 deposition by HIRA as suggested in human[26]. We found that *pcf1(KER*)*, *pcf1(ΔWHD)* or *pcf1(PIP*)* are co-lethal with *hip1* deletion (**Figure 6d**). Cells harboring *pcf1(ED*)* were viable when combined with *hip1* deletion, but exhibited a severe growth defect (**Figure 6a**), suggesting that CAF1(ED*) complexes can still perform some histone deposition *in vivo*. These genetic interactions indicate that binding of CAF-1 to PCNA, DNA and histones are critical determinants for its function *in vivo*, as well as the WHD C-terminal domain.

Figure 7 : The WHD domain of SpCAF-1 specifies CAF-1 function. a Top: Schematic representation of the silencing assay used. Otr: outer repeats, imr; inner repeats: cnt1; central core of the centromere 1. Bottom: Serial fivefold dilution on indicated strains on indicated media. **b** Example of Rad52-GFP foci in *WT* cells. **c** Quantification of Rad52-GFP foci in indicated strains. Values are means of at least 3 independent experiments ± standard error of the mean. P values are indicated as stars and were calculated with the student test. At least 1000 nuclei were analyzed per strain. **d**. Co-lethality assay. Tetrad dissections of cells deleted for *hip1* (*hip1Δ*) crossed with cells deleted for *pcf1* (*pcf1Δ*) (grey) or harbouring *pcf1(PIP*)* (magenta), *pcf1(ED*)* (red), *pcf1(KER*)* (purple) or *pcf1(ΔWHD*)* (orange). Spores with double mutations are surrounded.

Discussion

In the present work, we provide a comprehensive study of the histone chaperone CAF-1 from *S. pombe*. Despite the low sequence conservation between orthologues of the large subunit of CAF-1, Pcf1 from *Sp*CAF-1 mediates the heterotrimer complex that binds dimeric histones H3−H4, as does *Sc*CAF-1 [16,27,28]. Using AlphaFold2, we built a structural model of the *Sp*CAF-1, fully compatible with all our experimental data (**Figure 1**). This structure defines the 2BD and 3BD regions in Pcf1 as involved in the binding of Pcf2 and Pcf3, respectively. This matches remarkably the corresponding segments identified by HDX in *Sc*CAF-1 [28]. In line with previous observations in *Hs*CAF-1 [29] and *Sc*CAF-1 [17,23,28], the ED domain of *Sp*CAF-1 is crucial for histone binding. (**Figure 2**). Mainly disordered in the free chaperone, we show that this domain folds upon histone binding, promoting a conformational change with increased accessibility of the KER domain (**Figure S1**). *Sp*CAF-1 binds dsDNA longer than 40bp in the micromolar affinity range (**Figure 3, Table 1, Figure S3**) through the KER domain forming a long monomeric helix with a positively charged face. Interestingly, the helix length roughly corresponds to the size of 40bp dsDNA, suggesting that it could lie on DNA and act as a DNA ruler to sense free DNA for histone deposition [13,16,30]. Together, our findings highlight the conservation of CAF-1 properties in histone deposition mechanism *in vitro*, and thus unifies the current model [18].

This work revealed strong interdependency between histone deposition by CAF-1 and its association with PCNA. The PIP* mutation did not compromise DNA binding of *Sp*CAF-1 *in vitro* (**Figure S4**). Conversely, upon interaction with DNA, *Sp*CAF-1 interacted tighter with PCNA (**Figure 4**), consistently with a recent study in budding yeast [31]. We show that *Sp*CAF-1(PIP*) is still able to assemble histones *in vitro*, although slower than WT-*Sp*CAF-1. In contrast, *in vivo*, *pcf1(PIP*)* phenocopy the deletion of *pcf1*. From these results, we conclude that the binding of *Sp*CAF-1 to PCNA though the PIP motif is required for *Sp*CAF-1 functions *in vivo*, by allowing its recruitment and efficient histone deposition at DNA synthesis sites.

*Sp*CAF-1(ED*) showed a stronger interaction with PCNA than WT-*Sp*CAF-1 *in vivo*, and was more retained in replication foci (**Figure 6**). This default may not result from a direct competition between PCNA and histones for CAF-1 association since *Sp*CAF-1-ED* and WT-*Sp*CAF-1 show similar interaction with DNA and PCNA *in vitro* (**Figure 4, Figure S4**). In human cells lacking new histones, PCNA accumulates on newly synthetized DNA, and PCNA unloading has recently linked to histone deposition in budding yeast [32-34]. We propose that the accumulation of CAF-1 at replication foci in the ED* mutant may reflect PCNA recycling defects. This cannot be attributed to the inability of *Sp*CAF-1(ED*) to deposit histones otherwise similar accumulations we should have observed the same accumulation for *Sp*CAF-1(ΔWHD) also defective for histone deposition. The ED* mutation could rater interfere with other interactions, or with Post-Translational Modifications (PTMs) contributing to recycle PCNA.

Deletion of the WHD domain allowed separating *Sp*CAF-1 functions in chromatin assembly, heterochromatin maintenance and the prevention of DNA damage. Unlike *Sc*CAF-1[17,19,23,28] and *Hs*CAF-1[28], Pcf1_WHD did not bind DNA nor the ED domain (which remains fully disordered) in the free chaperone. Nevertheless, on the NMR spectra of the free and histone bound *Sp*CAF-1(^{15}N-Pcf1), the resonances of the isolated WHD domain are not present (**Figure 1d, S3n**), in agreement with a restricted movement of this domain that could likely interacts with other folded parts of the complex. *In vitro*, we found no impact of the WHD deletion on CAF-1 interaction with DNA, histones or PCNA, but the *Sp*CAF-1(ΔWHD) was deficient for histone deposition. Thus, the synthetic lethality of this mutant with *hip1* most likely reflects a replication-coupled assembly defect. Unexpectedly, this defect does not cause a problem of heterochromatin maintenance or damage accumulation, indicating that the WHD domain contributes to specify CAF-1 functions. Further investigations will be necessary to understand the role of this domain.

We reveal that disorder is a fundamental feature of Pcf1 supporting its molecular functions. First, the ED domain is disordered in the FL complex and folds upon histone binding. Second, four IDRs demarcate specific domains within Pcf1. We believe that these unfolded regions provide unique 'plasticity' properties to Pcf1 allowing these domains to bind concomitantly their multiple specific partners (Pcf1, Pcf3, PCNA, DNA and histones). We also reveal that although these domains individually bind their specific partners, there is an important crosstalk between them as exemplified by the fact that DNA stabilizes the CAF-1−PCNA interaction. Such plasticity and cross-talks provided by structurally disordered domains might be key for the multivalent CAF-1 functions.

Materials and Methods

Plasmid preparation for recombinant protein production

The cDNA sequence of WT Pcf1 (codon optimized for *E.coli* expression) was synthetized and

inserted into the pCM153 plasmid to obtain the recombinant MBP–6His-TEV cleavage site-Pcf1 protein

(named MBP-Pcf1 below). The cDNA sequence of WT Pcf2 and WT Pcf3 (codon optimized for insect

cells expression) were synthetized and introduced into a pKL plasmid for protein expression in insect

cells (MultiBac approach[35]) with either a C-terminal (for Pcf2) or a N-terminal (for Pcf3) 6His tag with

a TEV cleavage site between the protein and the His tag. Pcf1_ED (325-396) and Pcf1_WHD (471-544)

were sub cloned in frame into pET28A-B18R plasmid for expression with a N-terminal 6His-SUMO

tag. Pcf1_KER (56-170) and Pcf1-KER-PIP (56-185) were inserted in frame into pCM153 plasmid[36]

for expression with a N-terminal 6His-MBP-TEV tag. The cDNA sequence if *S. pombe* histones H3−H4

(codon optimized for *E.coli* expression) were introduced in the 6His-dAsf1 from the pET28 plasmid

(generous gift from R.N. Dutnall) in place of histones *Dm*H3-H4[37]. With this vector, histones H3−H4

are coexpressed with the chaperone ASF1, leading to soluble untagged free-histones, and ASF1-bound

histones. The cDNA of *Sp*PCNA (codon optimized for *E.coli* expression) was synthetized and inserted

into the pET28A-B18R plasmid for expression with an N-terminal 6His-SUMO. Pcf1 mutants were

generated by PCR. All plasmids for recombinant protein expression were constructed by GenScript.

Recombinant protein production

Pcf1 was overexpressed in *E.coli*. After fresh transformation of *E. coli* BL21 (DE3) Star cells

(Thermo Fisher Scientific), cells were grown in an auto-induction rich medium Terrific Broth (12 g/L

tryptone, 24 g/L yeast extract) containing 50 μg/mL of Kanamycin for 30 hours at 20°C, under agitation.

*Sp*Histones H3-H4, *Sp*PCNA and the all domains of Pcf1 were overexpressed in *E.coli*. The plasmid for

expressing the desired protein was freshly transformed in *E. coli* strain BL21 DE3 STAR (Thermo

Fisher Scientific). Cells were grown at 37 °C in a LB medium containing 50 μg/mL of Kanamycin until

OD reached 0.7 and recombinant protein expression was induced for 16 hours at 20 °C under agitation

by adding 1 mM isopropy β-D-1-thiogalactopyranoside IPTG, or cells were grown 30 hours at 20 °C in

a ZY auto-inducible medium. For ^{15}N or ^{13}C uniformly labeled proteins, the expression was made in

minimal media with 0.5g/L of ^{15}NH$_4$Cl and/or 2g/L of ^{13}C-glucose. Pcf2 and Pcf3 were produced in

insect cells. Sf9 Insect cells were infected with an MOI of 5*10-3 virus/cell and incubated for 5 days at

27 °C at 130 rpm. After centrifugation, cell pellets stored at -70°C until further use.

Protein purifications

Purification of Pcf1

Cells were pelleted by centrifugation and resuspended in the lysis buffer LB1 for 30 minutes

(50 mM Tris-HCl pH 8, 500 mM NaCl, 5% glycerol, 0.1% Triton X-100, 2 mM DTT, 5 mM MgCl2,

0.5 mM PMSF, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail, 1.2 mg/mL lysozyme and 70 U/mL of benzonase). Cells were lysed by sonication at 4°C, the lysate was clarified by centrifugation at 5°C at 18 500 rpm for 30min and loaded onto gravity flow amylose resin (NEB) previously equilibrated with buffer WB1_1 (50 mM Tris-HCl pH 8, 500 mM NaCl, 2 mM DTT). After loading the cell lysate onto the resin, the resin was washed with 5 column volumes of buffer WB1_1 to ensure complete passage of the cell lysate through the resin. Then, the resin was further washed with10 column volumes of buffer WB1_2 (50 mM Tris-HCl pH 8, 1000 mM NaCl, 2 mM DTT) to remove non-specific binding,before re-equilibration with 10 column volumes of buffer WB1_1. MBP-Pcf1 was eluted with 10 column volumes of buffer EB1 (50 mM Tris-HCl pH 8, 500 mM NaCl, 0.5 mM TCEP, 10 mM maltose and 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail,). After addition of 1 mM MgCl, the eluate containing MBP-Pcf1 was incubated 16 hours at 5°C with TEV protease (added with a ratio 1/20 in mass). The elate was then concentrated to 300µL (with Amicon® Ultra-15 30kDa filter concentrators), 2000 U of benzonase were added and incubated for 2 hours.The concentrated eluate was injected into a column Superose™ 6 increase 10/300 GL (Cytiva) previously equilibrated with the final buffer FB1 (50 mM Tris-HCl pH 8, 500 mM NaCl, 1 mM DTT). The Pcf1-containing fractions were pooled. 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail, 0,5 mM TCEP and 30% glycerol was added and samples were snap-frozen and stored at -70°C.

Purification of Pcf2

Cells pellets were resuspended into lysis buffer LB2 (50 mM Tris-HCl pH 8, 500 mM NaCl, 5% glycerol, 0.1% Triton X-100, 10 mM imidazole, 0.5 mM PMSF, cOmplete™ EDTA-free Protease Inhibitor Cocktail and 70 U/mL of benzonase) and sonicated at 4°C. Lysates were clarified by centrifugation at 5 °C at 18 500 rpm for 30 min and loaded to gravity flow Ni-NTA agarose resin (QIAGEN) previously equilibrated with wash buffer WB2_1 (50 mM Tris-HCl pH 8, 500 mM NaCl, 10 mM imidazole,). Resin was then washed with 10 column volumes of wash buffer WB2_1 followed by 10 column volumes of wash buffer WB2_2 (50 mM Tris-HCl pH 8, 1M NaCl, 10 mM imidazole). Pcf2-6His was eluted with 5 column volumes of the elution buffer EB2 (50 mM Tris-HCl pH 8, 500 mM NaCl, 250 mM imidazole, 0.5 mM TCEP, and 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail). The eluate was then concentrated to 300 µL (with Amicon® Ultra-15 30 kDa filter concentrators). 1 mM DTT, 1 mM MgCl and ≈2000 U of benzonase were added to the concentrated eluate and the sample was incubated 2 hours at 4 °C and injected into a Superdex 200 increase 10/300 (Cytiva) previously equilibrated with the final buffer FB2 (50 mM Tris-HCl pH 8, 500 mM NaCl and 1 mM DTT). Pcf2-6His containing fractions were pooled and directly used for CAF-1 reconstitution or stored at -70 °C with cOmplete™ EDTA-free Protease Inhibitor Cocktail, 0.5 mM TCEP and 30% glycerol.

Purification of Pcf3

Cells pellets were resuspended into lysis buffer LB3 (50 mM Tris-HCl pH 8, 200 mM NaCl, 5% glycerol, 0.1% Triton X-100, 10 mM imidazole, 0.5 mM PMSF, cOmplete™ EDTA-free Protease Inhibitor Cocktail and 70 U/mL of benzonase) and sonicated at 4°C. Lysate was clarified by centrifugation at 5 °C at 18500 rpm for 30 min and loaded to gravity flow Ni-NTA agarose resin (QIAGEN) previously equilibrated with wash buffer WB3_1 (50 mM Tris-HCl pH 8, 200 mM NaCl, 10 mM imidazole). Resin was then washed with 5 column volumes of wash buffer WB3_1 and 10 column volumes of wash buffer WB3_2 (50 mM Tris-HCl pH 8, 200 mM NaCl, 30 mM imidazole). his-Pcf3 was then eluted with EB3 (50 mM Tris-HCl pH 8, 200 mM NaCl, 250 mM imidazole, 1 mM DTT, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail). After addition of 1 mM MgCl2, 6His-TEV protease (with a ratio 1/10 in mass), ≈2000 U of benzonase the eluate was dialysed o/n at 5 °C in the final buffer FB3 (50 mM Tris-HCl pH 8, 200 mM NaCl and 1 mM DTT). Because of their similar size, Pcf3 and 6His-TEV protease cannot be completely separated by size-exclusion chromatography. Therefore, to remove the 6His-TEV protease and uncleaved His-Pcf3, 30 mM of imidazole was added to the dialysate, which was then loaded to gravity flow Ni-NTA agarose resin (QIAGEN) previously equilibrated with wash buffer WB3_2. The Flow through was concentrated to 300µL (with Amicon® Ultra-15 30 kDa filter concentrators) and injected into a Superdex 200 increase 10/300 (Cytiva) previously equilibrated with the final buffer FB3. Pcf3-containing fractions were pooled and stored at -70 °C after adding 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail, 0.5 mM TCEP and 30% glycerol.

Reconstitution of CAF-1 complexes

CAF-1 complexes were formed by mixing the isolated proteins Pcf1 (WT or mutant), Pcf2-6His and Pcf3 previously purified as described above. Isolated Pcf2-6His and Pcf3 were added in small excess compared to Pcf1. Tris 50 mM pH 8 was added to the Pcf1/Pcf2-his/Pcf3 mix to reach a final NaCl concentration of 150mM. After addition of 1 mM MgCl2, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail, 6His-TEV protease (with a ratio 1/10 in mass) and ≈2000 U of benzonase, the mixture was incubated over night at 4 °C and applied on a HiTrap® heparin FF column (Cytiva) previously equilibrated with EB4_1 (50 mM Tris-HCl pH 8, 100 mM NaCl). A gradient was applied with the high salt buffer EB4_2 (50 mM Tris-HCl pH 8, 1M NaCl). Fractions containing the full *Sp*CAF-1 were pooled, concentrated to 300µL (with Amicon® Ultra-15 30 kDa filter concentrators) and injected into a Superdex 200 increase 10/300 (Cytiva) previously equilibrated with the final buffer FB4_1 (50 mM Tris-HCl pH 8, 150 mM NaCl and 1 mM DTT). The *Sp*CAF-1-containing fractions were pooled and directly used for MST or EMSA analysis or stored at -70°C with 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail, 0,5 mM TCEP and 30% glycerol.

The *Sp*CAF-1(^{15}N-^{13}C-Pcf1) and *Sp*CAF-1(^{15}N-Pcf1) was reconstituted by co-lysing the pellets of ^{15}N-^{13}C-MBP-Pcf1 or ^{15}N-MBP-Pcf1 (WT or mutants), Pcf2-6His and his-Pcf3. Cell pellets from Pcf2-6His and Pcf3-his were added in excess compared to labeled MBP-Pcf1, based on the yield previously obtained for the isolated proteins. The pellets were resuspended and mixed in the lysis buffer LB4 (50 mM Tris-HCl pH 8, 150 mM NaCl, 5% glycerol, 0.1% Triton X-100, 10 mM imidazole, 0.5 mM PMSF, cOmplete™ EDTA-free Protease Inhibitor Cocktail and 70 U/mL of benzonase), sonicated and centrifuged as described before. The clarified lysate was applied to gravity flow Ni-NTA agarose resin (QIAGEN) previously equilibrated with wash buffer WB4_1 (50 mM Tris-HCl pH 8, 150 mM NaCl, 10 mM imidazole). Beads were washed with 5 column volume of WB4_1 buffer, followed by 10 column volumes of WB4_2 (50 mM Tris-HCl pH 8, 1 M NaCl, 10 mM imidazole). Elution was performed with EB4 (50 mM Tris-HCl pH 8, 150 mM NaCl, 250 mM imidazole, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail) and applied to an anion exchange column HiTrap® Q FF (Cytiva) previously equilibrated with buffer EB4_1. A gradient was applied with the high salt buffer EB4_2. The tagged CAF-1-containing fractions were pooled, and dialyzed overnight against buffer 4 DB4 (Tris 50 mM pH 8, 150 mM NaCl, 1 mM DTT) after addition of 1 mM DTT, 1 mM MgCl2, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail, 6His-TEV protease (with a ratio 1/10 in mass) and ≈2000 U of benzonase. The mixture was applied on a HiTrap® heparin FF column (Cytiva) using the same buffers (EB4_1 and EB4_2). The *Sp*CAF-1(^{15}N-^{13}C/^{15}N-Pcf1) -containing fractions were concentrated to 300μL (with Amicon® Ultra-15 30 kDa filter concentrators) and injected into a Superdex 200 increase 10/300 (Cytiva) previously equilibrated with buffer FB4_2 (10 mM Tris-HCl, 50 mM HEPES pH 7, 300 mM NaCl and 0.5 mM TCEP). The *Sp*CAF-1(^{15}N-^{13}C-Pcf1)-containing fractions were pooled and immediately used for NMR measurements.

Purification of histones *Sp*H3−*Sp*H4

Cells expressing *Sp*H3, *Sp*H4 with 6His-dAsf1 were pelleted by centrifugation and resuspended in the lysis buffer LB5 (50 mM Tris-HCl pH 8, 500 mM NaCl, 5% glycerol, 1% Triton X-100, 1 mM DTT, 10 mM MgCl2, 0.5 mM PMSF, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail) and flash frozen in liquid nitrogen. After thawing, lysozyme and benzonase were added at a final concentration of 0.25 mg/mL and 70 U/mL respectively. After incubation 20 minutes at 4°C, cells were lysed by sonication. Soluble 6His-Asf1 was removed on a NiNTA column (Qiagen) equilibrated in the LB5 buffer. The flow through (containing soluble free histones) was filtered with 0.22 μ filters and loaded on a cation exchange Resource S column (GE Healthcare) equilibrated with the dilution buffer EB5_1 (50 mM Tris-HCl pH8). Histones H3−H4 were eluted with a NaCl gradient in a buffer EB5_2 (50 mM Tris-HCl pH8, 2 M NaCl). The H3−H4-containing fractions were pooled, the salt concentration adjusted to 2 M NaCl, and concentrated in a 3 kDa concentrator (Millipore), flash freezed in liquid nitrogen and stored at -70°C.

Purification of Pcf1_ED and Pcf1_ED*

Cells expressing Pcf1_ED or Pcf1_ED* with a N-terminal 6His-SUMO tag were collected by

centrifugation, resuspended in lysis buffer LB6 (50 mM Tris-HCl pH8, 500 mM NaCl, 5% glycerol, 1%

Triton X-100, 1 mM PMSF, 1 μM aprotinin, 0.25 mM DTT) and flash frozen in liquid nitrogen. After

thawing, lysosyme was added at a final concentration of 1 mg/mL and cells were incubated 30 min at

4 °C and lysed by sonication. 6His-SUMO-Pcf1_ED was first purified on Histrap colums (Cytiva).

Fractions containing the protein were pulled. SUMO protease was added at a final concentration 1/10

and the mixture was dialyzed overnight at 4 °C against the buffer DB6 (50 mM Tris-HCl pH 8, 150 mM

NaCl, 10 mM imidazole) and applied on a NiNTA column (Qiagen) equilibrated in the DB6 buffer. The

flow-through fraction containing Pcf1_ED or Pcf1_ED*_was then purified by size exclusion

chromatography using a Superdex 75 increase 10/300 column (Cytiva) previsouly equilibrated with the

final buffer in FB6 (10 mM Tris-HCl, 50 mM HEPES pH 7, 300 mM NaCl). Fraction containing

Pcf1_ED or Pcf1_ED* were concentrated using Amicon centrifuge filter units of 3 kDa cutoff

(Millipore) flash freezed in liquid nitrogen and stored at -20 °C or -70 °C.

Purification of Pcf1_KER and Pcf1_KER-PIP

Cells expressing Pcf1_KER(56-170), or Pcf1-KER-PIP(56-185) (WT or mutant) with a N-

terminal 6His-MBP-TEV tag were collected by centrifugation, resuspended in lysis buffer LB7 (50 mM

Tris-HCl pH 8, 500 mM NaCl, 5% glycerol, 1% Triton X-100, 1 mM PMSF, 1 μM aprotinin, 0.25 mM

DTT, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail) and flash frozen in liquid nitrogen. After

thawing, 5 mM MgCl2, 1 mg/mL lysozyme and 70 U/mL of benzonase were added and cells were lysed

by sonication. Proteins were first purified on Histrap columns (Cytiva) including a wash step with WB7

(50 mM Tris-HCl pH 8, 1000 mM NaCl). 1 mM DTT and TEV protease (1/10 ratio) was added to the

fractions containing the 6His-MBP-TEV_ Pcf1_KER fragment and the mixture was incubated 2 hours

at room temperature and injected on a resource S column (Cytiva) previously equilibrated with EB7_1

(50 mM Tris-HCl pH 8). A gradient was applied with the high salt buffer EB7_2 (50 mM Tris-HCl pH

8, 2M NaCl). Fractions containing Pcf1_KER fragment were pooled and diluted to reach a concentration

of 150 mM NaCl and concentrated (with Amicon® Ultra-10 kDa filter concentrators).

Purification of Pcf1_WHD

Cells expressing Pcf1_WHD with a N-terminal 6His-SUMO tag were resuspended in lysis buffer

LB8 (50 mM Tris-HCl pH8, 500 mM NaCl, 5% glycerol, 1% Triton X-100, 1 mM PMSF, 1 μM

aprotinin, 0.25 mM DTT, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail) and flash frozen in

liquid nitrogen. After thawing, 5 mM MgCl2, 1 mg/mL lysozyme and 70 U/mL of benzonase was added

and cells were further lysed by sonication. The lysate was loaded onto gravity flow amylose resin (NEB)

previously equilibrated with buffer WB8_1 (50 mM Tris-HCl pH 8). Resin was then washed with 10

column volume of buffer WB8_1, 10 column volumes of buffer WB8_2 (50 mM Tris-HCl pH 8, 1000

mM NaCl), 10 column volumes of buffer WB8_1. 6His-SUMO- Pcf1_WHD was eluted with 10 column

volume of buffer EB8 (50 mM Tris-HCl pH 8, 500 mM NaCl, 250 mM Imidazole). SUMO protease was added at a final concentration 1/10 and the mixture was incubated overnight at 4°C. The mixture was concentrated (with Amicon® Ultra-3 kDa filter concentrators) and applied on a a Superdex 75 increase 10/300 size exclusion column (Cytiva) previously equilibrated with the final buffer FB8 (10 mM Tris-HCl, 50 mM HEPES pH7, 150 mM, NaCl). Finally, proteins were concentrated in a 3 kDa concentrator (Millipore).

Purification of *Sp*PCNA

Cells expressing *Sp*PCNA with a N-terminal 6His-SUMO tag were resuspended in lysis buffer LB9 (50 mM Tris-HCl pH8, 500 mM NaCl, 5% glycerol, 1% Triton X-100, 1 mM PMSF, 1 μM aprotinin, 0.25 mM DTT, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail) and flash frozen in liquid nitrogen. After thawing, 1 mM MgCl2, 1 mg/mL lysozyme and 70 U/mL of benzonase was added and cells were further lysed by sonication. The lysate was loaded onto gravity flow amylose resin (NEB) previously equilibrated with buffer WB9_1 (50 mM Tris-HCl pH 8). Resin was then washed with 10 column volume of buffer WB9_1, 10 column volumes of buffer WB8_2 (50 mM Tris-HCl pH 8, 2000 mM NaCl), 10 column volumes of buffer WB8_1. 6His-SUMO-*Sp*PCNA was eluted with 3 column volumes of buffer EB9 (50 mM Tris-HCl pH 8, 250 mM Imidazole). 1mM DTT and SUMO protease was added at a final concentration 1/10 and the mixture was dialyzed overnight at 4°C against the buffer DB9 (50 mM Tris-HCl pH 8, 150 mM NaCl, 10 mM imidazole). The mixture was applied on a Histrap column (Cytiva), the flow through (containing *Sp*PCNA) was concentrated (with Amicon® Ultra-15 3 kDa filter concentrators) and applied on a hiLoad 16/600 superdex 200 size exclusion column previously equilibrated with a FB9 (50 mM Tris-HCl pH8, 150 mM NaCl). In case the digestion of the tag was incomplete, the two last steps digestion with SUMO protease and gel filtration were repeated.

For all protein samples, depending on specific requirements of different techniques used, aliquots of concentrated protein were either maintained at 4°C or flash frozen in liquid nitrogen after addition or not of 30% glycerol and stored at -70°C for further use.

DNAs used to monitor protein-DNA interactions

The different DNAs were purchased from eurofins genomics. The sequences were derived from the 601 positioning sequence: ATCAATATCCACCTGCAGATACTACCAAAAGTGTATTTGG. For MST, the DNA were labeled with ALEXA488 at their 5' extremity. The ssDNA was annealed with the reverse-complementary sequence by heating at 90 °C and cooling slowly at room temperature.

Size-exclusion chromatography (SEC)

*Sp*CAF-1 subunits interaction was performed by mixing 2.2 nmoles of each isolated protein together in a final volume of 1.26 mL and left o/n at 5°C. The complexes were then concentrated to 300μL (with Amicon® Ultra-15 30 kDa filter concentrators) and injected into a Superdex 200 increase 10/300 (Cytiva) for separation by size-exclusion chromatography previously equilibrated with the

FB4_3 (50 mM Tris-HCl pH 7.5, 500 mM NaCl, 1 mM DTT). The different fractions were analyzed on mPAGE® 8% Bis-Tris Precast Gels (Sigma) with MOPS SDS running buffer. Interaction between CAF-1 and H3−H4 was carried out by incubating for 3 hours, 3 nmoles of *Sp*CAF-1 with 3 nmoles of *Sp*H3−H4 in a final buffer FB4_4 (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 4 mM DTT, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail) or FB4_5 (50 mM Tris-HCl pH 7.5, 1 M NaCl, 4 mM DTT, 1X cOmplete™ EDTA-free Protease Inhibitor Cocktail). Samples were then concentrated to 300µL (with Amicon® Ultra-15 30 kDa filter concentrators) and injected into a Superdex 200 increase 10/300 (Cytiva) for separation by size-exclusion chromatography with their corresponding buffers. The different fractions were analyzed on mPAGE® 4-20% Bis-Tris Precast Gels (Sigma) with MES SDS running buffer.

Electrophoretic Mobility Shift Assay (EMSA)

The proteins and DNA were mixed to be in a final EMSA buffer EMB (25 mM Tris-HCl pH 8, 1 mM EDTA pH 8.0, 150 mM NaCl) and incubated at 4 °C for 30 min and heated at 37 °C for 5 min prior to analysis on precast "any KD" Mini-PROTEAN TGX (Bio-rad, Cat #4569033) polyacrylamide gels using 1x TBE as running buffer. The gels were stained with 1x of SYBR Safe (Thermo Fisher Scientific, Waltham, MA) then visualized with BIORAD EZ Imager. A second identical the gel was Coomassie Blue before being visualized with the BIORAD EZ Imager. Band intensities were quantified by ImageJ.

MicroScale Thermophoresis (MST)

DNAs labeled with ALEXA488 at their 5' extremity. The final dilution buffer was FB_MST (10 mM Tris-HCl, 40 mM HEPES pH 7, 150 mM NaCl). The labeled DNA was adjusted to 20 nM. Freshly prepared proteins or complexes were diluted in the same buffer with 16 serial dilution 1:2. Each protein dilution was mixed with one volume of labeled DNA and filled into Monolith NT standard treated capillaries (NanoTemper Technologies GmbH). Thermophoresis was measured using a Monolith NT.115 instrument (NanoTemper Technologies GmbH) at an ambient temperature of 20 °C with 3-s/20-s/1-s laser off/on/off times, respectively. Instrument parameters were adjusted with 80% LED power and 40% MST power. Data of two measurements were analyzed (MO.Affinity Analysis software, NanoTemper Technologies) using the signal from thermophoresis at 5s.

Circular Dichroism (CD)

Circular Dichroism (CD) measurements were carried out at 20 °C on a JASCO J-810 spectro-polarimeter. Temperature was controlled by a Peltier. Spectra from 190 to 250 nm were obtained using a 2 mm optical path length quartz cell (Hellma #100-2-40) containing Pcf1_KER or Pcf1_KER* (5µM) in 10 mM of phosphate buffer (pH 7.4).

Nuclear Magnetic Resonance (NMR)

NMR experiments were carried out on Bruker DRX-600 MHz, 700 MHz or 950MHz spectrometers equipped with cryo-probes. All NMR data were processed using Topspin (Bruker) and analyzed using Sparky (T.D. Goddard and D.G. Kneller, UCSF). Samples were prepared in 3 mm NMR tubes, in solution containing 5% D2O, 0.1% NaN3, 0.1 mM DSS with different buffer appropriate for different complex formations or reactions. Heteronuclear Multiple Quantum Correlation (sofast-HMQC) or best-HSQC spectra were all recorded at 283°K. The protein concentrations were between 9 µM and 500 µM. For backbone resonances assignments, 3D data were collected at 283°K using standard Heteronuclear Single Quantum Correlation (HSQC) spectra ^{1}H-^{15}N HSQC, TOCSY-HSQC, HNCA, HBHA(CO)NH, CBCA(CO)NH, HN(CA)CO, HNCO, HN(CO)CA, CBCANH and HN(CA)CO experiments. Proton chemical shifts (in ppm) were referenced relative to internal DSS and ^{15}N and ^{13}C references were set indirectly relative to DSS using frequency ratios[38]. Chemical shift index were calculated according to the sequence-specific random coil chemical shifts[39,40].

Structural models of the *Sp*CAF-1 WHD domain were computed from NMR data with CS-ROSETTA[41] version 1.01. First, the MFR program from NMRpipe[42] was used to search a structural database for best matched fragments based on the protein backbone ^{15}N, ^{13}C, ^{13}CA, ^{13}CB and ^{1}HN chemical shifts. Then the ROSETTA 3.8 software was used to generate 27753 models by fragment assembly and full-atom relaxation. These models were rescored by comparing the experimental chemical shifts with the chemical shifts predicted by SPARTA[43] for each model. The best model after rescoring was chosen as a representative NMR model of the WHD domain.

Small Angle Xray Scattering (SAXS)

SAXS data were collected at the SWING beamline on a EigerX 4 M detector using the standard beamline setup in SEC mode[44]. Samples were injected into a Superdex 5/150 GL (Cytivia) column coupled to a high-performance liquid chromatography system, in front of the SAXS data collection capillary. The initial data processing steps including masking and azimuthal averaging were performed using the program FOXTROT[45] and completed using US-SOMO[46]. The final buffer subtracted and averaged SAXS profiles were analyzed using ATSAS v.3. software package[47]. To model the structures and improve the AlphaFold2 models, the program Dadimodo[48] (https://dadimodo.synchrotron-soleil.fr) that refines multidomain protein structures against experimental SAXS data was used (see **Table S1** in supplementary data for more information).

Structural Modeling

Sequences of *S. pombe* Pcf1 (Q1MTN9), Pcf2 (O13985), Pcf3 (Q9Y825), H3 (P09988), H4 (P09322) and PCNA (Q03392) were retrieved from UniProt database[49]. These sequences were used as input of mmseqs2 homology search program[50] used with three iterations to generate a multiple sequence alignment (MSA) against the uniref30_2103 database[51]. The resulting alignments were filtered using hhfilter[52] using parameters ('id'=100, 'qid'=25, 'cov'=50) and the taxonomy assigned to every sequence

keeping only one sequence per species. To increase the number of sequences in the alignment of *S. pombe* Pcf1 we independently generated MSA using mmseqs2 starting from the *S. cerevisiae* or the human homolog of Pcf1 (Q12495 and Q13111, respectively) and the resulting alignments were combined with the one of *Sp*Pcf1. Full-length sequences in the alignments were then retrieved and the sequences were realigned using MAFFT[53] with the default FFT-NS-2 protocol. To build the so-called mixed co-alignments, sequences in the alignment of individual partners were paired according to their assigned species and left unpaired in case no common species were found[51]. A first global model with full-length Pcf1, Pcf2 and Pcf3 was generated to map the regions of Pcf1 binding to Pcf2 and Pcf3 and to obtain the pLDDT scores shown in **Figure 1b** for Pcf1, **Figure S1f** for Pcf2 and Pcf3. Next, three models of the complex corresponding to independent modules of the complex were generated using different delimitations: model_1 (presented in **Figure S1g-i**) with Pcf1(403-450)-Pcf2(1-453) (MSA with 2180 species, 501 positions), model_2 (presented in **Figure S1j-l**) with Pcf1(200-335)-Pcf3(1-408) (MSA with 2148 species, 544 positions), model_3 Pcf1(352-383)-H3(60-136)-H4(25-103) (presented in **Figure 2f and S2d**) (MSA with 3530 species, 188 positions). Concatenated mixed MSAs were generated using the delimitations defined above and used as input to run 5 independent runs of the Alphafold2 algorithm with 6 iterations each[54]generating 5 structural models using a local version of the ColabFold interface[51] trained on the multimer dataset[55] on a local HPC equipped with NVIDIA Ampere A100 80Go GPU cards. The best models of each of the 5 runs converged toward similar conformations. They reached high confidence and quality scores with pLDDTs in the range [83.7, 84.3], [88.8, 89.8] and [86.5, 88.4] and the model confidence score (weighted combination of pTM- and ipTM-scores with a 20:80 ratio)[55] in the range [0.9, 0.93], [0.88, 0.89], [0.85, 0.87], for model_1, model_2 and model_3, respectively. The models with highest confidence score for each of the three models were relaxed using rosetta relax protocols to remove steric clashes[56] with constraints (std dev. of 2 Å for the interatomic distances) and were used for structural analysis. MSA web logos were generated with the weblogo server (https://weblogo.berkeley.edu/logo.cgi).

Nucleosome assembly assay

Mock- and p150CAF-1-depleted Xenopus high speed egg extract (HSE) were prepared as previoulsy[24]. Nucleosome assembly was performed on pBS plasmid damaged by UV (500J/m2) to promote DNA synthesis as previously described[24] except that the reaction mixed contained 3.2 µM of biotin-14-dCTP (Invitrogen, Ref 19518-0189) instead of [α^{32}P]-dCTP. The p150CAF-1-depleted extracts were complemented with 50 ng of isolated/reconstituted spCAF-1 complex composed of WT or mutated Pcf1. After DNA purification, samples were by processed for gel electrophoresis (1% agarose) to resolve topoisomers as previously described[24]. After staining with Ethidium bromide to visualize total DNA and gel transfer on a Nylon N+ membrane (GE Healthcare Ref RPN203B) (Qbiogen) for 45 min at 40 mbar in 10x SSC, the membrane was rinsed in PBS, air dried and DNA was crosslinked to the membrane using Stratalinker (Biorad). DNA synthesis was visualized by detecting

1 biotin with the Phototope-Star detection kit (New England Biolabs Ref N7020S) and images acquired

2 on a Chemidoc system (Biorad).

Standard yeast genetics

4 Yeast strains were freshly thawed from frozen stocks and grown at 30 ºC using standard yeast genetics

5 practices. The _pcf1_ mutants were obtained by classical genetic techniques. Yeast strains used in this

6 study are listed in Supplementary **Table S2**.

Peri-centromeric silencing assay

8 5-FOA (EUROMEDEX, 1555) resistant colonies were grown on uracil-containing liquid media

9 overnight and 10 µL of 5 fold serial dilutions (from 1.10^7 cells/mL to 1.10^5 cells/ml) were spotted on

10 indicated media.

Co-immunoprecipitation.

5.10^8 cells from exponentially growing cultures were harvested with 10% NaN_3 and 1 mM PMSF, final

concentration, and then washed twice in water and once in Lysis buffer (buffer (50 mM HEPES High

salt, 50 mM KoAc pH7.5, 5 mM EGTA, 1% triton X100, 0.01mg/mL AEBSF, EDTA-free protease

inhibitor cocktail). Cell pellets were resuspended in 800 µL of lysis buffer and were broken with a

Precellys homogenizer (twice 4 cycles at 10 000 rpm, 20 sec-2 min pause). After lysate clarification (30

minutes at 13 000 rpm, 4°C), 2.5 mg of proteins were incubated with pre-washed Dynabeads protein G

(Invitrogen, 10003D) coupled to anti-FLAG antibody (Sigma F7425) and incubated overnight at 4 °C

on a wheel. Beads were washed three times for 5 minutes at 4 °C with 800 µL of lysis buffer, and then

resuspended in 1X Laemmli buffer, and boiled at 95 °C for 10 minutes. INPUT and UNBOUND (both

10 % of initial protein extract) and BOUND (IP) fraction were resolved by electrophoresis on acrylamide

gels (4-12% Invitrogen) and the transferred onto nitrocellulose membrane that were saturated for 1 hour,

RT in TBS-0.075% tween-5% milk. Proteins of interest were detected with anti-FLAG antibody (Sigma

F1805, 1:1000) and anti-PCNA antibody (Santa Cruz sc-8349, 1:500).

Live cell imaging

All image acquisition was performed on the PICT-IBiSA Orsay Imaging facility of Institut Curie. For

snapshot microscopy, cells were grown in filtered supplemented EMM-glutamate, with or without

thiamine respectively, for 24 hours. Exponentially growing cultures were centrifuged and resuspended

in 50 µL of fresh medium. 2 µL from this concentrated solution was dropped onto a Thermo Scientific

slide (ER-201B-CE24) covered with a thin layer of 1.4 % agarose in filtered EMMg. 13 z-stack pictures

(each z step of 300 nm) were captured using a Spinning Disk Nikon inverted microscope equipped with

the Perfect Focus System, Yokogawa CSUX1 confocal unit, Photometrics Evolve512 EM-CCD camera,

100X/1.45-NA PlanApo oil immersion objective and a laser bench (Errol) with 491 (GFP) and 561

(MmCherry) nm diode lasers, 100 mX (Cobolt). Pictures were collected with METAMORPH software

and analyzed with ImageJ. For Pcf2-GFP and Rad52-GFP foci, a threshold (find maxima) was setup at the same level for each genetic background analyzed within the same experiment.

Data availability

We deposited structural models generated by AlphaFold2 at the modelarchive repository site (https://www.modelarchive.org/doi/10.5452/ma-1bb5w, https://www.modelarchive.org/doi/10.5452/ma-bxxkp, https://www.modelarchive.org/doi/10.5452/ma-htx0n)

Acknowledgements:

We thank the Alain LECOQ and Denis SERVENT from giving access to the CD spectro-polarimeter. This work was supported by grants from the INCA (2016-1-PL BIO-03-CEA-1, 2016-1-PLBIO-03-ICR-1), ANR (ANR-16-CE11-0028; ANR-20-CE18-0038; ANR-21-CE11-0027; ANR-21-CE44-0009-01) the program labeled by the ARC foundation 2016 (PGA1*20160203953), , the Fondation LIGUE "Equipe Labellisée 2020" (EL2020LNCC/Sal), and by french infrastructures, the Synchrotron Soleil (20191119; 20210745), the French Infrastructure for Integrated Structural Biology (FRISBI) ANR-10-INBS-0005 and the IR INFRANALYTICS FR2054. It benefited from the ERC-2015-ADG-694694 "ChromADICT", the Ligue Nationale contre le Cancer (Equipe labellisée Ligue) and ANR-11-LABX-0044 5. We also thank the PICT-IBiSA@Orsay Imaging Facility of the Institut Curie (particularly Laetitia Besse).

References

Luger, K., Mader, A. W., Richmond, R. K., Sargent, D. F. & Richmond, T. J. Crystal structure of the nucleosome core particle at 2.8 A resolution. *Nature* **389**, 251-260, doi:10.1038/38444 (1997).

Gurard-Levin, Z. A., Quivy, J. P. & Almouzni, G. Histone chaperones: assisting histone traffic and nucleosome dynamics. *Annu Rev Biochem* **83**, 487-517, doi:10.1146/annurev-biochem-060713-035536 (2014).

Ray-Gallet, D. & Almouzni, G. H3-H4 histone chaperones and cancer. *Curr Opin Genet Dev* **73**, 101900, doi:10.1016/j.gde.2022.101900 (2022).

Yi, S. J. & Kim, K. New Insights into the Role of Histone Changes in Aging. *Int J Mol Sci* **21**, doi:10.3390/ijms21218241 (2020).

Sultana, S., Zarreen, F. & Chakraborty, S. Insights into the roles of histone chaperones in nucleosome assembly and disassembly in virus infection. *Virus Res* **297**, 198395, doi:10.1016/j.virusres.2021.198395 (2021).

Smith, S. & Stillman, B. Purification and characterization of CAF-I, a human cell factor required for chromatin assembly during DNA replication in vitro. *Cell* **58**, 15-25 (1989).

Loyola, A. & Almouzni, G. Histone chaperones, a supporting role in the limelight. *Biochim Biophys Acta* **1677**, 3-11, doi:10.1016/j.bbaexp.2003.09.012 (2004).

Ridgway, P. & Almouzni, G. CAF-1 and the inheritance of chromatin states: at the crossroads of DNA replication and repair. *J Cell Sci* **113 (Pt 15)**, 2647-2658 (2000).

Martini, E., Roche, D. M., Marheineke, K., Verreault, A. & Almouzni, G. Recruitment of phosphorylated chromatin assembly factor 1 to chromatin after UV irradiation of human cells. *J Cell Biol* **143**, 563-575 (1998).

Moggs, J. G. *et al.* A CAF-1-PCNA-mediated chromatin assembly pathway triggered by sensing DNA damage. *Molecular and cellular biology* **20**, 1206-1218 (2000).

Rolef Ben-Shahar, T. *et al.* Two fundamentally distinct PCNA interaction peptides contribute to chromatin assembly factor 1 function. *Molecular and cellular biology* **29**, 6353-6365, doi:10.1128/MCB.01051-09 (2009).

Pietrobon, V. *et al.* The chromatin assembly factor 1 promotes Rad51-dependent template switches at replication forks by counteracting D-loop disassembly by the RecQ-type helicase Rqh1. *PLoS biology* **12**, e1001968, doi:10.1371/journal.pbio.1001968 (2014).

Gopinathan Nair, A. *et al.* Unorthodox PCNA Binding by Chromatin Assembly Factor 1. *Int J Mol Sci* **23**, doi:10.3390/ijms231911099 (2022).

Kaufman, P. D., Kobayashi, R. & Stillman, B. Ultraviolet radiation sensitivity and reduction of telomeric silencing in Saccharomyces cerevisiae cells lacking chromatin assembly factor-I. *Genes Dev* **11**, 345-357, doi:10.1101/gad.11.3.345 (1997).

Dohke, K. *et al.* Fission yeast chromatin assembly factor 1 assists in the replication-coupled maintenance of heterochromatin. *Genes Cells* **13**, 1027-1043, doi:10.1111/j.1365-2443.2008.01225.x (2008).

Sauer, P. V. *et al.* Insights into the molecular architecture and histone H3-H4 deposition mechanism of yeast Chromatin assembly factor 1. *eLife* **6**, doi:10.7554/eLife.23474 (2017).

Mattiroli, F. *et al.* DNA-mediated association of two histone-bound CAF-1 complexes drives tetrasome assembly in the wake of DNA replication. *eLife* **6**, doi:10.7554/eLife.22799 (2017).

Sauer, P. V. *et al.* Mechanistic insights into histone deposition and nucleosome assembly by the chromatin assembly factor-1. *Nucleic acids research*, doi:10.1093/nar/gky823 (2018).

Zhang, K. *et al.* A DNA binding winged helix domain in CAF-1 functions with PCNA to stabilize CAF-1 at replication forks. *Nucleic acids research* **44**, 5083-5094, doi:10.1093/nar/gkw106 (2016).

Richet, N. *et al.* Structural insight into how the human helicase subunit MCM2 may act as a histone chaperone together with ASF1 at the replication fork. *Nucleic acids research* **43**, 1905-1917, doi:10.1093/nar/gkv021 (2015).

Huang, H. *et al.* A unique binding mode enables MCM2 to chaperone histones H3-H4 at replication forks. *Nature structural & molecular biology* **22**, 618-626, doi:10.1038/nsmb.3055 (2015).

Ayoub, J., Buonanno, M., Di Fiore, A., De Simone, G. & Monti, S. M. Biochemical and Structural Insights into the Winged Helix Domain of P150, the Largest Subunit of the Chromatin Assembly Factor 1. *Int J Mol Sci* **23**, doi:10.3390/ijms23042160 (2022).

Liu, W. H. *et al.* The Cac1 subunit of histone chaperone CAF-1 organizes CAF-1-H3/H4 architecture and tetramerizes histones. *eLife* **5**, doi:10.7554/eLife.18023 (2016).

Ray-Gallet, D. & Almouzni, G. DNA synthesis-dependent and -independent chromatin assembly pathways in Xenopus egg extracts. *Methods Enzymol* **375**, 117-131, doi:10.1016/s0076-6879(03)75008-3 (2004).

Hardy, J. *et al.* Histone deposition promotes recombination-dependent replication at arrested forks. *PLoS genetics* **15**, e1008441, doi:10.1371/journal.pgen.1008441 (2019).

Ray-Gallet, D. *et al.* Dynamics of histone H3 deposition in vivo reveal a nucleosome gap-filling mechanism for H3.3 to maintain chromatin integrity. *Molecular cell* **44**, 928-941, doi:10.1016/j.molcel.2011.12.006 (2011).

Liu, W. H., Roemer, S. C., Port, A. M. & Churchill, M. E. CAF-1-induced oligomerization of histones H3/H4 and mutually exclusive interactions with Asf1 guide H3/H4 transitions among histone chaperones and DNA. *Nucleic acids research* **40**, 11229-11239, doi:10.1093/nar/gks906 (2012).

Mattiroli, F., Gu, Y., Balsbaugh, J. L., Ahn, N. G. & Luger, K. The Cac2 subunit is essential for productive histone binding and nucleosome assembly in CAF-1. *Sci Rep* **7**, 46274, doi:10.1038/srep46274 (2017).

Kaufman, P. D., Kobayashi, R., Kessler, N. & Stillman, B. The p150 and p60 subunits of chromatin assembly factor I: a molecular link between newly synthesized histones and DNA replication. *Cell* **81**, 1105-1114 (1995).

Rosas, R., Aguilar, R. R., Arslanovic, N., Tyler, J. K. & Churchill, M. E. A. A novel Single Alpha-Helix-DNA-binding domain in CAF-1 promotes gene silencing and DNA damage survival through tetrasome-length DNA selectivity and spacer function. *bioRxiv*, doi:10.1101/2022.10.11.511754 (2022).

Rouillon, C. *et al.* CAF-1 deposits newly synthesized histones during DNA replication using distinct mechanisms on the leading and lagging strands. *Nucleic acids research*, doi:10.1093/nar/gkad171 (2023).

Mejlvang, J. *et al.* New histone supply regulates replication fork speed and PCNA unloading. *J Cell Biol* **204**, 29-43, doi:10.1083/jcb.201305017 (2014).

Janke, R., King, G. A., Kupiec, M. & Rine, J. Pivotal roles of PCNA loading and unloading in heterochromatin function. *Proceedings of the National Academy of Sciences of the United States of America* **115**, E2030-E2039, doi:10.1073/pnas.1721573115 (2018).

Thakar, T. *et al.* Ubiquitinated-PCNA protects replication forks from DNA2-mediated degradation by regulating Okazaki fragment maturation and chromatin assembly. *Nat Commun* **11**, 2147, doi:10.1038/s41467-020-16096-w (2020).

Bieniossek, C., Imasaki, T., Takagi, Y. & Berger, I. MultiBac: expanding the research toolbox for multiprotein complexes. *Trends in biochemical sciences* **37**, 49-57, doi:10.1016/j.tibs.2011.10.005 (2012).

Miele, S. *et al.* The Xer activation factor of TLCPhi expands the possibilities for Xer recombination. *Nucleic acids research* **50**, 6368-6383, doi:10.1093/nar/gkac429 (2022).

Anderson, M. *et al.* Co-expression as a convenient method for the production and purification of core histones in bacteria. *Protein Expr Purif* **72**, 194-204, doi:10.1016/j.pep.2010.03.013 (2010).

Wishart, D. S. *et al.* 1H, 13C and 15N chemical shift referencing in biomolecular NMR. *Journal of biomolecular NMR* **6**, 135-140, doi:10.1007/bf00211777 (1995).

Tamiola, K., Acar, B. & Mulder, F. A. Sequence-specific random coil chemical shifts of intrinsically disordered proteins. *J Am Chem Soc* **132**, 18000-18003, doi:10.1021/ja105656t (2010).

Wishart, D. S., Bigam, C. G., Holm, A., Hodges, R. S. & Sykes, B. D. (1)H, (13)C and (15)N random coil NMR chemical shifts of the common amino acids. I. Investigations of nearest-neighbor effects. *Journal of biomolecular NMR* **5**, 332, doi:10.1007/BF00211764 (1995).

Shen, Y. *et al.* Consistent blind protein structure generation from NMR chemical shift data. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 4685-4690, doi:10.1073/pnas.0800256105 (2008).

Delaglio, F. *et al.* NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *Journal of biomolecular NMR* **6**, 277-293, doi:10.1007/BF00197809 (1995).

Shen, Y. & Bax, A. Protein backbone chemical shifts predicted from searching a database for torsion angle and sequence homology. *Journal of biomolecular NMR* **38**, 289-302, doi:10.1007/s10858-007-9166-6 (2007).

Thureau, A., Roblin, P. & Perez, J. BioSAXS on the SWING beamline at Synchrotron SOLEIL. *J Appl Crystallogr* **54**, 1698-1710, doi:10.1107/S1600576721008736 (2021).

Girardot, R., Viguier, G., Pérez, J. & Ounsy, M. FOXTROT: a Java-based application to reduce and analyse SAXS and WAXS piles of 2D data at Synchrotron SOLEIL. *canSAS-VIII* (2015).

Brookes, E., Vachette, P., Rocco, M. & Perez, J. US-SOMO HPLC-SAXS module: dealing with capillary fouling and extraction of pure component patterns from poorly resolved SEC-SAXS data. *J Appl Crystallogr* **49**, 1827-1841, doi:10.1107/S1600576716011201 (2016).

Manalastas-Cantos, K. *et al.* ATSAS 3.0: expanded functionality and new tools for small-angle scattering data analysis. *J Appl Crystallogr* **54**, 343-355, doi:10.1107/S1600576720013412 (2021).

Rudenko, O., Thureau, A. & Perez, J. Evolutionary refinement of the 3D structure of multi-domain protein complexes from Small Angle X-ray Scattering data. *Proceedings of the 2019*

Genetic and Evolutionary Computation Conference Companion (Geccco'19 Companion), 401-402, doi:10.1145/3319619.3322002 (2019).

UniProt, C. UniProt: the universal protein knowledgebase in 2021. *Nucleic acids research* **49**, D480-D489, doi:10.1093/nar/gkaa1100 (2021).

Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028, doi:10.1038/nbt.3988 (2017).

Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682, doi:10.1038/s41592-022-01488-1 (2022).

Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein annotation. *BMC Bioinformatics* **20**, 473, doi:10.1186/s12859-019-3019-7 (2019).

Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular biology and evolution* **30**, 772-780, doi:10.1093/molbev/mst010 (2013).

Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *BioRxiv*, doi:10.1101/2021.10.04.463034 (2022).

Leman, J. K. *et al.* Macromolecular modeling and design in Rosetta: recent methods and frameworks. *Nat Methods*, doi:10.1038/s41592-020-0848-2 (2020).