1   **Discovery of a new craniofacial gene influencing adaptive phenotypes in a non-model**

2   **pupfish radiation**

M. Fernanda Palominos[1,2], Vanessa Muhl[1,2], Emilie J. Richards[3], Christopher H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley

[2]Museum of Vertebrate Zoology, University of California, Berkeley

[3]Department of Ecology, Evolution, and Behavior, University of Minnesota

Keywords: gene expression, fluorescence in situ hybridization, galanin, trophic morphology,

craniofacial divergence, evo devo, speciation, adaptive radiation

Abstract

Understanding the origins of new species with novel adaptations is a fundamental question in biology that also provides an opportunity to uncover new genes and regulatory networks with potential clinical relevance. Here we demonstrate a new role for *galr2* in vertebrate craniofacial development using a non-model adaptive radiation of trophic specialist pupfishes endemic to San Salvador Island in the Bahamas. We confirmed the loss of a *Sry* transcription factor binding site in scale-eating pupfish and found significant spatial differences in *galr2* expression among pupfish species in the Meckel's cartilage and premaxilla using in situ hybridization. We then experimentally confirmed a novel function for Galr2 in craniofacial development and jaw elongation by exposing embryos to Galr2. Galr2-inhibition reduced Meckel's cartilage length and increased chondrocyte density in both trophic specialists but not in the generalist genetic background. We propose a mechanism for jaw elongation in scale-eaters based on the reduced expression of *galr2* putatively due to the loss of the *Sry* binding site that results in reduced opportunities for binding with their endogenous ligand during craniofacial chondrogenesis. Fewer Galr2 receptors in the scale-eater Meckel's cartilage may result in their enlarged jaw lengths as adults by limiting opportunities for Galr2 antagonists to bind to these receptors during development. Our findings illustrate the growing utility of linking candidate adaptive SNPs in non-model vertebrate systems with highly divergent phenotypes to novel gene functions with potential implications for human biology.

Introduction

Craniofacial developmental anomalies are the most common source of birth defects in humans, present in 1 out of 700 births (Hall et al., 2010; Meng et al., 2009; Palmer et al., 2016). While Mendelian craniofacial defects are well characterized (e.g. Treacher Collins Syndrome (Kadakia et al., 2014), Apert Syndrome (Wilkie et al., 1995), and Crouzon Syndrome (Helman et al., 2014; Reardon et al., 1994)), the developmental genetics of complex craniofacial defects, such as non-syndromic orofacial clefts and craniosynostosis, are poorly understood (Carlborg and Haley, 2004; Glazier, 2002; Hirschhorn and Daly, 2005; Hochheiser et al., 2011; Manolio et al., 2009; Monteiro and Podlaha, 2009; Weinberg et al., 2018). With the continued lowering costs of genomic sequencing and functional genetic tools, it is increasingly feasible to develop new genetic models for understanding human development and disease.

Current knowledge of craniofacial developmental genetics results mainly from mutagenic screens and forward genetic approaches in a few vertebrate model systems (primarily mice and zebrafish); however, these approaches are typically biased toward large Mendelian effects on easily detectable phenotypes, often missing the variation most relevant to common craniofacial birth defects in humans, such as small-effect variants, subtle differences in skeletal shape, and epistatic interactions among genes and the environment (Albertson et al., 2009; Powder and Albertson, 2016; Roberts et al., 2011). Conversely, complementary genome-wide association studies (GWAS) of craniofacial syndromes and normal-range facial variation in humans often reveal small-effect loci with no tractable possibilities for functional interrogation (Bonfante et al., 2021; Naqvi et al., 2022; Weinberg et al., 2018). Thus, understanding the genetic bases of naturally occurring, highly divergent adaptive phenotypes in non-model systems that parallel human clinical variation, known as 'evolutionary mutant' models (Albertson et al., 2009; Concannon and

Albertson, 2015; Powder and Albertson, 2016; Rohner, 2018), provides a powerful approach

combining the tractable functional investigations possible in model systems with genome-wide

association scans of small-effect regulatory loci underlying natural craniofacial diversity. For

example, novel insights into a broad diversity of human diseases have now arisen from studies of

diverse non-model species experiencing strong selective pressures in nature, including senescence

(i.e., naked mole rat (Buffenstein, 2008); turquoise killifish (Platzer and Englert, 2016), and

rockfish (Kolora et al., 2021)); eye degeneration, bone loss and enhanced lipogenesis (blind

cavefishes (Gross et al., 2016; Hofmann et al., 2009; Protas et al., 2007; Lam et al. 2022)); body

size (domesticated dogs (Sutter et al., 2007)); cholesterol (polar bears (Liu et al., 2014)); and

diabetes (dolphins (Venn-watson et al., 2011)). In particular, the most remarkable diversity of

vertebrate craniofacial morphology is represented in teleost fishes, often associated with their

diverse and sometimes highly specialized modes of feeding (e.g., Liem, 1991; Otten, 1983;

Albertson et al., 2003; Martin and Richards 2019; Konings et al., 2021; Evans et al. 2021).

Emerging fish model systems include the rapidly evolving East African and Cameroon

cichlid radiations, in which a small number of genetic changes underlie immense morphological

disparity in over 1,500 cichlid species (Brawand et al., 2014; Conith et al., 2019; Kocher et al.,

2003; Martin, 2012; Martin, 2013; Martin and Genner, 2009; Navon et al., 2020) and the repeated

parallel speciation of stickleback ecomorphs in glacial lakes (Chan et al., 2010; Erickson et al.,

2016; Jones et al., 2012; Miller et al., 2007; Ishikawa et al. 2019). These systems provide excellent

examples of leveraging naturally-occurring and highly divergent craniofacial phenotypes as

'evolutionary mutants' or 'evolutionary forward genetics' models to gain novel insights into the

genetics of natural human craniofacial variation (Erickson et al., 2014; Glazer et al., 2015; J.

Parsons et al., 2021; Kocher, 2004; Powder and Albertson, 2016; Roberts et al., 2009).

Figure 1. Divergent craniofacial morphology and development of the San Salvador Island *Cyprinodon* **pupfish radiation.** *C. variegatus* (first row) is a trophic generalist distributed across the western Atlantic and Caribbean; *C. brontotheroides* (second row) is a molluscivore, and *C. desquamator* (third row) is a scale-eater, both endemic to the hypersaline lakes of San Salvador Island (SSI), Bahamas. Left panel: development at 1-, 3-, and 8-days post-fertilization (dpf). Middle panel: lab-reared adult female pupfishes of each species. Right panel: Lateral and dorsal views of µCT scans of craniofacial morphology (modified from Hernández et al., 2017). The maxilla is colored in blue, premaxilla in red, dentary in green, articular in orange.

Here we demonstrate the utility of a non-model evolutionary radiation of *Cyprinodon* pupfishes for discovering and validating a new craniofacial gene of potential clinical relevance to normal human facial variation and craniofacial disorders. Pupfishes offer unique advantages over other evolutionary model systems because they 1) rapidly evolved highly divergent and unique craniofacial phenotypes not found in other non-model fish systems (Fig. 1) with minimal genetic differentiation among species (Hernandez et al., 2018; Lencer et al., 2016; Lencer et al., 2017; Martin and Wainwright, 2011; Martin and Wainwright, 2013a; Martin et al., 2019; St John et al., 2020a; St John et al., 2020b), 2) speciated in the face of ongoing gene flow after colonizing hypersaline lakes, resulting in very few highly differentiated genomic regions associated with

species-characteristic craniofacial traits (Martin and Feinstein 2014; Martin et al., 2017; McGirr and Martin, 2016; McGirr and Martin, 2021; Richards and Martin, 2017; Richards and Martin, 2022; Richards et al., 2021), and 3) are highly amenable to laboratory rearing and imaging due to their high fecundity, daily egg production, and egg transparency comparable to zebrafish (Martin and Gould, 2020; Martin and Wainwright, 2013b). This radiation contains the widespread algae-eating generalist pupfish, *Cyprinodon variegatus* (Fig. 1a), which is broadly distributed across the Caribbean and North American Atlantic coast and occurs in sympatry with two microendemic trophic specialist species found only in the hypersaline lakes of San Salvador Island (SSI), Bahamas. Each trophic specialist displays highly divergent behavior, pigmentation, and craniofacial morphology: the molluscivore *C. brontotheroides*, has a novel nasal protrusion that is a skeletal extension of the maxilla with bilateral nasal tissue clefting and foreshortened robust oral jaw (Fig. 1b); and the scale-eater, *C. desquamator*, exhibits two-fold larger oral jaws and overall brachycephalic features (Fig. 1c) (Martin, 2016; Martin and Wainwright, 2011; Martin and Wainwright, 2013a). There is also a fourth intermediate scale-eating ecotype in some lakes (Richards et al. 2022).

Previous genomic and transcriptomic work on the SSI pupfishes identified dozens of new candidate craniofacial genes never previously characterized or investigated in other systems (McGirr and Martin 2016, 2019, 2020; Richards et al. 2021, 2022; Richards and Martin 2017; Patton et al. 2021; St. John et al. 2021). One of the most promising candidates was galanin receptor 2a, the second receptor type for galanin (see discussion of genomic evidence in Methods and Materials). In humans, *galr2* is abundantly expressed within the hypothalamus and hippocampus of the central nervous system and in the heart, kidney, liver, colon, and small intestine, and has genetic associations with epilepsy and Alzheimer's (Li et al., 2013; Ma et al., 2018). Classified as

an orexigenic (appetite stimulant) gene, *galr2* is expressed in the human hypothalamus but also in

the ventral telencephalon of larval and adult zebrafish (Ahi et al., 2019; Li et al., 2013; Kim et al.,

2016). Although a role of *galr2* in craniofacial development has not previously been reported in

the literature, its ligand, the neuropeptide galanin (GAL), is highly expressed in bones from early

to post-embryonic development (Xu et al., 1996, Jones et al., 2009), with demonstrated effects on

bone mass (Idelevich et al., 2018; Idelevich et al., 2019), muscle contraction (Kakuyama et al.,

1997. EJP), and periodontal regeneration (Ma et al. 2021).

Here we confirmed the presence of two highly differentiated SNPs between SSI specialist

species affecting transcription factor binding sites in the *galr2* regulatory region, characterized the

divergent craniofacial expression of *galr2* across all three SSI pupfish species from two different

lake populations using *in situ* hybridization chain reaction (HCR) (Choi et al., 2018; Ibarra-Padilla-

García et al., 2020) at two key developmental timepoints, and demonstrated that receptor inhibition

of exclusively the Galr2 pathway reduces Meckel's cartilage length and increases chondrocyte

density dependent on the species' genetic backgrounds.

Results

Two highly differentiated SNPs are associated with *galr2a* transcription factor binding sites

Previous whole genome sequencing of over one hundred San Salvador Island (SSI) pupfishes

identified only two single nucleotide polymorphisms (SNP) nearly fixed between trophic

specialists across lake populations within 20 kb of *galr2a* (Richards et al. 2021). We designed

primers and used Sanger sequencing to verify these SNPs in a large panel of wild-caught specialists

from five lake populations on SSI (Table S5). We confirmed the presence of a transversion from

G to A approximately 11 kb upstream of the *galr2a* transcription starting site (TSS; Fig. 2a., S1;

Table S1), which changes the predicted CA**G**CAA *Elf1/Erg* transcription factor binding site

(TFBS) to an **A**GGGASW *Elf5* TFBS at this locus (using the Multiple Expectation maximizations

for Motif Elicitation (MEME) server and the motif database scanning algorithm TOMTOM

(Bailey et al., 2009)). This transversion was observed in 86.3% of scale-eaters across four lake

populations ($n = 51$) versus 20% of molluscivores across six lake populations on SSI ($n = 30$) (Fig.

2a., S1).

We confirmed the presence of a second transversion from C to T approximately 8 kb

upstream of *galr2a* TSS, which changes the predicted AGA**C**AA *Sry* TFBS to a YAGA**T**A *Znf146*

TFBS (Fig. 2b., S2; Table S1). This transversion was observed in 90.9% of scale-eaters across four

lake populations ($n = 55$) versus 33.9% of molluscivores across six lake populations on SSI ($n =$

53) (Figure S2). Notably, all scale-eaters sampled from Crescent Pond (CRP) contained the T

transversion ($n = 30$) (Supplementary Figure S3). The predicted TFBS changes in the *galr2a* cis-

regulatory region across pupfishes, combining with a previous genetic mapping study that found

a significant QTL in this region explaining 15% of phenotypic variance in oral jaw size (Martin et

al. 2017), suggests that different spatial or temporal *galr2* expression may underlie some of the

craniofacial divergence in SSI pupfishes.

Figure 2. The two most differentiated single-nucleotide polymorphisms (SNP) between trophic specialists within 20kb upstream of *galr2*. a. Putative ancestral TFBS for *Elf1/Erg* and *Sry*, respectively, in the generalist (G), *Cyprinodon variegatus* and the molluscivore (M) *Cyprinodon brontotheroides*. **b.** Derived changes in TFBS to *Elf5* and *Znf146* in the scale-eater (S) *Cyprinodon desquamator*. Pie charts indicate frequencies of ancestral (teal) and derived (orange) alleles at each locus. +1 represents the transcription starting site of *Galr2a* (Galanin receptor 2a). μCT scans of craniofacial morphology are modified from Hernández et al., 2017. Sequence logos of the Position Weight Matrixes (PWM) identified using MEME and TOMTOM.

Different spatiotemporal patterns of *galr2a* expression in craniofacial tissues

To determine if spatial or temporal changes in *galr2a* expression underlie SSI craniofacial divergence, we quantified *galr2a* expression in 2 dpf embryos and 8 dpf hatching larvae across all SSI species in two independent lake populations using hybridization chain reaction (HCR) in-situ hybridization for *galr2a* and *tropomyosin 3b* (*tpm3b*), a component of thin filaments of myofibrils expressed in fish skeletal muscles (Dube et al., 2016), as a positive control.

At 2 dpf, *galr2a* was expressed in more regions of the central nervous system of specialists (Fig. 3a,c) than in generalists (e.g. the posterior tectum and medial longitudinal fasciculus); however, we observed increased *galr2a* expression in the first pharyngeal arches and the surrounding mandibular mesenchyme in the molluscivore and generalists pupfishes than in the scale-eaters. In addition to an absence of *galr2a* expression in the mesenchyme surrounding the first pharyngeal arch, its expression was conserved in the first and second pharyngeal arches and clefts (Fig. 3c,e).

Using 3D reconstruction and volume rendering analysis of HCR data for whole-mounted pupfishes at hatching time (8 dpf), we found that *galr2a* expression was higher in the jaw of the molluscivores than in the generalists (Fig. 4; $P = 0.03$, Tukey's HSD). In contrast, *galr2a* showed no differences in expression among species in the brain and head (Fig. 4, Table S3). G*alr2a* was expressed in the Meckel's and palatoquadrate cartilages in all SSI pupfishes, in the premaxilla of the generalist and the scale-eater SSI specialist (Fig. 4 b-b', c-c'), the maxilla of the molluscivore SSI specialist (Fig. 4 a-a'), and in the intermandibular muscles of the scale-eaters (Fig. 4 c').

At the subcellular level, *galr2a* was expressed in the cytoplasm of chondrocytes at the Meckel's symphysis (Fig. 5a). In contrast, on the distal edge of the Meckel's cartilage it was expressed only in the cytoplasm of the elongated chondrocytes (Fig. 5a'). Towards the most

posterior region of the Meckel's closest to the palatoquadrate cartilage, we observed strong

membrane expression of *galr2a* in the cells surrounding the jaw joint (Fig. 5 b, c, d). We conclude

that g*alr2a* was significantly differentially expressed in specific and distinct craniofacial tissues in

the specialists at hatching time, suggesting an important role for craniofacial divergence in the SSI

radiation.

We further corroborated the differential spatial expression of *galr2a* at 2 dpf and 8 dpf by

quantifying transcript counts from the only existing RNAseq data set on SSI heads during

development (Lencer and McCune, 2020). Only *galr2a*, but not *galr1a, galr1b, galr2a, galr2b,* or

galanin, showed significantly higher mRNA expression in the molluscivores relative to generalists

(P = 0.003, Tukey's HSD test) and scale-eaters (P = 0.014, Tukey's HSD test) at 2 dpf (Table S2),

whereas *galr2a* showed overall similar levels of expression in the head from 4 to 15 dpf in all three

species (Table S2).

As the rapid evolution of jaw size in the SSI radiation is accompanied by rapid divergence

of the adductor muscle mass (Martin and Wainwright, 2011) and hypertrophy of the adductor

mandibulae complex (Hernandez et al., 2018), we selected a known skeletal muscle gene, *tpm3b*

(tropomyosin alpha-3) as a positive control for our HCR and to test for possible muscle differences

between species during jaw development. At 2 dpf that *tpm3b* was expressed in adaxial cells, the

first muscle precursor cells, in all SSI pupfishes (Fig. 3a''-c''). However, only in the scale-eaters,

Tpm3b was expressed in the eye's inferior oblique muscle primordial cells (Fig. 3c''). At hatching

time, *Tpm3b* labeled all larval head muscles (Fig. 4a-a', b-b', c-c'; 5b). The scale-eater and

molluscivore *Tpm3b* expression volume was significantly higher than in the generalists (Table S3;

P < 0.05; 1-way ANOVA, Tukey's HSD test).

Figure 3. Spatiotemporal patterns of *galr2a* expression differ among SSI species at 2 dpf.

Representative images of *Galr2a* (green) and *Tpm3b* (magenta) expression revealed by hybridization chain

reaction (HCR) and DAPI nuclear staining (gray) in the: **a.** molluscivore, *C. brontotheroides*; **b.** generalist,

C. variegatus; and **c.** scale-eater, *C. desquamator*. **d.** Galr2a reads per kilobase of transcript per million

reads (RPKM) from an existing mRNAseq study sampling the entire head of each species at four

developmental stages (Lencer and McCune, 2020). Galr2a showed significantly higher expression in the

molluscivore relative to generalists (P = 0.003, Tukey's HSD test) and scale-eaters (P = 0.014, Tukey's

HSD test) at 2 post fertilization (dpf). **e.** Mean fluorescence per area for galr2a quantified at 2 dpf in all three

species shows similar elevated levels of expression in the molluscivore for the mandibular and pharyngeal

arches. Images shown are representative Z-stack maximum projection images of 30 optical sections taken

every 15 μm from whole-mounted larvae imaged using a Zeiss LSM880 laser confocal microscope. PA:

pharyngeal arches, PC: pharyngeal cleft, ManM: mandibular mesenchyme, MLF: medial longitudinal

fasciculus, PTe: posterior tectum, Di: diencephalon; Mes: mesencephalon, Mye: myelencephalon, Y: yolk,

AdC: Adaxial cells, IO: inferior oblique muscle.

Figure 4. Spatiotemporal patterns of *galr2a* expression differ in molluscivores at 8 dpf. Representative images of *Galr2a* (green) and *Tpm3b* (magenta) expression revealed by hybridization chain reaction (HCR) and DAPI nuclear staining (gray). **a.** Lateral view of an 8 dpf molluscivore (M in white circle), **a'.** ventral view. **b.** Lateral view of an 8 dpf generalist (G in white circle), **b'.** ventral view. **c.** Lateral view of an 8 dpf scale-eater (S in white circle), **c'.** ventral view. The arrowhead in **b.** and the arrow in **c.** point to high expression of *galr2a* in the maxilla and premaxilla of the molluscivore and scale-eater specialists, respectively. White arrowheads in **a'.** point to high expression of *galr2a* in the palatoquadrate of the

molluscivore. Images shown here are representative Z-stack maximum projection images of 30 optical

sections taken every 15 µm from whole-mounted larvae imaged using a Zeiss LSM880 laser confocal

microscope. **d-d'.** 3D-Slicer view of the expression volume of *galr2a* (**d.**) and *tpm3b* (**d'.**) in a hatched

molluscivore pupfish larvae (*C. brontotheroides*). **e.** Volume of *galr2a* expression in the brain (left panel),

whole head (middle panel), and jaw (right panel) across SSI pupfishes. CNS: central nervous system, Max:

maxilla. LEVs: levator arcus palatini and operculi, AdM: adductor mandibular. IMA: intermandibular anterior.

Total volume rendered: 450 µm. All scale bars: 100 µm.

Figure 5. *Galr2a* is expressed in the chondrocytes of the Meckel's and palatoquadrate cartilages at

hatching (8 dpf) in SSI pupfishes. HCRs showing tissue expression of *Galr2a* in green, and the muscle

marker Tropomyosin 3b, (*Tpm3b*), in magenta. Nuclei are stained with DAPI (gray). **a-a'.** Scale-eater *galr2a*

expression in the chondrocytes of the Meckel's symphysis (Sym) and lateral side. **b.** Ventral view of a

molluscivore jaw at 8 dpf showing the expression of *Tpm3b* in the intermandibular muscles of the lower

jaw. *Galr2a* is expressed in the jaw joint and the most anterior region of the lower jaw. **c-d.** *Galr2a* is

expressed in the chondrocytes of the Meckel's and palatoquadrate cartilages and around the jaw joint. **e.**

Galr2a expression in the premaxilla of a scale-eater. Unlabeled scale-bars: 10 μm.

Chemical inhibition of Galr2 receptors affects Meckel's cartilage length and chondrocyte

density

To study the effect of Galr2 in craniofacial development, we inhibited the endogenous activation

of all four known galanin receptors in teleost fishes (*Galr1a* and b, *Galr2a* and b; Butler et al.

2020, Li et al. 2013) using M35 (Innopep, Inc.), a synthetic peptide antagonist of Galr1+2 galanin

receptors (Wiesenfeld-Hallin et al., 1993), and M871 (Abcam), a Galr2-specific synthetic peptide

antagonist (Sollenberg et al., 2006). Embryos of all three species from two different lake

populations were exposed from stages 24-25 (2 dpf, with the appearance of the first pharyngeal

arches) until hatching at stages 32-33 (8 dpf, Fig. 6a).

At approximate hatching time (8 dpf), both trophic specialist species raised under

laboratory common garden conditions exhibited increased Meckel's cartilage length (Fig. 6d),

consistent with the longer jaws of adult scale-eaters and more robust jaws of the molluscivore

relative to more gracile jaws of the generalist (Martin and Wainwright 2013). We found that

exposure to M871, the Galr2-specific antagonist, significantly reduced the length of the Meckel's

cartilage in both specialists relative to the generalists (Fig. 6d.; $P = 7.89 \times 10^{-5}$ for scale-eaters; P

= 0.001 for molluscivores; 2-way ANOVA, Tukey's HSD test). In contrast, M35 (Galr1 and Galr2

antagonist) only significantly reduced Meckel's length in the molluscivore (Fig. 6c). Meckel's

length of generalists was unaffected by exposure to M35 or M871 (Fig. 6d).

 Figure 6. Galr2 receptor inhibition reduced Meckel's cartilage length in both specialists and

 increased chondrocyte density in scale-eaters. a. Galr2 and Galr1+2 inhibition protocol during pupfish

 development. *C. desquamator* embryos are depicted. **b.** Alcian blue cartilage staining at Stage 32-33. IOD:

 interocular distance, MC: Meckel's cartilage. The MC length and IOD are shown in red. **c.** Chondrocytes at

 the symphysis. **d.** Changes in the log-transformed Meckel's length and the log-transformed interocular

 distance at Stage 32-33 across species and treatments. **e.** Chondrocyte numbers at the symphysis across

 species and treatments. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ Tukey's HSD test.

To understand the cellular effect of Galr1 and 2 antagonists on Meckel's length across

species, we quantified the number of chondrocytes 100 μm from the symphysis and the mean

width of ten chondrocytes nearest to the symphysis. We found no significant differences in the

mean chondrocyte width among pupfish species but observed increased chondrocyte density in

untreated scale-eaters relative to generalists (Fig. 6e; Table S4; $P = 0.009$, ANOVA, Tukey's HSD

test). Moreover, only scale-eaters responded to M871, but not M35, by further increasing

chondrocyte density relative to the control larvae (Fig. 6e; Table S4; $P = 0.03$, ANOVA, Tukey's

HSD test; mean ± 1 SE; control = 28.22 ± 0.38; M871 = 32.06 ± 1.06). Despite having shorter

jaws after treatment with M871, chondrocyte density was significantly increased (Fig. 6e; mean ±

1 SE; control = 28.22 ± 0.38; M871 = 32.06 ± 1.06).

Discussion

We used a non-model evolutionary radiation of trophic specialist pupfishes, endemic to the

Bahamas, to discover a novel function for *galr2a* in craniofacial divergence. Specifically, we

confirmed that only two transcription factor binding sites upstream of *galr2a* display highly

divergent allele frequencies between trophic specialist species, visualized *galr2a* expression in

craniofacial tissues in all three SSI pupfish species at two developmental stages, and demonstrated

a phenotypic effect on Meckel's cartilage length and chondrocyte density using synthetic peptides

to inhibit the activity of Galr2 and Galr1+2. Our findings demonstrate a crucial role for Galr2 in

craniofacial divergence within this pupfish radiation. Our study also provides a roadmap in a non-

model vertebrate system for rapidly identifying previously uncharacterized candidate genes

important for adaptation to novel ecological niches (e.g. trophic specialization) that can be quickly

validated through classic and state-of-the-art developmental biology tools. Overall, our research

contributes to understanding the genetic basis of phenotypic evolution and adaptation in non-

model organisms.

Putative loss of a transcription factor binding site for *galr2a* in scale-eating pupfish

One of the most common evolutionary changes associated with phenotypic changes among closely

related species is the gain or loss of cis-regulatory elements (Mack and Nachman, 2017; Signor

and Nuzhdin, 2018). More than 85% of scale-eaters carry two transversions in the regulatory

region of *galr2a* (Fig. 2). Combined with our observations of reduced *galr2a* expression in the

mandibular mesenchyme during early jaw development in this species (Fig. 3), we conclude that

a loss of a Sry transcription factor binding site in scale-eaters is the most likely explanation for

changes in gene expression, rather than a gain of a new TFBS for Znf416 at this locus (Fig. 2).

This is further supported by the critical role of the Sry-related HMG box (Sox) family of

transcription factors (especially the SoxE group including Sox8, Sox9, and Sox10) in craniofacial

development as Sry transcription factors specify the behavior, multipotency, and survival of neural

crest cells during vertebrate development (Leathers and Rogers, 2022; Simões-Costa and Bronner,

2015; Guo et al., 2015; Haldin and LaBonne, 2010; Betancur et al., 2010; Kamachi and Kondoh

2013).

Alternatively, we cannot rule out a gain of a TFBS upstream of *galr2a* in scale-eaters,

additive or epistatic effects of both upstream inversions, or more complex regulatory architectures,

such as many interacting functional cis- and trans-acting regulatory variants or combinations of

variants segregating at lower frequencies in trophic specialists that we have not prioritized (Boyle

et al., 2017; McGirr and Martin, 2021; Price et al., 2015). However, our mapping cross of a single

outbred pair of trophic specialists indicates that a single moderate-effect QTL containing *galr2a*

explains 15% of phenotypic variation in oral jaw length between these species, consistent with

causative variants affecting jaw size originating from this region (Martin et al. 2017).

Differential *Galr2a* expression during development is associated with craniofacial divergence

in SSI pupfishes

We found *galr2a* expression in our in situ hybridization experiments to be consistent with

previously published RNAseq data for craniofacial tissue in this radiation (Fig. 3; Lencer and

McCune, 2017). We observed distinctive spatial *galr2a* expression among SSI species at 2 and 8

dpf suggesting important time and tissue-specific regulation of *galr2a* expression during pupfish

development. At 2 dpf, we found a strong association between decreased expression of *galr2a* in

the mandibular mesenchyme anterior to the first pharyngeal arch with the future Meckel's cartilage

and oral jaw lengths in the adults of each species; with increasing *galr2* abundance in the

molluscivores associated with shorter, but more robust jaws in adults (Martin and Wainwright

2013). In contrast, the reduced *galr2a* expression in the mandibular mesenchyme is associated

with the development of longer oral jaws in scale-eaters, which is apparent as early as hatching

time (Fig. 6d; Lencer et al., 2016).

We noted expression of *galr2a* in the maxilla of only molluscivores at hatching (Fig. 4 a-

c), absent in the scale-eaters and generalists, consistent with the uniquely enlarged and anteriorly

protruding head of the maxilla in this species (Martin and Wainwright 2013; Hernandez et al. 2018;

Martin and Feinstein 2014; St. John et al. 2020). Furthermore, *galr2a* expression in the

intermandibular muscles of the scale-eaters at hatching time is consistent with the observed

hypertrophic musculature of the adductor mandibulae in the adult scale-eating pupfish (Hernandez

et al., 2018). Altogether, these interspecific differences in spatial expression support a novel role

for *galr2a* in musculoskeletal development accounting for the divergent craniofacial morphology

observed in SSI pupfishes.

Receptor inhibition reveals a novel function for GALR2 in craniofacial divergence of SSI

pupfishes

Interestingly, the response of Meckel's cartilage length and chondrocyte density to the inhibition

of Galr-receptor pathways was highly dependent on the species' genetic background. Scale-eaters

responded only to the Galr2-specific antagonist M871 with substantially reduced Meckel's

cartilage length and increased chondrocyte density, but were unaffected by the Galr1+2 antagonist

M35 (Fig. 6c). Importantly, M871 exhibits a higher binding constant for Galr2 receptors than the

Galr1+2 antagonist M35 (Sollenberg et al., 2006; Sollenberg et al., 2010). Previous transcriptomic

data from craniofacial tissues during early development (Lencer and McCune 2017) also indicates

that *galanin* and *Galr1* mRNA transcript abundance does not vary among the three SSI species

(Table S3).

Therefore, we conclude that even though scale-eaters expressed the lowest amount of *galr2a*

transcripts and presumably contain overall fewer Galr2 receptors in their craniofacial tissue,

M871's higher binding affinity was sufficient to inhibit these fewer receptors, resulting in reduced

Meckel's cartilage length (Fig. 7). In contrast, the presumably greater concentration of Galr2

receptors in molluscivore craniofacial tissue due to increased Galr2 expression during

development resulted in the inhibition of these receptors by both M871 and M35, despite its lower

binding affinity for Galr2 (Lang et al., 2007). We also conclude that inhibition of Galr1 receptors

by M35 does not affect Meckel's cartilage length or chondrocyte density and that the specific

inhibition of Galr2 by M871 was sufficient to drive equal Meckel's cartilage length shortening.

Finally, the generalists' lack of significant response to M35 is consistent with their decreased levels

of *galr2a* at 2 dpf, suggesting that, as in scale-eaters treated with M35, fewer Galr2 receptors may

be present for inhibition in these species' background. However, the response to M871 appears to

be specialist-specific, with little effect on the generalist Meckel's cartilage length, and thus, we

cannot rule out more complex species-specific factors interacting with Galr regulatory pathways.

Our work is consistent with previous work showing that *Galr2* activation inhibits cell

proliferation in neuronal cell lines (Lang et al., 2007; Berger et al., 2004), suggesting that the TFBS

allele frequency shifts observed in scale-eaters causing low *galr2a* abundance and, as a

consequence, have low Galr2 activity during craniofacial development, result in the increased

chondrocyte density and Meckel's cartilage length observed in scale-eaters. Thus, the increased

chondrocyte density observed in the M871-treated scale-eater larvae results likely from the strong

and continued inhibition of fewer Galr2 available during development as a consequence or a

synergy of/with their genetic background.

In conclusion, we propose that the inhibition of Galr2 and not Galr1 induces the reduction

of Meckel's cartilage length. Thus, a greater number of Galr2 receptors in the molluscivore

specialist, as a result of increased expression of *galr2a* during early development, may result in

their reduced oral jaw lengths as adults through increased opportunities for endogenous agonistic

binding interactions (Fig. 7). Fewer Galr2 receptors on the scale-eater jaw may result in their

enlarged jaw lengths as adults by limiting opportunities for Galr2 endogenous agonists to bind to

these receptors during development. In the generalists, a greater number of Galr2 receptors but

smaller overall Meckel's cartilage length may limit the sensitivity of this species to Galr2 and

Galr1+2 endogenous agonists, however, resulting in their intermediate length and least robust oral

jaws among the three species under control conditions (Hernandez et al. 2018; Martin and Wainwright 2011).

Figure 7. Proposed mechanism for how galanin receptor abundance affects Meckel's cartilage length in the two pupfish trophic specialists after exposure to M35 (weak Galr1+Galr2 antagonist) or M871 (strong Galr2 antagonist). Molluscivores (top row) express more *galr2a* at 2 and 8 dpf than scale-eaters (bottom row) in their Meckel's cartilage (see Figs. 3-4), resulting in more Galr2 receptors (open circles). Increased Galr2 receptor density in molluscivores results in their reduced Meckel's cartilage length

(see Fig. 6) due to inhibition by both the weak antagonist (M35: middle column; half-filled circles) and strong

antagonist (M871: right column; filled circles) of Galr2. In contrast, reduced Galr2 receptor density in scale-

eaters (bottom row) results in their reduced Meckel's cartilage length only after exposure to the strong

Galr2-specific antagonist M871.

Conclusion

Our results show a novel role of the second receptor for galanin, Galr2, as a craniofacial gene

important in controlling craniofacial development and interspecific divergence through modifying

its transcript abundance and receptor activity. *Galr2a* transcript abundance changes are associated

with genetic changes in the regulatory region of *galr2a*, consistent with the loss of a transcription

factor binding site in the scale-eating pupfish. We propose a model in which reduced Galr2

receptor density in the oral jaws of scale-eaters results in fewer endogenous agonistic interactions,

increasing Meckel's cartilage length presumably by decreased inhibition of chondrocyte

proliferation (as a downstream effect of endogenous Galr2 activation). We also acknowledge the

polygenic nature of jaw development and note that our previous genetic mapping experiments

place an upper bound of 15% of oral jaw variation that may be due to differences in *galr2a*

regulation among the myriad craniofacial genes driving the evolution of divergent craniofacial

traits in this system.

Materials and Methods

Genomic evidence for galr2 as a candidate craniofacial gene

Using a genome-wide association (GWA) test across 202 individuals from the SSI radiation and

outgroup populations, we previously found an association of the regulatory region of *galr2a* with

lower jaw length, confirming an earlier pilot study that found the *galr2a* region to be among the

top five strongest associations with lower jaw length containing highly differentiated SNPs

between trophic specialist species (Richards et al. 2021; McGirr and Martin 2017). An analysis of

hard selective sweeps using both site frequency spectrum (SweeD) and linkage disequilibrium

(Omegle) based summary statistics additionally found evidence of a putative adaptive allele in the

20 kb regulatory region upstream of *galr2* (Pavlidis et al., 2013; Richards et al., 2021). This allele

swept to fixation in the scale-eater *C. desquamator* population on SSI 696-1,008 (95% credible

interval) years ago, potentially providing a pivotal stage in adaptation to scale-eating (Richards et

al. 2021). An independent quantitative trait loci (QTL) mapping study in F2 intercross hybrids

between scale-eater and molluscivore parents found a significant QTL on linkage group 15

containing *galr2a* that accounted for 15.3% of the phenotypic variance in premaxilla length (*n* =

178; Martin et al. 2016). Similarly, a second independent study of F2 intercross hybrids from a

second lake found evidence of a QTL in this region explaining 8% of the phenotypic variance in

the length of the coronoid process on the articular bone of the lower jaw (jaw closing in-lever; *n* =

227; St. John et al. 2021). The combined strength of evidence for a role of *galr2a* in craniofacial

development across independent analyses of GWA, QTL, selective sweeps, and genetic

differentiation between species indicated that this was one of our highest priority candidates for

functional studies.

Pupfish husbandry and maintenance

Three sympatric *Cyprinodon* species (*C. variegatus, C. bronthotheroides,* and *C. desquamator*)

were collected from both Osprey Lake and Crescent Pond in San Salvador Island, Bahamas in

2018 and maintained in the lab for at least two generations in small breeding groups. All lab

colonies were maintained at a pH of 8.0 and 2-5 ppt salinity at 25-30°C on a 12:12 hours light:dark

cycle at the University of California, Berkeley. Newly hatched fry were raised on brine shrimp

nauplii for the first month and adults and juveniles were fed a combination of commercial pellet

food and frozen bloodworms. All protocols and procedures employed were reviewed and approved

by the University of California, Berkeley Animal Care and Use Committee (AUP-2015-01-7053)

and animals were collected and exported from the Bahamas with research and export permits from

the Bahamas Environmental, Science, and Technology commission through the Gerace Research

Centre.

Multiple breeding pairs for each population were housed together in 40-liter aquaria with

synthetic yarn spawning mops and fed ad libitum daily with frozen foods and commercial pellets.

Embryos were collected by handpicking and raised at 27°C in 5 ppt salinity water with the addition

of 10 mg/l gentamycin and 0.0001% methylene blue. Fertilization was confirmed after collection

and embryos were staged according to Lencer and McCune (2018). The day of collection was

noted as day "zero", and thus one day after collection was 1-day post-fertilization (dpf).

HCR FISH staining

Hybridization chain reaction (HCR) was used to visualize spatial expression of *galr2a* and *tpm3b*

at key development timepoints (Choi et al., 2018; Ibarra-García-Padilla et al., 2021). HCR

reagents, including probes, hairpins, and buffers, were purchased directly from Molecular

Instruments (Los Angeles, California, USA). Staining was performed according to a modified

protocol from Ibarra-García-Padilla et al., 2021, and Palominos and Whitlock, 2021.

Fixation and storage: 2 and 8 dpf larvae were fixed in 4% paraformaldehyde (Electron Microscopy

Science) 1X PBS (Invitrogen) overnight at 4°C. Samples were then serially dehydrated in

methanol/1X PBS (25%, 50%, 75%) and stored in 100% methanol at -20°C for at least 16 hours

before use. _Permeabilization:_ Rehydration was performed serially in methanol/PBST (1X PBS

0.1% Tween20, Fisher Scientific) up to 100% PBST. Samples were then incubated for 10 minutes

in pre-chilled acetone at -20°C for permeabilization. 2 and 8 dpf samples were then incubated for

20 and 80 minutes in 10 µg/ml Proteinase K (Thermofisher), respectively. Prehybridization:

Samples were then pre-hybridized in the pre-warmed (at 37°C) hybridization buffer for 30 minutes

at 37°C. _Hybridization:_ For _Galr2a_ and _Tpm3,_ 4 to 6 pmol and 2 to 3 pmol were used, respectively.

The probe mixture was pre-warmed for 30 minutes at 37°C before hybridization for 20 hours at

37°C. _Washing probe excess:_ After hybridization, we performed four washes in pre-warmed wash

buffer (15 minutes each) at 37°C, followed by two final washes in 5X SSCT (20X SSC from

Thermofisher; SSCT is SSC with 0.1% Tween20) for 5 minutes each. _Amplification:_

Preamplification involved incubating samples in pre-equilibrated (20-25°C) amplification buffer

for 30 minutes. Next, amplification was performed using 4 µl of each snap-cooled hairpin in 250

µl of amplification buffer for 16 hours in the dark at room temperature. _Galr2a_ expression was

confirmed independently using two amplifiers with different excitation/emission wavelengths: B1-

488 HCR amplifiers, and B1-594 HCR amplifiers. _Tpm3b_ amplifiers used were B2-647 HCR

amplifiers. _Washing hairpin excess:_ Excess hairpins were washed several times with 5X SSCT,

followed by a last wash with PBST. _Nuclei labeling:_ After rinsing, samples were incubated for 30

minutes in DAPI/1X PBST (1 µm/ml, Sigma) and then washed in PBST 3 times for 5 minutes

each. _Clearing and mounting:_ Samples were then serially cleared in glycerol solutions at room

temperature by sequentially removing and adding: 25% glycerol/75% PBST, 50% glycerol/50%

PBST, 75% glycerol/25% PBST. Samples were then mounted in a custom-made bridged

microscope slide (Fisher Scientific) in 75% glycerol/25% PBST.

Imaging: Samples were imaged using a Zeiss LSM880 (inverted) confocal microscope at the CNR

Biological Imaging Facility at the University of California, Berkeley. All HCR were volumetric z-

stack images of embryos or larvae positioned laterally or ventrally and taken every 3 μm. Images

were deconvoluted, analyzed, and edited in FIJI (Schindelin et al., 2012) and Imaris (Bitplane).

Galr2 inhibition during jaw development

According to the pupfish developmental series of Lencer and McCune (2018), the pharyngeal

arches of *C. variegatus* are visible as segmented pouches ventral to the developing head at Stage

25 (48-64 hours post-fertilization at 27 °C, or 2 – 2.5 dpf), while the presence of a stomodeum-

like oral cavity is evident by Stage 30 (100 – 130 hpf, or 4 – 5.5 dpf). We hypothesized that the

key stages for proper jaw development occur between Stages 24 and 25, and thus we dechorionated

Stage 24-25 embryos (or 2 to 2.5 dpf) pupfish by incubating them on 20 mg/ml Pronase (Roche)

in water for 2.5 hours. The remaining chorions were removed manually with forceps. After larvae

were dechorionated, we incubated them in 100 μM M35 (InnoPep Inc, San Diego, CA), a chimeric

peptide antagonist to all galanin receptors (Galr1 and Galr2 in teleost fishes; Butler et al. 2020)

and with a conserved 21 aminoacidic sequence to the core Galanin peptide sequence

(GWTLNSAGYLLGPPPGFSPFR-NH2), or M871 (Abcam, USA), a Galr2-specific antagonist

with the following aminoacidic sequence: WTLNSAGYLLGPEHPPPALALA-NH2), up to

hatching time (around 8 dpf, Stage 32-33).

Cartilage staining: a non-acid clearing and staining method was used based on Walker and

Kimmel, 2007, and Lencer and McCune, 2018. Fish larvae were cleared in 80% glycerol and

mounted in custom-made coverslips (Fisher Scientific). Meckel's length, inter-ocular distance, and

chondrocyte density were quantified using the segmented line tool and ROI manager in Fiji

(Schindelin et al., 2012). Imaging was done using a Zeiss Z1 AxioImager M2 microscope located

at the Biological Imagining Facility of the University of California, Berkeley.

Sequencing of Galr2a regulatory alleles

We used the DNeasy Blood and Tissue kit (Qiagen, Inc.) to extract DNA from the fins of wild-

caught adult fish and quantified it on a Qubit 3.0 fluorometer (Thermofisher Scientific, Inc.). All

specimens used in this study are kept in the Museum of Vertebrate Zoology at the University of

California, Berkeley. The primers used to amplify both *Galr2a* regulatory loci are listed in Table

S5. The sizes of the PCR products were confirmed by gel electrophoresis. PCR product clean-up

was done using ExoSap-IT (USB) and quantified on a Nanodrop ND-1000 (Thermo Scientific)

spectrophotometer. Sanger sequencing was performed at the UC Berkeley DNA Sequencing

Facility. Sequence alignment was performed using Geneious Prime (Dotmatics).

Prediction of transcription factors binding sites in *Galr2a-15kb* and *Galr2a-12kb*

We used the Multiple Expectation maximizations for Motif Elicitation (MEME) server

(http://meme-suite.org/) and the motif database scanning algorithm TOMTOM (Bailey et al.,

2009) to retrieve TFBS motifs from a 20kb conserved generalist, molluscivore, and scale-eater

sequences upstream *galr2a* obtained from Richards et al., 2021. MEME suite finds, in the given

sequences, the most statistically significant motifs first. We numbered the motifs following their

statistical significance (Table S1).

Statistics

Expression quantification of galr2a and tpm3b at 8 dpf: We segmented and quantified "jaw" *galr2a* expression in the maxilla, and the Meckel's and palatoquadrate cartilages and "brain" *galr2a* expression in the central nervous system from HCR confocal images using 3D-Slicer (Fedorov et al., 2012). *Tpm3b* expression was segmented and quantified in the intermandibular muscles (IMA), the protractor hyoideus (PR-H), the adductor mandibular complex (AddM), and the levator arcus palatini (LEVs). Muscles were named according to Diogo et al., 2008. Statistical tests comparing the expression volume among species and tissues were performed in R (R Core Team 2023). We tested for significant differences in means via ANOVA and contrasted groups post-hoc using Tukey's HSD test.

Expression quantification of garlr2a at 2 dpf: From the HCR confocal images, a total volume of 45 μm containing the pharyngeal arches was projected using the maximum intensity tool in Fiji. We selected two different tissues using ROI manager and quantified the relative fluorescence normalized by the background fluorescence of the image. First, we selected the mandibular mesenchyme plus the first and second pharyngeal arches (labeled in the figure "Pharyngeal Arches"). Then, we de-selected the second pharyngeal arch, quantifying only the expression of *galr2* in the mandibular mesenchyme and the first pharyngeal arch (labeled in the figure "Mandibular arch"). We then tested for significant differences in means via ANOVA and post-hoc analysis with Tukey's HSD in R.

mRNA transcript analysis: A table containing RPKM values of transcripts from Lencer and McCune, 2017, datasets was obtained from the following Dryad repository,

doi:10.5061/dryad.mb3gb. Genes of interest were found with the Gene ID listed on NCBI. Gene

IDs are as follows: 106090243, 107092788, 107090827, 107096257, 107098818, and 107092130,

for *galr2a, galr2b, galr1a, galr1b, gal,* and *sox21b*, respectively. Statistical tests comparing

transcripts RPKM expression between species and developmental stages were performed in R,

testing for significant differences in means via ANOVA and post-hoc analysis with Tukey's HSD.

Cartilage staining: We compared chondrocyte density for significant differences in means using

pairwise post-hoc Tukey's HSD tests. We used a generalized linear model accounting for species

and treatment and their interaction. We chose to measure chondrocytes within the Meckel's

cartilage because it allows for a clear view of each cell at the symphysis.

Author's contributions

Conceptualization: MFP, CHM; Data Collection: MFP, VM, EJR; Statistical analyses: MFP;

Resources: CHM; Microscopy: MFP, VM; Original draft: MFP; Revising: MFP, EJR, CHM.

Acknowledgments

We thank Michelle St. John, David Tian, Mara Van Tassell, Chloe Clair, Dylan Chau, and Heidi

Buratti for valuable comments and discussion of the results. We thank Lydia Smith for her help

throughout the development of this project in the Evolutionary Genetics Lab at the University of

California, Berkeley. We are particularly thankful to Austin H. Patton, who extracted the *C.*

brontotheroides mRNA reads to create the HCR probes tested here, and for his insightful ideas

and comments on the early stages of this manuscript. We thank the Gerace Research Centre and

Troy Day for logistical support and the government of the Bahamas for permission to collect and

export samples. This research was funded by the National Science Foundation DEB CAREER

grant #1749764, National Institutes of Health grant 5R01DE027052-02, the University of North

Carolina at Chapel Hill, and the University of California, Berkeley to CHM.

References

Albertson, R. C., Cresko, W., Detrich, H. W. and Postlethwait, J. H. (2009). Evolutionary

mutant models for human disease. *Trends in Genetics* **25**, 74–81.

Bonfante, B., Faux, P., Navarro, N., Mendoza-Revilla, J., Dubied, M., Montillot, C.,

Wentworth, E., Poloni, L., Varón-González, C., Jones, P., et al. (2021). A GWAS in

Latin Americans identifies novel face shape loci, implicating VPS13B and a Denisovan

introgressed region in facial variation. *Sci Adv* **7**,.

Boyle, E. A., Li, Y. I. and Pritchard, J. K. (2017). An Expanded View of Complex Traits: From

Polygenic to Omnigenic. *Cell* **169**, 1177–1186.

Brawand, D., Wagner, C. E., Li, Y. I., Malinsky, M., Keller, I., Fan, S., Simakov, O., Ng, A.

Y., Lim, Z. W., Bezault, E., et al. (2014). The genomic substrate for adaptive radiation in

African cichlid fish. *Nature* **513**, 375–381.

Buffenstein, R. (2008). Negligible senescence in the longest living rodent, the naked mole-rat:

Insights from a successfully aging species.

Carlborg, O. and Haley, C. S. (2004). Epistasis: too often neglected in complex trait studies?

Nature reviews. Genetics **5**, 618–625.

Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Shapiro, M. D., Brady, S. D.,

Southwick, A. M., Absher, D. M., Grimwood, J., Schmutz, J., et al. (2010). Adaptive

evolution of pelvic reduction in sticklebacks by recurrent deletion of a Pitx1 enhancer.

Science (New York, N.Y.) **327**, 302–305.

Concannon, M. R. and Albertson, R. C. (2015). The genetic and developmental basis of an

exaggerated craniofacial trait in East African cichlids. *J. Exp. Zool. B Mol. Dev. Evol.* **324**,

662–670.

Conith, A. J., Lam, D. T. and Albertson, R. C. (2019). Muscle-induced loading as an important

source of variation in craniofacial skeletal shape. *Genesis* **57**, e23263.

Erickson, P. A., Glazer, A. M., Cleves, P. A., Smith, A. S. and Miller, C. T. (2014). Two

developmentally temporal quantitative trait loci underlie convergent evolution of increased

branchial bone length in sticklebacks. *Proceedings of the Royal Society B: Biological*

Sciences **281**,.

Erickson, P. A., Ellis, N. A. and Miller, C. T. (2016). Microinjection for transgenesis and

genome editing in threespine sticklebacks. *J. Vis. Exp.* 10.3791/54055.

Glazer, A. M., Killingbeck, E. E., Mitros, T., Rokhsar, D. S. and Miller, C. T. (2015). Genome

assembly improvement and mapping convergently evolved skeletal traits in sticklebacks

with genotyping-by-sequencing. *Genes,Genomes,Genetics* **5**, 1463–1472.

Glazier, A. M. (2002). Finding genes that underlie complex traits. *Science (New York, N.Y.)* **298**,

2345–2349.

Gross, J. B., Stahl, B. a., Powers, A. K. and Carlson, B. M. (2016). Natural bone fragmentation
in the blind cave-dwelling fish, *Astyanax mexicanus*: Candidate gene identification through
integrative comparative genomics. *Evolution and Development* **18**, 7–18.

Hall, D., Tegstrom, C. and Ingvarsson, P. K. (2010). Using association mapping to dissect the
genetic basis of complex traits in plants. *Briefings in functional genomics* **9**, 157–165.

Hawes, J. J., Narasimhaiah, R. and Picciotto, M. R. (2006). Galanin and galanin-like peptide
modulate neurite outgrowth via protein kinase C-mediated activation of extracellular
signal-related kinase. *Eur. J. Neurosci.* **23**, 2937–2946.

Helman, S. N., Badhey, A., Kadakia, S. and Myers, E. (2014). Revisiting Crouzon syndrome:
reviewing the background and management of a multifaceted disease. *Oral and*
Maxillofacial Surgery **18**, 373–379.

Hernandez, L. P., Adriaens, D., Martin, C. H., Wainwright, P. C., Masschaele, B. and
Dierick, M. (2018). Building trophic specializations that result in substantial niche
partitioning within a young adaptive radiation. *J. Anat.* **232**, 173–185.

Hirschhorn, J. N. and Daly, M. J. (2005). Genome-wide association studies for common diseases
and complex traits. *Nature reviews* **6**, 95–108.

Hochheiser, H., Aronow, B. J., Artinger, K., Beaty, T. H., Brinkley, J. F., Chai, Y., Clouthier,
D., Cunningham, M. L., Dixon, M., Donahue, L. R., et al. (2011). The FaceBase
Consortium: A comprehensive program to facilitate craniofacial research. *Developmental*
Biology **355**, 175–182.

Hofmann, C. M., O'Quin, K. E., Justin Marshall, N., Cronin, T. W., Seehausen, O. and

Carleton, K. L. (2009). The eyes have it: Regulatory and structural changes both underlie

cichlid visual pigment diversity. *PLoS Biol.* **7**,.

Humphries, J. M. and Miller, R. R. (1981). A remarkable species flock of pupfishes, genus

Cyprinodon, from Yucatan, Mexico. *Copeia* **1981**, 52–64.

J. Parsons, K., McWhinnie, K. and Armstrong, T. (2021). An Evo-devo View of Post-genomic

African Cichlid Biology: Enhanced Models for Evolution and Biomedicine. In *The*

Behavior, Ecology and Evolution of Cichlid Fishes (ed. Abate, M. E.) and Noakes, D. L.

G.), pp. 779–802. Dordrecht: Springer Netherlands.

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., Swofford,

R., Pirun, M., Zody, M. C., White, S., et al. (2012). The genomic basis of adaptive

evolution in threespine sticklebacks. *Nature* **484**, 55–61.

Kadakia, S., Helman, S. N., Badhey, A. K., Saman, M. and Ducic, Y. (2014). Treacher Collins

Syndrome: The genetics of a craniofacial disease.

Kocher, T. D. (2004). Adaptive evolution and explosive speciation: The cichlid fish model. *Nat.*

Rev. Genet. **5**, 288–298.

Kocher, T. D., Albertson, R. C., Carleton, K. L. and Streelman, J. T. (2003). The genetic basis

of biodiversity, genomic studies of cichlid fishes. *Aquatic genomics: steps toward a great*

future. 34–44.

Kolora, S. R. R., Owens, G. L., Vazquez, J. M., Stubbs, A., Chatla, K., Jainese, C., Seeto, K.,
McCrea, M., Sandel, M. W., Vianna, J. A., et al. (2021). Origins and evolution of
extreme life span in Pacific Ocean rockfishes. *Science* **374**, 842–847.

Lencer, E. S., Riccio, M. L. and McCune, A. R. (2016). Changes in growth rates of oral jaw
elements produce evolutionary novelty in bahamian pupfish. *J. Morphol.* **277**, 935–947.

Lencer, E. S., Warren, W. C., Harrison, R. and McCune, A. R. (2017). The Cyprinodon
variegatus genome reveals gene expression changes underlying differences in skull
morphology among closely related species. *BMC Genomics* **18**, 424.

Li, L., Wei, S., Huang, Q., Feng, D., Zhang, S. and Liu, Z. (2013). A novel galanin receptor 1a
gene in zebrafish: tissue distribution, developmental expression roles in nutrition
regulation. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **164**, 159–167.

Liu, S., Lorenzen, E. D., Fumagalli, M., Li, B., Harris, K., Xiong, Z., Zhou, L., Korneliussen,
T. S., Somel, M., Babbitt, C., et al. (2014). Population genomics reveal recent speciation
and rapid evolutionary adaptation in polar bears. *Cell* **157**, 785–794.

Ma, A., Bai, J., He, M. and Wong, A. O. L. (2018). Spexin as a neuroendocrine signal with
emerging functions. *Gen. Comp. Endocrinol.* **265**, 90–96.

Mack, K. L. and Nachman, M. W. (2017). Gene Regulation and Speciation. *Trends Genet.* **33**,
68–80.

Man, P.-S. and Lawrence, C. B. (2008). The effects of galanin-like peptide on energy balance,

body temperature and brain activity in the mouse and rat are independent of the GALR2/3

receptor. *J. Neuroendocrinol.* **20**, 128–137.

Manolio, T. A., Collins, F. S., Cox, N. J., Goldstein, D. B., Hindorff, L. A., Hunter, D. J.,

McCarthy, M. I., Ramos, E. M., Cardon, L. R., Chakravarti, A., et al. (2009). Finding

the missing heritability of complex diseases. *Nature* **461**, 747–753.

Martin, C. H. (2012). Weak disruptive selection and incomplete phenotypic divergence in two

classic examples of sympatric speciation: Cameroon crater lake cichlids. *Am. Nat.* **180**,

E90–E109.

Martin, C. H. (2013). Strong assortative mating by diet, color, size, and morphology but limited

progress toward sympatric speciation in a classic example: Cameroon crater lake cichlids.

Evolution **67**, 2114–2123.

Martin, C. H. (2016). The cryptic origins of evolutionary novelty: 1000-fold faster trophic

diversification rates without increased ecological opportunity or hybrid swarm. *Evolution*.

Martin, C. H. and Genner, M. J. (2009). High niche overlap between two successfully coexisting

pairs of Lake Malawi cichlid fishes. *Canadian Journal of Fisheries and Aquatic Sciences*

66, 579–588.

Martin, C. H. and Gould, K. J. (2020). Surprising spatiotemporal stability of a multi-peak fitness

landscape revealed by independent field experiments measuring hybrid fitness. *Evol Lett*

4, 530–544.

Martin, C. H. and Wainwright, P. C. (2011). Trophic novelty is linked to exceptional rates of

morphological diversification in two adaptive radiations of *Cyprinodon* pupfishes.

Evolution **65**, 2197–2212.

Martin, C. H. and Wainwright, P. C. (2013a). A remarkable species flock of *Cyprinodon*

pupfishes endemic to San Salvador Island, Bahamas. *Bulletin of the Peabody Museum of*

Natural History **54**, 231–240.

Martin, C. H. and Wainwright, P. C. (2013b). Multiple fitness peaks on the adaptive landscape

drive adaptive radiation in the wild. *Science (New York, N.Y.)* **339**, 208–211.

Martin, C. H., Cutler, J. S., Friel, J. P., Dening, T., Coop, G. and Wainwright, P. C. (2015).

Complex histories of repeated colonization and hybridization cast doubt on the clearest

examples of sympatric speciation in the wild. *Evolution* **69**, 1406–1422.

Martin, C. H., Erickson, P. A. and Miller, C. T. (2017). The genetic architecture of novel trophic

specialists: higher effect sizes are associated with exceptional oral jaw diversification in a

pupfish adaptive radiation. *Molecular Ecology* **26**, 624–638.

Martin, C. H., McGirr, J. A., Richards, E. J. and St John, M. E. (2019). How to Investigate

the Origins of Novelty: Insights Gained from Genetic, Behavioral, and Fitness

Perspectives. *Integr Org Biol* **1**, obz018.

Matsumoto, Y., Watanabe, T., Adachi, Y., Itoh, T., Ohtaki, T., Onda, H., Kurokawa, T.,

Nishimura, O. and Fujino, M. (2002). Galanin-like peptide stimulates food intake in the

rat. *Neurosci. Lett.* **322**, 67–69.

McGirr, J. A. and Martin, C. H. (2016). Novel candidate genes underlying extreme trophic
specialization in Caribbean pupfishes. *Molecular Biology and Evolution* **34**, 873–888.

McGirr, J. A. and Martin, C. H. (2021). Few Fixed Variants between Trophic Specialist Pupfish
Species Reveal Candidate Cis-Regulatory Alleles Underlying Rapid Craniofacial
Divergence. *Mol. Biol. Evol.* **38**, 405–423.

Meng, L., Bian, Z., Torensma, R. and Von den Hoff, J. W. (2009). Biological mechanisms in
palatogenesis and cleft palate. *Journal of Dental Research* **88**, 22–23.

Miller, C. T., Beleza, S., Pollen, A. A., Schluter, D., Kittles, R. A., Shriver, M. D. and
Kingsley, D. M. (2007). cis-Regulatory changes in Kit ligand expression and parallel
evolution of pigmentation in sticklebacks and humans. *Cell* **131**, 1179–1189.

Monteiro, A. and Podlaha, O. (2009). Wings, horns, and butterfly eyespots: how do complex
traits evolve? *PLoS biology* **7**, e37.

Naqvi, S., Hoskens, H., Wilke, F., Weinberg, S. M., Shaffer, J. R., Walsh, S., Shriver, M. D.,
Wysocka, J. and Claes, P. (2022). Decoding the Human Face: Progress and Challenges
in Understanding the Genetics of Craniofacial Morphology. *Annu. Rev. Genomics Hum.*
Genet.

Navon, D., Male, I., Tetrault, E. R., Aaronson, B., Karlstrom, R. O. and Albertson, R. C.
(2020). Hedgehog signaling is necessary and sufficient to mediate craniofacial plasticity in
teleosts. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 19321–19327.

Palmer, K., Fairfield, H., Borgeia, S., Curtain, M., Hassan, M. G., Dionne, L., Yong Karst,
S., Coombs, H., Bronson, R. T., Reinholdt, L. G., et al. (2016). Discovery and
characterization of spontaneous mouse models of craniofacial dysmorphology.
Developmental Biology **415**, 216–227.

Pavlidis, P., ??ivkovi??, D., Stamatakis, A. and Alachiotis, N. (2013). SweeD: Likelihood-
based detection of selective sweeps in thousands of genomes. *Mol. Biol. Evol.* **30**, 2224–
2234.

Platzer, M. and Englert, C. (2016). *Nothobranchius furzeri*: A model for aging research and
more. *Trends in Genetics* **32**, 543–552.

Powder, K. E. and Albertson, R. C. (2016). Cichlid fishes as a model to understand normal and
clinical craniofacial variation. *Developmental Biology* **415**, 338–346.

Price, A. L., Spencer, C. C. A. and Donnelly, P. (2015). Progress and promise in understanding
the genetic basis of common diseases. *Proceedings of the Royal Society B: Biological*
Sciences **282**, 20151684.

Protas, M., Conrad, M., Gross, J. B., Tabin, C. and Borowsky, R. (2007). Regressive evolution
in the Mexican cave tetra, Astyanax mexicanus. *Current biology : CB* **17**, 452–454.

Reardon, W., Winter, R. M., Rutland, P., Pulleyn, L. J., Jones, B. M. and Malcolm, S. (1994).
Mutations in the fibroblast growth factor receptor 2 gene cause Crouzon syndrome. *Nature*
Genetics **8**, 98–103.

Richards, E. J. and Martin, C. H. (2017). Adaptive introgression from distant Caribbean islands
contributed to the diversification of a microendemic adaptive radiation of trophic specialist
pupfishes. *PLoS Genet.* **13**, 1–35.

Richards, E. J. and Martin, C. H. (2022). We get by with a little help from our friends: shared
adaptive variation provides a bridge to novel ecological specialists during adaptive
radiation. *Proc. Biol. Sci.* **289**, 20220613.

Richards, E. J., McGirr, J. A., Wang, J. R., St John, M. E., Poelstra, J. W., Solano, M. J.,
O'Connell, D. C., Turner, B. J. and Martin, C. H. (2021). A vertebrate adaptive
radiation is assembled from an ancient and disjunct spatiotemporal landscape. *Proc. Natl.*
Acad. Sci. U. S. A. **118**,.

Roberts, R. B., Ser, J. R. and Kocher, T. D. (2009). Sexual conflict resolved by invasion of a
novel sex determiner in Lake Malawi cichlid fishes. *Science (New York, N.Y.)* **326**, 998–
1001.

Roberts, R. B., Hu, Y., Albertson, R. C. and Kocher, T. D. (2011). Craniofacial divergence and
ongoing adaptation via the hedgehog pathway. *Proceedings of the National Academy of*
Sciences of the United States of America **108**, 13194–13199.

Rohner, N. (2018). Cavefish as an evolutionary mutant model system for human disease. *Dev.*
Biol. **441**, 355–357.

Signor, S. A. and Nuzhdin, S. V. (2018). The Evolution of Gene Expression in cis and trans.
Trends Genet. **34**, 532–544.

Sollenberg, U. E., Lundström, L., Bartfai, T. and Langel, Ü. (2006). M871—A Novel Peptide
Antagonist Selectively Recognizing the Galanin Receptor Type 2. *Int. J. Pept. Res. Ther.*
12, 115–119.

Sollenberg, U. E., Runesson, J., Sillard, R. and Langel, Ü. (2010). Binding of Chimeric Peptides
M617 and M871 to Galanin Receptor Type 3 Reveals Characteristics of Galanin Receptor–
Ligand Interaction. *Int. J. Pept. Res. Ther.* **16**, 17–22.

St John, M. E., Holzman, R. and Martin, C. H. (2020a). Rapid adaptive evolution of scale-
eating kinematics to a novel ecological niche. *J. Exp. Biol.* **223**,.

St John, M. E., Dixon, K. E. and Martin, C. H. (2020b). Oral shelling within an adaptive
radiation of pupfishes: Testing the adaptive function of a novel nasal protrusion and
behavioural preference. *J. Fish Biol.* **97**, 163–171.

St. John, M. E., Dunker, J. C., Richards, E. J., Romero, S. and Martin, C. H. (2021). Parallel
genetic changes underlie integrated craniofacial traits in an adaptive radiation of trophic
specialist pupfishes. *bioRxiv* 2021.07.01.450661.

Sutter, N. B., Bustamante, C. D., Chase, K., Gray, M. M., Zhao, K., Zhu, L.,
Padhukasahasram, B., Karlins, E., Davis, S., Jones, P. G., et al. (2007). A single IGF1
allele is a major determinant of small size in dogs. *Science (New York, N.Y.)* **316**, 112–115.

Venn-watson, S., Carlin, K. and Ridgway, S. (2011). General and Comparative Endocrinology
Dolphins as animal models for type 2 diabetes : Sustained , post-prandial hyperglycemia
and hyperinsulinemia. *General and Comparative Endocrinology* **170**, 193–199.

Weinberg, S. M., Cornell, R. and Leslie, E. J. (2018). Craniofacial genetics: Where have we

been and where are we going? *PLoS Genet.* **14**, e1007438.

Wilkie, A. O., Slaney, S. F., Oldridge, M., Poole, M. D., Ashworth, G. J., Hockley, A. D.,

Hayward, R. D., David, D. J., Pulleyn, L. J. and Rutland, P. (1995). Apert syndrome

results from localized mutations of FGFR2 and is allelic with Crouzon syndrome. *Nature*

genetics **9**, 165–172.