Jaw size variation is associated with a novel craniofacial function for galanin receptor 2 in

an adaptive radiation of pupfishes

M. Fernanda Palominos[1,2], Vanessa Muhl[1,2], Emilie J. Richards[3], Craig T. Miller[4], Christopher

H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley

[2]Museum of Vertebrate Zoology, University of California, Berkeley

[3]Department of Ecology, Evolution, and Behavior, University of Minnesota

[4]Department of Molecular & Cell Biology, University of California, Berkeley

Keywords: gene expression, fluorescence in situ hybridization, galanin, trophic morphology,

craniofacial divergence, evo devo, adaptive radiation

Abstract

Understanding the genetic basis of novel adaptations in new species is a fundamental question in

biology that also provides an opportunity to uncover new genes and regulatory networks with

potential clinical relevance. Here we demonstrate a new role for *galr2* in vertebrate craniofacial

development using an adaptive radiation of trophic specialist pupfishes endemic to San Salvador

Island in the Bahamas. We confirmed the loss of a putative *Sry* transcription factor binding site

in the upstream region of *galr2* in scale-eating pupfish and found significant spatial differences

in *galr2* expression among pupfish species in Meckel's cartilage and premaxilla using in situ

hybridization chain reaction (HCR). We then experimentally demonstrated a novel function for

Galr2 in craniofacial development and jaw elongation by exposing embryos to drugs that inhibit

Galr2 activity. Galr2-inhibition reduced Meckel's cartilage length and increased chondrocyte

density in both trophic specialists but not in the generalist genetic background. We propose a

mechanism for jaw elongation in scale-eaters based on the reduced expression of *galr2* due to the

loss of a putative *Sry* binding site. Fewer Galr2 receptors in the scale-eater Meckel's cartilage

may result in their enlarged jaw lengths as adults by limiting opportunities for a postulated Galr2

agonist to bind to these receptors during development. Our findings illustrate the growing utility

of linking candidate adaptive SNPs in non-model systems with highly divergent phenotypes to

novel vertebrate gene functions.

Introduction

Craniofacial developmental anomalies are the most common source of birth defects in humans, present in 1 out of 700 births (Hall et al., 2010; Meng et al., 2009; Palmer et al., 2016). While Mendelian craniofacial defects are well characterized (e.g. Treacher Collins Syndrome (Kadakia et al., 2014), Apert Syndrome (Wilkie et al., 1995), and Crouzon Syndrome (Helman et al., 2014; Reardon et al., 1994)), the developmental genetics of complex craniofacial defects, such as micrognathia, are poorly understood (Carlborg and Haley, 2004; Glazier, 2002; Hirschhorn and Daly, 2005; Hochheiser et al., 2011; Manolio et al., 2009; Monteiro and Podlaha, 2009; Weinberg et al., 2018). With the continued lowering costs of genomic sequencing and functional genetic tools, it is increasingly feasible to develop new genetic models for understanding human development and disease.

Current knowledge of craniofacial developmental genetics results mainly from mutagenic screens and forward genetic approaches in a few vertebrate model systems (primarily mice and zebrafish); however, these approaches are typically biased toward large Mendelian effects on easily detectable phenotypes, often missing the variation most relevant to common craniofacial birth defects in humans, such as small-effect variants, subtle differences in skeletal shape, and epistatic interactions among genes and the environment (Albertson et al., 2009; Powder and Albertson, 2016; Roberts et al., 2011). Conversely, complementary genome-wide association studies (GWAS) of craniofacial syndromes and normal-range facial variation in humans often reveal small-effect loci with no tractable possibilities for functional interrogation (Bonfante et al., 2021; Naqvi et al., 2022; Weinberg et al., 2018). Thus, understanding the genetic bases of naturally occurring, highly divergent adaptive phenotypes in novel systems that parallel human clinical variation, such as 'evolutionary mutant' models (Albertson et al., 2009; Concannon and

Albertson, 2015; Powder and Albertson, 2016; Rohner, 2018), provides a powerful approach

combining the tractable functional investigations possible in vertebrate model systems with

genome-wide association scans of small-effect regulatory loci underlying natural craniofacial

diversity. For example, novel insights into a broad diversity of human diseases have now arisen

from studies of diverse species experiencing strong selective pressures in nature, including

senescence (i.e., naked mole rat (Buffenstein, 2008); turquoise killifish (Platzer and Englert,

2016), and rockfish (Kolora et al., 2021)); eye degeneration, bone loss and enhanced lipogenesis

(blind cavefishes (Gross et al., 2016; Hofmann et al., 2009; Protas et al., 2007; Lam et al. 2022));

body size (domesticated dogs (Sutter et al., 2007). In particular, the most remarkable diversity of

vertebrate craniofacial morphology is represented in teleost fishes, often associated with their

diverse and sometimes highly specialized modes of feeding (e.g., Liem, 1991; Otten, 1983;

Albertson et al., 2003; Martin and Richards 2019; Konings et al., 2021; Evans et al. 2021).

Emerging fish model systems include the rapidly evolving East African and Cameroon

cichlid radiations, in which a small number of genetic changes underlie immense morphological

disparity in over 1,500 cichlid species (Brawand et al., 2014; Conith et al., 2019; Kocher et al.,

2003; Martin, 2012; Martin, 2013; Martin and Genner, 2009; Navon et al., 2020) and the

repeated parallel speciation of stickleback ecomorphs in glacial lakes (Chan et al., 2010;

Erickson et al., 2016; Jones et al., 2012; Miller et al., 2007; Ishikawa et al. 2019). These systems

provide excellent examples of leveraging naturally-occurring and highly divergent craniofacial

phenotypes as 'evolutionary mutants' or 'evolutionary forward genetics' models to gain novel

insights into the genetics of natural human craniofacial variation (Erickson et al., 2014; Glazer et

al., 2015; J. Parsons et al., 2021; Kocher, 2004; Powder and Albertson, 2016; Roberts et al.,

2009).

Figure 1. Divergent craniofacial morphology and development of the San Salvador Island *Cyprinodon* **pupfish radiation.** *C. variegatus* (first row) is a trophic generalist distributed across the western Atlantic and Caribbean; *C. brontotheroides* (second row) is a molluscivore, and *C. desquamator* (third row) is a scale-eater, both endemic to the hypersaline lakes of San Salvador Island (SSI), Bahamas. Left panel: development at 1-, 3-, and 8-days post-fertilization (dpf). Middle panel: lab-reared adult female pupfishes of each species. Right panel: Lateral and dorsal views of µCT scans of craniofacial morphology (modified from Hernández et al., 2017). The maxilla is colored in blue, premaxilla in red, dentary in green, articular in orange.

Here we demonstrate the utility of a an evolutionary radiation of *Cyprinodon* pupfishes for discovering and validating the craniofacial function of a new gene associated with jaw evolution and identified by QTL and GWAS analyses. Pupfishes offer some advantages over other evolutionary fish systems because they 1) rapidly evolved highly divergent and unique craniofacial phenotypes (Fig. 1) with minimal genetic differentiation among species (Hernandez et al., 2018; Lencer et al., 2016; Lencer et al., 2017; Martin and Wainwright, 2011; Martin and Wainwright, 2013a; Martin et al., 2019; St John et al., 2020a; St John et al., 2020b), 2) speciated in the face of ongoing gene flow resulting in very few highly differentiated genomic regions associated with species-specific craniofacial traits (Martin and Feinstein 2014; Martin et al.,

2017; McGirr and Martin, 2016; McGirr and Martin, 2021; Richards and Martin, 2017; Richards

and Martin, 2022; Richards et al., 2021), and 3) are highly amenable to laboratory rearing and

imaging due to their high fecundity, daily egg production, and egg transparency comparable to

zebrafish (Martin and Gould, 2020; Martin and Wainwright, 2013b). This radiation contains the

widespread algae-eating generalist pupfish, *Cyprinodon variegatus* (Fig. 1a), which is broadly

distributed across the Caribbean and North American Atlantic coast and occurs in sympatry with

two microendemic trophic specialist species found only in the hypersaline lakes of San Salvador

Island (SSI), Bahamas. Each trophic specialist displays highly divergent behavior, pigmentation,

and craniofacial morphology: the molluscivore *C. brontotheroides*, has a novel nasal protrusion

that is a skeletal extension of the maxilla and foreshortened robust oral jaw (Fig. 1b); and the

scale-eater, *C. desquamator*, exhibits two-fold larger oral jaws and overall brachycephalic

features (Fig. 1c) (Martin, 2016; Martin and Wainwright, 2011; Martin and Wainwright, 2013a).

There is also a fourth intermediate scale-eating ecotype in some lakes (Richards et al. 2022).

Previous genomic and transcriptomic work on the SSI pupfishes identified dozens of new

candidate craniofacial genes never previously characterized as craniofacial or directly

investigated in other systems (McGirr and Martin 2016, 2019, 2020; Richards et al. 2021, 2022;

Richards and Martin 2017; Patton et al. 2021; St. John et al. 2021). One of the most promising

candidates was galanin receptor 2a, the second receptor type for galanin. Using a genome-wide

association (GWA) test across 202 individuals from the SSI radiation and outgroup populations,

we previously found an association of the regulatory region of *galr2a* with lower jaw length,

confirming an earlier pilot study that found the *galr2a* region to be among the top five strongest

associations with lower jaw length, containing highly differentiated SNPs between trophic

specialist species (Richards et al. 2021; McGirr and Martin 2017). An analysis of hard selective

sweeps using both site frequency spectrum (SweeD) and linkage disequilibrium (Omegle) based

summary statistics additionally found evidence of a putative adaptive allele in the 20 kb

regulatory region upstream of *galr2* that swept to fixation in the scale-eater *C. desquamator*

population on SSI 696-1,008 (95% credible interval) years ago, potentially providing a pivotal

stage in adaptation to scale-eating (Richards et al. 2021). Furthermore, an independent

quantitative trait loci (QTL) mapping study in F2 intercross hybrids between scale-eater and

molluscivore parents found a significant QTL on linkage group 15 containing *galr2a* that

accounted for 15.3% of the phenotypic variance in premaxilla length (n = 178; Martin et al.

2016), with a positive effect on jaw length in the scale-eater genotype. Similarly, a second

independent study of F2 hybrid intercross from a second lake found evidence of a QTL in this

region explaining 8% of the phenotypic variance in the length of the coronoid process on the

articular bone of the lower jaw (jaw closing in-lever; n = 227; St. John et al. 2021). The

combined strength of evidence for a role of *galr2a* in craniofacial development across

independent analyses of GWA, QTL, selective sweeps, and genetic differentiation between

species indicated that this was one of our highest priority candidates for functional studies.

In humans, *galr2* is abundantly expressed within the hypothalamus and hippocampus of

the central nervous system and in the heart, kidney, liver, colon, and small intestine, and has

genetic associations with epilepsy and Alzheimer's (Li et al., 2013; Ma et al., 2018). Classified

as an orexigenic (appetite stimulant) gene, *galr2* is expressed in the human hypothalamus but

also in the ventral telencephalon of larval and adult zebrafish (Ahi et al., 2019; Li et al., 2013;

Kim et al., 2016). Although a role of *galr2* in craniofacial development has not previously been

reported in the literature, its ligand, the neuropeptide galanin (GAL), is highly expressed in

bones from early to post-embryonic development (Xu et al., 1996, Jones et al., 2009), with

demonstrated effects on bone mass (Idelevich et al., 2018; Idelevich et al., 2019), muscle

contraction (Kakuyama et al., 1997. EJP), and periodontal regeneration (Ma et al. 2021).

Here we used Sanger sequencing to confirm two highly differentiated SNPs between SSI

specialist species detected in our previous genomic studies affecting two predicted transcription

factor binding sites in the *galr2* regulatory region, characterized the divergent craniofacial

expression of *galr2* across all three SSI pupfish species using *in situ* hybridization chain reaction

(HCR) (Choi et al., 2018; Ibarra-Padilla-García et al., 2020) at two key developmental

timepoints, and demonstrated that treatment with two different Galr2 receptor inhibitors reduced

Meckel's cartilage length and increased chondrocyte density dependent on the species' genetic

backgrounds.

Results

Two highly differentiated SNPs are associated with *galr2a* transcription factor binding sites

Previous whole genome resequencing of over one hundred San Salvador Island (SSI) pupfishes

identified only two highly differentiated single nucleotide polymorphisms (SNP) within the 20

kb regulatory region of *galr2a* between trophic specialists across different lake populations on

SSI (Richards et al. 2021). We designed primers (Table S5) and used Sanger sequencing to

further genotype these SNPs in a large panel of wild-caught specialists from six lake populations.

We confirmed the presence of a transversion from G to A approximately 11 kb upstream of the

galr2a transcription starting site (TSS; Fig. 2a., S1; Table S1), which changes a predicted

CA**G**CAA *Elf1/Erg* transcription factor binding site (TFBS) to a predicted **A**GGGASW *Elf5*

TFBS at this locus (using the Multiple Expectation maximizations for Motif Elicitation (MEME)

server and the motif database scanning algorithm TOMTOM (Bailey et al., 2009)). This

transversion was observed in 86.3% of scale-eaters across four lake populations ($n = 51$) versus

20% of molluscivores across six lake populations on SSI ($n = 30$) (Fig. 2a., S1).

Using Sanger sequencing, we genotyped a second transversion from C to T

approximately 8 kb upstream of *galr2a* TSS, which changes the predicted AGA**C**AA *Sry* TFBS

to a predicted YAGA**T**A *Znf146* TFBS (Fig. 2b., S2; Table S1). This transversion was observed

in 90.9% of scale-eaters across six lake populations ($n = 55$) versus 33.9% of molluscivores

across six lake populations on SSI ($n = 53$) (Figure S2). Notably, all scale-eaters sampled from

Crescent Pond (CRP) contained the T transversion ($n = 30$) (Figure S2). The predicted TFBS

changes in the *galr2a* cis-regulatory region across pupfishes, combined with a previous genetic

mapping study that found a significant QTL in this region explaining 15% of phenotypic

variance in oral jaw size (Martin et al. 2017), suggests that different spatial or temporal *galr2a*

expression may underlie some of the craniofacial divergence in SSI pupfishes. Previous studies

of allele-specific expression in SSI pupfishes were inconclusive due to lack of heterozygous sites

in the *galr2a* transcripts (McGirr and Martin 2021).

Figure 2. The two most differentiated single-nucleotide polymorphisms (SNP) between trophic specialists lie within the 20kb regulatory region of *galr2a*. a. Putative ancestral TFBS for *Elf1/Erg* and *Sry*, respectively, in the generalist (G), *Cyprinodon variegatus* and the molluscivore (M) *Cyprinodon brontotheroides*. **B.** Derived changes in predicted TFBS to *Elf5* and *Znf146* in the scale-eater (S) *Cyprinodon desquamator*. Pie charts indicate frequencies of ancestral (teal) and derived (orange) alleles at each locus. +1 represents the transcription starting site of *galr2a* (Galanin receptor 2a). μCT scans of craniofacial morphology are modified from Hernández et al., 2017. Sequence logos of the Position Weight Matrixes (PWM) identified using MEME and TOMTOM.

Different spatiotemporal patterns of *galr2a* expression in craniofacial tissues

To determine if spatial or temporal changes in *galr2a* expression underlie SSI craniofacial

divergence, we assayed *galr2a* expression in 2 dpf embryos and 8 dpf larvae from two

independent lake populations for each of the three SSI species using hybridization chain reaction

(HCR) in-situ hybridization for *galr2a* and *tropomyosin 3b* (*tpm3b*), a component of thin

filaments of myofibrils expressed in fish skeletal muscles (Dube et al., 2016), to visualize jaw

and other cranial muscles. We also tested *galr2a* expression using orthogonal amplifiers labeled

with three distinct fluorophores (see Methods for details) to ensure reliable detection of *galr2a*

transcripts *in situ* across species and developmental stages (Supp. Video S1).

At 2 dpf, *galr2a* expression was detected in broad regions of the central nervous system

(e.g. the posterior tectum and medial longitudinal fasciculus) of specialists (Fig. 3a,c) and

generalists, with apparent higher expression in both specialist species than in generalists. We

also observed an apparent increase of *galr2a* expression anterior to the first pharyngeal arches in

the molluscivore and generalist pupfishes relative to the scale-eaters.

Using 3D reconstruction and volume rendering analysis of HCR data for whole-mounted

pupfishes at hatching time (8 dpf), we found that *galr2a* expression domain was expanded in the

jaws of the molluscivores than in the generalists (Fig. 4; $P = 0.03$, Tukey's HSD), consistent

with either greater tissue volume or an increased gene expression domain. In contrast, *galr2a*

showed no differences in expression volume among species in the brain and head (Fig. 4, Table

S3). G*alr2a* expression was detected in the Meckel's and palatoquadrate cartilages in all SSI

pupfishes, in the premaxilla of the generalist and the scale-eater SSI specialist (Fig. 4 b-b', c-c'),

the maxilla of the molluscivore SSI specialist (Fig. 4 a-a'), and in the intermandibular muscles of

the scale-eaters (Fig. 4 c').

At the subcellular level, *galr2a* mRNA was detected in the cytoplasm of chondrocytes at

the Meckel's symphysis (Fig. 5a). In contrast, on the distal edge of the Meckel's cartilage

expression was detected only in the cytoplasm of the elongated chondrocytes (Fig. 5a'). Towards

the most posterior region of the Meckel's cartilage closest to the palatoquadrate cartilage, we

observed *galr2a* expression in the cells surrounding the jaw joint (Fig. 5 b, c, d; Supplementary

Video S1). We conclude that g*alr2a* was significantly differentially expressed in specific and

distinct craniofacial tissues in the specialists at hatching time, suggesting an important role for

craniofacial divergence in the SSI radiation.

We further tested for quantitative differences *in galr2a* expression at 2 dpf and 8 dpf by

quantifying transcript counts from the only existing RNAseq data set on SSI pupfishes heads

during development (Lencer and McCune, 2020). Only *galr2a*, but not *galr1a, galr1b, galr2a,*

galr2b, or *galanin*, showed significantly higher mRNA expression in the molluscivores relative

to generalists (P = 0.003, Tukey's HSD test) and scale-eaters (P = 0.014, Tukey's HSD test) at 2

dpf (Table S2), whereas *galr2a* showed overall similar levels of expression in the head from 4 to

15 dpf in all three species (Table S2).

By using *Tpm3b* expression to label muscle cells, we observed at 2 dpf that *Tpm3b*

expression was detected in somitic myofibers in all SSI pupfishes (Fig. 3a''-c''). However, only

in the scale-eaters, *Tpm3b* expression was detected in the eye's inferior oblique muscle

primordial cells (Fig. 3c''). At hatching time, *Tpm3b* expression was detected in all larval head

muscles (Fig. 4a-a', b-b', c-c'; 5b). The scale-eater and molluscivore *Tpm3b* expression volume

was significantly larger than in the generalists (Table S3; P < 0.05; 1-way ANOVA, Tukey's

HSD test).

Figure 3. Spatiotemporal patterns of *galr2a* expression differ among SSI species at 2 dpf.

Representative images of *Galr2a* (green) and *Tpm3b* (magenta) expression revealed by hybridization

chain reaction (HCR) and DAPI nuclear staining (gray) in the: **a.** molluscivore, *C. brontotheroides*; **b.**

generalist, *C. variegatus*; and **c.** scale-eater, *C. desquamator*. **d.** *Galr2a* reads per kilobase of transcript

per million reads (RPKM) from an existing mRNAseq study sampling the entire head of each species at

four developmental stages (Lencer and McCune, 2020). Galr2a showed significantly higher expression in

the molluscivore relative to generalists (P = 0.003, Tukey's HSD test) and scale-eaters (P = 0.014,

Tukey's HSD test) at 2 post fertilization (dpf). **e.** Mean fluorescence per area for *galr2a* quantified at 2 dpf

in all three species shows similarly elevated levels of expression in the molluscivore for the mandibular

and pharyngeal arches. Images shown are representative Z-stack maximum projection images of 30

optical sections taken every 3 µm from whole-mounted embryos imaged using a Zeiss LSM880 laser

confocal microscope. PA: pharyngeal arches, PE: pharyngeal endoderm, ManM: mandibular

mesenchyme, MLF: medial longitudinal fasciculus, PTe: posterior tectum, Di: diencephalon; Mb: midbrain,

Hb: hindbrain, Y: yolk, Myo: somitic myofibers, IO: inferior oblique muscle.

Figure 4. Spatiotemporal patterns of *galr2a* expression differ in molluscivores at 8 dpf.
Representative images of *Galr2a* (green) and *Tpm3b* (magenta) expression revealed by hybridization chain reaction (HCR) and DAPI nuclear staining (gray). **a.** Lateral view of an 8 dpf molluscivore (M in white circle), **a'.** ventral view. **b.** Lateral view of an 8 dpf generalist (G in white circle), **b'.** ventral view. **c.** Lateral view of an 8 dpf scale-eater (S in white circle), **c'.** ventral view. The arrowhead in **b.** and the arrow in **c.** point to high expression of *galr2a* in the maxilla and premaxilla of the molluscivore and scale-eater specialists, respectively. White arrowheads in **a'.** point to high expression of *galr2a* in the palatoquadrate

of the molluscivore. *Galr2a* expression in the IMA was found only in the scale-eaters (**c'**). Images shown

here are representative Z-stack maximum projection images of 30 optical sections taken every 15 µm

from whole-mounted larvae imaged using a Zeiss LSM880 laser confocal microscope. **d-d'.** 3D-Slicer

view of the expression volume of *galr2a* (**d.**) and *tpm3b* (**d'.**) in an 8 dpf molluscivore pupfish larvae (*C.*

brontotheroides). **e.** Volume of *galr2a* expression in the brain (left panel), whole head (middle panel), and

jaw (right panel) across SSI pupfishes. CNS: central nervous system, Max: maxilla. LEVs: levator arcus

palatini and operculi, AdM: adductor mandibular. IMA: intermandibular anterior. Total volume rendered:

450 µm. All scale bars: 100 µm.

Figure 5. *Galr2a* is expressed in the chondrocytes of the Meckel's and palatoquadrate cartilages at

hatching (8 dpf) in SSI pupfishes. HCRs showing tissue expression of *Galr2a* in green, and the muscle

marker Tropomyosin 3b, (*Tpm3b*), in magenta. Nuclei are stained with DAPI (gray). **a-a'.** Scale-eater

galr2a expression in the chondrocytes of the Meckel's symphysis (Sym) and lateral side. **b.** Ventral view

of a molluscivore jaw at 8 dpf showing the expression of *Tpm3b* in the intermandibular muscles of the

lower jaw. *Galr2a* is expressed in the jaw joint and the most anterior region of the lower jaw. **c-d.** *Galr2a*

is expressed in the chondrocytes of the Meckel's and palatoquadrate cartilages and around the jaw joint.

e. *Galr2a* expression in the premaxilla of a scale-eater. Unlabeled scale-bars: 10 μm.

Chemical inhibition of Galr2 receptors affects Meckel's cartilage length and chondrocyte density

To study the effect of Galr2 in craniofacial development, we inhibited the endogenous activation of all four known galanin receptors in teleost fishes (*Galr1a* and b, *Galr2a* and b; Butler et al. 2020, Li et al. 2013) using M35 (Innopep, Inc.), a synthetic peptide antagonist of Galr1+2 galanin receptors (Wiesenfeld-Hallin et al., 1993), and M871 (Abcam), a Galr2-specific synthetic peptide antagonist (Sollenberg et al., 2006). Embryos of all three species from two different lake populations were exposed from stages 24-25 (2 dpf, with the appearance of the first pharyngeal arches) until hatching at stages 32-33 (8 dpf, Fig. 6a).

At approximate hatching time (8 dpf), both trophic specialist species raised under laboratory common garden conditions exhibited increased Meckel's cartilage length (Fig. 6d), consistent with the longer jaws of adult scale-eaters and more robust jaws of the molluscivore relative to more gracile jaws of the generalist (Martin and Wainwright 2013). We found that exposure to M871, the Galr2-specific antagonist, significantly reduced the length of the Meckel's cartilage in both specialists relative to the generalists (Fig. 6d.; $P = 7.89 \times 10^{-5}$ for scale-eaters; $P = 0.001$ for molluscivores; 2-way ANOVA, Tukey's HSD test) while the interocular distance remained unchanged between control and treated larvae (Fig. S4). In contrast, M35 (Galr1 and Galr2 antagonist) only significantly reduced Meckel's length in the molluscivore (Fig. 6c). Meckel's length of generalists was unaffected by exposure to M35 or M871 (Fig. 6d).

a.

GalR2

M35 and M871 incubation

Stage 24-25   Stage 32-33

b.

MC

IOD

c.

d.

Molluscivore   Generalist   Scale-eater

log Meckel's cartilage length

**   ***
                   **

Control M35 M871   Control M35 M871   Control M35 M871

Treatments

e.

Molluscivore   Generalist   Scale-eater

Chondrocyte Density

*
***
*

Control M35 M871   Control M35 M871   Control M35 M871

Treatments

Treatments
● Control
▲ M35
■ M871

Species
Molluscivore
Generalist
Scale-eater

f.

control   M35   M871

M   M   M

control   M35   M871

G   G   G

control   M35   M871

S   S   S

Figure 6. Galr2 receptor inhibition reduced Meckel's cartilage length in both specialists and

increased chondrocyte density in scale-eaters. a. Galr2 and Galr1+2 inhibition protocol during pupfish

development. *C. desquamator* embryos are depicted. **b.** Alcian blue cartilage staining at Stage 32-33.

IOD: interocular distance, MC: Meckel's cartilage. The MC length and IOD are shown in red. **c.**

Chondrocytes at the symphysis. **d.** Changes in the log-transformed Meckel's length and the log-

transformed interocular distance at stage 32-33 across species and treatments. **e.** Chondrocyte numbers

at the symphysis across species and treatments. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ Tukey's HSD test. **f.**

Representative control and treated larvae of each species at stage 32-33. Arrows point to abnormal lower

jaws. M: molluscivores, G: generalist, S: scale-eaters. Scale-bars = 0.4 mm. Galr1 and Galr2 cartoons

were modified from Lang et al., 2007.

To understand the cellular effect of Galr1 and 2 antagonists on Meckel's length across

species, we quantified the number of chondrocytes 100 µm from the symphysis and the mean

width of ten chondrocytes nearest to the symphysis. We found no significant differences in the

mean chondrocyte width among pupfish species but observed increased chondrocyte density in

untreated scale-eaters relative to generalists (Fig. 6e; Table S4; $P = 0.009$, ANOVA, Tukey's

HSD test). Moreover, only scale-eaters responded to M871, but not M35, by further increasing

chondrocyte density relative to the control larvae (Fig. 6e; Table S4; $P = 0.03$, ANOVA, Tukey's

HSD test; mean ± 1 SE; control = 28.22 ± 0.38; M871 = 32.06 ± 1.06). Despite having shorter

jaws after treatment with M871, chondrocyte density was significantly increased (Fig. 6e; mean

± 1 SE; control = 28.22 ± 0.38; M871 = 32.06 ± 1.06).

Discussion

We used an evolutionary radiation of trophic specialist pupfishes, endemic to the Bahamas, to

discover a novel function for *galr2a* in craniofacial divergence. Specifically, we confirmed that

two transcription factor binding sites upstream of *galr2a* display highly divergent allele

frequencies between trophic specialist species, visualized *galr2a* expression in craniofacial

tissues in all three SSI pupfish species at two developmental stages and demonstrated a

phenotypic effect on Meckel's cartilage length and chondrocyte density using synthetic peptides

to inhibit the activity of Galr2 and Galr1+2. Our findings demonstrate a crucial role for Galr2 in

craniofacial divergence within this pupfish radiation. Our study also provides a roadmap in a

non-model vertebrate system for rapidly identifying previously uncharacterized candidate genes

important for adaptation to novel ecological niches (e.g., trophic specialization) that can be

quickly validated through classic and state-of-the-art developmental biology tools. Overall, our

research contributes to understanding the genetic basis of phenotypic evolution and adaptation in

non-model organisms.

Putative loss of a transcription factor binding site for *galr2a* in scale-eating pupfish

One of the most common evolutionary changes associated with phenotypic changes among

closely related species is the gain or loss of cis-regulatory elements (Mack and Nachman, 2017;

Signor and Nuzhdin, 2018). More than 85% of scale-eaters carry two transversions in the

regulatory region of *galr2a* (Fig. 2). Combined with our observations of reduced *galr2a*

expression in the mandibular mesenchyme during early jaw development in this species (Fig. 3),

we conclude that the putative loss of a predicted *Sry* transcription factor binding site in scale-

eaters is the most likely explanation for changes in gene expression, rather than a gain of a new

predicted TFBS for *Znf416* at this locus (Fig. 2). This is further supported by the critical role of

the *Sry*-related HMG box (Sox) family of transcription factors (especially the SoxE group

including *Sox8, Sox9,* and *Sox10*) in craniofacial development as *Sry* transcription factors specify

the behavior, multipotency, and survival of neural crest cells during vertebrate development

(Leathers and Rogers, 2022; Simões-Costa and Bronner, 2015; Guo et al., 2015; Haldin and

LaBonne, 2010; Betancur et al., 2010; Kamachi and Kondoh 2013).

Alternatively, we cannot rule out a gain of a TFBS upstream of *galr2a* in scale-eaters,

additive or epistatic effects of both upstream transversions or more complex regulatory

architectures, such as many interacting functional cis- and trans-acting regulatory variants or

combinations of variants segregating at lower frequencies in trophic specialists that we have not

prioritized (Boyle et al., 2017; McGirr and Martin, 2021; Price et al., 2015). However, our

mapping cross of a single outbred pair of trophic specialists indicates that a single moderate-

effect QTL containing *galr2a* explains 15% of phenotypic variation in oral jaw length between

these species, consistent with causative variants affecting jaw size originating from this region

(Martin et al. 2017).

Differential *Galr2a* expression during development is associated with craniofacial

divergence in SSI pupfishes

We found *galr2a* expression in our in situ hybridization experiments to be consistent with

previously published RNAseq data for craniofacial tissue in this radiation (Fig. 3; Lencer and

McCune, 2017; McGirr and Martin, 2018; McGirr and Martin, 2019; McGirr and Martin, 2021),

with *galr2a* being differentially expressed only at 2 dpf. We observed distinctive spatial *galr2a*

expression among SSI species at 2 and 8 dpf suggesting important time and tissue-specific

regulation of *galr2a* expression during pupfish development. At 2 dpf, we found a strong

association between decreased expression of *galr2a* in the mandibular mesenchyme anterior to

the first pharyngeal arch with the future Meckel's cartilage and oral jaw lengths in the adults of

each species; with increasing *galr2* abundance in the molluscivores associated with shorter, but

more robust jaws in adults (Martin and Wainwright 2013). In contrast, the reduced *galr2a*

expression in the mandibular mesenchyme is associated with the development of longer oral

jaws in scale-eaters, which is apparent as early as hatching time (Fig. 6d; Lencer et al., 2016).

We noted expression of *galr2a* in the maxilla of only molluscivores at hatching (Fig. 4 a-

c), absent in the scale-eaters and generalists, consistent with the uniquely enlarged and anteriorly

protruding head of the maxilla in this species (Martin and Wainwright 2013; Hernandez et al.

2018; Martin and Feinstein 2014; St. John et al. 2020). Furthermore, *galr2a* expression in the

intermandibular muscles of the scale-eaters at hatching time (Fig. 4c) suggests that *galr2a*

expression can also modulate the development of the observed hypertrophic musculature of the

adductor mandibulae in the adult scale-eating pupfish (Hernandez et al., 2018). Altogether, these

interspecific differences in spatial expression support a novel role for *galr2a* in musculoskeletal

development and may contribute to the divergent craniofacial morphology observed in SSI

pupfishes.

Receptor inhibition supports a novel function for GALR2 in craniofacial divergence of SSI

pupfishes

Interestingly, the response of Meckel's cartilage length and chondrocyte density to the inhibition

of Galr-receptor pathways was highly dependent on the species' genetic background. Scale-

eaters responded only to the Galr2-specific antagonist M871 with substantially reduced Meckel's

cartilage length and increased chondrocyte density but were unaffected by the Galr1+2

antagonist M35 (Fig. 6c). Importantly, M871 exhibits a higher binding constant for Galr2

receptors than the Galr1+2 antagonist M35 (Sollenberg et al., 2006; Sollenberg et al., 2010).

Previous transcriptomic data from craniofacial tissues during early development (Lencer and

McCune 2017) also indicates that *galanin* and *Galr1* mRNA transcript abundance does not vary

among the three SSI species (Table S3).

Therefore, we conclude that even though scale-eaters expressed the lowest amount of

galr2a transcripts and presumably contain overall fewer Galr2 receptors in their craniofacial

tissue, M871's higher binding affinity (Lang et al., 2007) was sufficient to inhibit these fewer

receptors, resulting in reduced Meckel's cartilage length (Fig. 7). In contrast, the presumably

greater concentration of Galr2 receptors in molluscivore craniofacial tissue due to increased

Galr2 expression during development resulted in the inhibition of these receptors by both M871

and M35, despite its lower binding affinity for Galr2 (Lang et al., 2007). We also conclude that

inhibition of Galr1 receptors by M35 does not affect Meckel's cartilage length or chondrocyte

density and that the specific inhibition of Galr2 by M871 was sufficient to drive equal Meckel's

cartilage length shortening. Finally, the generalists' lack of significant response to M35 is

consistent with their decreased levels of *galr2a* at 2 dpf, suggesting that, as in scale-eaters treated

with M35, fewer Galr2 receptors may be present for inhibition in these species' background.

However, the response to M871 appears to be specialist-specific, with little effect on the

generalist Meckel's cartilage length, and thus, we cannot rule out more complex species-specific

factors interacting with Galr regulatory pathways.

Our work is consistent with previous work showing that *Galr2* activation inhibits cell

proliferation in neuronal cell lines (Lang et al., 2007; Berger et al., 2004), suggesting that the

TFBS allele frequency shifts observed in scale-eaters are the cause for low *galr2a* abundance

and, as a consequence, have low Galr2 activity during craniofacial development resulting in

increased chondrocyte density and Meckel's cartilage length. Thus, the increased chondrocyte

density observed in the M871-treated scale-eater larvae result likely from the strong and

continued inhibition of the fewer Galr2 receptors available during development as a consequence

or a synergy of/with their genetic background.

In conclusion, we propose that the inhibition of Galr2 and not Galr1 induces the

reduction of Meckel's cartilage length. Thus, a greater number of Galr2 receptors in the

molluscivore specialist, as a result of increased expression of *galr2a* during early development,

may result in their reduced oral jaw lengths as adults through increased opportunities for

endogenous agonistic binding interactions (Fig. 7). Fewer Galr2 receptors on the scale-eater jaw

may result in their enlarged jaw lengths as adults by limiting opportunities for Galr2 endogenous

agonists to bind to these receptors during development. In the generalists, a greater number of

Galr2 receptors but smaller overall Meckel's cartilage length may limit the sensitivity of this

species to Galr2 and Galr1+2 endogenous agonists, however, resulting in their intermediate

length and least robust oral jaws among the three species under control conditions (Hernandez et

al. 2018; Martin and Wainwright 2011).

Meckel's cartilage length

Figure 7. Proposed mechanism for how galanin receptor abundance affects Meckel's cartilage length in the two pupfish trophic specialists after exposure to M35 (weak Galr1+Galr2 antagonist) or M871 (strong Galr2 antagonist). Molluscivores (top row) express more *galr2a* at 2 and 8 dpf than scale-eaters (bottom row) in their Meckel's cartilage (see Figs. 3-4), resulting in more Galr2 receptors (open circles). Increased Galr2 receptor density in molluscivores results in their reduced Meckel's cartilage length (see Fig. 6) due to inhibition by both the weak antagonist (M35: middle column; half-filled circles) and strong antagonist (M871: right column; filled circles) of Galr2. In contrast, reduced Galr2

receptor density in scale-eaters (bottom row) results in their reduced Meckel's cartilage length only after

exposure to the strong Galr2-specific antagonist M871.

Conclusion

Our results support a novel role of the second receptor for galanin, Galr2, as a craniofacial

modulator gene important in controlling craniofacial development and interspecific divergence

through modifying its transcript abundance and receptor activity. *Galr2a* transcript abundance

changes between species are associated with genetic changes in the regulatory region of *galr2a*,

consistent with the loss of a transcription factor binding site in the scale-eating pupfish. We

propose a model in which reduced Galr2 receptor density in the oral jaws of scale-eaters results

in fewer endogenous agonistic interactions, increasing Meckel's cartilage length presumably by

decreased inhibition of chondrocyte proliferation (as a downstream effect of endogenous Galr2

activation). We also acknowledge the polygenic nature of jaw development and note that our

previous genetic mapping experiments place an upper bound of 15% of oral jaw variation that

may be due to differences in *galr2a* regulation among the myriad craniofacial genes driving the

evolution of divergent craniofacial traits in this system.

Materials and Methods

Pupfish husbandry and maintenance

Three sympatric *Cyprinodon* species (*C. variegatus, C. brontotheroides,* and *C. desquamator*)

were collected from both Osprey Lake and Crescent Pond in San Salvador Island, Bahamas in

2018 and maintained in the lab for at least two generations in small breeding groups. All lab

colonies were maintained at a pH of 8.0 and 2-5 ppt salinity at 25-30°C on a 12:12 hours

light:dark cycle at the University of California, Berkeley. Newly hatched fry were raised on brine

shrimp nauplii for the first month and adults and juveniles were fed a combination of commercial

pellet food and frozen bloodworms. All protocols and procedures employed were reviewed and

approved by the University of California, Berkeley Animal Care and Use Committee (AUP-

2015-01-7053) and animals were collected and exported from the Bahamas with research and

export permits from the Bahamas Environmental, Science, and Technology commission through

the Gerace Research Centre.

Multiple breeding pairs for each population were housed together in 40-liter aquaria with

synthetic yarn spawning mops and fed ad libitum daily with frozen foods and commercial

pellets. Embryos were collected by handpicking and raised at 27°C in 5 ppt salinity water with

the addition of 10 mg/l gentamycin and 0.0001% methylene blue. Fertilization was confirmed

after collection and embryos were staged according to Lencer and McCune (2018). The day of

collection was noted as day "zero", and thus one day after collection was 1-day post-fertilization

(dpf).

HCR FISH staining

Hybridization chain reaction (HCR) was used to visualize spatial expression of *galr2a* and *tpm3b*

at key development timepoints (Choi et al., 2018; Ibarra-García-Padilla et al., 2021). HCR

reagents, including probes, hairpins, and buffers, were purchased directly from Molecular

Instruments (Los Angeles, California, USA). Staining was performed according to a modified

protocol from Ibarra-García-Padilla et al., 2021, and Palominos and Whitlock, 2021.

Fixation and storage: 2 and 8 dpf larvae were fixed in 4% paraformaldehyde (Electron

Microscopy Science) 1X PBS (Invitrogen) overnight at 4°C. Samples were then serially

dehydrated in methanol/1X PBS (25%, 50%, 75%) and stored in 100% methanol at -20°C for at

least 16 hours before use. *Permeabilization:* Rehydration was performed serially in

methanol/PBST (1X PBS 0.1% Tween20, Fisher Scientific) up to 100% PBST. Samples were

then incubated for 10 minutes in pre-chilled acetone at -20°C for permeabilization. 2 and 8 dpf

samples were then incubated for 20 and 80 minutes in 10 µg/ml Proteinase K (Thermofisher),

respectively. Prehybridization: Samples were then pre-hybridized in the pre-warmed (at 37°C)

hybridization buffer for 30 minutes at 37°C. *Hybridization:* For *Galr2a* and *Tpm3,* 4 to 6 pmol

and 2 to 3 pmol were used, respectively. The probe mixture was pre-warmed for 30 minutes at

37°C before hybridization for 20 hours at 37°C. *Washing probe excess:* After hybridization, we

performed four washes in pre-warmed wash buffer (15 minutes each) at 37°C, followed by two

final washes in 5X SSCT (20X SSC from Thermofisher; SSCT is SSC with 0.1% Tween20) for

5 minutes each. *Amplification:* Preamplification involved incubating samples in pre-equilibrated

(20-25°C) amplification buffer for 30 minutes. Next, amplification was performed using 4 µl of

each snap-cooled hairpin in 250 µl of amplification buffer for 16 hours in the dark at room

temperature. *Galr2a* expression was confirmed independently using two amplifiers with different

excitation/emission wavelengths: B1-488 HCR amplifiers, and B1-594 HCR amplifiers. *Tpm3b*

amplifiers used were B2-647 HCR amplifiers. *Washing hairpin excess:* Excess hairpins were

washed several times with 5X SSCT, followed by a last wash with PBST. *Nuclei labeling:* After

rinsing, samples were incubated for 30 minutes in DAPI/1X PBST (1 µm/ml, Sigma) and then

washed in PBST 3 times for 5 minutes each. *Clearing and mounting:* Samples were then serially

cleared in glycerol solutions at room temperature by sequentially removing and adding: 25%

glycerol/75% PBST, 50% glycerol/50% PBST, 75% glycerol/25% PBST. Samples were then

mounted in a custom-made bridged microscope slide (Fisher Scientific) in 75% glycerol/25%

PBST.

Imaging: Samples were imaged using a Zeiss LSM880 (inverted) confocal microscope at the

CNR Biological Imaging Facility at the University of California, Berkeley. All HCR were

volumetric z-stack images of embryos or larvae positioned laterally or ventrally and taken every

3 µm. Images were deconvoluted, analyzed, and edited in FIJI (Schindelin et al., 2012) and

Imaris (Bitplane).

Galr2 inhibition during jaw development

According to the pupfish developmental series of Lencer and McCune (2018), the pharyngeal

arches of *C. variegatus* are visible as segmented pouches ventral to the developing head at Stage

25 (48-64 hours post-fertilization at 27 °C, or 2 – 2.5 dpf), while the presence of a stomodeum-

like oral cavity is evident by Stage 30 (100 – 130 hpf, or 4 – 5.5 dpf). We hypothesized that the

key stages for proper jaw development occur between Stages 24 and 25, and thus we

dechorionated Stage 24-25 embryos (or 2 to 2.5 dpf) pupfish by incubating them on 20 mg/ml

Pronase (Roche) in water for 2.5 hours. The remaining chorions were removed manually with

forceps. After larvae were dechorionated, we incubated them in 100 µM M35 (InnoPep Inc, San

Diego, CA), a chimeric peptide antagonist to all galanin receptors (Galr1 and Galr2 in teleost

fishes; Butler et al. 2020) and with a conserved 21 aminoacidic sequence to the core Galanin

peptide sequence (GWTLNSAGYLLGPPPGFSPFR-NH2), or M871 (Abcam, USA), a Galr2-

specific   antagonist   with   the   following   aminoacidic   sequence:

WTLNSAGYLLGPEHPPPALALA-NH2, up to hatching time (Stage 33), looking for the

following developmental events that are present in Stg 33 but not Stg 32: partial yolk absorption,

frequent mouth movements with visible buccal pumping, frequent pectoral fin movements, and

the ability to propel when touched. We also confirmed the formation of the ethmoid plate in both

control and treated embryos after cartilage staining.

Cartilage staining: a non-acid clearing and staining method was used based on Walker and

Kimmel, 2007, and Lencer and McCune, 2018. Fish larvae were cleared in 80% glycerol and

mounted in custom-made coverslips (Fisher Scientific). Meckel's length, inter-ocular distance,

and chondrocyte density were quantified using the segmented line tool and ROI manager in Fiji

(Schindelin et al., 2012). Inter-ocular distance was done by drawing a line from the center of one

lens to the other and measuring the distance between the eyes (from the edge of the retinal

pigmented epithelium). Imaging was done using a Zeiss Z1 AxioImager M2 microscope located

at the Biological Imagining Facility of the University of California, Berkeley.

Sequencing of *Galr2a* regulatory alleles

We used the DNeasy Blood and Tissue kit (Qiagen, Inc.) to extract DNA from the fins of wild-

caught adult fish and quantified it on a Qubit 3.0 fluorometer (Thermofisher Scientific, Inc.). All

specimens used in this study are kept in the Museum of Vertebrate Zoology at the University of

California, Berkeley. The primers used to amplify both *Galr2a* regulatory loci are listed in Table

S5. The sizes of the PCR products were confirmed by gel electrophoresis. PCR product clean-up

was done using ExoSap-IT (USB) and quantified on a Nanodrop ND-1000 (Thermo Scientific)

spectrophotometer. Sanger sequencing was performed at the UC Berkeley DNA Sequencing

Facility. Sequence alignment was performed using Geneious Prime (Dotmatics).

Prediction of transcription factors binding sites in *Galr2a-15kb* and *Galr2a-12kb*

We used the Multiple Expectation maximizations for Motif Elicitation (MEME) server (http://meme-suite.org/) and the motif database scanning algorithm TOMTOM (Bailey et al., 2009) to retrieve TFBS motifs from a 20kb conserved generalist, molluscivore, and scale-eater sequences upstream *galr2a* obtained from Richards et al., 2021. MEME suite finds, in the given sequences, the most statistically significant motifs first. We numbered the motifs following their statistical significance (Table S1).

Statistics

Expression quantification of galr2a and tpm3b at 8 dpf: We segmented and quantified "jaw" *galr2a* expression in the maxilla, and the Meckel's and palatoquadrate cartilages and "brain" *galr2a* expression in the central nervous system from HCR confocal images using 3D-Slicer (Fedorov et al., 2012). *Tpm3b* expression was segmented and quantified in the intermandibular muscles (IMA), the protractor hyoideus (PR-H), the adductor mandibular complex (AddM), and the levator arcus palatini (LEVs). Muscles were named according to Diogo et al., 2008. Statistical tests comparing the expression volume among species and tissues were performed in R (R Core Team 2023). We tested for significant differences in means via ANOVA and contrasted groups post-hoc using Tukey's HSD test.

Expression quantification of garlr2a at 2 dpf: From the HCR confocal images, a total volume of 45 µm containing the pharyngeal arches was projected using the maximum intensity tool in Fiji. We selected two different tissues using ROI manager and quantified the relative fluorescence normalized by the background fluorescence of the image. First, we selected the mandibular

mesenchyme plus the first and second pharyngeal arches (labeled in the figure "Pharyngeal

Arches"). Then, we de-selected the second pharyngeal arch, quantifying only the expression of

galr2 in the mandibular mesenchyme and the first pharyngeal arch (labeled in the figure

"Mandibular arch"). We then tested for significant differences in means via ANOVA and post-

hoc analysis with Tukey's HSD in R.

mRNA transcript analysis: A table containing RPKM values of transcripts from Lencer and

McCune, 2017, datasets was obtained from the following Dryad repository,

doi:10.5061/dryad.mb3gb. Genes of interest were found with the Gene ID listed on NCBI. Gene

IDs are as follows: 106090243, 107092788, 107090827, 107096257, 107098818, and

107092130, for *galr2a, galr2b, galr1a, galr1b, gal,* and *sox21b*, respectively. Statistical tests

comparing transcripts RPKM expression between species and developmental stages were

performed in R, testing for significant differences in means via ANOVA and post-hoc analysis

with Tukey's HSD.

Cartilage staining: We compared chondrocyte density for significant differences in means using

pairwise post-hoc Tukey's HSD tests. We used a generalized linear model accounting for species

and treatment and their interaction. We chose to measure chondrocytes within the Meckel's

cartilage because it allows for a clear view of each cell at the symphysis.

Author's contributions

Conceptualization: MFP, CTM, CHM; Data Collection: MFP, VM; Statistical analyses: MFP;

Resources: CHM; Microscopy: MFP, VM; Original draft: MFP; Revising: MFP, EJR, CTM,

CHM

Acknowledgments

We thank Michelle St. John, David Tian, Mara Van Tassell, Chloe Clair, Dylan Chau, and Heidi

Buratti for valuable comments and discussion of the results. We thank Lydia Smith for her help

throughout the development of this project in the Evolutionary Genetics Lab at the University of

California, Berkeley. We are particularly thankful to Austin H. Patton, who extracted the *C.*

brontotheroides mRNA reads to create the HCR probes tested here, and for his insightful ideas

and comments on the early stages of this manuscript. We thank the Gerace Research Centre and

Troy Day for logistical support and the government of the Bahamas for permission to collect and

export samples. This research was funded by the National Science Foundation DEB CAREER

grant #1749764, National Institutes of Health grant 5R01DE027052-02, the University of North

Carolina at Chapel Hill, and the University of California, Berkeley to CHM.

References

Albertson, R. C., Cresko, W., Detrich, H. W. and Postlethwait, J. H. (2009). Evolutionary

mutant models for human disease. *Trends in Genetics* **25**, 74–81.

Bonfante, B., Faux, P., Navarro, N., Mendoza-Revilla, J., Dubied, M., Montillot, C.,

Wentworth, E., Poloni, L., Varón-González, C., Jones, P., et al. (2021). A GWAS in

Latin Americans identifies novel face shape loci, implicating VPS13B and a Denisovan

introgressed region in facial variation. *Sci Adv* **7**,.

Boyle, E. A., Li, Y. I. and Pritchard, J. K. (2017). An Expanded View of Complex Traits:
From Polygenic to Omnigenic. *Cell* **169**, 1177–1186.

Brawand, D., Wagner, C. E., Li, Y. I., Malinsky, M., Keller, I., Fan, S., Simakov, O., Ng, A.
Y., Lim, Z. W., Bezault, E., et al. (2014). The genomic substrate for adaptive radiation
in African cichlid fish. *Nature* **513**, 375–381.

Buffenstein, R. (2008). Negligible senescence in the longest living rodent, the naked mole-rat:
Insights from a successfully aging species.

Carlborg, O. and Haley, C. S. (2004). Epistasis: too often neglected in complex trait studies?
Nature reviews. Genetics **5**, 618–625.

Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Shapiro, M. D., Brady, S. D.,
Southwick, A. M., Absher, D. M., Grimwood, J., Schmutz, J., et al. (2010). Adaptive
evolution of pelvic reduction in sticklebacks by recurrent deletion of a Pitx1 enhancer.
Science (New York, N.Y.) **327**, 302–305.

Concannon, M. R. and Albertson, R. C. (2015). The genetic and developmental basis of an
exaggerated craniofacial trait in East African cichlids. *J. Exp. Zool. B Mol. Dev. Evol.*
324, 662–670.

Conith, A. J., Lam, D. T. and Albertson, R. C. (2019). Muscle-induced loading as an
important source of variation in craniofacial skeletal shape. *Genesis* **57**, e23263.

Erickson, P. A., Glazer, A. M., Cleves, P. A., Smith, A. S. and Miller, C. T. (2014). Two
developmentally temporal quantitative trait loci underlie convergent evolution of

increased branchial bone length in sticklebacks. *Proceedings of the Royal Society B:*

Biological Sciences **281**,.

Erickson, P. A., Ellis, N. A. and Miller, C. T. (2016). Microinjection for transgenesis and

genome editing in threespine sticklebacks. *J. Vis. Exp.* 10.3791/54055.

Glazer, A. M., Killingbeck, E. E., Mitros, T., Rokhsar, D. S. and Miller, C. T. (2015).

Genome assembly improvement and mapping convergently evolved skeletal traits in

sticklebacks with genotyping-by-sequencing. *Genes,Genomes,Genetics* **5**, 1463–1472.

Glazier, A. M. (2002). Finding genes that underlie complex traits. *Science (New York, N.Y.)* **298**,

2345–2349.

Gross, J. B., Stahl, B. a., Powers, A. K. and Carlson, B. M. (2016). Natural bone

fragmentation in the blind cave-dwelling fish, *Astyanax mexicanus*: Candidate gene

identification through integrative comparative genomics. *Evolution and Development* **18**,

7–18.

Hall, D., Tegstrom, C. and Ingvarsson, P. K. (2010). Using association mapping to dissect the

genetic basis of complex traits in plants. *Briefings in functional genomics* **9**, 157–165.

Hawes, J. J., Narasimhaiah, R. and Picciotto, M. R. (2006). Galanin and galanin-like peptide

modulate neurite outgrowth via protein kinase C-mediated activation of extracellular

signal-related kinase. *Eur. J. Neurosci.* **23**, 2937–2946.

Helman, S. N., Badhey, A., Kadakia, S. and Myers, E. (2014). Revisiting Crouzon syndrome:

reviewing the background and management of a multifaceted disease. *Oral and*

Maxillofacial Surgery **18**, 373–379.

Hernandez, L. P., Adriaens, D., Martin, C. H., Wainwright, P. C., Masschaele, B. and

Dierick, M. (2018). Building trophic specializations that result in substantial niche

partitioning within a young adaptive radiation. *J. Anat.* **232**, 173–185.

Hirschhorn, J. N. and Daly, M. J. (2005). Genome-wide association studies for common

diseases and complex traits. *Nature reviews* **6**, 95–108.

Hochheiser, H., Aronow, B. J., Artinger, K., Beaty, T. H., Brinkley, J. F., Chai, Y.,

Clouthier, D., Cunningham, M. L., Dixon, M., Donahue, L. R., et al. (2011). The

FaceBase Consortium: A comprehensive program to facilitate craniofacial research.

Developmental Biology **355**, 175–182.

Hofmann, C. M., O'Quin, K. E., Justin Marshall, N., Cronin, T. W., Seehausen, O. and

Carleton, K. L. (2009). The eyes have it: Regulatory and structural changes both

underlie cichlid visual pigment diversity. *PLoS Biol.* **7**,.

Humphries, J. M. and Miller, R. R. (1981). A remarkable species flock of pupfishes, genus

Cyprinodon, from Yucatan, Mexico. *Copeia* **1981**, 52–64.

J. Parsons, K., McWhinnie, K. and Armstrong, T. (2021). An Evo-devo View of Post-

genomic African Cichlid Biology: Enhanced Models for Evolution and Biomedicine. In

The Behavior, Ecology and Evolution of Cichlid Fishes (ed. Abate, M. E.) and Noakes,

D. L. G.), pp. 779–802. Dordrecht: Springer Netherlands.

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., Swofford,
R., Pirun, M., Zody, M. C., White, S., et al. (2012). The genomic basis of adaptive
evolution in threespine sticklebacks. *Nature* **484**, 55–61.

Kadakia, S., Helman, S. N., Badhey, A. K., Saman, M. and Ducic, Y. (2014). Treacher
Collins Syndrome: The genetics of a craniofacial disease.

Kocher, T. D. (2004). Adaptive evolution and explosive speciation: The cichlid fish model. *Nat.*
Rev. Genet. **5**, 288–298.

Kocher, T. D., Albertson, R. C., Carleton, K. L. and Streelman, J. T. (2003). The genetic
basis of biodiversity, genomic studies of cichlid fishes. *Aquatic genomics: steps toward a*
great future. 34–44.

Kolora, S. R. R., Owens, G. L., Vazquez, J. M., Stubbs, A., Chatla, K., Jainese, C., Seeto,
K., McCrea, M., Sandel, M. W., Vianna, J. A., et al. (2021). Origins and evolution of
extreme life span in Pacific Ocean rockfishes. *Science* **374**, 842–847.

Lencer, E. S., Riccio, M. L. and McCune, A. R. (2016). Changes in growth rates of oral jaw
elements produce evolutionary novelty in bahamian pupfish. *J. Morphol.* **277**, 935–947.

Lencer, E. S., Warren, W. C., Harrison, R. and McCune, A. R. (2017). The Cyprinodon
variegatus genome reveals gene expression changes underlying differences in skull
morphology among closely related species. *BMC Genomics* **18**, 424.

Li, L., Wei, S., Huang, Q., Feng, D., Zhang, S. and Liu, Z. (2013). A novel galanin receptor

1a gene in zebrafish: tissue distribution, developmental expression roles in nutrition

regulation. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **164**, 159–167.

Liu, S., Lorenzen, E. D., Fumagalli, M., Li, B., Harris, K., Xiong, Z., Zhou, L.,

Korneliussen, T. S., Somel, M., Babbitt, C., et al. (2014). Population genomics reveal

recent speciation and rapid evolutionary adaptation in polar bears. *Cell* **157**, 785–794.

Ma, A., Bai, J., He, M. and Wong, A. O. L. (2018). Spexin as a neuroendocrine signal with

emerging functions. *Gen. Comp. Endocrinol.* **265**, 90–96.

Mack, K. L. and Nachman, M. W. (2017). Gene Regulation and Speciation. *Trends Genet.* **33**,

68–80.

Man, P.-S. and Lawrence, C. B. (2008). The effects of galanin-like peptide on energy balance,

body temperature and brain activity in the mouse and rat are independent of the

GALR2/3 receptor. *J. Neuroendocrinol.* **20**, 128–137.

Manolio, T. A., Collins, F. S., Cox, N. J., Goldstein, D. B., Hindorff, L. A., Hunter, D. J.,

McCarthy, M. I., Ramos, E. M., Cardon, L. R., Chakravarti, A., et al. (2009).

Finding the missing heritability of complex diseases. *Nature* **461**, 747–753.

Martin, C. H. (2012). Weak disruptive selection and incomplete phenotypic divergence in two

classic examples of sympatric speciation: Cameroon crater lake cichlids. *Am. Nat.* **180**,

E90–E109.

Martin, C. H. (2013). Strong assortative mating by diet, color, size, and morphology but limited

progress toward sympatric speciation in a classic example: Cameroon crater lake cichlids.

Evolution **67**, 2114–2123.

Martin, C. H. (2016). The cryptic origins of evolutionary novelty: 1000-fold faster trophic

diversification rates without increased ecological opportunity or hybrid swarm.

Evolution.

Martin, C. H. and Genner, M. J. (2009). High niche overlap between two successfully

coexisting pairs of Lake Malawi cichlid fishes. *Canadian Journal of Fisheries and*

Aquatic Sciences **66**, 579–588.

Martin, C. H. and Gould, K. J. (2020). Surprising spatiotemporal stability of a multi-peak

fitness landscape revealed by independent field experiments measuring hybrid fitness.

Evol Lett **4**, 530–544.

Martin, C. H. and Wainwright, P. C. (2011). Trophic novelty is linked to exceptional rates of

morphological diversification in two adaptive radiations of *Cyprinodon* pupfishes.

Evolution **65**, 2197–2212.

Martin, C. H. and Wainwright, P. C. (2013a). A remarkable species flock of *Cyprinodon*

pupfishes endemic to San Salvador Island, Bahamas. *Bulletin of the Peabody Museum of*

Natural History **54**, 231–240.

Martin, C. H. and Wainwright, P. C. (2013b). Multiple fitness peaks on the adaptive landscape

drive adaptive radiation in the wild. *Science (New York, N.Y.)* **339**, 208–211.

Martin, C. H., Cutler, J. S., Friel, J. P., Dening, T., Coop, G. and Wainwright, P. C. (2015).

Complex histories of repeated colonization and hybridization cast doubt on the clearest

examples of sympatric speciation in the wild. *Evolution* **69**, 1406–1422.

Martin, C. H., Erickson, P. A. and Miller, C. T. (2017). The genetic architecture of novel

trophic specialists: higher effect sizes are associated with exceptional oral jaw

diversification in a pupfish adaptive radiation. *Molecular Ecology* **26**, 624–638.

Martin, C. H., McGirr, J. A., Richards, E. J. and St John, M. E. (2019). How to Investigate

the Origins of Novelty: Insights Gained from Genetic, Behavioral, and Fitness

Perspectives. *Integr Org Biol* **1**, obz018.

Matsumoto, Y., Watanabe, T., Adachi, Y., Itoh, T., Ohtaki, T., Onda, H., Kurokawa, T.,

Nishimura, O. and Fujino, M. (2002). Galanin-like peptide stimulates food intake in the

rat. *Neurosci. Lett.* **322**, 67–69.

McGirr, J. A. and Martin, C. H. (2016). Novel candidate genes underlying extreme trophic

specialization in Caribbean pupfishes. *Molecular Biology and Evolution* **34**, 873–888.

McGirr, J. A. and Martin, C. H. (2021). Few Fixed Variants between Trophic Specialist

Pupfish Species Reveal Candidate Cis-Regulatory Alleles Underlying Rapid Craniofacial

Divergence. *Mol. Biol. Evol.* **38**, 405–423.

Meng, L., Bian, Z., Torensma, R. and Von den Hoff, J. W. (2009). Biological mechanisms in

palatogenesis and cleft palate. *Journal of Dental Research* **88**, 22–23.

Miller, C. T., Beleza, S., Pollen, A. A., Schluter, D., Kittles, R. A., Shriver, M. D. and
Kingsley, D. M. (2007). cis-Regulatory changes in Kit ligand expression and parallel
evolution of pigmentation in sticklebacks and humans. *Cell* **131**, 1179–1189.

Monteiro, A. and Podlaha, O. (2009). Wings, horns, and butterfly eyespots: how do complex
traits evolve? *PLoS biology* **7**, e37.

Naqvi, S., Hoskens, H., Wilke, F., Weinberg, S. M., Shaffer, J. R., Walsh, S., Shriver, M. D.,
Wysocka, J. and Claes, P. (2022). Decoding the Human Face: Progress and Challenges
in Understanding the Genetics of Craniofacial Morphology. *Annu. Rev. Genomics Hum.*
Genet.

Navon, D., Male, I., Tetrault, E. R., Aaronson, B., Karlstrom, R. O. and Albertson, R. C.
(2020). Hedgehog signaling is necessary and sufficient to mediate craniofacial plasticity
in teleosts. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 19321–19327.

Palmer, K., Fairfield, H., Borgeia, S., Curtain, M., Hassan, M. G., Dionne, L., Yong Karst,
S., Coombs, H., Bronson, R. T., Reinholdt, L. G., et al. (2016). Discovery and
characterization of spontaneous mouse models of craniofacial dysmorphology.
Developmental Biology **415**, 216–227.

Pavlidis, P., ??ivkovi??, D., Stamatakis, A. and Alachiotis, N. (2013). SweeD: Likelihood-
based detection of selective sweeps in thousands of genomes. *Mol. Biol. Evol.* **30**, 2224–
2234.

Platzer, M. and Englert, C. (2016). *Nothobranchius furzeri*: A model for aging research and
more. *Trends in Genetics* **32**, 543–552.

Powder, K. E. and Albertson, R. C. (2016). Cichlid fishes as a model to understand normal and

clinical craniofacial variation. *Developmental Biology* **415**, 338–346.

Price, A. L., Spencer, C. C. A. and Donnelly, P. (2015). Progress and promise in understanding

the genetic basis of common diseases. *Proceedings of the Royal Society B: Biological*

Sciences **282**, 20151684.

Protas, M., Conrad, M., Gross, J. B., Tabin, C. and Borowsky, R. (2007). Regressive

evolution in the Mexican cave tetra, Astyanax mexicanus. *Current biology : CB* **17**, 452–

454.

Reardon, W., Winter, R. M., Rutland, P., Pulleyn, L. J., Jones, B. M. and Malcolm, S.

(1994). Mutations in the fibroblast growth factor receptor 2 gene cause Crouzon

syndrome. *Nature Genetics* **8**, 98–103.

Richards, E. J. and Martin, C. H. (2017). Adaptive introgression from distant Caribbean

islands contributed to the diversification of a microendemic adaptive radiation of trophic

specialist pupfishes. *PLoS Genet.* **13**, 1–35.

Richards, E. J. and Martin, C. H. (2022). We get by with a little help from our friends: shared

adaptive variation provides a bridge to novel ecological specialists during adaptive

radiation. *Proc. Biol. Sci.* **289**, 20220613.

Richards, E. J., McGirr, J. A., Wang, J. R., St John, M. E., Poelstra, J. W., Solano, M. J.,

O'Connell, D. C., Turner, B. J. and Martin, C. H. (2021). A vertebrate adaptive

radiation is assembled from an ancient and disjunct spatiotemporal landscape. *Proc. Natl.*

Acad. Sci. U. S. A. **118**,.

Roberts, R. B., Ser, J. R. and Kocher, T. D. (2009). Sexual conflict resolved by invasion of a

novel sex determiner in Lake Malawi cichlid fishes. *Science (New York, N.Y.)* **326**, 998–

1001.

Roberts, R. B., Hu, Y., Albertson, R. C. and Kocher, T. D. (2011). Craniofacial divergence

and ongoing adaptation via the hedgehog pathway. *Proceedings of the National Academy*

of Sciences of the United States of America **108**, 13194–13199.

Rohner, N. (2018). Cavefish as an evolutionary mutant model system for human disease. *Dev.*

Biol. **441**, 355–357.

Signor, S. A. and Nuzhdin, S. V. (2018). The Evolution of Gene Expression in cis and trans.

Trends Genet. **34**, 532–544.

Sollenberg, U. E., Lundström, L., Bartfai, T. and Langel, Ü. (2006). M871—A Novel Peptide

Antagonist Selectively Recognizing the Galanin Receptor Type 2. *Int. J. Pept. Res. Ther.*

12, 115–119.

Sollenberg, U. E., Runesson, J., Sillard, R. and Langel, Ü. (2010). Binding of Chimeric

Peptides M617 and M871 to Galanin Receptor Type 3 Reveals Characteristics of Galanin

Receptor–Ligand Interaction. *Int. J. Pept. Res. Ther.* **16**, 17–22.

St John, M. E., Holzman, R. and Martin, C. H. (2020a). Rapid adaptive evolution of scale-

eating kinematics to a novel ecological niche. *J. Exp. Biol.* **223**,.

St John, M. E., Dixon, K. E. and Martin, C. H. (2020b). Oral shelling within an adaptive

radiation of pupfishes: Testing the adaptive function of a novel nasal protrusion and

behavioural preference. *J. Fish Biol.* **97**, 163–171.

St. John, M. E., Dunker, J. C., Richards, E. J., Romero, S. and Martin, C. H. (2021).

Parallel genetic changes underlie integrated craniofacial traits in an adaptive radiation of

trophic specialist pupfishes. *bioRxiv* 2021.07.01.450661.

Sutter, N. B., Bustamante, C. D., Chase, K., Gray, M. M., Zhao, K., Zhu, L.,

Padhukasahasram, B., Karlins, E., Davis, S., Jones, P. G., et al. (2007). A single IGF1

allele is a major determinant of small size in dogs. *Science (New York, N.Y.)* **316**, 112–

115.

Venn-watson, S., Carlin, K. and Ridgway, S. (2011). General and Comparative Endocrinology

Dolphins as animal models for type 2 diabetes : Sustained , post-prandial hyperglycemia

and hyperinsulinemia. *General and Comparative Endocrinology* **170**, 193–199.

Weinberg, S. M., Cornell, R. and Leslie, E. J. (2018). Craniofacial genetics: Where have we

been and where are we going? *PLoS Genet.* **14**, e1007438.

Wilkie, A. O., Slaney, S. F., Oldridge, M., Poole, M. D., Ashworth, G. J., Hockley, A. D.,

Hayward, R. D., David, D. J., Pulleyn, L. J. and Rutland, P. (1995). Apert syndrome

results from localized mutations of FGFR2 and is allelic with Crouzon syndrome. *Nature*

genetics **9**, 165–172.