Comparative Brain-Wide Mapping of Ketamine and Isoflurane-Activated Nuclei and Functional Networks

Yue Hu[1†], **Wenjie Du**[1†], **Jiangtao Qi**[1†], **Huoqing Luo**[2], **Zhao Zhang**[1], **Mengqiang Luo**[1*], **Yingwei Wang**[1*]

*For correspondence:
luomq16@fudan.edu.cn (FS);
wangyw@fudan.edu.cn (FMS)

[†]These authors contributed equally to this work

[1]Department of Anesthesiology, Huashan Hospital, Fudan University, 12 Middle Wulumuqi Road, Shanghai 200040, China; [2]School of Life Science and Technology, ShanghaiTech University; Shanghai 201210, China.

Abstract   Ketamine (KET) and isoflurane (ISO) are two widely used general anesthetics, yet their distinct and shared neurophysiological mechanisms remain elusive. In this study, we conducted a comparative analysis of the effects of KET and ISO on c-Fos expression across the brain, utilizing hierarchical clustering and c-Fos-based functional network analysis to evaluate the responses of individual brain regions to each anesthetic. Our findings reveal that KET activates a wide range of brain regions, notably in the cortical and subcortical nuclei involved in sensory, motor, emotional, and reward processing, with the temporal association areas (TEa) as a strong hub, suggesting a top-down mechanism affecting consciousness by primarily targeting higher-order cortical networks. In contrast, ISO predominantly influences brain regions in the hypothalamus, impacting neuroendocrine control, autonomic function, and homeostasis, with the locus coeruleus (LC) as a connector hub, indicating a bottom-up mechanism in anesthetic-induced unconsciousness. KET and ISO both activate brain areas involved in sensory processing, memory and cognition, reward and motivation, as well as autonomic and homeostatic control, highlighting their shared effects on various neural pathways. In conclusion, our results highlight the distinct but overlapping effects of KET and ISO, enriching our understanding of the mechanisms underlying general anesthesia.

Introduction

Despite considerable investigation into molecular targets, neural circuits, and functional connectivity associated with various anesthetics, our understanding of their effects on overall brain activity continues to be limited and incomplete (*Hemmings et al., 2019*). At the molecular level, ketamine (KET) modulates neuronal excitability predominantly through N-methyl-D-aspartate (NMDA) receptors, whereas isoflurane (ISO) primarily affects gamma-aminobutyric acid type A (GABAa) receptors. Their distinct mechanisms of action can each contribute to the loss of consciousness. (*Franks, 2008*). In systems neuroscience, the neural mechanisms of anesthetic induced unconsciousness involve both top-down and bottom-up processes (*Mashour, 2014*; *Mashour and Hudetz, 2017*). As evidenced by in vivo electrophysiology or functional magnetic resonance imaging (fMRI) studies, the top-down paradigm demonstrates that anesthetics induce unconsciousness by disrupting corticocortical and corticothalamic circuits responsible for neural information integration, while peripheral sensory information can still be conveyed to the primary sensory cortex (*Schroeder et al., 2016*; *Lee et al., 2013*). The bottom-up approach, exemplified by ISO, involves the activation of

sleep-promoting nuclei like ventrolateral preoptic nucleus (VLPO) and inhibition of arousal centers in the brainstem and diencephalon, supporting the shared circuits of sleep and anesthesia (*Moore et al., 2012*; *Nelson et al., 2002*). However, the limited spatial resolution of fMRI studies and the inability of electroencephalography (EEG) to capture specific brainstem nuclei significantly hinder the acquisition of comprehensive whole-brain information. Although a substantial body of knowledge has been gathered, our understanding of the reciprocal responses among different brain regions during general anesthesia remains relatively sparse and fragmented. To bridge these gaps, further investigation using advanced techniques that can capture whole-brain dynamics is needed to elucidate the complex interactions and shared mechanisms between various anesthetics.

Neuronal extracellular stimulation typically results in the elevation of adenosine 3',5'-cyclic monophosphate (cAMP) levels and calcium influx, ultimately leading to the upregulation of immediate early genes (IEGs) such as c-fos (*Yap and Greenberg, 2018*; *Morgan and Curran, 1989*). The translation product of c-fos, c-Fos protein, offers single-cell spatial resolution and has been utilized as a biomarker to identify anesthetic-activated brain regions (*Zhang et al., 2022*). Previous investigations of c-Fos expression throughout the brain demonstrated that GABAergic agents inhibited cortical activity while concurrently activating subcortical brain regions, including VLPO, median preoptic nucleus (MEPO), lateral septal nucleus (LS), Edinger-Westphal nucleus (EW), and locus coeruleus (LC) (*Smith et al., 2016*; *Lu et al., 2008*; *Yatziv et al., 2020*; *Han et al., 2014*). In contrast, KET was shown to provoke wake-like c-Fos expression and intense augmentation of c-Fos expression in various brain regions at clinical dosages (75-100 mg/kg) (*Lu et al., 2008*). It is important to note that these experiments administered KET at lights-on and GABAa receptor agonists at lights-off, potentially introducing circadian influences for direct comparison of KET and ISO. Moreover, it has been revealed that the state of general anesthesia is not determined by activity in individual brain areas but emerges as a global change within the brain. This change involves the activation of lateral habenular nucleus (LHb), VLPO, supraoptic nucleus (SO), and central amygdaloid nucleus (CeA), which are essential for anesthetic induced sedation, unconsciousness, or analgesia (*Moore et al., 2012*; *Gelegen et al., 2018*; *Hua et al., 2020*; *Jiang-Xie et al., 2019*). However, comprehensive brain-wide mapping and comparison of distinct anesthetic (KET and ISO) activated nuclei at the cellular level has not been fully elucidated.

In this study, we initially investigated the distribution of nuclei activated by KET and ISO across 984 brain regions. This was accomplished through immunochemical labeling and utilizing a customized MATLAB software package, which aided in signal detection and alignment with the Allen Mouse Brain Atlas reference (*Ma et al., 2021*). Subsequently, we analyzed whole-brain c-Fos expression elicited by KET and ISO using hierarchical clustering, and assessed inter-regional correlations by calculating the covariance across subjects for each brain region pair. Regions with significantly positive correlations were identified to construct functional networks, and graph theory-based methods were applied to identify hub nodes. Our findings revealed distinct yet overlapping whole-brain activation patterns for KET and ISO.

Results

Identification of brain regions activated by KET

To investigate the pattern of c-Fos expression throughout the brain, either 1.5% ISO was continuously ventilated or 100 mg/kg KET was given 90 minutes before harvesting (Figure 1A). Sample collection for all mice was uniformly conducted at 14:30 (ZT7.5), and the c-Fos labeling and imaging procedures were performed using consistent parameters across all experiments. To quantitatively assess brain states during anesthesia, we utilized EEG and electromyography (EMG) recordings. The depth of anesthesia was evaluated based on the power ratio of the EEG signals within the delta (0.5–4 Hz) and theta (6–10 Hz) frequency bands, combined with the changes in the EMG (*Luo et al., 2023*). Our results indicated that the administration of either KET or ISO elevated the EEG delta/theta power ratio while reducing EMG power (Figure 1—figure supplement 1), suggesting

that each drug induced a loss of consciousness. The raw images of brain slices were aligned to the Allen Mouse Brain Atlas, followed by signal detection and analysis (Figure 1B). Building upon the framework established in previous literature, our study first categorizes the whole brain of mice into 53 distinct subregions (*Do et al., 2016*). To assess the brain regions activated by KET and ISO in comparison to their respective controls, Z-scores were calculated for a statistical comparison of c-Fos expression differences. This process entailed normalizing the mean expression difference against the standard deviation of the control group. Our findings revealed that KET elicited significant activation in almost all 53 brain regions when compared to the saline group. Conversely, ISO showed substantial activation across various regions, excluding the cortical areas, when compared to the home cage group (Figure 2—figure supplement 1).

To enable a more detailed examination and facilitate clearer differentiation and comparison of the effects caused by KET and ISO, we subdivided the 53 brain regions into 201 distinct areas. This approach, guided by the standard mouse atlas available at http://atlas.brain-map.org/atlas, allowed for an in-depth analysis of the responses in various brain regions. Compared to the saline group, KET activated 141 out of a total of 201 brain regions (Figure 2). To further identify the brain regions that are most significantly affected by KET, we calculated Cohen's d for each region to quantify the magnitude of activation and subsequently focused on those regions that had a corrected p-value below 0.05 and effect size in the top 40% (Figure 3, Figure 3—figure supplement 1). Our findings were in line with previous studies: numerous cortical regions associated with somatosensory, auditory, visual, and movement were activated (Figure 3—source data 1). Notably, KET activated sensory areas, such as visual areas (VIS), auditory areas (AUD), somatosensory areas (SS), gustatory areas (GU), and higher-order cortical areas, including the anterior cingulate area (ACA), temporal association areas (TEa), retrosplenial area (RSP), and dorsal peduncular area (DP). This contrasts with the usual inhibiting effect of anesthetics on these neural circuits and suggests the unique ability of KET to sustain certain neuronal circuits in an active state (*Brown et al., 2011*). Furthermore, its impact on the olfactory system, involving key structures such as the anterior olfactory nucleus (AON), main olfactory bulb (MOB), olfactory tubercle (OT), perirhinal area (PERI), piriform area (PIR), and ectorhinal area (ECT), was distinct compared to other types of general anesthetics (*Zhao et al., 2020*). Moreover, KET affected emotional and reward processing regions such as the nucleus accumbens (ACB) (*Floresco, 2015*), orbital area (ORB) (*Rolls et al., 2020*), infralimbic area (ILA), and prelimbic area (PL) (*Pfarr et al., 2018*), suggesting an impact on the circuitry responsible for modulating reward and affective states. The thalamus, including the paraventricular nucleus of the thalamus (PVT) (*Ren et al., 2018*) and lateral group of the dorsal thalamus (LAT) (*Suzuki and Larkum, 2020*), which are vital in arousal regulation, also showed enhanced activity under the influence of KET. Hindbrain regions under KET anesthesia, including the magnocellular reticular nucleus (MARN), paragigantocellular reticular nucleus (PGRN), lateral reticular nucleus (LRN), nucleus raphe obscurus (RO), and sublaterodorsal nucleus (SLD), showed notable activity. In particular, the activation in regions like PGRN and SLD suggests that KET modulates brain regions critical for the regulation of rapid eye movement (REM) sleep (*Goutagny et al., 2008*). The activation of diverse regions such as the fundus of striatum (FS), substantia innominata (SI), and agranular insular area (AI), underscores that the effects of KET are not confined to classical neural pathways, but rather encompass a wide array of brain networks. Overall, KET induces marked activation across various brain regions, encompassing not only higher-order cortical areas but also those involved in sensory processing, emotional regulation, reward systems, and arousal mechanisms, thereby highlighting its distinct and widespread influence on diverse neural networks.

Identification of brain regions activated by ISO

To identify brain regions activated by ISO, we first reviewed nuclei previously reported to be activated by ISO. These encompass the PIR in the cortex, the lateral septal nucleus (LS), and the central amygdaloid nucleus (CEA) in the striatum. Additionally, activations were observed in hypothalamic regions such as the SO, VLPO, and MPO. Activations were also observed in the EW, the nucleus of

Figure 1. Experimental design and workflow for data analysis. (A) Experimental timeline for anesthetic exposure. Mice were handled for four days, subsequently exposed to KET (100 mg/kg, i.p.) or 1.5% ISO for 90 minutes, and then euthanized for immunostaining and data analysis. (B) Steps for data processing. Example of brain section registration to a corresponding coronal section from the Allen Brain Atlas. For Atlas rotation, the Allen reference atlas was rotated to mimic the slice angle of the experimental brain. Image registration maps the original coronal image (upper panel) to the corresponding Allen Mouse Brain Atlas (lower panel). The registration module applies several geometric transformations (translation, rotation, and scaling) to optimize the matching of the original image to the anatomical structures. Fluorescence signals detected in the original image were projected onto the Allen Mouse Brain Atlas for quantification. Finally, the processed data undergo hierarchical clustering and network analysis to investigate the patterns of c-Fos expression and central network nodes.

Figure 1—figure supplement 1. EEG and EMG power change after each treatment. The box represents the 25th-75th percentiles; the central line denotes the median; whiskers indicate maximum and minimum values. n = 6, 6, 8, 6 for the home cage, ISO, saline, and KET, respectively. (A) Normalized change in EEG power: ISO vs KET, $p > 0.99$; Saline vs KET, $p = 0.01$; Home cage vs ISO, $p = 0.11$. (B) Normalized change in EMG power: ISO vs KET, $p = 0.36$; Saline vs KET, $p = 0.30$; Home cage vs ISO, $p = 0.02$. Analyses were conducted using the Kruskal-Wallis test, followed by Dunn's multiple comparisons test.

Figure 2. KET and ISO induced c-Fos expression relative to their respective control group across 201 distinct brain regions. Z-scores represent the normalized c-Fos expression in the KET and ISO groups, calculated against the mean and standard deviation from their respective control groups. Statistical analysis involved the comparison of Z-scores to a null distribution with a zero mean and adjustment for multiple comparisons using the Benjamini–Hochberg method at a 5% false discovery rate (*p < 0.05, **p < 0.01, ***p < 0.001). n = 6, 6, 8, 6 for the home cage, ISO, saline, and KET, respectively. Missing values resulted from zero standard deviations in control groups. Brain regions are categorized into major anatomical subdivisions, as shown on the left side of the graph.

Figure 2—figure supplement 1. KET and ISO induced c-Fos expression relative to their respective control group across 53 distinct brain regions. Z-scores for c-Fos expression in the KET and ISO groups were normalized to the mean and standard deviation of their respective control groups. Statistical analysis involved the comparison of Z-scores to a null distribution with a zero mean and adjustment for multiple comparisons using the Benjamini–Hochberg method at a 5% false discovery rate (*p < 0.05, **p < 0.01, ***p < 0.001). Brain regions are organized into major anatomical subdivisions, as indicated on the left side of the graph.

Figure 3. Brain regions exhibiting significant activation by KET. (A) Fifty-five brain regions exhibited significant KET activation. These were chosen from the 201 regions analyzed in Figure 2, focusing on the top 40% ranked by effect size among those with corrected p values less than 0.05. Data are presented as mean ± SEM, with p-values adjusted for multiple comparisons (*p < 0.05, **p < 0.01, ***p < 0.001). (B) Representative immunohistochemical staining of brain regions identified in Figure 3A, with control group staining available in Figure 3—figure supplement 1. Scale bar: 200 μm.

Figure 3—figure supplement 1. c-Fos expression in distinct brain regions following saline administration. Representative immunohistochemical staining of c-Fos-positive cells in the specified brain regions is provided as a control for the c-Fos expression observed in brain regions after KET exposure, as detailed in Figure 3. Scale bar: 200 μm.

Table 1. Figure 3—source data 1. Summary of prior studies on KET activated brain regions identified via c-Fos immunostaining.

the solitary tract (NTS), locus coeruleus (LC), the ventral group of the dorsal thalamus (VeN), and the area postrema (AP) (Figure 4—source data 1). Building upon these established findings, our study further explored the effects of ISO. Using the same criteria applied to KET, which involved selecting regions with Cohen's d values in the top 40% of significantly activated areas from Figure 2, we identified 32 key brain regions impacted by ISO (Figure 4, Figure 4—figure supplement 1). Our findings not only confirmed the activation of known regions like CeA, SO, NTS, and LC, but also broadened the scope of understanding regarding the effects of ISO, revealing its impact on several previously unexplored areas. This included the paraventricular hypothalamic nucleus (PVH), arcuate hypothalamic nucleus (ARH), lateral hypothalamic area (LHA), tuberal nucleus (TU), median preoptic nucleus (MEPO), lateral preoptic area (LPO), retrochiasmatic area (RCH), periventricular hypothalamic nucleus intermediate part (PVi), dorsomedial nucleus of the hypothalamus (DMH), and anteroventral periventricular nucleus (AVPV). The activation of these regions suggested a substantial influence of ISO on hypothalamic regulation, including neuroendocrine control, autonomic function, and homeostasis (*Gu et al., 1999*; *Adamantidis and de Lecea, 2023*). Additionally, ISO affected the areas governing circadian rhythms and sleep-wake regulation, as indicated by activation in the suprachiasmatic nucleus (SCH) and subparaventricular zone (SBPV) (*Lu et al., 2001*). Moreover, sensory processing and reward circuitry, including the OT (*Giessel and Datta, 2014*), SI (*Cui et al., 2022*), and ACB (*Zhou et al., 2022*), demonstrated significant activation. Activation of the perirhinal area (PERI) (*Yang and Naya, 2023*), and nucleus of reuniens (RE) (*Huang et al., 2021*) pointed to a potential impact on cognitive processes and memory systems. Our study also revealed ISO activates the EW, a finding that aligned with observations under sevoflurane anesthesia (*Yi et al., 2023*). The bed nuclei of the stria terminalis (BST), associated with stress responses and emotional regulation, exhibited significant activation (*Daniel and Rainnie, 2016*). In summary, our findings suggest that ISO predominantly influences the hypothalamus, a key area for neuroendocrine regulation, autonomic control, and homeostasis, while also influencing brain functions associated with sensory processing, reward mechanisms, stress response, and cognitive functions.

The co-activation of multiple brain regions by KET and ISO indicates that they have overlapping effects on brain functions. Examples of these effects include impacts on sensory processing, as evidenced by the activation of the PIR, ENT (*Chapuis et al., 2013*), and OT(*Giessel and Datta, 2014*), pointing to changes in sensory perception typical of anesthetics. Memory and cognitive functions are influenced, as indicated by the activation of the subiculum (SUB) (*Roy et al., 2017*), dentate gyrus (DG) (*Sun et al., 2020*), and RE (*Huang et al., 2021*). The reward and motivational systems are engaged, involving the ACB and ventral tegmental area (VTA), signaling the modulation of reward pathways (*Al-Hasani et al., 2021*). Autonomic and homeostatic control are also affected, as shown by areas like the lateral hypothalamic area (LHA) (*Mickelsen et al., 2019*) and medial preoptic area (MPO) (*McGinty and Szymusiak, 1990*), emphasizing effects on functions such as feeding and thermoregulation. Stress and arousal responses are impacted through the activation of the paraventricular hypothalamic nucleus (PVH) (*Rasiah et al., 2023*; *Islam et al., 2022*) and LC (*Ross and Van Bockstaele, 2020*). This broad activation pattern highlights the overlap in drug effects and the complexity of brain networks in anesthesia.

Comparison of c-Fos activation patterns in response to KET and ISO

To reveal common activation patterns and potential associations among the brain regions responding to KET and ISO, hierarchical clustering was employed on 53 brain regions as delineated in Figure 2–figure supplement 1. We normalized the c-Fos densities in each of the 53 regions to the average values of that region in the control group and then log-transformed the data. Hierarchical clustering of these values revealed distinct patterns of drug-induced changes (Figure 5A), with both KET and ISO showing optimal clustering at a 0.5 cut-off value, as confirmed by high silhouette coefficients (Figure 5B). Concurrently, we computed the number of clusters formed at each potential cut-off value and found that a 0.5 threshold resulted in four distinct clusters for both KET and ISO

Figure 4. Brain regions exhibiting significant activation by ISO. (A) Brain regions significantly activated by ISO were initially identified using a corrected p-value below 0.05. From these, the top 40% in effect size (Cohen's d) were further selected, resulting in 32 key areas. p-values are adjusted for multiple comparisons (**p < 0.01, ***p < 0.001). (B) Representative immunohistochemical staining of brain regions identified in Figure 4A. Control group staining is available in Figure 4—figure supplement 1. Scale bar: 200 µm. Scale bar: 200 µm. (C) A Venn diagram displays 43 brain regions co-activated by KET and ISO, identified by the adjusted p-values (p < 0.05) for both KET and ISO. CTX: cerebral cortex; CNU: cerebral nuclei; TH: thalamus; HY: hypothalamus; MB: midbrain; HB: hindbrain.

Figure 4—figure supplement 1. c-Fos expression in home cage group. Representative immunohistochemical staining of c-Fos-positive cells in the home cage group, serving as a baseline comparison for the ISO groups. Scale bar: 200 µm.

Table 2. Figure 4—source data 1. Summary of prior studies on brain regions activated by ISO, as detected through c-Fos immunostaining.

treatments (Figure 5C). After hierarchical clustering, we preserved the sequence of brain regions and replaced log relative c-Fos densities with Z values, calculated by dividing the log relative c-Fos density in each brain region by the standard error, for both KET and ISO conditions (Figure 5E). KET administration elicited a significant activation in a broad array of brain regions, with cluster 4 displaying the most substantial upregulation in expression. This cluster incorporates cortical areas, including the visceral area (VISC), GU, ORB, somatomotor areas (MO), ACA, and SS, and cerebral nuclei that are integral for motor coordination and motivational behaviors, specifically, the dorsolateral and ventrolateral pallidum (PALd, PALv) and striatum (STRd, STRv) (*Grillner et al., 2005*) (Figures 5D and 5E). No significantly inhibited brain regions were detected in the ISO group (as no Z value fell below -2); clusters 3 and 4 expression levels were similar to or lower than those in the control group, with cluster 3 having a mean Z value of 0.27 and cluster 4 a mean Z value of -1.1, as shown in Figures 5D and 5E. The most pronounced upregulation of ISO-induced c-Fos expression was observed in the second cluster (mean Z value=4.7), which includes brain regions responsible for a variety of functions: the periventricular zone (PVZ) and lateral zone (LZ) of the hypothalamus, vital for endocrine and autonomic system regulation (*Gu et al., 1999*); the ventral striatum (STRv), pallidum (PAL), and lateral septal complex (LSX) of the striatum, key areas in reward and motivation (*Cox and Witten, 2019*); the taenia tecta (TT), PIR, essential for olfactory processing (*Bekkers and Suzuki, 2013*; *Kataoka et al., 2020*). This collective upregulation in cluster 2, indicates a consistent activation pattern and drug response among diverse functional brain regions. It underscores the impact of ISO on hypothalamic regulation, reward mechanisms, and olfactory processing, with marked activation observed in the hypothalamic and striatal regions. In contrast, KET induces widespread activation across the brain, with heightened activity in a key cluster that includes cortical and cerebral nuclei governing complex sensory, emotional, and motor processes, thus highlighting its notable influence on higher-order functions.

Similarities and differences in KET and ISO-activated c-Fos brain areas

To achieve a more granular analysis and better discern the differences in responses between KET and ISO, we expanded our study from the initial 53 brain regions to 201 distinct subregions. After log-transforming the c-Fos densities relative to controls, we applied hierarchical clustering to uncover patterns of brain activity (Figure 6A). ISO demonstrates a more concentrated influence on brain activity with consistently higher clustering quality and fewer clusters than KET, suggesting a focused impact on specific brain functions (Figures 6B and 6C). Figure 6E maintains the order from hierarchical clustering and marks the most activated brain regions in the top 10% of Z values with red boxes and areas of potential suppression with Z values below -2 with white boxes. In the hierarchical clustering analysis of KET exposure, cluster 2 with a mean Z value of -0.01 was similar to the control group in terms of activation, while the other four clusters showed increased activation. Cluster 5, showing the most significant activation, is composed mainly of the cortical and striatal regions crucial for sensory processing, spatial memory, reward, and cognitive functions (*Haber, 2016*). In addition, brain regions within the thalamus and hindbrain, including RE, inferior olivary complex (IO), nucleus raphe obscurus (RO), MARN, and PGRN, also display similar activation patterns. This composition, centered on cortical regions, implies that KET primarily targets higher-order brain functions while also influencing fundamental neurological processes through its effect on cerebral nuclei, the thalamus, and the hindbrain. ISO hierarchical clustering identified three distinct clusters, with cluster 3 exhibiting the highest mean Z value of 7.27. This cluster encompasses the SO, VLPO, TU, and CEA (Figure 6E). Notably, the SO, VLPO, and CEA have been previously reported to play critical roles in the mechanisms of anesthesia (*Moore et al., 2012*; *Hua et al., 2020*; *Jiang-Xie et al., 2019*). Cluster 1 showed a relative decrease in expression (n= 91, mean Z value: -0.54), indicating that nearly half of the brain areas had reduced c-Fos expression compared to the control group. Cluster 2, characterized by overall moderate activation, was particularly significant in the top 10% of its regions based on Z value ranking, as highlighted by the red box (Figure 6E). This segment includes key regions such as ARH and PVH, which were also prominently activated

Figure 5. Comparison of c-Fos activation patterns in response to KET and ISO. (A) Hierarchical clustering of log-relative c-Fos density data for KET and ISO was conducted using the complete linkage method, based on the Euclidean distance matrix. Clusters were identified using a dendrogram cut-off ratio of 0.5, with numerical labels indicating distinct clusters. (B) Silhouette values plotted against the dendrogram tree height ratio for KET and ISO reveal higher Silhouette values at a 0.5 ratio (dashed line), suggesting optimal clustering at this level. (C) The number of clusters for each treatment condition at different dendrogram tree height ratios is shown, with a cut-off level of 0.5 yielding 4 clusters for both KET and ISO (marked by the dashed line). (D) Bar graphs represent the Z values for clusters under both KET and ISO conditions. Data are presented as mean ± SEM. One-way ANOVA with Tukey's post hoc multiple comparisons. ns: no significance; ***p < 0.001. (E) Heatmaps illustrate Z values of the log relative c-Fos expression density in clustered brain regions. The arrangement and abbreviations of the brain regions, along with the numerical labels, correspond to those in Figure 5A. Red box in each heatmap emphasizes the cluster with the highest mean Z value in its respective condition. CTX: cerebral cortex; CNU: cerebral nuclei; TH: thalamus; HY: hypothalamus; MB: midbrain; HB: hindbrain.

as shown in Figure 4. It furthermore encompasses cortical areas such as ILA, DP, and TT cortices involved in emotional and reward processing (*Pfarr et al., 2018*; *Kataoka et al., 2020*) and integrates striatal and pallidal regions like OT and ACB and midbrain and hindbrain structures such as AP and NTS. The collective inclusion of these diverse regions, exhibiting pronounced and similar activation patterns, suggests a unified response to ISO, indicative of potential functional connectivity and interaction across these areas. In conclusion, the activation of specific areas such as the SO, VLPO, TU, and CEA by ISO in cluster 3 suggests a targeted influence. In contrast, the extensive engagement of cortical areas by KET in cluster 5, along with its effects on striatal and thalamic regions, underscores a profound and widespread modulation of both higher-order and fundamental neural functions.

Network generation and Hub identification

Previous research has established that general anesthesia is mediated by various brain regions (*Moore et al., 2012*; *Gelegen et al., 2018*; *Hua et al., 2020*; *Jiang-Xie et al., 2019*). The expression of c-Fos serves as an indicator of neuronal activity, providing a single index of activation per region per animal. By examining the covariance of c-Fos expression across animals within each group, we can infer interactions between brain regions and identify functional networks engaged during general anesthesia (*Wheeler et al., 2013*). Initially, 63 brain regions were incorporated, including those significantly activated by KET and ISO (as indicated in Figures 3 and 4) and others previously associated with general anesthesia and sleep-wakefulness mechanisms (*Leung et al., 2014*). A comprehensive interregional correlation for four groups was calculated (Figure 7A). Correlation analysis revealed similar network connectivity and densities between the ISO and home cage groups (mean interregional correlations: Fisher Z = -0.018, p = 0.98; network densities: 0.10 vs. 0.09). In contrast, KET significantly increased network density (0.46 vs. 0.06) and mean interregional correlations compared to the saline group (Fisher Z = 3.54, p < 0.001; Figure 7–figure supplement 1D), indicating enhanced interregional correlation with KET administration. Functional networks were segmented into modules based on hierarchical clustering of correlation coefficients (Figure 7–figure supplement 1A), with dendrogram cutting at a value of 0.7 for high Silhouette scores (Figure 7–figure supplement 1B). This resulted in two modules for the home cage and KET groups, and four for the saline and ISO groups (Figure 7–figure supplement 1C). Networks were formed using Pearson's coefficients over 0.82 and significant correlations (p < 0.05), yielding a single module for KET and three modules for ISO (Figure 7B).

Hubs are nodes that occupy critical central positions within the network and enable the network to function properly. Due to the singular module structure of the KET network and the sparsity of intermodular connections in the home cage and saline networks, the assessment of network hub nodes did not employ within-module degree Z-score and participation coefficients as these measures predominantly underscore the importance of connections within and between modules (*Kimbrough et al., 2020*). The analysis evaluated nodal importance using degree (number of direct connections), betweenness centrality (proportion of shortest network paths passing through a node), and eigenvector centrality (influence based on the centrality of connected nodes), identifying brain regions that scored highly in all three as hub nodes (Figure 7–figure supplement 2). The LC, characterized by its high degree and eigenvector centrality, as well as its notably higher betweenness centrality (Figure 7–figure supplement 2) underscores its pivotal role in propagating and integrating neural signals within the ISO-influenced network. In the KET group, the TEa serves as a hub, emphasizing its integrative role in KET-induced dissociative anesthesia. Meanwhile, the periaqueductal gray (PAG) and LSr function as connector hubs in the home cage and saline groups, respectively.

Discussion

In this study, we conducted a comparative analysis to examine the effects of two general anesthetics, KET and ISO, on c-Fos expression throughout various brain regions. Our results revealed that

Figure 6. Similarities and differences in KET and ISO-activated c-Fos brain areas. (A) Heatmaps display hierarchical clustering of brain regions based on log-relative c-Fos density for KET and ISO groups, using complete linkage on Euclidean distances. Clusters are demarcated with a 0.5 dendrogram cut-off. (B) Silhouette values are plotted against the dendrogram tree height ratios in Figure 6A. (C) The number of clusters identified at different dendrogram tree height ratios, with a cut-off ratio of 0.5 resulting in 5 for KET and 3 clusters for ISO (indicated by the dashed line). (D) Z values for identified clusters under KET and ISO conditions. Data are presented as mean ± SEM. One-way ANOVA with Tukey's post hoc multiple comparisons. ns: no significance; ***$p < 0.001$. (E) Heatmaps illustrate Z values of log-relative c-Fos densities in brain region clusters, arranged and labeled as in Figure 6A. Red boxes highlight regions ranking in the top 10 percent within their respective clusters, and white boxes indicate regions with Z values less than -2.

Figure 6—figure supplement 1. Region labels for the hierarchical clustering of the KET group in Figure 6A.

Figure 6—figure supplement 2. Region labels for the hierarchical clustering of the ISO group in Figure 6A.

Figure 7. Generation of anesthetics-induced networks and identification of hub regions. (A) Heatmaps display the correlations of log c-Fos densities within brain regions (CTX, CNU, TH, HY, MB, and HB) for various states (home cage, ISO, saline, KET). Correlations are color-coded according to Pearson's coefficients. The brain regions within each anatomical category are organized by hierarchical clustering of their correlation coefficients. Full names and expression levels for each brain region are detailed in Figure 3—figure supplement 1. (B) Network diagrams illustrate significant positive correlations (P < 0.05) between regions, with Pearson's r exceeding 0.82. Edge thickness indicates correlation magnitude, and node size reflects the number of connections (degree). Node color denotes betweenness centrality, with a spectrum ranging from dark blue (lowest) to dark red (highest). The networks are organized into modules consistent with the clustering depicted in Figure 6—figure supplement 1A.

Figure 7—figure supplement 1. Hierarchical clustering across different conditions. (A) Heatmaps depict the hierarchical clustering of brain regions within the home cage, ISO, saline, and KET groups, using Euclidean distance matrices calculated from correlation coefficients. Modules are demarcated within each dendrogram at a cut-off threshold of 0.7. (B) Silhouette scores are plotted against the dendrogram tree height ratio for each condition, with optimal cluster definition indicated by a dashed line at a 0.7 ratio. (C) The number of clusters formed at different cutoff levels. At a ratio of 0.7, ISO and saline treatments result in three clusters, whereas home cage and KET conditions yield two clusters. (D) The mean Pearson's correlation coefficient (r) was computed from interregional correlations displayed in Figure 7A. Data were analyzed using one-way ANOVA with Tukey's post hoc test, ***p < 0.001.

Figure 7—figure supplement 2. Hub region characterization across different conditions: home cage (A), ISO (B), saline (C), and KET (D) treatments. Brain regions are sorted by degree, betweenness centrality, and eigenvector centrality, with each metric presented in separate bar graphs. Bars to the left of the dashed line indicate the top 20% of regions by rank, highlighting the most central nodes within the network. Red bars signify regions that consistently appear within the top rankings for both degree and betweenness centrality across the metrics.

Figure 8. The possible mechanism for KET and ISO-induced unconsciousness. The distinct pathways of KET and ISO-induced unconsciousness can be explained by two contrasting mechanisms. The 'top-down' process attributes KET's effect to widespread cortical activation (represented in yellow), with the temporal association areas (TEa) acting as the central node in the functional network (depicted in blue). Conversely, the 'bottom-up' approach suggests that ISO-induced unconsciousness stems from the activation of certain hypothalamic regions (highlighted in yellow), with the locus coeruleus (LC) acting as the hub node within the ISO-induced functional network. Green denotes representative regions co-activated by both ISO and KET. Adapted from (*Mashour, 2014*; *Mashour and Hudetz, 2017*; *Reimann and Niendorf, 2020*). PL, prelimbic area; ILA, infralimbic areas; LS, lateral septal nucleus; ACB, nucleus accumbens; OT, olfactory tubercle; PVT, paraventricular nucleus of the thalamus; EW, Edinger-Westphal nucleus; SO, supraoptic nucleus; PVH, paraventricular hypothalamic nucleus; NTS, nucleus of the solitary tract; CTX: cerebral cortex; TH: thalamus; HY, hypothalamus; MB; midbrain; HB, hindbrain.

KET predominantly activates the cerebral cortex, with the TEa emerging as the central node in its functional network. This pattern suggests a top-down mechanism of action for KET. In contrast, ISO primarily stimulates subcortical regions, notably the hypothalamus, with the LC identified as the hub node within the ISO-induced functional network. This finding is consistent with a bottom-up approach (Figure 7) (*Mashour, 2014*; *Mashour and Hudetz, 2017*). Additionally, our analysis revealed the brain nuclei that were co-activated by both KET and ISO, indicating shared neural pathways involved in general anesthesia.

Studies utilizing functional MRI, PET scans, EEG, and computational modeling have indicated that the emergence of consciousness results from active and coherent connections and communication among various brain regions, enabling global information availability and integration within a neural network (*Lamme, 2006*; *Dehaene and Changeux, 2011*). Our findings revealed a significant activation throughout the cortex-dominated brain following KET administration, accompanied by a notable increase in network density (Figure 7A). The observed hyperactivity and heightened functional connectivity could potentially compromise the effective integration of information within the network, possibly leading to the loss of consciousness. This finding aligns with previous studies which indicate that KET preferentially inhibits GABAergic interneurons through NMDA receptor mediation, leading to abnormal excitatory activity in the cortex and limbic system and potentially disrupting consciousness (*Seamans, 2008*; *Brown et al., 2010*). Furthermore, our analysis identified the TEa as a connector hub within this hyperactivated network (Figure 7). This finding, when combined with the extensive cortical activation induced by KET, support the hypothesis that consciousness alterations from KET are a result of a top-down mechanism, primarily attributed to the disruption of high-level cognitive processes regulating sensory input and consequently impairing corticothalamic circuits (Figure 8A) (*Mashour, 2014*).

Moreover, our study employed a hierarchical clustering method to categorize brain regions by their relative activation levels, uncovering potential functional similarities and connections among these areas. Notably, during the administration of ISO, regions such as the SO, VLPO, TU, and CeA demonstrated similar and significant activation (Figure 6E). This aligns with previous research highlighting the involvement of SO (*Jiang-Xie et al., 2019*), VLPO (*Moore et al., 2012*), and CeA (*Hua et al., 2020*) in anesthesia mechanisms. Additionally, we noted activation in several pallidum and striatum-related brain regions under ISO (Figure 4), providing a new perspective for understanding the mechanism of action of ISO. Additionally, our analysis utilizing c-Fos-based functional network analysis identified the LC as a key connector hub, highlighting the importance of brainstem neurotransmitter-related nuclei during the unconscious state induced by ISO. The LC, which is the primary source of noradrenaline in the brain, plays a pivotal role in regulating vigilance, as well as respiratory and cardiovascular responses (*Magalhães et al., 2018*; *Wood and Valentino, 2017*). During ISO anesthesia, key autonomic functions are typically suppressed, including the regulation of normal body temperature, respiration, blood pressure, and heart rate (*Skovsted and Sapthavichaikul, 1977*). In this context, the activation of the LC during the administration of ISO may act as a compensatory mechanism to maintain a necessary level of autonomic response. This mechanism helps prevent the excessive suppression of crucial autonomic functions induced by ISO. While it is known that activation of the LC during ISO anesthesia can have a wake-promoting effect (*Vazey and Aston-Jones, 2014*), this phenomenon might contribute to a balance between inducing loss of consciousness and preserving essential autonomic functions during anesthesia. However, the exact role of LC activation during ISO induced anesthesia warrants further investigation. Nevertheless, this observation, combined with the significant activation of hypothalamic-related nuclei under ISO, supports the bottom-up model of unconsciousness (*Mashour and Hudetz, 2017*). This model underscores the influence of anesthetics on the hypothalamus and brainstem and focuses on how the interruption of sensory information transmission from subcortical areas to the cortex alters levels of consciousness (Figure 8B). Together, these findings suggest a pivotal role for the hypothalamus and brainstem regions in the loss of consciousness induced by ISO.

Identifying shared neural features between KET and ISO is essential for understanding their

roles in anesthetic-induced unconsciousness, analgesia, and amnesia. The concurrent activation of the ACB and VTA alongside homeostatic control nuclei such as the LHA and MPO might suggest a shared mechanism in reward processing and homeostatic regulation (*Zhou et al., 2022*; *Al-Hasani et al., 2021*; *Mickelsen et al., 2019*; *McGinty and Szymusiak, 1990*). Additionally, the activation of neuroendocrine-related nuclei in the hypothalamus, specifically the PVH, SO, and AVPV, suggests anesthetic effects on hormonal release and homeostasis (*Gu et al., 1999*; *Adamantidis and de Lecea, 2023*). The exploration of other coactivated nuclei, including the PVT, RE, ARH, SCH, and RCH, is vital to clarify their specific anesthesia-related functions. Notably, each activated region may play multifaceted roles in anesthesia, affecting various physiological processes. The roles of these brain regions can be state-dependent and intricate, particularly in their contribution to unconsciousness, analgesia, and amnesia, underscoring the necessity for more focused research.

Our results demonstrate that c-Fos expression is significantly higher in the KET group compared to the ISO group, and in the saline group compared to the home cage group, as shown in Figures 2 and 3. Despite four days of acclimation, including handling and injections, intraperitoneal injections in the saline and KET groups might still elicit pain and stress responses in mice. This point is corroborated by the subtle yet measurable variations in brain states between the home cage and saline groups, characterized by changes in normalized EEG delta/theta power (home cage: 0.05±0.09; saline: -0.03±0.11) and EMG power (home cage: -0.37±0.34; saline: 0.04±0.13), as shown in Figure 1–figure supplement 1. These changes suggest a relative increase in brain activity in the saline group compared to the home cage group, potentially contributing to the higher c-Fos expression. Additionally, despite the use of consistent parameters for c-Fos labeling and imaging across all experiments, the substantial differences observed between the saline and home cage groups might be partly attributed to the fact that the operations were conducted in separate sessions. While the difference in EEG power between the ISO and home cage groups was not significant, the ISO group exhibited a similar increase in EEG power to the KET group (0.47 ± 0.07 vs 0.59 ± 0.10), suggesting that both agents induce loss of consciousness in mice. In terms of EMG power, ISO showed a notable decrease compared to its control group, whereas the KET group exhibited a smaller reduction (ISO: -1.815± 0.10; KET: -0.96 ± 0.21), possibly contributing to the higher c-Fos expression in the KET group. This aligns with previous research indicating that KET doses up to 150 mg/kg increase delta power and induce a wakefulness-like c-Fos expression pattern in the brain (*Lu et al., 2008*). Additionally, the differences in c-Fos expression may stem from dosages, administration routes, and distinct pharmacokinetic profiles, compounded by the absence of detailed physiological monitoring like blood pressure, heart rate, and respiration. Future research should include comprehensive physiological monitoring and controlled dosing to better understand the effects of anesthetics on brain activity.

Methods and Materials

Animals

All animal experiments were conducted in accordance with the National Institutes of Health guidelines and were approved by the Chinese Academy of Sciences' Institute of Neuroscience. Adult male wild-type (WT) mice (C57BL/6J) (8-10 weeks old, weight from 22 to 26 g) were purchased from institute-approved vendors (LingChang Experiment Animal Co., China). Mice were individually housed and maintained on a 12 h:12 h light/dark cycle (lights on at 07:00 a.m. and off at 07:00 p.m.) with food and water available ad libitum.

Drug administration

All experiments were conducted between 13:00-14:30 (ZT6–ZT7.5). We adapted mice to handling and the anesthesia chamber (10×15×15 cm) for four days to minimize experimental confound-induced c-Fos induction. For KET administration, adult male mice were handled for 10 minutes per day, and normal saline (NS) was injected intraperitoneally (i.p.) for four consecutive days at

13:00. On day five, a randomly chosen mouse received an injection of ketamine (Gutian Medicine,
H35020148) (n=6), and the control groups (n=8) received the same volume of saline. ISO group
(n=6) mice were handled and inhaled 1.5% isoflurane (RWD Life Science, 1903715) at 13:00 on day
four in the chamber. Meanwhile, the control groups (n=6) were left undisturbed in their home
cages before sampling. We confirmed the loss of righting reflex at 5 min after anesthetic exposure.
For 90 min after KET injection or ISO inhalation, mice were deeply anesthetized with 5% ISO and
transcardially perfused with 35 ml 0.1 M phosphate-buffered saline (PBS) followed by 35 ml 4%
paraformaldehyde (PFA). The brains were then removed and postfixed overnight with 4% PFA. Fol-
lowing fixation, the brains were dehydrated for 48 hours with 30% sucrose (wt/vol) in PBS. Coronal
sections (50 μm) of the whole brain were cut using a cryostat (HM525 NX, Thermo Scientific) after
being embedded with OCT compound (NEG-50, Thermo Scientific) and freezing.

Anesthesia depth measurement

EEG and EMG recordings were conducted as previously described (*Luo et al., 2023*). Specifically, a
stainless-steel electrode was positioned over the frontal cortex for EEG measurements, while an
insulated EMG electrode was inserted into the neck muscles. Additionally, a reference electrode
was placed above the cerebellum. The EEG/EMG data were captured using TDT system-3 amplifiers,
specifically RZ5 + RA16PA and RZ2 + PZ5 configurations. To measure the depth of anesthesia using
EEG/EMG signals, a fast Fourier transform spectral analysis was employed, featuring a frequency
resolution of 0.18 Hz. Data was noted every 5 seconds through a MATLAB tool and subsequently
verified manually by experts. The assessment was based on EMG power and the ratio of EEG's
delta power (0.5-4 Hz) to theta power (6-10 Hz). EEG and EMG power values within 30 minutes
post-administration were normalized to a 5-minute pre-administration baseline.

Immunohistochemistry

One out of every three brain slices (100 μm intervals) of each whole brain was washed three times
with 0.1 M phosphate-buffered saline (PBS) for 10 min and permeabilized for 30 minutes at room
temperature with 0.3% Triton X-100 in PBS (PBST). Slices were incubated for 2 hours at room tem-
perature with 2% normal donkey serum (Sigma, G6767) in PBS overnight at 4 °C with c-Fos primary
antibodies (226003, Synaptic Systems; 1:500) diluted in PBS with 1% donkey serum. After three
washes with PBST, slices were incubated with the Cy3 donkey anti-rabbit (711165152, Jackson;
1:200) secondary antibody for 2 hours at room temperature. Immunostained brain slices were
first mounted using VECTASHIELD mounting medium containing DAPI. Comprehensive imaging of
these slices was initially performed with a 10× objective on a fluorescent microscope (VS120, Olym-
pus). Regions showing statistically significant differences were then imaged in greater detail using
a 20× objective on a confocal microscope (FV300, Olympus). Due to the limited scope of a single
field of view, either 2 or 4 adjacent fields of view were stitched together to offer a comprehensive
representation of specific brain regions, such as the PL, ILA, LSc, LSr, SS, and VISC.

Quantification of c-Fos positive cells

425 The procedures used for c-Fos analysis were based on previous research (*Ma et al., 2021*). A
426 custom-written software package was employed for cellular quantification within brain images.
427 The software consists of three components: atlas rotation, image registration, and signal detec-
428 tion.

Atlas rotation module

430 Utilizing the Allen Mouse Brain Atlas, this module allows rotations in both horizontal and vertical
431 dimensions to align with mouse brain samples. To determine the appropriate rotation angles, we
432 manually pinpointed anatomical landmarks corresponding to the samples. For the left-right axis
433 rotation, we chose points from the CA3 pyramidal layer in both hemispheres and the posterior slice

of the dentate gyrus, specifically the granule cell layer. For the dorsal-ventral axis rotation, we iden-
tified key anatomical landmarks. These include the initial connections of the anterior commissure
and corpus callosum between the hemispheres, as well as ventral regions like the interpeduncular
nucleus and the suprachiasmatic nucleus. After determining these rotation angles, we adjusted
the reference atlas to match our samples.

Image registration module
This module uses a tool that aligns brain slice images with a reference atlas, facilitating the align-
ment of overall brain region distribution. Registration starts by matching the coronal plane of the
sample sections to the atlas. After defining the external boundaries of the brain section, the system
performs geometric transformations on the section to optimize its fit with the atlas.

Signal detection module
The detection module is specifically designed to automatically label c-Fos positive cells. Following
detection, each digitized brain section image underwent manual verification to ensure the accuracy
and precision of the c-Fos positive cell labeling.

Hierarchical clustering
Prior to hierarchical clustering in Figures 5 and 6, we calculated the relative c-Fos densities by divid-
ing the c-Fos densities of each brain region in the experimental groups by their respective controls
and then performed a log transformation on these values to obtain the log relative c-Fos densities.
These log ratios, which normalize the data and reduce variance, indicate differential expression
with a value of zero denoting no change compared to control. After normalizing the data, we per-
formed hierarchical clustering by first computing the pairwise Euclidean distances among brain
regions. Regions with the smallest distances, indicating high similarity, were grouped iteratively.
Cluster boundaries were defined using complete linkage, ensuring homogeneity within clusters
by considering the largest distance between members. In Figure 7A, hierarchical clustering was
performed within each of the cerebral cortex (CTX), thalamus (TH), cerebral nuclei (CNU), hypotha-
lamus (HY), midbrain (MB), and hindbrain (HB) regions based on the log-transformed c-Fos density
correlations. In Figure 7–figure supplement 1, hierarchical clustering was performed based on the
interregional log c-Fos density correlations to identify modules of coactivation within each treat-
ment group, revealing underlying functional connectivity networks (*Kimbrough et al., 2020*). To
determine the statistical significance of c-Fos expression differences, we computed the Z value for
each treatment condition—KET and ISO—by dividing the log relative c-Fos density by the standard
error. Positive Z values indicate higher values than the control, and negative Z values indicate lower
values than the control.

Network generation
To evaluate how functional connectivity changed under general anesthesia in WT mice, we ex-
tracted 63 brain regions from major brain subdivisions (CTX, CNU, TH, HY, MB, and HB) listed
in Figure 7A. Correlation matrices were generated by computing Pearson correlation coefficients
from interregional c-Fos expression in the 63 regions. Mean correlations were calculated to assess
changes in functional connectivity between these major subdivisions of the brain. Weighted undi-
rected networks were constructed by considering correlations with Pearson's $r \geq 0.82$, correspond-
ing to a one-tailed significance level of $P<0.05$ (uncorrected). The nodes in the networks represent
brain regions, and the correlations that survived thresholding were considered connections. Theo-
retical graph analysis was performed using Brain Connectivity Toolbox (https://sites.google.com/site/bctnet/
version 2019-03-03) in MATLAB R2021 (The MathWorks Inc.) (*Rubinov and Sporns, 2010*). Network
visualization was performed using Cytoscape (version 3.2.1) (*Shannon et al., 2003*).

Hub identification

Network centrality was evaluated using degree, betweenness, and eigenvector centrality measures to identify potential hub regions (*Sciarra et al., 2018*). Degree counts the number of edges connected to a node, signifying its immediate influence. Betweenness centrality is gauged by the number of shortest paths passing through a node, indicating its role as a connector within the network. Eigenvector centrality measures a node's influence by the centrality of its connections, valuing nodes linked to well-connected neighbors. High eigenvector centrality indicates significant influence through these high-quality connections within the network.

Statistical analysis

Sample size was determined based on prior studies (*Lu et al., 2008*; *Yatziv et al., 2020*). In Figure 2 and Figure 2–figure supplement 1, c-Fos densities in both experimental and control groups were log-transformed. Z-scores were calculated for each brain region by normalizing these log-transformed values against the mean and standard deviation of its respective control group. This involved subtracting the control mean from the experimental value and dividing the result by the control standard deviation. For statistical analysis, Z-scores were compared to a null distribution with a zero mean, and adjustments were made for multiple comparisons using the Benjamini–Hochberg method with a 5% false discovery rate (Q) (*Benjamini and Hochberg, 1995*). Pearson correlation coefficients (R) were transformed into Z-scores using Fisher's Z transformation before computing group means and making statistical comparisons in Figure 7–figure supplement 1D. All statistical analyses were conducted using GraphPad Prism 9.0 (GraphPad Software, USA) and MATLAB R2021 (Mathworks Inc.).

Acknowledgments

We express our gratitude to our interns, Chuhang Wong from Imperial College London and Jiale Huang from ShanghaiTech University, for their assistance in cell counting.

Funding

This study was funded by the National Natural Science Foundation of China (grants 82271292, 81730031 to Yingwei Wang; 82371286, 82101350 to Mengqiang Luo), the Shanghai Municipal Key Clinical Specialty (grant shslczdzk06901 to Yingwei Wang), and the Foundation of Shanghai Municipal Science and Technology Medical Innovation Research Project (23Y21900600 to Yingwei Wang).

Author contributions

Yue Hu, Conceptualization, Formal analysis, Investigation, Visualization, Methodology, Writing—original draft, Writing—review and editing; Wenjie Du, Conceptualization, Formal analysis, Visualization, Writing—original draft, Writing—review and editing; Jiangtao Qi, Conceptualization, Software, Formal analysis; Huoqing Luo, Formal analysis, Methodology, Writing—review and editing; Zhao Zhang, Resources, Investigation, Supervision, Writing—review and editing; Mengqiang Luo, Resources, Supervision, Funding acquisition, Project administration, Writing—review and editing; Yingwei Wang, Conceptualization, Resources, Supervision, Funding acquisition, Project administration, Writing—review and editing.

Competing Interest Statement

All authors declare no competing financial interest.

References

Adamantidis AR, de Lecea L. Sleep and the hypothalamus. Science. 2023; 382(6669):405–412. doi: 10.1126/science.adh8285.

Al-Hasani R, Gowrishankar R, Schmitz GP, Pedersen CE, Marcus DJ, Shirley SE, Hobbs TE, Elerding AJ, Renaud SJ, Jing M, Li Y, Alvarez VA, Lemos JC, Bruchas MR. Ventral tegmental area GABAergic inhibition of cholinergic interneurons in the ventral nucleus accumbens shell promotes reward reinforcement. Nat Neurosci. 2021; 24(10):1414–1428. doi: 10.1038/s41593-021-00898-2.

Bekkers JM, Suzuki N. Neurons and circuits for odor processing in the piriform cortex. Trends Neurosci. 2013; 36(7):429–38. https://www.ncbi.nlm.nih.gov/pubmed/23648377, doi: 10.1016/j.tins.2013.04.005.

Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B (Methodological). 1995; 57(1):289–300. http://www.jstor.org/stable/2346101.

Brown EN, Lydic R, Schiff ND. General anesthesia, sleep, and coma. N Engl J Med. 2010; 363(27):2638–50. https://www.nejm.org/doi/10.1056/NEJMra0808281?url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org&rfr_dat=cr_pub%3Dpubmed, doi: 10.1056/NEJMra0808281.

Brown E, Purdon P, Van Dort C. General anesthesia and altered states of arousal: a systems neuroscience analysis. Annual review of neuroscience. 2011; 34:601–28. https://www.annualreviews.org/doi/full/10.1146/annurev-neuro-060909-153200?url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org&rfr_dat=cr_pub%3Dpubmedhttps://www.ncbi.nlm.nih.gov/pmc/articles/PMC3390788/pdf/nihms315743.pdf, doi: 10.1146/annurev-neuro-060909-153200.

Chapuis J, Cohen Y, He X, Zhang Z, Jin S, Xu F, Wilson DA. Lateral entorhinal modulation of piriform cortical activity and fine odor discrimination. J Neurosci. 2013; 33(33):13449–59. doi: 10.1523/jneurosci.1387-13.2013.

Cox J, Witten IB. Striatal circuits for reward learning and decision-making. Nat Rev Neurosci. 2019; 20(8):482–494. doi: 10.1038/s41583-019-0189-2.

Cui Y, Huang X, Huang P, Huang L, Feng Z, Xiang X, Chen X, Li A, Ren C, Li H. Reward ameliorates depressive-like behaviors via inhibition of the substantia innominata to the lateral habenula projection. Sci Adv. 2022; 8(27):eabn0193. doi: 10.1126/sciadv.abn0193.

Daniel SE, Rainnie DG. Stress Modulation of Opposing Circuits in the Bed Nucleus of the Stria Terminalis. Neuropsychopharmacology. 2016; 41(1):103–25. doi: 10.1038/npp.2015.178.

Dehaene S, Changeux JP. Experimental and theoretical approaches to conscious processing. Neuron. 2011; 70(2):200–27. doi: 10.1016/j.neuron.2011.03.018.

Do JP, Xu M, Lee SH, Chang WC, Zhang S, Chung S, Yung TJ, Fan JL, Miyamichi K, Luo L, Dan Y. Cell type-specific long-range connections of basal forebrain circuit. Elife. 2016; 5. https://www.ncbi.nlm.nih.gov/pubmed/27642784, doi: 10.7554/eLife.13214.

Floresco SB. The nucleus accumbens: an interface between cognition, emotion, and action. Annu Rev Psychol. 2015; 66:25–52. doi: 10.1146/annurev-psych-010213-115159.

Franks NP. General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. Nat Rev Neurosci. 2008; 9(5):370–86. https://www.nature.com/articles/nrn2372.pdf, doi: 10.1038/nrn2372.

Gelegen C, Miracca G, Ran MZ, Harding EC, Ye Z, Yu X, Tossell K, Houston CM, Yustos R, Hawkins ED, Vyssotski AL, Dong HL, Wisden W, Franks NP. Excitatory Pathways from the Lateral Habenula Enable Propofol-Induced Sedation. Curr Biol. 2018; 28(4):580–587.e5. doi: 10.1016/j.cub.2017.12.050.

Giessel AJ, Datta SR. Olfactory maps, circuits and computations. Curr Opin Neurobiol. 2014; 24(1):120–32. doi: 10.1016/j.conb.2013.09.010.

Goutagny R, Luppi PH, Salvert D, Lapray D, Gervasoni D, Fort P. Role of the dorsal paragigantocellular reticular nucleus in paradoxical (rapid eye movement) sleep generation: a combined electrophysiological and anatomical study in the rat. Neuroscience. 2008; http://dx.doi.org/10.1016/j.neuroscience.2007.12.014http://www.ncbi.nlm.nih.gov/pubmed/18308473.

Grillner S, Hellgren J, Ménard A, Saitoh K, Wikström MA. Mechanisms for selection of basic motor programs–roles for the striatum and pallidum. Trends Neurosci. 2005; 28(7):364–70. doi: 10.1016/j.tins.2005.05.004.

Gu G, Varoqueaux F, Simerly RB. Hormonal regulation of glutamate receptor gene expression in the anteroventral periventricular nucleus of the hypothalamus. J Neurosci. 1999; 19(8):3213–22. doi: 10.1523/jneurosci.19-08-03213.1999.

Gupta RG, Schafer C, Ramaroson Y, Sciullo MG, Horn CC. Role of the abdominal vagus and hindbrain in inhalational anesthesia-induced vomiting. Auton Neurosci. 2017; 202:114–121. https://www.ncbi.nlm.nih.gov/pubmed/27396693, doi: 10.1016/j.autneu.2016.06.007.

Haber SN. Corticostriatal circuitry. Dialogues Clin Neurosci. 2016; 18(1):7–21. doi: 10.31887/DCNS.2016.18.1/shaber.

Han B, McCarren HS, O'Neill D, Kelz MB. Distinctive recruitment of endogenous sleep-promoting neurons by volatile anesthetics and a nonimmobilizer. Anesthesiology. 2014; 121(5):999–1009. doi: 10.1097/aln.0000000000000383.

Hase T, Hashimoto T, Saito H, Uchida Y, Kato R, Tsuruga K, Takita K, Morimoto Y. Isoflurane induces c-Fos expression in the area postrema of the rat. J Anesth. 2019; 33(4):562–566. doi: 10.1007/s00540-019-02662-0.

Hemmings J H C, Riegelhaupt PM, Kelz MB, Solt K, Eckenhoff RG, Orser BA, Goldstein PA. Towards a Comprehensive Understanding of Anesthetic Mechanisms of Action: A Decade of Discovery. Trends Pharmacol Sci. 2019; 40(7):464–481. doi: 10.1016/j.tips.2019.05.001.

Hua T, Chen B, Lu D, Sakurai K, Zhao S, Han BX, Kim J, Yin L, Chen Y, Lu J, Wang F. General anesthetics activate a potent central pain-suppression circuit in the amygdala. Nat Neurosci. 2020; 23(7):854–868. https://www.nature.com/articles/s41593-020-0632-8.pdf, doi: 10.1038/s41593-020-0632-8.

Huang X, Huang P, Huang L, Hu Z, Liu X, Shen J, Xi Y, Yang Y, Fu Y, Tao Q, Lin S, Xu A, Xu F, Xue T, So KF, Li H, Ren C. A Visual Circuit Related to the Nucleus Reuniens for the Spatial-Memory-Promoting Effects of Light Treatment. Neuron. 2021; http://dx.doi.org/10.1016/j.neuron.2020.10.023http://www.ncbi.nlm.nih.gov/pubmed/33171117.

Inta D, Trusel M, Riva MA, Sprengel R, Gass P. Differential c-Fos induction by different NMDA receptor antagonists with antidepressant efficacy: potential clinical implications. Int J Neuropsychopharmacol. 2009; 12(8):1133–6. https://www.ncbi.nlm.nih.gov/pubmed/19638253, doi: 10.1017/S1461145709990319.

Islam MT, Rumpf F, Tsuno Y, Kodani S, Sakurai T, Matsui A, Maejima T, Mieda M. Vasopressin neurons in the paraventricular hypothalamus promote wakefulness via lateral hypothalamic orexin neurons. Curr Biol. 2022; 32(18):3871–3885.e4. doi: 10.1016/j.cub.2022.07.020.

Jiang-Xie LF, Yin L, Zhao S, Prevosto V, Han BX, Dzirasa K, Wang F. A Common Neuroendocrine Substrate for Diverse General Anesthetics and Sleep. Neuron. 2019; 102(5):1053–1065.e4. doi: 10.1016/j.neuron.2019.03.033.

Kataoka N, Shima Y, Nakajima K, Nakamura K. A central master driver of psychosocial stress responses in the rat. Science. 2020; 367(6482):1105–1112. doi: 10.1126/science.aaz4639.

Kimbrough A, Lurie DJ, Collazo A, Kreifeldt M, Sidhu H, Macedo GC, D'Esposito M, Contet C, George O. Brain-wide functional architecture remodeling by alcohol dependence and abstinence. Proc Natl Acad Sci U S A. 2020; 117(4):2149–2159. https://www.ncbi.nlm.nih.gov/pubmed/31937658, doi: 10.1073/pnas.1909915117.

Lamme VA. Towards a true neural stance on consciousness. Trends Cogn Sci. 2006; 10(11):494–501. doi: 10.1016/j.tics.2006.09.001.

Lee U, Ku S, Noh G, Baek S, Choi B, Mashour GA. Disruption of frontal-parietal communication by ketamine, propofol, and sevoflurane. Anesthesiology. 2013; 118(6):1264–75. https://www.ncbi.nlm.nih.gov/pubmed/23695090, doi: 10.1097/ALN.0b013e31829103f5.

Leung LS, Luo T, Ma J, Herrick I. Brain areas that influence general anesthesia. Prog Neurobiol. 2014; 122:24–44. https://pubmed.ncbi.nlm.nih.gov/25172271, doi: 10.1016/j.pneurobio.2014.08.001.

Lu J, Nelson LE, Franks N, Maze M, Chamberlin NL, Saper CB. Role of endogenous sleep-wake and analgesic systems in anesthesia. J Comp Neurol. 2008; 508(4):648–62. https://onlinelibrary.wiley.com/doi/abs/10.1002/cne.21685, doi: 10.1002/cne.21685.

Lu J, Zhang YH, Chou TC, Gaus SE, Elmquist JK, Shiromani P, Saper CB. Contrasting effects of ibotenate lesions of the paraventricular nucleus and subparaventricular zone on sleep-wake cycle and temperature regulation. J Neurosci. 2001; 21(13):4864–74. doi: 10.1523/jneurosci.21-13-04864.2001.

Luo M, Fei X, Liu X, Jin Z, Wang Y, Xu M. Divergent Neural Activity in the VLPO During Anesthesia and Sleep. Adv Sci (Weinh). 2023; 10(2):e2203395. https://www.ncbi.nlm.nih.gov/pubmed/36461756, doi: 10.1002/advs.202203395.

Ma G, Liu Y, Wang L, Xiao Z, Song K, Wang Y, Peng W, Liu X, Wang Z, Jin S. Hierarchy in sensory processing
reflected by innervation balance on cortical interneurons. Science Advances. 2021; 7(20):eabf5676.

Magalhães KS, Spiller PF, da Silva MP, Kuntze LB, Paton JFR, Machado BH, Moraes DJA. Locus Coeruleus as a
vigilance centre for active inspiration and expiration in rats. Sci Rep. 2018; 8(1):15654. doi: 10.1038/s41598-
018-34047-w.

Mashour GA. Top-down mechanisms of anesthetic-induced unconsciousness. Front Syst Neu-
rosci. 2014; 8:115. https://fjfsdata01prod.blob.core.windows.net/articles/files/97687/pubmed-zip/
.versions/1/.package-entries/fnsys-08-00115/fnsys-08-00115.pdf?sv=2015-12-11&sr=b&sig=
DQhXzk4GgU12ckemlquDAU7hP2bisbegP0NFL7%2BjkWQ%3D&se=2020-07-20T11%3A30%3A05Z&sp=r&
rscd=attachment%3B%20filename%2A%3DUTF-8%27%27fnsys-08-00115.pdf, doi: 10.3389/fnsys.2014.00115.

Mashour GA, Hudetz AG. Bottom-Up and Top-Down Mechanisms of General Anesthetics Modulate Different
Dimensions of Consciousness. Front Neural Circuits. 2017; 11:44. https://www.ncbi.nlm.nih.gov/pubmed/
28676745, doi: 10.3389/fncir.2017.00044.

McGinty D, Szymusiak R. Keeping cool: a hypothesis about the mechanisms and functions of slow-wave sleep.
Trends Neurosci. 1990; 13(12):480–7. doi: 10.1016/0166-2236(90)90081-k.

Mickelsen LE, Bolisetty M, Chimileski BR, Fujita A, Beltrami EJ, Costanzo JT, Naparstek JR, Robson P, Jackson AC.
Single-cell transcriptomic analysis of the lateral hypothalamic area reveals molecularly distinct populations
of inhibitory and excitatory neurons. Nat Neurosci. 2019; 22(4):642–656. doi: 10.1038/s41593-019-0349-8.

Moore JT, Chen J, Han B, Meng QC, Veasey SC, Beck SG, Kelz MB. Direct activation of sleep-promoting VLPO
neurons by volatile anesthetics contributes to anesthetic hypnosis. Curr Biol. 2012; 22(21):2008–16. doi:
10.1016/j.cub.2012.08.042.

Morgan JI, Curran T. Stimulus-transcription coupling in neurons: role of cellular immediate-early genes. Trends
Neurosci. 1989; 12(11):459–62. doi: 10.1016/0166-2236(89)90096-9.

Nagata A, Nakao S, Miyamoto E, Inada T, Tooyama I, Kimura H, Shingu K. Propofol inhibits ketamine-induced
c-fos expression in the rat posterior cingulate cortex. Anesth Analg. 1998; 87(6):1416–20. https://www.ncbi.
nlm.nih.gov/pubmed/9842840, doi: 10.1097/00000539-199812000-00040.

Nagata A, Nakao Si S, Nishizawa N, Masuzawa M, Inada T, Murao K, Miyamoto E, Shingu K. Xenon inhibits but
N(2)O enhances ketamine-induced c-Fos expression in the rat posterior cingulate and retrosplenial cortices.
Anesth Analg. 2001; 92(2):362–8. doi: 10.1097/00000539-200102000-00016.

Nakao S, Arai T, Mori K, Yasuhara O, Tooyama I, Kimura H. High-dose ketamine does not induce c-Fos protein
expression in rat hippocampus. Neurosci Lett. 1993; 151(1):33–6. doi: 10.1016/0304-3940(93)90038-m.

Nakao S, Miyamoto E, Masuzawa M, Kambara T, Shingu K. Ketamine-induced c-Fos expression in the mouse
posterior cingulate and retrosplenial cortices is mediated not only via NMDA receptors but also via sigma
receptors. Brain Res. 2002; 926(1-2):191–6. doi: 10.1016/s0006-8993(01)03338-8.

Nelson LE, Guo TZ, Lu J, Saper CB, Franks NP, Maze M. The sedative component of anesthesia is mediated by
GABA(A) receptors in an endogenous sleep pathway. Nat Neurosci. 2002; 5(10):979–84. doi: 10.1038/nn913.

Pfarr S, Schaaf L, Reinert JK, Paul E, Herrmannsdörfer F, Roßmanith M, Kuner T, Hansson AC, Spanagel R, Körber
C, Sommer WH. Choice for Drug or Natural Reward Engages Largely Overlapping Neuronal Ensembles in the
Infralimbic Prefrontal Cortex. J Neurosci. 2018; 38(14):3507–3519. doi: 10.1523/jneurosci.0026-18.2018.

Rasiah NP, Loewen SP, Bains JS. Windows into stress: a glimpse at emerging roles for CRH(PVN) neurons.
Physiol Rev. 2023; 103(2):1667–1691. doi: 10.1152/physrev.00056.2021.

Reimann HM, Niendorf T. The (Un)Conscious Mouse as a Model for Human Brain Functions: Key Principles
of Anesthesia and Their Impact on Translational Neuroimaging. Front Syst Neurosci. 2020; 14:8. https:
//www.ncbi.nlm.nih.gov/pubmed/32508601, doi: 10.3389/fnsys.2020.00008.

Ren S, Wang Y, Yue F, Cheng X, Dang R, Qiao Q, Sun X, Li X, Jiang Q, Yao J, Qin H, Wang G, Liao X, Gao D,
Xia J, Zhang J, Hu B, Yan J, Wang Y, Xu M, et al. The paraventricular thalamus is a critical thalamic area for
wakefulness. Science. 2018; 362(6413):429–434. https://science.sciencemag.org/content/sci/362/6413/429.
full.pdf, doi: 10.1126/science.aat2512.

Rolls ET, Cheng W, Feng J. The orbitofrontal cortex: reward, emotion and depression. Brain Commun. 2020;
2(2):fcaa196. doi: 10.1093/braincomms/fcaa196.

Ross JA, Van Bockstaele EJ. The Locus Coeruleus- Norepinephrine System in Stress and Arousal: Unraveling Historical, Current, and Future Perspectives. Front Psychiatry. 2020; 11:601519. doi: 10.3389/fpsyt.2020.601519.

Roy DS, Kitamura T, Okuyama T, Ogawa SK, Sun C, Obata Y, Yoshiki A, Tonegawa S. Distinct Neural Circuits for the Formation and Retrieval of Episodic Memories. Cell. 2017; 170(5):1000–1012.e19. doi: 10.1016/j.cell.2017.07.013.

Rubinov M, Sporns O. Complex network measures of brain connectivity: uses and interpretations. Neuroimage. 2010; 52(3):1059–69. https://www.ncbi.nlm.nih.gov/pubmed/19819337, doi: 10.1016/j.neuroimage.2009.10.003.

Schroeder KE, Irwin ZT, Gaidica M, Nicole Bentley J, Patil PG, Mashour GA, Chestek CA. Disruption of corticocortical information transfer during ketamine anesthesia in the primate brain. Neuroimage. 2016; 134:459–465. doi: 10.1016/j.neuroimage.2016.04.039.

Sciarra C, Chiarotti G, Laio F, Ridolfi L. A change of perspective in network centrality. Sci Rep. 2018; 8(1):15269. doi: 10.1038/s41598-018-33336-8.

Seamans J. Losing inhibition with ketamine. Nat Chem Biol. 2008; 4(2):91–3. https://www.ncbi.nlm.nih.gov/pubmed/18202677, doi: 10.1038/nchembio0208-91.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome research. 2003; 13(11):2498–2504. https://pubmed.ncbi.nlm.nih.gov/14597658https://www.ncbi.nlm.nih.gov/pmc/articles/PMC403769/, doi: 10.1101/gr.1239303.

Skovsted P, Sapthavichaikul S. The effects of isoflurane on arterial pressure, pulse rate, autonomic nervous activity, and barostatic reflexes. Can Anaesth Soc J. 1977; 24(3):304–14. doi: 10.1007/bf03005103.

Smith ML, Li J, Cote DM, Ryabinin AE. Effects of isoflurane and ethanol administration on c-Fos immunoreactivity in mice. Neuroscience. 2016; 316:337–43. https://www.ncbi.nlm.nih.gov/pubmed/26742790, doi: 10.1016/j.neuroscience.2015.12.047.

Sun X, Bernstein MJ, Meng M, Rao S, Sørensen AT, Yao L, Zhang X, Anikeeva PO, Lin Y. Functionally Distinct Neuronal Ensembles within the Memory Engram. Cell. 2020; 181(2):410–423.e17. doi: 10.1016/j.cell.2020.02.055.

Suzuki M, Larkum ME. General Anesthesia Decouples Cortical Pyramidal Neurons. Cell. 2020; 180(4):666–676 e13. https://www.ncbi.nlm.nih.gov/pubmed/32084339, doi: 10.1016/j.cell.2020.01.024.

Vazey EM, Aston-Jones G. Designer receptor manipulations reveal a role of the locus coeruleus noradrenergic system in isoflurane general anesthesia. Proceedings of the National Academy of Sciences. 2014; 111(10):3859–3864. https://doi.org/10.1073/pnas.1310025111, doi: 10.1073/pnas.1310025111.

Wheeler AL, Teixeira CM, Wang AH, Xiong X, Kovacevic N, Lerch JP, McIntosh AR, Parkinson J, Frankland PW. Identification of a functional connectome for long-term fear memory in mice. PLoS Comput Biol. 2013; 9(1):e1002853. https://www.ncbi.nlm.nih.gov/pubmed/23300432, doi: 10.1371/journal.pcbi.1002853.

Wood SK, Valentino RJ. The brain norepinephrine system, stress and cardiovascular vulnerability. Neurosci Biobehav Rev. 2017; 74(Pt B):393–400. doi: 10.1016/j.neubiorev.2016.04.018.

Yang C, Naya Y. Sequential involvements of the perirhinal cortex and hippocampus in the recall of item-location associative memory in macaques. PLoS Biol. 2023; 21(6):e3002145. doi: 10.1371/journal.pbio.3002145.

Yap EL, Greenberg ME. Activity-Regulated Transcription: Bridging the Gap between Neural Activity and Behavior. Neuron. 2018; 100(2):330–348. https://www.ncbi.nlm.nih.gov/pubmed/30359600, doi: 10.1016/j.neuron.2018.10.013.

Yatziv SL, Yudco O, Dickmann S, Devor M. Patterns of neural activity in the mouse brain: Wakefulness vs. General anesthesia. Neurosci Lett. 2020; 735:135212. https://www.ncbi.nlm.nih.gov/pubmed/32593772, doi: 10.1016/j.neulet.2020.135212.

Yi T, Wang N, Huang J, Wang Y, Ren S, Hu Y, Xia J, Liao Y, Li X, Luo F, Ouyang Q, Li Y, Zheng Z, Xiao Q, Ren R, Yao Z, Tang X, Wang Y, Chen X, He C, et al. A Sleep-Specific Midbrain Target for Sevoflurane Anesthesia. Adv Sci (Weinh). 2023; p. e2300189. https://www.ncbi.nlm.nih.gov/pubmed/36961096, doi: 10.1002/advs.202300189.

Zhang D, Liu J, Zhu T, Zhou C. Identifying c-fos Expression as a Strategy to Investigate the Actions of General
Anesthetics on the Central Nervous System. Curr Neuropharmacol. 2022; 20(1):55–71. https://www.ncbi.nlm.
nih.gov/pubmed/34503426, doi: 10.2174/1570159X19666210909150200.

Zhao F, Meng X, Lu S, Hyde LA, Kennedy ME, Houghton AK, Evelhoch JL, Hines CDG. fMRI study of olfactory pro-
cessing in mice under three anesthesia protocols: Insight into the effect of ketamine on olfactory processing.
Neuroimage. 2020; 213:116725. doi: 10.1016/j.neuroimage.2020.116725.

Zhou K, Xu H, Lu S, Jiang S, Hou G, Deng X, He M, Zhu Y. Reward and aversion processing by input-defined
parallel nucleus accumbens circuits in mice. Nat Commun. 2022; 13(1):6244. doi: 10.1038/s41467-022-
33843-3.

Table 1. Figure 3—source data 1. Summary of prior studies on KET activated brain regions identified via c-Fos immunostaining.

Brain regions	Function	Treatment Regimen	Duration	Time of day	Effect	References
CeA	Pain Suppression	100 mg/kg and 10 mg/kgxylazine i.p.	2 h	/	Up-regulated	Hua T, et al.,(2020) (*Hua et al., 2020*)
SO	Natural sleep/General anesthesia	100 mg/kg with or without xylazine(10 mg/kg) i.p.	2 h	/	Up-regulated	Jiang-Xie LF, et al.,(2019) (*Jiang-Xie et al., 2019*)
pCC	/	100 mg/kg i.p.	2 h	/	Up-regulated	Nagata A, et al.,(1998) (*Nagata et al., 1998*)
pCC, RSC	/	100 mg/kg i.p.	2 h	/	Up-regulated	Nagata A, et al.,(2001) (*Nagata et al., 2001*)
RSP, CeA, Midline thalamic nuclei	Antidepressant	10 mg/kg i.p.	1.5 h	/	Up-regulated	Inta D, et al.,(2009) (*Inta et al., 2009*)
Cg, BF, Orx, TMN, vlPAG, LC, LHb, EW, midline thalamus	LC/vPAG engaged in analgesia	150 mg/kg; 60 mg/kg; 30 mg/kg i.p.	2 h	ZT3	Up-regulated	Lu J, et al.,(2008) (*Lu et al., 2008*)
CC, PVT	/	100 mg/kg i.p.	/	/		Nakao S, et al.,(1993) (*Nakao et al., 1993*)
PC, RSP	/	50 mg/kg i.p.	2 h	ZT3-9	Up-regulated	Nakao S, et al.,(2002) (*Nakao et al., 2002*)

Table 2. Figure 4—source data 1. Summary of prior studies on brain regions activated by ISO, as detected through c-Fos immunostaining.

Brain regions	Function	Concentration of ISO	Duration	Time of day		References
VLPO	Sleep-Promoting, Hypnosis	1.2%, 0.6%, 0.3%	2 h	ZT12-14/ZT4-6	Up-regulated	Moore JT,.et al.,(2012) (*Moore et al., 2012*)
PIR, LSD, LSV	/	4% 5mins+1%	2 h	ZT2-7	Up-regulated	Smith ML, et al.,(2016) (*Smith et al., 2016*)
NTS, VeN	Vomiting	1% and 3%	1.5 h	/	Up-regulated	Gupta RG, et al.,(2017) (*Gupta et al., 2017*)
CeA	Pain Suppression	1.5%	2 h	/	Up-regulated	Hua T, et al.,(2020) (*Hua et al., 2020*)
SO	Natural sleep, General anesthesia	1%,1.2%	2 h	/	Up-regulated	Jiang-Xie LF, et al.,(2019) (*Jiang-Xie et al., 2019*)
VLPO, MPO	Sleep and unconsciousness	1.2%	2 h	ZT14-17	Up-regulated	Han B, et al.,(2014) (*Han et al., 2014*)
EW, VLPO, LC	Sleep-Promoting, analgesia	2%	2 h	ZT12	Up-regulated	Lu J, et al.,(2008) (*Lu et al., 2008*; *Yi et al., 2023*)
AP	Nausea, vomiting	1.3%,2.6%	2 h	/	Up-regulated	Hase T, et al.,(2019) (*Hase et al., 2019*)

Figure 1—figure supplement 1. EEG and EMG power change after each treatment. The box represents the 25th-75th percentiles; the central line denotes the median; whiskers indicate maximum and minimum values. n = 6, 6, 8, 6 for the home cage, ISO, saline, and KET, respectively. (A) Normalized change in EEG power: ISO vs KET, p > 0.99; Saline vs KET, p = 0.01; Home cage vs ISO, p = 0.11. (B) Normalized change in EMG power: ISO vs KET, p = 0.36; Saline vs KET, p = 0.30; Home cage vs ISO, p = 0.02. Analyses were conducted using the Kruskal-Wallis test, followed by Dunn's multiple comparisons test.

Figure 2—figure supplement 1. KET and ISO induced c-Fos expression relative to their respective control group across 53 distinct brain regions. Z-scores for c-Fos expression in the KET and ISO groups were normalized to the mean and standard deviation of their respective control groups. Statistical analysis involved the comparison of Z-scores to a null distribution with a zero mean and adjustment for multiple comparisons using the Benjamini–Hochberg method at a 5% false discovery rate (*p < 0.05, **p < 0.01, ***p < 0.001). Brain regions are organized into major anatomical subdivisions, as indicated on the left side of the graph.

Figure 3—figure supplement 1. c-Fos expression in distinct brain regions following saline administration. Representative immunohistochemical staining of c-Fos-positive cells in the specified brain regions is provided as a control for the c-Fos expression observed in brain regions after KET exposure, as detailed in Figure 3. Scale bar: 200 μm.

Figure 4—figure supplement 1. c-Fos expression in home cage group. Representative immuno-histochemical staining of c-Fos-positive cells in the home cage group, serving as a baseline comparison for the ISO groups. Scale bar: 200 μm.

Figure 6—figure supplement 1. Region labels for the hierarchical clustering of the KET group in Figure 6A.

Figure 6—figure supplement 2. Region labels for the hierarchical clustering of the ISO group in Figure 6A.

Figure 7—figure supplement 1. Hierarchical clustering across different conditions. (A) Heatmaps depict the hierarchical clustering of brain regions within the home cage, ISO, saline, and KET groups, using Euclidean distance matrices calculated from correlation coefficients. Modules are demarcated within each dendrogram at a cut-off threshold of 0.7. (B) Silhouette scores are plotted against the dendrogram tree height ratio for each condition, with optimal cluster definition indicated by a dashed line at a 0.7 ratio. (C) The number of clusters formed at different cutoff levels. At a ratio of 0.7, ISO and saline treatments result in three clusters, whereas home cage and KET conditions yield two clusters. (D) The mean Pearson's correlation coefficient (r) was computed from interregional correlations displayed in Figure 7A. Data were analyzed using one-way ANOVA with Tukey's post hoc test, ***p < 0.001.

Figure 7—figure supplement 2. Hub region characterization across different conditions: home cage (A), ISO (B), saline (C), and KET (D) treatments. Brain regions are sorted by degree, betweenness centrality, and eigenvector centrality, with each metric presented in separate bar graphs. Bars to the left of the dashed line indicate the top 20% of regions by rank, highlighting the most central nodes within the network. Red bars signify regions that consistently appear within the top rankings for both degree and betweenness centrality across the metrics.