Allosteric activation and inhibition of glycogen phosphorylase share common transient structural features

Monika Kish[1], Dylan P. Ivory[1], Jonathan J. Phillips[1,2]

[1]Living Systems Institute, Department of Biosciences, University of Exeter, Stocker Road, Exeter, EX4 4QD, UK

[2]Alan Turing Institute, British Library, London, NW1 2DB, UK

ABSTRACT

It remains a major challenge to ascertain the specific structurally dynamic changes that underpin protein functional switching. Now with the routine ability to determine and predict accurate structural models of proteins with high resolution, there is a growing need to complement this with the ability to determine the structural changes that occur as these proteins are regulated and function. The archetypal allosteric enzyme, glycogen phosphorylase is one of the most studied proteins and is a clinical target of much interest to treat type II diabetes and metastatic cancers. However, a lack of understanding of its complex regulation, mediated by dynamic structural changes, hinder its exploitation as a drug target. Here, we precisely locate dynamic structural changes upon allosteric activation and inhibition of glycogen phosphorylase, by developing a time-resolved non-equilibrium millisecond hydrogen/deuterium-exchange mass spectrometry (HDX-MS) approach. We resolved obligate transient changes in localized structure that are absent when directly comparing active/inactive states of the enzyme and which are common to allosteric activation by AMP and inhibition by caffeine, operating at different sites. This approach has broad application to determine the structural kinetic mechanisms by which proteins are regulated.

INTRODUCTION

Allostery refers to the transfer of a signal between two sites of a protein resulting in a change in the enzymatic/catalytic activity or binding affinity to a ligand at a remote site of the protein. These subtle conformational and structural transitions are fundamental to protein functional regulation in receptor response[1], cell signalling[2], and metabolic regulation[3]. While allostery is typically associated with ligand binding to the allosteric site, which prompts the conformational and dynamic change at a distant site of the protein[4], covalent modification (e.g., phosphorylation)[5], or proteolytic cleavage[6] can also trigger the allosteric transition. Despite extensive studies since its inception[7-9], very little is known about the core dynamic process of allosteric communication and how this long-range signal and communication transmits between particularly remote protein sites. This is in large part due to the lack of biophysical approaches to measure at high resolution in both time and space. Recently, important advances have been made towards this goal in single molecule FRET[10,11,17,18], NMR[12], time resolved cryo-EM[13], time resolved crystallography[14], dynamic simulations[15] and Double Electron Electron Resonance (DEER)[19,20]. These begin to reveal non-equilibrium protein structural dynamics at high structural and temporal resolution, first described by Monod and colleagues in 1965[16]. These methods are impressive, though still require considerable adaptation per sample and are not yet broadly applicable to cases of ligand-induced allostery.

Glycogen phosphorylase (GlyP) is the archetypal allosteric enzyme whose regulation is tightly coupled to solid tumor metastasis, type II diabetes and adaptive immunity (early memory CD8+ T-cell recall response)[21]. GlyP catalyses the first committed energetic step of carbohydrate metabolism from glycogen stores and as such is one of the most highly regulated enzymes known, with at least six allosteric sites that sense natural ligands and drugs. GlyP has been intensively studied since the 1930s[22], thus there is abundance of structural and

kinetic data on this protein system[23-32], with detailed studies on the activation mechanism[33]
and stabilization of R-/T-state by allostery[34] and phosphorylation[35]. GlyP is, therefore, an ideal
system to challenge our ability to discern mechanisms of allosteric regulation as it is alternately
activated by AMP at the 'nucleotide site' and inhibited by caffeine at the 'inhibitor site'[36,37],
whilst also being constitutively active by Ser 14 phosphorylation[16]. Binding of AMP at the
allosteric effector site can bring similar changes as Ser 14 phosphorylation, but these are
believed to act by different mechanisms, supported by the fact that GlyPa can be further
activated by AMP binding[38]. How a single protein is able to support multiple such conflicting
structural pathways to underpin allosteric activation and inhibition has not been elucidated.

Hydrogen-deuterium exchange mass spectrometry (HDX-MS) has several advantages and
can infer hydrogen-bonding networks which are steered by local structural dynamics and to a
smaller measure the solvent accessibility in large complexes up to a single amino acid
resolution. We aimed to exploit these advantages to compare and provide differential analysis
of data from an equilibrium and non-equilibrium experiment upon activation/inhibition with
allosteric modification. Conventionally proteins are mixed with a ligand and allowed to reach
equilibrium, here, we used a non-equilibrium millisecond HDX approach to identify specific
dynamic changes associated with ligand binding and long-range allosteric communication.
Comparing the difference in D labeling between the states provides highly structurally resolved
determination of dynamic reconfiguration during allosteric regulation. Equilibrium states reflect
local minima, while non-equilibrium uptake data reveal transitions between local minima on
the conformational landscape[39]. We were able to clearly identify amino acids that go through
conformational dynamical changes and the nature of those changes upon allosteric ligand
binding – an approach that could be used broadly to identify such features in almost any
ligand-induced allosteric transition of a protein.

RESULTS
73

Non-equilibrium allosteric activation/inhibition of GlyP
74

To observe the kinetically populated transition state of glycogen phosphorylase activation and
inhibition, we first established the chemical conditions that fully activate or inhibit the enzyme.
The catalytic activity for the fully active form phosphorylated at Ser14 (GlyPa) was used as a
positive control reference in a glycogen hydrolysis colorimetric assay (**Figure 1A,E**; Error!
Reference source not found.**).** Against this level, the maximal extent of GlyPb allosteric
activation by AMP was then established[24,38]. This showed that full activation is achieved with
AMP ≥25 mM, the lowest concentration of AMP tested (range 25-100 mM), given no significant
difference between the activity of GlyPb+AMP compared with GlyPa (**Figure 1A,E**). We also
sought to verify the requirement of ammonium sulfate (AS) for full activation, known to mimic
the presence of phosphate by binding GlyP at the AMP effector site, catalytic site and Ser14
phosphorylation site[16]. There are conflicting reports on the requirement of AS for full activation
of GlyPb cooperatively with AMP: whilst GlyPb affinity for AMP is reported to increase 50-fold
and activity to increase by 50% with sulfate[23,40-42], here we found activation by sulfate alone
was not achieved (15% of maximum), nor was a cooperative effect observed on activity
(**Figure 1E,C** and **Figure S2**). However, there was structural cooperativity between AMP and
AS, as evidenced by concentration-dependent changes in hydrogen-exchange kinetics for
amino acids site-localized to the nucleotide site and tower helix (**Figure 1D**). Local structural
perturbations were observed for GlyPb samples supplemented with 0-100 mM AMP and/or 0-
100 mM AS by measuring HDX-MS at equilibrium. Binding of AMP at the nucleotide site
results in local protection against D-labeling that is positively cooperative with AS (**Figure 1D**).
Corresponding changes were observed in the tower helix, which shows only weak protection
with AS alone, but stronger protection with AMP (similar at all AMP concentrations tested) that
increases with [AS] (**Figure 1D**). The 380 loop shows cooperativity between AS and AMP,
resulting in deprotection against D-labeling as the loop becomes disordered in the active
state[16] (**Figure 1D**). This identified cooperativity in GlyPb structural activation by AMP/AS that
is not readily observed in the activity assay (**Figure 1D,E**)[23,38,43]. One-way ANOVA tests

proved that higher concentrations of AMP or AS were not significant for either catalytic activity
or structural perturbation (**Supplementary table 2**): Therefore, the lowest concentrations
tested of 25 mM AMP and 25 mM AS were considered to fully activate GlyPb catalytically and
structurally and these conditions were used for the experiments to study non-equilibrium
activation of GlyPb. Caffeine inhibits the active GlyPa R-state, stabilising the T-state, therefore
we also sought to determine the minimal caffeine concentration required to fully inhibit 12 μM
GlyPa – the concentration for subsequent non-equilibrium HDX-MS experiments. This was
achieved with 32 mM caffeine, given by equivalence in specific activity to inactive GlyPb by
ANOVA these conditions were used for the experiments to study non-equilibrium inhibition of
GlyPa (**Figure 1F**).

In order to observe any kinetically populated transient structural ensembles, it is critical for the
non-equilibrium experiments that the protein:ligand complex is fully saturated within the dead
time of the experiment (here 50 ms), governed by the second-order rate constant of
association with units $M^{-1}s^{-1}$. While T-state GlyPb is allosterically activated by the binding of
AMP at the nucleotide site, active phosphorylated R-state GlyPa is allosterically inhibited by
the binding of caffeine between the 280s active site gate loop and the L40 loop. Therefore,
the degree of AMP saturation at the nucleotide site was determined by protection against D-
labeling at amino acids T38-T47, whereas the degree of caffeine saturation was determined
by protection against D-labelling at amino acids N274-F285. In separate experiments, a range
of AMP concentrations (2.5, 25 and 100 mM) and a range of caffeine concentrations (3, 10
and 32 mM) were assessed by rapid mixing with GlyPb or GlyPa, respectively, under non-
equilibrium conditions (**Figure 1G-H,** Error! Reference source not found.**).** To confirm that a
1:1 complex with ligand had been reached, these were compared with the end-state of the
respective process: activated GlyPb fully-equilibrated with 25 mM AMP, or inhibited GlyPa
fully-equilibrated with caffeine. After 50 ms of incubation with ligand in deuterated buffer, 32
mM of caffeine achieved equivalent protection against D-labelling compared with the fully-
equilibrated inhibited state (GlyPa + 32 mM caffeine) within the 50 ms dead-time of the HDX-
MS experiment. The case of GlyPb activation was more complicated, as it was important to
consider the contribution of both AMP and AS to ternary complex formation under these rapid-
mixing conditions. Incubation of GlyPb with AS fully equilibrated (*AS) or at non-equilibrium
(^AS) showed a lack of protection against D-labeling relative to GlyPb alone, signifying
absence of binding to the putative sites for AS interaction. 25 mM AMP alone did not fully bind
in 50 ms, whereas full protection is achieved within error for the non-eq addition of 25 mM
AMP and AS. This confirms that the presence of AS enhances the k_a for GlyP:AMP association
and these conditions yield full complex formation inside the dead time of the non-eq HDX-MS
experiment. As we and others have recently reported, the presence of salts can significantly
alter the intrinsic rates of HDX, thus masking – or artefactually creating – protein structural
dynamics that are inferred by the HDX data[44]. This is not a systemic alteration of HDX kinetics
by the presence of AS, but rather it is confirmed to be a local effect at the nucleotide site, as
peptides covering other regions of the protein in the same HDX-MS data set do not show the
same behavior: for example the tower helix (reported by peptide segment V266-I275) shows
the opposite relationship with the k_{obs} for D-labeling being significantly faster in the presence
of 25 mM AS/AMP under equilibrium and non-eq conditions (**Figure 1I,** Error! Reference
source not found.). We concluded that the ligand binding sites for inhibition and activation of
GlyP by caffeine and AMP, respectively, are achieved within the dead-time of the HDX-MS
experiment at times post-mixing >= 50 ms. Therefore, we proceeded to evaluate the transient
structural kinetics that occur upon allosteric regulation across the whole enzyme.

Transient structural dynamics during GlyPb allosteric activation by AMP

To identify specific dynamic changes associated with long-range allosteric communication
induced by ligand binding, we developed a non-equilibrium approach to millisecond
hydrogen/deuterium-exchange that reveals the local structural perturbations that result from
the conditions established above (**Figure S1**). As the D-labeling of polypeptide backbone
amide groups is a sensitive, but convoluted measure of H-bonding and solvent accessibility,

first we sought to categorize the local differences in the protein ensemble during the non-
equilibrium phase following allosteric activation by AMP/AS. Subtle changes in dynamics and
structural changes can be derived from HDX-MS data, reflected in the observed kinetics of
deuterium uptake plots at specific positions within the protein, given by short peptide segments
in the 'bottom-up' experiments used here[39]. We calculated the local hydrogen exchange rate
(k_{obs}) by fitting HDX-MS data to a stretched exponential model[44]. The data comprised
measurements at 31 D-labeling time points over four orders of magnitude from 50 ms – 300 s
for 219 peptide segments that cover 90.5% of the GlyP amino acid sequence (Error!
Reference source not found.). This was done for three conditions: (i) inactive GlyPb (apo), (ii)
fully activated GlyPb (termed GlyPb*) equilibrated for one hour with 25 mM AMP, 25 mM AS
and (iii) GlyPb activated at non-equilibrium by rapid mixing with 25 mM AMP, 25 mM AS
(termed GlyPb^). It is clear that for many peptides in a variety of regions of the protein the D-
labeling kinetics under non-eq conditions are qualitatively different from both the apo and the
activated states (Error! Reference source not found.**-6** and Error! Reference source not
found.). There are peptides for which the D-labeling values fall outside of the range measured
for either the initial (apo) or equilibrated active (GlyPb*) forms, which indicates a degree of
non-linearity in the structural interpolation (**Figure 3**). However, the sub-molecular location,
magnitude and nature of relative (de)protection of these show a variety of differences.
Therefore, we next sought to classify these structural dynamic changes occurring during
allosteric activation to better identify clear behaviors in each part of the enzyme.

To categorize the local structural changes that occur during non-equilibrium
activation/inhibition into correlated transient dynamics, we developed a simple, but robust
quantitative analysis of the HDX-MS data that served to represent the non-equilibrium HDX
kinetics for the pool of peptide segments and to act as input to a clustering algorithm[45,46]. Each
peptide segment of the protein was represented spatially in a manner that reflects the
relationship of its non-equilibrium state to the starting and ending states of the protein (Error!
Reference source not found.**A**, Error! Reference source not found.). We identified the
archetypes of non-equilibrium structural kinetics that might be possible to observe in this plot
in Error! Reference source not found.. In order to categorize the continuum of diverse D-
labeling kinetic differences that we observed, we performed a k-means clustering analysis to
identify amino acids with correlated HDX kinetics. (Error! Reference source not found.). Seven
clusters of amino acids within GlyP comprises a common structural response to activation by
AMP/AS (**Figure 2**). **Figure 3** shows the D-labeling uptake plot for a representative peptide
segment from each cluster, alongside the structural location of the cluster.

The hierarchical relationship between the seven clusters was determined based on the
observed transient structural changes (**Figure 2A**)[47]. This reveals that there are two closely
related clusters (**a** and **g**) that show little or no observable transient structural change during
allosteric activation (i.e. close to the origin on **Figure 2A**). For most peptides, k_{obs} was low (k_{obs}
$< 1 \times 10^{-5}$ s^{-1} for 130 of 219 peptides measured) in all three conditions (Error! Reference source
not found.Error! Reference source not found.), as GlyP is highly structured with many amino
acids buried in the core of the enzyme with low solvent accessible surface area (SASA) and
stable H-bond networks, consistent with previous studies[16,28,44,48-51]. This validates that much
of the enzyme is structurally identical before, during and after allosteric activation, Error!
Reference source not found. and Error! Reference source not found.. A third related cluster
(**e**), representing the nucleotide binding site for AMP, shows no deviation from the active state
and only transient relative protection compared to the inactive state that is on-pathway from
the apo- to AMP-bound state (i.e. on y=0 in **Figure 2A**). This group includes residues from α2
(residues 47-78), the cap' region (residues 38-47) and residues from the nucleotide binding
domain (Error! Reference source not found.). The α8 helix is also essential for AMP binding,
but unfortunately our coverage was lacking peptides in that region. There are two closely
related clusters which represent the 250' loop (**f**) and the tower helix (**c**) that show transiently
increased D-labeling compared to one or both of the equilibrium states. There are also two
related clusters, representing the pyridoxal phosphate cofactor binding residues (**b**) and the
606 loop that are distinct from all others which show protection against D-labeling compared

to both the inactive and the active state. The amino acids surrounding the PLP cofactor retain
apo GlyPb HDX kinetics, indicating either that this region has a structural transition with a low
probability or that the induced protection against HDX occurs too slowly to prevent the
observed labeling.

A feature of the HDX-MS data during non-equilibrium AMP activation is that there are transient
structural changes that are distinct from both the inactive/active states (i.e. amino acids that
do not lie on either x=0 or y=0 in **Figure 2A**) and therefore are not represented by high-
resolution structural models derived from the equilibrium states. Activation of GlyPb by AMP
binding to the nucleotide site induces the well-established structural changes in the 250' and
280s loops, together with rotation of the tower helix (α2). The non-equilibrium HDX-MS reveals
previously unobserved transient changes that facilitate this. The 250' loop (A248-G260)
exhibits unique behavior during allosteric activation (cluster **f** in **Figure 3)**. The D-labeling
kinetics interpolate smoothly between the GlyPb state (at t=50 ms) and the active GlyPb* state
(at t=10 s). This indicates a likely two-state process for helical lengthening and that a gain of
structure has fully equilibrated in the molecular population by 10 s of D-labeling, consistent
with the tower helix in the active state lengthening at the N-terminal end by one turn in the
GlyPb* state (3E3N.pdb[28]). Clusters **c** (the tower helix) and **d** (the 606 loop) show opposite
structural behaviors: the 606 loop shows transient protection (increased H-bonding; reduced
solvent accessibility), whereas the tower helix shows transient deprotection (i.e. local
unfolding). The tower helix (G261-S276) is a contiguous alpha helical structure in all known
experimental models of apo, active and inactive forms of GlyP. However, unexpectedly here
we observe that it transiently loses protection against hydrogen/exchange, with increased
observed D-labeling during allosteric activation following AMP binding at the nucleotide site
(Error! Reference source not found.**A**).

We sought an atomistic rationale for this transient unfolding, so an all-atom interpolation
method was used to generate structures on a non-linear, energy minimized trajectory between
apo GlyPb and GlyPb*. The Climber method has been shown to sample intermediate
structures effectively, though has been difficult to validate until now, owing to the experimental
challenge of determining transient local structural changes[52]. Here a trajectory was created
which interpolates between two crystal models that have intact tower helices with predicted
hydrogen-bonds in conventional geometries, yet several of these hydrogen-bonds are shown
to break and then later reform as the helix refolds and rotates by ~10° to adopt the established
active R-state, GlyPb* (**Figure 4**; **Supplementary movies**)[53]. The interpolation is consistent
with the non-equilibrium HDX-MS data in which there is relative deprotection compared to both
GlyPb and GlyPb* precisely at D268-R269, resolved by multiple overlapping peptide
segments in the data, supporting an interpretation that there is a transient unfolding in the
tower helix during allosteric activation.

Transient structural dynamics during GlyPa allosteric inhibition by caffeine

We next considered whether this transient unfolding of the tower helix seen during activation
is a feature common to the GlyP inhibition pathway. GlyPa phosphorylated at serine14 is
basally active but is itself allosterically inhibited by binding of caffeine, which stabilizes a T-
state that is similar to the GlyPb apo state. The caffeine binding site is formed between the
280s and 606 loops, 3.5 nm from the nucleotide site where AMP binding stabilizes the reverse
transition to the R-state. Therefore, we next sought to identify the transient structural features
on the inhibition pathway by performing non-equilibrium HDX-MS of GlyPa in the presence of
caffeine. As for the AMP-activation experiments, this comprised measurements at 31 D-
labeling time points 50 ms – 300 s resulting in 171 peptide segments that cover 91.4% of the
GlyP amino acid sequence (Error! Reference source not found.). This was done for three
conditions: (i) active GlyPa (apo), (ii) fully inhibited GlyPa (termed GlyPa') equilibrated for one
hour with 32 mM caffeine and (iii) GlyPa inhibited at non-equilibrium by rapid mixing with 32
mM caffeine (termed GlyPa^).

From a comparison of the observed HDX kinetics between GlyPa^ and each of the two
equilibrated states, it is clear that the inhibition pathway involves a number of amino acids that
show a transient structure distinct from both the start/end states (**Figure 2C;** Error! Reference
source not found.). Based on this, a clustering analysis identified amino acids with correlated
HDX kinetics (Error! Reference source not found.). As for AMP-activation, caffeine-inhibition
resulted in seven clusters of amino acids within GlyP with a common structural response to
ligand (**Figure 2C; Figure 3B**). Again, much of the protein is relatively unchanged before,
during and after allosteric inhibition (clusters **h** and **j**, **Figure 2C-D**), which is consistent with
the available crystal structures. The 280s and 606 loops form the binding site for caffeine
which intercalates between them in the GlyPa T-state crystal model. These amino acids are
co-located in cluster **l** which shows HDX equivalent to the GlyPa' and protected compared to
the GlyPa states, consistent with direct binding at this site without any observed transitional
behavior (cluster **l, Figure 2C-D**). During caffeine-inhibition these three clusters (**h/j/l**) have
minimal transient structural features and are found to be hierarchically unrelated to all other
amino acids in the other four clusters (**Figure 2C**). During caffeine-inhibition, the 250' loop is
highly deprotected against hydrogen-exchange and labels more than the initial GlyPa state,
but similarly to the inhibited GlyPa' state (cluster **k**). This is consistent with an interpretation
that it is a two-state process, where the caffeine-inhibited T-state is observed immediately
upon binding, in which one turn of the tower helix unwinds to extend the 250' loop, as seen in
the crystal models. The tower helix itself (cluster **m**) is labeled more with deuterium under non-
equilibrium conditions (GlyPa^) than in either GlyPa or GlyPa' states. As this is a highly
solvent-exposed alpha helix in the initial R-state, it is most consistent with an interpretation of
transient unfolding, again resolved by overlapping peptide segments to be most pronounced
at amino acids D268-R269. The 280s loop shows a bifurcated behavior, with the more C-
terminal part forming part of the direct ligand binding site (cluster **l**) and with the N-terminal
stretch showing transiently reduced hydrogen-exchange compared to the initial apo GlyPa
state but increased when compared to GlyPa'. This is consistent with the initial binding of
caffeine to F285 and a subsequent propagation of induced helical structure further upstream
in the 280s loop as the tower helix lengthens by one turn at the C-terminal end.

So, it is clear that a number of highly resolved transient structurally dynamic features are
observed by non-equilibrium HDX-MS upon allosteric inhibition of GlyP by caffeine.

Transient unfolding of the tower helix is common to allosteric activation and inhibition
pathways

We next sought to compare the transient structural features observed upon AMP-activation
and caffeine-inhibition to determine whether these different ligands (and different allosteric
sites) appear to act via different pathways, or the same one.

The 250' and 280s loops show opposite neHDX-MS behaviour in response to AMP and
caffeine, which is logical given that these motifs are structurally different in the R/T-states and
necessarily interpolate between them. However, there is an unexpected common feature
during both allosteric activation and inhibition with transiently increased HDX observed in the
tower helix that sits in between these two loops. Non-linear interpolation between the atomistic
models for GlyPa R-state (apo) and caffeine-bound (T-state) also predicts breakage, then
reforming, of hydrogen-bonds in the tower helix, as it does for AMP-activation. It also predicts
that the two structural pathways are closely related with low rmsd in the 250' loop, tower helix
and 280s loop between structures close to the T-state and much larger differences in rmsd
close to the R-state (Figure 4B-C). As the tower helix is highly solvent-exposed, the transiently
lower protection factor would likely be the result of H-bond breakage which is known to be a
major determinant of hydrogen-exchange rate. This is suggested by the structural
interpolation, which predicts these hydrogen-bonds in the tower helix to be volatile in both
pathways.

These are the major transient structural features observed by non-equilibrium HDX-MS and
appear common to both pathways, with a consistent atomistic rationalization given by the
Climber structural interpolations.

DISCUSSION

Transient features of allosteric pathways revealed

Here we show that allosteric regulation of the central metabolic enzyme, glycogen
phosphorylase, involves a number of dynamic changes in local structure that are not observed
in static structural models of the R- and T-states. We established a new protocol exploiting
millisecond hydrogen/deuterium-exchange at non-equilibrium to characterize transient
changes in protection in response to ligand, resolved at near amino acid level throughout a
194 kDa enzyme. This revealed that AMP-activation induces a transient increase in HDX in
the tower helix. We rationalized this by interpolating the known structures for this allosteric
transition using an atomistic method based on the Energy Calculation and Dynamics (ENCAD)
molecular-mechanics force-field, Climber. This predicts that the well-established lengthening
and 10° rotation of the tower helix by first shortening, then bending it, wholly consistent with
the increased hydrogen-exchange observed during the process. Unexpectedly, caffeine-
inhibition – binding to a different allosteric site 3.5 nm away from AMP - also resulted in a
transient increase in HDX in the tower helix. The structural interpolation predicts that this is
due to a similar extent of transient breaking of hydrogen-bonds in the tower helix. So, both
experiments are consistent with the interpretation suggested by the simulation that breaking
of the tower helix is a common requirement of both AMP-activation and caffeine-inhibition
pathways.

Slow (millisecond) labeling can observe transient structural dynamics of the molecular population

It is perhaps unexpected that differences in non-equilibrium HDX-MS kinetics were observed
across a range of time domains from milliseconds to seconds. Structural kinetics were
observed even at the fastest labeling time of 50 ms (**Error! Reference source not found.**),
in several significant regions, including: 250 ɣ turn loop, β3 and α6, 280s loop and another ɣ
turn with the central residue Tyr 613 said to be forming a hydrophobic sandwich of the
nucleotide inhibitor site[54]. At slower deuterium labeling times, significant changes in protection
between the three states was observed throughout the α7 tower helix (residues 261-274), the
β11b (residues 276-279), α23 (residues 714-725) and α24 (residues 728-735). We interpret
this range of observed non-equilibrium structural kinetics to indicate that the probability of the
local structural changes is very low (i.e. bounded by a high energy transition state), resulting
in observation of the populated transient species (presumably transition states), even
milliseconds after ligand binding. The wide range of local protection factors against hydrogen-
exchange (due to existence of stable – or transient - local structure) would then effectively
stretch the observation of structural changes across multiple orders of magnitude, as seen
here.

We believe that this approach, while critically important to establish rapid saturation of the
ligand binding site, provides a straightforward way to directly observe allosteric structural
dynamics at non-equilibrium which can be applied broadly to proteins irrespective of size or
stability and with, in principle, any chemical or protein ligand. In this manner, we expect it to
be important in future to understand not only what has changed in protein structure, but how
the changes come about and propagate through the molecule.

METHODS

Sample preparation

356 Glycogen phosphorylase b and glycogen phosphorylase a were dissolved in phosphate buffered saline (PBS), pH
357 7.4 to 10.3 pmol/µL. For the equilibrium and non-equilibrium activated samples, the required amount of AMP/AS
358 and caffeine were added, in the stock solution or the labelling buffer, respectively. During the screening experiments

359 (HDX screen and enzyme activity assay) each sample was prepared simultaneously, with 10.3 pmol/μL
360 GlyPb/GlyPa and with varying concentrations of AMP, AS or caffeine, **Error! Reference source not found.**.

Screening of conditions to activate GlyP by enzyme activity assay

The enzyme activity of GlyPb and GlyPa was screened for AMP/AS and caffeine dependence**Error! Reference**
source not found.. A colorimetric assay kit was used (ab273271), with only one adjustment of the protocol for
GlyP sample preparation. A suitable protein concentration was determined by pre-testing a range of concentrations
and plate reader settings, then all 25 samples were prepared in parallel at 1 mg/mL GlyP (**Error! Reference source**
not found.). Three relevant concentrations of caffeine were screened (3, 10 and 32 mM) to determine the extent
of inhibition. Reaction kinetic curves and specific activity (mU/mG) were plotted and calculated according to the
suggested protocol.

Screening of conditions to activate GlyP by HDX-MS

The observable perturbations in HDX were screened for AMP/AS dependence. A CTC PAL sample handling robot
(LEAP Technologies, USA) was used to mix samples (**Error! Reference source not found.**) with deuterated PBS
buffer in 1:20 ratio, then quench the labeling reaction by 1:1 mixing with 100 mM potassium phosphate to a final
pH of 2.55 at 0 °C. Samples were immediately digested online with an Enzymate immobilized pepsin column
(Waters) at 12 °C, the derived peptides trapped on a VanGuard 2.1 x 5 mm ACQUITY BEH C18 column (Waters)
for 3 minutes at 125 μL/min at 0.5 °C and separated on a 1 × 100 mm ACQUITY BEH 1.7 μm C18 column (Waters)
with a 7 min linear gradient of acetonitrile (5-40%) supplemented with 0.1% formic acid. Peptides were eluted into
a Synapt G2-Si mass spectrometer (Waters).

Ammonium sulfate influence and maximum velocity kinetics

The maximum velocity kinetics and the AS influence on the HDX were determined prior continuing with the non-
equilibrium experiments. HDX labelling experiments were performed at 11 time points, ranging from 50 ms to 300
s. AMP activation was tested at 3 different concentrations, including 2.5, 25 and 100 mM under non-equilibrium
conditions with 25 mM AS. The AS influence on the equilibrated sample and the pre-equilibrated AS sample, was
also determined. Analyzed protein states include: GlyPb - apo, * - equilibrium activated with 25 mM AMP and 25
mM AS, *AS - equilibrium activated with 25 mM AS, ^ eq AS – equilibrated with 25 mM AS and then non- equilibrium
activated with AMP, ^ - non-equilibrium activated with 25 mM AMP and 25 mM AS, ^AMP – non-equilibrium
activated with 25 mM AMP, ^AS – non- equilibrium activated with 25 mM AS.

Hydrogen-deuterium exchange mass spectrometry

HDX was performed using a fully-automated, millisecond HDX labelling and online quench-flow instrument, ms2min
(Applied Photophysics Ltd), connected to an HDX manager (Waters). During the labelling experiments, 14 μL of
either apo or equilibrated GlyPb/GlyPa were delivered to the labelling mixer. A 20-fold dilution with labelling buffer
at 20˚C commenced the HDX. Depending on the desired experiment, the labelling buffer differed significantly. For
apo and equilibrium experiments, the labelling buffer was 1 x PBS, pH_{read} =7.00 at 20°C. During non-equilibrium
experiments and other assessments (see Error! Reference source not found.), the labelling buffer consisted of 1 x
PBS, pH =7.00 at 20°C combined with 25 mM AMP/25 mM AS or 32 mM caffeine. All GlyPb samples were labelled
at 30 time points, ranging from 50 ms to 300 s. All GlyPa samples were labelled at 20 time points, ranging from 50
ms to 300 s. Immediately after, the HDX reaction was quenched by mixing with quench buffer (100 mM Potassium
phosphate, pH =2.5 at 0˚C) at 1:1 ratio, and sample digested online with an Enzymate immobilized pepsin column
(Waters), the derived peptides trapped on a VanGuard 2.1 x 5 mm ACQUITY BEH C18 column (Waters) for 3
minutes at 125 μL/min and separated on a 1 × 100mm ACQUITY BEH 1.7 μm C18 column (Waters) with a 7-
minute linear gradient of acetonitrile (5-40%) supplemented with 0.1% formic acid. Peptides were eluted into a
Synapt G2-Si mass spectrometer (Waters). Mass spectra were obtained using the Waters HDMS[E] mode within a
mass range of 50 to 2000 m/z. Instrument parameters were configured as follows: a capillary voltage of 3.0 kV, a
cone voltage of 50 V, a trap collision energy of 4 V, a traveling wave ion mobility separation at a velocity of 475
m/s, a wave amplitude of 36.5 V, and a nitrogen pressure of 2.75 mbar. For low-energy scans, a transfer collision
energy of 4 V was applied, while high-energy scans utilized four separate collision energy ramps ranging from 15
to 55 V.

Data and statistical analysis

PLGS (ProteinLynx Global Server 2.5.1, Waters) was used for the analysis of MS[E] reference data and identification
of all discoverable peptic peptides. DynamX 3.0 (Waters, USA) was used for processing and assignments of
isotopic distributions of all raw data files. All HDX-MS experiments were performed in triplicate (technical
replicates). The means and standard deviations (SD) of these replicates were used to determine the significance
of changes, global significance threshold, as previously described[51]. In-house developed scripts in Python 3.4.0
(Python Software Foundation) were used for all post-processing analyses, clustering analyses, and plotting in
PyMOL (Schrödinger, Inc.).

A clustering analysis was employed to investigate the behavior of the peptide during non-equilibrium
activation/inhibition. The D-labeling data for each peptide segment was normalized to the theoretical maximum, to
account for varying peptide length and presence of proline for which there is no signal. The sum of deuterium

uptake over all time points for each peptide was calculated for all three protein conditions (states) of GlyPa or
GlyPb, in each case. The values for the non-eq data were then subtracted from both other states for every peptide
segment in the data set, resulting in an [n x 2] vector of 2-d coordinates. With the non-equilibrium measurements
defined as 2-d coordinates relative to a start/end state, this constitutes a well-formed input for a k-means clustering
analysis. The 2-d coordinates for HDX-MS differences were then *k-means* clustered with *k* clusters indicated to be
optimal by both an elbow method and a Silhouette analysis.

Non-linear atomistic structural interpolation

The Climber program was utilized for stepwise and non-linear structural interpolation between the active and
inactive conformations using default paramters. This involved dynamic evaluation of RMSD between backbone
and side chain atom pairs, with continual updates. Atom pair sets were created for each generated conformation
between the initial (A) and final (B) states, based on criteria of distance between alpha carbons (>10 Å) and side
chain atom pairs (<10 Å). Harmonic restraints were applied to steer inter-residue distances towards the target
conformation. The force constant for conformational transformation was adjusted based on RMSD progress.
Morphed conformations underwent iterative minimization using the ENCAD potential energy function. Initial
coordinates for the allosteric activation and inactivation were taken from crystallographic structures of glycogen
phosphorylase b and glycogen phosphorylase a. PDB ID 1gpb and 3e3n were used for GlyPb transition, and PDB
ID 1gpa and 1gfz for GlyP inactivation. Missing residues (251-259, 315-324 in 1gfz; 251-260, 281-287 in 3e3n)
were built and optimized using MODELLER (version 10.4). For each loop region, a total of 100 conformations were
generated, and the model exhibiting the lowest value for MODELLER's[55,56] objective was chosen as the final protein
structure.

Graph network analysis of protein trajectories

The n PDB files output by Climber were compared recursively with each other to calculate [n x n] pairwise rmsd
values. For plotting, the desired amino acids (i.e. the 250' loop – tower helix – 280s loop) were extracted beforehand
and the calculation was run only on those and the values normalised to the observed maximum. Plotting of the
rmsd trajectory was done in Matplotlib. A graph network was constructed by selecting ten pdb files equidistant on
the trajectory (with respect to number of steps taken to complete interpolation in Climber) and a Spring_layout
model was used to plot the 2-d projection with the network Python library.

DATA AVAILABILITY STATEMENT

446 The authors declare that the data supporting the findings of this study are available in this paper and its
447 supplementary information files. All additional data is available upon reasonable request. Source data are provided
448 with this paper. All mass spectrometry .raw files will be available from the PRIDE repository [accession pending].

REFERENCES

1   Fernandes, R. A. *et al.* Immune receptor inhibition through enforced phosphatase recruitment. *Nature* **586**, 779-784,
doi:10.1038/s41586-020-2851-2 (2020).
2   Gaud, G., Lesourne, R. & Love, P. E. Regulatory mechanisms in T cell receptor signalling. *Nat Rev Immunol* **18**, 485-
497, doi:10.1038/s41577-018-0020-8 (2018).
3   Mulukutla, B. C., Yongky, A., Le, T., Mashek, D. G. & Hu, W. S. Regulation of Glucose Metabolism - A Perspective
From Cell Bioprocessing. *Trends Biotechnol* **34**, 638-651, doi:10.1016/j.tibtech.2016.04.012 (2016).
4   Kruse, A. C. *et al.* Activation and allosteric modulation of a muscarinic acetylcholine receptor. *Nature* **504**, 101-106,
doi:10.1038/nature12735 (2013).
5   Oesch-Bartlomowicz, B. & Oesch, F. Cytochrome-P450 phosphorylation as a functional switch. *Archives of
Biochemistry and Biophysics* **409**, 228-234, doi:https://doi.org/10.1016/S0003-9861(02)00558-1 (2003).
6   Rojas, B. E., Hartman, M. D., Figueroa, C. M. & Iglesias, A. A. Proteolytic cleavage of Arabidopsis thaliana
phosphoenolpyruvate carboxykinase-1 modifies its allosteric regulation. *J Exp Bot* **72**, 2514-2524,
doi:10.1093/jxb/eraa583 (2021).
7   Guo, J. & Zhou, H. X. Protein Allostery and Conformational Dynamics. *Chem Rev* **116**, 6503-6515,
doi:10.1021/acs.chemrev.5b00590 (2016).
8   Wodak, S. J. *et al.* Allostery in Its Many Disguises: From Theory to Applications. *Structure* **27**, 566-578,
doi:10.1016/j.str.2019.01.003 (2019).
9   Motlagh, H. N., Wrabl, J. O., Li, J. & Hilser, V. J. The ensemble nature of allostery. *Nature* **508**, 331-339,
doi:10.1038/nature13001 (2014).
10   Willkomm, S. *et al.* Single-molecule FRET uncovers hidden conformations and dynamics of human Argonaute 2. *Nat
Commun* **13**, 3825, doi:10.1038/s41467-022-31480-4 (2022).
11   Asher, W. B. *et al.* GPCR-mediated β-arrestin activation deconvoluted with single-molecule precision. *Cell* **185**, 1661-
1675.e1616, doi:10.1016/j.cell.2022.03.042 (2022).
12   Kaplan, M. *et al.* EGFR Dynamics Change during Activation in Native Membranes as Revealed by NMR. *Cell* **167**,
1241-1251 e1211, doi:10.1016/j.cell.2016.10.038 (2016).
13   Zhang, S. *et al.* USP14-regulated allostery of the human proteasome by time-resolved cryo-EM. *Nature* **605**, 567-574,
doi:10.1038/s41586-022-04671-8 (2022).
14   Gruhl, T. *et al.* Ultrafast structural changes direct the first molecular events of vision. *Nature* **615**, 939-944,
doi:10.1038/s41586-023-05863-6 (2023).
15   Silva, D. A. *et al.* Millisecond dynamics of RNA polymerase II translocation at atomic resolution. *Proc Natl Acad Sci U
S A* **111**, 7665-7670, doi:10.1073/pnas.1315751111 (2014).

16 Barford, D. & Johnson, L. N. The allosteric transition of glycogen phosphorylase. *Nature* **340**, 609-616,
doi:10.1038/340609a0 (1989).
17 Hellenkamp, B. *et al.* Precision and accuracy of single-molecule FRET measurements—a multi-laboratory benchmark
study. *Nature Methods* **15**, 669-676, doi:10.1038/s41592-018-0085-0 (2018).
18 Ha, T. *et al.* Probing the interaction between two single molecules: fluorescence resonance energy transfer between a
single donor and a single acceptor. *Proc Natl Acad Sci U S A* **93**, 6264-6268, doi:10.1073/pnas.93.13.6264 (1996).
19 Elgeti, M. & Hubbell, W. L. DEER Analysis of GPCR Conformational Heterogeneity. *Biomolecules* **11**,
doi:10.3390/biom11060778 (2021).
20 Seal, M. *et al.* Gd(III)-19F Distance Measurements of Proteins in Cells by Electron-Nuclear Double Resonance. *Angew
Chem Int Ed Engl*, e202218780, doi:10.1002/anie.202218780 (2023).
21 Longo, J., Watson, M. J., Vos, M. J., Williams, K. S. & Jones, R. G. PYGBacking on glycogen metabolism to fuel early
memory T cell recall responses. *Mol Cell* **82**, 2918-2921, doi:10.1016/j.molcel.2022.07.016 (2022).
22 Cori, G. T., Colowick, S. P. & Cori, C. F. The Activity of the Phosphorylating Enzyme in Muscle Extract. *Journal of
Biological Chemistry* **127**, 771-782, doi:10.1016/s0021-9258(18)73783-6 (1939).
23 Barford, D., Hu, S. H. & Johnson, L. N. Structural mechanism for glycogen phosphorylase control by phosphorylation
and AMP. *Journal of Molecular Biology* **218**, 233-260, doi:https://doi.org/10.1016/0022-2836(91)90887-C (1991).
24 Klinov, S. V. & Kurganov, B. I. Kinetic Mechanism of Activation of Muscle Glycogen Phosphorylase b by Adenosine 5′-
Monophosphate. *Archives of Biochemistry and Biophysics* **312**, 14-21, doi:https://doi.org/10.1006/abbi.1994.1274
(1994).
25 Dombrádi, V. Structural aspects of the catalytic and regulatory function of glycogen phosphorylase. *International
Journal of Biochemistry* **13**, 125-139, doi:https://doi.org/10.1016/0020-711X(81)90147-6 (1981).
26 Fletterick, R. J. & Sprang, S. R. Glycogen phosphorylase structures and function. *Accounts of Chemical Research* **15**,
361-369, doi:10.1021/ar00083a004 (1982).
27 Johnson, L. N. Glycogen phosphorylase: control by phosphorylation and allosteric effectors. *Faseb j* **6**, 2274-2282,
doi:10.1096/fasebj.6.6.1544539 (1992).
28 Leonidas, D. D. *et al.* Glycogen phosphorylase revisited: extending the resolution of the R- and T-state structures of
the free enzyme and in complex with allosteric activators. *Acta Crystallogr F Struct Biol Commun* **77**, 303-311,
doi:10.1107/S2053230X21008542 (2021).
29 Sprang, Withers, S., Goldsmith, E., Fletterick, R. & Madsen, N. Structural basis for the activation of glycogen
phosphorylase b by adenosine monophosphate. *Science* **254**, 1367-1371, doi:10.1126/science.1962195 (1991).
30 Sprang, S. R. *et al.* Structural changes in glycogen phosphorylase induced by phosphorylation. *Nature* **336**, 215-221,
doi:10.1038/336215a0 (1988).
31 Sprang, S. R., Goldsmith, E. J., Fletterick, R. J., Withers, S. G. & Madsen, N. B. Catalytic site of glycogen
phosphorylase: structure of the T state and specificity for .alpha.-D-glucose. *Biochemistry* **21**, 5364-5371,
doi:10.1021/bi00264a038 (1982).
32 Klinov, S. V. & Kurganov, B. I. Combined kinetic mechanism describing activation and inhibition of muscle glycogen
phosphorylase b by adenosine 5′-monophosphate. *Biophysical Chemistry* **92**, 89-102,
doi:https://doi.org/10.1016/S0301-4622(01)00189-2 (2001).
33 Helmreich, E. & Cori, C. F. The effects of ph and temperature on the kinetics of the phosphorylase reaction. *Proc Natl
Acad Sci U S A* **52**, 647-654, doi:10.1073/pnas.52.3.647 (1964).
34 Monod, J., Wyman, J. & Changeux, J.-P. On the nature of allosteric transitions: A plausible model. *Journal of Molecular
Biology* **12**, 88-118, doi:https://doi.org/10.1016/S0022-2836(65)80285-6 (1965).
35 Krebs, E. G. & Fischer, E. H. The phosphorylase b to a converting enzyme of rabbit skeletal muscle. *Biochim Biophys
Acta* **20**, 150-157, doi:10.1016/0006-3002(56)90273-6 (1956).
36 Oikonomakos, N. G. *et al.* Flavopiridol inhibits glycogen phosphorylase by binding at the inhibitor site. *Journal of
Biological Chemistry* **275**, 34566-34573 (2000).
37 Oikonomakos, N. G. *et al.* Allosteric inhibition of glycogen phosphorylase a by the potential antidiabetic drug 3-isopropyl
4-(2-chlorophenyl)-1,4-dihydro-1-ethyl-2-methyl-pyridine-3,5,6-tricarbo   xylate. *Protein Sci* **8**, 1930-1945,
doi:10.1110/ps.8.10.1930 (1999).
38 Walcott, S. & Lehman, S. L. Enzyme Kinetics of Muscle Glycogen Phosphorylase b. *Biochemistry* **46**, 11957-11968,
doi:10.1021/bi7005527 (2007).
39 Knox, R., Lento, C. & Wilson, D. J. Mapping Conformational Dynamics to Individual Steps in the TEM-1 beta-Lactamase
Catalytic Mechanism. *J Mol Biol* **430**, 3311-3322, doi:10.1016/j.jmb.2018.06.045 (2018).
40 Sotiroudis, T. G., Oikonomakos, N. G. & Evangelopoulos, A. E. Effect of sulfated polysaccharides and sulfate anions
on the AMP-dependent activity of phosphorylase b. *Biochemical and Biophysical Research Communications* **90**, 234-
239, doi:https://doi.org/10.1016/0006-291X(79)91615-2 (1979).
41 Engers, H. D. & Madsen, N. B. The effect of anions on the activity of phosphorylase b. *Biochemical and Biophysical
Research Communications* **33**, 49-54, doi:https://doi.org/10.1016/0006-291X(68)90253-2 (1968).
42 Leonidas, D. D. *et al.* The ammonium sulfate activation of phosphorylase b. *FEBS Letters* **261**, 23-27,
doi:https://doi.org/10.1016/0014-5793(90)80627-U (1990).
43 Schreiber, W. E. & Bowling, S. An automated assay of glycogen phosphorylase in the direction of phosphorolysis. *Ann
Clin Biochem* **27 (Pt 2)**, 129-132, doi:10.1177/000456329002700207 (1990).
44 Seetaloo, N., Kish, M. & Phillips, J. J. HDfleX: Software for Flexible High Structural Resolution of Hydrogen/Deuterium-
Exchange Mass Spectrometry Data. *Anal Chem* **94**, 4557-4564, doi:10.1021/acs.analchem.1c05339 (2022).
45 Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825-2830 (2011).
46 Guo, Z. *et al.* Engineering and exploiting synthetic allostery of NanoLuc luciferase. *Nat Commun* **13**, 789,
doi:10.1038/s41467-022-28425-2 (2022).
47 Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods* **17**, 261-272,
doi:10.1038/s41592-019-0686-2 (2020).
48 Acharya, K. R., Stuart, D. I., Varvill, K. M. & Johnson, L. N.
49 Mitchell, E. P. *et al.* Ternary complex crystal structures of glycogen phosphorylase with the transition state analogue
nojirimycin tetrazole and phosphate in the T and R states. *Biochemistry* **35**, 7341-7355, doi:10.1021/bi960072w (1996).
50 Huang, J. *et al.* Insights into Phosphorylation-Induced Protein Allostery and Conformational Dynamics of Glycogen
Phosphorylase via Integrative Structural Mass Spectrometry and In Silico Modeling. *ACS Chem Biol* **17**, 1951-1962,
doi:10.1021/acschembio.2c00393 (2022).

51 Hageman, T. S. & Weis, D. D. Reliable Identification of Significant Differences in Differential Hydrogen Exchange-Mass
Spectrometry Measurements Using a Hybrid Significance Testing Approach. *Anal Chem* **91**, 8008-8016,
doi:10.1021/acs.analchem.9b01325 (2019).
52 Weiss, D. R. & Levitt, M. Can morphing methods predict intermediate structures? *J Mol Biol* **385**, 665-674,
doi:10.1016/j.jmb.2008.10.064 (2009).
53 Tam, B., Sinha, S. & Wang, S. M. Combining Ramachandran plot and molecular dynamics simulation for structural-
based variant classification: Using TP53 variants as model. *Comput Struct Biotechnol J* **18**, 4033-4039,
doi:10.1016/j.csbj.2020.11.041 (2020).
54 Kasvinsky, P. J., Madsen, N. B., Sygusch, J. & Fletterick, R. J. The regulation of glycogen phosphorylase alpha by
nucleotide derivatives. Kinetic and x-ray crystallographic studies. *Journal of Biological Chemistry* **253**, 3343-3351,
doi:https://doi.org/10.1016/S0021-9258(17)40842-8 (1978).
55 Fiser, A., Do, R. K. G. & Šali, A. Modeling of loops in protein structures. *Protein Science* **9**, 1753-1773,
doi:https://doi.org/10.1110/ps.9.9.1753 (2000).
56 Sali, A. & Blundell, T. L. Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol* **234**, 779-815,
doi:10.1006/jmbi.1993.1626 (1993).

ACKNOWLEDGEMENTS

J.J.P. and M.K. are supported by a UKRI Future Leaders Fellowship [Grant Number MR/T02223X/1]. DI is funded by a BBSRC SWBioDTP studentship. For the purpose of open access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

AUTHOR INFORMATION

AUTHORS AND AFFILIATIONS

Living Systems Institute, Department of Biosciences, University of Exeter, Stocker Road, Exeter, EX4 4QD, UK

Monika Kish, Dylan P. Ivory, Jonathan J. Phillips

Alan Turing Institute, British Library, London, NW1 2DB, UK

Jonathan J. Phillips

CONTRIBUTIONS

JJP conceived the study. MK and JJP designed the experiments and performed computational structural interpolations. MK, DI and JJP performed mass spectrometry experiments. MK and DI performed spectrophotometry assays, constructed atomistic models, analyzed mass spectrometry data, and performed statistical analysis. MK performed clustering analysis. All authors wrote the manuscript.

CORRESPONDING AUTHOR

Correspondence to JJP: jj.phillips@exeter.ac.uk

CODE AVAILABILITY STATEMENT

This study uses in-house developed scripts in Python. Available upon request.

ETHICS DECLARATIONS

COMPETING INTERESTS

The authors declare no competing interests.

FIGURES

597 **Figure 1**

598
599

600   **Figure 2**

601
602

Figure 3

A.

B.

606 **Figure 4**

FIGURE LEGENDS

Figure 1: Allosteric activation of glycogen phosphorylase b and inactivation of glycogen phosphorylase a. (A) Glycogen hydrolysis kinetics for GlyPa (black), GlyPb (blue) and GlyPb* (orange; GlyPb equilibrated with 25 mM AMP and 25 mM AS). Absorbance (A.U.) at 30 °C for +60 min at 450 nm. Mean and SD from n=2. Plots for all conditions in Figure S1A. **(B)** Glycogen hydrolysis kinetics for GlyPa (black), GlyPb (blue) and GlyPa* (orange; GlyPa equilibrated with 32 mM caffeine. Absorbance (A.U.) at 30 °C for +60 min at 450 nm. Mean and SD from n=3. Plots for all conditions in Figure S1B. **(C)** Heat map of measured activity for combinations of AMP/AS concentrations in range 0-100 mM. Inactive GlyPb (black); highest activity achieved (white). Activity was calculated by a standard curve for n=2 (Figure S1A). **(D)** Heat maps for structural perturbations in the presence of AMP/AS, detected by equilibrium HDX-MS for the nucleotide site **(i)**, tower helix **(ii)** and 380s loop **(iii)**. Deuterium uptake data from the HDX screen was used to produce these heat maps, n=3. The relative deuteration level for each sample is color coded shown on the scale right; note different normalization. **(E)** Catalytic activity of GlyPb supplemented with AS and AMP. Shaded region – GlyPa 95% confidence intervals. **(F)** Catalytic activity of GlyPa supplemented with caffeine. Shaded region – GlyPb 95% confidence intervals. **(G)** AMP binding saturation during the dead time of the HDX labelling (50 ms). Non-equilibrium addition of 25 mM AMP/25 mM AS is optimal to fully saturate the nucleotide site. **(H)** Caffeine binding saturation during the dead time of the HDX labelling (50 ms). Non-equilibrium addition of 32 mM caffeine is optimal to fully saturate the inhibitor site. **(I)** Ammonium sulfate influence on the tower helix and nucleotide site. HDX labelling data was fitted as previously described and plotted versus the state (**Figure S2**). States annotated include: GlyPb - apo, * - equilibrium activated with 25 mM AMP and 25 mM AS, *AS - equilibrium activated with 25 mM AS, ^ eq AS – equilibrated with 25 mM AS and then non- equilibrium activated with AMP, ^ - non-equilibrium activated with 25 mM AMP and 25 mM AS, ^AMP – non-equilibrium activated with 25 mM AMP, ^AS – non- equilibrium activated with 25 mM AS.

Figure 2: Clusters of structural kinetic behavior in GlyP. (A) The sum difference of the D uptake per peptide segment between non-equilibrium and GlyPb/active state plotted against x/y axes, respectively. Peptides were grouped into seven clusters (a-g) resulting from a k-means hierarchical analysis. The centroid of each cluster is denoted by the letter connected to each member by an edge. Raw D-labelling uptake data was centroided, corrected for maximum exchangeable amide hydrogens, normalized and the different states were pairwise subtracted. Dendrogram of clustering in A is shown on the right. Y-axis is Euclidean distance between the clusters (arbitrary units). The number of peptide members in each cluster is indicated at the bottom of each cluster (truncated for clarity; **Figure S9A**). **(B)** Clusters (excluding g and a) represented on GlyPb active state structure (PDB:3E3N). Notable features are indicated on the left subunit molecular surface; Notable secondary structural elements indicated. **(C)** The sum difference of the D uptake per peptide segment between non-equilibrium and GlyPa/inactive state plotted against x/y axes, respectively. Peptides were grouped into seven clusters (h-n) resulting from a k-means hierarchical analysis. The centroid of each cluster is denoted by the letter connected to each member by an edge. Raw D-labelling uptake data was centroided, corrected for maximum exchangeable amide hydrogens, normalized and the different states were pairwise subtracted. Dendrogram of clustering in C is shown on the right. Y-axis is Euclidean distance between the clusters (arbitrary units). The number of peptide members in each cluster is indicated at the bottom of each cluster (truncated for clarity; **Figure S9B**). **(D)** Clusters (excluding h, n and j) represented on GlyPa inactive state structure (PDB:1GFZ). Notable features are indicated on the left subunit molecular surface; Notable secondary structural elements indicated.

Figure 3: Local structural kinetics upon allosteric activation and inactivation revealed by non-equilibrium D-labeling. (A) Hydrogen exchange kinetics of representative peptide segments for each cluster. Mean of n=3; error bars mostly contained within data points. GlyPb (blue); fully equilibrated activated with 25 mM AMP and 25 mM AS (orange); non-equilibrium

activated with 25 mM AMP and 25 mM AS (green). **(B)** Clusters mapped on the crystal
structure of inactive GlyPb (3E3N.pdb). Detailed annotation is represented on each structure
per cluster. **(C)** Hydrogen exchange kinetics of representative peptide segments for each
cluster. Mean of n=3; error bars mostly contained within data points. GlyPa (blue); fully
equilibrated inactivated with 32 mM caffeine (orange); non-equilibrium inactivated with 32 mM
caffeine (green). **(D)** Clusters mapped on the crystal structure of inactive GlyPa (1GFZ.pdb).
Detailed annotation is represented on each structure per cluster.

Figure 4: Structural interpolation between (in)active states of GlyP. (A) Tower helix region
in inactive (left panel), transitional (middle) and active (right) conformations. Transitional
structures with multiple broken hydrogen-bonds in the alpha7 helix are observed upon
interpolation between apo GlyPb – GlyPb-AMP (activation) and between GlyPa-caffeine – apo
GlyPa (inhibition). Representative structure shown; full trajectories provided as supplementary
files. **(B)** Interpolation trajectory represented as rmsd between each structure for the tower
helix region highlighted in panel a (amino acids 256:288), normalized to maximum values from
the AMP-activation (lower triangle) and caffeine-inhibition (upper triangle), respectively. (c)
Interpolation trajectory rmsd values between each step visualized as a 2d projection of the
graph network. Values were calculated for the tower helix region (amino acids 256-288) as for
panel b. Proximity indicates similarity in structure between the structural interpolations for
AMP-activation (pink) and caffeine-inhibition (cyan). Active and inactive states as black circles.
Ten equidistant steps along the trajectory shown in each case.