Allosteric activation and inhibition of glycogen phosphorylase share common transient structural features

Monika Kish[1], Dylan P. Ivory[1], Jonathan J. Phillips[1,2]

[1]Living Systems Institute, Department of Biosciences, University of Exeter, Stocker Road, Exeter, EX4 4QD, UK

[2]Alan Turing Institute, British Library, London, NW1 2DB, UK

ABSTRACT

It remains a major challenge to ascertain the specific structurally dynamic changes that underpin protein functional switching. There is a growing need to complement structural models with the ability to determine the dynamic structural changes that occur as these proteins are regulated and function. The archetypal allosteric enzyme, glycogen phosphorylase is one of the most studied proteins and is a clinical target of much interest to treat type II diabetes and metastatic cancers. However, a lack of understanding of its complex regulation, mediated by dynamic structural changes, hinder its exploitation as a drug target. Here, we precisely locate dynamic structural changes upon allosteric activation and inhibition of glycogen phosphorylase, by developing a time-resolved non-equilibrium millisecond hydrogen/deuterium-exchange mass spectrometry (HDX-MS) approach. We resolved obligate transient changes in localized structure that are absent when directly comparing active/inactive states of the enzyme and show that they are common to allosteric activation by AMP and inhibition by caffeine, operating at different sites. This indicates that opposing allosteric regulation by inhibitor and activator ligands is mediated by pathways that intersect at a common structurally dynamic motif. This approach has broad application to determine the structural kinetic mechanisms by which proteins are regulated.

INTRODUCTION

Allostery refers to the transfer of a signal between two sites of a protein resulting in a change in the catalytic activity to a substrate or binding affinity to a ligand at a remote site of the protein. These sophisticated structural transitions are fundamental to receptor transduction[1], cell signalling[2], and metabolic regulation[3]. While allostery is typically associated with ligand binding to the allosteric site, connected to conformational change[4], these structural transitions can also be triggered by covalent modification (e.g., phosphorylation)[5], or proteolytic cleavage[6]. Despite extensive studies since its inception[7-9], very little is known about the core dynamic process of allosteric communication, precisely how these signals are transmitted long distances through a protein molecule. This is in large part due to the lack of biophysical approaches to measure these signals at high resolution in both time and space. Recently, important advances have been made towards this goal in single molecule FRET[10,11,12,13], NMR[14], time resolved cryo-EM[15], time resolved crystallography[16], dynamic simulations[17] and Double Electron Electron Resonance (DEER)[18,19]. Perhaps the most direct evidence for an intramolecular mechanism comes from infrared laser absorbance and emission experiments of conjugated peptides, which strongly argues that the vibrational energy transfer (VET) is mediated by hydrogen-bonds. Collectively, these studies begin to reveal non-equilibrium protein structural dynamics at high structural and temporal resolution, first described by Monod and colleagues in 1965[20]. Though impressive, they still require considerable adaptation per sample and are not yet broadly applicable to cases of ligand-induced allostery in proteins.

Glycogen phosphorylase (GlyP) is the archetypal allosteric enzyme whose regulation is tightly coupled to solid tumor metastasis[21], type II diabetes[22] and adaptive immunity (early

memory CD8+ T-cell recall response)[23]. GlyP catalyzes the first committed energetic step of
carbohydrate metabolism from glycogen stores and as such is one of the most highly
regulated enzymes known, with at least six allosteric sites that sense natural ligands and
drugs. GlyP has been intensively studied since the 1930s[24], thus there is abundance of
structural and kinetic data on this protein system[25-34], with detailed studies on the activation
mechanism[35] and stabilization of R/T-states (active/inactive-states) by allosteric ligands[36]
and phosphorylation[37]. GlyP is, therefore, an ideal system to challenge our ability to discern
dynamic structural mechanisms of allosteric regulation as it is alternately activated by AMP
at the 'nucleotide site' and inhibited by caffeine at the 'inhibitor site'[38,39], whilst also being
constitutively active by Ser 14 phosphorylation[20]. Binding of AMP at the allosteric effector
site brings similar changes to phosphorylation but believed to be through somewhat different
mechanisms[40]. How a single protein can support allosteric activation and inhibition by
multiple intersecting structural pathways has not been elucidated.

Hydrogen-deuterium exchange mass spectrometry (HDX-MS) is a sensitive measure of
solvent accessible hydrogen-bonding of polypeptide backbone amide groups. The amide H-
bonds directly relate to local structural dynamics and now can be routinely determined in
large protein systems[41], with millisecond time resolution[42] and at or near single amino acid
structural resolution[43]. Here, we used a non-equilibrium millisecond HDX-MS approach to
identify specific dynamic changes that follow ligand binding. The difference in deuterium
labeling before, during and after ligand binding provides a highly structurally resolved
measurement of dynamic reconfiguration during allosteric regulation. Equilibrium
experiments reflect local minima of R/T-states, while non-equilibrium data reveal transition
state ensembles between local minima on the energy landscape[44]. Here, we clearly identify
amino acids that participate in transient conformational dynamics and we propose a
structural pathway for those changes upon allosteric ligand binding. This approach stands to
uniquely discover local structural dynamics and could be used broadly to identify such
features in almost any ligand-induced allosteric transition of a protein.

RESULTS

Non-equilibrium allosteric activation/inhibition of GlyP

To observe the kinetically populated transition state of glycogen phosphorylase activation
and inhibition, we first established the chemical conditions that fully activate or inhibit the
enzyme. The catalytic activity for the fully active form phosphorylated at Ser14 (GlyPa) was
used as a positive control reference in a glycogen hydrolysis colorimetric assay of GlyPb
activation (**Figure 1A,E**; **Figure S2A**). Against this level, the maximal extent of GlyPb
allosteric activation by AMP was then established[26,40]. This showed that full activation is
achieved with AMP ≥25 mM, the lowest concentration of AMP tested (range 25-100 mM),
given no significant difference between the activity of GlyPb+AMP compared with GlyPa
(**Figure 1A,E**). We also sought to verify the requirement of ammonium sulfate (AS) for full
activation, known to mimic the presence of phosphate by binding GlyP at the AMP effector
site, catalytic site and Ser14 phosphorylation site[20]. There are conflicting reports on the
requirement of AS for full activation of GlyPb cooperatively with AMP: whilst GlyPb affinity for
AMP is reported to increase 50-fold and activity to increase by 50% with sulfate[25,45-47], here
we found activation by sulfate alone was not achieved (15% of maximum), nor was a
cooperative effect observed on activity (**Figure 1E,C** and **Figure S2A**). However, there was
structural cooperativity between AMP and AS, as evidenced by concentration-dependent
changes in hydrogen-exchange kinetics for amino acids site-localized to the nucleotide site
and tower helix (**Figure 1D**). Local structural perturbations were observed for GlyPb samples
supplemented with 0-100 mM AMP and/or 0-100 mM AS by measuring HDX-MS at
equilibrium. Binding of AMP at the nucleotide site results in local protection against D-
labeling that is positively cooperative with AS (**Figure 1D**). Corresponding changes were
observed in the tower helix, which shows only weak protection with AS alone, but stronger
protection with AMP (similar at all AMP concentrations tested) that increases with [AS]

(**Figure 1D**). The 380 loop shows cooperativity between AS and AMP, resulting in
deprotection against D-labeling as the loop becomes disordered in the active state[20] (**Figure**
1D). This identified cooperativity in GlyPb structural activation by AMP/AS that is not readily
observed in the activity assay (**Figure 1D,E**)[25,40,48]. One-way ANOVA tests proved that
higher concentrations of AMP or AS were not significant for either catalytic activity or
structural perturbation. Therefore, the lowest concentrations tested of 25 mM AMP and 25
mM AS were considered to fully activate GlyPb catalytically and structurally and these
conditions were used for the experiments to study non-equilibrium activation of GlyPb.
Caffeine inhibits the active GlyPa R-state, stabilising the T-state, therefore we also sought to
determine the minimal caffeine concentration required to fully inhibit 12 μM GlyPa – the
concentration for subsequent non-equilibrium HDX-MS experiments. This was achieved with
32 mM caffeine, given by equivalence in specific activity to inactive GlyPb by ANOVA these
conditions were used for the experiments to study non-equilibrium inhibition of GlyPa
(**Figure 1F** and **Figure S2B**).

In order to observe any kinetically populated transient structural ensembles, it is critical for
the non-equilibrium experiments that the protein:ligand complex is fully saturated within 50
ms (dead time of the non-equilibrium experiment), governed by the second-order rate
constant of association with units $M^{-1}s^{-1}$. While T-state GlyPb is allosterically activated by the
binding of AMP at the nucleotide site, active phosphorylated R-state GlyPa is allosterically
inhibited by the binding of caffeine between the 280s active site gate loop and the L40 (Tyr
613) loop. Therefore, the degree of AMP saturation at the nucleotide site was determined by
protection against D-labeling at amino acids T38-T47, whereas the degree of caffeine
saturation was determined by protection against D-labelling at amino acids N274-F285. In
separate experiments, a range of AMP concentrations (2.5, 25 and 100 mM) and a range of
caffeine concentrations (3, 10 and 32 mM) were assessed by rapid mixing with GlyPb or
GlyPa, respectively, under non-equilibrium conditions (**Figure 1G-H**). To confirm that a 1:1
complex with ligand had been reached, these were compared with the end-state of the
respective process: activated GlyPb fully-equilibrated with 25 mM AMP, or inhibited GlyPa
fully-equilibrated with caffeine. After 50 ms of incubation with ligand in deuterated buffer, 32
mM of caffeine achieved equivalent protection against D-labelling compared with the fully-
equilibrated inhibited state (GlyPa + 32 mM caffeine) within the 50 ms dead-time of the HDX-
MS experiment. The case of GlyPb activation was more complicated, as it was important to
consider the contribution of both AMP and AS to ternary complex formation under these
rapid-mixing conditions (**Figure S3**Error! Reference source not found.). Incubation of GlyPb
with AS fully equilibrated (*AS) or at non-equilibrium (^AS) showed a lack of protection
against D-labeling relative to GlyPb alone, signifying absence of binding to the putative sites
for AS interaction. 25 mM AMP alone did not fully bind in 50 ms, whereas full protection is
achieved within error for the non-eq addition of 25 mM AMP and AS. This confirms that the
presence of AS enhances the k_a for GlyP:AMP association and these conditions yield full
complex formation inside the dead time of the non-eq HDX-MS experiment. As we and
others have recently reported, the presence of salts can significantly alter the intrinsic rates
of HDX, thus masking – or artefactually creating – protein structural dynamics that are
inferred by the HDX data[49]. This is not a systemic alteration of HDX kinetics by the presence
of AS, but rather it is confirmed to be a local effect at the nucleotide site, as peptides
covering other regions of the protein in the same HDX-MS data set do not show the same
behavior: for example the tower helix (reported by peptide segment V266-I275) shows the
opposite relationship with the k_{obs} for D-labeling being significantly faster in the presence of
25 mM AS/AMP under equilibrium and non-eq conditions (**Figure 1I, Figure S3**Error!
Reference source not found.). We concluded that the ligand binding sites for inhibition and
activation of GlyP by caffeine and AMP, respectively, are achieved within the dead-time of
the HDX-MS experiment at times post-mixing >= 50 ms. Therefore, we proceeded to
evaluate the transient structural kinetics that occur upon allosteric regulation across the
whole enzyme.

Transient structural dynamics during GlyPb allosteric activation by AMP

To identify specific dynamic changes associated with long-range allosteric communication
induced by ligand binding, we developed a non-equilibrium approach to millisecond
hydrogen/deuterium-exchange that reveals the local structural perturbations that result from
the conditions established above (**Figure S1**). As the D-labeling of polypeptide backbone
amide groups is a sensitive, but convoluted measure of H-bonding and solvent accessibility,
first we sought to categorize the local differences in the protein ensemble during the non-
equilibrium phase following allosteric activation by AMP/AS. Subtle changes in dynamics
and structural changes can be derived from HDX-MS data, reflected in the observed kinetics
of deuterium uptake plots at specific positions within the protein, given by short peptide
segments in the 'bottom-up' experiments used here[44]. The data comprised measurements at
31 D-labeling time points over four orders of magnitude from 50 ms – 300 s for 219 peptide
segments that cover 90.5% of the GlyP amino acid sequence (**Figure S4A**). This was done
for three conditions: (i) inactive GlyPb (apo), (ii) fully activated GlyPb (termed GlyPb*)
equilibrated for one hour with 25 mM AMP, 25 mM AS and (iii) GlyPb activated at non-
equilibrium by rapid mixing with 25 mM AMP, 25 mM AS (termed GlyPb^). It is clear that for
many peptides in a variety of regions of the protein the D-labeling kinetics under non-eq
conditions are qualitatively different from both the apo and the activated states (**Figure S5-6**
and **Figure S16**). There are peptides for which the D-labeling values fall outside of the range
measured for either the initial (apo) or equilibrated active (GlyPb*) forms, which indicates a
degree of non-linearity in the structural interpolation (**Figure 3, Figure S16;17**). However,
the sub-molecular location, magnitude and nature of relative (de)protection of these show a
variety of differences. Therefore, we next sought to classify these structural dynamic
changes occurring during allosteric activation to better identify clear behaviors in each part of
the enzyme.

To categorize the local structural changes that occur during non-equilibrium
activation/inhibition into correlated transient dynamics, we developed a simple, but robust
quantitative analysis of the HDX-MS data that served to represent the non-equilibrium HDX
kinetics for the pool of peptide segments and to act as input to a clustering algorithm[50,51].
Each amino acid of the protein was represented spatially in a manner that reflects the
relationship of its non-equilibrium state to the starting and ending states of the protein
(**Figure 2**). We identified the archetypes of non-equilibrium structural kinetics that might be
possible to observe in this plot (**Figure S9)**. In order to categorize the continuum of diverse
D-labeling kinetic differences that we observed, we performed a k-means clustering analysis
to identify amino acids with correlated HDX kinetics. Seven clusters of amino acids within
GlyP comprises a common structural response to activation by AMP/AS (**Figure 2, Figure
S10**). **Figure 3A** shows the D-labeling uptake plot for representative amino acids from each
cluster, alongside the structural location of the cluster.

The hierarchical relationship between the seven clusters was determined based on the
observed transient structural changes (**Figure 2A**)[52]. This reveals that there are two closely
related clusters (**a** and **g)** that show little or no observable transient structural change during
allosteric activation (i.e. close to the origin on **Figure 2A** and **Figure SI15**). We calculated
the local hydrogen exchange rate (k_{obs}) by fitting HDX-MS data to a stretched exponential
model[49]. For most peptides, k_{obs} was low (k_{obs} < 1 x 10^{-5} s^{-1} for 130 of 219 peptides
measured) in all three conditions (Error! Reference source not found.**Table S2** and **Table
S3**), as GlyP is highly structured with many amino acids buried in the core of the enzyme
with low solvent accessible surface area (SASA) and stable H-bond networks, consistent
with previous studies[20,30,49,53-56]. This validates that much of the enzyme is structurally
identical before, during and after allosteric activation, **Figure S5-6** and **Figure S16**. A third
related cluster (**e**), representing the nucleotide binding site for AMP, shows no deviation from
the active state and only transient relative protection compared to the inactive state that is
on-pathway from the apo- to AMP-bound state (i.e. on y=0 in **Figure 2A**). This group
includes residues from α2 (residues 47-78), the cap' region (residues 38-47) and residues
from the nucleotide binding domain (**Figure 3A** and **Figure S16**). The α8 helix is also
essential for AMP binding, but unfortunately our coverage was lacking peptides in that

region. There are two closely related clusters which represent the 250' loop (**f**) and the tower
helix (**c**) that show transiently increased D-labeling compared to one or both equilibrium
states. There are also two related clusters, representing the pyridoxal phosphate cofactor
binding residues (**b**) and the 606 loop that are distinct from all others which show protection
against D-labeling compared to both the inactive and the active state. The amino acids
surrounding the PLP cofactor retain apo GlyPb HDX kinetics, indicating either that this
region has a structural transition with a low probability or that the induced protection against
HDX occurs too slowly to prevent the observed labeling.

A feature of the HDX-MS data during non-equilibrium AMP activation is that there are
transient structural changes that are distinct from both the inactive/active states (i.e. amino
acids that do not lie on either x=0 or y=0 in **Figure 2A**) and therefore are not represented by
high-resolution structural models derived from the equilibrium states. Activation of GlyPb by
AMP binding to the nucleotide site induces the well-established structural changes in the
250' and 280s loops, together with rotation of the tower helix (α7) (**Figure S12**). The non-
equilibrium HDX-MS reveals previously unobserved transient changes that facilitate this. The
250' loop (A248-G260) exhibits unique behavior during allosteric activation (cluster **f** in
Figure 3A). The D-labeling kinetics interpolate smoothly between the GlyPb state (at t=50
ms) and the active GlyPb* state (at t=10 s). This indicates a likely two-state process for
helical lengthening and that a gain of structure has fully equilibrated in the molecular
population by 10 s of D-labeling, consistent with the tower helix in the active state
lengthening at the N-terminal end by one turn in the GlyPb* state (3E3N.pdb[30]). Clusters **c**
(the tower helix) and **d** (the 606 loop) show opposite structural behaviors: the 606 loop
shows transient protection (increased H-bonding; reduced solvent accessibility), whereas the
tower helix shows transient deprotection (i.e. local unfolding). The tower helix (G261-S276)
is a contiguous alpha helical structure in all known experimental models of apo, active and
inactive forms of GlyP. However, unexpectedly here we observe that it transiently loses
protection against hydrogen/exchange, with increased observed D-labeling during allosteric
activation following AMP binding at the nucleotide site (Error! Reference source not found.**A**
and **Figure S12**).

We sought an atomistic rationale for this transient unfolding, so an all-atom interpolation
method was used to generate structures on a non-linear, energy minimized trajectory
between apo GlyPb and GlyPb*. The Climber method has been shown to sample
intermediate structures effectively, though has been difficult to validate until now, owing to
the experimental challenge of determining transient local structural changes[57]. Here a
trajectory was created which interpolates between two crystal models that have intact tower
helices with predicted hydrogen-bonds in conventional geometries, yet several of these
hydrogen-bonds are shown to undergo dynamic remodeling as the helices refold and rotate
by ~40° relative to each other to adopt the established active R-state, GlyPb* (**Figure 4A**
and **Figure S12**; **Supplementary movies**)[58]. The interpolation is consistent with the non-
equilibrium HDX-MS data in which there is relative deprotection compared to both GlyPb and
GlyPb* precisely at I263-D268, resolved by multiple overlapping peptide segments in the
data, supporting an interpretation that there is a transient unfolding in the tower helix during
allosteric activation.

Transient structural dynamics during GlyPa allosteric inhibition by caffeine

We next considered whether this transient unfolding of the tower helix seen during activation
is a feature common to the GlyP inhibition pathway. GlyPa phosphorylated at serine14 is
basally active but is itself allosterically inhibited by binding of caffeine, which stabilizes a T-
state that is similar to the GlyPb apo state. The caffeine binding site is formed between the
280s and 606 loops, 3.5 nm from the nucleotide site where AMP binding stabilizes the
reverse transition to the R-state. Therefore, we next sought to identify the transient structural
features on the inhibition pathway by performing non-equilibrium HDX-MS of GlyPa in the
presence of caffeine. As for the AMP-activation experiments, this comprised measurements
at 20 D-labeling time points 50 ms – 300 s resulting in 171 peptide segments that cover

91.4% of the GlyP amino acid sequence (Error! Reference source not found.**B**). This was
done for three conditions: (i) active GlyPa (apo), (ii) fully inhibited GlyPa (termed GlyPa')
equilibrated for one hour with 32 mM caffeine and (iii) GlyPa inhibited at non-equilibrium by
rapid mixing with 32 mM caffeine (termed GlyPa^).

From a comparison of the observed HDX kinetics between GlyPa^ and each of the two
equilibrated states, it is clear that the inhibition pathway involves a number of amino acids
that show a transient structure distinct from both the start/end states (**Figure 2C-D; Figure
S7-8**). Based on this, a clustering analysis identified amino acids with correlated HDX
kinetics. As for AMP-activation, caffeine-inhibition resulted in seven clusters (**Figure S11;15**)
of amino acids within GlyP with a common structural response to ligand (**Figure 2C; Figure
3B**). Again, much of the protein is relatively unchanged before, during and after allosteric
inhibition (clusters **h** and **j**, **Figure 2C**), which is consistent with the available crystal
structures. The 280s and 606 loops form the binding site for caffeine which intercalates
between them in the GlyPa T-state crystal model. These amino acids are co-located in
cluster **l** which shows HDX equivalent to the GlyPa' and protected compared to the GlyPa
states, consistent with direct binding at this site without any observed transitional behavior
(cluster **l, Figure 2C-D;Figure 3B**). During caffeine-inhibition these three clusters (**h/j/l**) have
minimal transient structural features and are found to be hierarchically unrelated to all other
amino acids in the other four clusters (**Figure 2C**). Furthermore, the 250' loop is highly
deprotected against hydrogen-exchange and labels more than the initial GlyPa state, but
similarly to the inhibited GlyPa' state (cluster **k**). This is consistent with an interpretation that
it is a two-state process, where the caffeine-inhibited T-state is observed immediately upon
binding, in which one turn of the tower helix unwinds to extend the 250' loop, as seen in the
crystal models. The tower helix itself (cluster **m**) is labeled more with deuterium under non-
equilibrium conditions (GlyPa^) than in either GlyPa or GlyPa' states. As this is a highly
solvent-exposed alpha helix in the initial R-state, it is most consistent with an interpretation
of transient unfolding (**Figure S13**), again resolved by overlapping peptide segments to be
most pronounced at amino acids L254-L271. The 280s loop shows a diverged behavior, with
the more C-terminal part forming part of the direct ligand binding site (cluster **l**) and with the
N-terminal stretch showing transiently reduced hydrogen-exchange compared to the initial
apo GlyPa state but increased when compared to GlyPa'. This is consistent with the initial
binding of caffeine to F285 and a subsequent propagation of induced helical structure further
upstream in the 280s loop as the tower helix lengthens by one turn at the C-terminal end.

It is apparent that a number of highly resolved transient structurally dynamic features are
observed by non-equilibrium HDX-MS upon allosteric inhibition of GlyP by caffeine.

**Transient unfolding of the tower helix is common to allosteric activation and
inhibition pathways**

We next sought to compare the transient structural features observed upon AMP-activation
and caffeine-inhibition to determine whether these different ligands (and different allosteric
sites) appear to act via different pathways, or the same one.

The 250' and 280s loops show opposite neHDX-MS behaviour in response to AMP and
caffeine, which is logical given that these motifs are structurally different in the R/T-states
and necessarily interpolate between them. However, there is an unexpected common
feature during both allosteric activation and inhibition with transiently increased HDX
observed in the tower helix that sits in between these two loops. Non-linear interpolation
between the atomistic models for GlyPa R-state (apo) and caffeine-bound (T-state) also
predicts breakage, then reforming, of hydrogen-bonds in the tower helix, as it does for AMP-
activation (**Figure 4A**). It also predicts that the two structural pathways are closely related
with low rmsd in the 250' loop, tower helix and 280s loop between structures close to the T-
state and much larger differences in rmsd close to the R-state (**Figure 4B**). As the tower
helix is highly solvent-exposed, the transiently lower protection factor would likely be the
result of H-bond breakage which is known to be a major determinant of hydrogen-exchange

rate. This is suggested by the structural interpolation, which predicts these hydrogen-bonds
in the tower helix to be volatile in both pathways (**Figure 4A**).

These are the major transient structural features observed by non-equilibrium HDX-MS and
appear common to both pathways, with a consistent atomistic rationalization given by the
Climber structural interpolations.

DISCUSSION

Transient features of allosteric pathways revealed

Here we show that allosteric regulation of the central metabolic enzyme, glycogen
phosphorylase, involves a number of dynamic changes in local structure that are not
observed in static structural models of the R- and T-states. We established a new protocol
exploiting millisecond hydrogen/deuterium-exchange at non-equilibrium to characterize
transient changes in protection in response to ligand, resolved at near amino acid level
throughout a 194 kDa enzyme. This revealed that AMP-activation induces a transient
increase in HDX in the tower helix. We rationalized this by interpolating the known structures
for this allosteric transition using an atomistic method based on the Energy Calculation and
Dynamics (ENCAD) molecular-mechanics force-field, Climber. This predicts that the well-
established lengthening and 40° relative rotation of the tower helices by first shortening, then
bending it, wholly consistent with the increased hydrogen-exchange observed during the
process. Unexpectedly, caffeine-inhibition – binding to a different allosteric site 3.5 nm away
from AMP - also resulted in a transient increase in HDX in the tower helix. The structural
interpolation predicts that this is due to a similar extent of transient breaking of hydrogen-
bonds in the tower helix (**Figure 4A**). So, both experiments are consistent with the
interpretation suggested by the simulation that breaking of the tower helix is a common
requirement of both AMP-activation and caffeine-inhibition pathways (**Figure 4B-C**).

Slow (millisecond) labeling can observe transient structural dynamics of the molecular population

It is perhaps unexpected that differences in non-equilibrium HDX-MS kinetics were observed
across a range of time domains from milliseconds to seconds. Structural kinetics were
observed even at the fastest labeling time of 50 ms (**Figure S5;7**), in several significant
regions, including: 250 □ turn loop, β3 and α6, 280s loop and another □ turn with the central
residue Tyr 613 said to be forming a hydrophobic sandwich of the nucleotide inhibitor site[59].
At slower deuterium labeling times, significant changes in protection between the three
states was observed throughout the α7 tower helix (residues 261-274), the β11b (residues
276-279), α23 (residues 714-725) and α24 (residues 728-735) (**Figure S6;8**). We interpret
this range of observed non-equilibrium structural kinetics to indicate that the probability of
the local structural changes is very low (i.e. bounded by a high energy transition state),
resulting in observation of the populated transient species (presumably transition states),
even milliseconds after ligand binding. The wide range of local protection factors against
hydrogen-exchange (due to existence of stable – or transient - local structure) would then
effectively stretch the observation of structural changes across multiple orders of magnitude,
as seen here.

Hydrogen/deuterium-exchange may directly observe vibrational energy transfer upon ligand binding

The HDX-MS intrinsic rate constants indicate a half time of 100's of milliseconds for which a
protein structural state would need to exist in order for it to be appreciably detectable by this
technique[60]. However, our non-equilibrium HDX-MS data appear to observe transient
species whose lifetime would normally be considerably shorter than this. One explanation for
this is that the vibrational energy transfer (VET), derived from the enthalpy of ligand binding,
is anisotropically focused into a 'hotspot' which can undergo the HDX reaction faster than
would be predicted because of an elevated local temperature. This would agree with our

simulated trajectories where hydrogen bond stability is locally reduced in the 'hotspot' motif
and with recent experimental evidence that VET is mediated by hydrogen-bond networks[61]
and with coarse-grained models of specific energy dissipation pathways in proteins[62].

We believe that this approach, while critically important to establish rapid saturation of the
ligand binding site, provides a straightforward way to directly observe allosteric structural
dynamics at non-equilibrium which can be applied broadly to proteins irrespective of size or
stability and with, in principle, any chemical or protein ligand. In this manner, we expect it to
be important in future to understand not only what has changed in protein structure, but how
the changes come about and propagate through the molecule.

METHODS

Sample preparation

Glycogen phosphorylase b and glycogen phosphorylase a were dissolved in phosphate buffered saline (PBS),
pH 7.4 to 10.3 pmol/µL. For the equilibrium and non-equilibrium activated samples, the required amount of
AMP/AS and caffeine were added, in the stock solution or the labelling buffer, respectively. During the screening
experiments (HDX screen and enzyme activity assay) each sample was prepared simultaneously, with 10.3
pmol/µL GlyPb/GlyPa and with varying concentrations of AMP, AS or caffeine, **Error! Reference source not**
found..

Screening of conditions to activate GlyP by enzyme activity assay

The enzyme activity of GlyPb and GlyPa was screened for AMP/AS and caffeine dependence**Error! Reference**
source not found.. A colorimetric assay kit was used (ab273271), with only one adjustment of the protocol for
GlyP sample preparation. A suitable protein concentration was determined by pre-testing a range of
concentrations and plate reader settings, then all 25 samples were prepared in parallel at 1 mg/mL GlyP (**Error!**
Reference source not found.). Three relevant concentrations of caffeine were screened (3, 10 and 32 mM) to
determine the extent of inhibition. Reaction kinetic curves and specific activity (mU/mG) were plotted and
calculated according to the suggested protocol.

Screening of conditions to activate GlyP by HDX-MS

The observable perturbations in HDX were screened for AMP/AS dependence. A CTC PAL sample handling
robot (LEAP Technologies, USA) was used to mix samples (**Error! Reference source not found.**) with
deuterated PBS buffer in 1:20 ratio, then quench the labeling reaction by 1:1 mixing with 100 mM potassium
phosphate to a final pH of 2.55 at 0 °C. Samples were immediately digested online with an Enzymate
immobilized pepsin column (Waters) at 12 °C, the derived peptides trapped on a VanGuard 2.1 x 5 mm
ACQUITY BEH C18 column (Waters) for 3 minutes at 125 µL/min at 0.5 °C and separated on a 1 × 100 mm
ACQUITY BEH 1.7 µm C18 column (Waters) with a 7 min linear gradient of acetonitrile (5-40%) supplemented
with 0.1% formic acid. Peptides were eluted into a Synapt G2-Si mass spectrometer (Waters).

Ammonium sulfate influence and maximum velocity kinetics

The maximum velocity kinetics and the AS influence on the HDX were determined prior continuing with the non-
equilibrium experiments. HDX labelling experiments were performed at 11 time points, ranging from 50 ms to 300
s. AMP activation was tested at 3 different concentrations, including 2.5, 25 and 100 mM under non-equilibrium
conditions with 25 mM AS. The AS influence on the equilibrated sample and the pre-equilibrated AS sample, was
also determined. Analyzed protein states include: GlyPb - apo, * - equilibrium activated with 25 mM AMP and 25
mM AS, *AS - equilibrium activated with 25 mM AS, ^ eq AS – equilibrated with 25 mM AS and then non-
equilibrium activated with AMP, ^ - non-equilibrium activated with 25 mM AMP and 25 mM AS, ^AMP – non-
equilibrium activated with 25 mM AMP, ^AS – non- equilibrium activated with 25 mM AS.

Hydrogen-deuterium exchange mass spectrometry

HDX was performed using a fully-automated, millisecond HDX labelling and online quench-flow instrument,
ms2min (Applied Photophysics Ltd), connected to an HDX manager (Waters). During the labelling experiments,
14 µL of either apo or equilibrated GlyPb/GlyPa were delivered to the labelling mixer. A 20-fold dilution with
labelling buffer at 20˚C commenced the HDX. Depending on the desired experiment, the labelling buffer differed
significantly. For apo and equilibrium experiments, the labelling buffer was 1 x PBS, pH$_{read}$ =7.00 at 20˚C. During
non-equilibrium experiments and other assessments (see Error! Reference source not found.), the labelling buffer
consisted of 1 x PBS, pH =7.00 at 20˚C combined with 25 mM AMP/25 mM AS or 32 mM caffeine. All GlyPb
samples were labelled at 30 time points, ranging from 50 ms to 300 s. All GlyPa samples were labelled at 20 time
points, ranging from 50 ms to 300 s. Immediately after, the HDX reaction was quenched by mixing with quench
buffer (100 mM Potassium phosphate, pH =2.5 at 0˚C) at 1:1 ratio, and sample digested online with an Enzymate
immobilized pepsin column (Waters), the derived peptides trapped on a VanGuard 2.1 x 5 mm ACQUITY BEH
C18 column (Waters) for 3 minutes at 125 µL/min and separated on a 1 × 100mm ACQUITY BEH 1.7 µm C18

column (Waters) with a 7-minute linear gradient of acetonitrile (5-40%) supplemented with 0.1% formic acid.
Peptides were eluted into a Synapt G2-Si mass spectrometer (Waters). Mass spectra were obtained using the
Waters HDMS^E mode within a mass range of 50 to 2000 m/z. Instrument parameters were configured as follows:
a capillary voltage of 3.0 kV, a cone voltage of 50 V, a trap collision energy of 4 V, a traveling wave ion mobility
separation at a velocity of 475 m/s, a wave amplitude of 36.5 V, and a nitrogen pressure of 2.75 mbar. For low-
energy scans, a transfer collision energy of 4 V was applied, while high-energy scans utilized four separate
collision energy ramps ranging from 15 to 55 V.

Data and statistical analysis

PLGS (ProteinLynx Global Server 2.5.1, Waters) was used for the analysis of MS^E reference data and
identification of all discoverable peptic peptides. DynamX 3.0 (Waters, USA) was used for processing and
assignments of isotopic distributions of all raw data files. All HDX-MS experiments were performed in triplicate
(technical replicates). The means and standard deviations (SD) of these replicates were used to determine the
significance of changes, global significance threshold, as previously described[56]. In-house developed scripts in
Python 3.4.0 (Python Software Foundation) were used for all post-processing analyses, clustering analyses, and
plotting in PyMOL (Schrödinger, Inc.).

A clustering analysis was employed to investigate the behavior of the peptide during non-equilibrium
activation/inhibition. The D-labeling data for each peptide segment was normalized to the theoretical maximum,
to account for varying peptide length and presence of proline for which there is no signal. The sum of deuterium
uptake over all time points for each peptide was calculated for all three protein conditions (states) of GlyPa or
GlyPb, in each case. The values for the non-eq data were then subtracted from both other states for every
peptide segment in the data set, resulting in an [n x 2] vector of 2-d coordinates. With the non-equilibrium
measurements defined as 2-d coordinates relative to a start/end state, this constitutes a well-formed input for a k-
means clustering analysis. The 2-d coordinates for HDX-MS differences were then *k-means* clustered with *k*
clusters indicated to be optimal by both an elbow method and a Silhouette analysis.

Non-linear atomistic structural interpolation

The Climber program was utilized for stepwise and non-linear structural interpolation between the active and
inactive conformations using default paramters. This involved dynamic evaluation of RMSD between backbone
and side chain atom pairs, with continual updates. Atom pair sets were created for each generated conformation
between the initial (A) and final (B) states, based on criteria of distance between alpha carbons (>10 Å) and side
chain atom pairs (<10 Å). Harmonic restraints were applied to steer inter-residue distances towards the target
conformation. The force constant for conformational transformation was adjusted based on RMSD progress.
Morphed conformations underwent iterative minimization using the ENCAD potential energy function. Initial
coordinates for the allosteric activation and inhibition were taken from crystallographic structures of glycogen
phosphorylase b and glycogen phosphorylase a. PDB ID 1gpb and 3e3n were used for GlyPb transition, and
PDB ID 1gpa and 1gfz for GlyP inhibition. Missing residues (251-259, 315-324 in 1gfz; 251-260, 281-287 in 3e3n)
were built and optimized using MODELLER (version 10.4). For each loop region, a total of 100 conformations
were generated, and the model exhibiting the lowest value for MODELLER's[63,64] objective was chosen as the
final protein structure.

Graph network analysis of protein trajectories

The n PDB files output by Climber were compared recursively with each other to calculate [n x n] pairwise rmsd
values. For plotting, the desired amino acids (i.e. the 250' loop – tower helix – 280s loop) were extracted
beforehand and the calculation was run only on those and the values normalised to the observed maximum.
Plotting of the rmsd trajectory was done in Matplotlib. A graph network was constructed by selecting ten pdb files
equidistant on the trajectory (with respect to number of steps taken to complete interpolation in Climber) and a
Spring_layout model was used to plot the 2-d projection with the network Python library.

DATA AVAILABILITY STATEMENT

466 The authors declare that the data supporting the findings of this study are available in this paper and its
467 supplementary information files. All additional data is available upon reasonable request. Source data are
468 provided with this paper. All mass spectrometry .raw files will be available from the PRIDE repository [accession
469 pending].

REFERENCES

1   Fernandes, R. A. *et al.* Immune receptor inhibition through enforced phosphatase recruitment. *Nature*
586, 779-784, doi:10.1038/s41586-020-2851-2 (2020).
2   Gaud, G., Lesourne, R. & Love, P. E. Regulatory mechanisms in T cell receptor signalling. *Nat Rev*
Immunol **18**, 485-497, doi:10.1038/s41577-018-0020-8 (2018).
3   Mulukutla, B. C., Yongky, A., Le, T., Mashek, D. G. & Hu, W. S. Regulation of Glucose Metabolism - A
Perspective From Cell Bioprocessing. *Trends Biotechnol* **34**, 638-651, doi:10.1016/j.tibtech.2016.04.012
(2016).

4   Kruse, A. C. *et al.* Activation and allosteric modulation of a muscarinic acetylcholine receptor. *Nature*
504, 101-106, doi:10.1038/nature12735 (2013).
5   Oesch-Bartlomowicz, B. & Oesch, F. Cytochrome-P450 phosphorylation as a functional switch. *Archives*
of Biochemistry and Biophysics **409**, 228-234, doi:https://doi.org/10.1016/S0003-9861(02)00558-1
(2003).
6   Rojas, B. E., Hartman, M. D., Figueroa, C. M. & Iglesias, A. A. Proteolytic cleavage of Arabidopsis
thaliana phosphoenolpyruvate carboxykinase-1 modifies its allosteric regulation. *J Exp Bot* **72**, 2514-
2524, doi:10.1093/jxb/eraa583 (2021).
7   Guo, J. & Zhou, H. X. Protein Allostery and Conformational Dynamics. *Chem Rev* **116**, 6503-6515,
doi:10.1021/acs.chemrev.5b00590 (2016).
8   Wodak, S. J. *et al.* Allostery in Its Many Disguises: From Theory to Applications. *Structure* **27**, 566-578,
doi:10.1016/j.str.2019.01.003 (2019).
9   Motlagh, H. N., Wrabl, J. O., Li, J. & Hilser, V. J. The ensemble nature of allostery. *Nature* **508**, 331-339,
doi:10.1038/nature13001 (2014).
10   Willkomm, S. *et al.* Single-molecule FRET uncovers hidden conformations and dynamics of human
Argonaute 2. *Nat Commun* **13**, 3825, doi:10.1038/s41467-022-31480-4 (2022).
11   Asher, W. B. *et al.* GPCR-mediated β-arrestin activation deconvoluted with single-molecule precision.
Cell **185**, 1661-1675.e1616, doi:10.1016/j.cell.2022.03.042 (2022).
12   Hellenkamp, B. *et al.* Precision and accuracy of single-molecule FRET measurements—a multi-
laboratory benchmark study. *Nature Methods* **15**, 669-676, doi:10.1038/s41592-018-0085-0 (2018).
13   Ha, T. *et al.* Probing the interaction between two single molecules: fluorescence resonance energy
transfer between a single donor and a single acceptor. *Proc Natl Acad Sci U S A* **93**, 6264-6268,
doi:10.1073/pnas.93.13.6264 (1996).
14   Kaplan, M. *et al.* EGFR Dynamics Change during Activation in Native Membranes as Revealed by NMR.
Cell **167**, 1241-1251 e1211, doi:10.1016/j.cell.2016.10.038 (2016).
15   Zhang, S. *et al.* USP14-regulated allostery of the human proteasome by time-resolved cryo-EM. *Nature*
605, 567-574, doi:10.1038/s41586-022-04671-8 (2022).
16   Gruhl, T. *et al.* Ultrafast structural changes direct the first molecular events of vision. *Nature* **615**, 939-
944, doi:10.1038/s41586-023-05863-6 (2023).
17   Silva, D. A. *et al.* Millisecond dynamics of RNA polymerase II translocation at atomic resolution. *Proc*
Natl Acad Sci U S A **111**, 7665-7670, doi:10.1073/pnas.1315751111 (2014).
18   Elgeti, M. & Hubbell, W. L. DEER Analysis of GPCR Conformational Heterogeneity. *Biomolecules* **11**,
doi:10.3390/biom11060778 (2021).
19   Seal, M. *et al.* Gd(III)-19F Distance Measurements of Proteins in Cells by Electron-Nuclear Double
Resonance. *Angew Chem Int Ed Engl*, e202218780, doi:10.1002/anie.202218780 (2023).
20   Barford, D. & Johnson, L. N. The allosteric transition of glycogen phosphorylase. *Nature* **340**, 609-616,
doi:10.1038/340609a0 (1989).
21   Massagué, J. & Obenauf, A. C. Metastatic colonization by circulating tumour cells. *Nature* **529**, 298-306,
doi:10.1038/nature17038 (2016).
22   Kantsadi, A. L. *et al.* Glycogen phosphorylase as a target for type 2 diabetes: synthetic, biochemical,
structural and computational evaluation of novel N-acyl-N´-(β-D-glucopyranosyl) urea inhibitors. *Curr*
Top Med Chem **15**, 2373-2389, doi:10.2174/1568026615666150619142253 (2015).
23   Longo, J., Watson, M. J., Vos, M. J., Williams, K. S. & Jones, R. G. PYGBacking on glycogen
metabolism to fuel early memory T cell recall responses. *Mol Cell* **82**, 2918-2921,
doi:10.1016/j.molcel.2022.07.016 (2022).
24   Cori, G. T., Colowick, S. P. & Cori, C. F. The Activity of the Phosphorylating Enzyme in Muscle Extract.
Journal of Biological Chemistry **127**, 771-782, doi:10.1016/s0021-9258(18)73783-6 (1939).
25   Barford, D., Hu, S. H. & Johnson, L. N. Structural mechanism for glycogen phosphorylase control by
phosphorylation and AMP. *Journal of Molecular Biology* **218**, 233-260, doi:https://doi.org/10.1016/0022-
2836(91)90887-C (1991).
26   Klinov, S. V. & Kurganov, B. I. Kinetic Mechanism of Activation of Muscle Glycogen Phosphorylase b by
Adenosine   5′-Monophosphate.   *Archives   of   Biochemistry   and   Biophysics*   **312**,   14-21,
doi:https://doi.org/10.1006/abbi.1994.1274 (1994).
27   Dombrádi, V. Structural aspects of the catalytic and regulatory function of glycogen phosphorylase.
International Journal of Biochemistry **13**, 125-139, doi:https://doi.org/10.1016/0020-711X(81)90147-6
(1981).
28   Fletterick, R. J. & Sprang, S. R. Glycogen phosphorylase structures and function. *Accounts of Chemical*
Research **15**, 361-369, doi:10.1021/ar00083a004 (1982).
29   Johnson, L. N. Glycogen phosphorylase: control by phosphorylation and allosteric effectors. *Faseb j* **6**,
2274-2282, doi:10.1096/fasebj.6.6.1544539 (1992).
30   Leonidas, D. D. *et al.* Glycogen phosphorylase revisited: extending the resolution of the R- and T-state
structures of the free enzyme and in complex with allosteric activators. *Acta Crystallogr F Struct Biol*
Commun **77**, 303-311, doi:10.1107/S2053230X21008542 (2021).
31   Sprang, Withers, S., Goldsmith, E., Fletterick, R. & Madsen, N. Structural basis for the activation of
glycogen   phosphorylase   b   by   adenosine   monophosphate.   *Science*   **254**,   1367-1371,
doi:10.1126/science.1962195 (1991).

32 Sprang, S. R. *et al.* Structural changes in glycogen phosphorylase induced by phosphorylation. *Nature*
336, 215-221, doi:10.1038/336215a0 (1988).
33 Sprang, S. R., Goldsmith, E. J., Fletterick, R. J., Withers, S. G. & Madsen, N. B. Catalytic site of
glycogen phosphorylase: structure of the T state and specificity for .alpha.-D-glucose. *Biochemistry* **21**,
5364-5371, doi:10.1021/bi00264a038 (1982).
34 Klinov, S. V. & Kurganov, B. I. Combined kinetic mechanism describing activation and inhibition of
muscle glycogen phosphorylase b by adenosine 5′-monophosphate. *Biophysical Chemistry* **92**, 89-102,
doi:https://doi.org/10.1016/S0301-4622(01)00189-2 (2001).
35 Helmreich, E. & Cori, C. F. The effects of ph and temperature on the kinetics of the phosphorylase
reaction. *Proc Natl Acad Sci U S A* **52**, 647-654, doi:10.1073/pnas.52.3.647 (1964).
36 Monod, J., Wyman, J. & Changeux, J.-P. On the nature of allosteric transitions: A plausible model.
Journal of Molecular Biology **12**, 88-118, doi:https://doi.org/10.1016/S0022-2836(65)80285-6 (1965).
37 Krebs, E. G. & Fischer, E. H. The phosphorylase b to a converting enzyme of rabbit skeletal muscle.
Biochim Biophys Acta **20**, 150-157, doi:10.1016/0006-3002(56)90273-6 (1956).
38 Oikonomakos, N. G. *et al.* Flavopiridol inhibits glycogen phosphorylase by binding at the inhibitor site.
Journal of Biological Chemistry **275**, 34566-34573 (2000).
39 Oikonomakos, N. G. *et al.* Allosteric inhibition of glycogen phosphorylase a by the potential antidiabetic
drug 3-isopropyl 4-(2-chlorophenyl)-1,4-dihydro-1-ethyl-2-methyl-pyridine-3,5,6-tricarbo xylate. *Protein*
Sci **8**, 1930-1945, doi:10.1110/ps.8.10.1930 (1999).
40 Walcott, S. & Lehman, S. L. Enzyme Kinetics of Muscle Glycogen Phosphorylase b. *Biochemistry* **46**,
11957-11968, doi:10.1021/bi7005527 (2007).
41 Lim, X.-X. *et al.* Conformational changes in intact dengue virus reveal serotype-specific expansion.
Nature Communications **8**, 14339, doi:10.1038/ncomms14339 (2017).
42 Kish, M. *et al.* Online Fully Automated System for Hydrogen/Deuterium-Exchange Mass Spectrometry
with   Millisecond   Time   Resolution.   *Analytical   Chemistry*   **95**,   5000-5008,
doi:10.1021/acs.analchem.2c05310 (2023).
43 Zheng, J. *et al.* Comparative Analysis of Cleavage Specificities of Immobilized Porcine Pepsin and
Nepenthesin II under Hydrogen/Deuterium Exchange Conditions. *Anal Chem* **92**, 11018-11028,
doi:10.1021/acs.analchem.9b05694 (2020).
44 Knox, R., Lento, C. & Wilson, D. J. Mapping Conformational Dynamics to Individual Steps in the TEM-1
beta-Lactamase Catalytic Mechanism. *J Mol Biol* **430**, 3311-3322, doi:10.1016/j.jmb.2018.06.045
(2018).
45 Sotiroudis, T. G., Oikonomakos, N. G. & Evangelopoulos, A. E. Effect of sulfated polysaccharides and
sulfate anions on the AMP-dependent activity of phosphorylase b. *Biochemical and Biophysical*
Research Communications **90**, 234-239, doi:https://doi.org/10.1016/0006-291X(79)91615-2 (1979).
46 Engers, H. D. & Madsen, N. B. The effect of anions on the activity of phosphorylase b. *Biochemical and*
Biophysical Research Communications **33**, 49-54, doi:https://doi.org/10.1016/0006-291X(68)90253-2
(1968).
47 Leonidas, D. D. *et al.* The ammonium sulfate activation of phosphorylase b. *FEBS Letters* **261**, 23-27,
doi:https://doi.org/10.1016/0014-5793(90)80627-U (1990).
48 Schreiber, W. E. & Bowling, S. An automated assay of glycogen phosphorylase in the direction of
phosphorolysis. *Ann Clin Biochem* **27 (Pt 2)**, 129-132, doi:10.1177/000456329002700207 (1990).
49 Seetaloo, N., Kish, M. & Phillips, J. J. HDfleX: Software for Flexible High Structural Resolution of
Hydrogen/Deuterium-Exchange   Mass   Spectrometry   Data.   *Anal   Chem*   **94**,   4557-4564,
doi:10.1021/acs.analchem.1c05339 (2022).
50 Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825-2830
(2011).
51 Guo, Z. *et al.* Engineering and exploiting synthetic allostery of NanoLuc luciferase. *Nat Commun* **13**,
789, doi:10.1038/s41467-022-28425-2 (2022).
52 Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods*
17, 261-272, doi:10.1038/s41592-019-0686-2 (2020).
53 Acharya, K. R., Stuart, D. I., Varvill, K. M. & Johnson, L. N.
54 Mitchell, E. P. *et al.* Ternary complex crystal structures of glycogen phosphorylase with the transition
state analogue nojirimycin tetrazole and phosphate in the T and R states. *Biochemistry* **35**, 7341-7355,
doi:10.1021/bi960072w (1996).
55 Huang, J. *et al.* Insights into Phosphorylation-Induced Protein Allostery and Conformational Dynamics of
Glycogen Phosphorylase via Integrative Structural Mass Spectrometry and In Silico Modeling. *ACS*
Chem Biol **17**, 1951-1962, doi:10.1021/acschembio.2c00393 (2022).
56 Hageman, T. S. & Weis, D. D. Reliable Identification of Significant Differences in Differential Hydrogen
Exchange-Mass Spectrometry Measurements Using a Hybrid Significance Testing Approach. *Anal*
Chem **91**, 8008-8016, doi:10.1021/acs.analchem.9b01325 (2019).
57 Weiss, D. R. & Levitt, M. Can morphing methods predict intermediate structures? *J Mol Biol* **385**, 665-
674, doi:10.1016/j.jmb.2008.10.064 (2009).
58 Tam, B., Sinha, S. & Wang, S. M. Combining Ramachandran plot and molecular dynamics simulation
for structural-based variant classification: Using TP53 variants as model. *Comput Struct Biotechnol J* **18**,
4033-4039, doi:10.1016/j.csbj.2020.11.041 (2020).

59   Kasvinsky, P. J., Madsen, N. B., Sygusch, J. & Fletterick, R. J. The regulation of glycogen
phosphorylase alpha by nucleotide derivatives. Kinetic and x-ray crystallographic studies. *Journal of*
Biological Chemistry **253**, 3343-3351, doi:https://doi.org/10.1016/S0021-9258(17)40842-8 (1978).
60   Bai, Y., Milne, J. S., Mayne, L. & Englander, S. W. Primary structure effects on peptide group hydrogen
exchange. *Proteins* **17**, 75-86, doi:10.1002/prot.340170110 (1993).
61   Deniz, E. *et al.* Through bonds or contacts? Mapping protein vibrational energy transfer using non-
canonical amino acids. *Nature Communications* **12**, 3284, doi:10.1038/s41467-021-23591-1 (2021).
62   Rezaei, H. *et al.* Sequential generation of two structurally distinct ovine prion protein soluble oligomers
displaying different biochemical reactivities. *J Mol Biol* **347**, 665-679, doi:10.1016/j.jmb.2005.01.043
(2005).
63   Fiser, A., Do, R. K. G. & Šali, A. Modeling of loops in protein structures. *Protein Science* **9**, 1753-1773,
doi:https://doi.org/10.1110/ps.9.9.1753 (2000).
64   Sali, A. & Blundell, T. L. Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol*
234, 779-815, doi:10.1006/jmbi.1993.1626 (1993).

ACKNOWLEDGEMENTS

626 J.J.P. and M.K. are supported by a UKRI Future Leaders Fellowship [Grant Number MR/T02223X/1]. DI is
627 funded by a BBSRC SWBioDTP studentship. For the purpose of open access, the author has applied a CC BY
628 public copyright license to any Author Accepted Manuscript version arising from this submission.

AUTHOR INFORMATION

AUTHORS AND AFFILIATIONS

Living Systems Institute, Department of Biosciences, University of Exeter, Stocker Road, Exeter, EX4 4QD, UK

Monika Kish, Dylan P. Ivory, Jonathan J. Phillips

Alan Turing Institute, British Library, London, NW1 2DB, UK

Jonathan J. Phillips

CONTRIBUTIONS

637 JJP conceived the study. MK and JJP designed the experiments and performed computational structural
638 interpolations. MK, DI and JJP performed mass spectrometry experiments. MK and DI performed
639 spectrophotometry assays, constructed atomistic models. MK performed clustering analysis, analyzed mass
640 spectrometry data, and performed data and statistical analysis. All authors wrote the manuscript.

CORRESPONDING AUTHOR

Correspondence to JJP: jj.phillips@exeter.ac.uk

CODE AVAILABILITY STATEMENT

This study uses in-house developed scripts in Python. Available upon reasonable request.

ETHICS DECLARATIONS

COMPETING INTERESTS

The authors declare no competing interests.

FIGURES

 Figure 1

Figure 1: Allosteric activation of glycogen phosphorylase b and inhibition of glycogen phosphorylase a.
(A) Glycogen hydrolysis kinetics for GlyPa (black), GlyPb (blue) and GlyPb* (orange; GlyPb equilibrated with 25
mM AMP and 25 mM AS). Absorbance (A.U.) at 30 ˚C for +60 min at 450 nm. Mean and SD from n=2. Plots for
all conditions in Figure S2A. **(B)** Glycogen hydrolysis kinetics for GlyPa (black), GlyPb (blue) and GlyPa* (orange;
GlyPa equilibrated with 32 mM caffeine. Absorbance (A.U.) at 30 ˚C for +60 min at 450 nm. Mean and SD from
n=3. Plots for all conditions in Figure S2B. **(C)** Heat map of measured activity for combinations of AMP/AS
concentrations in range 0-100 mM. Inactive GlyPb (black); highest activity achieved (white). Activity was
calculated by a standard curve for n=2. **(D)** Heat maps for structural perturbations in the presence of AMP/AS,
detected by equilibrium HDX-MS for the nucleotide site **(T38-Y51)**, tower helix **(V266-I275)** and 380s loop **(Y274-**
L384). Deuterium uptake data from the HDX screen was used to produce these heat maps, n=3. The relative
deuteration level for each sample is color coded shown on the scale right; note different normalization. **(E)**
Catalytic activity of GlyPb supplemented with AS and AMP. Shaded region – GlyPa 95% confidence intervals. **(F)**
Catalytic activity of GlyPa supplemented with caffeine. Shaded region – GlyPb 95% confidence intervals. **(G)**
AMP binding saturation during the dead time of the HDX labelling (50 ms). Non-equilibrium addition of 25 mM
AMP/25 mM AS is optimal to fully saturate the nucleotide site. **(H)** Caffeine binding saturation during the dead
time of the HDX labelling (50 ms). Non-equilibrium addition of 32 mM caffeine is optimal to fully saturate the
inhibitor site. **(I)** Ammonium sulfate influence on the tower helix and nucleotide site. HDX labelling data was fitted
and plotted versus the state (Figure S3). States annotated include: GlyPb - apo, * - equilibrium activated with 25
mM AMP and 25 mM AS, *AS - equilibrium activated with 25 mM AS, ^ eq AS – equilibrated with 25 mM AS and
then non- equilibrium activated with AMP, ^ - non-equilibrium activated with 25 mM AMP and 25 mM AS, ^AMP –
non-equilibrium activated with 25 mM AMP, ^AS – non- equilibrium activated with 25 mM AS.

Figure 2

Figure 2: Clusters of structural kinetic behavior in GlyP. (A) The sum difference of the D uptake per amino
acid between non-equilibrium and GlyPb/active state plotted against x/y axes, respectively. Amino acids were
grouped into seven clusters (a-g) resulting from a k-means hierarchical analysis. The centroid of each cluster is
denoted by the letter connected to each member by an edge. Raw D-labelling uptake data was centroided,
corrected for maximum exchangeable amide hydrogens, normalized and the different states were pairwise
subtracted. Dendrogram of clustering in A is shown on the right. Y-axis is Euclidean distance between the
clusters (arbitrary units). The number of amino acid members in each cluster is indicated at the bottom of each
cluster (truncated for clarity; Figure S15A). **(B)** Clusters (excluding g and a) represented on GlyPb active state
structure (PDB:3E3N). Notable features are indicated on the left subunit molecular surface; Notable secondary
structural elements indicated. **(C)** The sum difference of the D uptake per amino acid between non-equilibrium
and GlyPa/inactive state plotted against x/y axes, respectively. Amino acids were grouped into seven clusters (h-
n) resulting from a k-means hierarchical analysis. The centroid of each cluster is denoted by the letter connected
to each member by an edge. Raw D-labelling uptake data was centroided, corrected for maximum exchangeable
amide hydrogens, normalized and the different states were pairwise subtracted. Dendrogram of clustering in C is
shown on the right. Y-axis is Euclidean distance between the clusters (arbitrary units). The number of amino acid
members in each cluster is indicated at the bottom of each cluster (truncated for clarity; Figure S15B). **(D)**
Clusters (excluding h and j) represented on GlyPa inactive state structure (PDB:1GFZ). Notable features are
indicated on the left subunit molecular surface; Notable secondary structural elements indicated.

Figure 3: Local structural kinetics upon allosteric activation and inhibition revealed by non-equilibrium D-
labeling. (A) Hydrogen exchange kinetics of representative peptide segments for each cluster. Mean of n=3;
error bars mostly contained within data points. GlyPb (blue); fully equilibrated activated with 25 mM AMP and 25
mM AS (orange); non-equilibrium activated with 25 mM AMP and 25 mM AS (green). **(B)** Clusters mapped on the
crystal structure of inactive GlyPb (3E3N.pdb). Detailed annotation is represented on each structure per cluster.
(C) Hydrogen exchange kinetics of representative peptide segments for each cluster. Mean of n=3; error bars
mostly contained within data points. GlyPa (blue); fully equilibrated inactivated with 32 mM caffeine (orange);
non-equilibrium inactivated with 32 mM caffeine (green). **(D)** Clusters mapped on the crystal structure of inactive
GlyPa (1GFZ.pdb). Detailed annotation is represented on each structure per cluster.

Figure 4

Figure 4: Structural interpolation between (in)active states of GlyP. (A) Tower helix and adjacent loops in
inactive (left panel), transitional (middle) and active (right) conformations. Transitional structures with multiple
broken hydrogen-bonds (pink lines) in the α7 helix are observed upon interpolation between apo GlyPb – GlyPb-
AMP (activation) and between apo GlyPa - GlyPa-caffeine (inhibition). Representative structure shown; full
trajectories provided as supplementary files. Bottom and top heat maps represent the hydrogen-bond network of
each structure upon activation (top) and inhibition (bottom) in this region of the enzyme, gray: H-bond; white: no
H-bond. **(B)** Interpolation trajectory represented as rmsd between each structure. Calculated only for the tower
helix region (amino acids 256:288). Normalized to maximum values from the AMP-activation (lower triangle) and
caffeine-inhibition (upper triangle), respectively. **(C)** Proposed model of the allosteric modification of GlyP. Both
modifications, activation (AMP) and inhibition (CFF) of GlyPb and GlyPa, respectively, undergo similar transition
mechanism, with common structural features and proximal to the active state.

 FIGURE LEGENDS

Figure 1: Allosteric activation of glycogen phosphorylase b and inhibition of glycogen phosphorylase a.
(A) Glycogen hydrolysis kinetics for GlyPa (black), GlyPb (blue) and GlyPb* (orange; GlyPb equilibrated with 25
mM AMP and 25 mM AS). Absorbance (A.U.) at 30 ˚C for +60 min at 450 nm. Mean and SD from n=2. Plots for
all conditions in Figure S2A. **(B)** Glycogen hydrolysis kinetics for GlyPa (black), GlyPb (blue) and GlyPa* (orange;
GlyPa equilibrated with 32 mM caffeine. Absorbance (A.U.) at 30 ˚C for +60 min at 450 nm. Mean and SD from
n=3. Plots for all conditions in Figure S2B. **(C)** Heat map of measured activity for combinations of AMP/AS
concentrations in range 0-100 mM. Inactive GlyPb (black); highest activity achieved (white). Activity was
calculated by a standard curve for n=2. **(D)** Heat maps for structural perturbations in the presence of AMP/AS,
detected by equilibrium HDX-MS for the nucleotide site **(T38-Y51)**, tower helix **(V266-I275)** and 380s loop **(Y274-**
L384). Deuterium uptake data from the HDX screen was used to produce these heat maps, n=3. The relative
deuteration level for each sample is color coded shown on the scale right; note different normalization. **(E)**
Catalytic activity of GlyPb supplemented with AS and AMP. Shaded region – GlyPa 95% confidence intervals. **(F)**
Catalytic activity of GlyPa supplemented with caffeine. Shaded region – GlyPb 95% confidence intervals. **(G)**
AMP binding saturation during the dead time of the HDX labelling (50 ms). Non-equilibrium addition of 25 mM
AMP/25 mM AS is optimal to fully saturate the nucleotide site. **(H)** Caffeine binding saturation during the dead
time of the HDX labelling (50 ms). Non-equilibrium addition of 32 mM caffeine is optimal to fully saturate the
inhibitor site. **(I)** Ammonium sulfate influence on the tower helix and nucleotide site. HDX labelling data was fitted
and plotted versus the state (Figure S3). States annotated include: GlyPb - apo, * - equilibrium activated with 25
mM AMP and 25 mM AS, *AS - equilibrium activated with 25 mM AS, ^ eq AS – equilibrated with 25 mM AS and
then non- equilibrium activated with AMP, ^ - non-equilibrium activated with 25 mM AMP and 25 mM AS, ^AMP –
non-equilibrium activated with 25 mM AMP, ^AS – non- equilibrium activated with 25 mM AS.

Figure 2: Clusters of structural kinetic behavior in GlyP. (A) The sum difference of the D uptake per amino
acid between non-equilibrium and GlyPb/active state plotted against x/y axes, respectively. Amino acids were
grouped into seven clusters (a-g) resulting from a k-means hierarchical analysis. The centroid of each cluster is
denoted by the letter connected to each member by an edge. Raw D-labelling uptake data was centroided,
corrected for maximum exchangeable amide hydrogens, normalized and the different states were pairwise
subtracted. Dendrogram of clustering in A is shown on the right. Y-axis is Euclidean distance between the
clusters (arbitrary units). The number of amino acid members in each cluster is indicated at the bottom of each
cluster (truncated for clarity; Figure S15A). **(B)** Clusters (excluding g and a) represented on GlyPb active state
structure (PDB:3E3N). Notable features are indicated on the left subunit molecular surface; Notable secondary
structural elements indicated. **(C)** The sum difference of the D uptake per amino acid between non-equilibrium
and GlyPa/inactive state plotted against x/y axes, respectively. Amino acids were grouped into seven clusters (h-
n) resulting from a k-means hierarchical analysis. The centroid of each cluster is denoted by the letter connected
to each member by an edge. Raw D-labelling uptake data was centroided, corrected for maximum exchangeable
amide hydrogens, normalized and the different states were pairwise subtracted. Dendrogram of clustering in C is
shown on the right. Y-axis is Euclidean distance between the clusters (arbitrary units). The number of amino acid
members in each cluster is indicated at the bottom of each cluster (truncated for clarity; Figure S15B). **(D)**
Clusters (excluding h and j) represented on GlyPa inactive state structure (PDB:1GFZ). Notable features are
indicated on the left subunit molecular surface; Notable secondary structural elements indicated.

Figure 3: Local structural kinetics upon allosteric activation and inhibition revealed by non-equilibrium D-
labeling. (A) Hydrogen exchange kinetics of representative peptide segments for each cluster. Mean of n=3;
error bars mostly contained within data points. GlyPb (blue); fully equilibrated activated with 25 mM AMP and 25
mM AS (orange); non-equilibrium activated with 25 mM AMP and 25 mM AS (green). **(B)** Clusters mapped on the
crystal structure of inactive GlyPb (3E3N.pdb). Detailed annotation is represented on each structure per cluster.
(C) Hydrogen exchange kinetics of representative peptide segments for each cluster. Mean of n=3; error bars
mostly contained within data points. GlyPa (blue); fully equilibrated inactivated with 32 mM caffeine (orange);
non-equilibrium inactivated with 32 mM caffeine (green). **(D)** Clusters mapped on the crystal structure of inactive
GlyPa (1GFZ.pdb). Detailed annotation is represented on each structure per cluster.

Figure 4: Structural interpolation between (in)active states of GlyP. (A) Tower helix and adjacent loops in
inactive (left panel), transitional (middle) and active (right) conformations. Transitional structures with multiple
broken hydrogen-bonds (pink lines) in the α7 helix are observed upon interpolation between apo GlyPb – GlyPb-
AMP (activation) and between apo GlyPa - GlyPa-caffeine (inhibition). Representative structure shown; full
trajectories provided as supplementary files. Bottom and top heat maps represent the hydrogen-bond network of
each structure upon activation (top) and inhibition (bottom) in this region of the enzyme, gray: H-bond; white: no
H-bond. **(B)** Interpolation trajectory represented as rmsd between each structure. Calculated only for the tower
helix region (amino acids 256:288). Normalized to maximum values from the AMP-activation (lower triangle) and
caffeine-inhibition (upper triangle), respectively. **(C)** Proposed model of the allosteric modification of GlyP. Both
modifications, activation (AMP) and inhibition (CFF) of GlyPb and GlyPa, respectively, undergo similar transition
mechanism, with common structural features and proximal to the active state.