Contrasting responses to aridity by different-sized decomposers cause similar decomposition rates across a precipitation gradient

Viraj R. Torsekar*[1,2,]†, Nevo Sagi*[1,]†, J Alfred Daniel[3], Yael Hawlena[1], Efrat Gavish-Regev[3] & Dror Hawlena[1,3]

[1] Risk-Management Ecology Lab, Department of Ecology, Evolution & Behavior,
  The Alexander Silberman Institute of Life Sciences, The Hebrew University of Jerusalem.

[2] Theoretical Ecology and Evolution Lab, Centre for Ecological Sciences, Indian Institute of Science.

[3] The National Natural History Collections, The Hebrew University of Jerusalem.

*Contact information:

Viraj R. Torsekar - viraj.torsekar@mail.huji.ac.il

Nevo Sagi - nevo.sagi@mail.huji.ac.il

† Viraj R. Torsekar and Nevo Sagi contributed equally to the manuscript and are both corresponding authors.

Authors' contributions

D.H., N.S. and V.T. conceived this study and designed the experiments. V.T, N.S. and D.H performed the field experiment. N.S. and V.T. performed lab analyses. J.A.D., Y.H. and E.G.R.

sorted and identified animal samples. N.S. analyzed the data. N.S., V.T and D.H wrote the

manuscript, and all authors provided critical feedback and approved submission.

Data sharing plans

All data used in the manuscript has been deposited in an open repository and can be found on the

following link https://figshare.com/s/96ff438c98ae4a6afbc4

Type of article: Letter

Abstract

Litter decomposition is expected to be positively associated with precipitation despite evidence that decomposers of varying sizes have different moisture dependencies. We hypothesized that higher tolerance of macro-decomposers to aridity may counterbalance the effect of smaller decomposers, leading to similar decomposition rates across climatic gradients. We tested this hypothesis by placing plant litter baskets of different mesh sizes in seven sites along a sharp precipitation gradient, and by characterizing the macro-decomposer assemblages using pitfall trapping. We found that decomposers responded differently to precipitation levels based on their size. Microbial decomposition increased with precipitation in the winter while macro-decomposition peaked in arid sites during the summer. This led to similar overall decomposition rates across the gradient except in hyper-arid sites. Macro-decomposer richness, abundance and biomass peaked in arid environments. Our findings highlight the importance of macro-decomposition in arid-lands, possibly resolving the dryland decomposition conundrum, and emphasizing the need to contemplate decomposer size when investigating zoogeochemical processes.

Keywords

litter decomposition, climate, detritivores, dryland, zoogeochemistry, dryland decomposition conundrum

Introduction

Litter decomposition is a key process determining elemental cycling in terrestrial ecosystems[1]. Decomposition is controlled by climate, litter quality and origin, and the identity and abundance of microbial and faunal decomposers[2-4]. Climate regulates decomposition rates directly, but also indirectly by influencing food-web structure and dynamics[5]. Thus, understanding how climate and decomposers interact is a key step in explaining variation in plant litter decomposition across ecosystems and seasons, and in forecasting the consequences of climate change and biodiversity loss on elemental cycling.

Theory suggests that decomposition is positively associated with moisture and temperature[4]. Cross-site studies, reviews, and meta-analyses verified this global pattern, showing that plant litter decomposition in microbial litter bags is indeed faster under warm and wet conditions than under cool and dry conditions[6-11]. This well accepted realization implicitly assumes that microorganisms dominate plant litter decomposition, largely ignoring the growing recognition that animals may play an important role in litter cycling. This role includes mineralizing and excreting assimilated plant nutrients, fragmenting and partly decomposing plant material, and transporting detritus to microbial havens[12-19].

Attempts to explore how climate affects faunal decomposition revealed a similar positive association with temperature and precipitation[20,21]. This global pattern, however, may be confounded by compiling the effect of all decomposer fauna together, ignoring the well-established understanding that soil animals of various size groups respond differently to climate[4,22]. Specifically, larger arthropods can survive and remain active during hot and dry periods when smaller organisms cannot[23]. Indeed, handful evidence shows that macro-

detritivorous arthropods dominate litter and wood decomposition in warm drylands, especially

during warm and dry seasons[24–26]. This suggests that the conceptualization of how animals and

climate interact to regulate decomposition rates requires considering the effects of meso-

decomposers and those of macro-decomposers separately, particularly in warm drylands.

Detritivorous animals are expected to be exceptionally abundant in arid ecosystems

where plant detritus is prevalent year around but green plant material is available predominantly

in short pulses following precipitation events[27]. Macrofauna are physiologically and

morphologically more adapted to aridity than mesofauna[23]. Moreover, their large size enables

them to remain active during long dry periods by shuttling between existing and self-engineered

climatic havens and the hostile foraging grounds on the surface[28]. Consequently, macro-

decomposition should be especially important in hot moisture-deprived habitats and periods,

whereas the activity of microbes and mesofauna is expected to be minimal.

The predicted negative association between moisture and macro-decomposition in

drylands may be reversed in hyper-arid environments. In these environments, the extreme

climatic conditions and scarce and unpredictable plant litter availability may limit macro-

decomposer populations, diminishing macro-decomposition rates with increasing aridity.

Consequently, and in sharp contrast to smaller organisms, macro-decomposition should follow a

hump-shaped response to precipitation that peaks in arid ecosystems.

To test this novel hypothesis, we examined the climate dependency of plant litter

decomposition by microorganisms, meso-decomposers and macro-decomposers along a sharp

aridity gradient spanning from mean annual precipitation of only 22 mm to 526 mm. This

gradient represents hyper-arid, arid, semiarid and dry sub-humid Mediterranean climates. We

repeated the experiment during hot summer with no precipitation and again during cooler and

wetter winter. We hypothesized that both microbial and mesofaunal decomposition should

increase with increasing precipitation during the winter, but during the dry summer contribute

only minimally to plant litter decomposition across the aridity gradient. In the dry summer,

macrofaunal decomposition should follow a hump-shaped response to precipitation, increasing

from hyper-arid to arid sites and decreasing gradually in more mesic semi-arid and

Mediterranean sites. We also predicted that the opposing climatic dependencies of macrofauna

and microorganisms and mesofauna should lead to similar overall decomposition rates across the

precipitation gradient except in the hyper-arid sites in which decomposers activity is predicted to

be minimal regardless of organism size (**Fig. 1**). To reveal the mechanism, we sampled macro-

decomposers across the aridity gradient and the two seasons, using pitfall traps. We predicted

hump-shaped relationships between precipitation and the abundance, richness, and biomass of

macro-decomposers that peak in arid ecosystems.

Figure 1 - Hypothetical climate dependence of litter decomposition by microorganisms and
mesofauna (dotted orange curve), by macrofauna (grey – dash), and by the whole decomposer
community (maroon - solid).

Methods

Methods

We performed a manipulative litter mass loss experiment across seven sites representing

a sharp mean annual precipitation (MAP) gradient ranging from hyper-arid desert to

Mediterranean maquis (Fig. 2A, Table 1). All sites were chosen to be on calcareous soils formed

upon sedimentary limestone rock in natural habitats. The mean annual temperature varies only

slightly across sites from 18.7ºC to 22.3 ºC. The exact study sites were determined to ensure

minimal human disturbance during the year-long experiment. In each of the seven sites we

installed litter baskets of three different mesh sizes that control organismal access to litter:

Micro-baskets allowing entry of only microorganisms (<200 μm), meso-baskets allowing entry

of microorganisms and mesofauna (<2 mm), and macro-baskets that were identical to the meso-

baskets but with side openings that allow entry of macrofauna (<2 cm). Litter baskets were filled

with leaf litter belonging to the annual grass *Stipa capensis* Thunb. that is native to all seven

study sites. This approach allowed us to compare decomposition rates across sites without using

exogenous litter that may decompose in a very different rate than local litter[3]. Twenty-five

blocks, each including the three basket types (Fig. 2B), were installed in each site for two

consecutive experimental periods - a wet cool winter and a dry hot summer (2 periods x 7 sites x

3 treatments x 25 blocks = 1050 baskets in total). We also characterized the macro-decomposer

assemblage in each site during the two seasons using pitfall trapping.

Table 1 - Properties of the seven experimental sites

Site	Abb.	Coordinates	MAT* [□]	MAP* [mm]	AI$_U$* (MAP/PET)	Climate	Winter experiment	Summer experiment
Ramat Hanadiv	RH	32°33'22.4"N 34°56'26.6"E	20.2	526	0.518	Dry subhumid Mediterranean	3.12.2020 – 27.6.2021	27.6 – 27.10.2021
Bet Guvrin	BG	31°35'54.7"N 34°54'14.2"E	20.9	403	0.370	Semiarid	2.12.2020 – 13.6.2021	13.6 – 21.10.2021
Havat Shikmim	HS	31°30'49.7"N 34°41'18.8"E	19.8	367	0.364	Semiarid	2.12.2020 – 13.6.2021	13.6 – 21.10.2021
Sayeret Shaked	SS	31°16'05.7"N 34°39'12.9"E	20.0	148	0.145	Arid	26.11.2020 – 23.5.2021	23.5 – 21.10.2021
Avdat	AV	30°47'02.3"N 34°46'13.3"E	18.7	84	0.089	Arid	26.11.2020 – 23.5.2021	23.5 – 21.10.2021
Meishar	MS	30°27'04.2"N 34°56'03.0"E	20.8	33	0.029	Hyperarid	10.12.2020 – 12.7.2021	12.7 – 8.11.2021
Nahal Shita	NS	30°08'29.4"N 35°07'36.6"E	22.3	22	0.017	Hyperarid	10.12.2020 – 12.7.2021	12.7 – 8.11.2021

129 * Climatic data extracted from http://www.meteo-tech.co.il/hanadiv_new/hanadiv_en.asp (RH),
130   courtesy of Shaily Dor-Haim (SS), and extracted from https://ims.gov.il/en (all other sites).

Figure 2 - (A) Locations and landscapes of the seven experimental sites across a precipitation gradient from 22 to 526 mm mean annual precipitation (MAP). (B) A block of three litter baskets in the Sayeret Shaked site. Macro-basket in front, meso-basket on the right and micro-basket on the left. The precipitation map courtesy of the Hebrew University GIS center.

Litter basket experiment

We collected *S. capensis* litter from the Avdat site in the summer of 2020 and air dried it.

We sorted the litter to remove litter belonging to any other species and assigned 3 ± 0.0001 g

(Mettler Toledo MS105DU) to each litter basket. Thirty additional litter samples were oven dried

at 60 ℃ for 48 hours and weighed again for determination of initial moisture content. The 14 x

13 x 3.6 cm litter baskets were prepared of a 12 mm mesh galvanized welded metal, lined at the

bottom with a 1.5 mm fiberglass mesh to prevent litter loss, and covered from all sides (including

top and bottom) with a 2 mm metal mesh to exclude termites. In the macro-baskets, three 2 x 2

cm windows were cut at each of the four sides. The windows were cut approximately 1 cm above

ground level to allow macrofaunal access but prevent accidental litter spill. This may slightly

reduce macrofaunal access, making our estimations of the macrofaunal effect conservative. In

the micro-baskets, we placed the litter within a polyethylene 200 μm mesh bag. In the macro-

and meso-baskets, we laid a 2 cm heavy metal mesh over the litter to minimize litter loss due to

wind.

We installed the first batch of 525 litter baskets in the field in November-December 2020.

All blocks were placed around similarly sized bushes of locally distributed species and tethered

to the ground using metal stakes. We collected the baskets in May-July 2021 and replaced them

with a new similar batch that was later collected in October-November 2021. At the end of each

season, the collected baskets were transported to the laboratory in sealed Ziplock bags. Any litter

spilled during transportation was weighed and the weight loss was incorporated in the

calculations. Leaf litter in each basket was first screened for adulteration from leaf litter of other

species, following which the *S. capensis* litter was oven dried at 60 ℃ for 48 hours and weighed.

To account for dust accumulation on the litter we applied an ash correction procedure[29]. We

burned and weighed five sub-samples from each site-season-treatment combination (550 ℃ for 5

hours) and calculated the combination-specific mean ash content. The final litter mass was

corrected for ash content based on these calculations. We burned and weighed 15 additional

samples of *S. capensis* litter that were not placed in the field and calculated the mean ash content

of the initial litter. The initial litter mass was corrected accordingly. The rate of litter removal

from each basket was calculated as the difference between the ash corrected final dry litter mass

and the ash and moisture corrected initial litter mass, divided by the number of days the litter

spent in the field.

Pitfall trapping

We characterized the macro-decomposer assemblages by setting up 20 pitfall traps for 5-

7 days at each site during each experimental period. Wet season traps were opened in February

2021, whereas the dry season traps were opened between late August and early October. We

installed traps by placing two 10 cm diameter X 7.5 cm deep plastic containers one inside

another such that the opening was flushed with the ground. We added to each trap 150 ml of

preservative, which comprised of 40% absolute ethanol, 40% distilled water and 20% Propylene

glycol. Traps were covered with steel mesh of large mesh size to prevent small mammals and

reptiles from falling inside. At the end of the 5-7 days, samples were collected and transferred to

70% ethanol. Samples were sorted and identified to morphospecies level in the lab. Only animals

larger than 2 mm were included in the analysis. Sub-samples were freeze-dried and weighed

(Mettler Toledo MS105DU) for biomass estimation of each morphospecies.

Analytical procedures

We first fitted a linear mixed model (LMM) to the litter removal rate data, including

experimental site, experimental season, mesh size and all interaction terms as fixed effects. The

random effect of the experimental spatial blocks was found insignificant using a simulation test

with 9999 simulations (RLRT=2.1, P-value=0.07). Therefore, we assessed the effects of the site,

season and mesh size on the litter removal rate using a full factorial analysis of variance,

followed by Tukey HSD pairwise comparisons. We calculated the contribution of each size

group to litter mass loss by block. Microbial contribution was defined as the mass loss from

micro-baskets; mesofaunal contribution was calculated as the difference between mass loss from

meso- and micro-baskets; macrofaunal contribution was calculated as the difference between

mass loss from macro- and meso-baskets; whole-community decomposition was defined as the

mass loss from macro-baskets. We modeled the relationship between MAP and each of these

contributions using Locally Estimated Scatterplot Smoothing (LOESS). We used the same

method to model the relationship between MAP and the macro-decomposer abundance, biomass

and morphospecies richness in each season. We assessed differences in the macro-decomposer

assemblage among experimental sites and seasons using a principal coordinates analysis (PCoA)

with individual traps as the sampling units and Bray-Curtis index (BC) as the dissimilarity

metric. We tested for differences across sites and seasons in macro-decomposer assemblage

using a permutational multivariate analysis of variance (PERMANOVA), followed by pairwise

comparisons between sites using the Benjamini-Hochberg P-value adjustment. BC indices

between site-season combinations were calculated as well, based on the summed abundances

across traps. To explore which macro-decomposer groups dominate the different sites and

seasons, we classified the identified morphospecies to ten macro-decomposer taxa:

Archaeognatha (bristletails), Coleoptera (beetles), Diplopoda (millipedes), Formicidae (ants),

Gastropoda (snails and slugs), Grylloidea (crickets), Isoptera (termites), Lumbricina

(earthworms), Oniscidea (woodlice) and Zygentoma. Then we summed the abundance, richness

and biomass from each group in each trap. We used the abundance data to fit the group scores

onto the PCoA ordination. Litter removal data was analyzed using the 'stats', 'lme4'[30] and

'RLRsim'[31] packages from R software (version 4.3.0)[32], whereas assemblage data was analyzed

using the 'vegan' package[33].

Results

Litter removal rate differed across seasons, sites and mesh sizes, and all interactions

between these factors were found significant as well (**Table 2**). On average, the litter removal

rate was 2.6-fold higher in winter than in summer, 1.6-fold higher in meso- than in micro-baskets

and 1.3-fold higher in macro- than in meso-baskets. Litter removal was negligible in the

hyperarid sites during both seasons, while it was highest in the arid sites during summer and in

the more mesic sites during winter (**Fig. 3**). Within site and season comparisons between mesh

sizes yielded significant differences only in Avdat, Sayeret Shaked and Havat Shikmim,

indicating that faunal effects on decomposition were found only under arid to dry-semiarid

conditions (**Fig. 3**). Both macro- and mesofaunal effects were detected in the arid sites (Avdat,

Sayeret Shaked), whereas the semiarid Havat Shikmim site exhibited only a mesofaunal effect

during both seasons (**Fig. 3**). The macrofaunal, mesofaunal and microorganismal contributions to

litter mass loss peaked under arid, semiarid and Mediterranean climate conditions, respectively

(**Fig. 4**). Whole-community litter removal rates were dictated by microorganisms in winter and

by macrofauna in summer, resulting in comparable rates across the aridity gradient from

Mediterranean to arid climate at the annual scale (**Fig. 4**). In total, the whole-community litter

removal rate peaked in Sayeret Shaked (MAP=148 mm) and significantly decreased under drier

and wetter conditions (**Fig. 4**).

Macro-decomposer abundance, biomass and morphospecies richness peaked in the arid

sites during both seasons (**Fig. 5**). The macro-decomposer assemblage differed significantly

across sites ($F_{6,229}$=10.6, P-value<0.001) and across seasons ($F_{1,229}$=13.1, P-value<0.001), where

woodlice, millipedes and snails were substituted by ants and termites with increasing aridity

(**Fig. 6A, B**). Assemblage was significantly affected by the interaction between site and season

too ($F_{6,229}$=5.4, P-value<0.001). The experimental site explained much of the assemblage

variability across traps (R^2=0.18), whereas experimental season accounted for a smaller fraction

(R^2=0.04), and site-season interaction played an intermediate role (R^2=0.10). All pairwise

comparisons across sites yielded significant differences in assemblage (**Table S1**). In general,

ants were the most abundant group, whereas beetles accounted for most of the biomass.

However, under mesic conditions, woodlice (Ramat Hanadiv site), millipedes (Ramat Hanadiv

and Bet Guvrin) and snails (Havat Shikmim) were dominant (**Fig. 6C**). The Ramat Hanadiv

assemblage was distinctively different from all other sites (**Fig. 6A**), as demonstrated by very

high Bray-Curtis dissimilarity indices compared to the other sites, regardless of the season

(**Table S2**). There were parallels between the spatial and temporal axes of aridity, as winter

communities in the most arid sites (Nahal Shita, Meishar and Avdat) were relatively similar to

the summer communities of the more mesic sites (Sayeret Shaked, Havat Shikmim and Bet

Guvrin) (**Fig. 6A,B**; **Table S2**). The arid sites, where macro-decomposer assemblages flourished

and were responsible for the highest litter mass loss, showed interesting seasonal dynamics.

Bray-Curtis dissimilarity across seasons was higher in Sayeret Shaked than in Avdat (BC=0.79

and 0.72, respectively). Cross-site dissimilarity between Sayeret Shaked and Avdat was higher in

winter than in summer (BC=0.85 and 0.72, respectively).

Table 2 - Results of a full-factorial analysis of variance in litter removal rate across mesh sizes,
experimental sites and seasons.

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	6	0.002092	0.000349	171.946	<0.001
Season	1	0.000679	0.000679	334.94	<0.001
Mesh size	2	0.000267	0.000133	65.77	<0.001
Site : Season	6	0.00112	0.000187	92.044	<0.001
Site : Mesh size	12	0.000392	3.27E-05	16.11	<0.001
Season : Mesh size	2	1.77E-05	8.9E-06	4.368	0.0129
Site : Season : Mesh size	12	0.000129	1.08E-05	5.306	<0.001
Residuals	1008	0.002044	0.000002		

252
Figure 3 - Litter removal rate (mean ± se) from baskets with different mesh sizes across sites
and seasons. Asterisks represent significant differences between mesh sizes within site and
season: * P-value<0.05, **P-value<0.01, ***P-value<0.001. NS – Nahal Shita; MS – Meishar;
AV – Avdat; SS – Sayeret Shaked; HS – Havat Shikmim; BG – Bet Guvrin; RH – Ramat
Hanadiv. Negative values may represent cases in which physical cleaning and ash correction
failed to correct for all dust accumulation on the litter or cases in which exogenous litter may
have penetrated the baskets. Our findings were not sensitive to these negative values.

Figure 4 - Contribution of different organism size classes to litter removal (mean ± se) across the precipitation gradient during summer, winter and both seasons combined. Macrofaunal contribution was calculated as the within-block difference between macro- and meso-baskets; mesofaunal contribution as the difference between meso- and micro-baskets; microbial and whole community contributions represent litter removal rates in the micro- and macro-baskets, respectively. Curves were fitted to data using local estimation scatterplot smoothing (LOESS). Negative values may represent cases in which physical cleaning and ash correction failed to correct for all dust accumulation on the litter or cases in which exogenous litter may have penetrated the baskets. Our findings were not sensitive to these negative values.

Figure 5 - Macro-decomposer abundance, biomass and alpha morphospecies richness across the
precipitation gradient in the two experimental seasons (mean ± se). Values are averaged across
traps and divided by the number of trapping days. NS – Nahal Shita; MS – Meishar; AV –
Avdat; SS – Sayeret Shaked; HS – Havat Shikmim; BG – Bet Guvrin; RH – Ramat Hanadiv.

Figure 6 - (A) and (B) Graphical representation of the first two axes of a principal coordinate analysis on the macro-decomposer assemblage data. Colors represent experimental sites in A and seasons in B. Arrows represent taxonomic group scores fitted onto the PCoA ordination. (C) Distribution of abundance, biomass and morphospecies richness among macro-decomposer taxonomic groups in each site across the aridity gradient. NS – Nahal Shita; MS – Meishar; AV – Avdat; SS – Sayeret Shaked; HS – Havat Shikmim; BG – Bet Guvrin; RH – Ramat Hanadiv. Color codes (left to right in panel C): grey – Zygentoma, burgundy – termites, olive green – millipedes, turquoise – woodlice, pink – snails and slugs, purple – earthworms, dark green – crickets, pale yellow – bristletails, red – beetles, pale blue – ants.

Discussion

Our goal was to investigate how climate interacts with soil biota of different size categories to influence litter decomposition. We used litter baskets of different mesh sizes that were placed along a sharp precipitation gradient during hot-dry summer and again during colder-wetter winter. Our results suggest that decomposers respond differently to precipitation levels based on their size, leading to similar overall decomposition rates across the gradient, except in hyper-arid sites. We found that microbial decomposition was minimal during the summer. In the winter, microbial decomposition was positively associated with precipitation, governing the whole-community decomposition. In both seasons, mesofaunal decomposition was moderate and followed a hump-shaped response to precipitation, peaking in semiarid sites. Macro-decomposition contributed minimally to whole-community decomposition during the winter, but during the summer dominated decomposition in the two arid sites. Using pitfall trapping, we found that macro-decomposer richness, abundance and biomass followed a hump-shaped response to precipitation, peaking in arid environments.

The puzzle of why plant litter decomposition in arid-lands is decoupled from annual precipitation and is faster than expected based on microbial decomposition models has bothered scientists for half a century[7,34,35], and was later termed the desert decomposition conundrum[36]. Attempts to resolve this conundrum have focused predominantly on abiotic weathering agents, such as photodegradation[37,38] and thermal degradation[39], alternative sources of moisture such as fog, dew and atmospheric water vapor[40] and soil–litter mixing[36,41]. We, in turn, hypothesized that the opposing climatic dependencies of macrofauna and that of microorganisms and mesofauna should lead to similar overall decomposition rates across precipitation gradients, except in hyperarid environments in which decomposers activity is predicted to be minimal regardless of

organism size. Our results largely agree with this hypothesis. Whole-community decomposition

was minimal in hyper-arid sites in both summer and winter. In the winter, microbial

decomposition dominated the whole-community decomposition, demonstrating a positive

response to precipitation that reached a maximum in the most mesic Mediterranean site. In

contrast, macro-decomposition has contributed only little to the whole-community

decomposition during the winter, but dominated the arid sites' decomposition in the summer.

These findings supported the long-suggested but largely overlooked hypothesis that macro-

decomposition governs plant litter decomposition in deserts[7,34,35]. The opposing climatic

dependencies of micro- and macro-decomposers have led to similar or even higher annual

decomposition rates in arid sites compared to those measured in more mesic sites. Consequently,

we highlighted that differential climatic dependencies of different-sized decomposers rather than

abiotic factors explain the discrepancy between classic decomposition models and the observed

decomposition rates in drylands. This realization provided a plausible resolution to the

longstanding desert decomposition conundrum, and exposed a hidden mechanism that may

account for unexplained variation in plant litter decomposition across biomes.

Canonically, faunal decomposition is expected to be positively associated with

temperature and moisture[20]. We, however, hypothesized that climate dependencies of mesofauna

and macrofauna should differ due to the lower sensitivity of macrofauna to high temperature and

low moisture, and the ability of macro-decomposers to shuttle between the hostile environment

aboveground and the climatic havens belowground[28]. We also hypothesized that low and

unpredictable resource availability in hyper-arid environments should limit macro-decomposer

populations. Consequently, we predicted that macro-decomposers should be more prevalent in

arid environments in comparison to hyper-arid or more mesic environments. Furthermore, ample

resource availability may increase niche space[42], resulting in higher macro-decomposer diversity,

which in turn can facilitate decomposition through synergistic effects of functionally

complementary species[43–46]. Thus, we predicted that macro-decomposition should reflect the

variation in the abundance, richness and biomass of macro-decomposers across the precipitation

gradient. Our findings supported these predictions. The richness, abundance and biomass of

macro-decomposers followed a hump-shaped relationship with precipitation, peaking in arid

environments and diminishing toward hyper-arid or semiarid and Mediterranean sites. Macro-

decomposer assemblages were dominated by ants and beetles across the aridity gradient except

in the Mediterranean site that was dominated by isopods and millipedes. During the summer, the

observed hump-shaped relationship between macro-decomposition and precipitation tightly

echoed the variation in richness, abundance, and biomass of macro-decomposers, revealing the

mechanistic foundation for the cross-system variation in macro-decomposition.

In winter, macro-decomposer abundance, richness, and biomass were similar to or even

higher than those measured during the summer across all sites. Despite this resemblance, macro-

decomposition did not reflect the observed variation in macro-decomposer assemblages. This

discrepancy could be explained by between-seasons differences in the structure of the macro-

decomposer assemblages (**Fig. 6B**). The macro-decomposer summer assemblage in Sayeret

Shaked was more similar to the Avdat summer assemblage than to the Sayeret Shaked winter

assemblage. The Avdat assemblages were more similar to each other across seasons than the

Sayeret Shaked assemblages. This may explain why macro-decomposition in winter was higher

in Avdat than in Sayeret Shaked. Termites (*Hodotermitidae* sp.), that play an important role in

decomposition, were more abundant in summer compared to the winter in Sayeret Shaked but

not in Avdat. Moreover, our data revealed that several beetle taxa (*Adelostoma* sp., *Akis reflexa*

(Fabricius, 1775), *Dailognatha crenata* (Reiche & Saulcy, 1857), *Tentyrina* sp., *Zophosis* sp.)

were prevalent in both arid sites during the summer but were absent or very scarce in the winter.

Phenological differences in the behavior of dominant macro-decomposers may also contribute to

the seasonal differences[47]. For instance, *Hemilepistus reaumuri* (Milne Edwards, 1840), an

abundant isopod species in Avdat, and *Anacanthotermes ubachi* (Navás, 1911), a common

termite species in Sayeret Shaked, consume detritus predominantly during the summer and

autumn and disperse and reproduce during the winter[48,49]. Future studies should explicitly test

these explanations. Regardless, the whole-year association between macro-decomposition and

the abundance, richness, and biomass of macro-decomposers strongly support our hypothesis.

Theory suggests that plant litter decomposition by meso-decomposers should increase

with moisture. This pattern was supported by a cross-biome experiment[50]. Thus, we

hypothesized that both meso-decomposition and microbial decomposition should increase with

precipitation and be more prominent in the winter than in the summer. Our results did not

coincide with these hypotheses. Litter decomposition by mesofauna followed a unimodal pattern

across the precipitation gradient, peaking under semiarid conditions in both seasons (**Fig. 4**).

Meso-decomposition and macro-decomposition were similar in the hyper-arid and

Mediterranean sites. However, meso-decomposition was higher than macro-decomposition in the

semiarid sites and much lower than macro-decomposition in the arid sites. These results suggest

that meso-decomposers, like macro-decomposers, have adaptations that allow them to strive in

moisture-deprived environments. Yet, meso-decomposition peaked in more mesic conditions

than macro-decomposition, implying higher moisture dependency. Our findings highlight the

need to explore the moisture dependency of mesofauna in greater details, and to generate

different predictions for how mesofauna and microorganisms may affect plant litter

decomposition.

Faunal decomposition in our study peaked in arid environments, contrasting the positive

association between faunal decomposition and precipitation that was found in recent global

meta-analyses[20,21]. This discrepancy may reflect underestimation of faunal decomposition rates

in drylands, possibly because these studies either deliberately grouped cold and dry

environments together[20,50], or focused solely on precipitation without accounting for differences

in temperature[21]. In cold water-deprived environments and seasons, low temperatures may limit

the populations and activity of ectotherm animals[4,22]. Therefore, ignoring the effect of

temperature may lead to falsely smaller faunal effects on decomposition in drylands. This bias

may contribute to the positive association between precipitation and faunal decomposition. To

reveal the realistic relationships, future studies on faunal decomposition should explore the

effects of temperature, precipitation and the interaction between them. It is important noting that

temperature could affect decomposition both directly by determining the activity of ectotherms

and indirectly by regulating moisture availability. Thus, using aridity indices that aim to correct

for moisture availability cannot resolve the need to account also for temperature *per se*.

We used litter of one plant species that is common to all seven experimental sites. In this

way, we were able to compare decomposition rates across sites without using litter that is

exogenous to one or more sites. Decomposer communities are assumed to perceive litter quality

based on the chemical composition of the focal litter, and that of past resource inputs that these

communities have experienced[51]. Plant communities differ substantially between our sites,

suggesting that each decomposer community has experienced very different resource inputs, and

may have access to alternative litter of various qualities. Thus, it is likely that the perceived

quality of *S. capensis* litter differs between sites, possibly confounding our results. Moreover,

our sites were distributed along a single precipitation gradient. Thus, our results may be

confounded by other environmental factors that were not accounted for. To determine how

general our findings are, future research should repeat our experimental approach using different

litter sources, and across multiple climatic gradients.

In conclusion, our work revealed that decomposers of varying size categories have

different moisture dependencies. This suggests that microorganisms, meso-decomposers and

macro-decomposers should be considered separately in decomposition models, and emphasizes

the need to contemplate animals' physiology and behavior when investigating zoogeochemical

processes. Warm drylands cover 19% of the land surface worldwide and expand rapidly due to

unsustainable land-use and climate change[52]. We highlight the importance of macro-

decomposition in arid-lands that compensates for the minimal microbial decomposition,

providing a plausible resolution to the long-debated dryland decomposition conundrum.

Understanding the mechanisms that regulate decomposition in drylands is key for conserving and

restoring fundamental ecosystem processes in these ever-growing areas, and in improving our

understanding of global processes like C cycling. To date, the general conceptualization of

decomposition is largely based on ample research from temperate ecosystems. Thus, prevailing

theory centers on focal processes that dominate decomposition in these systems. Our work

highlights that in other less studied ecosystems additional processes like the role of animal

decomposers may be dominant, opening the door for new exciting research that may shake our

conceptualization of decomposition processes.

References

1. Schlesinger, W. H. & Bernhardt, E. Biogeochemistry: An Analysis of Global Change. (Academic Press, 2020) doi:10.1016/B978-0-12-814608-8.09991-6.

2. Bradford, M. A., Berg, B., Maynard, D. S., Wieder, W. R. & Wood, S. A. Understanding the dominant controls on litter decomposition. *J. Ecol.* **104**, 229–238 (2016).

3. Joly, F. X., Scherer-Lorenzen, M. & Hättenschwiler, S. Resolving the intricate role of climate in litter decomposition. *Nat. Ecol. Evol.* **7**, 214–223 (2023).

4. Swift, M. J., Heal, O. W. & Anderson, J. M. *Decomposition in Terrestrial Ecosystems*. (University of California Press, 1979).

5. Wu, X., Niklas, K. J. & Sun, S. Climate change affects detritus decomposition rates by modifying arthropod performance and species interactions. *Curr. Opin. Insect Sci.* **47**, 62–66 (2021).

6. Berg, B. *et al.* Litter mass loss rates in pine forests of Europe and Eastern United States: some relationships with climate and litter quality. *Biogeochemistry* **20**, 127–159 (1993).

7. Meentemeyer, V. Macroclimate and lignin controls of litter decomposition rates. *Ecology* **59**, 465–472 (1978).

8. Bradford, M. A. *et al.* A test of the hierarchical model of litter decomposition. *Nat. Ecol. Evol.* **1**, 1836–1845 (2017).

9. Parton, W. *et al.* Global-scale similarities in nitrogen release patterns during long-term decomposition. *Science* **315**, 361–364 (2007).

10. Aerts, R. Climate, leaf litter chemistry and leaf litter decomposition in terrestrial ecosystems: A triangular relationship. *Oikos* **79**, 439 (1997).

11. Zhang, D., Hui, D., Luo, Y. & Zhou, G. Rates of litter decomposition in terrestrial ecosystems: Global patterns and controlling factors. *J. Plant Ecol.* **1**, 85–93 (2008).

12. David, J. F. The role of litter-feeding macroarthropods in decomposition processes: A reappraisal of common views. *Soil Biol. Biochem.* **76**, 109–118 (2014).

13. Frouz, J. Effects of soil macro- and mesofauna on litter decomposition and soil organic matter stabilization. *Geoderma* **332**, 161–172 (2018).

14. Griffiths, H. M., Ashton, L. A., Parr, C. L. & Eggleton, P. The impact of invertebrate decomposers on plants and soil. *New Phytol.* **231**, 2142–2149 (2021).

15. Auclerc, A. *et al.* Fostering the use of soil invertebrate traits to restore ecosystem functioning. *Geoderma* **424**, 116019 (2022).

16. Sagi, N., Zaguri, M. & Hawlena, D. Macro-detritivores assist resolving the Dryland Decomposition Conundrum by engineering an underworld heaven for decomposers. *Ecosystems* **24**, 56–67 (2021).

17. Coulis, M., Hättenschwiler, S., Coq, S. & David, J. F. Leaf litter consumption by macroarthropods and burial of their faeces enhance decomposition in a Mediterranean ecosystem. *Ecosystems* **19**, 1104–1115 (2016).

458   18.   Joly, F. X., Coq, S., Coulis, M., Nahmani, J. & Hättenschwiler, S. Litter conversion into detritivore faeces

459   reshuffles the quality control over C and N dynamics during decomposition. *Funct. Ecol.* **32**, 2605–2614

460   (2018).

461   19.   Joly, F. X. *et al.* Detritivore conversion of litter into faeces accelerates organic matter turnover. *Commun.*

462   *Biol.* **3**, 660 (2020).

463   20.   García-Palacios, P., Maestre, F. T., Kattge, J. & Wall, D. H. Climate and litter quality differently modulate

464   the effects of soil fauna on litter decomposition across biomes. *Ecol. Lett.* **16**, 1045–1053 (2013).

465   21.   Xu, X. *et al.* Cellulose dominantly affects soil fauna in the decomposition of forest litter: A meta-analysis.

466   *Geoderma* **378**, 114620 (2020).

467   22.   Johnston, A. S. A. & Sibly, R. M. Multiple environmental controls explain global patterns in soil animal

468   communities. *Oecologia* **192**, 1047–1056 (2020).

469   23.   Cloudsley-Thompson, J. L. Adaptations of arthropoda to arid environments. *Annu. Rev. Entomol.* **20**, 261–

470   283 (1975).

471   24.   Veldhuis, M. P., Laso, F. J., Olff, H. & Berg, M. P. Termites promote resistance of decomposition to

472   spatiotemporal variability in rainfall. *Ecology* **98**, 467-477 (2017)

473   25.   Sagi, N., Grünzweig, J. M. & Hawlena, D. Burrowing detritivores regulate nutrient cycling in a desert

474   ecosystem. *Proc. R. Soc. B Biol. Sci.* **286**, 20191647 (2019).

475   26.   Zanne, A. E. *et al.* Termite sensitivity to temperature affects global wood decay rates. *Science* **377**, 1440–

476   1444 (2022).

477   27.   Ayal, Y., Polis, G. A., Lubin, Y. & Goldberg, D. How can high animal diversity be supported in low-

478   productivity deserts? The role of macrodetritivory and habitat physiognomy. in *Biodiversity in drylands:*

479   *towards a unified framework* (eds. Shachack, M., Gosz, J. R., Perevolotsky, A. & Pickett, S. T. A.) 34–45

480   (Oxford University Press, 2005).

481   28.   Sagi, N. & Hawlena, D. Arthropods as the engine of nutrient cycling in arid ecosystems. *Insects* **12**, 726

482   (2021).

483   29.   Barney, J. N. *et al.* Global Invader Impact Network (GIIN): toward standardized evaluation of the ecological

484   impacts of invasive plants. *Ecol. Evol.* **5**, 2878–2889 (2015).

485   30.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4. *J. Stat.*

486   *Softw.* **67**, 1-48 (2015).

487   31.   Scheipl, F., Greven, S. & Kuechenhoff, H., Size and power of tests for a zero random effect variance or

488   polynomial regression in additive and linear mixed models. *Comput. Stat. Data Anal.* **52**, 3283-3299 (2008).

489   32.   R Core Team. R: A language and environment for statistical computing. (2022).

490   33.   Oksanen, J. *et al.* vegan: Community Ecology Package. (2022).

491   34.   Noy-Meir, I. Desert ecosystems : Higher trophic levels. *Annu. Rev. Ecol. Syst.* **5**, 195–214 (1974).

492   35.   Whitford, W. G. Exceptions to the AET model: Deserts and clear-cut forest. *Ecology* **62**, 275–277 (1981).

493   36.   Throop, H. L. & Archer, S. R. Resolving the Dryland Decomposition Conundrum: Some new perspectives

494   on potential drivers. in *Progress in Botany* (eds. Lüttge, U., Beyschlag, W., Büdel, B. & Francis, D.) **70**,

495   171–194 (Springer-Verlag Berlin Heidelberg, 2009).

496 37. Austin, A. T. & Vivanco, L. Plant litter decomposition in a semi-arid ecosystem controlled by
497   photodegradation. *Nature* **442**, 555–558 (2006).

498 38. Austin, A. T. Has water limited our imagination for aridland biogeochemistry. *Trends Ecol. Evol.* **26**, 229–
499   235 (2011).

500 39. Day, T. A., Bliss, M. S., Placek, S. K., Tomes, A. R. & Guénon, R. Thermal abiotic emission of CO_2 and
501   CH_4 from leaf litter and its significance in a photodegradation assessment. *Ecosphere* **10**, (2019).

502 40. Evans, S., Todd-Brown, K. E. O., Jacobson, K. & Jacobson, P. Non-rainfall moisture: A key driver of
503   microbial respiration from standing litter in arid, semiarid, and mesic grasslands. *Ecosystems* **23**, 1154–1169
504   (2020).

505 41. Throop, H. L. & Belnap, J. Connectivity dynamics in dryland litter cycles: Moving decomposition beyond
506   spatial stasis. *Bioscience* **69**, 602–614 (2019).

507 42. Macarthur, R. H. Patterns of species diversity. *Biol. Rev.* **40**, 510–533 (1965).

508 43. Heemsbergen, D. A. *et al.* Biodiversity effects on soil processes explained by interspecific functional
509   dissimilarity. *Science* **306**, 1019–1020 (2004).

510 44. Gessner, M. O. *et al.* Diversity meets decomposition. *Trends Ecol. Evol.* **25**, 372–380 (2010).

511 45. Boyero, L. *et al.* Impacts of detritivore diversity loss on instream decomposition are greatest in the tropics.
512   *Nat. Commun.* **12**, 1–11 (2021).

513 46. Zeng, X. *et al.* Soil invertebrates are the key drivers of litter decomposition in tropical forests. *Authorea*
514   *Prepr.* (2023) doi:10.22541/AU.167776750.08748069/V1.

515 47. Bonato Asato, A. E., Wirth, C., Eisenhauer, N. & Hines, J. On the phenology of soil organisms: Current
516   knowledge and future steps. *Ecol. Evol.* **13**, e10022 (2023).

517 48. Zaady, E., Groffman, P. M., Shachak, M. & Wilby, A. Consumption and release of nitrogen by the harvester
518   termite *Anacanthotermes ubachi navas* in the northern Negev desert, Israel. *Soil Biol. Biochem.* **35**, 1299–
519   1303 (2003).

520 49. Shachak, M., Chapman, E. A. & Steinberger, Y. Feeding, energy flow and soil turnover in the desert isopod,
521   Hemilepistus reaumuri. *Oecologia* **24**, 57–69 (1976).

522 50. Wall, D. H. *et al.* Global decomposition experiment shows soil animal impacts on decomposition are
523   climate-dependent. *Glob. Chang. Biol.* **14**, 2661–2677 (2008).

524 51. Strickland, M. S., Osburn, E., Lauber, C., Fierer, N. & Bradford, M. A. Litter quality is in the eye of the
525   beholder: initial decomposition rates as a function of inoculum characteristics. *Func. Ecol.* **23**, 627-636
526   (2009).

527 52. Cartereau, M. *et al.* Global bioregionalization of warm drylands based on tree assemblages mined from
528   occurrence big data. *Front. Biogeogr.* **14**, e56435 (2022).

Acknowledgements

We thank Liat Hadar, Ronen Kadmon, Ronen Ron, Omri Sharon, Shayli Dor-Haim, Nitzan Segev and Rachel Armoza-Zvuloni for facilitating our work in the experimental sites; Igor Armiach Steinpress, Laibale Friedman and Armin Ionescu for help with morphospecies identification; and Ariel Malinsky-Buller, Netanel Paz, Omri Sherman, Dor Gabay, Coral Ben-Lulu, Ofer Frumkin, Aparna Lajmi, and Atar, Alon and Ela Hawlena for lab and field assistance.

Funding

The research was funded by the Israel Science Foundation (ISF-No. 1391/19) to D.H.

Competing interests

The authors declare no conflict of interest.