Amyloid β Induces Lipid Droplet-Mediated Microglial Dysfunction in Alzheimer's Disease

Priya Prakash[1#], Palak Manchanda[1#], Evi Paouri[2], Kanchan Bisht[1], Kaushik Sharma[1], Jitika Rajpoot[1], Victoria Wendt[1], Ahad Hossain[1], Prageeth R. Wijewardhane[1], Caitlin E. Randolph[1], Yihao Chen[1], Sarah Stanko[2], Nadia Gasmi[2], Anxhela Gjojdeshi[2], Sophie Card[2], Jonathan Fine[1], Krupal P. Jethava[1], Matthew G. Clark[1], Bin Dong[1], Seohee Ma[1], Alexis Crockett[2], Elizabeth A. Thayer[1], Marlo Nicolas[3], Ryann Davis[1], Dhruv Hardikar[1], Daniela Allende[3], Richard A. Prayson[3], Chi Zhang[1], Dimitrios Davalos[2,4*], Gaurav Chopra[1,5,6,7,8*]

[1]Department of Chemistry, Purdue University, West Lafayette, IN 47907, USA
[2]Department of Neurosciences, Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44106, USA
[3]Department of Anatomic Pathology, Cleveland Clinic, Cleveland, OH 44106, USA
[4]Department of Molecular Medicine, Cleveland Clinic Lerner College of Medicine of Case, Western Reserve University, Cleveland, OH 44106, USA
[5]Purdue Institute for Integrative Neuroscience, Purdue University, West Lafayette, IN 47907, USA
[6]Purdue Institute for Drug Discovery, Purdue University, West Lafayette, IN 47907, USA
[7]Purdue Center for Cancer Research, Purdue University, West Lafayette, IN 47907, USA
[8]Purdue Institute of Inflammation, Immunology and Infectious Disease, Purdue University, West Lafayette, IN 47907, USA

[#]These authors contributed equally to this work

[*]Correspondence:
Dimitrios Davalos, Ph.D.; davalod@ccf.org
Gaurav Chopra, Ph.D.; gchopra@purdue.edu

Summary

Several microglia-expressed genes have emerged as top risk variants for Alzheimer's disease (AD). Impaired microglial phagocytosis is one of the main proposed outcomes by which these AD-risk genes may contribute to neurodegeneration, but the mechanisms translating genetic association to cellular dysfunction remain unknown. Here we show that microglia form lipid droplets (LDs) upon exposure to amyloid-beta (Aβ), and that their LD load increases with proximity to amyloid plaques in brains from human patients and the AD mouse model 5xFAD. LD formation is dependent on age and disease progression and is prominent in the hippocampus in mice and humans. Despite differences in microglial LD load between brain regions and sexes in mice, LD-laden microglia exhibited a deficit in Aβ phagocytosis. Unbiased lipidomic analysis identified a decrease in free fatty acids (FFAs) and a parallel increase in triacylglycerols (TGs) as the key metabolic transition underlying LD formation. DGAT2, a key enzyme for converting FFAs to TGs, promotes microglial LD formation and is increased in 5xFAD and human AD brains. Inhibition or degradation of DGAT2 improved microglial uptake of Aβ and drastically reduced plaque load in 5xFAD mice, respectively. These findings identify a new lipid-mediated mechanism underlying microglial dysfunction that could become a novel therapeutic target for AD.

Keywords

Microglia, lipids, lipidomics, lipid droplets, Alzheimer's disease, metabolism, neurodegeneration, phagocytosis, glia

Introduction

Alzheimer's disease (AD) is a progressive neurodegenerative disorder of the aging human population. Accumulation of amyloid-beta (Aβ) is a defining histological hallmark of the AD brain[1]. However, failures in Aβ-targeted clinical trials[2], combined with findings showing no apparent correlation between cognitive decline and overall plaque load in AD patients[3], suggest that additional mechanisms may be crucially involved in AD etiopathogenesis. Genome-wide association studies (GWAS) identified many AD risk genes related to the immune response and microglia, including the phagocytic receptors CD33 and TREM2[4,5,6,7,8]. Single-cell RNA sequencing identified gene signatures characteristic of prominent protective and dysfunctional microglial subpopulations in AD[9,10]. Moreover, a study that focused on human brain disease-associated variants of non-coding regulatory regions in a cell-specific manner identified multiple sporadic AD risk variants specifically in microglial transcriptional enhancers[11], further highlighting the involvement of microglia in AD pathogenesis.

In the AD brain, reactive microglia cluster around Aβ plaques[12] and form a physical barrier believed to restrict plaque propagation[13,14]. During early stages of AD, microglial reactivity is considered as beneficial for Aβ clearance[15]; however, sustained inflammation likely contributes to neurodegeneration[12]. Increased pro-inflammatory gene expression in response to accumulating Aβ in older AD mice leads to decreased microglial Aβ clearance receptors or Aβ-degrading enzymes, thereby promoting further Aβ accumulation[16]. Furthermore, the ability of microglia to remove Aβ declines over time, supporting that the progression of amyloid pathology correlates with impaired capacity of microglia to phagocytose Aβ[17,16].

In addition to genes and proteins, changes in microglial lipid content can also affect their state and function[18,19,20,21]. Cellular lipids regulate functions like migration and phagocytosis[22], are important for immune cell modulation and signaling[23], and their dysregulation has been linked to neurodegenerative disorders, including AD[24]. Top AD-risk genes such as TREM2, APOE, and INPP5D are directly related to lipid metabolism[25,26,27]. Although lipid droplets (LDs, first described as fat particles by Alois Alzheimer[28]) were initially considered to be passive fat deposits[29], they are dynamic intracellular organelles (diameter <1–100 μm) that regulate lipid metabolism. LDs consist of a phospholipid monolayer containing a core of neutral lipids like triacylglycerols (TAGs) and cholesteryl esters (CEs)[30]. Inflammatory triggers like lipopolysaccharide (LPS)[31,19], fatty acids[32], and aging[20] can result in LD accumulation in primary microglial cells and cell lines. Further, human iPSC-derived microglia were enriched in LDs in a chimeric mouse model of AD[33]. However, it is not known if Aβ can directly induce LD formation in microglia, and if/how changes to their lipid or metabolite composition can affect microglial function in AD.

Here we show that plaque-associated microglia closely associated with Aβ plaques contained more LDs and had larger cell bodies and shorter processes in both mouse and human brains, highlighting a unique LD-laden microglial subtype in AD. Extensive lipidomic and metabolomic profiling in microglia revealed specific types of lipids and metabolic pathways likely responsible for their LD-laden phenotype in the 5xFAD mouse model, which was more pronounced in microglia from females compared to males. Functionally, LD-laden microglia showed reduced phagocytosis of

Aβ compared to age-matched WT microglia. Mechanistically, we found that Aβ treatment alone
promotes a drastic shift in lipid content in microglia isolated from WT brains, even within only 24
hours of Aβ exposure. Extensive lipidomic characterization of these microglia revealed a gradual
increase in TAG content following Aβ treatment, which was similar to the lipid composition
changes we detected in microglia from 5xFAD mice. Based upon this finding, we identified
diacylglycerol O-acyltransferase 2 (DGAT2), a key enzyme for the conversion of FFAs to TAGs,
as also being the key catalyst for Aβ-induced LD formation in microglia. DGAT2 protein
expression was increased in both mouse and human AD brains, and inhibiting DGAT2 increased
the phagocytosis of Aβ by 5xFAD microglia. Importantly, degradation of the DGAT2 enzyme in
the brain of 5xFAD mice drastically reduced amyloid plaque load. Our study thus highlights
DGAT2 as a promising target for preventing or reversing phagocytic dysfunction in LD-laden
microglia in the AD brain.
131
132
133 **Microglia accumulate lipid droplets in an age-, sex-, and region-dependent manner in the**
134 **5xFAD model of Alzheimer's disease**
135
Prior literature has linked the accumulation of LDs in microglia with inflammation, aging, and
impaired phagocytosis, all of which are also hallmarks of AD[31,20,33]. We asked if/how exposure to
Aβ could induce LD formation in microglia, and/or alter their lipid or metabolite composition in a
way that affects their function in AD. We used a mouse model with five familial AD mutations
(5xFAD)[34], which progressively accumulates extensive Aβ plaques (especially in the subiculum
and deep cortical layers) starting around 2-3 months of age. 5xFAD mice also develop prominent
gliosis, inflammation, neuronal loss, behavioral impairments, and sex-distinct systemic metabolic
changes[35,34,36]. We isolated microglia from 5xFAD and WT mice and performed flow cytometry
analysis after staining for their neutral lipids (**Fig. 1a; Fig. S1a**). Although there was no difference
in younger mice (**Fig. 1b**), microglia from 5-7-month-old female 5xFAD mice showed
significantly higher LD content (1.58-fold) compared to cells from age-matched controls. Among
LD$^+$ cells, almost all 5xFAD microglia had high LD content, whereas WT microglia had lower LD
content (**Fig. 1c**). Microglia from 5-7-month-old male 5xFAD mice also showed a significant but
modest increase (1.15-fold) in LDs compared to WT (**Fig. 1d**), however a higher number of LD$^+$
microglia had high LD content in cells from 5xFAD compared to WT male mice (**Fig. 1e**). These
findings demonstrate that microglia show increased LD load, in a mouse model of AD with
extensive Aβ brain deposition, albeit with some age- and sex-dependent variability.
153
Next, we investigated if acutely-isolated microglia from 5xFAD mice showed alterations in their
global lipid and metabolite profiles as they experienced a drastic increase in Aβ plaque load with
age. We performed unbiased mass spectrometry lipidomic and metabolomic profiling using a
modified multiple reaction monitoring (MRM)[37] approach that allowed us to screen 1380 lipid
species, categorized into 10 main classes (**Fig. S1b**), and over 700 metabolites. Such broad
coverage and depth of profiling allowed us to explore possible differences in the lipid profiles of
microglia from 5-7-month-old male and female 5xFAD mice and compare them to those from age-
and sex-matched WT microglia in an unbiased manner (**Supplementary Tables ST1-ST2**).
Female 5xFAD microglia showed clearly distinct profiles, (indicated by their clear separation in
the principal component analysis (PCA) space, accounting for over 95% of the variation in the

data) (**Fig. 1f**)**,** with 105 differentially-regulated lipids that were primarily downregulated,
compared to cells from WT controls (**Fig. S1c-e**).
166
We further evaluated triacylglycerol (TAG) profiles in female microglia and found an increase in
several long-chain species, including TAG(48:0), TAG(50:2), TAG(50:0), TAG(52:2),
TAG(52:0) in 5xFAD compared to WT cells (**Fig. 1g, Supplementary Table ST3**). In contrast,
the male lipidome showed less variability (poor separation in the PCA space) (**Fig. S1f**), in
agreement with the more limited variation in LD content in male microglia when analyzed at the
single-cell level (**Fig. 1d**). We also compared microglial metabolite profiles from the same tissues
and found them to be distinctly different (**Fig. S2a**). Metabolites like fructose 6-phosphate and
lactose were downregulated in 5xFAD male and female microglia compared to cells from WT
controls, affecting several key metabolic pathways such as the citrate TCA cycle, glutamine and
glutamate metabolism, arginine biosynthesis, etc. (**Fig S2b-c, Supplementary Tables ST4, ST5**).
These results illustrate that following prolonged Aβ exposure, microglia exhibit: i) a unique lipid
signature that likely facilitates the formation and accumulation of LDs within them in a sex-distinct
manner, and ii) dysregulation in several key metabolic pathways, that could also impact their
functions in the context of AD.
181
182   **Microglia have increased LD content in the hippocampus of 5xFAD mice and AD patients**
183
We next asked if/how the increased LD content and related metabolic changes related to brain
amyloid deposition, one of the prominent pathological hallmarks of 5xFAD mice and AD brains.
We examined the spatial distribution of LDs by performing label-free stimulated Raman scattering
(SRS) microscopy[38] in hippocampal brain slices of 5xFAD mice. We found a significantly higher
number of LDs, and a higher percentage of LD area overall, in chronic 5xFAD compared to age-
matched WT hippocampi (**Fig. 1h**). Interestingly, some LDs appeared in close proximity to Aβ
plaques in the 5xFAD tissue, as identified by their respective spectral signatures (**Fig. S3**).
191
To further profile the spatial distribution of LD-laden microglia relative to amyloid plaques, we
combined histological staining with methoxy O4 (amyloid plaques) and LipidTox (lipids) with
immunohistochemistry for IBA1 (microglia) in 5-7-month-old female 5xFAD brain sections. Even
though aged WT brains had a high number of LDs in microglia (as also previously reported[20]), and
in other cells (**Fig. 1i; S4a**), 5xFAD brains had significantly higher overall LD density and total
LD area in cortex and parts of the hippocampus (CA1 and subiculum) compared to WT (**Fig. S4b-
c**). The mean cell body area of LD$^+$ microglia was not significantly different than that of LD$^-$ cells
(**Fig. S4d**), nor was the mean cell body area of LD$^+$ microglia in 5-7-month-old female 5xFAD
mice compared to LD$^+$ cells in age and sex-matched WT controls (**Fig. S4e**). Similarly, the overall
proportion of LD$^+$ microglia was similar across cortex, CA1, and subiculum (**Fig. S4f**), but the
proportion of LD area within microglia was significantly increased in the subiculum of 5xFAD
mice (**Fig. 1j**), indicating a preferential increase in LD load within microglia in this brain region.
204
Since the hippocampus is significantly affected by amyloid pathology in AD patients[39], we also
stained postmortem human hippocampal sections from non-symptomatic (NS) and AD patients
for Aβ plaques, LDs, and microglia (**Fig. 1k**). After 3D reconstruction and segmentation of each
stained structure imaged by high-resolution confocal microscopy, we performed volumetric co-

localization analysis using Imaris (**Fig. S5**). We measured significantly higher (5.7-fold increase)
overall LD density (**Fig. 1l; Fig. S6**), and—similar to the mouse model findings—a significantly
higher percentage of microglial volume occupied by LDs in the hippocampus of AD patients
compared to NS controls in both males and females (**Fig. 1m**).
213
214   **Chronic exposure of microglia to Aβ plaques promotes LD accumulation, alters microglial**
215   **morphology, and impairs their phagocytic ability**
216
We next examined the spatial relationship between plaque load and LD density and found them to
be positively correlated in the cortex, CA1, and subiculum (**Fig. 2a**). Interestingly, the vast
majority of plaque-proximal microglia were laden with LDs in all three brain regions (**Fig. 2b;**
Fig. S7a). Particularly in the subiculum, plaque-proximal microglia demonstrated an amoeboid
morphology, while plaque-distant cells typically had smaller cell bodies and longer processes (**Fig.**
2c-d; Fig. S7b, c). Importantly, this also translated in human microglia in the hippocampus of AD
patients (both male and female), where plaque-proximal microglia had a significantly higher
number of LDs than plaque-distant microglia, which decreased with distance from the nearest
plaque (**Fig. 2e, f; Fig. S7d; Supplementary Movie 1**). In addition, LD$^+$ microglia closer to
plaques (0-10μm) contained larger LDs, and their total LD load also decreased progressively with
distance from the nearest plaque (**Fig. 2g; Fig. S7e**). The methodological approach implemented
in Imaris to identify PLIN2$^+$IBA1$^+$ cells and their LD load as a function of distance from the nearest
plaque is schematically represented in **Fig S8**. Overall, these results suggest that a
morphologically-distinct plaque-associated microglial phenotype characterized by larger LD load
exists in both the human AD brain and the amyloid-rich 5xFAD animal model.
232
Plaque-associated microglia have been previously shown to exhibit morphological and molecular
differences compared to non-plaque-associated cells in AD[40,41,42]. We specifically asked if direct
exposure to Aβ leads to LD formation and changes to phagocytic function in microglia. Microglia
isolated from 5-7-month-old female 5xFAD and WT mice were acutely seeded (1 hour) and treated
with Aβ^{pH}—a pH-dependent fluorescent probe that emits green fluorescence in the acidic
lysosomes upon phagocytosis[43]; LDs were then stained with LipidTox, and all cells were analyzed
by flow cytometry (**Fig. 2h, Fig. S9a**). Microglia isolated from WT mice showed a significant
increase in LD content (4.5-fold) upon direct exposure to Aβ^{pH} (**Fig. 2i**). However, Aβ^{pH} treatment
did not cause a significant increase in LDs in microglia isolated from 5xFAD brains (**Fig. 2j**). This
could indicate that since 5xFAD mice progressively develop Aβ plaques starting at 2 months of
age, chronic microglial exposure to Aβ possibly alters their functional abilities and overall state
by the age of 5-7 months, when amyloid deposition is extensive throughout the brain.
245
Microglial phagocytosis of Aβ is a critical clearance mechanism, and alterations of microglial
phagocytic capacity have been reported in chronic inflammation and AD mouse models[15,16]. We
therefore investigated if the increase in microglial LDs affects their phagocytic capacity. Live
microglia from 5xFAD brains showed a significant (40%) reduction in Aβ^{pH} phagocytosis
compared to cells from WT brains (**Fig. S9b-c**). Specifically, out of all microglia, 63.55% and
47.92% were Aβ^{pH+} in WT and 5xFAD, respectively (**Fig. 2k**). Interestingly, LD$^+$ 5xFAD
microglia showed impaired phagocytosis compared to LD$^+$ WT cells (**Fig. 2l**). Surprisingly, WT
microglia showed an increase in LDs due to acute Aβ^{pH} but did not exhibit reduced phagocytic

capacity. In conclusion, these experiments demonstrate that LD-laden microglia that are
chronically exposed to Aβ exhibit a dysfunctional phagocytic phenotype.
256
257

Direct Aβ exposure is sufficient to significantly alter the microglial lipidome towards lipid
258
droplet formation, independently of inflammatory factors
259
260
We were surprised to discover that in microglia from 5-7-month-old WT brains, even acute direct
exposure to Aβ *ex vivo* was sufficient to induce large LD formation. Thus, we asked if direct Aβ
exposure is sufficient to promote LD formation and/or regulate broad metabolic changes in
microglia independently of other inflammatory factors. We isolated CD11b$^+$ primary microglia
from 5-7-month-old WT mice, cultured them *in vitro* for 7-10 days and treated them with Aβ
aggregates for 1, 12, and 24 hours before assaying both cellular and secretory lipids and
metabolites by mass spectrometry-based MRM-profiling (**Fig. 3a**). We again screened over 1370
lipid species categorized into 10 main classes including FFAs, ceramides, and TAGs
(**Supplementary Tables ST6, ST7**) as well as over 700 metabolites (**Supplementary Tables ST8,**
S9). We found considerable differences in the microglial lipidome between Aβ- and vehicle-
treated cells, at 1 hour and 24 hours of Aβ treatment, respectively, suggesting a distinct lipidomic
transition occurring within microglia following exposure to Aβ (**Fig. 3b**).
273
While we observed dramatic changes in several lipid classes, FFAs were the most differentially-
regulated lipid class with acute (1 hour) Aβ exposure (**Fig. 3c-d**), with very long-chain saturated
FFAs, C20:0, C22:0, and C19:0 being the most upregulated lipids at this time point (**Fig. 3e; S10a**).
We also used novel gas-phase ion/ion chemistry (**Supplementary Results**) to structurally validate
these highly-expressed saturated FFAs and confirmed that these specific saturated FFA structures
were indeed directly synthesized in microglia with acute (1 hour) Aβ exposure (**Fig. 3f, S11a-b**).
Interestingly, the cells did not maintain their saturated FFA repertoire with prolonged Aβ exposure
(24 hours) and transitioned to TAGs, which were the most differentially-regulated lipid class at
this time point, with TAG(52:3), TAG(54:3), and TAG(52:2) being the most upregulated species
(**Fig. 3c-e; S10b**). In addition, MRM-profiling confirmed the increase of specific TAGs in Aβ-
treated microglia at the 24-hour time point compared to vehicle-treated microglia (**Fig. 3e,**
Supplementary Table ST10). Furthermore, at 12 hours of Aβ treatment, in addition to FFAs,
cholesteryl esters (CEs) were the second most differentially-regulated lipid class, with very long
chain CEs 20:2, 24:1, and 16:3 being the top upregulated CEs in microglia (**Fig. S10c-e**). Neutral
lipids like TAGs and CEs form the core of LDs and are involved in energy storage and fatty acid
metabolism in cells. Importantly, we quantified the total amount of upregulated FFAs, CEs, and
TAGs as percent change of maximum ion intensity and verified that the reduction in FFAs from 1
hour to 24 hours was followed by a concomitant increase in TAGs and CEs at 24 hours (**Fig. 3g**).
This increase in core LD components suggested that the cells likely activated metabolic pathways
towards LD formation. Taken together with acute treatment of Aβ *ex vivo* (**Fig. 2b**), these results
suggest a direct effect of Aβ-aggregate exposure to promote LD formation in microglia. In contrast
to the cellular lipidome, we did not find many lipids differentially regulated in the microglial
conditioned media (secretory lipids) with Aβ treatment - none at 1 hour, five at 12 hours, and one
at 24 hours, respectively (**Supplementary Table ST7**). Given that both lipids and metabolites
work together to activate cellular metabolic pathways, we also evaluated changes in microglial

metabolite profiles with Aβ exposure. Metabolites corresponding to alanine, aspartate, and
glutamate metabolism, arginine biosynthesis, etc. were differentially regulated in microglia
exposed to Aβ (**Fig. S12, Supplementary Table ST11**). The metabolome of the microglial
conditioned media exhibited pronounced differences following 12 and 24 hours of Aβ exposure,
as seen in the PCA plots (**Fig. S13a**). Metabolites related to phenylalanine/tyrosine/tryptophan
biosynthesis and glycine/serine/threonine biosynthesis pathways were differentially regulated in
the microglial conditioned media due to Aβ (**Fig. S13b-c, Supplementary Table ST12**). Taken
together, the Aβ-treated cellular and secreted metabolites overlapped with metabolic pathways in
5xFAD compared to WT (**Fig. S2**). This suggests that Aβ plays a direct role in upregulating TAGs,
CEs and associated metabolites towards LD formation in AD. TAGs can be synthesized via two
major pathways involving the conversion of FFAs to TAGs via several intermediates: 1) the
glycerol phosphatase pathway or 2) the monoacylglycerol pathway (**Fig. 3h**)[44]. Since TAGs
constitute the major neutral lipid core of LDs and are integral to their structure and function, we
hypothesized that these pathways could be directly involved with the dramatic increase in TAGs
following Aβ exposure and likely also with LD formation in AD.

The DGAT2 pathway is required for Aβ-induced lipid droplet formation in AD microglia,
and inhibiting it rescues microglial phagocytic impairment and reduced plaque load in AD

The diacylglycerol O-acyltransferase (DGAT) enzymes catalyze the final rate-determining step in
the biosynthesis of TAGs. The DGAT2 enzyme is evolutionarily conserved across eukaryotes and
has the predominant and ancient function for mediating TAG synthesis from fatty acids[45]. In
addition, DGAT2 localizes around LDs, where it is required for LD-localized TAG
synthesis[46,47,48]. Thus, we investigated whether DGAT2 is involved in the formation of Aβ-induced
LDs in microglia (**Fig. 4a**). We immunostained 5xFAD and WT brain tissues for DGAT2, LDs,
Aβ plaques, and microglia. We found that LD-laden microglia surrounding the Aβ plaques
expressed DGAT2 in 5xFAD tissue (**Fig. 4b**), despite *Dgat2* mRNA being downregulated in
5xFAD microglia (**Fig. S14a**), we detected abundant DGAT2[+] microglia in the subiculum region
of the 5xFAD brain (**Fig. S14b, Fig. 4c**). We also immunostained human AD and NS hippocampal
tissue for Aβ plaques, LD, DGAT2, and microglia. We found that plaque-associated microglia
with increased LD content exhibited higher DGAT2 levels in AD brains compared to microglia
from NS brains (**Fig. 4d**).

Further, we used a DGAT2 inhibitor (D2i) to determine if reducing DGAT2 function affects
microglial LD content and Aβ-specific phagocytosis. Microglia from 5-7-month-old female
5xFAD and WT mice were acutely seeded (1 hour) and treated with Aβ[pH] and LipidTox
sequentially, in the presence of D2i or vehicle, after which cells were analyzed by flow cytometry
(**Fig. 5a**). Both WT and 5xFAD microglia showed a significant decrease in LDs upon D2i
treatment *in vitro* (approx. 51% and 57% decrease, respectively, **Fig. 5b**). Further, there was a
significant reduction in Aβ-induced LDs with D2i treatment in WT microglia, thereby confirming
that Aβ-induced LD formation requires the DGAT2 pathway (**Fig. 5c-d, Fig. S15a**). Interestingly,
we did not observe any significant differences in Aβ-induced LDs with D2i treatment in 5xFAD
microglia (**Fig. 5c-d, Fig. S15b**), similar to our results showing that Aβ[pH] treatment did not affect
LD load in microglia isolated from 5xFAD brains. Together, these results suggest an underlying
Aβ-induced LD saturation mechanism that allows DGAT2-mediated microglial LD formation in

5xFAD brain after chronic amyloid exposure *in vivo*, but prevents further LD accumulation upon
additional *in vitro* Aβ exposure.
346
Next, we examined if D2i affected microglial phagocytic capacity. LD$^+$ microglia from WT brains
showed a slight but non-significant increase in Aβ^{pH} uptake with D2i compared to vehicle-treated
cells (**Fig. 5e; Fig. S15c**). In contrast, LD$^+$ 5xFAD microglia showed a significant increase (1.41-
fold) in Aβ^{pH} uptake with D2i, which was similar to WT microglia in the presence of the inhibitor
(**Fig. 5e-f; Fig. S15f**). Furthermore, the overall phagocytic capacity of 5xFAD microglia was
improved with D2i treatment (**Fig. S15c**) compared to vehicle-treated cells, although again there
was no significant difference between WT and 5xFAD microglia in the presence of the inhibitor
(**Fig. S15d-e**). Thus, the phagocytic dysfunction of LD-laden 5xFAD microglia was attenuated in
the presence of D2i.
356
To investigate how targeting DGAT2 *in vivo* might affect amyloid pathology, we developed a
degrader of DGAT2[49] and continuously infused it into the lateral ventricles of 18-24 month old
5xFAD mice (**Fig. 5g**). Animals that received the DGAT2 degrader over a period of 1 week
showed a significant reduction in total LD area [36%] in the subicular hippocampal region
compared to the age-matched vehicle-treated control animals (**Fig. 5h,i, Fig. S16a**). Strikingly, the
DGAT2 degrader also drastically reduced the amyloid load in the 5xFAD mice by 63% compared
to mice that received the vehicle treatment (**Fig. 5i, Fig. S16b**). Taken together, these findings
indicate that degrading DGAT2 profoundly reduces LD content and improves amyloid pathology
even in chronically inflamed 18-24 month old 5xFAD mice. which typically presents itself with
an excessive amount of brain amyloid deposition and advanced disease.
367
368 In summary, we have discovered a novel Aβ-mediated mechanism that promotes LD formation in
369 microglia and identified the DGAT2 pathway as a novel target for therapeutic intervention to
370 reduce phagocytic impairment of microglia and promote amyloid clearance or limit amyloid
371 accumulation in the AD brain (**Fig. 5j**).
372
373 **DISCUSSION**
374
Over the past two decades, a plethora of studies have revealed important microglial contributions
to practically every neurological disorder or disease, with roles ranging from protective to
harmful[50]. This collective body of work, together with an increasing number of microglial
transcriptomic profiling studies, have underscored their multidimensional nature and their
impressive capacity to react to any signal presented to them in an age–, sex–, disease stage–, and
overall context–specific manner[50]. In this study, we showed that in the context of AD pathology,
chronic exposure to Aβ promotes specific metabolic changes in microglia that make them
progressively accumulate LDs and render them incapable of contributing to Aβ clearance. LD-
laden microglia increase in numbers in an age- (5-7-month old), sex- (female), and brain region-
specific manner (subiculum) in 5xFAD mice. These microglia are closely associated with Aβ
plaques and have a unique morphology comprised of larger cell bodies and shorter processes in
both 5xFAD and human AD brains. Importantly, LD-laden microglia exhibit reduced phagocytosis
of Aβ, a critical functional consequence of LD accumulation. Further, by challenging primary WT
microglia directly with Aβ, we discovered that even acute exposure to Aβ was sufficient to shift

their lipidomic composition towards reduced FFA and increased TAG content – a major
component of LDs. We thus identified DGAT2, a key enzyme for the conversion of FFAs to TAGs,
as an essential mediator of LD formation in microglia, and showed its increased abundance in
5xFAD as well as human AD brain tissue. Crucially, pharmacological inhibition of DGAT2
improved Aβ phagocytosis *ex vivo*, and DGAT2 degradation drastically reduced plaque load *in*
vivo, thus unraveling a new molecular target and mechanism related to microglial dysfunction in
AD.
396
Recent studies focused on transcriptomic changes occurring in microglia at the single-cell level
due to Aβ. While such studies demonstrate the extent of glial heterogeneity in a specific
environment based upon highly-abundant and actively-transcribed genes in each subpopulation,
they do not fully capture the functional aspects of these cell populations, which are directly
determined by bioactive molecules such as proteins, lipids, and other metabolites. We performed
the first-ever large-scale lipidomic and metabolomic characterization of microglia with acute
exposure to Aβ *in vitro* and in chronic 5xFAD mice, profiling over 1300 lipid species and 700
metabolites. Our data, which will be made available online (http://microgliaomics-
chopralab.appspot.com), showed an increase in long-chain saturated FFAs (specifically C20:0,
C19:0, and C22:0) upon 1h Aβ exposure; however, longer treatment with Aβ (24h) resulted in an
abundance of long-chain TAGs. Previous studies showed that saturated FFAs are extremely
cytotoxic to cells[51]. One way that cells overcome FFA-induced cytotoxicity is by metabolizing
them into TAGs[52], which are integral components of the LD core. Microglia are thus able to
circumvent lipid-mediated cytotoxicity via the production and accumulation of LDs. Interestingly,
we recently found that reactive astrocytes secrete long-chain saturated FFAs that kill injured
neurons and oligodendrocytes *in vitro* and *in vivo*[37]. While injured neurons and oligodendrocytes
typically have a low capacity for consuming FFAs and sequestering them into LDs[53], making them
more vulnerable to lipid-induced cytotoxicity in chronic inflammation[54,55], our findings
demonstrate that microglia can metabolize these toxic molecules into TAGs. This response could
be one way that microglia protect themselves and neurons from the cytotoxic environment of the
chronically-inflamed AD brain; however, it seems to affect their own morphology and functional
state as well.
419
Changes in microglial morphology have been described in numerous inflammatory, degenerative,
and injury paradigms as indications of changes in their overall state and reactivity. In our study,
LD-laden microglia were closely associated with Aβ plaques and exhibited shorter processes and
larger cell bodies in both 5xFAD mouse and human AD brains. These unique LD-laden microglia
were primarily located in the hippocampal subicular regions of 5xFAD mice and in postmortem
human hippocampi from AD patients, which are the sites of the earliest atrophy seen during AD
pathogenesis[56]. In addition to the protective roles of some lipids[57], their accumulation is also linked
to cellular senescence[58]. For example, in an obesity mouse model, senescent glia were found to
accumulate LDs, which contributed to impaired neurogenesis[59]. Whether these LD-laden
microglia are ultimately detrimental or protective to the tissue due to their dysfunctional, hyper-
reactive, and/or senescent state in the context of chronic inflammation remains elusive and may
well depend upon the extent or nature of metabolic changes that microglia experience in the
context of different challenges. Overall, these observations open new avenues for exploring the
functional roles of plaque-associated LD-laden microglia and their involvement in AD

progression, as well as the functional significance of LD accumulation in microglia in other
neuropathologies.
436
Phagocytosis is a major innate immune mechanism contributing to Aβ clearance, and its failure
favors Aβ accumulation in the brain. Late-onset AD, which accounts for ~95% of AD cases, is
associated with impaired Aβ clearance[60]. Microglia directly engulf and degrade Aβ[61,62], but their
capacity for efficient phagocytosis declines with aging, chronic exposure to Aβ, and during
sustained inflammation in the AD brain[63,16,17]. Indeed, acute exposure of primary microglia to Aβ
increased their inflammatory cytokine secretion and phagocytic performance, while chronic
exposure reduced phagocytosis and induced immune tolerance as well as a metabolic shift from
oxidative phosphorylation to glycolysis[64]. Interestingly, restoration of microglial phagocytosis
improved cognitive function in aged mice and promoted a homeostatic microglial transcriptional
signature[65].
447
Plaque–associated microglia have direct access to remove aggregated amyloid through
phagocytosis. Prior studies have proposed that these cells are compacting amyloid plaques through
phagocytic receptors such as TREM2[14], (a receptor expressed on myeloid cells, for which gene
variants have been associated to increased AD risk)[14,8], Axl, and Mer[66]. Genetic deletion of any of
these receptors in mouse models of AD resulted in less compact/more diffuse plaques, and was
associated with increased neuritic dystrophy around Aβ deposits[14,66]. These studies suggested that
microglia build a neuroprotective barrier that limits plaque growth through their phagocytic
receptors, and further proposed microglia as plaque "compactors" that uptake Aβ and condense it
in their lysosomes to eventually re-deposit it as dense core plaques[66]. We found that plaque–
associated microglia progressively accumulated more LDs, and that cells with increased LD
content were less capable of Aβ phagocytosis. LD-laden microglia found in the aged brain also
demonstrated phagocytic deficits[20], but the molecular mechanisms linking the appearance of these
organelles with functional deficits in microglia were unclear. We discovered that Aβ alone can
shift the metabolic equilibrium within microglia to convert FFAs to TAGs, one of the main
components of LDs. To decipher if this catalytic conversion that is known to contribute to LD
formation[52,67] is also responsible for microglial dysfunction, we blocked DGAT2, a key enzyme
for this pathway. Although we found the DGAT2 mRNA significantly decreased in 5xFAD
microglia, the protein was abundant in plaque-associated LD-laden microglia in both mouse and
human AD brains, underscoring how relying solely on transcript levels might mask important
ongoing cellular functions. Importantly, blocking DGAT2 was sufficient to restore phagocytosis
in 5xFAD microglia, suggesting that phagocytic dysfunction is the consequence, rather than the
cause, of chronic LD accumulation. In support of our findings, a study in macrophages showed
that their phagocytic function is also dependent upon the availability of FFAs that are generated
upon degradation of LDs, suggesting that LD accumulation somehow limits the phagocytic
capacity of all tissue-resident macrophages[68]. Furthermore, using a complementary DGAT2
protein degrader approach also reduced LD load *in vivo* in 18-24 month 5xFAD mice, underscoring
the importance of this molecular target for LD accumulation. Crucially, its very rapid (only 1 week
of treatment) and profound reduction (63%) of amyloid load in older animals highlights DGAT2
as a promising target to restore microglia-mediated Aβ clearance and decrease amyloid deposition
even in AD patients of advanced age and/or with increased amyloid pathology.
478

Delineating the molecular mechanisms that drive LD accumulation and their downstream impact on cellular functions and overall tissue health is paramount for the design of novel therapeutic strategies for AD and other neurodegenerative diseases. Here, we have identified one such druggable molecule that regulates TAG formation and LD accumulation in microglia while also directly impacting their function. Unlike other acyltransferases, such as GPAT and AGPAT, that are upstream and involved in the glycerol phosphatase pathway, the activity of DGAT is the direct rate-limiting step for the biosynthesis of TAGs[44]. Therefore, since disrupting DGAT2 alone was sufficient to improve microglial phagocytosis of Aβ, and reduce brain amyloid load, targeting DGAT2 emerges as a prime target for regulating the phagocytic activity of LD-laden microglia compared to other candidate enzymes. Our study thus provides the first proof-of-principle that disrupting DGAT2 can be a highly effective strategy for promoting the protective role of microglia in AD, and possibly in other neurodegenerative diseases with excessive protein aggregation and LD deposition.

MAIN REFERENCES

1. Serrano-Pozo, A., Frosch, M. P., Masliah, E. & Hyman, B. T. Neuropathological alterations in Alzheimer disease. *Cold Spring Harb. Perspect. Med.* **1**, (2011).
2. Plascencia-Villa, G. & Perry, G. Status and future directions of clinical trials in Alzheimer's disease. in *International Review of Neurobiology* vol. 154 3–50 (2020).
3. Morris, G. P., Clark, I. A. & Vissel, B. Inconsistencies and Controversies Surrounding the Amyloid Hypothesis of Alzheimer's Disease. *Acta Neuropathologica Communications* vol. 2 1–21 (2014).
4. Bertram, L. *et al.* Genome-wide Association Analysis Reveals Putative Alzheimer's Disease Susceptibility Loci in Addition to APOE. *Am. J. Hum. Genet.* **83**, 623–632 (2008).
5. Hollingworth, P. *et al.* Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat. Genet.* **43**, 429–436 (2011).
6. Naj, A. C. *et al.* Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nat. Genet.* **43**, 436–443 (2011).
7. Guerreiro, R. *et al.* TREM2 Variants in Alzheimer's Disease . *N. Engl. J. Med.* **368**, 117–127 (2013).
8. Jonsson, T. *et al.* Variant of TREM2 Associated with the Risk of Alzheimer's Disease . *N. Engl. J. Med.* **368**, 107–116 (2013).
9. Keren-Shaul, H. *et al.* A Unique Microglia Type Associated with Restricting Development of Alzheimer's Disease. *Cell* **169**, 1276-1290.e17 (2017).
10. Krasemann, S. *et al.* The TREM2-APOE Pathway Drives the Transcriptional Phenotype of Dysfunctional Microglia in Neurodegenerative Diseases. *Immunity* **47**, 566-581.e9 (2017).
11. Nott, A. *et al.* Brain cell type–specific enhancer–promoter interactome maps and disease-risk association. *Science (80-.).* **366**, 1134–1139 (2019).
12. Wyss-Coray, T. & Rogers, J. Inflammation in Alzheimer disease-A brief review of the basic science and clinical literature. *Cold Spring Harb. Perspect. Med.* **2**, a006346 (2012).
13. Condello, C., Yuan, P., Schain, A. & Grutzendler, J. Microglia constitute a barrier that

prevents neurotoxic protofibrillar Aβ42 hotspots around plaques. *Nat. Commun.* **6**, 1–14
(2015).
14. Yuan, P. *et al.* TREM2 Haplodeficiency in Mice and Humans Impairs the Microglia
Barrier Function Leading to Decreased Amyloid Compaction and Severe Axonal
Dystrophy. *Neuron* **90**, 724–739 (2016).
15. Heneka, M. T. *et al.* Neuroinflammation in Alzheimer's disease. *Lancet Neurol.* **14**, 388–
405 (2015).
16. Hickman, S. E., Allison, E. K. & El Khoury, J. Microglial dysfunction and defective β-
amyloid clearance pathways in aging alzheimer's disease mice. *J. Neurosci.* **28**, 8354–
8360 (2008).
17. Krabbe, G. *et al.* Functional Impairment of Microglia Coincides with Beta-Amyloid
Deposition in Mice with Alzheimer-Like Pathology. *PLoS One* **8**, (2013).
18. Leyrolle, Q., Layé, S. & Nadjar, A. Direct and indirect effects of lipids on microglia
function. *Neuroscience Letters* vol. 708 134–348 (2019).
19. Tremblay, M. E. *et al.* Remodeling of lipid bodies by docosahexaenoic acid in activated
microglial cells. *J. Neuroinflammation* **13**, 1–18 (2016).
20. Marschallinger, J. *et al.* Lipid-droplet-accumulating microglia represent a dysfunctional
and proinflammatory state in the aging brain. *Nat. Neurosci.* **23**, 194–208 (2020).
21. Chausse, B., Kakimoto, P. A. & Kann, O. Microglia and lipids: how metabolism controls
brain innate immunity. *Seminars in Cell and Developmental Biology* vol. 112 137–144
(2021).
22. Bourre, J. M. Brain lipids and ageing. in *Food for the Ageing Population: A volume in
Woodhead Publishing Series in Food Science, Technology and Nutrition* 219–251 (2008).
doi:10.1533/9781845695484.2.219.
23. Cantuti-Castelvetri, L. *et al.* Defective cholesterol clearance limits remyelination in the
aged central nervous system. *Science (80-.).* **359**, 684–688 (2018).
24. Wong, M. W. *et al.* Dysregulation of lipids in Alzheimer's disease and their role as
potential biomarkers. *Alzheimer's and Dementia* vol. 13 810–827 (2017).
25. Tsai, A. P. *et al.* INPP5D expression is associated with risk for Alzheimer's disease and
induced by plaque-associated microglia. *Neurobiol. Dis.* **153**, 105303 (2021).
26. Wang, Y. *et al.* TREM2 lipid sensing sustains the microglial response in an Alzheimer's
disease model. *Cell* **160**, 1061–1071 (2015).
27. Fitz, N. F. *et al.* Phospholipids of APOE lipoproteins activate microglia in an isoform-
specific manner in preclinical models of Alzheimer's disease. *Nat. Commun.* **12**, (2021).
28. Stelzmann, R. A., Norman Schnitzlein, H. & Reed Murtagh, F. An english translation of
alzheimer's 1907 paper, "über eine eigenartige erkankung der hirnrinde". *Clin. Anat.* **8**,
429–431 (1995).
29. Farese, R. V. & Walther, T. C. Lipid Droplets Finally Get a Little R-E-S-P-E-C-T. *Cell*
vol. 139 855–860 (2009).
30. Fujimoto, T. & Parton, R. G. Not Just Fat: The Structure and Function of the Lipid
Droplet. *Cold Spring Harb. Perspect. Biol.* **3**, 1–17 (2011).
31. Khatchadourian, A., Bourque, S. D., Richard, V. R., Titorenko, V. I. & Maysinger, D.
Dynamics and regulation of lipid droplet formation in lipopolysaccharide (LPS)-
stimulated microglia. *Biochim. Biophys. Acta* **1821**, 607–17 (2012).
32. Urso, C. J. & Zhou, H. Palmitic acid lipotoxicity in microglia cells is ameliorated by

unsaturated fatty acids. *Int. J. Mol. Sci.* **22**, 90–93 (2021).

33.   Claes, C. *et al.* Plaque-associated human microglia accumulate lipid droplets in a chimeric
model of Alzheimer's disease. *Mol. Neurodegener.* **16**, 50 (2021).

34.   Oakley, H. *et al.* Intraneuronal β-amyloid aggregates, neurodegeneration, and neuron loss
in transgenic mice with five familial Alzheimer's disease mutations: Potential factors in
amyloid plaque formation. *J. Neurosci.* **26**, 10129–10140 (2006).

35.   Oblak, A. L. *et al.* Comprehensive Evaluation of the 5XFAD Mouse Model for Preclinical
Testing Applications: A MODEL-AD Study. *Front. Aging Neurosci.* **13**, (2021).

36.   Merlini, M. *et al.* Fibrinogen Induces Microglia-Mediated Spine Elimination and
Cognitive Impairment in an Alzheimer's Disease Model. *Neuron* **101**, 1099-1108.e6
(2019).

37.   Guttenplan, K. A. *et al.* Neurotoxic reactive astrocytes induce cell death via saturated
lipids. *Nature* **599**, 102–107 (2021).

38.   Yu, Y., Ramachandran, P. V. & Wang, M. C. Shedding new light on lipid functions with
CARS and SRS microscopy. *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* **1841**, 1120–
1129 (2014).

39.   Deture, M. A. & Dickson, D. W. The neuropathological diagnosis of Alzheimer's disease.
Molecular Neurodegeneration (2019) doi:10.1186/s13024-019-0333-5.

40.   Itagaki, S., McGeer, P. L., Akiyama, H., Zhu, S. & Selkoe, D. Relationship of microglia
and astrocytes to amyloid deposits of Alzheimer disease. *J. Neuroimmunol.* **24**, 173–182
(1989).

41.   Baron, R., Babcock, A. A., Nemirovsky, A., Finsen, B. & Monsonego, A. Accelerated
microglial pathology is associated with Aβ plaques in mouse models of Alzheimer's
disease. *Aging Cell* **13**, 584–595 (2014).

42.   Orre, M. *et al.* Isolation of glia from Alzheimer's mice reveals inflammation
anddysfunction. *Neurobiol. Aging* **35**, 2746–2760 (2014).

43.   Prakash, P. *et al.* Monitoring phagocytic uptake of amyloid β into glial cell lysosomes in
real time. *Chem. Sci.* **12**, 10901–10918 (2021).

44.   Yen, C.-L. E., Stone, S. J., Koliwad, S., Harris, C. & Farese, R. V. Thematic review
series: glycerolipids. DGAT enzymes and triacylglycerol biosynthesis. *J. Lipid Res.* **49**,
2283–301 (2008).

45.   Chitraju, C., Walther, T. C. & Farese, R. V. The triglyceride synthesis enzymes DGAT1
and DGAT2 have distinct and overlapping functions in adipocytes. *J. Lipid Res.* **60**, 1112–
1120 (2019).

46.   Wilfling, F. *et al.* Triacylglycerol synthesis enzymes mediate lipid droplet growth by
relocalizing from the ER to lipid droplets. *Dev. Cell* **24**, 384–399 (2013).

47.   Kuerschner, L., Moessinger, C. & Thiele, C. Imaging of lipid biosynthesis: How a neutral
lipid enters lipid droplets. *Traffic* **9**, 338–352 (2008).

48.   Stone, S. J. *et al.* The endoplasmic reticulum enzyme DGAT2 is found in mitochondria-
associated membranes and has a mitochondrial targeting signal that promotes its
association with mitochondria. *J. Biol. Chem.* **284**, 5352–5361 (2009).

49.   Hossain A.*, Rajpoot J.*, Wendt V.*, Saklani P., Hembram K.C., Manchanda P., Bisht
K., Sharma K., Paouri E., Yang G., Chen Y., Bai L., Virani S., Fu J., Arora H., Davalos
D., Stone S.J., Chopra G.. Targeting DGAT2 Degradation for Reducing Lipid Droplets to
Combat Alzheimer's Disease. *In preparation*

50.   Uriarte Huarte, O., Richart, L., Mittelbronn, M. & Michelucci, A. Microglia in Health and
615   Disease: The Strength to Be Diverse and Reactive. *Frontiers in Cellular Neuroscience*
616   vol. 15 (2021).
617   51.   Gordon, G. B. Saturated free fatty acid toxicity. II. Lipid accumulation, ultrastructural
618   alterations, and toxicity in mammalian cells in culture. *Exp. Mol. Pathol.* **27**, 262–276
619   (1977).
620   52.   Listenberger, L. L. *et al.* Triglyceride accumulation protects against fatty acid-induced
621   lipotoxicity. *Proc. Natl. Acad. Sci.* **100**, 3077–3082 (2003).
622   53.   Schönfeld, P. & Reiser, G. Why does brain metabolism not favor burning of fatty acids to
623   provide energy-Reflections on disadvantages of the use of free fatty acids as fuel for brain.
624   *Journal of Cerebral Blood Flow and Metabolism* vol. 33 1493–1499 (2013).
625   54.   Sultana, R., Perluigi, M. & Butterfield, D. A. Lipid peroxidation triggers
626   neurodegeneration: A redox proteomics view into the Alzheimer disease brain. *Free*
627   *Radical Biology and Medicine* vol. 62 157–169 (2013).
628   55.   Reynolds, I. J. & Hastings, T. G. Glutamate induces the production of reactive oxygen
629   species in cultured forebrain neurons following NMDA receptor activation. *J. Neurosci.*
630   **15**, 3318–3327 (1995).
631   56.   Carlesimo, G. A. *et al.* Atrophy of presubiculum and subiculum is the earliest
632   hippocampal anatomical marker of Alzheimer's disease. *Alzheimer's Dement.*
633   *(Amsterdam, Netherlands)* **1**, 24–32 (2015).
634   57.   Li, N., Lizardo, D. Y. & Atilla-Gokcumen, G. E. Specific Triacylglycerols Accumulate
635   via Increased Lipogenesis during 5-FU-Induced Apoptosis. *ACS Chem. Biol.* **11**, 2583–
636   2587 (2016).
637   58.   Lizardo, D. Y., Lin, Y. L., Gokcumen, O. & Atilla-Gokcumen, G. E. Regulation of lipids
638   is central to replicative senescence. *Mol. Biosyst.* **13**, 498–509 (2017).
639   59.   Ogrodnik, M. *et al.* Obesity-Induced Cellular Senescence Drives Anxiety and Impairs
640   Neurogenesis. *Cell Metab.* **29**, 1061-1077.e8 (2019).
641   60.   Mawuenyega, K. G. *et al.* Decreased clearance of CNS β-amyloid in Alzheimer's disease.
642   *Science (80-.).* **330**, 1774 (2010).
643   61.   Liu, Z., Condello, C., Schain, A., Harb, R. & Grutzendler, J. CX3CR1 in microglia
644   regulates brain amyloid deposition through selective protofibrillar amyloid-β
645   phagocytosis. *J. Neurosci.* **30**, 17091–17101 (2010).
646   62.   Mandrekar, S. *et al.* Microglia mediate the clearance of soluble aβ through fluid phase
647   macropinocytosis. *J. Neurosci.* **29**, 4252–4262 (2009).
648   63.   Njie, e. M. G. *et al.* Ex vivo cultures of microglia from young and aged rodent brain
649   reveal age-related changes in microglial function. *Neurobiol. Aging* **33**, 195.e1-195.e12
650   (2012).
651   64.   Baik, S. H. *et al.* A Breakdown in Metabolic Reprogramming Causes Microglia
652   Dysfunction in Alzheimer's Disease. *Cell Metab.* **30**, 493-507.e6 (2019).
653   65.   Pluvinage, J. V. *et al.* CD22 blockade restores homeostatic microglial phagocytosis in
654   ageing brains. *Nature* **568**, 187–192 (2019).
655   66.   Huang, Y. *et al.* Microglia use TAM receptors to detect and engulf amyloid β plaques.
656   *Nat. Immunol.* **22**, 586–594 (2021).
657   67.   Castoldi, A. *et al.* Triacylglycerol synthesis enhances macrophage inflammatory function.
658   *Nat. Commun.* **11**, (2020).

68. Chandak, P. G. *et al.* Efficient phagocytosis requires triacylglycerol hydrolysis by adipose
triglyceride lipase. *J. Biol. Chem.* **285**, 20192–20201 (2010).
69. Bohlen, C. J., Bennett, F. C. & Bennett, M. L. Isolation and Culture of Microglia. *Curr.*
Protoc. Immunol. **125**, e70 (2019).
70. Prakash, P., Lantz, T. C., Jethava, K. P. & Chopra, G. Rapid, Refined, and Robust Method
for Expression, Purification, and Characterization of Recombinant Human Amyloid beta
1-42. *Methods Protoc.* **2**, 48 (2019).
71. Bligh E. G. and Dyer W. J. A Rapid Method Of Total Lipid Extraction and Purification.
Can. J. Biochem. Physiol. **37**, 911–917 (1959).
72. Xie, Z., Ferreira, C. R., Virequ, A. A. & Cooks, R. G. Multiple reaction monitoring
profiling (MRM profiling): Small molecule exploratory analysis guided by chemical
functionality. *Chem. Phys. Lipids* **235**, 105048 (2021).
73. Cordeiro, F. B. *et al.* Multiple reaction monitoring (MRM)-profiling for biomarker
discovery applied to human polycystic ovarian syndrome. *Rapid Commun. Mass*
Spectrom. **31**, 1462–1470 (2017).
74. de Lima, C. B. *et al.* Comprehensive lipid profiling of early stage oocytes and embryos by
MRM profiling. *J. Mass Spectrom.* **53**, 1247–1252 (2018).
75. Fahy, E. *et al.* Update of the LIPID MAPS comprehensive classification system for lipids.
J. Lipid Res. **50**, S9–S14 (2009).
76. Guttenplan, K. A. *et al.* Neurotoxic reactive astrocytes induce cell death via saturated
lipids. *Nature* 1–6 (2021).
77. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–
140 (2009).
78. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* **40**, 4288–
4297 (2012).
79. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* **57**, 289–300 (1995).
80. Randolph, C. E., Foreman, D. J., Blanksby, S. J. & McLuckey, S. A. Generating Fatty
Acid Profiles in the Gas Phase: Fatty Acid Identification and Relative Quantitation Using
Ion/Ion Charge Inversion Chemistry. *Anal. Chem.* **91**, 9032–9040 (2019).
81. Randolph, C. E., Marshall, D. L., Blanksby, S. J. & McLuckey, S. A. Charge-switch
derivatization of fatty acid esters of hydroxy fatty acids via gas-phase ion/ion reactions.
Anal. Chim. Acta **1129**, 31–39 (2020).
82. Xia, Y., Wu, J., McLuckey, S. A., Londry, F. A. & Hager, J. W. Mutual storage mode
ion/ion reactions in a hybrid linear ion trap. *J. Am. Soc. Mass Spectrom.* **16**, 71–81 (2005).
83. Xia, Y., Liang, X. & McLuckey, S. A. Pulsed dual electrospray ionization for ion/ion
reactions. *J. Am. Soc. Mass Spectrom.* **16**, 1750–1756 (2005).
84. Londry, F. A. & Hager, J. W. Mass selective axial ion ejection from a linear quadrupole
ion trap. *J. Am. Soc. Mass Spectrom. 2003 1410* **14**, 1130–1147 (2003).
85. Xia, J., Wishart, D. S. & Valencia, A. MetPA: a web-based metabolomics tool for
pathway analysis and visualization. *Bioinformatics* **26**, 2342–2344 (2010).
86. DeVos, S. L. & Miller, T. M. Direct Intraventricular Delivery of Drugs to the Rodent
Central Nervous System. *J. Vis. Exp.* (2013) doi:10.3791/50326-v.

87.   Spandl, J., White, D. J., Peychl, J. & Thiele, C. Live cell multicolor imaging of lipid
       droplets with a new dye, LD540. *Traffic* (2009) doi:10.1111/j.1600-0854.2009.00980.x.

DATA AVAILABILITY

Supplemental tables, figures, and associated content is available with the manuscript. All data analysis is available on GitHub (https://github.com/chopralab/microglia_omics). A web application has been developed for exploring lipid and metabolite mass spectrometry data that will be publicly available at http://microgliaomics-chopralab.appspot.com (for review, username: admin, password: Review). The accession information for raw lipid and metabolite mass spectrometry data is MassIVE MSV000089458:
https://massive.ucsd.edu/ProteoSAFe/dataset.jsp?task=0f7bd7cfaf504869bfac786e4184105e.

CODE AVAILIBILITY

All of the analysis codes for the lipidomics and metabolomics experiments are available on Github at https://github.com/chopralab/microglia_omics.

ACKNOWLEDGEMENTS

We thank the following individuals for their input and assistance: Ms. Anisa Dunham for help with animal breeding and maintenance, as well as PCR experiments; Dr. Christina Ferreira at the Purdue Metabolomics Facility for assistance with lipid mass spectrometry; Dr. Scott McLuckey for access to a modified Sciex QTRAP 4000 triple quadrupole/linear ion trap mass spectrometer to perform ion/ion reactions; Dr. Shane Tichy for his support, and Agilent Technologies for their gift of the Triple Quadrupole LC/MS to the Chopra Laboratory; Dr. J. Paul Robinson and Ms. Kathy Ragheb at the Purdue University Cytometry Laboratories for flow cytometry services; Dr. Chris Nelson in the Department of Neurosciences of the Lerner Research Institute of Cleveland Clinic for editorial assistance. This work was supported by the United States Department of Defense USAMRAA award W81XWH2010665 through the Peer Reviewed Alzheimer's Research Program, NIH National Institute of Mental Health award R01MH128866 and National Center for Advancing Translational Sciences ASPIRE awards to G.C.; the NIH National Institute for Neurological Disorders and Stroke award R01NS112526, and NIH National Institute on Alcohol Abuse and Alcoholism award P50AA024333 to D.D. Additional support, in part, by the Stark Neurosciences Research Institute, the Indiana Alzheimer Disease Center, Eli Lilly and Company; the Indiana Clinical and Translational Sciences Institute grant UL1TR002529 from the NIH, National Center for Advancing Translational Sciences. We also acknowledge the Pathology Research Core in the Robert J. Tomsich Pathology and Laboratory Medicine Institute of Cleveland Clinic for their human tissue services; the Clinical Core of Cleveland Clinic's Northern Ohio Alcohol Center funded by NIH grant P50AA024333; the Purdue University Center for Cancer Research funded by NIH grant P30 CA023168. We also thank current and prior members of the Davalos and Chopra laboratories for critical discussions and day-to-day assistance with experiments included in this manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Select illustrations in figures were

749 made using BioRender.
750
751
DECLARATION OF INTERESTS

753 G.C. is the Director of the Merck-Purdue Center for Measurement Science funded by Merck Sharp
754 & Dohme, a subsidiary of Merck and a co-founder of Meditati Inc., a startup developing smart
755 drugs for mental health indications. The remaining authors declare no competing interests.

FIGURES

Fig. 1: LD abundance in microglia is age-, sex-, and region-dependent in the AD brain
a. Experimental design for labeling and quantifying LDs in acutely-isolated microglia from
5xFAD and WT male and female mice at 3-4 or 5-7 months old. LDs were labeled with BODIPY
dye and quantified using flow cytometry. **b.** Representative graph (left) and quantification (right)
of median fluorescence intensity (MFI) of LDs in live microglia (CD11b$^+$DAPI$^-$) from 3-4-month-
old female mice show no increase in LD content in cells from 5xFAD (red line and bar) compared
to WT animals (blue line and bar). Data represent mean ± SD. Unpaired t-test, N=3 separate
experiments, each including one WT and one 5xFAD mouse. **c.** Quantification of LDs in microglia
from 5-7-month-old female mice shows an increase (shift towards higher BODIPY fluorescence
intensity) in LDs from 5xFAD (red) compared to WT (blue) microglia. Dot plot shows a
homogeneous population of BODIPY^{hi} microglia within the LD$^+$ cell subset from 5xFAD mice
(red dots) compared to microglia from WT mice (blue dots); **P= 0.0068. **d.** Quantification of
LDs in microglia from 5-7-month-old male mice shows an increase in LD content in 5xFAD
microglia compared to WT; ****P= 0.00003. **e.** Comparison between BODIPY^{hi} and BODIPY^{lo}
cell populations within LD$^+$ microglia from 5-7-month-old male mice shows more BODIPY^{hi}
microglial cells in 5xFAD mice (red dots) compared to microglia from WT mice (blue dots).
Quantification shows the relative frequency of microglia containing LDs in the BODIPY^{hi} gate in

5xFAD and WT microglia; **P= 0.0039. For **c-e**: Data represent mean ± SD. Unpaired t-test, N=3

separate experiments, each including two WT and two 5xFAD mice. **f.** Principal component

analysis (PCA) plot depicts a clear separation based upon variation in microglial lipidomes from

5-7-month-old WT and 5xFAD female mice. **g.** Graph shows the identification and relative

amounts of specific TAG lipid species that were increased in microglia from 5-7-month-old female

5xFAD mice compared to cells from WT controls. For **f-g:** N=3 separate experiments, each from

one WT and one 5xFAD mouse. **h.** Label-free SRS imaging of LDs and Aβ plaques in 5xFAD and

WT brain hippocampal slices. Increased LDs were observed and quantified in 5xFAD brain

sections, often associated with Aβ plaques (3, 4), compared to WT tissues (1, 2). Twelve areas

were quantified for each group (WT and 5xFAD), from the same brain section. Unpaired t-test,

N=2 separate experiments, each including one WT and one 5xFAD mouse. **i.** Immunofluorescence

of IBA1, and counter-staining for LDs (LipidTox), and Aβ plaques (Methoxy XO4; MO4) in the

cortex, CA1, and subiculum regions from 5xFAD and WT mice. Quantification showed a trend

towards increased % LD area within microglia in the cortex and CA1 regions, which was

statistically significant in the subiculum (*P= 0.0147) from 5xFAD compared to WT mice. Data

represent mean ± SD. Unpaired t-test, N=3 separate experiments, each including one WT and one

5xFAD mouse. **j.** Detection of lipid droplets in human hippocampal formalin-fixed paraffin-

embedded (FFPE) tissue from AD and non-symptomatic (NS) cases. Immunofluorescence was

performed on 15µm sections for the detection of lipid droplets (PLIN2), Aβ plaques (AmyloGlo),

and microglia (IBA1). Representative images show an increase in the density of PLIN2$^+$ LDs in

AD compared to non-symptomatic cases. Higher magnification inserts show an increase of LDs

in microglia from AD patients compared to NS individuals. **k.** Quantification of the number of

PLIN2$^+$ LDs per mm^3 of imaged volume (LD density) shows a significant increase in AD

compared to NS cases; ****P= 0.000007. **l.** Quantification of percentage of microglial volume

occupied by LDs over the total microglial volume per imaged volume of hippocampal tissue shows

an increase in AD compared to NS cases; ***P= 0.0002. For **k**, **l**: quantification was performed in

3D reconstructed confocal z-stacks using Imaris; Data represent mean ± SEM. Unpaired t-test,

N=6 (3 males and 3 females) per group.

Fig. 2: LD-laden microglia are in close proximity to amyloid plaques in mice and humans and exhibit phagocytosis deficits.

a. Parallel quantification in three different 5xFAD brain regions, shows that LD density seems to correlate with plaque density, with the subiculum area of the hippocampus (Sub) where plaque density is highest also demonstrating the highest LD density compared to CA1 or cortex (Ctx). Data represent mean ± SD. N=3 5xFAD mice. **b.** Quantification of % LD$^+$ microglia that are plaque-proximal or -distant in the subiculum of 5xFAD mice. Out of all microglia, 39% were in contact with plaques, while 61% were away from plaques. Out of plaque-proximal microglia, 83%

were LD$^+$, whereas only 17% were LD$^-$. N=3 5xFAD mice. **c.** In the subiculum of 5xFAD mice,
microglia (IBA1, green) exhibited larger cell bodies, shorter processes, and higher LD content
(LipidTox, red) when in close proximity to Aβ plaques (MO4, blue), compared to plaque-distant
microglia. **d.** Quantification showed a significantly higher morphological index in plaque-
proximal compared to plaque-distant microglia in the 5xFAD subiculum; ***P= 0.0003. Data
represent mean ± SD. Unpaired t-test, N=3 5xFAD mice. **e.** Immunofluorescence for lipid droplets
(PLIN2), amyloid plaques (AmyloGlo), and microglia (IBA1) revealed larger LD volume in
plaque-proximal microglia in the hippocampus of AD patients compared to plaque-distant
microglia. **f.** The average number of IBA1$^+$ microglial fragments containing LDs per AD patient,
was significantly increased within 10μm from the closest amyloid plaque compared to LD$^+$
microglial fragments detected 10-20μm (**P= 0.003095), 20-30μm (***P= 0.000455), 30-40μm
(P= 0.000095) or >40µm (**P= 0.008647) from the closest amyloid plaque. **g.** The sum volume
of all LDs within LD$^+$ microglial fragments was larger in cells located within 10μm from the
closest amyloid plaque compared to LD$^+$ microglial fragments detected 10-20μm (**P=
0.001158), 20-30μm (***P= 0.000241), 30-40μm (****P= 0.000065) or >40µm (**P= 0.001545)
from the closest amyloid plaque. For **f-g**: Data represent mean ± SEM. One-way ANOVA with
Tukey's multiple comparison tests, N=6 AD cases (3 males and 3 females). Individual values
shown were averaged from 4 z-stacks imaged per patient. **h.** Experimental design for determining
the phagocytic capacity and LD load of microglia from 5xFAD and WT female mice (5-7 months
old). Microglia were isolated from mouse brains, acutely seeded onto the culture plates for 1 hour,
treated with the Aβ^{pH} probe for 30 mins, and with the LipidTox dye for another 30 mins before
flow cytometry analysis. **i.** Quantification of LDs in Aβ^{pH}- (blue) or vehicle-treated (cyan)
microglia from WT mice with fluorescence minus one (FMO) Aβ^{pH} only control (grey). Aβ^{pH}
treatment induced an increase in LDs in WT microglia; ***P= 0.0002. **j.** Quantification of LDs in
Aβ^{pH}- (red) or vehicle-treated (pink) microglia from 5xFAD mice with FMO Aβ^{pH} only control
(charcoal). Aβ^{pH} treatment did not induce an increase in LDs in 5xFAD microglia. **k.**
Representative dot plots showing LD and Aβ^{pH} uptake by microglia from WT and 5xFAD mice.
Microglia from 5xFAD mice showed reduced Aβ^{pH} uptake compared to microglia from WT mice.
l. Quantification of Aβ^{pH} uptake showed a phagocytic deficit in LD$^+$ microglia from 5xFAD
compared to LD$^+$ microglia from WT mice; **P= 0.0019. For **i, j** and **l**: Data represent mean ± SD.
Unpaired t-tests, cells were pooled from 3 mice per group (3 WT and 3 5xFAD mice) for each of
the N=3 experiments.

Fig. 3: Aβ induces profound changes to the microglial lipidome and metabolome *in vitro*,
resulting in LD formation.
a. Experimental design for the global lipidomic profiling experiment performed on Aβ- and
vehicle-treated primary mouse microglia. Cells were isolated from ~7-month-old C56BL/6J
perfused mouse brains and cultured in growth medium containing TGF-β, IL-34, and cholesterol.
Cells isolated from each brain were split and treated with 500 nM Aβ or vehicle for 1, 12, and 24
hours, followed by lipid and metabolite extraction from conditioned media and cell pellets, which
were run on the Agilent triple quadrupole mass spectrometer. Lipids and metabolites were
identified in the samples using MRM-profiling. Each experiment was repeated 5 or 6 times

resulting in: N=5 mice were used for 1-hour treatments, N=6 mice for 12 hours, and N=6 mice
for 24 hours treatments. **b.** PCA demonstrating the variation in microglial lipidomes both within
and between groups (Aβ or vehicle treated microglia) at 1 and 24 hours of treatment. **c.** The
distribution of significantly different lipid classes identified in microglia at 1 and 24 hours of Aβ
treatment, compared to vehicle treated cells. 32% of the differentially regulated lipids at 1 hour
were FFA, whereas 59% of the differentially regulated lipids at 24 hours were TAGs. **d.**
Upregulated lipid classes at 1 and 24 hours of Aβ treatment compared to vehicle, showed FFAs
and TAGs were the most abundant lipids, respectively. **e.** Individual lipid species belonging to
FFAs and TAGs that were upregulated at 1 and 24 hours of Aβ treatment, compared to vehicle.
Long-chain saturated FFAs C20:0, C22:0, and C19:0 were the top 3 upregulated FFAs within the
first 1-hour of Aβ treatment, while neutral lipids TAG(52:3)_FA 18:1, TAG(54:3)_FA 18:1, and
TAG(52:2)_FA 18:1 were the top three upregulated TAGs with prolonged 24-hour Aβ treatment,
both compared to vehicle. **f.** Structural identification and confirmation of the C20:0 and C22:0
lipids in the 1-hour Aβ-treated microglial samples, using the gas-phase ion/ion chemistry (see
Supplementary Results). **g.** Percentage changes of maximum ion intensities as a quantitative
measure of changes in the respective amounts of FFAs (green), TAGs (magenta), and CEs (black)
in microglial cells at 1, 12, and 24 hours of Aβ treatment. The reduction in FFAs was followed by
an increase in TAGs and CEs – major components of LDs – suggesting a gradual conversion of
FFAs to TAGs towards LD formation. **h.** Convergent pathways for TAG biosynthesis. Glycerol-
3-phosphate acyltransferase (GPAT); acylglycerol-3-phosphate acyltransferase (AGPAT);
phosphatidic acid phosphatase (PAP); monoacylglycerol acyltransferase (MGAT); and the final
rate-limiting enzyme diacylglycerol acyltransferase (DGAT) that is needed for TAGs production
and is involved in LD formation.
884

885
886
Fig. 4: DGAT2 enzyme is highly expressed in LD-laden plaque proximal microglia in mouse and human AD brain.

a. Proposed mechanism for Aβ-induced LD formation in microglia. Microglial exposure to Aβ induces an upregulation of FFAs that are converted to TAGs within LDs via the DGAT2 pathway. **b.** Immunofluorescence staining of microglia (IBA1), LDs (LipidTox), DGAT2, and Aβ plaques (MO4) in the hippocampal subicular region of 5xFAD and WT mouse brains. Increased DGAT2 is shown in microglia associated with amyloid plaques. **c.** Quantification showing a significant increase in % of DGAT2[+] microglia out of all microglia and out of all DGAT2[+] cells in the mouse subiculum in the 5xFAD tissue vs. WT; *P= 0.0181, ***P= 0.0002. Data represent mean ± SD. Unpaired t-test, N=3 mice per group. **d.** DGAT2 expression in LD[+] microglia in close proximity to amyloid plaques in hippocampal FFPE tissue from human AD and NS cases (N=4 per group). Immunofluorescence was performed on 15µm-thick human hippocampal sections for the detection of DGAT2, lipid droplets (PLIN2), amyloid plaques (AmyloGlo) and microglia (IBA1). Increased DGAT2 signal (yellow) was detected in plaque-proximal LD[+] microglia in AD cases (arrows), compared to NS controls. Cross-sections of the selected microglial cells in the white boxes demonstrate representative example of increased DGAT2 signal in close proximity to a large PLIN2-labeled LD inside a plaque-proximal microglial cell in AD, and of a cell from a non-symptomatic case.

Fig. 5: DGAT2 enzyme is required for Aβ-induced LD formation, and inhibiting or degrading it restores Aβ phagocytosis, decreases LD and plaque burden in microglia.

a. Experimental design for determining the phagocytic capacity and LD load of microglia from 5xFAD and WT female mice (5-7 months old). Microglia were isolated from mouse brains, acutely seeded onto the culture plates with D2i or Veh for 1 hour, followed by sequential treatment with Aβ^{pH} probe and LipidTox dye, in presence of D2i or Veh, before flow cytometry analysis. **b.** DGAT2 inhibitor (D2i) treatment reduced LDs in cultured microglia from WT and 5xFAD brains; ***P= 0.0001, **P= 0.0029. **c.** Quantification showed that D2i treatment reduced LD formation

upon Aβ exposure in microglia from WT mice but not in cells from 5xFAD mice; ***P= 0.0001.
d. Representative dot plots showing LD and Aβ^{pH} uptake by microglia treated with D2i from WT
and 5xFAD mice. **e.** LD$^+$ microglia from WT mice showed a slight but non-significant increase
in Aβ^{pH} uptake with D2i, while LD+ microglia from 5xFAD mice showed a significant increase
in Aβ^{pH} uptake with D2i; ****P= 0.000014. **f.** Direct comparison of the effect of D2i treatment on
Aβ^{pH} uptake by LD$^+$ microglia from WT and 5xFAD showed that inhibiting DGAT2 restored the
phagocytic performance of 5xFAD microglia, making it comparable to that of WT cells. For **c, d,**
e, and **f**: Data represent mean ± SD. Unpaired t-tests, cells were pooled from 3 mice per group (3
WT and 3 5xFAD mice) for each of the N=3 experiments. **g.** Schematic showing the delivery of
DGAT2 degrader into the lateral ventricles of 18-24 months old 5xFAD mice using
subcutaneously implanted osmotic pumps. **h.** Immunofluorescence for microglia (IBA1), LDs
(LD540), Aβ plaques (MO4) in the hippocampal subicular region of vehicle and DGAT2 degrader-
treated 5xFAD mice showed evident LDs and Aβ plaques reduction following DGAT2 degrader
treatment. **i.** Quantification showed a significant decrease in total LD area and Aβ plaque area in
the subiculum of 5xFAD mice treated with DGAT2 degrader vs. vehicle; *P<0.05. Data represent
mean ± SD. Unpaired t-test, N=5 mice received DGAT2 degrader and N=3 mice received vehicle
treatment. **j.** Model proposing DGAT2 as the target in AD for Aβ-induced LD formation and
phagocytic dysfunction in microglia. Inhibition or degradation of DGAT2 resulted in increased
Aβ uptake and reduced plaque burden, respectively, while reducing LD load.
935

SUPPLEMENTARY TABLE LEGENDS

Supplementary Table ST1: Analyzed lipidomics data of 5xFAD vs. WT microglia, including female and male samples. Tables show the differential lipid profiles in 5xFAD vs. WT microglia samples, organized by most to least differentially-expressed lipids. Samples N1, N2, N3 belong to female mice and samples N4, N5 belong to male mice. P and FDR values are also provided and lipids with FDR<0.1 were considered to be significant.

Supplementary Table ST2: Analyzed lipidomics data of 5xFAD vs. WT microglia, female only. Table shows the differential lipid in 5xFAD vs. WT female microglia samples, organized by most to least differentially-expressed lipids. P and FDR values are also provided and lipids with FDR<0.1 were considered to be significant.

Supplementary Table ST3: Raw MRM lipidomics data of 5xFAD vs. WT microglia. Table lists the MRM transitions screened for TAGs and the respective ion intensity values, organized by MRM transition.

Supplementary Table ST4: Analyzed metabolomics data for 5xFAD vs. WT microglia (male and female). Tables show the differential metabolites in 5xFAD vs. WT microglia, organized by most to least differentially-expressed metabolites. P and FDR values are also provided and metabolites with FDR<0.1 were considered to be significant.

Supplementary Table ST5: Pathway analysis for 5xFAD vs. WT microglial metabolites. A list of all pathways matched to the differentially-expressed metabolites (FDR<0.1) is provided in the table. The pathways with P<0.05 were considered to be significant and are highlighted in the scatter plots in the supplementary figures.

Supplementary Table ST6: Analyzed lipidomics data for Aβ- vs. vehicle-treated primary microglial cells at 1, 12, and 24 hours. Tables show the differential lipid profiles in Aβ- vs. vehicle-treated cells at the three different time points, organized by most to least differentially-expressed lipids. P and FDR values are also provided and lipids with FDR<0.1 were considered to be significant.

Supplementary Table ST7: Analyzed lipidomics data of Aβ- vs. vehicle-treated primary microglial cell conditioned media at 1, 12, and 24 hours. Tables show the differential lipid profiles in Aβ- vs. vehicle-treated conditioned media samples at the three different time points, organized by most to least differentially-expressed lipids. P and FDR values are also provided and lipids with FDR<0.1 were considered to be significant.

Supplementary Table ST8: Analyzed metabolomics data of Aβ- vs. vehicle-treated primary microglial cells at 1, 12, and 24 hours. Tables show the differential metabolites in Aβ- vs. vehicle-treated cells at the three different time points, organized by most to least differentially-expressed metabolites. P and FDR values are also provided and metabolites with FDR<0.1 were considered to be significant.

Supplementary Table ST9: Analyzed metabolomics data of Aβ- vs. vehicle-treated primary microglial cell conditioned media at 1, 12, and 24 hours. Tables show the differential metabolites in Aβ- vs. vehicle-treated conditioned media samples at the three different time points, organized by most to least differentially-expressed metabolites. P and FDR values are also provided and metabolites with FDR<0.1 were considered to be significant.

Supplementary Table ST10: Raw MRM lipidomics data of Aβ- vs. vehicle-treated primary microglial cell conditioned media at 1 and 24 hours. The table lists the MRM transitions screened for TAGs and the respective ion intensity values at two different time points, organized by MRM transition.

Supplementary Table ST11: Pathway analysis for cell metabolites at 1, 12, and 24 hours of Aβ- vs. vehicle-treated primary microglia cultures. List of all pathways matched to the differentially-expressed metabolites (FDR<0.1) are highlighted in the tables. The pathways with P<0.05 were considered to be significant and are highlighted in the scatter plots in the supplementary figures.

Supplementary Table ST12: Pathway analysis for media metabolites at 1, 12, and 24 hours of Aβ- vs. vehicle-treated primary microglia cultures. List of all pathways matched to the differentially-expressed metabolites (FDR<0.1) are highlighted in the tables. The pathways with P<0.05 were considered to be significant and are highlighted in the scatter plots in the supplementary figures.

SUPPLEMENTARY MOVIE LEGEND

Supplementary Movie 1: 3D rendered plaque-proximal microglia show increased LDs in human AD patients. Representative confocal z-stack for which 3D "Surfaces" were made in Imaris for IBA1, PLIN2 and AmyloGlo channels. Only LDs in microglia (red) are shown. Plaque-proximal microglia (yellow) defined by a 0-10μm distance from the closest plaque (purple) contain increased LDs (red) compared to plaque-distant (>10μm from closest plaque) LD+ microglia (green). Plaque-proximal LD- microglia are also shown in orange.

STAR Methods

Animals
C57BL/6J and 5xFAD mice were obtained from the Jackson Laboratory and were maintained in a pathogen free facility. All experiments involving mice were performed in accordance with the Purdue University's Institutional Animal Care and Use Committee (IACUC) guidelines.

Primary mouse microglia isolation and culture from adult mouse brains
Primary microglia from adult mouse brains were isolated and cultured per a previously-described protocol[43]. Briefly, CD11b+ primary microglia were isolated from adult mice (both male and female) and cultured as described previously[43]. Mice were euthanized with CO_2 according to

IACUC guidelines, and perfused brains were removed and cut into small pieces before
homogenization in 1x Dulbecco's phosphate-buffered saline with Calcium and Magnesium
(DPBS^{++}) with 0.4% DNase-I on the tissue dissociator at 37 °C. After filtering the cells through a
70-μm filter, myelin was first removed using Percoll PLUS reagent (GE Healthcare #45001754),
then again by using myelin removal beads. After myelin removal, CD11b$^+$ cells were selected from
the single-cell suspension using the CD11b beads (Miltenyi) as per the manufacturer's instructions.
The CD11b$^+$ cells were finally resuspended in microglia growth media[69], further diluted in TIC
(TGF-β, IL-34, and cholesterol containing) media with 2% FBS (Atlanta Biologics #S11150, Lot
#H17115) before seeding 1x10^5 cells per 500 μL in a well of a 24-well plate (Falcon). The cells
were maintained in TIC media at 37°C and 10% CO_2, with media being changed every other day
until the day of Aβ treatment (around 12-14 days *in vitro* (d.i.v.)).
1037
Aβ preparation
The solid human Aβ1-42 peptide (Anaspec #20276) was prepared per our previously-described
protocol[70]. Briefly, the peptide was dissolved in 20 mM NaOH, pH=10.5 to make ~100 μM stock
solution. Peptide aggregation was initiated by incubating the solution at 37°C for 24 hours. The
peptide was then either stored at -80°C or used directly on the cells after being diluted in culture
medium and filtered through a 0.22 μm syringe filter.
1044
Treatment of Aβ
Primary microglia were treated with 500 μL/well of 500 nM Aβ1-42 or vehicle for 1, 12, or 24
hours. After the respective time points, the conditioned media (CM) from each well was collected
and stored at -80°C. The cells were then detached from wells with 0.25% trypsin and collected in
1x phosphate-buffered saline (PBS) before pelleting at 500 x*g* for 6 mins at 4°C. The supernatant
was aspirated, and pellets were also stored at -80°C along with the CM for direct injection-MS/MS
and MRM-profiling.
1052
Lipid extraction by the Bligh & Dyer method
Lipid and metabolite extracts were prepared using a slightly modified Bligh & Dyer extraction
procedure[71]. The frozen cell pellets from primary microglia were thawed at room temperature, and
ultrapure water, methanol, and HPLC-grade chloroform were added to the pellets. The samples
were vortexed, resulting in a one-phase solution that was then incubated at 4°C for 15 mins. Next,
ultrapure water and chloroform were added, resulting in a biphasic solution. The samples were
centrifuged at 16,000 x*g* for 10 mins, resulting in 3 phases in each tube. The bottom organic phase
containing the lipids were transferred to new tubes, while the middle phase consisting of proteins
and the upper polar phase were discarded. The solvents from the organic phase were evaporated
in a speed-vac. The dried lipid extracts were dissolved in acetonitrile/methanol/300 mM
ammonium acetate (3:6.65:0.35 v/v/v). The lipid extract solutions were diluted further 50 times
before running them on a mass spectrometer.
1065
Unbiased lipidomic and TAG-species profiling using MRM-profiling
To determine if there were differences in lipid profiles that occur with Aβ activation over 24 hours,
lipids extracted from cell lysates as well as conditioned medium were processed for Multiple
Reaction Monitoring (MRM)-profiling[72,73]. Instruments used in these experiments are listed in the
supplemental material. The detailed methodology of MRM-profiling for targeted lipid profiling

has been described previously[74]. This method enabled the interrogation of the relative amounts of
numerous lipid species within ten major classes of lipids based upon the LipidMaps database[75].
The lipid classes, along with the total number of MRM transitions screened, are presented in **Fig.**
S1b. Due to the large number of triacylglycerol (TAG) lipid species interrogated in this study,
TAGs were run in two separate methods (TAG 1 and TAG 2), each measuring relative ion
transitions for different TAG species. The TAG species measured in each method were arbitrarily
divided. For sample preparation, dried lipid extracts were diluted in methanol:chloroform (3:1 v/v)
and injection solvent to obtain a stock solution. Then, the diluted lipid extract was delivered to the
ESI source of an Agilent 6410 triple quadrupole mass spectrometer to acquire the mass
spectrometry data by flow injection (no chromatographic separation). The raw MS data obtained
were processed using an in-house script, and the lists containing MRM transitions along with the
respective ion intensity values were exported to Microsoft Excel for statistical analyses to identify
the significant lipids and metabolites in Aβ-treated versus vehicle-treated microglia. Individual
TAG species were also profiled in primary cultured microglial cells treated with Aβ and vehicle
controls, as well as 5xFAD and WT acutely-isolated microglia using MRM-profiling methodology
as described above. Briefly, diluted lipid extracts were directly infused into the Agilent Jet Stream
ion source of an Agilent 6495C triple quadrupole mass spectrometer. TAG molecular species were
identified based upon previously-established MRMs.
1089
Statistical analyses for lipidomics and metabolomics
Statistical analyses were performed according to our recently-published study[76]. The edgeR
package[77] was used for Aβ- versus vehicle-treated microglia comparisons, as well as comparisons
of 5xFAD vs. WT microglia. Abbreviation "s" was used to denote sample, which represent
different replicates of an analyte class and "b" was used to indicate a biomarker such as a single
lipid or metabolite. Ion counts of a biomarker were denoted using these two subscripts. The
experimental blank that was done with the injection media was modeled as an 'intercept' sample
in the analysis to make certain that comparisons were significant with respect to the blank. A
generalized linear model was fitted using the edgeR package[77] for the mean variance as follows:

$$\log \mu_{bs} = X_s^T \beta_b + \log N_s$$

1102 Here, N_s represents the sum of all ion intensities for the sample s. The coefficient of variance (CV)
1103 for a biomarker ion counts in a sample (y_{bs}) can be calculated using the following equation:

$$CV^2(y_{bs}) = 1/\mu_{bs} + \Phi_b$$

1107 Dispersion of the biomarker was denoted as Φ_b and it was estimated using the common dispersion
1108 method[78]. The associated log-fold change was calculated between the Aβ-treated and vehicle-
1109 treated microglia and p-values were obtained using the likelihood ratio test. The BH method was
1110 used to calculate p-values to acquire false discovery rates (FDRs)[79] and a lipid or a metabolite was
1111 considered to be significant if fold change > 0.5 and FDR < 0.1.
1112
1113 **Lipid droplet staining of acutely-isolated microglia in suspension**
1114 CD11b[+] cells were isolated from male and female mice (3-4 or 5-7 months old; 5xFAD or WT)
1115 and resuspended in 1x PBS, counted using a hemocytometer (1:10 ratio of trypan blue to cell

suspension) and stained with 2 μM BODIPY in 1x PBS for 1 hour at 37°C. The cells were then washed once, resuspended in 1x PBS and taken for analysis on an Attune NxT flow cytometer (Invitrogen) after staining of dead cells with DAPI.

Perfusion and tissue processing
Mice were euthanized with CO_2 and transcardially perfused with PBS and 4% paraformaldehyde (PFA). Brains were extracted and coronally sectioned (50 μm) using a vibratome and stored in antigel solution (30% glycerol, 30% ethylene glycol, in PBS) at -20°C until use for IHC staining.

Immunohistochemistry and staining
Free-floating sections were washed five times in PBS, followed by incubation with Methoxy X04 (10μM, Tocris Bioscience,) solution in PBS for 15 minutes. Sections were then stained with HCS LipidTox Green Neutral Lipid Stain (1:1000, ThermoFisher) in PBS for 15 min, followed by incubation with antigen retrieval buffer (10mM sodium citrate, 0.05% Tween20, pH=6.0) at 70°C for 40 min. The sections were allowed to cool, washed, and treated with 0.1% NaBH4 for 30 min. Following antigen retrieval, sections were blocked with blocking buffer (5% NGS, 0.01% Triton X-100, in PBS) for 1 hr at room temperature. The sections were then incubated overnight with the following primary antibodies: anti-IBA1 (1:150, Millipore Sigma) anti-DGAT2 (1:150, ThermoFisher) in blocking buffer at 4°C. Post-incubation, sections were washed thoroughly with PBS + 0.01% Triton X-100 and incubated with secondary antibodies: Goat anti-Mouse Alexa Fluor 594), Goat anti-Mouse Alexa Fluor 647 and Goat anti-Rabbit Alexa Fluor 594 (all from Invitrogen and diluted at 1:500) for 1.5 h at room temperature. Following washes with PBS + 0.01% Triton X-100, sections were mounted on slides, allowed to dry, and coverslipped using Fluoromount-G anti-fade mounting medium (Southern Biotech).

***Ex-vivo* Lipid droplet staining and Aβ phagocytosis assay of acutely-seeded microglia**
Following the protocol described above for cell isolation, 100K CD11b$^+$ cells from 5-7-month-old female C57BL/6J mice were seeded in TIC media for 1 hour, followed by 500 nM Aβ^{pH} (in TIC media) / vehicle treatment for 30 minutes. Post-treatment, the cells were stained with LipidTox (1:200) in 1xPBS for 30 minutes at 37°C. After staining, the cells were washed with 1x PBS, detached, and collected in ice-cold 1x PBS. Three minutes before analysis of each sample, DAPI was used to stain dead cells. Single positive controls, gating strategies and analyses were done as described above, but for this experiment live LipidTox$^+$ or LipidTox$^-$ cells were identified on the Alexa-Fluor 647 channel, while Aβ^{pH+} and Aβ^{pH-} cells were identified on the FITC channel on an Attune NxT flow cytometer (Invitrogen).

Treatment of DGAT2 inhibitor (D2i) and *ex-vivo* lipid droplet staining and Aβ phagocytosis assay of acutely-seeded microglia
The above protocol was followed with slight modifications for D2i (PZ0233, Sigma) treatment. Briefly, CD11b$^+$ cells were seeded TIC media containing 15 μM D2i / vehicle for 1 hour, followed by 500 nM Aβ^{pH} (in TIC media) / vehicle containing 15 μM D2i / vehicle treatment for 30 minutes. Post-treatment, the cells were washed with 1xPBS and stained with LipidTox (1:200) containing 15 μM D2i / vehicle in 1xPBS for 30 minutes at 37°C. After staining, the cells were washed with 1x PBS, detached, and collected in ice-cold 1x PBS containing 15 μM D2i / vehicle and analyzed on the flow cytometer.

1161

Sample	Aβ^{pH}	LipidTox	D2i
LipidTox only	-	+	-
Aβ^{pH} only	+	-	-
Aβ^{pH} + LipidTox	+	+	-
LipidTox+ D2i	-	+	+
Aβ^{pH} + LipidTox+ D2i	+	+	+

1162

***Dgat2* mRNA expression by PCR**
CD11b$^+$ cells isolated from 5-7-month-old female mice (5xFAD and WT) were resuspended in 1x
PBS, counted using a hemocytometer (1:10 ratio of trypan blue to cell suspension) and stored at -
80 °C until next step. Total RNA was isolated and purified by using Quick-RNA Miniprep Kit
(Zymo Research) following the manufacturer's protocol. RNA quantification and purity were
assessed using Varioskan LUX imaging multi-mode reader (Thermo Scientific). Real-time PCR
was conducted using TaqMan probes (Applied Biosystems, Foster City, CA) for Dgat2
(Mm00499536_m1) and the mouse housekeeping gene β-actin (Mm00607939_s1) as an
endogenous control. For PCR amplification, an initial denaturation at 95°C for 15 min was
followed by 40 cycles of denaturation at 95°C for 10 s and annealing at 60°C for 1 min. Reactions
were run in duplicate. Open qPCR software version 1.0.2 (Chai Biotechnologies Inc.) was used
for post-amplification analysis. Cq and Tm values were calculated directly by the instrument
software and used for finding fold change for *Dgat2* gene expression in 5xFAD relative to WT by
the comparative $2^{-\Delta\Delta CT}$ method.

1177

Stimulated Raman Scattering (SRS) microscopy for label-free lipid droplet imaging
A dual-output 80-MHz femtosecond pulsed laser source (InSight X3+, Spectra-Physics) was used
for the excitation of SRS. The wavelength-tunable output (680-1300 nm) was used as the pump
beam and the 1045 nm fixed-wavelength output was used as the Stokes beam. The pump beam
was tuned to 800 nm to image CH vibrations in brain samples. Both beams had a pulse duration
of ~120 fs. The Stokes beam was directed into an acousto-optic modulator (ISOMET, M1205-
P80L-0.5), which was controlled by a radio frequency driver (ISOMET, 522B-L) and modulated
by a function generator (DG1022Z, Rigol). A square wave of 2.5 MHz and 50% duty cycle was
used for laser intensity modulation. The 0th order laser beam from the AOM was used for
excitation. The beams were combined spatially by a dichroic beam splitter and were chirped using
glass rods (SF57, Lattice Electro-Optics). One 150 mm rod was placed only in the probe beam
pathway, while two 150 mm rods were used after combining the two beams. We bent the optical
beam path to double-pass the two chirping rods to increase the chirping. This gives a 1+4 (Stokes
+ combined) chirping configuration, which chirps the pump beam to 3.4 ps and the Stokes beam
to 1.8 ps. The laser power used on the sample was ~ 15 mW for the pump and ~30 mW for the
Stokes beam. A motorized linear translational stage (X-LSM050A, Zaber Technology) was used
to scan the optical delay between pump and Stokes beams, which were converted to Raman shifts
by spectral focusing. The optical delay scanning steps were 10 µm per step. The combined beams
were scanned by a 2D galvo scanner set (GVS002, Thorlabs) installed to an upright microscope
for imaging. A 60x/1.2 NA water immersion objective lens (UPLSAPO 60X, Olympus) was used
to focus the beams onto the sample. The SRS signal was collected by a 1.4 NA oil-immersion
condenser. The pump beam was detected with a photodiode detector (S3994, Hamamatsu) with a

short-pass filter (980 SP, Chroma technology) to reject the Stokes beam. The alternate voltage signal was amplified using a lab-built tuned amplifier centered at 2.5 MHz. The SRS signal was extracted using a lock-in amplifier (HF2LI, Zurich Instruments). A 2D translation stage (H101, ProScan III, Prior Technology) was used to control sample positions and to perform automated large-area image acquisition and stitching. Data acquisition was enabled using a high-speed data acquisition card (PCIe 6363, National Instruments). The laser scanning and image acquisition was performed by custom-written software based on LabVIEW.

Analysis of SRS microscopy data

For analysis, the images were saved in .txt files and processed using ImageJ. Pseudocolors were used to display different chemical compositions. Hyperspectral images were used to obtain spectra of lipid droplets and other chemical compositions of the tissue. The spectral profiles were normalized by using a laser intensity profile obtained from cross-phase modulation. For quantitative image analysis a Gaussian blur filter (r=3) was used to process the original image. Then, the processed image was subtracted from the original image to highlight the lipid droplets. Intensity thresholding and particle analysis were then performed using ImageJ built-in functions for quantitative lipid droplet analysis. For quantitative analysis results, areas from the WT and 5xFAD sections were selected and the lipid droplets were analyzed within each area. Merging of different image channels was performed using ImageJ. The percentage was calculated by dividing the number of pixels corresponding to LD signal by the total number of pixels of the entire image.

Saturated fatty acid structure elucidation using gas-phase ion/ion chemistry

Utilizing gas-phase ion/ion chemistries, the detailed structural elucidation of complex lipids in biological mixtures has been demonstrated previously[80,81]. Here, a charge inversion ion/ion reaction strategy was employed to examine the structure of saturated fatty acids. All experiments were conducted on a Sciex QTRAP 4000 triple quadrupole/linear ion trap mass spectrometer (SCIEX, Concord, ON, Canada) that has been modified to perform ion/ion reactions[82]. To facilitate the mutual storage of oppositely-charged ions, the key instrumental modifications involve the ability to apply AC voltages to the end plates of the q2 reaction cell. Alternately, pulsed nanoelectrospray ionization (nESI) emitters permit the sequential injection of tris-phenanthroline magnesium reagent dications and fatty acid analyte anions[83]. Singly-deprotonated fatty acid anions, denoted $[FA - H]^-$ anions, generated via direct negative nESI of the lipid extract or authentic reference standard were mass-selected with unit resolution during transient through Q1 and subsequently transferred to the high-pressure collision cell, q2, for storage. Next, positive nESI produced tris-phenanthroline magnesium dications, denoted $[MgPhen_3]^{2+}$, which were isolated in Q1 prior to accumulation in the reaction cell q2. The $[FA - H]^-$ anions and $[MgPhen3]2+$ reagent dications were then mutually stored in q2, yielding the $[FA - H + MgPhen_2]^+$ complex cation. Energetic transfer from the reaction cell q2 to the linear ion trap (LIT), Q3, resulted in the neutral loss of a single phenanthroline ligand and the generation of the charge-inverted complex cation referred to as $[FA - H + MgPhen]^+$. Following mass-selection in Q3, the analysis of charge inverted product ions was performed using single-frequency resonance excitation, commonly referred to as ion-trap collision-induced dissociation (CID) (q = 0.383). In summary, reproducible spectral patterns facilitate fatty acid identification[80]. In all cases, mass analysis was performed using mass-selective axial ejection (MSAE)[84].

Pathway analysis for metabolomics data

Metabolomic pathway analysis was performed using the MetPA[85] (metabolomics pathway analysis) tool on MetaboAnalyst 5.0: a free, web-based tool for metabolomics data analysis that uses the KEGG metabolic pathways as the backend knowledge-base. The differentially-regulated metabolites (FDR<0.1) were uploaded into the compound list with hypergeometric test as the enrichment method and relative-betweenness centrality for topology analysis. The KEGG pathway library for *Mus musculus* was chosen as the reference database. All of the matched pathways according to the p values from the pathway enrichment analysis and pathway impact values from the pathway topology analysis were visualized using the "metabolome view" scatter plot.

Human brain tissue staining

Human hippocampal formalin-fixed paraffin-embedded (FFPE) tissue sections from autopsy samples of both male and female Alzheimer's disease (AD) patients (>74 years, n=3 per sex) and non-symptomatic (NS) cases (>62 years, n=3 per sex) were used. NS cases were obtained from individuals without any neurological or psychiatric diagnosis, and no chronic systemic inflammatory or infectious condition. All human post-mortem tissue was obtained from the Pathology Research Core in the Robert J. Tomsich Pathology and Laboratory Medicine Institute of the Cleveland Clinic. Institutional ethical guidelines were followed for the appropriate use of these fully de-identified samples for research purposes, after IRB approval. All tissue samples were cut at 15 μm and standard de-paraffinization procedures in xylene and decreasing concentration ethanol solutions were utilized. Sections were stained for amyloid plaques using Amylo-Glo RTD Amyloid Plaque Stain Reagent (Biosensis) according to the manufacturer's instructions. Then antigen retrieval was performed in 10mM Tris / 1mM EDTA buffer (pH=8.0) for 20 min at 97°C. After cooling down to room temperature, sections were rinsed with distilled water and blocked in 10% normal donkey serum in PBS-Tween-20 0.05% (v/v) for 1 hour. Primary antibodies were added in blocking buffer and sections were incubated for 72hr at 4°C (anti-Adipophilin (PLIN2 Fitzgerald Industries International, 1:200); rabbit anti-DGAT2, (ThermoFisher, 1:200); anti-IBA1, (Millipore, 1:200). Sections were washed with PBS-T 0.05% (v/v) and incubated with secondary antibodies in blocking buffer for 2h at room temperature (Jackson Immunoresearch: donkey anti-guinea pig AF594, 1:500; donkey anti-rabbit FITC, 1:1000; alpaca anti-mouse Cy5, 1:100). Autofluorescence was quenched with TrueBlack-Lipofuscin autofluorescence quencher (Biotium) according to the manufacturer's instructions and sections were coverslipped with anti-fade fluorescence mounting medium (Abcam). Imaging was performed using a Zeiss LSM 800 confocal microscope using a 40x 1.3NA oil immersion lens and quantification of lipid droplets in relation to microglia and amyloid plaques was performed using the surfaces module in Imaris 9.8.2 (Bitplane).

Image processing

Confocal microscopy images in Figures **1i**, **2c** and **4b** were processed using Imaris 9.8.2 (Bitplane) to reduce noise by applying the Gaussian or Median filters. For **1i** and **2c**, remaining non-specific speckles in the IBA1 channel (likely produced during the antigen retrieval and immunofluorescence protocol) were removed by size exclusion of 3D objects smaller than 1 or 2 μm^3 using the Imaris tool "surfaces". A similar approach was also used for confocal microscopy images acquired from human post-mortem FFPE tissue sections that often show artefacts from the deparaffinization and antigen retrieval protocols. 3D objects smaller than 2 or 3 μm^3 (for IBA1)

and 1 µm^3 (for PLIN2) were removed using Imaris. This size exclusion of speckles and artifacts
also ensured that only true LD particles were selected for visualization and quantification in the
PLIN2 channel. For the DGAT2 channel, only the Gaussian filter was applied to reduce noise. The
mouse brain confocal microscopy images were processed using ImageJ and the despeckle tool was
used to remove the fine noise / grainy speckles from the images prior to being used for LD
quantification using the 'Analyze particles' function coupled with the ROI manager tool.
1296
***In vivo* administration of DGAT2 degrader and immunohistochemistry**
The DGAT2 degrader, synthesized in-house, was dissolved in a mixture of DMSO, propylene
glycol, and saline to create a sterile solution. The final solution contained 120 µM of the degrader,
0.6% DMSO, 10% propylene glycol, and 89.4% buffer (0.9% saline solution). Solvents were
added to the protein degrader in the precise order specified and thoroughly mixed before
proceeding to the next step to avoid any possible precipitation. The DGAT2 degrader or vehicle
solutions was then administered to the lateral ventricles of each mouse brain using an
intracerebroventricular cannula connected to subcutaneously implanted mini-osmotic pumps
(Alzet #2001, 7-day pump, 1µl/hr flowrate). The cannula was implanted at the following
coordinates from bregma: -0.5 mm Posterior, -1.1 mm Lateral (Right), and -2.5 mm Ventral (length
of catheter)[86].
1308
After a 7-day treatment with either the DGAT degrader or vehicle, mice were euthanized using
CO2 and transcardially perfused with PBS and 4% PFA. Brains were then extracted and coronally
sectioned (50 µm thick slices) with a Leica VT1200 vibratome. The sections were stored in an
antigel solution (30% glycerol, 30% ethylene glycol in PBS) at -20°C until used for
immunohistochemical staining. Coronal brain sections, which included regions of the dorsal
hippocampus, were used for analyses. Free-floating sections were washed five times in PBS and
then incubated with 2% H_2O_2 in 70% methanol for 5 minutes to eliminate autofluorescence. After
thoroughly washing them with PBS, the sections were treated with 0.1% $NaBH_4$ dissolved in PBS
for 30 minutes to remove free aldehyde groups and autofluorescence. After using blocking buffer
(10% FBS, 3% BSA, 0.5% Triton X-100 in tris-buffered saline, TBS) for 1 hour at room
temperature, the sections were incubated overnight with rabbit anti-IBA1 primary antibody (1:800,
Wako # 019-19741) in the blocking buffer at 4°C. Next, the sections were thoroughly washed with
TBS + 0.01% Triton X-100 and incubated with goat anti-rabbit Alexa Fluor 488 secondary
antibody (Invitrogen, 1:500) for 1.5 hours at room temperature on the following day. After washing
with TBS + 0.01% Triton X-100, the sections were incubated with Methoxy X04 (10 µM, Tocris
Bioscience) and LD540 (1:500)[87] in TBS for 15 minutes and mounted on slides, allowed to dry,
coverslipped using Fluoromount-G anti-fade mounting medium (Southern Biotech), and imaged
using a Zeiss LSM900 confocal microscope. The dorsal hippocampus subicular region was imaged
at 20X and used for analyzing LD and Aβ plaques area. The confocal microscopy images
were processed using ImageJ for LD quantification using the 'Analyze particles' function coupled
with the ROI manager tool. The subtract background tool and despeckle tool were utilized to
remove the fine noise and grainy speckles from the images before being analyzed.
1331
Statistical analyses
Data collection was randomized for all experiments and experimenters were blinded for imaging
and data analyses. All statistical analyses were performed using GraphPad Prism version 8.2.1 or

on R version 4.1.2. Mean between two groups were compared using two-tailed unpaired Student's
t-test. Data from multiple groups were analyzed by one-way analysis of variance (ANOVA) with
Tukey's multiple comparison tests. Information on the sample size, numbers of replicates and
statistical test used for each experiment is included in the figure legends.