Title

The impact of common variants on gene expression in the human brain: from RNA to protein to schizophrenia risk

Authors

Qiuman Liang[1†], Yi Jiang[1,2†], Annie W. Shieh[3], Dan Zhou[4,5], Rui Chen[5], Feiran Wang[1], Meng Xu[1], Mingming Niu[6], Xusheng Wang[7], Dalila Pinto[8,9,10,11], Yue Wang[12], Lijun Cheng[13], Ramu Vadukapuram[14], Chunling Zhang[15], Kay Grennan[16], Gina Giase[17], The PsychENCODE Consortium[‡], Kevin P White[18], Junmin Peng[6], Bingshan Li[5], Chunyu Liu[1,16*], Chao Chen[1,19*], Sidney H. Wang[3*]

Affiliations

1 MOE Key Laboratory of Rare Pediatric Diseases & Hunan Key Laboratory of Medical Genetics, School of Life Sciences, and Department of Psychiatry, The Second Xiangya Hospital, Central South University, Changsha, Hunan 410000, China.

2 Department of Epidemiology and Biostatistics, Ministry of Education Key Laboratory of Environment and Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430000, China.

3 Center for Human Genetics, The Brown foundation Institute of Molecular Medicine,

The University of Texas Health Science Center at Houston, Houston, TX 77030, USA.

4 School of Public Health and the Second Affiliated Hospital, Zhejiang University School

of Medicine, Hangzhou, Zhejiang 310058, China.

5 Department of Molecular Physiology and Biophysics, Vanderbilt Genetics Institute,

Vanderbilt University, Nashville, TN 37232, USA.

6 Department of Structural Biology, Department of Developmental Neurobiology, Center

for Proteomics and Metabolomics, St. Jude Children's Research Hospital, Memphis, TN 38105,

USA.

7 Department of Genetics, Genomics, and Informatics, University of Tennessee Health

Science Center, Memphis, TN 38163, USA.

8 Department of Psychiatry, and Seaver Autism Center for Research and Treatment, Icahn

School of Medicine at Mount Sinai, New York, NY 10029, USA.

9 Department of Genetics and Genomic Sciences, and Icahn Institute for Data Science and

Genomic Technology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA.

10 The Mindich Child Health and Development Institute, Icahn School of Medicine at

Mount Sinai, New York, NY 10029, USA.

11 Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY

10029, USA.

12 Department of Electrical and Computer Engineering, Virginia Polytechnic Institute and

State University, Arlington, VA 22203, USA.

13 Institute for Genomics and Systems Biology, University of Chicago, Chicago, IL

60637, USA.

14 Department of Psychiatry, The University of Texas Rio Grande Valley, Harlingen, TX

78550, USA.

15 Department of Neuroscience and Physiology, SUNY Upstate Medical University,

Syracuse, NY 13210, USA.

16 Department of Psychiatry, SUNY Upstate Medical University, Syracuse, NY 13210,

USA.

17 The Feinberg School of Medicine, Northwestern University, Chicago, IL 60611, USA.

18 Department of Biochemistry, Yong Loo Lin School of Medicine, National University

of Singapore, Singapore 117596, Singapore.

19 Furong Laboratory, Changsha, Hunan 410000, China.

† These authors contributed equally to this work.

‡ PsychENCODE Consortium authors and affiliations are listed in the supplementary

materials.

* Corresponding author email: Hsi.Ming.S.Wang@uth.tmc.edu (S. H. Wang),

chenchao@sklmg.edu.cn (C. Chen), LiuCh@upstate.edu (C. Liu)

Abstract

Background

The impact of genetic variants on gene expression has been intensely studied at the transcription level, yielding invaluable insights into the association between genes and the risk of complex disorders, such as schizophrenia (SCZ). However, the downstream impact of these variants and the molecular mechanisms connecting transcription variation to disease risk are not well understood.

Results

We quantitated ribosome occupancy in prefrontal cortex samples of the BrainGVEX cohort. Together with transcriptomics and proteomics data from the same cohort, we performed cis-Quantitative Trait Locus (QTL) mapping and identified 3,253 expression QTLs (eQTLs), 1,344 ribosome occupancy QTLs (rQTLs), and 657 protein QTLs (pQTLs) out of 7,458 genes from 185 samples. Of the eQTLs identified, only 34% have their effects propagated to the protein level. Further analysis on the effect size of prefrontal cortex eQTLs identified from an independent dataset clearly replicated the post-transcriptional attenuation of eQTL effects. We identified omics-specific QTLs and investigated their potential in driving disease risks. Using a variant based approach, we found expression-specific QTLs (esQTLs) for 1,553 genes, ribosome-occupancy-specific QTLs (rsQTLs) for 155 genes, and protein-specific QTLs (psQTLs) for 161 genes. Among these omics-specific QTL, 38 showed strong colocalization with brain associated disorder GWAS signals, 29 of them are esQTLs. Because a gene could contain multiple QTL signals, each could either be shared across omics or omics-specific, we aggregated QTL signals from each omics for each gene and found 11 brain associated disorder risk genes that are driven predominantly by omics-specific QTL, all of them are driven by variants impacting transcriptional regulation. This gene-based approach also enabled us to categorize risk genes

containing both omics-specific and shared QTL signals. The limited number of GWAS

colocalization discoveries from gene-based omics-specific mapping, however, prompted us to

take a complementary approach to investigate the functional relevance of genes driven

predominantly by attenuated eQTL signals. Using S-PrediXcan we identified 74 SCZ risk genes

across the three omics, 30% of which were novel, and 67% of these risk genes were confirmed to

be causal in a MR-Egger test using data from the corresponding omics. Notably, 52 out of the 74

risk genes were identified using eQTL data and 68% of these SCZ-risk-gene-driving eQTLs show

little to no evidence of driving corresponding variations at the protein level.

Conclusion

The effect of eQTLs on gene expression in the prefrontal cortex is commonly attenuated post-

transcriptionally. Many of the attenuated eQTLs still correlate with GWAS signals of brain

associated complex disorders, indicating the possibility that these eQTL variants drive disease

risk through mechanisms other than regulating protein expression level. Further investigation is

needed to elucidate the mechanistic link between attenuated eQTLs and brain associated complex

disorders.

Introduction

Complex diseases such as neuropsychiatric disorders are multi-factorial with genetic components

(*1*, *2*). Large scale Genome-Wide Association Studies (GWAS) have uncovered thousands of

disease associated loci, signaling a promising era ahead for causal variant identification (*3*).

However, efforts in fine mapping these disease risk loci often narrowed down the underlying

causal signals to non-coding regions of the genome (*4–6*). Regulatory variants in such non-coding

regions are therefore the prime candidates for driving the genetic risk of disease etiology.

Consequently, integrating gene expression information to pinpoint causal variants or to identify

risk genes has become a staple of genetic studies of complex diseases (*7*), with multiple consortia

efforts facilitating large scale gene expression profiling and regulatory element mapping (*8–10*).

Many powerful methods, such as *coloc*, PrediXcan, SMR/HEIDI, to name a few, have also been

developed to leverage gene expression information for fine mapping GWAS signals or for

identifying underlying risk genes (*11–13*).

Brain associated complex disorders, such as neuropsychiatric disorders, are a group of complex

diseases that can be highly heritable (*14*). For example, schizophrenia (SCZ), a psychiatric

disorder affecting ~1% of the world-wide population (*15*), has a heritability estimate ranged

between 60% to 80%, indicating a strong genetic component (*16*, *17*). Accordingly, recent SCZ

GWAS study reported by Trubetskoy et al. identified 287 significant risk loci and prioritized 120

risk genes using functional genomics (*18*). The use of RNA-Seq data from the brain was

instrumental for risk gene prioritization by Trubetskoy et al., however, information from

downstream gene regulation processes, such as translation rate and protein abundance, was either

not utilized or unavailable.

Measuring transcriptional changes as a proxy for gene activity has a long history in molecular

biology (*19*). In the context of human genetics, buffering of downstream effects of genetic

variants impacting gene expression (i.e. an eQTL) has been shown to be prevalent (*20*). In

addition, QTLs specific to protein level have also been reported (*20-22*). Together, these

observations indicated the importance and potential benefits of including downstream omics

types, such as proteomics data, as information sources for fine mapping disease regulatory

variants. Indeed, recent studies using multi-omics approaches have demonstrated increased power

for risk gene identification among other benefits (*23*, *24*). Of note, our recent work on genetic

variants associated with protein expression level in prefrontal cortex of the human brain indicated

the contribution from non-synonymous coding variants to the changes in protein expression level,

and the utility of these protein QTL variants in prioritizing GWAS risk genes for psychiatric

disorders (*22*).

Another potential benefit of taking a multi-omics approach for identifying disease risk genes rests

in the potential to dissect the fine details of regulatory mechanisms driving the disease-genotype

association. Having relevant datasets to illuminate the origin and propagation of genetic impact

could potentially arrive at a driver regulatory process from a risk variant to a risk gene. Ribo-seq

is a technology that fills in the gap between transcript and protein expression. By adapting RNA-

Seq to a ribosome footprinting method, ribo-seq provides transcriptome-wide quantification of

ribosome occupancy (*25, 26*), which can serve as a proxy for the amount of active translation

synthesizing proteins from each mRNA transcript. When analyzed in conjunction with RNA-Seq

and quantitative proteomics data, ribo-seq enables identification of translational and post-

translational regulatory events (*20, 27*), both major steps defining the Central Dogma of

molecular biology.

As a part of the consortium efforts to improve our understanding of the genetic basis of

neuropsychiatric disorders (*28*), we generated multiple data modalities that included SNP

genotyping, RNA-Seq, ribo-seq, and proteomics of postmortem cortical tissue samples of the

BrainGVEX cohort, which altogether covered the major steps of the Central Dogma. In

conjunction with the quantitative proteomics and transcriptome profiling results that we

previously published (*22, 29*), here we integrated ribo-seq data as our operational definition for

protein synthesis, based on the ribosome occupancy level across the transcriptome, to advance our

multi-omics investigation. Our results reveal regulatory properties of common variants and their

utility in identifying the regulatory processes in the human brain driving disease risk for brain-

associated complex disorders. It offers an opportunity to dissect and appreciate the regulatory

information flow in the biological processes from a population perspective. Additional "rules"

including an effect attenuation process is recognized.

156 Results

Measuring transcriptome-wide ribosome occupancy level in prefrontal cortex of adult human

brain to quantify the level of protein synthesis

To investigate regulatory impact of genetic variants on protein synthesis in the prefrontal cortex

of the human brain, we performed ribosome profiling on 211 prefrontal cortex samples from the

BrainGVEX collection. In total we collected ~62 billion ribosome profiling reads. Consistent with

the expected ribosome footprint size, we found the average insert size of our ribo-seq libraries to

range between 27.4 and 29.5 nucleotides. Similar to prior published studies (*30*), we found on

average 74 % of unwanted reads from ribosomal RNA, tRNA, and snoRNA, which contributed

no information to the translation of protein coding genes. After removing these unwanted reads,

we obtained an average of 30.3 million uniquely mapped informative reads per sample (inter-

quartile range: 20.5 ~ 37.6 million reads) (Fig. S1A). Focusing on the informative reads, we

found the majority (i.e. 84%) of our ribo-seq reads mapped to coding sequence (CDS) exons (Fig.

S1B). Moreover, when visualized in aggregate across annotated coding genes, we found our ribo-

seq data to show strong sub-codon periodicity at the expected positions (Fig. S1C). High

proportion of CDS exon reads and strong sub-codon periodicity indicates the enrichment of

footprints from ribosomes actively engaged in translating mature mRNA and reflects the quality

of the dataset.

Multi-omics cis-QTL mapping identified candidate regulatory variants and revealed translational

and post-translational attenuation of eQTL effects

To identify variants associated with inter-individual expression differences, we perform cis-QTL

mapping for each omics type independently. From the full dataset we filtered out low coverage

genes and removed potentially low quality samples (see details in Data processing section in

Materials and Methods) to reach a mapping dataset of 416 RNA-Seq samples, 195 ribo-seq

samples, 268 proteomics samples, and the corresponding genotype data. Using the mapping

dataset, we identified 12,411 eQTLs (out of 16,540 genes we deem sufficiently quantitated),

2,849 rQTLs (out of 14,572 genes), and 1,036 pQTLs (out of 8,013 genes) at FDR < 0.1. Overall,

we observed similar genomic feature enrichment patterns of QTL variants across the three omics

types (Fig. S2). On the other hand, when focusing on coding variants, we found pQTL to have the

largest proportion of coding variants (pQTL: 11.5% vs. eQTL: 3.4% vs. rQTL: 5.1%). Among the

coding variants, we found 82.4% were missense variants for pQTL comparing to 56.2% for rQTL

and 46.4% for eQTL (Fig. 2A). Of note, we found three start loss variants (i.e. nonsynonymous

substitutions in start codons) in rQTL, which was not found in eQTL.

Intriguingly, we found drastic differences between omics types in the numbers of QTLs mapped

with far fewer rQTLs and pQTLs identified, suggesting reduction of genetic effect along the path

of Central Dogma. However, the differences in the number of genes tested between omics types

and the differences in sample size complicate the interpretation. To better compare the effects of

genetic regulation across data modalities, we identified 185 samples with 7,458 genes that were

sufficiently quantitated across all three omics types (Fig. S3). Using this subset of data, we found

3,253 eQTLs, 1,344 rQTLs, and 657 pQTLs at FDR < 0.1 (Fig. 1A, Fig. S4, Table S1, S2, S3).

Consistently fewer significant QTLs were identified as we moved downstream the Central Dogma

of molecular biology.

A challenge in comparing between the numbers of QTLs identified from each omics type rests in

the fact that not all true effects were identified. While the majority of the QTLs identified here

were replicated in independent datasets (Table S4, S5; Fig. S5), which provides an experimental

support for our discoveries, the level of false negative rates is less clear. Tests replicating QTLs

identified from one omics type in the other omics types can better capture the proportion of

genetic effects shared between QTL types. We performed replication tests using $\pi 1$ estimates

from the qvalue method (*31*). Overall, we found high proportion of QTLs replicated in other

omics types (Fig. 1B). However, when considering the replication rates with the direction of

genetic information flow, we found asymmetric replication rates, with the downstream omics

types to replicate less than the upstream omics types. More specifically, we found 84.7% of the

rQTLs were replicated at the transcript level, but only 60.2% of the eQTLs were replicated at the

ribosome occupancy level. Moreover, while 75.9% of the pQTLs were replicated at the transcript

level, only 34.0% of the eQTLs were replicated at the protein level (Fig. 1B). A similar

asymmetry in proportion replicated between upstream and downstream omics types was observed

when using a direction-aware cutoff-based approach across a wide range of significance cutoffs

(Fig. S6). The lower percentages of eQTLs and rQTLs replicated at the protein level indicate

potential effect attenuation (i.e. either the inter-individual variation in gene expression becoming

smaller and therefore harder to detect or a lack of such effect in the downstream omics types).

Such effect attenuation at the protein level has been previously reported in lymphoblastoid cell

lines (LCLs) (*20*). Our brain data revealed an additional layer of complexity, where effect

attenuation was also observed at the level of mRNA translation.

While our replication tests revealed a trend of effect attenuation for eQTL variants in the

downstream phenotypes (Fig. 1B), these same observations could alternatively be explained by

differences in statistical power between technologies. An independent piece of evidence that is

not sensitive to measurement precision is needed to reach a solid conclusion. Fold change

estimates from less precise measurements are noisier but not biased. In other words, by directly

comparing QTL impact on expression level change across omics types we could gain a direct

view of effect size attenuation. One issue that remains is that using eQTLs identified from our

dataset could bias the results by artificially inflating the variant pool with large eQTL effect sizes,

which might arise simply from applying the QTL mapping procedure to the same dataset (i.e., a

technical ascertainment bias). Using eQTLs independently identified from prefrontal cortex

samples by the CommonMind Consortium (CMC) (*32*), we avoid the ascertainment bias and can

therefore compare the effects size of eQTL variants between the three omics types using our

dataset. A similar approach was successfully implemented to address this power issue in previous

work in LCLs (*20*). Using 5,915 CMC eQTLs that were also quantitated in our dataset, we found

the eQTL effects on the transcript expression level to be significantly larger than their effects on

ribosome occupancy level (per allele $\log_2$ fold differences: mRNA 0.2433 [95% CI=

0.2381~0.2486] vs. ribosome occupancy 0.1836 [95% CI= 0.1794~0.1878]), which were in turn

significantly larger than their effects on protein level (per allele $\log_2$ fold differences: 0.1486

[95% CI= 0.1451~0.1522]) (Fig. 1C, t-test $P < 2.2e^{-16}$ for all pairwise comparisons). Moreover,

for these CMC eQTLs, we found translational regulation to account for more effect size reduction

than post-translational regulation (Fig. 1C). By focusing on the effect sizes of independently

identified eQTLs, our results strongly support the presence of downstream mechanisms

attenuating eQTL effects both at the ribosome occupancy level (translationally) and at the protein

level (post-translationally). A complementary analysis looking at the effect size of pQTL variants

identified from an independent study (*33*) further supported our conclusion. We found, in our

dataset, larger effect size at transcript level when comparing to the fold change at protein level for

these independently identified pQTL SNPs (Fig. S7). We noticed that for these independently

identified pQTL SNPs, the eQTL effect attenuation appears to be predominantly translationally

regulated (Fig. S7B; mRNA vs. ribosome occupancy t-test $P = 6.6e^{-15}$ and ribosome occupancy

vs. protein t-test P = 0.4). Similar results were observed across a wide range of significance levels

used to define pQTL (Fig. S7).

In contrast to prior studies in LCL (*20*), we found clear translational attenuation of eQTL effects

in prefrontal cortex. To further investigate this apparent discrepancy, we cross referenced the

genes associated with eQTL identified in prefrontal cortex samples to those identified in LCL. We

found rather limited overlaps: of the 3,253 eQTL associated genes identified in our brain data

only 598 (i.e. ~18%) were found associated with a significant eQTL in LCL (Fig. S8A),

indicating the possibility that the between tissue differences in gene expression, and their inter-

individual variations, could play a role in the observed differences in translational attenuation of

eQTL effects between brain and LCL. Indeed, gene set enrichment analysis of eQTL associated

genes identified distinct enrichment between eQTLs found only in brain and those found only in

LCL (Fig. S8B, S8C). Importantly, when inspecting the downstream effect of LCL specific eQTL

versus a set of effect size matched brain specific eQTL in our brain dataset we found no

attenuation of eQTL effect at the ribosome occupancy level among LCL specific eQTL while

translational attenuation of eQTL effect remains significant among brain specific eQTLs (Fig.

S8D, S8E), again supporting the notion that the distinct groups of genes analyzed between

cell/tissue types drive the observed discrepancy.

Identifying omics specific QTLs and their signal colocalization with brain disorder GWAS

The prevalent effect size reduction of eQTLs raised the question of the relevance of these genetic

regulation at the organismal level. Because most cellular tasks are executed by proteins, the

genetic regulatory effects not reaching the protein level are, presumably, less likely to have an

impact on organismal traits. More specifically, we seek the biological relevance of expression

specific and ribosome occupancy level specific QTLs (i.e., esQTL and rsQTL). To answer this

question, we first set out to investigate the relevance of different QTL types in brain associated

complex traits such as neuropsychiatric and neurodevelopmental disorders. By applying mediated

expression score regression (MESC) (*34*) to summary statistics from GWAS of four different

complex brain disorders and our three types of molecular QTL summary statistics we estimated

for each disorder the proportion of heritability mediated by each of our molecular QTL types

(Table S6 and Fig. S9). We noticed that for schizophrenia (SCZ), bipolar disorder (BD), and

autism spectrum disorder (ASD), eQTL mediated the most heritability, followed by rQTL, which

is then followed by pQTL. For example, using the Trubetskoy et al. SCZ GWAS (*18*), we found

our eQTLs to mediate 7.09%, rQTLs to mediate 4.06%, and pQTLs to mediate 2.17% of SCZ

heritability (Table S6). On the other hand, for major depressive disorder (MDD) all three QTL

types mediated similar proportion of heritability (~2.5%).

After establishing the relevance for each of the three QTL types in brain associated disorders, we

next sought to identify omics-specific QTLs, in order to further evaluate their relevance in driving

disease risk. For example, we aim to identify expression specific QTLs (i.e. genetic variants that

impact transcript level of the linked genes but not the downstream ribosome occupancy level nor

protein level) that colocalize with GWAS signals of brain associated disorders. To identify omics-

specific QTLs we took a multiple regression approach by first removing effects of the two other

omics-types and then test for genotype association with the residual. To further reduce the number

of false positives, for each potential omics-specific QTL identified at 10% FDR we further test

genotype association with the other two omics-types and remove any variant that has nominally

significant association. We identified 1,553 loci/genes with expression specific QTL (esQTL),

155 genes with ribosome occupancy specific QTL (rsQTL), and 161 genes with protein specific

QTL (psQTL) (Fig. S10). Similar to standard QTLs we found enrichment of esQTL and psQTL in

genomic features such as the exons and the 3'UTRs (Fig. S2). In addition, we found even higher

proportion of coding variants and missense variants among psQTLs comparing to the enrichment

of these categories reported above for pQTL (Fig. 2A). Of note, we found a clear downward shift

in 5'UTR enrichment from pQTL to psQTL (odds ratio 5.62 vs. 1.66, $P << 2.2e^{-16}$). Similar

decrease in 5'UTR variant enrichment was also observed among the other two pairs of standard

and omics-specific QTL (i.e., eQTL vs. esQTL and rQTL vs. rsQTL) indicating the possibility

that QTL variants in 5'UTR are more likely to be shared across omics types, presumably because

they impact protein expression level via transcript level and/or ribosome occupancy level.

To identify genes with omics-specific QTL effects driving disease risk, we perform, for each gene

containing omics-specific QTL signal, *coloc* analyses with default prior (*11*) between the nominal

p value from *cis*-QTL mapping and each brain associated disorder GWAS. We performed

separate analysis using either the nominal p value from the omics-specific *cis*-QTL mapping or

the standard *cis*-QTL mapping. At a colocalization posterior probability cutoff of 70% we

identified risk genes for each omics-type by brain disorder combination for both approaches

(Table S7). We found the most risk genes from esQTL loci, with 14 genes showing signal

colocalization between SCZ GWAS signals and either standard or omics-specific eQTL signals,

and 14 genes for BD, while the remaining combinations resulted in single digit colocalizations

(Table S7). For the majority of the risk genes, we observed consistent results between the two

approaches (Table S7). On the other hand, we found a few risk genes that have GWAS

colocalization with only standard QTL signals, indicating the possibility of GWAS signals

colocalizing with QTL signals that were shared with, and therefore regressed out by, other omics

types at these loci. In other words, some of these omics-specific QTL were identified from loci

containing shared QTL signals that colocalized with GWAS signal.

Because regulation of a gene is often modulated by multiple genetic variants, to evaluate the

consequence of overall *cis*-QTL impact on gene expression, we use PrediXcan to estimate

aggregate genetic regulatory effects for each gene. To identify genes that have omics-specific

QTL effects, for each omics type we built PrediXcan models using normalized quantifications

with and without regressing out signals from the other omics types. A significant PrediXcan

model fit on residuals indicates the potential presence of omics-specific QTL signals (i.e., QTL

signals remained after regression). Using a high PrediXcan model fit R^2 cutoff of 0.1 we

identified candidate genes containing omics-specific QTL signals. To further identify genes with

predominantly omics-specific QTL effects we then remove, from the candidate gene list, genes

that have either shared QTL effects or additional QTL effects from other omics types by filtering

out genes with significant PrediXcan model built from any of the two other omics-types. Using

this approach we identified 382 esQTL genes, 36 rsQTL genes, and 51 psQTL genes that have

predominantly omics-specific QTL effect. Consistently, these predominantly omics-specific QTL

effect genes identified using the PrediXcan gene-based approach are enriched of omics-specific

QTL variants identified from the SNP based multiple regression approach (Fig. S11).

Colocalization with brain-associated complex disorder GWAS identified 11 risk genes with

predominantly omics-specific QTL effects (Table S8). We found most signal colocalization with

SCZ, which resulted in 7 SCZ risk genes with predominantly esQTL effects (Fig. 2B).

For the remaining 629 candidate genes that potentially contain both omics-specific and shared

QTL signals or contain independent omics-specific signals from multiple omics types (i.e. the

candidate omics-specific QTL effect genes filtered out from the above step), we use SuSiE to

identify independent QTL signals from each omics for each gene (i.e. SuSiE 95% confidence

sets). By separating out independent signals form the same gene using SuSiE we can then operate

on the confidence set level to identify additional omics-specific QTL signals that colocalize with

complex disorder GWAS. We identified a total of 829 confidence sets from 594 candidate genes

(Fig. S12). For each confidence set, we perform QTL signal colocalization between omics-types

using *coloc*. We found 404 confidence sets to contain QTL signals that are shared with at least

one other omics types (i.e. PPH4 > 0.7) while 96 contain QTL signals that are shared across all

three omics types (Table S9 full table). Of note, of the 594 candidate genes we found 170 genes to

have at least two confidence sets. Among these genes with multiple confidence sets we found

examples of separate omics-specific QTL signals from different omics types (Fig. 2C) and

example genes that contain both omics-specific and -shared QTL signals (Fig. 2D).

To further investigate the relevance of these QTL confidence sets in brain-associated disorders we

search for colocalization with brain-associated disorder GWAS signals. Of the 829 confidence

sets we found 25 (from 25 genes) to have signal colocalization with at least one of the four brain-

associated complex disorders tested. Of these 25 GWAS colocalization signals 13 have QTL

signals shared between at least two omics types. Upon visually inspecting the nominal p values

from QTL mapping and brain associated disorder GWAS signals at these risk loci we found a few

genes with apparently differential colocalization results between confidence sets. For example, we

found *RCBTB1* to have two eQTL confidence sets while only one of the two confidence sets

contain eQTL signals colocalizing with SCZ GWAS (PPH4 = 0.8). Interestingly, the GWAS

colocalizing eQTL for the *RCBTB1* locus appears to be omics specific while the other confidence

set appears to contain a QTL signal that is shared between transcript expression and ribosome

occupancy (PPH4 = 0.7) (Fig. 2E).

Functional genomics identification of Schizophrenia risk genes

To further investigate the relevance of attenuated eQTLs in brain disease risk, we next took a

complementary approach by first identifying risk genes from each omics type separately and then

investigating the relevant regulatory processes driving disease risks. Given that we observed the

highest number of risk genes for SCZ from our omics-specific QTL analyses, here we focused on

SCZ. We applied S-PrediXcan (*35*) on the Trubetskoy et al SCZ GWAS data and our subset of

multi-omics data. At 5% family-wise error rate, we found 52, 29, and 16 SCZ risk genes,

respectively from RNA-Seq, ribo-seq, and proteomics data (Fig. 3A, Fig. S13). Consistently,

when comparing the corresponding GWAS-QTL *coloc* results between genes that passed and

genes that failed the S-PrediXcan tests, we found S-PrediXcan risk genes to have higher

colocalization posterior probability (Fig. S14). Among S-PrediXcan risk genes only four genes,

NEK4, *KHK*, *CNNM2* and *DARS2* were consistently identified as SCZ risk genes from all three

omics types (Fig. 3B). The majority (74.3%) of the risk genes were identified only from one of

the three omics types. This limited sharing in risk gene identification between omics types is

consistently observed across multiple significance levels and is in clear contrast to the amount of

signal sharing found between QTL types (Fig. 1B; Fig. S15; Table S11, S12, S13).

Among the 74 risk genes we identified using S-PrediXcan (i.e. the union of the risk genes

identified from each of the three omics types), 44 have previously been reported in GWAS studies

as either the mapped genes or as one of the nearby genes under the GWAS peak (Table S14) (*18*,

23, 29, *36–39*). Of these previously reported genes, 27 matched the mapped genes while the

remaining 17 nominated an alternative candidate gene for each GWAS locus (Note that for some

of these 17 loci, the original GWAS signals were mapped to an intergenic region). Comparing our

results to other published multi-omics SCZ risk gene identification studies, we found 15 matched

to the risk genes identified by Giambartolomei et al., which used RNA-Seq and DNA methylation

data from prefrontal cortex of the human brain (*23*), and 12 matched to the 120 prioritized SCZ

risk genes from Trubetskoy et al., which used colocalization with eQTL and Hi-C data (*18*).

When comparing our results to other published TWAS SCZ risk gene identification studies, we

found 19 matched to the risk genes identified by Gandal et al. (*29*), and 21 matched to the risk

genes identified by Gusev et al. (*39*) (See Supplemental Table S14 for an extended list of risk

gene comparisons to prior TWAS studies). On the other hand, for 22 of our 74 risk genes (30%),

we found no match to the known risk gene list (Table S14, bottom rows), which we compiled

based on previous GWAS, TWAS, and other functional genomics risk gene identification studies

(*18*, *23*, *29*, *36–39*). These "no match" novel risk genes have relatively weak SCZ GWAS signals

and are therefore challenging to identify without the additional information provided by our

multi-omics QTL dataset. For example, we found strong colocalization between a modest SCZ

GWAS signal at *2p23.3* and all three types of molecular QTLs of the gene *KHK* (Fig. 3D).

Intriguingly, the *KHK* pQTL is in opposite direction of the *KHK* eQTL and rQTL (Fig. 3C),

suggesting a strong post-translational process regulating its protein level in the opposite direction

to the transcriptional regulation. *KHK*, Ketohexokinase, plays a pivotal role in fructose

metabolism and has been hypothesized to contribute to neuronal glycolysis and the loss of

neuronal functions (*40*). In addition to *KHK*, other novel SCZ risk genes have also been reported

to participate in neuronal/ cognitive functions: For example, in neuroglial cell lines, silencing of

SNX32 leads to defects in neuronal differentiation (*41*). As another example, *NSF*, N-

Ethylmaleimide Sensitive Factor, which encodes a vesicle fusing ATPase, was reported as a

causal factor in intelligence traits (*42*).

Analyses of multi-omics dataset reveal regulatory mechanisms of schizophrenia risk genes

While PrediXcan is a powerful tool for GWAS risk gene identification, it does not control for

potential horizontal pleiotropy (*43*). To this end, we performed two-sample Mendelian

Randomization (MR) with Egger regression to replicate the risk genes we identified from using S-

PrediXcan. Egger regression includes an intercept term, which can be used to evaluate the level of

horizontal pleiotropy (*44*). For each risk gene we first used LD-clumping (*45*) to identify the top

QTL SNP from each clump (i.e. a group of linked variants defined by the LD cutoff) to serve as

strong genetic instruments (*46*). We then tested for the causal relationship between gene

regulation (i.e. QTL signal) and SCZ (i.e. GWAS). We used an operational definition of a causal

effect based on the MR test results (see Methods). At 10% FDR, of the 97 gene-by-omics

combinations (i.e. a total of 74 risk genes including some discovered from more than one omics

type), 67.0% (65/97) passed the MR test. Of those 33.0% that failed the MR test, 37.5% (12 out of 32) failed because of horizontal pleiotropy identified by the Egger intercept test (Table S15). A total of 53 risk genes were replicated in at least one of the three omics types. Note that in order to increase our power, we used a relaxed R^2 cutoff of 0.5 for LD-clumping and adjusted for the linkage between instrument variables in our MR tests using generalized IVW and Egger (*47*). Similar causal effects were observed when using either a stringent R^2 cutoff of 0.01 or using SuSiE (the Sum of Single Effects) (*48*) as an alternative approach to select independent instrument SNPs (Fig. S16). However, these more stringent criteria on instrument variable independence led to lower power to detect and fewer genes available to test (Table S16). For example, when using SuSiE for selecting instrument variables, only 42 genes were tested because some of the risk genes had no fine mapped QTL SNPs according to SuSiE.

A key strength of using a multi-omics QTL approach to identify GWAS risk genes rests in the possibility of further narrowing down the potential regulatory mechanisms. To this end, we further examined the likely causal mechanisms for the 53 replicated SCZ risk genes using one-sample Mendelian Randomization to infer causality between QTL types. We focused our analysis on independently testing for causal effects between neighboring QTL types following the direction of information flow of the Central Dogma (i.e. mRNA -> ribosome occupancy, and ribosome occupancy -> protein). Here we used fine-mapped QTL SNPs identified from the exposure omics types (i.e. the upstream omics types) as instrument variables for one-sample MR analysis. Among the 53 MR-Egger replicated SCZ risk genes, we found 21 genes with significant causal effects both from eQTL to rQTL (i.e. the upstream pathway) and from rQTL to pQTL (i.e. the downstream pathway) (both-passed risk genes; Fig. 4A). Significant causal effects detected from both pathways suggest transcriptionally regulated protein level differences as the potential mechanisms driving these risk genes in SCZ etiology (see an example in Fig. 4B). On the other hand, 25 and 7 replicated risk genes have either significant causal effects only in one of the two

pathways (single-passed risk genes; Fig. 4A) or have no significant causal effects (none-passed

risk genes; Fig. 4A), respectively.

Of note, for 23 of the 25 single-passed risk genes, we found significant causal effects in the

upstream pathway (mRNA -> ribosome occupancy). This asymmetry is reminiscent of the eQTL

effect attenuation described in the prior sections. A failed test could indicate either a true lack of

effect or a lack of statistical power. To take a closer look, we directly assessed the effect size, the

noise level, and instrument strength of the one-sample MR test results. When comparing to the 21

both-passed risk genes, we found clearly smaller effect sizes for the failed tests of both the 7

none-passed risk genes (average 0.577 vs 0.058, t-test $P < 10^{-3}$, Fig. S17A) and the 23 single-

passed risk genes (average 0.524 vs 0.116, t-test $P < 10^{-3}$, Fig. S18A). Note that in both cases the

inter-quartile range of the estimated causal effects for the failed tests covered zero (Fig. S17A,

Fig. S18A). For these same comparison groups, we found no significant differences in instrument

strength (Fig. S17C and S18C) and found slightly higher noise level in the failed MR tests (Fig.

S17B and S18B). When considered together, these observations (i.e. clearly smaller effect size

and slightly elevated noise level for the failed tests) indicate that many of these test results reflect

a true lack of causal effect while in some cases weak effects were obscured by noise. In other

words, many of these risk genes are likely to be driven by omics-specific QTLs. Case in point, we

found *SF3B1* to have strong signal colocalization between eQTLs and SCZ GWAS without clear

QTL signals in either ribo-seq or proteomics data (Fig. 4C).

Discussion

Using a panel of postmortem prefrontal cortex samples, we found clear evidence of post-

transcriptional attenuation of eQTL effects in the human brain. Many of the differences found in

transcriptomes were not present in proteomes. This observation echoes earlier work in HapMap

lymphoblastoid cell lines (*20*) and extends the prior conclusion from *in vitro* cell lines to complex

human tissues. Importantly, distinct from the earlier work in lymphoblastoid cell lines, which

found translation to mostly track with transcription, we found clear attenuation of eQTL effect in

ribo-seq data indicating that translational regulatory processes are involved in eQTL effect

attenuation in the human brain. Prevalent translational attenuation of variant impact on

transcriptional gene expression level has previously been reported in budding yeast (*49*, *50*).

However, results from follow up studies (*51*) appear to present an inconsistent picture. Here,

using replication tests for individual eQTLs and testing for aggregate effect size of eQTLs

independently identified from CMC, we provided strong evidences supporting prevalent

translational attenuation of eQTL effects in the human brain. Importantly, by focusing on

comparing QTL effects (i.e. between genotype fold change) in aggregate, our analyses are robust

to power issues resulted from measurement imprecision (random noise). On the other hand, our

conclusions could still be sensitive to potential biases in the dataset, which could result from

technical issues that cannot be randomized. For this reason we paid close attention to the impact

of sequencing coverage (for sequencing data) and ratio compression (for quantitative mass

spectrometry data) on our major conclusion. Of note, our quantitative mass spectrometry study

employed an analytical procedure that adjusts for contaminating background signals in order to

correct for potential ratio compression in quantification (*52*). Consistently, among low abundance

proteins, which are more susceptible to the influence of ratio compression, we did not find a

disproportional decrease in pQTL effect size (Fig. S19C). Similarly, while genes with low

expression level are more sensitive to issues related to low sequencing coverage, we found little

to no correlation between eQTL/rQTL effect size and expression level (Fig. S19A, S19B).

Although ample precautions and multiple distinct approaches were taken to confirm the observed

effect size attenuation, it remains possible that unknown, unaccounted for omics-specific features

could potentially confound the results. In other words, although operating at the molecular level,
our study remained observational. Future work replicating this observation and elucidating
molecular mechanisms of post-transcriptional attenuation of eQTL effects are needed to provide a
clear understanding of the phenomenon.

Although our study was not designed to answer the key questions regarding the underlying
mechanisms of the observed translational and post-translational attenuation of eQTL effect, the
high prevalence of the observed eQTL effect attenuation when considered in conjunction with the
low prevalence of coding mutations in human population provides some hints. Since a
compensatory variant driving translational or post-translational attenuation is expected to be
either a coding mutation that impact translation elongation rates or protein turnover rates or a
UTR variant that impacts translation initiation rate, the apparent mismatch in incident prevalence
indicated that compensatory variants are unlikely to explain the majority of the effect size
attenuation and therefore implies a broad-spectrum mechanism that reduce variation at the protein
level across genes. Consistently, we observed significant translational and post-translational
attenuation of eQTL effect across all PhastCons score strata on top of a clear trend of decreased
QTL effect size as the evolution conservation scores increase (Fig. S20). Our current work
focuses on the other important follow-up question of whether the attenuated eQTLs are
functionally (biologically) relevant. We attempted to address this question by exploring the
relevance of attenuated eQTLs in brain associated disorders, such as SCZ, a neuropsychiatric
disorder that is highly heritable. We took multiple approaches to identify risk genes that have
either omics-specific QTL signals or attenuated eQTLs. In one approach, we used regression
based methods to identify omics-specific QTL performing separate analysis either at the SNP
level or using PrediXcan to aggregate variant impact at the gene level. In both cases, we found the
most number of colocalization between esQTL and complex disorders (i.e. comparing to rsQTL
and psQTL). From the gene level analysis we further identified interesting example genes

containing multiple QTL signals with varying level of omics-specificity, including an example

gene, *RCBTB1*. We found two strong eQTL signals at *RCBTB1*, while one potentially shared and

one potentially omics-specific. Only the apparently omics-specific eQTL was found to colocalize

with SCZ GWAS. These examples illustrate the complexity of variant impact on gene regulation

and disease risks. Regression based approach, however, could produce false positive omics-

specific QTL simply from noisy predictors. Although we implemented checks and filters to

remove these likely false positives, this caveat is important to keep in mind when interpreting

these results.

In another approach, we conducted our search by first identified risk genes from all quantitated

genes using a TWAS approach, S-PrediXcan method. We then tested for causal regulatory

mechanisms driving the risk gene using QTL SNPs as instrumental variable in a one-sample

Mendelian Randomization test. Before testing for causal regulatory mechanisms, we first

replicated the TWAS risk genes using two-sample Mendelian Randomization with Egger

intercept tests. About 30% of the risk genes that we identified here have not previously been

reported as SCZ risk genes. The use of our functional genomics dataset is key to the discovery of

these novel SCZ risk genes because of their relatively weak GWAS signals (see Fig. 3D for an

example). As a reminder, we would like to emphasize the fact that individual risk gene

identification is dependent on the choice of significance cutoffs, which is to some extend

arbitrary. We used a rather stringent family-wised error rate for risk gene identification in order to

reduce the number of false positive results that could mislead researchers interested in molecular

biology follow-up studies.

To identify causal regulatory mechanisms for each replicated risk gene, we further tested

causality between QTL types using one-sample MR. We found 21 risk genes that are likely

contributing to SCZ risk through transcriptionally regulated protein level. On the other hand, we

also identify 7 genes that show no significant results from MR tests (candidate omics-specific risk

genes) and 23 genes that have significant MR test results only from transcription to translation

(candidate post-translationally-attenuated risk genes). In essence, attempts to identify omics-

specific risk genes (or omics-specific QTLs) are dealing with the challenge of separating true

negatives from false negatives. As such, the interpretation of the failed MR tests is challenging.

Our subsequent analyses looking at comparing instrument strength, noise level, and effect size

between passed and failed MR tests, indicated comparable instrument strength, slightly elevated

noise level in the failed tests, and clearly smaller effect size in the failed tests. In other words, it

remains possible that some of the failed tests could be reflecting a small effect size obscured by

the elevated noise level. In addition to the issues with false negative results, some of the

discoveries could be false positives to begin with. Although we have good confidence with our

FDR and FWER estimates, given that our test statistics for QTL mapping and risk gene

identification are well calibrated (Fig. S4, S13), pleiotropy could introduce positive results from a

different underlying cause (i.e. true effects on SCZ risk but false positive risk genes).

Power issues, however, do not explain the whole story. As was consistently observed throughout

our study, when viewed in aggregate, we see clear effect size differences between omics types,

both for QTLs and for causal effects from MR tests. These effect size estimates are not influenced

by significance cutoffs and are not biased by power differences. Such general trends are also

unlikely to simply be a result of spurious associations or made up entirely of false positives. Our

results therefore bring forth an interesting mechanistic question: how do attenuated eQTL variants

impact SCZ without changing protein levels? One interesting possibility is that the biologically

relevant traits here are protein synthesis rates and protein turnover rates. An association between

genotype and protein synthesis rates could manifest in the form of an attenuated eQTL, if the

differences in per transcript protein synthesis rate (e.g. differences in ribosome elongation rate or

translation initiation rate) appears to offset the differences introduced at the transcript level, which

in turn could resulted in a lack of differences in ribosome occupancy level. Similarly a protein

turnover QTL could also manifest in the form of a rQTL attenuated at the protein level. In other

words, colocalization between an attenuated rQTL and a GWAS signal could actually be

reflecting a colocalization of the GWAS signal with a protein turnover QTL. Testing this

hypothesis requires a direct measurement of protein turnover rate. We hope by presenting the

current results, our findings can inspire future studies on this topic to better understand the

regulatory processes from DNA to brain associated disorders.

Materials and Methods

Data sources

The SNP genotypes (*22*), RNA-Seq (*29*), and quantitative mass spectrometry (*22*) data generated

for prefrontal cortex tissue samples of the BrainGVEX cohort were downloaded from the

PsychENCODE Synapse portal (https://www.synapse.org/#!Synapse:syn5613798) (See Table

S18 for a summary on the number of samples in each dataset and their respective overlap with the

samples in the genotype data). The BrainGVEX cohort includes 420 Caucasians, 2 Hispanics, 1

African American, 3 Asian Americans, and 14 unknown-origin individuals. We also used GWAS

summary statistics data of schizophrenia (*18*), bipolar disorder (*53*), autism spectrum disorder

(*54*), and major depression disorder (*55*) obtained from the Psychiatric Genomics Consortium.

Ribosome profiling

Ribosome Profiling experiments were performed using Illumina's TrueSeq Ribo Profile

(Mammalian) Kit. TrueSeq Ribo Profile (Mammalian) Kit was developed for cell lines. We

adapted the protocol to frozen tissue samples with MP Biomedicals™ FastPrep-24™ Classic

Bead Beating Grinder and Lysis System. Specifically, 60-80 mg of frozen brain tissue was

homogenized in Lysing Matrix D tubes containing 800 µl polysome lysis buffer. The Lysing

tubes were placed on the FastPrep-24™ homogenizer set at 4.0 m/s with 20 s increments until no

visible chunks of tissue remained. Tissue lysate was incubated on ice for 10 min followed by

centrifugation at 20,000g at 4 °C for 10 minutes to pellet cell debris. Clarified lysate was

subsequently used for ribo-seq library preparation following TrueSeq Ribo Profile (Mammalian)

Kit instructions. Indexed libraries were pooled and then sequenced on an Illumina HiSeq 4000.

Note that the experimental protocol for TrueSeq Ribo Profile (Mammalian) Kit that we followed

is a modified version of the previous ribo-seq protocol published by Ingolia and colleagues (*26*),

and it has the following key modifications: Monosome isolation was performed using Sephacryl

S400 spin columns (GE; 27–5140-01) on a tabletop centrifuge instead of ultra-high speed

centrifugation in sucrose cushion. Ribosomal RNA depletion was carried out by using Ribo-Zero

Magnetic Kits and this step is moved up to right after the monosomal RNA isolation step and

before the Ribosome Protected Fragment gel isolation step.

Data processing, gene expression quantification, and normalization

For Genotype data, we obtained Genotype VCF files from PsychENCODE BrainGVEX Synapse

portal (https://www.synapse.org/#!Synapse:syn5613798). The VCF file includes data from 420

individuals genotyped using a combination of 3 different platforms. Details on genotyping

experiments and data processing can be found in our companion manuscripts (*22, 29*). Briefly,

data from whole-genome sequencing (WGS), Human PsychChip platform (a custom version of

the Illumina Infinium CoreExome-24 v1.1 BeadChip), and Affymetrix GeneChip Mapping 5.0K

Array were separately imputed using Minimac3 with Haplotype Reference Consortium (HRC)

reference panel and then merged using PLINK with filters for variant call quality, imputation

quality, minor allele frequency, and relatedness. Only variants with genotype (or imputed) $R^2 >$

0.3, Hardy-Weinberg Equilibrium $> 10^{-6}$ and minor allele frequency (MAF) > 0.01 are kept. The

combined and filtered VCF file contains genotype data of 8,108,028 SNPs for 420 individuals.

We further evaluated potential effects of additional confounding factors on genotype data (Fig.

S21).

For RNA-Seq data, we obtained raw reads in FASTQ format from the PsychENCODE

BrainGVEX project (https://www.synapse.org/#!Synapse:syn5613798). We then process the raw

reads by removing the adapter sequence using cutadapt (v1.12) code "cutadapt –minimum-

length=5 –quality-base=33 -q 30 -a AGATCGGAAGAGCACACGTCTGAACTCCAGTCA -A

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT". The cutadapt processed reads were

then mapped onto GENCODE Release 19 (GRCh37.p13) genome with transcript annotations

from the corresponding version of GTF files using STAR (v2.4.2a). We used RSEM software

(v1.2.29) to quantify read counts for each gene (*56*). The cpm function in the R package "limma"

was used to calculate the log-transformed counts per million (CPM). We filtered out genes with

CPM < 1 in more than 75% of the samples and samples with network connectivity (*57*) z score $<$

-5 (Fig. S22A), which resulted in a total of 17,207 genes from 426 individuals in the

quantification table. We then used normalize.quantiles function in the R package

"preprocessCore" (*58*) to normalize expression level for each sample. DRAMS software was used

to detect and correct mixed-up samples (*59*), which resulted in a final count of 423 individuals.

For ribo-seq data, we used cutadapt (v1.12) to remove adapter sequence from raw reads with code

"cutadapt -a AGATCGGAAGAGCACACGTCT –quality-base=33 –minimum-length=25 –

discard-untrimmed". The processed reads were then mapped against a FASTA file of rRNA,

tRNA, and snoRNA sequence downloaded from NCBI using bowtie2 (*60*) to filter out

uninformative reads. The filtered reads were mapped to GENCODE Release 19 (GRCh37.p13)

genome with corresponding transcript annotation GTF file using STAR (v2.4.2a). The uniquely

mapped reads, as defined by the "NH:i:1" flag of the alignment files, were kept for subsequent

analysis. We used the featureCounts function in the R package "subreads" (*61*) to calculate gene

level read counts for ribosome occupancy. The cpm function in the R package "limma" was used

to calculate log-transformed CPM value. We filtered out genes with CPM < 1 in more than 75%

of the samples and samples with network connectivity (*57*) z score < -3.5 (Fig. S22B), which

resulted in a total of 15,171 genes quantitated from 209 individuals in the quantification table. We

then used the normalize.quantiles function in the R package "preprocessCore" (*58*) to normalize

ribosome occupancy level for each sample. DRAMS software was used to detect and correct

mixed-up samples (*59*), which resulted in a final count of 199 individuals.

For quantitative mass spectrometry data, we obtained protein quantification table from the

PsychENCODE BrainGVEX Synapse portal (https://www.synapse.org/#!Synapse:syn5613798).

This table includes abundance quantification for 11,572 proteins from 268 individuals. The

MS/MS raw data were searched against a composite target/decoy human proteome database. The

target database contains 83,955 entries, including both canonical proteins and isoforms

downloaded from the UniProt database. Since we employed shotgun proteomics, we identified

peptides rather than full-length proteins. In cases where a peptide could be mapped to multiple

homologous proteins, we followed the rule of parsimony: the peptide was first assigned to the

canonical protein. If no canonical form was defined, the peptide was assigned to the protein

isoform with the highest uniquely mapped PSM (Peptide-Spectrum Match). For quantification,

we summarized all peptides that are mapped to the representative protein. These data processing

steps for producing the mass spectrometry quantification table is detailed in Luo et al. (*28*). We

further log-transformed protein abundance for each sample. We filtered out genes with log-

transformed protein abundance < 1 in more than 75% of the samples and samples with network

connectivity (*57*) z score < -6 (Fig. S22C, note that all genes and samples passed the filtering

cutoff). We then used the normalize.quantiles function in the R package "preprocessCore" (*58*) to

normalize protein level for each sample. DRAMS software was used to detect mixed-up samples

(*59*) and found none. We matched protein IDs to gene IDs according to the UCSC database (hg19

version), which resulted in protein level quantifications for 8,330 genes. When matching pQTL

signals to genes for comparison with eQTLs/rQTLs, we used the pQTL signals associated with

the protein isoform that has the highest median abundance.

QTL mapping

Estimating and adjusting for unwanted factors

We used the R package "PEER" (*62*) to estimate hidden factors for RNA-Seq, ribo-seq, and

proteomics data separately. The principle of selecting unwanted hidden factors was to maximize

the variance explained with the least number of factors. We identified 30, 29, and 19 hidden

factors to remove from RNA-Seq, ribo-seq, and mass spectrometry data, respectively (Fig. S23).

For each gene from each omics type, we adjusted the expression level by fitting the selected

hidden factors as predictors in a linear model and taking the residuals as the adjusted expression

level. The adjusted expression levels were then further centered by mean and scaled by standard

deviation.

Genotype association tests

We identified *cis*-region expression QTLs (eQTLs), ribosome occupancy QTLs (rQTLs), and

protein QTLs (pQTLs) separately using QTLTools (1.3.1) (*63*). Because each gene can encode

several protein isoforms, we selected the protein isoform with the highest median abundance as

the representative protein. For each gene, we defined *cis*-region as the region ranging from 1Mb

upstream of the 5' end of the gene to 1Mb downstream of the 3' end of the gene (i.e. the length of

the gene body plus 2 Mb in size). We tested all common SNPs (MAF > 0.05) within the *cis*-

region for each gene. We performed 10,000 permutations to create null distributions to obtain

permutation-based p values and further adjusted permutation-based p values used the beta-

approximation approach implemented in QTLTools to calculate empirical p values. For each

gene, we selected the most significant SNP to represent the QTL signal (i.e. the variant with the

smallest empirical p value) and then calculated a genome-wide FDR using the qvalue function of

the R package "qvalue".

Annotation and enrichment analyses of QTL variants

We used Ensembl Variant Effect Predictor (VEP) version 105 to annotate QTL SNPs. Each

variant was assigned multiple annotations as each variant could map to multiple transcript

isoforms of the same gene. We perform three separate analyses with different counting schemes,

counting all annotations, counting only most frequent annotations, counting only the annotation

with the most severe consequence, to ensure the percentage estimates and conclusions of

enrichment are robust.

For genomic feature enrichment analysis, we extracted 5' UTRs, exons, introns, 3' UTRs, and

intergenic regions from the GENCODE v19 GTF file. The exon regions were extracted directly

from entries in the third column of the GTF file (column label: "exon"). Similarly, gene regions

were extracted directly from the GTF file. Intron regions were calculated by removing exon

regions from gene regions, while intergenic regions were obtained by removing gene regions

from the genome. The 5' UTR and 3' UTR coordinates were generated using the

fiveUTRsByTranscript() and threeUTRsByTranscript() functions from the R package

"GenomicFeatures", respectively. We used bedtools intersect -wa to identify QTL SNPs

overlapping each genomic feature. For each variant we used the most frequent genomic feature

prioritizing exon over UTR and UTR over intergenic when ties occurred. We used Fisher's exact

test to evaluate statistical significance of QTL SNP enrichment in each genomic feature. For genic

features (e.g., 5' UTRs, exons, introns, 3' UTRs) total genic counts were used as the background;

for intergenic features, total cis-region (i.e., mapping window) counts were used as the

background.

Estimating mediated heritability of each omics

We used MESC to estimate the proportion of heritability mediated by different omics separately

for four different brain-associated disorders (*33*). In the first step, we calculated the overall

expression score using the unwanted-factor-adjusted-expression data (see QTL mapping section)

as the individual-level gene expression data and the corresponding BrainGVEX genotype data.

We used the 1000 genome phase 3 genotype data to calculate LD scores in order to match

BrainGVEX genotype data to GWAS genotype data. In the second step, we used the overall

expression score calculated in the previous step and GWAS summary statistics to estimate

heritability mediated by each omics.

Identifying omics-specific QTL

SNP-based approach

We used QTLTools to identify omics-specific QTL variants following the same procedure as

standard QTL mapping, which is described in the above section, except that the gene expression

quantification for the omics-type of interest was adjusted by regressing out signals from the other

two omics types. The regression was achieved by including peer adjusted normalized

quantification of the other two omics types as covariates in QTLTools. For each gene, we selected

the SNP with the most significant genotype-expression association as the candidate omics-

specific QTL SNP and used Benjamini-Hochberg procedure to control for false discovery rates.

To further remove false positives resulted from ineffective removal of signals from the other two

omics types, likely due to poor linear model fit, we removed candidate omics-specific QTLs with

nominal significant association (p values < 0.01) with any of the other two omics types.

Gene-based PrediXcan approach

Building prediction models: We used the PrediXcan software (*13*) to separately build gene

expression prediction models based on RNA-Seq, ribo-seq, and quantitative mass spectrometry

data. We built these models for each gene for each omics type with and without regressing out

signals from the other two omics types (i.e. for each gene using data of each omics type we build

both a standard model and a residual model). For each model, an Elastic Net algorithm was used

for feature selection from SNPs located within the *cis*-region defined for each gene (i.e. gene

body +/- 1Mb flanking regions) based on results from a ten-fold cross-validation. Weights were

produced for every selected SNP of each gene, which were used in the prediction models. For

each model, we calculated the Pearson correlation between the predicted and the observed gene

expression (Cross-validation R, R_{cv}), which was considered as a metric for model prediction

accuracy. Unless otherwise specified, we consider models with R_{cv}^2 above the PrediXcan default

0.01 threshold as models describing genes with significant genetic contribution to expression

regulation (*13*). We produced prediction models for the unified set of 7,458 genes from 185

samples.

Identifying and categorizing genes containing omics-specific genetic effects: We defined genes

with significant residual model as candidate genes containing omics-specific QTL signals using a

high R_{cv}^2 cutoff of 0.1 in conjunction with a FDR cutoff of 0.05 for the model fit. Of these

candidate genes we further classify them by the model fit results from the other omics types. If no

significant standard models were built from any of the two other omics-types, we consider the

gene as a predominantly omics-specific QTL gene. For example, for a gene with significant

residual model built from RNA-Seq data (eQTL signal), if there are no significant standard model

built from either ribo-seq nor proteomics data we consider the gene as a predominantly transcript

specific QTL gene (esQTL). For candidate genes that have significant model built from any of the

other two omics types we consider these candidate genes as potentially harboring multiple QTL

signals from different omics types (either omics-specific or shared QTL effects). To separate out

independent signals from each gene we used SuSiE to identify 95% confidence sets representing

each individual signals. To categorize these independent signals as shared or omics-specific we

then perform *coloc* analysis with data from the other omics-types for each confidence set using

SNPs in modest LD (squared correlation > 0.01) with the confidence set of interest to ensure

sufficient background SNPs and excluding variants that are in linkage (squared correlation > 0.1)

with any other confidence sets. For each confidence set we also performed *coloc* analysis with

brain-associated disorder GWAS, in order to identify QTL signals driving disease risks among

these confidence sets. See the next section for details on signal colocalization analysis using

coloc.

Colocalization

We used *coloc* (*11*) to detect signal colocalization between brain-associated disorder GWAS and

each QTL type at the cis-region of each QTL gene. For each QTL gene, for all common SNPs

(MAF > 0.05) within the cis-region, we use QTL effects and GWAS summary statistics of brain

associated disorders as input for *coloc* analysis. To estimate QTL effects, we calculated, from a

linear model fit using SNP genotype as the predictor for gene quantifications, the predictor

regression coefficient and its corresponding square of standard error. We then used the coloc.abf

function in the R package "*coloc*" (*11*) to calculate the posterior probability of each hypothesis

using the default prior. We use posterior probability of 70% for the colocalization hypothesis (i.e.

PPH4) as the cutoff for reporting our colocalization findings.

TWAS identification of SCZ risk genes

We used S-PrediXcan (*34*) to perform gene-level association tests based on the prediction models

built using our brain prefrontal cortex multi-omics data, which is described in the "*Gene-based*

PrediXcan approach" section, and the SCZ GWAS summary statistics data from PGC3 (*18*). The

association tests were performed separately for each omics. For protein data, we performed

omnibus test to incorporate p values of all protein isoforms together to calculate a single p value

for the corresponding gene. The family wise error rates for SCZ risk genes were calculated using

Bonferroni correction of nominal p values.

Two-sample Mendelian randomization (MR)

To identify causal relationships between each omics type and SCZ we used MR analysis. Here we

used fine mapped QTL SNPs as instruments, gene expression quantification at each omics type as

exposure, and SCZ GWAS signal as outcome.

More specifically, we took the following steps to test for causal relationship between gene

regulation at each omics level and SCZ:

Step 1: Here we used two methods for selecting instrument SNPs: an LD clumping approach and

a SuSiE approach. For the LD clumping approach, for each gene of each QTL type, we used

PLINK with "–clump-kb 1000 –clump-r2 0.5" parameters and a QTL nominal p value cutoff to

define instrument SNPs from separate QTL signals. Since our goal is to test as many risk genes as

we reasonably can, we used a dynamic (greedy) approach in defining the QTL p value cutoff for

instrument SNP selection. We first set a stringent p value cutoff at 10^{-4}, which identified the

majority of our instrument SNPs. For genes that don't have at least 3 instrument SNPs identified

we incrementally relaxed the p value cutoff first to 10^{-3} and then to 10^{-2}. See Supplemental Table

15 for details on the p value cutoffs used for each omics type by risk gene combination.

Independent of the above mentioned, we performed a separate MR analysis using LD clumping

with $r^2 < 0.01$ to ensure independence between instrument SNPs and a p value cutoff of 0.05. For

the SuSiE approach, we used susie_rss() function with default parameters in the R package

"SuSiE" (*47*).

Step 2: We then used the mr_steiger() function in the R package "TwoSampleMR" to test the

causal direction of selected instrument SNPs. For genes failing the Steiger test, we applied the

steiger_filtering() function to verify the direction of each instrument SNP and excluded those with

incorrect directions, ensuring the explained variance by the instrument SNPs was greater for the

exposure than for the outcome.

Step 3: For each gene, we used harmonise_data() function in R package "TwoSampleMR" (*64*) to

harmonize QTLs of each omics type and SCZ GWAS SNPs to be in the same direction (i.e. effect

relative to the same allele).

Step 4: We then performed two-sample MR for each gene by omics type combination separately.

Two-sample MR analysis was done using the R package "MendelianRandomization" (*65*). We

used mr_ivw(data, correl = TRUE) function to conduct generalized IVW and

mr_egger(data,correl = TRUE) function to conduct generalized Egger (see next step). Generalized

IVW accounts for linkage between instrument SNPs, which enables the MR analysis to use linked

instrument SNPs.

Step 5: We used the intercept test (i.e. the Egger method) to test for horizontal pleiotropy, and

used predictor coefficient and its corresponding p value from IVW to determine the effect size

and significance of causal effects for each omics type on SCZ.

Step 6: We used Benjamini-Hochberg control procedure to adjust the nominal p values from IVW

for multiple testing.

Step 7: We define a gene by omics type combination as causal for SCZ if the causal effect test

passed the significance cutoff of FDR < 0.1 and the Egger intercept test passed the intercept p

value > 0.05 cutoff.

One-sample Mendelian randomization

For one-sample MR we used Two-stage predictor substitution (TSPS) approach, which is a

method to find causal relationships between omics types. We performed the following analysis in

two iterations, both following the direction of genetic information flow. In the first iteration, we

tested causal relationships between transcript level and ribosome occupancy level (i.e. mRNA ->

ribosome occupancy). In the second iteration, we tested causal relationships between ribosome

occupancy level and protein level (i.e. ribosome occupancy -> protein).

Step 1: We used LD clumping for instrument SNP selection as described in the previous section.

Here we tested two pathways (mRNA -> ribosome occupancy and ribosome occupancy ->

protein). For mRNA -> ribosome occupancy, we used the SNPs of eQTL with p value $< 10^{-4}$ and

$R^2 < 0.5$ from LD. In order to test as many risk genes as we reasonably can, the same dynamic p

value cutoff approach (i.e. the same as the above described for two-sample MR) was used for

instrument SNP selection. See Supplemental Table 17 for details on the p value cutoffs used for

each risk gene.

Step 2: We then used the mr_steiger() function in the R package "TwoSampleMR" to test the

causal direction of selected instrument SNPs. For genes failing the Steiger test, we applied the

steiger_filtering() function to verify the direction of each instrument SNP and excluded those with

incorrect directions, ensuring the explained variance by the instrument SNPs was greater for the

exposure than for the outcome.

Step 3: We then combined genotype and quantification data of the relevant omics types into a

dataframe: dat_{mr} for mRNA -> ribosome occupancy pathway, dat_{rp} for ribosome occupancy ->

protein pathway.

Step 4: For each gene, we then set formula in R package "OneSampleMR" (*66*): tsps ("ribosome

occupancy ~ mRNA | SNP_1+SNP_2+SNP_3+…+SNP_n", data = dat_{mr}) and tsps ("protein ~ ribosome

occupancy | SNP_1+SNP_2+SNP_3+…+SNP_m", data = dat_{rp}).

Step 5: We used the summary() function to get the slope (i.e. the regression coefficient) and the

corresponding p value of the slope from each tsps object.

Step 6: We used Benjamini-Hochberg control procedure to adjust the p value of the slope for

multiple testing.

Step 7: We defined a relationship for each gene by pathway combination as causal if the test

results passed the significance cutoff of FDR < 0.1.

Step 8: We also used R package "ivreg" with the same dataset and formula as the above described

for tsps objects to calculate F-statistics for each gene. The calculated F-statistics was used to

check instrument strength.

Data availability

All raw data from psychENCODE BrainGVEX project are available through

https://www.synapse.org/#!Synapse:syn5613798.

Code availability

All code, and processed data used in the analyses are available at

https://github.com/qiumanL/multi-omics-qtl.

References

1. A. Fialkowski, T. M. Beasley, H. K. Tiwari, *Emery and Rimoin's Principles and Practice of Medical Genetics and Genomics (Seventh Edition) Ch. 11*, (Academic Press, 2019).

2. A. S. Brown, The environment and susceptibility to schizophrenia. *Prog. Neurobiol.* **93**, 23–58 (2011).

3. N. Solovieff, C. Cotsapas, P. H. Lee, S. M. Purcell, J. W. Smoller, Pleiotropy in complex traits: challenges and strategies. *Nat. Rev. Genet.* **14**, 483–495 (2013).

4. K. Watanabe, E. Taskesen, A. van Bochoven, D. Posthuma, Functional mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

5.   M. G. Heckman, et al., Fine-mapping of the non-coding variation driving the Caucasian

LRRK2 GWAS signal in Parkinson's disease. *Parkinsonism Relat. Disord*. **83**, 22–30

(2021).

6.   D. J. Schaid, W. Chen, N. B. Larson, From genome-wide associations to candidate causal

variants by statistical fine-mapping. *Nat. Rev. Genet.* **19**, 491–504 (2018).

7.   A. Gusev et al., Integrative approaches for large-scale transcriptome-wide association

studies. *Nat. Genet.* **48**, 245–252 (2016).

8.   Wang, L. et al. Brain Banks Spur New Frontiers in Neuropsychiatric Research and

Strategies for Analysis and Validation. *Genomics Proteomics Bioinformatics.* **17**, 402–414

(2019).

9.   ENCODE Project Consortium et al., Expanded encyclopaedias of DNA elements in the

human and mouse genomes. *Nature*. **583**, 699–710 (2020).

10. GTEx Consortium, The GTEx Consortium atlas of genetic regulatory effects across human

tissues. *Science*. **369**, 1318–1330 (2020).

11. C. Giambartolomei et al., Bayesian test for Colocalisation between pairs of genetic

association studies using summary statistics. *PloS Genet*. **10**, e1004383 (2014).

12. E. R. Gamazon et al., A gene-based association method for mapping traits using reference

transcriptome data. *Nat. Genet*. **47**, 1091–1098 (2015).

13.   Z. Zhu et al., Integration of summary data from GWAS and eQTL studies predicts complex

trait gene targets. *Nat. Genet*. **48**, 481–487 (2016).

14.   Bray, N. J. & O'Donovan, M. C. The genetics of neuropsychiatric disorders. *Brain Neurosci.*

Adv. **2**, 2398212818799271 (2018).

15.   R. S. Kahn et al., Schizophrenia. *Nat. Rev. Dis. Primers*. **1**, 15067 (2015).

16.   P. Lichtenstein et al., Common genetic determinants of schizophrenia and bipolar disorder in

Swedish families: a population-based study. *Lancet*. **373**, 234–239 (2009).

17.   P. F. Sullivan, K. S. Kendler, M. C. Neale, Schizophrenia as a complex trait: evidence from

a meta-analysis of twin studies. *Arch Gen Psychiatry*. **60**, 1187–1192 (2003).

18.   V. Trubetskoy et al., Mapping genomic loci implicates genes and synaptic biology in

schizophrenia. *Nature*. **604**, 502–508 (2022).

19.   J. B. W. Wolf, Principles of transcriptome analysis and gene expression quantification: an

RNA-seq tutorial. *Mol Ecol Resour*. **13**, 559–572 (2013).

20.   A. Battle, Z. Khan, S. H. Wang, A. Mitrano, M. J. Ford, J. K. Pritchard, Y. Gilad, Impact of

regulatory variation from RNA to protein. *Science*. **347**, 664–667 (2015).

21.   E. Ferkingstad et al., Large-scale integration of the plasma proteome with genetics and

disease. *Nat. Genet*. **53**, 1712–1721 (2021).

22. Luo, J. et al. Genetic regulation of human brain proteome reveals proteins implicated in

psychiatric disorders. *Mol Psychiatry.* (2024).

23. C. Giambartolomei et al., A Bayesian framework for multiple trait colocalization from

summary association statistics. *Bioinformatics*. **34**, 2538–2545 (2018).

24. C. N. Foley et al., A fast and efficient colocalization algorithm for identifying shared genetic

risk factors across multiple traits. *Nat. Commun*. **12**, 764 (2021).

25. N. T. Ingolia, S. Ghaemmaghami, J. R. S. Newman, J. S. Weissman, Genome-wide analysis

in vivo of translation with nucleotide resolution using ribosome profiling. *Science*. **324**, 218–

223 (2009).

26. N. T. Ingolia, G. A. Brar, S. Rouskin, A. M. McGeachy, J. S. Weissman, The ribosome

profiling strategy for monitoring translation in vivo by deep sequencing of ribosome-

protected mRNA fragments. *Nat. Protoc*. **7**, 1534–1550 (2012).

27. Cenik C et al., Integrative analysis of RNA, translation, and protein levels reveals distinct

regulatory variation across humans. *Genome Res.* **25**, 1610-1621 (2015).

28. S. Akbarian et al., The PsychENCODE project. *Nat. Neurosci*. **18**, 1707–1712 (2015).

29. M. J. Gandal et al., Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia,

and bipolar disorder. *Science*. **362**, eaat8127 (2018).

30. G. A. Brar, J. S. Weissman, Ribosome profiling reveals the what, when, where and how of

protein synthesis. *Nat. Rev. Mol. Cell Biol*. **16**, 651–664 (2015).

31. J. D. Storey, R. Tibshirani, Statistical significance for genomewide studies. *Proc. Natl.*

Acad. Sci. U.S.A. **100**, 9440–9445 (2003).

32. G. E. Hoffman et al., CommonMind Consortium provides transcriptomic and epigenomic

data for Schizophrenia and Bipolar Disorder. *Sci. Data*. **6**, 180 (2019).

33. D. W. Yao, L. 1. Robins, C. et al. Genetic control of the human brain proteome. The

American Journal of Human Genetics 108, 400–410 (2021).

34. J. OʻConnor, A. L. Price, A. Gusev, Quantifying genetic effects on disease mediated by

assayed gene expression levels. *Nat. Genet*. **52**, 626–633 (2020).

35. A. N. Barbeira et al., Exploring the phenotypic consequences of tissue specific gene

expression variation inferred from GWAS summary statistics. *Nat. Commun*. **9**, 1825

(2018).

36. S. Ripke et al., Biological insights from 108 schizophrenia-associated genetic loci. *Nature*.

511, 421–427 (2014).

37. A. F. Pardiñas et al., Common schizophrenia alleles are enriched in mutation-intolerant

genes and in regions under strong background selection. *Nat. Genet*. **50**, 381–389 (2018).

38. M. Lam et al., Comparative genetic architectures of schizophrenia in East Asian and

European populations. *Nat. Genet*. **51**, 1670–1678 (2019).

39. Gusev, A. et al. Transcriptome-wide association study of schizophrenia and chromatin

activity yields mechanistic disease insights. *Nat Genet*. **50**, 538–548 (2018).

40. Johnson RJ et al., Cerebral Fructose Metabolism as a Potential Mechanism Driving

Alzheimer's Disease. *Front Aging Neurosci*. **2**, 560865 (2020).

41. Sugatha, J. et al. Insights into cargo sorting by SNX32 and its role in neurite outgrowth.

eLife. **12**, e84396 (2023).

42. Korologou-Linden R, Leyden GM, Relton CL, Richmond RC, Richardson TG. Multi-omics

analyses of cognitive traits and psychiatric disorders highlights brain-dependent

mechanisms. *Hum Mol Genet*. **32**, 885–96 (2021).

43. D. Zhou, Y. Jiang, X. Zhong, N. J. Cox, C. Liu, E. R. Gamazon, A unified framework for

joint-tissue transcriptome-wide association and Mendelian randomization analysis. *Nat.*

Genet. **52**, 1239–1246 (2020).

44. S. Burgess, S. G. Thompson, Interpreting findings from Mendelian randomization using the

MR-Egger method. *Eur. J. Epidemiol*. **32**, 377–389 (2017).

45. S. Purcell et al., PLINK: A Tool Set for Whole-Genome Association and Population-Based

Linkage Analyses. *Am. J. Hum. Genet*. **81**, 559–575 (2007).

46. A. Gkatzionis, S. Burgess, P. J. Newcombe, Statistical methods for cis-Mendelian

randomization with two-sample summary-level data. *Genet. Epidemiol*. **47**, 3–25 (2023).

47. Burgess, S., Dudbridge, F., Thompson, S. G. Combining information on multiple

instrumental variables in Mendelian randomization: comparison of allele score and

summarized data methods. *Statist. Med*. **35**, 1880–1906 (2016).

48.   Zou Y, Carbonetto P, Wang G, Stephens M. Fine-mapping from summary data with the

"Sum of Single Effects" model. *PloS Genet*. **18**, e1010299 (2022).

49.   C. G. Artieri, H. B. Fraser, Evolution at two levels of gene expression in yeast. *Genome Res.*

24 (2014).

50.   C. J. McManus, G. E. May, P. Spealman, A. Shteyman, Ribosome profiling reveals post-

transcriptional buffering of divergent gene expression in yeast. *Genome Res*. **24**, 422–430

(2014).

51.   F. W. Albert, D. Muzzey, J. S. Weissman, L. Kruglyak, Genetic influences on translation in

yeast. *PloS Genet*. **10**, e1004692 (2014).

52.   Niu M, Cho JH, Kodali K, Pagala V, High AA, Wang H et al. Extensive peptide

fractionation and y 1 ion-based interference detection method for enabling accurate

quantification by isobaric labeling and mass spectrometry. *Anal Chem*. **89**(5), 2956-2963

(2017).

53.   Mullins, N. et al. Genome-wide association study of more than 40,000 bipolar disorder cases

provides new insights into the underlying biology. *Nat Genet*. **53**, 817–829 (2021).

54.   Autism Spectrum Disorders Working Group of The Psychiatric Genomics Consortium.

Meta-analysis of GWAS of over 16,000 individuals with autism spectrum disorder

highlights a novel locus at 10q24.32 and a significant overlap with schizophrenia. *Mol*

Autism **8**, 21 (2017).

55. Meng, X. *et al.* Multi-ancestry genome-wide association study of major depression aids

locus discovery, fine mapping, gene prioritization and causal inference. *Nat Genet* **56**, 222–

233 (2024).

56. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics*. **12**, 323 (2011).

57. M. C. Oldham, P. Langfelder, S. Horvath, Network methods for describing sample

relationships in genomic datasets: application to Huntington's disease. *BMC Syst. Biol.* **6**, 63

(2012).

58. B. M. Bolstad, R. A. Irizarry, M. Astrand, T. P. Speed, A comparison of normalization

methods for high density oligonucleotide array data based on variance and bias.

Bioinformatics. **19**, 185–193 (2003).

59. Y. Jiang et al., DRAMS: A tool to detect and re-align mixed-up samples for integrative

studies of multi-omics data. *PloS Comput. Biol.* **16**, e1007522 (2020).

60. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods*. **9**,

357–359 (2012).

61. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics*. **30**, 923–930 (2014).

62. O. Stegle, L. Parts, M. Piipari, J. Winn, R. Durbin, Using probabilistic estimation of

expression residuals (PEER) to obtain increased power and interpretability of gene

expression analyses. *Nat. Protoc*. **7**, 500–507 (2012).

63.   O. Delaneau, H. Ongen, A. A. Brown, A. Fort, N. I. Panousis, E. T. Dermitzakis, A

complete tool set for molecular QTL discovery and analysis. *Nat. Commun*. **8**, 15452 (2017).

64.   B. L. Pierce, S. Burgess, Efficient Design for Mendelian Randomization Studies: Subsample

and 2-Sample Instrumental Variable Estimators. *Am. J. Epidemiol.* **178**, 1177–1184 (2013).

65.   Yavorska O, Burgess S. MendelianRandomization: an R package for performing Mendelian

randomization analyses using summarized data. *Int J Epidemiol*. **46**(6) (2017).

66.   Terza, J. V., Basu, A. Rathouz, P. J. Two-stage residual inclusion estimation: Addressing

endogeneity in health econometric modeling. *Journal of Health Economics.* **27**, 531–543

(2008).

1015

Acknowledgments

We thank the three anonymous reviewers for their valuable comments, which resulted in

substantial improvement of the manuscript. We thank Majd Alsayed, Miguel Brown, Dominic

Fitzgerald, Amber Thomas, and Mimi Brown for their assistance with RNA-Seq data production

and processing. We thank Richard Kopp at SUNY Upstate Medical University for his help with

wordsmithing. We thank all of the donors and their relatives participated in the brain collections

at Stanley Medical Research Institute and Banner Sun Health Research Institute. We thank Drs.

Maree J. Webster and Thomas Beach for their supports to make the tissue available. We thank the

High Performance Computing Center of Central South University for partial support of

calculation. This project has received funding from National Natural Science Foundation of China

(Grants No. 82022024), National Key R&D Project of China (Grants No. 2024YFA1108002),

Author contributions

1038

1039   **Competing interests**

1040   Authors declare that they have no competing interests.

1041

Figures and Tables

Fig. 1. Genetic regulation of gene expression in the human brain. (A) P-value quantile-

quantile plot between the observed (Y-axis) and the expected based on null distribution (X-axis).

The black line indicates the expected distribution of p values when there are no real QTL signals.

The number of cis-QTLs (i.e. the most significantly associated SNP for each gene) identified at

10% FDR is labeled in the top left inset. **(B)** Replication rate between QTL types. Proportions of

QTLs replicated in the other two omics types are listed in the 3 × 3 matrix. Each row is a

discovery omics type and each element of the row correspond to the proportion QTL signals

1052   replicated in the omics type specified by the column label. For example, only 34% of the eQTL

1053   signals were replicated in the protein data. Ribo: ribosome occupancy. **(C)** Effect size of CMC

1054   eQTL SNPs in BrainGVEX data. Mean and 95% confidence interval of absolute per allele effect

1055   across all CMC eQTL SNPs that were also analyzed in the BrainGVEX union set is shown.

1056
Fig. 2. Omics-specific genetic regulations. (**A**) Categorizing QTL SNPs according to genic

feature annotations. Bar plots showing the percentage of QTL SNPs in each category out of the

total number of QTL SNPs considered. Note the high percentage (28%) of coding variants in

psQTL SNPs out of all psQTL SNPs and the dominating percentage (97.8%) of missense variants

in psQTL SNPs out of coding psQTL SNPs. (**B**) An example SCZ risk gene, *GATAD2A,* with

predominantly transcript specific QTL effects. The red line indicates the top signal colocalization

SNP between eQTL and SCZ GWAS on a Manhattan plot. Boxplots to the right summarize the

normalized gene expression level stratified by eQTL SNP genotypes. Ribo: ribosome occupancy.

(**C, D, E**) Example genes containing multiple QTL signals of varying omics specificity.

Individual confidence set SNPs containing 95% of the QTL signal are color coded on a

Manhattan plot with a vertical line indicating the top SNP that contributes the most signal.

Boxplots to the right summarize the normalized gene expression level stratified by genotypes of

the top SNP. Note that the example gene in (E) is a SCZ risk gene with two confidence sets. Only

one of the two confidence sets' QTL signal colocalizes with SCZ GWAS signal. The bar plots

underneath the expression level boxplots show the log odds ratio +/- standard errors for the

association between the confidence set top SNP genotype and SCZ, which is calculated based on

GWAS summary statistics from Trubetskoy et al.

1074
1075

1076

1077

Fig. 3. Schizophrenia risk genes identified from each of the three omics types, RNA-Seq (mRNA), ribo-seq (ribosome occupancy), and quantitative mass spectrometry (protein), using S-PrediXcan. (A) Manhattan plots showing significance level (i.e. -log10 FWER from S-PrediXcan) of gene expression-schizophrenia association across the genome for genes that pass the 5% FWER significance cutoff for each omics type. The black horizontal dotted line indicated the significance cutoff (5% Bonferroni-adjusted p value). Ribo: ribosome occupancy. **(B)** Venn diagram illustrating the number of overlapping risk genes identified between omics types. **(C)** Boxplots summarizing normalized gene expression level stratified by eQTL SNP genotypes for *KHK*. **(D)** Manhattan plots showing nominal p value distribution for each QTL type and

1089   schizophrenia GWAS for the 1Mb QTL mapping window flanking *KHK*. The red line indicates

1090   the position of the lead colocalization SNP between eQTLs and schizophrenia GWAS.

1091

Fig. 4. Identifying driver regulatory mechanism for SCZ risk genes using Mendelian

Randomization (MR). **(A)** Summaries of MR test results. Left panel: Two-sample MR

replication of S-PrediXcan SCZ risk gene discoveries. Each circle represents a S-PrediXcan

discovery. A solid-colored circle indicates successful replication in a MR test. Right panel: One-

sample MR test results for causality between omics types. Shown are MR effect size +/- the

corresponding standard errors. Red data points indicate significant test results. Ribo: ribosome

occupancy. **(B, C)** Example SCZ risk genes driven by either transcriptional regulation of protein

expression (B) or transcript specific effects (C). Boxplots on the left summarize the normalized

gene expression level stratified by QTL SNP genotypes. The effect size estimates from the

corresponding one-sample MR causality test between each pair of omics are shown in the plot

beneath. Manhattan plots on the right show nominal p values for all variants in the QTL mapping

window for each omics and SCZ GWAS with the red vertical line indicating the top signal

colocalization SNP between QTLs and GWAS. Directly underneath the Manhattan plots are

effect size estimates from the corresponding two-sample MR tests replicating the risk gene

discoveries.