A neuromorphic model of active vision shows how spatiotemporal encoding in lobula neurons can aid pattern recognition in bees

HaDi MaBouDi[1,2,3,4*], Mark Roper[4,5], Marie-Genevieve Guiraud[4,6], Mikko Juusola[2,3], Lars Chittka[4], James A.R. Marshall[1,3]

[1]Department of Computer Science, University of Sheffield, Sheffield, UK
[2]School of Biosciences, University of Sheffield, Sheffield, UK
[3]Neuroscience Institute, University of Sheffield, Sheffield, UK
[4]School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK
[5]Drone Development Lab, Ben Thorns Ltd, Colchester, Essex, UK
[6]School of Natural Sciences, Macquarie University, North Ryde, NSW, Australia

*Corresponding author: HaDi MaBouDi, maboudi@gmail.com

Short title: *Insect-inspired model for active vision*

Abstract: Bees' remarkable visual learning abilities make them ideal for studying active information acquisition and representation. Here, we develop a biologically inspired model to examine how flight behaviours during visual scanning shape neural representation in the insect brain, exploring the interplay between scanning behaviour, neural connectivity, and visual encoding efficiency. Incorporating non-associative learning—adaptive changes without reinforcement—and exposing the model to sequential natural images during scanning, we obtain results that closely match neurobiological observations. Active scanning and non-associative learning dynamically shape neural activity, optimising information flow and representation. Lobula neurons, crucial for visual integration, self-organise into orientation-selective cells with sparse, decorrelated responses to orthogonal bar movements. They encode a range of orientations, biased by input speed and contrast, suggesting co-evolution with scanning behaviour to enhance visual representation and support efficient coding. To assess the significance of this spatiotemporal coding, we extend the model with circuitry analogous to the mushroom body, a region linked to associative learning. The model demonstrates robust performance in pattern recognition, implying a similar encoding mechanism in insects. Integrating behavioural, neurobiological, and computational insights, this study highlights how spatiotemporal coding in the lobula efficiently compresses visual features, offering broader insights into active vision strategies and bio-inspired automation.

Impact statements: Active vision dynamically refines spatiotemporal neural representations, optimising visual processing through scanning behaviour and non-associative learning, providing insights into efficient sensory encoding in dynamic environments.

Keywords: active vision, image statistics, lobula, mushroom bodies, non-associative learning, orientation selective neurons, scanning behaviour, visual recognition.

Introduction

Bees are capable of remarkable cognitive feats, particularly in visual learning (Chittka, 2022; Srinivasan, 2010; Turner, 1911; Von Frisch, 1914; Wehner, 1967). They can not only learn to associate a colour or orientation of a bar with reward (Dyer et al., 2011; Guiraud et al., 2025b; MaBouDi et al., 2020b; Srinivasan, 1994; Stach et al., 2004) but are also able to identify specific features to categorise visual patterns, by finding the relevant stimuli properties (Benard et al., 2006; Guiraud et al., 2025a; Stach et al., 2004). Furthermore, bees have demonstrated the capacity to grasp abstract concepts (Avarguès-Weber et al., 2011; Giurfa et al., 2001; Guiraud et al., 2018; MaBouDi et al., 2020c; Menzel, 2012) and solve numerosity tasks by sequentially scanning the elements within a stimulus (MaBouDi et al., 2020a). These exceptional capabilities position bees as a valuable animal model for investigating the principles of visual learning through the analysis of their behavioural responses (Menzel and Giurfa, 2006; Srinivasan, 2010). Nevertheless, it remains unclear how bees, despite their supposedly low visual acuity (Gribakin, 1975; Spaethe and Chittka, 2003a; Srinivasan and Lehrer, 1988) and limited neural resources, recognise complex patterns and perceive the intricacies of the natural world encountered during foraging (Chittka and Niven, 2009; Giurfa, 2013).

The natural scene that animals encounter is structured differently from random or artificial ones (Clark et al., 2014a; Ruderman and Bialek, 1994; Simoncelli and Olshausen, 2001; Zimmermann et al., 2018). It has been hypothesised that visual sensory neurons evolve to exploit statistical regularities in natural scenes, efficiently encoding information through their spatiotemporal structures (Barlow, 1961). Over evolutionary time, insect visual neurons have developed mechanisms that provide robust and efficient responses to naturalistic inputs (Dyakova et al., 2019, 2015; Dyakova and Nordström, 2017; Juusola et al., 2025; Song and Juusola, 2014; Zheng et al., 2006). For instance, Song and Juusola (2014) showed that fly photoreceptors extract more information from naturalistic time series than from artificial stimuli or white noise, yielding stronger responses with a higher signal-to-noise ratio (Song and Juusola, 2014). Additionally, numerous studies have demonstrated that insect sensory pathways and their associated behaviours dynamically adapt to varying environmental conditions, adjusting their responses based on input parameters such as contrast, spatial frequency, and spatiotemporal correlations (Arenz et al., 2017; Brinkworth and O'Carroll, 2009; Clark et al., 2014b; Dyakova et al., 2019; Dyakova and Nordström, 2017; Juusola et al., 2025; Juusola and Song, 2017; Schwegmann et al., 2014; Serbe et al., 2016; Song and Juusola, 2014, 2014; van Hateren, 1997; van Hateren, 1992). Experience-dependent adaptation has been observed in fly photoreceptors and motion-sensitive neurons in the lobula plate, enabling efficient visual processing under varying conditions. For instance, photoreceptors adapt their response dynamics to different light intensities, optimizing sensitivity to natural stimuli (Juusola and Hardie, 2001a, 2001b). Similarly, motion-sensitive neurons such as the H1 neuron adjust their response properties based on prior motion exposure,

enhancing motion detection in dynamic environments (Maddess et al., 1997). This dynamic plasticity
allows insects to process ecologically relevant information in real time. However, the precise neural
mechanisms underlying natural scene processing remain elusive and require further investigation.
Here we examine how insect visual circuitry has adapted to regularities in natural scenes, focusing
on the efficient coding strategies and robust response mechanisms that enhance visual pattern
recognition.
82
In animal vision, active sampling strategies—wherein animals actively scan their environment to
extract visual information over time—are widely observed across species (Land, 1999; Land and
Nilsson, 2012; Severance and Washburn, 1907; Varella et al., 2024; Washburn, 1926, 1916; Yarbus,
2013). Primates employ eye movements, including saccades and microsaccades, to enhance fine
spatial resolution and improve the encoding of natural stimuli (Anderson et al., 2020; Land, 1999;
Näher et al., 2023; Rucci et al., 2007; Rucci and Victor, 2015). Similarly, insects utilise active vision
strategies, incorporating characteristic head and body movements or specific approach trajectories to
optimise visual processing during behavioural tasks (Bertrand et al., 2021; Chittka and Skorupski,
2017; Dawkins and Woodington, 2000; Egelhaaf et al., 2009; Land, 1973; Land and Nilsson, 2012;
Langridge et al., 2021; MaBouDi et al., 2021b). Recent studies have shown that *Drosophila* generate
photomechanical photoreceptor microsaccades and can move their retinas to stabilise their retinal
images, achieving hyperacute vision and enhancing depth perception (Fenk et al., 2022; Hardie and
Franze, 2012; Juusola et al., 2017; Kemppainen et al., 2022b, 2022a). Likewise, honeybee vision
may require sequential sampling and integration of colour information due to their limited ability to
discriminate between similar hues in brief flashes (<50 ms) (Nityananda et al., 2014). To overcome
this constraint, bees engage in systematic scanning movements, continuously sampling their
surroundings to construct an internal neural representation of their environment (Boeddeker et al.,
2015; Collett et al., 1993; Doussot et al., 2021; Guiraud et al., 2018; Kemppainen et al., 2022a;
Langridge et al., 2018; Lehrer and Collett, 1994; MaBouDi et al., 2020a; Werner et al., 2016). For
instance, bumblebees enumerate visual elements sequentially rather than processing them in
parallel, suggesting a reliance on scanning behaviour for feature extraction parallel (MaBouDi et al.,
2020a), and their flight trajectories further indicate that they prioritise specific pattern regions before
making a decision, rather than processing the entire pattern globally (Langridge et al., 2021; MaBouDi
et al., 2021b). Given the low-resolution nature of compound eyes and the potentially reduced parallel
processing capacity in insects compared to vertebrates, it is likely that bees rely on active vision and
sequential sampling to construct a more robust neural representation of their (Chittka and Skorupski,
2017; Nityananda et al., 2014). These active strategies, akin to primate eye movements, play a crucial
role in early visual processing, redundancy reduction, and efficient encoding of visual stimuli (Doussot
et al., 2021; Kuang et al., 2012; Odenthal et al., 2021). However, it remains poorly understood how
such mechanisms allow bees to overcome representational constraints, detect visual regularities, and

solve complex discrimination tasks. Understanding these strategies is key to uncovering the
fundamental principles of insect vision and their broader implications for visual processing across
biological and artificial systems.

Building on our previous work analysing bee flight paths during a simple visual task (MaBouDi et al.,
2021b), we further investigated the main circuit elements that contribute to active vision in achromatic
pattern recognition, focusing on a simplified yet biologically plausible model. Our primary objective
was to determine how bees' scanning behaviour contributes to the functional organisation and
connectivity of neurons in the visual lobe. We hypothesised that the bees' scanning behaviours have
co-adapted to sample complex visual features in a way that efficiently encodes them into
spatiotemporal patterns of activity in the lobula neurons, facilitating distinct and specific
representations that support learning in the bees' compact brain. To test this, we developed a
neuromorphic model of the bee optic lobes incorporating efficient coding principles via a novel model
of non-associative plasticity. This model demonstrates how spatial scanning behaviour in response
to naturalistic visual inputs has shaped the connectivity within the medulla (the 2^{nd} optic ganglion) to
facilitate an efficient representation of these inputs in the lobula (the 3^{rd} optic ganglion). This efficiency
is achieved through the self-organisation of a specific set of orientation-selective neuron in the lobula,
highlighting the combined impact of scanning behaviour and non-associative learning on shaping the
neural circuitry within the bees' optic lobes.

To evaluate the proposed visual network, we enhance our visual processing framework by
incorporating a secondary decision-making module inspired by insect associative learning
mechanisms, grounded in previous neurobiological evidence (Cassenaer and Laurent, 2012; Fiala
and Kaun, 2024; Fisher et al., 2015; Li et al., 2017; Okada et al., 2007; Paulk et al., 2009; Paulk and
Gronenberg, 2008). Visual input and flight dynamics for the model were derived from our observations
of bee behaviour during a visual discrimination task (MaBouDi et al., 2021b). This allowed us to
evaluate and test the hypothesis of active sampling from our model against real-world behaviour
results (MaBouDi et al., 2021b), as well as other published visual discrimination tasks performed by
bees (Benard et al., 2006; Dyer et al., 2005; Guiraud et al., 2022, 2025a; Srinivasan, 2010, 1994;
Zhang and Horridge, 1992). Furthermore, we conducted a detailed analysis comparing the neural
response features emerging from our model with existing neurobiological findings (James and Osorio,
1996; Paulk et al., 2008; Seelig and Jayaraman, 2013; Yang and Maddess, 1997). This alignment
enhances the credibility of our model in capturing essential aspects of neural processing underlying
active vision.

Results

A bio-inspired neural network for active vision

To investigate how bee scanning behaviour optimises neural activity in the visual lobes and enhances visual information processing for efficient pattern recognition, we developed a neural network model inspired by the key morphological and functional characteristics of the bee brain (Figure 1A, C). The network abstracts the circuitry responsible for the initial processing of visual input in the bee's lamina and medulla (the first and second optic ganglia). To mimic temporal encoding during scans (Figure 1B), we introduced a progressive time delay of 1–5 'temporal instances' between the outputs of medulla neurons and wide-field neurons in the lobula (the third optic ganglion; Figure 1D, see Methods). This temporal structuring facilitates sequential sampling of specific locations along the scan trajectory, gradually integrating visual information into a coherent internal representation that emerges as the final output of the lobula neurons.

Building on findings from bee scanning behaviour (MaBouDi et al., 2021b), the model extracts image input in five sequential patches of 75 × 75 pixels, sampled at a speed of 0.1 m/s, corresponding to a lateral displacement of 15 pixels between consecutive patches (Figure 1B; see Methods for details). The green pixel intensities of each patch modulate the membrane potentials of 5,625 (75 × 75) grid photoreceptors within the simulated bee's single eye. These photoreceptor responses converge onto 625 lamina neurons via recurrent neural connectivity, providing a feedforward mechanism for transferring visual information. The lamina neurons then project to 250 small-field medulla neurons through a simple feedforward pathway (Figure 1C, see Methods).

For each of the five sequential patches that compose a full scan, medulla neuron responses are computed using a spiking neural model. These responses are progressively integrated into the synapses of their corresponding lobula neuron with a structured time delay. As depicted in Figure 1D, the synaptic weights dynamically encode the visual information at different temporal instances (T, 2T, 3T, 4T, 5T), effectively aligning sequentially sampled spatial information into a temporally coherent representation. This ensures that the lobula neuron accumulates and processes the underlying medulla input signals at a synchronised time point, mirroring mechanisms that may occur in biological systems (see Discussion). Additionally, lateral inhibitory connections (red connections in Figure 1C) are proposed between lobula neurons to reduce correlation between them, enhancing redundancy reduction in the process.

It is important to note that this proposed spatiotemporal coding is a simplification. In the bee brain, similar processes are likely mediated through dendritic and synaptic latencies, as well as intermediate neuron transmission within the medulla, influenced by non-associative learning in the visual lobe (Figure 1C, D). We hypothesise that connectivity in the medulla and lobula can be refined through exposure to sequences of time-varying images, incorporating non-associative learning rules and efficient coding principles. These mechanisms are optimised and shaped through a generative

183 learning process to align with the statistical properties of natural scenes, enhancing the system's
184 capacity for processing complex visual inputs (see next section).
185

Figure 1. Neural network of active vision inspired by neurobiology and flight dynamics of bees. (A) The right side displays the front view of the bumblebee head showing the component eye and antenna. Left hand side presents a schematic view of the bee's brain regions. Part of neural pathways from the retina to the mushroom bodies are also represented. Labels: AL – Antennal lobe; LH – Lateral horn; CC – Central complex; La – Lamina; Me – Medulla; Lo – Lobula; MB – Mushroom body. Figure was designed by Alice Bridges **(B)** A representation of the modelled bee's scanning behaviour of a flower demonstrating how a sequence of patches project to the simulated bee's eye with lateral movement from left to right. Below are five image patches sampled by the simulated bee. **(C)** Representation of the neural network model of active vision inspired by micromorphology of the bee brain that underlie learning, memory and experience-

dependent control of behaviour. The photoreceptors located in the eye are excited by the input pattern. The activities of photoreceptors change the membrane potential of a neuron in the next layer, Lamina. The lamina neurons send signals (through *W* connectivity matrix) to the medulla neurons to generate spikes in this layer. Each wide-field lobula neuron integrates the synaptic output of five small-field medulla neurons. The lobula neurons are laterally inhibited by local lobula interconnections (via *Q* connectivity matrix). Lobula neurons project their axons into the mushroom body, forming connections with Kenyon Cells (KCs) through a randomly weighted connectivity matrix, *S*. The KCs all connect to a single mushroom body output neuron (MBON) through random synaptic connections *D*. A single reinforcement neuron (yellow neuron) modulates the synaptic weights between KCs and MBON by simulating the release of octopamine or dopamine when presented with specific visual stimuli (see Method section). **(D)** A temporal coding model that is proposed as the connectivity between medulla and lobula neurons. Each matrix shows the inhibitory (blue) and excitatory (red) connectivity between lamina neurons to a medulla neuron at a given time delay. In this model, the five small-field medulla neurons that are activated by the locally visual input, at different times of scanning, send their activities to a wide-field lobula neuron with a synaptic delay such that the lobula neuron receives all medulla input signals at the same instance (*i.e.* in the presented simulation the lobula neuron is maximally activated by the black vertical bar passing across the visual field from the left to right. Each underlying medulla neuron encodes the vertical bar in a different location of the visual field).

The neural representation of the visual inputs was subsequently transmitted and processed in the mushroom body - the learning centre of the bee brain (Ehmer and Gronenberg, 2002; Li et al., 2017; Paulk et al., 2008; Paulk and Gronenberg, 2008; Schmalz et al., 2022) (Figure 1C). To simplify the model, we incorporated a single mushroom body output neuron (MBON), whose firing rate reflects the simulated bee's preference for a given visual input. By adjusting synaptic weights within the mushroom body, the network was trained to classify visual patterns as either positive (low MBON firing rates) or negative (high MBON firing rates; see Discussion). Following non-associative learning and extensive exposure to natural images, the entire network was trained and tested on various pattern recognition tasks commonly used in experimental studies (Benard et al., 2006; Dyer et al., 2005; Srinivasan, 2010, 1994; Zhang and Horridge, 1992), including the discrimination of "plus" and "multiplication sign" patterns, as previously examined in real bumblebees (MaBouDi et al., 2021b) (Figure 4A).

To evaluate the performance of the active vision model, we analysed MBON activity, which functions as a decision-making unit. A lower MBON response from its baseline activity to a particular pattern indicates preference, whereas a higher response suggests rejection. After multiple training trials through a novel associative learning (see Methods and Discussion), the MBON exhibited a distinct response pattern, with reduced activity towards the chosen visual stimulus and increased activity towards the rejected one. This suggests that the MBON plays a role akin to a decision neuron in pattern recognition tasks. Importantly, no reinforcement learning, or synaptic updates were applied during the testing phase, ensuring that the observed responses reflected the network's learned capacity for visual discrimination rather than online adaptation.

Figure 2. Neural responses of the simulated bee model to visual patterns. (A) Top: each square in the matrix corresponds to a single time slice of the obtained spatiotemporal receptive field of a lobula neuron (5x10 lobula neurons) that emerged from non-associative learning in the visual lobes after exposing the model to images of flowers and nature scenes (see Video 3). Bottom: spatiotemporal receptive field of two example lobula neurons are visualised in the five-time delay slices of the matrices of synaptic connectivity between lamina and five medulla neurons (See Figure 1D). The lobula neuron integrates signals from these medulla neurons at each of five time periods as the simulated bees scan a pattern (time goes from left to right). Blue and red cells show inhibitory and excitatory synaptic connectivity, respectively. The first example lobula neuron (#1) encodes the 150° angled bar moving from lower left to the upper right of the visual field. The second example lobula neuron (#48) encodes the movement of the horizontal bar moving up in the visual field. **(B)** An example of an image sequence projected to the simulated bee' eye with lateral movement from left to right. Below shows the five images patched sampled by the simulated bee. The right side presents the firing rate of all lobula neurons responding to the image sequence. The spatiotemporal receptive field of two highest active neurons to the image sequence are highlighted in purple. **(C)** The polar plot shows the average orientation selectivity of one example lobula neuron (#1) to differently angled bars moving across the visual field in a direction orthogonal to their axis (average of 50 simulations). This neuron is most sensitive to movement when the bar orientation is at 150°. **(D)** The spiking response of the lobula neuron to the preferred orientation raised as the contrast was increased, whereas the response of the lobula neuron to a non-preferred orientation is maintained irrespective of contrast. **(E)** The average velocity-sensitivity curve (±SEM) of the orientation-sensitive lobula neuron (#1) is obtained from the responses of the lobula neuron to optimal (angle of maximum sensitivity) moving stimuli presented to the model at different velocities. The red line shows the Gamma function fitted to the data.

Video 1: Spatiotemporal dynamics of receptive fields in 50 lobula neurons emerging from non-associative learning and active scanning. Each square in the matrix represents a single time slice of the spatiotemporal receptive field for a lobula neuron (5 × 10 array of neurons). These receptive fields illustrate the connectivity matrix between five medulla neurons and their corresponding lobula neuron, operating under a temporal coding structure. In this framework, each of the five medulla neurons sequentially transfers a portion of the visual input to the lobula neuron through excitatory (red) and inhibitory (blue) synaptic connections. These receptive fields develop within the visual lobes after the model is exposed to natural images, including flowers and scenery. As the simulated bee scans a visual pattern, lobula neurons dynamically integrate inputs from medulla neurons over time, forming a temporally structured neural representation of the visual scene.

Non-associative learning shapes spatiotemporal coding in the lobula to align with the
statistical features of natural scenes.

The synaptic weights in the optic lobe were updated through exposure to natural images during the
model's lateral scanning process (see Methods, Figure 1B). While lamina-to-medulla connections are
structured based on temporal coding (Figure 1D), lobula neurons are configured to laterally inhibit
each other, facilitating competitive interactions. Synaptic connections were updated using Oja's
implementation of Hebb's rule (Oja, 1982). Simultaneously, a symmetric inhibitory spike-timing-
dependent plasticity (iSTDP) rule was applied to lateral inhibitory connections among lobula neurons
(Vogels et al., 2011). These local synaptic plasticity rules, which govern interactions between lamina,
medulla, and lobula neurons, support non-associative learning—i.e., synaptic modifications occurring
in the absence of reward (see Methods). Together, these plasticity mechanisms drive the network
toward an efficient representation of visual input, reducing redundancy while preserving essential
visual information.

Figure 2A illustrates the receptive fields of lobula neurons, which exhibit spatiotemporal orientation
selectivity after training on 100 flower and natural images (comprising 50,000 time-varying image
patches). Each square in the figure represents one of the 50 lobula neurons, with the heat map
indicating the synaptic weights of the corresponding lamina neurons (connected via the medulla
neurons). To aid interpretation, the lower portion of Figure 2A provides examples of two individual
lobula neurons, detailing their lamina synaptic weights for each of the five medulla neurons. For a
more dynamic representation of these receptive fields over time, see Video 1. The receptive fields of
lobula neurons are characterised by an elongated "on" area (regions representing positive synaptic
weights) adjacent to an antagonistic "off" area (regions with negative synaptic weights). These regions
are generally aligned along a specific orientation, and their balance changes dynamically across time-
delayed instances of medulla responses. For instance, one lobula neuron responds most strongly to
a 135° bar moving orthogonally to its "on" or "off" areas while exhibiting little or no response to other
orientations (Figure 2C). The population of 50 lobula neurons demonstrates specificity to both
orientation and direction, closely resembling neuronal responses observed in bees and other insects

(James and Osorio, 1996; Paulk et al., 2008; Seelig and Jayaraman, 2013; Yang and Maddess,
1997).

A 100 lobula neurons trained to natural images

B 50 lobula neurons trained to shuffled natural images

Figure 2- figure supplement 1: (A) Spatiotemporal receptive fields of lobula neurons emerging from non-associative learning when the number of lobula neurons is set to 100 (see Video 2). The details follow those described in Figure 2A. **(B)** Spatiotemporal receptive fields of 50 lobula neurons trained with shuffled natural images. The neurons fail to develop meaningful connections, resulting in random synaptic weight distributions, indicating that spatial coherence in training images is essential for efficient feature extraction.

Video 2: Spatiotemporal dynamics of receptive fields in 100 lobula neurons emerging from non-associative learning and active scanning. This follows the same structure as Video 1, depicting the receptive field evolution in a larger population of 100 lobula neurons under the temporal coding framework and non-associative learning.

To illustrate how lobula neurons process natural visual inputs, Figure 2B depicts the sequence of
image patches scanned during a simulated horizontal movement. The results show that only a small
subset of lobula neurons respond at any given moment, indicating that their activity is decorrelated
and relatively selective—an outcome of non-associative learning mechanisms in the visual lobe (see
Discussion). Notably, the two most active lobula neurons captured distinct structural features of the
flower petal: one neuron's receptive field aligned with the left 45° edge of the petal, while another
matched the right-angled edge. This suggests that the model effectively extracts distinct visual
features with a minimal number of filters (lobula neurons).
To examine the selectivity of lobula neurons further, we analysed the spiking activity of a
representative neuron tuned to a 150° orientation. As expected, the neuron showed maximal firing

(26 spikes/sec) when presented with a 150° moving bar. It also exhibited moderate responses to a

horizontal bar and a 120° moving bar (18 spikes/sec) but remained largely unresponsive to other

orientations (Figure 2C). Consistent with experimental findings (Yang and Maddess, 1997), the firing

rate of lobula neurons increased with contrast at their preferred orientations, whereas responses to

non-preferred orientations remained unchanged across contrast levels (Figure 2D).

Finally, Figure 2E highlights the velocity sensitivity of lobula neurons. Each neuron responds

maximally at a specific velocity, demonstrating a tuning curve that aligns with known insect neural

responses (Paulk et al., 2008; Yang and Maddess, 1997). This reinforces that our model successfully

captures key quantitative properties of lobula edge detector neurons, including their joint selectivity

for orientation, contrast, and motion velocity.

The model demonstrates robustness in generating spatiotemporal receptive fields, even as the

number of lobula neurons increases. Training the non-associative learning model with a larger lobula

neuronal population while maintaining the same underlying structure from photoreceptors to the

medulla, enhances the diversity of orientation-selective responses (see Figure 2 -Supplementary

figure 1A and Video 2). As the number of lobula neurons increases, their tuning properties become

more distributed, enabling a finer and more precise encoding of different orientations and motion

patterns. This scalability highlights the model's ability to generalise its representation of natural scene

statistics while achieving varying levels of resolution in visual encoding.

Lobula neuron responses become sparse and decorrelated through non-associative learning

with natural images

To assess the impact of training on the population activity of lobula neurons, we quantified their

response sparsity and decorrelation before and after learning. Figure 3A presents the correlation

matrix of lobula neurons in response to 10,000 sequential scans of natural images after training. The

results reveal a highly decorrelated response pattern, with a strong diagonal structure indicating that

each lobula neuron maintains a distinct response profile. This demonstrates that non-associative

learning enhances both the selectivity and independence of neural representations, allowing the

network to develop more efficient and diverse feature encoding.

Further supporting this, Figure 3B displays the sparseness index of lobula neurons under different

training conditions. Before training, lobula neuron activity was broadly distributed, as reflected in the

high sparseness index (see Methods). Training on natural images significantly reduced the

sparseness index, indicating that lobula neurons developed a more selective and efficient coding

scheme, where only a small subset of neurons responded to any given input. In contrast, training on

shuffled natural images led to only a moderate reduction in sparseness, suggesting that the

implemented local synaptic plasticity rules alone are insufficient for optimal feature encoding without

exposure to structured natural inputs. Notably, while shuffling natural images preserves pixel intensity

and overall distribution, it disrupts spatial correlations and higher-order structures present in natural

scenes. Consequently, training on these shuffled datasets results in non-structured receptive fields,
leading to broadly distributed and less selective coding in the lobula neurons (Figure 2-
Supplementary figure 1B).

Figure 3. Effect of non-associative learning on lobula neuron activity and response sparseness. (A) Correlation matrix of lobula neuron responses after training with natural images. The near-diagonal structure indicates that neurons develop distinct and strongly uncorrelated responses, suggesting an efficient, decorrelated representation of visual input. (B) Sparseness index of lobula neurons before and after training with different image sets. Before training, neural responses are broadly distributed. Training with shuffled natural images does not change the sparseness of lobula population, whereas training with natural images significantly increases response sparseness, indicating that exposure to structured visual inputs enhances efficient coding. Error bars represent SEM. Asterisks (*) indicate p-values < 0.05, while 'n.s.' denotes non-significant results.

These findings underscore the role of non-associative learning in shaping neural representations,
fostering both sparsity and decorrelation in lobula neurons. Such sparse coding is crucial for efficient
sensory processing, as it minimises redundancy while preserving essential visual information.
Moreover, it optimises metabolic efficiency in neural networks, aligning with coding strategies
observed in biological visual systems (see Discussions).

Active vision enhances visual discrimination through sequential scanning
To replicate bee behavioural findings reported in the literature (Benard et al., 2006; Dyer et al., 2005;
MaBouDi et al., 2021b; Srinivasan, 1994, 2010; Zhang and Horridge, 1992), we implemented
computational plasticity within the mushroom body circuitry, leveraging sparse lobula neurons that
emerge through non-associative learning—critical for encoding both appetitive and aversive values.
Specifically, we incorporated classical spike-timing-dependent plasticity (STDP), modulated by
dopamine, to regulate synaptic modifications between mushroom body Kenyon cells (KCs) and
extrinsic MBONs in response to negative (unrewarded) patterns. Additionally, we introduced a novel

STDP-based plasticity rule, modulated by octopamine (See Methods, Figure 9), which we hypothesise
induces synaptic depression among KC-MBON connections in response to positive (rewarded)
patterns. These plasticity mechanisms allowed us to explore synaptic dynamics underlying the
discrimination of rewarded and non-rewarded stimuli (see Discussion).

Figure 4. Simulated bees' performance in a pattern recognition task using different scanning strategies. Twenty simulated bees, with random initial neuronal connectivity in mushroom bodies (see Methods) and a fixed connectivity in the visual lobe that were shaped from the non-associative learning, were trained to discriminate plus from a multiplication symbol (100 random training exposures per pattern). The simulated bees scanned different regions of the patterns at different speeds. **(A)** Top and below panels show the five image patches sampled from the plus and multiplication symbols by simulated bees, respectively. It is assumed that the simulated bees scanned the lower half of the patterns with lateral movement from left to right with normal speed (0.1 m/s). **(B)** The plot shows the average responses of the MBON to rewarding multiplication and punishing plus patterns during training procedure (multiplication symbol rewarding, producing an Octopamine release by the reinforcement neuron, and the plus symbol inducing a Dopamine release). This shows how the response of the MBON to the rewarding plus was decreased while its response to the punishing multiplication pattern was increased during the training. The MBON equally responded to both multiplication and plus before the training (at number of visits =0). **(C)** The performance of the simulated bees in discriminating the right-angled plus and a 45° rotated version of the same cross (i.e. multiplication

symbol) (MaBouDi et al., 2021b; Srinivasan, 1994), when the stimulated bees scanned different regions of the pattern (left corner, lower half, whole pattern) at different speeds: no speed 0.0m/s (i.e. all medulla to lobula temporal slices observed the same visual input), normal speed at 0.1 m/s and fast speed at 0.3m/s), and from a simulated distance of 2cm from stimuli (default) and 10cm (distal view). The optimal model parameters were for the stimulated bees at the default distance when only a local region of the pattern (bottom half or lower left quadrant) was scanned at a normal speed. **(D)** Mean performance (±SEM) of two groups of simulated bees in discriminating the plus from multiplication patterns when their inhibitory connectivity between lobula neurons were not modified by non-associative learning rules. Asterisks (*) indicate p-values < 0.05, while 'n.s.' denotes non-significant results.

The model was trained using a differential conditioning paradigm, where the correct stimulus (S⁺) was

paired with a reward and the incorrect stimulus (S⁻) was associated with punishment. For simplicity,

we denote positive patterns as S⁺ and negative patterns as S⁻. During associative learning

simulations, only the synaptic weights between KCs and MBONs corresponding to the presented

patterns were updated, while randomly weighted connections between the lobula and KCs were

incorporated to ensure sparse activity in KCs (Figure 1C; see Methods). To capture individual

variability observed in bees, we repeated the simulations with different initial conditions, including

random neural connectivity between lobula neurons and Kenyon cells, as well as between KCs and

MBONs. The model's performance was then assessed across multiple visual discrimination

paradigms (Figures 4, 6 and 7).

In the initial implementation, simulated bees were trained to distinguish a plus sign from a

multiplication sign, with training focused on the lower half of the plus pattern (Figure 4A). Following

training, MBON activity decreased in response to the plus (S⁺) while increasing in response to the

multiplication sign (S⁻), whereas prior to training, MBON responses to both patterns were similar

(Figure 4B). This demonstrates that the model successfully discriminated between S⁺ and S⁻ through

temporal coding and sequential scanning of the visual pattern (Figure 4C). In contrast, a model with

fixed, random connectivity in the visual lobe failed to differentiate between the plus and multiplication

patterns (Figure 4D). This underscores the importance of structured connectivity that emerges in the

bee visual lobes through non-associative learning—specifically, the development of spatiotemporal

receptive fields in lobula neurons—for successful visual learning.

Our model further revealed that rewarding patterns elicited a reduction in extrinsic neuron responses,

while punished patterns led to increased responses (Figure 4B), a phenomenon consistent with neural

activity recorded from alpha lobe PE1 neurons in the mushroom body (Okada et al., 2007). Initially,

the simulated bees performed worse than real bees in the plus versus multiplication discrimination

task (Figure 4C, last pair of bars). Using experimentally derived parameters from bumblebee

studies—including an average scanning speed of 0.1 m/s (referred to as normal speed) for whole-

pattern scanning and a viewing distance of 20 mm—simulated bees achieved a correct choice rate

of only 63% for the plus stimulus and 60% for the reciprocal cross-protocol (averaged over 20

simulations; see Figure 4C, fifth pair of bars).

However, when the experimental conditions were adjusted so that simulated bees scanned only the

lower half of the patterns or focused on the lower left corner—consistent with real bee behaviour

(MaBouDi et al., 2021)—correct choice performance improved significantly, reaching ≥96% and

≥98%, respectively (Figure 4C, first two pairs of bars). Conversely, increasing scanning speed

(resulting in larger separations between sampled image patches) reduced accuracy to 70%, while

stationary simulated bees—those that did not actively scan—achieved only 60% correct choices.

In additional experiments where the model bees were trained from a greater distance (>100 mm),

they failed to discriminate between patterns. This aligns with behavioural findings in bees, where they

initially select a pattern at random and move closer to scan it before making a decision (MaBouDi et

al., 2021). Studies show that bumblebees approach stimuli to scan patterns at close range, and their

initial approach choices are random (Guiraud et al., 2018; MaBouDi et al., 2021b). These findings

underscore the critical role of active vision in visual learning and discrimination, demonstrating that

the model effectively captures key aspects of biological visual processing, including sequential

sampling, spatial integration, and plasticity-driven adaptation.

Scanning strategy modulates lobula neural representations: Enhanced selectivity with

localised sampling

To further investigate how scanning behaviour influences the distinctiveness of neural representations

in the lobula and impacts performance in visual learning tasks, we analysed activity patterns and

response magnitudes under two different scanning conditions from the first experiment (Figure 4C).

Heatmaps of lobula neuron activity (Figure 5A, B) illustrate neural responses to two stimulus

conditions: scanning only the lower half of the pattern at normal speed (Panel A) and scanning the

entire pattern at normal speed (Panel B).

The activity maps indicate that different subsets of lobula neurons responded preferentially to the

lower half of the stimulus, exhibiting stronger activation in a smaller neuronal subset. In contrast,

scanning the entire pattern resulted in a more widespread and overlapping activation across the lobula

neuron population. Corresponding spike rate plots (Figure 5, right panels) further highlight these

differences, revealing that neurons displayed stronger selectivity when scanning was restricted to the

lower half at normal speed compared to whole-pattern scanning. These findings suggest that lobula

neurons exhibit stimulus-specific selectivity that is modulated by the spatial dynamics of scanning

behaviour.

To quantify the separability of these population responses, we computed the angular distance (θ)

between their activity vectors over 20 stimulations using the cosine similarity metric (see Methods).

The results (Figure 5C) indicate that neural responses exhibited a significantly larger angular

distance when the model scanned only the lower half of the stimuli at normal speed compared to

scanning the entire of the same pattern. This suggests that neural activity population were more

distinct when scanning was confined to a localised region of these visual patterns, reinforcing the
idea that restricted sampling enhances neural discriminability.

Figure 5. The effect of scanning behaviour on the spatiotemporal encoding of visual patterns in lobula neurons responses. (A, B) Neural responses of simulated lobula neurons to different scanning conditions. Left panels: heatmaps showing the spiking activity of 50 lobula neurons in response to visual patterns when scanning either the lower half of the pattern at normal speed **(A)** or the whole pattern at normal speed **(B)**. Right panels: the mean and standard deviation of the spike rate responses of individual lobula neurons to two distinct visual patterns (plus and multiplication), with colour-coded responses (purple for ⊕, green for ⊗). Scanning behaviour significantly alters the neural responses, with distinct sets of neurons preferentially responding to each stimulus. **(C)** Mean angular distance between lobula neuron responses for different scanning conditions. Lower half-normal speed scanning results in greater separation between neural representations, suggesting that scanning of local region enhances feature selectivity. Error bars represent SEM. Asterisks (*) indicate p-values < 0.05, while 'n.s.' denotes non-significant results.

These findings demonstrate that lobula neurons encode visual stimuli in a structured manner, with

response contrast influenced by both spatial and temporal properties of scanning behaviour

(Figures 4, 5). By selectively sampling specific regions of a stimulus, the system enhances the

differentiation of these visual patterns, supporting the hypothesis that active vision plays a crucial

role in effective neural coding and discrimination.

Neural network model of active vision bee behaviours across various visual experiments

In this study, we evaluated our model - using scans from the lower half of the visual field - by

comparing its performance with results from bee experiments reported in the literature. (Note: bees

may exhibit variations in scanning behaviour under different patterns and training conditions; see

Discussion). Our simulated bees successfully discriminated between angled bars (Hateren et al.,

1990), a 22.5° angled cross from a 90° rotated version (Srinivasan, 1994), and spiral patterns (Zhang

and Horridge, 1992) (Figure 6A). When trained on grating patterns with -45° versus +45° orientations,

the simulated bees successfully identified the correct pattern. Moreover, it demonstrated the ability to

transfer the learned rule to novel patterns the model had never encountered during training, including

single-bar patterns (Figure 6B). This suggests that the model captures key aspects of visual

generalisation observed in real bees. Figure 6C shows that the proposed model not only learned to

identify the correctly oriented bar pattern but also to distinguish the rewarding pattern from a novel

one (two circles). Notably, the model exhibited a 22% lower preference for the negatively trained

pattern compared to the novel pattern, validating the implementation of the rejection behaviour and

demonstrating that the model can simultaneously learn rewarding and aversive stimuli.

This was further explored by training the network with patterns containing two oriented bars in each

lower quadrant (Figure 6D) (Benard et al., 2006; Stach et al., 2004; Zhang and Horridge, 1992). The

simulated bees discriminated these training patterns with over 99% accuracy, but performance

dropped to an average of 61% when presented with a simplified variant. When tested with the original

positive pattern and novel patterns containing only one correct orientation, the bees showed a high

preference for the correct stimulus. Similarly, the simulated bees exhibited a clear preference for a

pattern with a single correct feature over the trained negative pattern, indicating that the model can

extract multiple features during scanning.

Figure 6. Proposed neural network performance to published bee pattern experiments. Twenty simulated bees, with random initial neuronal connectivity in mushroom bodies (see Methods), were trained to discriminate a positive target pattern from a negative distractor pattern (50 training exposures per pattern). The simulated bees' performances were examined via unrewarded tests, where synaptic weights were not updated (average of 20 simulated pattern pair tests per bee). All simulations were conducted under the assumption that model bees viewed the targets from a distance of 2 cm while flying at a normal speed of 0.1 m/s. During this process, the bees scanned the lower half of the pattern. **(A)** Mean percentage of correct choices (±SEM) in discriminating bars oriented at 90° to each other, 25.5° angled cross with a 45° rotated version of the same cross, and a pair of mirrored spiral patterns (MaBouDi et al., 2021b; Srinivasan, 1994). The simulated bees achieved greater than chance performances. **(B)** Performance of

simulated bees trained with a generalisation protocol (Benard et al., 2006). Trained to 6 pairs of perpendicular oriented gratings (10 exposures per grating). Simulated bees then tested with a novel gating pair, and a single oriented bar pair. The simulated bees performed well in distinguishing between the novel pair of gratings; less well, but still significantly above chance, to the single bars. This indicates that the model can generalise the orientation of the training patterns to distinguish the novel patterns. **(C)** Mean performance (±SEM) of the simulated bees in discriminating the positive orientation from negative orientation. Additionally, the performance in recognising the positive orientation from the novel pattern, and preference for the negative pattern from a novel pattern. Simulated bees learnt to prefer positive patterns, but also reject negative patterns, in this case preferring novel stimuli. **(D)** Performance of simulated bees trained to a horizontal and -45° bar in the lower pattern half versus a vertical and +45° bar (Stach et al., 2004). The simulated bees could easily discriminate between the trained bars, and a colour inverted version of the patterns. They performed less well when the bars were replaced with similarly oriented gratings, but still significantly above chance. When tested on the positive pattern vs. a novel pattern with one correctly and one incorrectly oriented bar, the simulated bees chose the positive patterns (fourth and fifth bars), whereas with the negative pattern versus this same novel pattern the simulated bees rejected the negative pattern in preference for the novel pattern with single positive oriented bar (two last bars). **(E)** The graph shows the mean percentage of correct choices for the 20 simulated bees during a facial recognition task (Dyer et al., 2005). Simulated bees were trained to the positive (rewarded) face image versus a negative (non-rewarded) distractor face. The model bee is able to recognise the target face from distractors after training, and also to recognise the positive face from novel faces even if the novel face is similar to the target face (fourth bar). However, it failed to discriminate between the positive and negative faces rotated by 180°. **(F)** The model was trained on spatially structured patterns from Stach et al. (2004), requiring recognition of orientation arrangements across four quadrants. Unlike bees, the model failed to discriminate these patterns, highlighting its limitations in integrating local features into a coherent global representation. Asterisks (*) indicate p-values < 0.05, while 'n.s.' denotes non-significant results.

To present a more complex pattern recognition task, we replicated a facial recognition experiment

performed on honeybees (Dyer et al., 2005) by training the neural network with images of two human

faces (Figure 6E). As with honeybees, our simulated bees were able to identify the positive trained

face from the negative one, as well as distinguish two novel faces and a caricature. Both the real and

simulated bees failed to discriminate the faces when rotated through 180°. These results demonstrate

that complex visual features can be condensed through spatiotemporal encoding in the lobula

neurons into specific and distinct neuronal representations that are critical for learning in the miniature

bee brain.

To evaluate the model's ability to discriminate more complex visual stimuli, we trained it on a set of

patterns from (Stach et al., 2004), which require recognising the spatial arrangement of orientations

across four quadrants (Figure 6F). Unlike previous experiments where scanning behaviour was

confined to specific regions, the complexity of these stimuli necessitated training the model using

whole-pattern exposure. The results indicate that the model failed to replicate the bees' ability to

discriminate these patterns (Figure 6F), despite successfully distinguishing between the plus and

multiplication signs when scanning the entire pattern (Figure 4C), even if its performance remained

lower than that of real bees. Moreover, the model was unable to generalise this capability to more

configurations testing patterns, suggesting that while it can process simple spatial features through

sequential scanning, it lacks the longer and more dynamic scanning required for assembling and

integrating local features into a coherent global representation, as observed in bees (see Discussion).

What is the minimally sufficient number of lobula neurons and the necessary connectivity for

active vision in bees?

As reported above, our model successfully accomplishes various pattern recognition tasks (Figures

4, 6). We then asked whether our neural networks could perform with a very limited number of lobula

neurons that transfer visual information to the mushroom body. To investigate this, we ran the non-

associative learning process with different numbers of lobula neurons – specifically, 4, 16, or 36

neurons (the original model had 50). The visual network was subsequently trained using the same

set of natural images and protocol as the original model (Figure 7A).

Figure 7. Minimum number of lobula neurons that are necessary for pattern recognition.
(A) Obtained spatiotemporal receptive field of lobula neurons when the number of lobula neurons were set at 36, 16 or 4 during the non-associative learning in the visual lobe (See Figure 5A). This shows the models with lower number of lobula neurons encode less variability of orientations and temporal coding of the visual inputs (see Videos 5,6) **(B, C & D)** The average correct choices of the three models with 36, 16 or 4 lobula neurons after training to a pair of plus and multiplication patterns (B) mirrored spiral patterns (C) and human face discrimination (D). The model with 36 lobula neurons still can solve pattern recognition tasks at a level above chance. It indicates that only 36 lobula neurons that provide all inputs to mushroom bodies are sufficient for the simulated bees to be able to discriminate between patterns. Asterisks (*) indicate p-values < 0.05, while 'n.s.' denotes non-significant results.

Video 3: Spatiotemporal dynamics of receptive fields in 36 lobula neurons emerging from non-associative learning and active scanning. This follows the same structure as Video 1, depicting the receptive field evolution in a larger population of 36 lobula neurons under the temporal coding framework and non-associative learning.

Video 4: Spatiotemporal dynamics of receptive fields in 16 lobula neurons emerging from non-associative learning and active scanning. This follows the same structure as Video 1, depicting the receptive field evolution in a larger population of 16 lobula neurons under the temporal coding framework and non-associative learning (compare to Videos 1, 3).

Video 5: Spatiotemporal dynamics of receptive fields in only 4 lobula neurons emerging from non-associative learning and active scanning. This follows the same structure as Video 1, depicting the receptive field evolution in a larger population of 4 lobula neurons under the temporal coding framework and non-associative learning (compare to Videos 1, 3 & 4).

Interestingly, the non-associative learning process led to the emergence of distinct spatiotemporal

structures in the lobula neurons. We found that reducing the number of lobula neurons decreased the

variability in their spatiotemporal receptive (Figure 7A and Videos 3, 4 ,5). In particular, when the

network was limited to four neurons, it could not encode the full spatiotemporal structure of the training

patterns obtained with the model with 50 lobula neurons - only vertical and horizontal receptive fields

were produced (Video 1, 5). As expected, the overall performance of the model decreased as the

number of lobula neurons was reduced. Although the model with 16 lobula neurons demonstrated the

ability to discriminate more complex patterns beyond the plus and multiplication signs (Figure 7B, C),

it remained insufficient for recognising highly complex stimuli such as human faces (Figure 7D, 7E).

Moreover, to investigate the effect of inhibitory neurons within the visual lobe on lobula neuron output,

we trained the model using the same protocol but fixed the synaptic weights of the inhibitory

connections (i.e., these weights were not updated during exposure to the training images). These

fixed inhibitory connections limited the ability of the lobula neuron population to encode moving

orientations (Figure 8 and Video 6), indicating that the plasticity of inhibitory interneurons in the visual

lobe plays a crucial role in facilitating an efficient representation of the visual environment. While this

suggests that increasing the network size can enhance the model's capacity for discriminating

complex visual patterns, the results (Figure 4) indicate that scanning behaviour plays a crucial role in

overcoming this limitation. Specifically, adopting a more targeted scanning strategy can improve

discriminability by directing visual sampling to the most informative regions of the stimulus (see

Discussion).

Taken together, these findings demonstrate that our assumption regarding non-associative plasticity

in the visual lobe successfully replicates the neural responses of lobula neurons across various

patterns and conditions. This plasticity yields a sparse, uncorrelated representation of the visual input

that benefits subsequent learning processes in the mushroom body. Importantly, these results closely

align with theoretical studies (see Discussion), further supporting the effectiveness of the active vision

in capturing the underlying principles of information encoding in the insect visual system.

Figure 8. The role of lateral inhibitory connections between lobula neurons. Obtained spatiotemporal receptive field of lobula neurons (B) when the lateral inhibitory connectivity between lobula neurons is fixed (A) during the non-associative learning (see Video 6 and Figure 2A & Video 1).

Video 6: Spatiotemporal dynamics of receptive fields in 50 lobula neurons emerging from non-associative learning and active scanning. This follows the same structure as Video 1, but with fixed lateral inhibitory connectivity between lobula neurons during non-associative learning. The video illustrates how receptive fields evolve under the temporal coding framework, providing a comparison to Video 1, where lateral inhibition was plastic.

Discussion

In this study, we investigated the core computational requirements for visual pattern recognition by

examined a minimal neural network Inspired by the active scanning flight behaviours of bees

(MaBouDi et al., 2021b). We developed a novel model based on the insect visual system, simulating

how a small population of lobula neurons encodes the visual environment through spatiotemporal

responses. By incorporating non-associative learning, the model self-organises its connectivity within

the visual lobe, generating efficient environmental representations (Figures 2, 3, 5). The process leads

to the emergence of orientation-selective cells in the lobula, which are essential for encoding complex

visual scenes (Figure 2).

Our simulations reveal that a small subset of lobula neurons, sensitive to specific orientations and

velocities, can condense complex visual environments into spatiotemporal representations expressed

as firing rates (Figure 2). These sparse representations effectively discriminate between the plus and
multiplication patterns used in behavioural experiments (MaBouDi et al., 2021b) while also
generalising to novel stimuli—including successful recognition of human faces—highlighting the
model's broader applicability (Figures 4, 6).

Furthermore, our findings highlight the crucial role of bee movement, or active vision, in optimising
the analysis and encoding of environmental information (Figure 4). Spatiotemporal encoding in the
visual lobe emerges as a key mechanism driving the efficiency of minimal intelligent systems. Our
study underscores the fundamental computational principles of visual pattern recognition, particularly
the interplay between active vision and spatiotemporal encoding in insect information processing.
These insights not only advance our understanding of biological vision but also inspire the
development of novel computational models for visual recognition tasks (Figures 3, 4, 5).

The question of how animals cope with a noisy and complex natural world has long been a central
topic in neuroscience and behavioural ecology (Barlow, 1961; Gibson, 1979; Menzel and Giurfa,
2006; Srinivasan, 2010). One key theoretical framework addressing this challenge is the efficient
coding hypothesis, which posits that early sensory systems compress incoming information into a
more efficient format, optimising the transmission of relevant signals to higher brain regions (Barlow,
1961). According to this hypothesis, individual visual neurons should maximise their output capacity
(e.g., reaching their maximum firing rate) when responding to natural stimuli while population
responses should exhibit statistical independence (Simoncelli and Olshausen, 2001). Despite the
relative simplicity of our model compared to recently available full *Drosophila* connectome data (Lin
et al., 2024; Schlegel et al., 2024), our findings suggest that insects optimise visual coding through
non-associative learning while actively exploring their environment (see below). Specifically, we
demonstrate that neural features of the insect brain, combined with active vision, facilitate this
optimisation by developing uncorrelated and sparse coding in the lobula (Figures 2, 3, 5). This
supports the idea that efficient coding is not merely a passive process but one actively shaped by an
animal's interactions with its surroundings. However, bees exhibit a remarkably diverse behavioural
repertoire despite their small brains—ranging from fine-scale object inspections to long-distance
navigation (Chittka, 2022; Juusola et al., 2025; Menzel, 2012; Srinivasan, 2010). This behavioural
diversity makes them an excellent model for investigating how ecological constraints shape neural
computation and, ultimately, efficient coding. Understanding how insects dynamically refine sensory
representations in response to environmental demands offers broader insights into the fundamental
principles of neural information processing in biological systems.

The non-associative model presented in this study operates as a linear generative model that
effectively captures the receptive fields of lobula neurons, linking the spatiotemporal statistics of
natural environments to principles of efficient neural coding (Barlow, 1961; Olshausen, 2003).

Following training, lobula neuron activity in response to naturalistic spatiotemporal signals becomes
highly decorrelated, with only a limited subset of neurons selectively responding to specific visual
stimuli (Figure 5). This sparse coding strategy enhances energy efficiency by minimising overall
neural activity while maintaining distinct stimulus encoding. By ensuring that only a small fraction of
neurons is active at any given moment, this mechanism optimises information processing, reduces
redundancy, and enhances metabolic efficiency, reinforcing the adaptive advantages of efficient
coding in visual systems. Our model introduces a novel generative framework that can be extended
to other species, including primates, to investigate how movement contributes to visual-spatial
encoding in larger brains. Given the ubiquity of active vision across the animal kingdom (Land, 1973;
Land and Nilsson, 2012; Washburn, 1926, 1916; Yarbus, 2013), the principles identified in this study
may be broadly conserved across different taxa, underscoring the fundamental role of sensorimotor
interactions in shaping neural representations. These insights provide a valuable foundation for future
comparative studies on the interplay between movement, efficient coding, and sensory processing
across diverse neural architectures.

Our findings align with previous studies on bumblebees' discrimination of plus and multiplication sign
patterns (MaBouDi et al., 2021b), demonstrating improved model performance when scanning the
lower half of patterns at specific velocities (Figure 3). However, bees exhibit variations in scanning
behaviour depending on pattern complexity and training (Giurfa et al., 1999; Guiraud et al., 2018,
2025b). Research has shown that both honeybees and bumblebees solve visual tasks by extracting
localised or elemental features within patterns, adapting their discrimination strategies accordingly
(Giurfa et al., 1999; MaBouDi et al., 2021b; Stach et al., 2004; Stach and Giurfa, 2005). This suggests
that bees develop tailored flight manoeuvres during training, optimising their scanning behaviour to
maximise visual information extraction. Although our model simplifies visual flight dynamics by
employing a five-step constant-speed horizontal scan (Figure 1B), this abstraction was useful for
isolating key computational principles. However, its failure to solve the complex pattern recognition
task in Figure 6F underscores its limitations, suggesting that incorporating longer and more dynamic
scanning strategies could enhance visual processing capacity. Extending the scanning duration while
integrating multiple visual features across quadrants—alongside mechanisms such as working
memory and sequential learning—could improve performance by enabling the model to retain and
integrate previously acquired visual information. Real-world insect vision, however, relies on more
flexible and adaptive scanning behaviours shaped by flight speed, head movements, and
environmental feedback. Future work should leverage recent advances in insect connectomics, which
reveal a diverse range of neuron types—including small object-detecting neurons, motion-sensitive
neurons, and colour-processing cells—alongside machine learning techniques for analysing animal
movement to develop a more comprehensive flight dynamics model. Incorporating variable scan

trajectories and real-time sensorimotor feedback will offer deeper insights into how active vision
optimises information acquisition and enhances learning in dynamic environments.

A key advantage of our model lies in its ability to leverage sparsity and selectivity to efficiently process
sequential visual data (Figures 2B, 3, 6), in contrast to models that rely on pixel-wise image
representation for training. Traditional models that directly process raw pixel values often require
substantial computational resources and struggle with scalability, particularly when handling large-
scale visual inputs (Amin et al., 2024, 2024; Ardin et al., 2016; Baddeley et al., 2012). In contrast, our
model extracts and encodes sequential visual information within a small population of lobula neurons
(Figures 2B, 6), significantly compressing the visual input while preserving essential features. This
sparse representation reduces redundancy, enhances computational efficiency, and ensures that only
the most informative aspects of the scene are processed for pattern recognition. Furthermore, our
model demonstrates adaptive selectivity, dynamically adjusting to different visual inputs by learning
lobula responses optimised to the statistical features of the scene. Unlike pixel-based models, which
require processing every individual pixel in an image, our approach extracts compact, high-contrast
signals that are more robust to noise and enhance generalisability. This is particularly relevant for bio-
inspired visual processing, as it aligns with known sparse and decorrelated representations in
biological vision systems. By integrating biologically inspired sparse coding, adaptive selectivity, and
motion-driven encoding, our model provides a robust alternative to conventional pixel-based
architectures. This not only improves computational efficiency but also enhances discriminability and
generalisation, making it well-suited for real-world applications, including robotic vision and
autonomous navigation, where rapid adaptation to dynamic environments is crucial.

The results of our model suggest that passive visual exposure to natural images alters the connectivity
in the visual lobes, leading to enhanced pattern recognition abilities (Figures 4, 6). Notably, these
synaptic connections develop independently of the initial connectivity profiles of the simulated bees.
We propose that, beyond the general gross neuroanatomy of the insect optic lobes—which has been
preserved since the Cambrian period for efficient neural representation (Ma et al., 2012)—the specific
visual experiences encountered by real bees during early life play a crucial role in shaping their
individual visual representations. This, in turn, may influence their subsequent performance in
behavioural tasks (Hertel, 1983, 1982; MaBouDi et al., 2017a; Vetter and Visscher, n.d.). There is
direct empirical evidence for such neural developmental processes in olfactory systems of bees,
where early passive exposure improves subsequent odour discrimination (Arenas and Farina, 2008;
Locatelli et al., 2013). Our previous research in olfactory coding demonstrated that the iSTDP learning
rule can establish specific connectivity in the sensory system and enhance the separability of odour
representations in antennal lobe outputs (MaBouDi et al., 2017a). A similar mechanism was observed
here among lobula neurons, where only a limited subset is activated by specific visual inputs, resulting

in sparse and distinct outputs to the mushroom body learning centres (Figures 2B, 5). The receptive

fields of lobula cells, maintained with the fixed lateral connectivity, shows that inhibition is required for

orientation selectivity and temporal coding in the visual lobe (Fisher et al., 2015). These findings

highlight the critical role of inhibitory connections within the visual lobes. Accordingly, our model

predicts that bees with limited early-life visual experiences will perform worse in visual learning and

memory compared to bees with rich visual experiences. Further behavioural and neurobiological

studies are needed to test this prediction.

Mushroom bodies are critical centres for associative learning and memory in insects (Fiala and Kaun,

2024; Heisenberg, 2003; Menzel, 2022, 2012). Synapses between Kenyon Cells and extrinsic

mushroom body neurons follow a Hebbian STDP rule (Aso et al., 2014; Cassenaer and Laurent, 2007;

Markram et al., 1997); however, the STDP rule alone cannot maintain associative learning (Abbott

and Nelson, 2000; Meeks and Holy, 2008). In insects, associative learning appears to rely on the

neurotransmitters octopamine and dopamine to signal unconditioned appetitive and aversive values

(Cognigni et al., 2018; Davidson et al., 2023; Fisher, 2024; Hammer, 1993; Hammer and Menzel,

1995; Matsumoto et al., 2015; Mohammad et al., 2024; Perry and Barron, 2013; Schwaerzel et al.,

2003; Selcho, 2024). These neurotransmitters are released into the mushroom body lobes, where

Kenyon cells connect to mushroom body output neurons (MBON) (Burke et al., 2012; Menzel, 2022;

Okada et al., 2007; Strube-Bloss et al., 2011). Using in vivo electrophysiology in locusts, Cassenaer

and Laurent (2012) reported that octopamine depresses synapses underlying STDP rule, leading to

a lower response in MBONs when octopamine is present.

Following this observation, we modelled associative learning by pairing the positive pattern with the

reward via octopamine-modulated STDP (Equation 4; Figure 1). In this formulation, the temporal

ordering of pre- or postsynaptic spikes depresses the synaptic connection between Kenyon cells and

the MBONs. Conversely, synapses are updated according to classical STDP when negative patterns

are paired with the punishment (Equation 3; Figure 9). This combination produces a complex interplay

between synaptic changes and reinforcer signals, enabling the model to not only learn to select the

positive patterns but to reject incorrect ones (Figure 4B, 6C). The resulting changes in MBON

response to positive patterns during associative learning are consistent with the PE1 extrinsic neuron

in the honeybee brain, which exhibits a lower response to the positive patterns (Ce et al., 2022; Fiala

and Kaun, 2024; Okada et al., 2007) (Figure 4B).

However, further studies are required to investigate the novel combination of octopamine and

dopamine modulation of STDP that is introduced in this study. Combining non-associative learning in

the optic lobes with supervised learning in the mushroom bodies produced a model capable of not

only discriminating simple patterns but also generalisation (Figure 6B), and correct judgments in

conflicting stimulus experiments (Figure 6D). The real power of this approach is exemplified in the
facial recognition task (Figure 6E), where the complexity of the human face is reduced to a set of
sparse lobula neuron activations that can be learnt by the mushroom bodies. Moreover, the
spatiotemporal receptive fields formed during non-associative learning respond differently for different
faces, allowing fine differences to be encoded. Although real bees rarely have to discriminate between
human faces, these processes likely enable bees to select rewarding flowers without requiring a
complex visual memory within their miniature brains.

We used natural scenes with a statistical structure similar to those that visual systems have adapted
to over evolutionary time (Geisler, 2008; Hyvärinen et al., 2009; MaBouDi et al., 2016; Simoncelli and
Olshausen, 2001). Because bee navigation and foraging primarily involve locating food among a
variety of flowers, our non-associative network was trained with a set of different flower images. As
with all theoretical models, this is a simplification, since real bees navigate a 3D environment with a
large field of view. Here, we assume that receptive field formation in real bees is comparable to our
2D simulations. Nevertheless, further studies are necessary to refine and expand our model based
on a more comprehensive understanding of the function and structure of the bee eye components
(Juston et al., 2013; Juusola et al., 2017; Kemppainen et al., 2022b; Viollet and Franceschini, 2010).
Moreover, investigating the neural mechanisms underlying visual learning in the bee brain will allow
us to fine-tune our model's architecture and parameters, leading to a more faithful representation of
the bee visual system.

In this study, we restricted the model's input to green photoreceptors to align with the known visual
processing mechanisms of honeybees. This decision was based on the hypothesis that bee pattern
recognition primarily relies on the green component of visual input, as green-sensitive photoreceptors
are the most abundant, comprising approximately two-thirds of the ommatidia in the compound eye
(Briscoe and Chittka, 2001; Giger and Srinivasan, 1996; Spaethe et al., 2001; Spaethe and Briscoe,
2004). There is also empirical evidence that the green channel provides the predominant input to
movement and edge detection, as well as detailed spatial information for visual discrimination.
Moreover, natural images exhibit a strong correlation among colour channels, meaning that excluding
certain channels does not substantially alter the structure of the visual scene. By focusing on the
green photoreceptor input, our model remains biologically plausible while ensuring computational
efficiency. Future work could explore the contributions of other photoreceptor types to assess their
impact on visual pattern recognition and potential interactions between colour and spatial information
processing in the insect visual system.

Our model provides a functional abstraction of the insect visual system, focusing on core
computational principles rather than replicating the detailed structural connectivity available from

recent connectome studies (Lin et al., 2024; Schlegel et al., 2024). By prioritising the identification of
fundamental mechanisms underlying active vision and visual learning, our approach avoids the
challenges associated with highly parameterised models that can be difficult to interpret
mechanistically. Integrating known physiological properties and behavioural findings from bees, our
model generates testable hypotheses on how motion-driven visual processing enhances pattern
recognition. This functional simplification allows us to isolate key mechanisms that might otherwise
be obscured in large-scale anatomical reconstructions. Additionally, our study emphasises the critical
role of motion in visual recognition—an aspect often overlooked in static-image studies—and
demonstrates how sequential visual input dynamically shapes neural encoding, reinforcing the
importance of active vision in efficient sensory processing.

Recent studies have shown that bees often employ efficient, low-cost strategies to solve cognitive
tasks (Cope et al., 2018; Guiraud et al., 2018; Langridge et al., 2021; MaBouDi et al., 2023a, 2023b,
2021a, 2020b; Roper et al., 2017; Vasas et al., 2019).Understanding these cognitive strategies not
only advances our knowledge of neural computation in miniature brains but also provides a framework
for improving artificial intelligence and autonomous systems. Our study highlights the minimal neural
architectures required for visual learning and lays the foundation for bio-inspired, unsupervised
machine learning algorithms. By emphasising active vision through movement-driven pattern
recognition, our model offers insights into solutions for key AI challenges, such as visual invariance
and robust 3D environmental understanding. Moreover, engineering implementations of eye
micromovements have been shown to enhance edge and bar discrimination, improving the visual
processing efficiency of flying robots (Juston et al., 2013; Viollet and Franceschini, 2010). Additionally,
the non-associative learning model and local plasticity rules explored in this work closely align with
unsupervised learning techniques, particularly sparse coding models, where sparsity constraints
enhance efficiency by reducing redundancy and promoting selective coding. This bio-inspired
framework enables the extraction of latent structures in high-dimensional temporal data, with
applications ranging from sensory signal processing to more adaptive and robust autonomous
perception. Bridging biological and machine intelligence through evolutionarily optimised
computational strategies paves the way for the next generation of AI, driving advancements in
robotics, autonomous navigation, and real-world learning systems (de Croon et al., 2022;
Manoonpong et al., 2021; Serres and Viollet, 2018; Webb, 2020).

Materials and Methods

Network topology of active vision model

The model architecture of the bee visual pathway is illustrated in Figure 1A. The bumblebee has a
pair of compound eyes that are composed of ~5,500 ommatidia (Spaethe and Chittka, 2003b;

Streinzer et al., 2013). Each eye contains three different types of photoreceptors, short, medium and

long wavelength sensitive peaking in the UV, blue and the green respectively (Menzel and Blakers,

1976; Skorupski et al., 2007). Since the green photoreceptors are those that predominantly mediate

visual pattern recognition (Giger and Srinivasan, 1996; Spaethe et al., 2001), we modelled that 75x75

green photoreceptors in one eye component are activated by the pixel values of the input pattern.

Photoreceptors then project to 625 (25x25) neurons in the lamina, which is the first centre of visual

processing. In this model, each lamina neuron, r_l^{La}, receives input from a non-overlapping 3x3 grid of

neighbouring photoreceptors, corresponding to adjust ommatidia. The response of a lamina neuron

is computed as $r_l^{La} = f(\sum_{p=1}^{P=9} r_p \; ; \; A_0, m , b)$. Here, the activation function f is defined as:

$$f(r; A_0, m, b) = A_0 / (1 + exp(mr + b),$$

where r is the input to the lamina neuron La, and A_0 represent the maximum possible activity of

lamina neurons. The parameters m and b define the shape of the activation function, controlling its

steepness and midpoint, respectively. For simplicity, the activation function in our model is fixed with

$A_0 = 1$, $m = -1$, and $b = 0.5$. This function imposes a constraint where weak inputs result in low

activity, while stronger inputs drive the response toward its maximum value in a sigmoidal manner.

Each photoreceptor's output, $r_{p,}$ is derived directly from the pixel intensity at the corresponding location

in the input image, representing the green channel's brightness. The values are normalised between

0 and 1, ensuring a continuous response that reflects the natural variation in luminance.

In this study, each spiking neuron operates according to the integrate-and-fire model. The dynamics

of the subthreshold membrane potential of a neuron, $u(t)$ is described by the following standard

conductance- based leaky integrate-and-fire model: $\tau \frac{du(t)}{dt} = -u(t) + R.I(t)$, where $R = 10$ and $\tau =$

$10ms$ are the resistance and membrane time constant of the neuron respectively. Here, the input $I(t)$

exhibit the total synaptic input to the cell from presynaptic neurons. The membrane potential is reset

to the base activity, $v_0 = -80 \; mV$, if it exceeds the threshold, $V_T = 0 \; mV$.

Each medulla neuron is activated by the summed activity of lamina neurons through the synaptic

connectivity matrix W. The input of the $m-$the medulla neuron, I_m^{Me}, is calculated $I_m^{Me} =$

$\sum_{l=1}^{L} W_{m,l} \; r_l^{La}$. The value $W_{l,m}$ specifies the strength of a synaptic input from the $l-$th lamina neuron

to the $m-$the medulla neuron. To account for the inherent variability in neural responses, we

incorporated stochasticity by adding signal noise generated from a Poisson distribution to the output

of each neuron. The Poisson distribution was chosen because it closely models the statistical

fluctuations observed in biological neural firing, where the variance of the response scales with the

mean activity. This noise was applied independently to each neuron, ensuring that the variability in

responses remains biologically plausible while preserving the overall signal structure. By introducing

this element, the model better reflects the natural dynamics of neural processing, capturing the

probabilistic nature of spike generation and sensory encoding in biological systems.

We propose a temporal coding model that captures the interaction between medulla and lobula

neurons in the visual pathway, incorporating sequential scanning of visual stimuli to optimise

information processing. In this model, each wide-field lobula neuron receives synaptic input from M

small-field medulla neurons, with a structured progressive delay T (Figure 1D). Here, $M = 5$

represents the number of temporal instances within the model's input sequence. Each medulla neuron

is activated by visual information sampled from one of M overlapping segments of an image patch,

determined by the scanning speed, and follows the hierarchical processing pathway from

photoreceptors to lamina neurons (Figure 1C). While individual medulla neurons encode only a

fraction of the visual stimulus, the lobula neuron integrates input from all medulla neurons to generate

spiking activity, thereby forming a holistic representation of the entire visual scene. The synaptic

transmission between medulla and lobula neurons incorporates structured temporal delays at distinct

instances $(T, 2T, 3T, 4T, 5T)$, ensuring that sequentially acquired visual information is temporally

aligned. This results in the synchronized activation of the lobula neuron at a single unified time point,

effectively integrating spatially and temporally structured input into a cohesive internal representation.

By simulating the dynamic interplay between spatial sampling and temporal integration, this model

mirrors the way bees may optimise visual processing through active vision. The resulting alignment

of visual signals enhances feature extraction and pattern recognition, providing a biologically plausible

mechanism for encoding complex scenes efficiently (Figures 2, 3, 6).

The model incorporates lateral inhibitory connections between lobula neurons (Figure 1C, depicted

in red), $Q = [q_{i,j}]$, where $q_{i,j}$ represents the lateral connectivity between $i-$th and $j-$th lobula

neurons. This connectivity along with $W_{m,l}$ are updated during a non-associative learning process, to

reduce redundancy and decorrelate overlapping inputs (see next subsection). This inhibitory

mechanism enhances contrast and improves pattern recognition by selectively amplifying novel

spatial features (Figures 3, 6).

The processed visual information is then transformed to the Kenyon cells (KCs) in the mushroom

body. The synaptic connectivity matrix $S^{Lo \rightarrow KC} = [s_{o,k}]$ determines the excitatory connections

between lobula neurons and KCs in the mushroom body, following previous findings on sparse,

random connectivity (Caron et al., 2013; Szyszka et al., 2005). Kenyon cells exhibit sparse activity,

with fewer than 5% of KCs activated per stimulus, ensuring high selectivity for particular image

features (Honegger et al., 2011). This sparsity emerges naturally from the random connectivity and

thresholder activation dynamics, reinforcing the sparseness in the model. Each connection weight

$s_{o,k}$ is randomly initialized from a uniform distribution in the range $[0, w_{max}]$, where , w_{max} is a scaling

factor ensuring the limitation of input to the KCs. For each simulated bee, the connectivity matrix

$S^{Lo \to KC}$ is randomly reinitialised, ensuring that each instance of the model has a unique but statistically

comparable connectivity structure. This reinitialization reflects individual variations in synaptic wiring

and allows us to assess the robustness of the model's pattern recognition ability across different

randomly generated network configurations.

All Kenyon cells project to a single mushroom body output neuron (MBON), which is the final output

of the model. The input of the of the MBON, I_{MBON}, is computed by the KC-MBON connections D such

that $I_{MBON} = \sum_{k=1}^{K} D_k r_k^{KC}$, where r_k^{KC} is the spiking activity of the k-th KCs. Finally, a reinforcement

neuron makes reinforcement-modulated connections with the KCs and MBON in the presence of the

positive and negative patterns (see the next section).

Each neuron type is denoted with a superscript corresponding to its respective processing stage. For

example, r_p, represents the response of an individual photoreceptor, while r^{La} denotes the activity of

a lamina neuron, where 'La' refers to the lamina layer. Similarly, other superscripts—Me, Lo, KC, and

MBON—correspond to specific network components: the medulla, lobula, Kenyon cells, and

mushroom body output neurons, respectively. This ensures consistency in notation throughout the

model description.

Training the network via a non-associative learning

We trained the model using 50,000 time-varying image patches randomly sampled from a dataset of

100 natural flowers and scene images. During each training step, the model received an input

sequence of five sequential 75 × 75-pixel patches, extracted by shifting 15 pixels across the image

from the left or right or the reverse orientation (Figures 1B, 2B). This patchwise input simulates the

sequential scanning behaviour of bees as they explore visual stimuli.

Using the described network architecture, each time-varying patch dynamically drives spiking activity

in lobula neurons as the simulated movement progresses. At the start of training, all inhibitory

connection strengths Q were initialized with values randomly drawn from a uniform distribution

between 0 and 1. The feedforward synaptic weights W were initialized using Gaussian white noise

$\mathcal{N}(0,1)$. As training progressed, the evoked neural responses of lobula neurons to time-varying

patches were used to iteratively update both the inhibitory weights (Q) and feedforward connections

(W) simultaneously (see Discussion section).

After the image presentation, the feed-forward weight W is updated according to Oja's

implementation of the Hebbian learning rule (MaBouDi et al., 2017a; Oja, 1982) via

802
$$\Delta W_{i,j} = \gamma r_j^{Me}(r_i^{La} - r_j^{Me} W_{i,j}) \qquad \text{(Equation 1)}$$

Here, the r_j^{Me} and r_i^{La} represent the activities of the j-th medulla and i-th lamina neurons, respectively.

The positive constant γ defines the learning rate.

At the same time of processing, the lateral inhibitory connectivity in the lobula is modified by inhibitory

spike-time-dependent plasticity (iSTDP) (Vogels et al., 2011). Here, we model *non-associative*

learning in lobula by a symmetric iSTDP between presynaptic of the inhibitory neurons and

postsynaptic lobula neurons. In this learning rule, both temporal ordering of pre- or post- synaptic

spikes potentiates the connectivity and the synaptic strength of j-th inhibitory neuron onto i-th lobula

neuron $(Q_{i,j})$ is updated as follows:

$$\Delta Q_{i,j} = \eta(r_i^{Lo} * r_j^{In} - \alpha) \qquad \text{(Equation 2)}$$

where r_i^{Lo} and r_j^{In} exhibit the mean firing rate of the lobula and inhibitory neurons, respectively. The

depression factor α controls the target activity rate of the lobula neurons. Here, η is the learning rate.

To simplify, a one-to-one connection between the inhibitory and lobula neuron is assumed in the

model such that the activity of the j-th inhibitory neuron is equal to the activity of the j-th lobula neuron.

The training is terminated when the synaptic weights over time are changed less than a small

threshold (0.001). In the training process, synaptic weights were constrained within the range $[-1, +1]$

for the W and $[0, +1]$ for Q to ensure stable convergence and to reflect biological limitations in

synaptic transmission strength.

Associative learning in Mushroom Bodies

To verify if the lobula neurons can reproduce empirical behavioural results in different visual tasks,

the model is enriched with associative learning process in the mushroom bodies (a bio-inspired

supervised learning). When the training process of the non-associative learning is terminated, we use

a reward-based synaptic wright modification rule in KCs-MBON connection (D), such that, if a stimulus

is rewarding (*i.e.* positive), the corresponding synapses between activated neurons will be weakened

while for a stimulus paired with punishment (*i.e.* negative), activated synapses are strengthened

(Cassenaer and Laurent, 2012) (see Discussion section). The model behaves as the activity of

mushroom body neurons in decreasing their firing rate in responding to the positive stimuli during

training (Okada et al., 2007). In this model, two reinforcement neurons modulated strengths of

synaptic connectivity at the output of the KCs in response to both reward and punishment. In the

presence of the negative patterns, the synaptic strengths from the KCs to the MBON are modified,

and modulated by dopamine, based on the classical STDP (Song et al., 2000; Zhang et al., 1998)

(Figure S3A):

$$STDP_{Dop}(\Delta t) = \begin{cases} A\,e^{-\Delta t/\tau}, & \Delta t > 0 \\ -A\,e^{\Delta t/\tau}, & \Delta t < 0 \end{cases}, \qquad \text{(Equation 3)}$$

where $\Delta t = t_{post} - t_{pre}$ implies the difference between the spike time of pre- and post- synaptic

neurons. Further, applying the synaptic plasticity rule modulated by octopamine (octopamine

modulated STDP) observed in the presence of rewarding stimuli to the synapses between KCs and

MBON (Cassenaer and Laurent, 2012), the change in synaptic weight can be summarised as (Figure

4B):

840
$$STDP_{OCT}(\Delta t) = \begin{cases} -A\, e^{-\Delta t/\tau}, & \Delta t > 0 \\ -A\, e^{\Delta t/\tau}, & \Delta t < 0 \end{cases}, \qquad \text{(Equation 4)}.$$

841 Here, $A = 0.01$ and $\tau = 20\, ms$ exhibit the maximum magnitude and time constant of the STDP

842 function for the synaptic potentiation or depression.

Figure 9. STPS curves. (A) Classical STDP curve showing relationship between synaptic weight change and the precise time difference between the Kenyon Cells and MBON spikes. The synaptic weight can be either depressed or potentiated. **(B)** STDP curve modulated by octopamine in the insect mushroom body. The Synaptic weights are depressed. The formula of these curves was described in Equations 3 and 4.

To train the model in different conditions of scanning, the flight-scan forms of the positive and negative

patterns were presented to the model. Each set of flight-scan input contained a set of five patches

with size 75x75 pixels were selected from the test patterns by shifting 15 pixels over each pattern

from the left to right (Figure 4A). The numbers of shifted pixels control the speed of scanning. The

activity of the MBON was used to assess the performance of the model. Following the training, the

performance of the model was calculated from a decrease in firing rate of the MBON to a pattern that

had been rewarding and/or an increase in firing rate of MBON to a pattern that had been punishing in

training. The bee's final behavioural decision is proposed to come from a simple integration of these

different valence-encoding neurons.

Quantifying neural population sparseness and response separability

To quantify population sparseness in the lobula following training on natural images, we employed

the Treves-Rolls sparseness index (MaBouDi et al., 2017b; Willmore and Tolhurst, 2001), defined as:

$$SI = \frac{\left(\sum_{j=1}^{N} r_j/N\right)^2}{\left(\sum_{j=1}^{N} r_j^2\right)/N}$$

where r_j represents the firing rate of the j-th lobula neuron, and N is the total number of lobula

neurons.

This metric provides insight into the population coding strategy of lobula neurons, distinguishing

between broad distributed representations and sparse selective responses:

• *Maximum SI = 1 (low sparseness):* Achieved when all neurons respond equally, indicating a

fully distributed code where the entire population is uniformly active across all stimuli.

• *Minimum SI = 1/N (high sparseness):* Occurs when only a single neuron is active while all

others remain silent, reflecting a highly selective encoding scheme.

To further assess the separability between response population of lobula neurons in response to

different stimuli, we computed the angular distance (θ) using the cosine similarity formula: $\theta =$

$\cos^{-1}(R_1.R_2/(|R_1||R_1|))$, where R_1 and R_2 represent the activity vectors of two neural populations,

and $|R_1|$ and $|R_2|$ denote their responses Euclidean norms. This measure captures the geometric

distinction between response patterns, with larger angles indicating greater separability between

neural representations of different stimuli. A higher angular distance suggests that the population

responses are more distinct, reflecting improved stimulus discrimination within the lobula.

Simulation and Statistical Analysis

To assess the model's performance across experiments, we conducted 20 independent simulations

for each condition, ensuring statistical reliability and robustness. In each simulation, the synaptic

connectivity matrix between lobula neurons and Kenyon cells was randomly initialised using a uniform

distribution within a biologically plausible range, mimicking the individual variability observed in real

bees. This stochastic initialisation prevented bias in learning outcomes and allowed us to examine

how the model generalises across different neural configurations.

Each simulated bee underwent multiple training exposures to visual stimuli. Following training, once

the model parameters were fixed at the final stage of training, the model was tested across 50

repetitions per condition with testing patterns to account for variability in responses. The model's

performance was evaluated by averaging across simulations, with standard error of the mean (SEM)

reported in the figures to provide a statistically robust representation of discrimination accuracy.

Statistical analyses were conducted to compare pattern discrimination across conditions. To maintain

clarity and focus on the modelling findings, detailed significance values are not reported in the main

text. Instead, figures indicate p-values < 0.05 with '*' and non-significant results with 'n.s.'. Since data

distributions did not always meet normality assumptions, we used the Wilcoxon signed-rank test for

matched data and the Wilcoxon rank-sum test (Mann-Whitney U test) for independent samples. For

comparisons across multiple groups, we applied the Kruskal-Wallis test, followed by Dunn's post-hoc

test when necessary.

Computing Environment: All modelling and visualisation were performed using MATLAB

(RRID:SCR_001622) and Python (RRID:SCR_008394). MATLAB was also used for statistical

analysis.

898 Acknowledgements

We thank Paul Graham and Andrew Barron for valuable comments on manuscript and Alice Bridges

for drawing the front view of the bumblebee presented in Figure 1A. This research was financed by

HFSP programme grant RGP0022/2014, EPSRC program grant Brains-on-Board EP/Poo6094/1 and

by Horizon Europe Framework Programme grant NimbleAI. MJ was supported by BBSRC

(BB/F012071/1 and BB/X006247/1), EPSRC (EP/X019705/1) and Leverhulme (RPG-2024-016).

MGG was supported by ARC Discovery Projects DP230100006 and DP210100740 and Templeton

World Charity Foundation Project Grant TWCF-2020-20539.

907 Conflict of interest statement

908 All authors declare no conflict of interest.

910 Data and code accessibility

911 The code developed for this research project has been made openly accessible on GitHub:

912 https://github.com/hadiimaboudi/neuromorphic_model_active-vision-

914 Reference

Abbott LF, Nelson SB. 2000. Synaptic plasticity: taming the beast. *Nature neuroscience* **3**:1178–
1183.
Amin AA, Kagioulis E, Domcsek N, Nowotny T, Graham P, Philippides A. 2024. Investigating the
limits of familiarity-based navigation. *Artificial Life* 1–17.
Anderson AG, Ratnam K, Roorda A, Olshausen BA. 2020. High-acuity vision from retinal image
motion. *Journal of vision* **20**:34–34.
Ardin P, Peng F, Mangan M, Lagogiannis K, Webb B. 2016. Using an insect mushroom body circuit
to encode route memory in complex natural environments. *PLoS computational biology*
12:e1004683.
Arenas A, Farina WM. 2008. Age and rearing environment interact in the retention of early olfactory
memories in honeybees. *J Comp Physiol A* **194**:629–640. doi:10.1007/s00359-008-0337-z

Arenz A, Drews MS, Richter FG, Ammer G, Borst A. 2017. The temporal tuning of the *Drosophila*
motion detectors is determined by the dynamics of their input elements. *Current Biology* **27**:929–
944.
Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo T-T, Dionne H, Abbott LF, Axel R, Tanimoto H.
2014. The neuronal architecture of the mushroom body provides a logic for associative learning.
elife **3**:e04577.
Avarguès-Weber A, Dyer AG, Giurfa M. 2011. Conceptualization of above and below relationships
by an insect. *Proceedings of the Royal Society of London B: Biological Sciences* **278**:898–905.
doi:10.1098/rspb.2010.1891
Baddeley B, Graham P, Husbands P, Philippides A. 2012. A model of ant route navigation driven by
scene familiarity. *PLoS computational biology* **8**:e1002336.
Barlow HB. 1961. Possible principles underlying the transformation of sensory messages. *Sensory*
communication **1**.
Benard J, Stach S, Giurfa M. 2006. Categorization of visual stimuli in the honeybee *Apis mellifera*.
Anim Cogn **9**:257–270. doi:10.1007/s10071-006-0032-9
Bertrand OJ, Doussot C, Siesenop T, Ravi S, Egelhaaf M. 2021. Visual and movement memories
steer foraging bumblebees along habitual routes. *Journal of Experimental Biology* **224**:jeb237867.
Boeddeker N, Mertes M, Dittmar L, Egelhaaf M. 2015. Bumblebee Homing: The Fine Structure of
Head Turning Movements. *PLOS ONE* **10**:e0135020. doi:10.1371/journal.pone.0135020
Brinkworth RS, O'Carroll DC. 2009. Robust models for optic flow coding in natural scenes inspired
by insect biology. *PLoS computational biology* **5**:e1000555.
Briscoe AD, Chittka L. 2001. THE EVOLUTION OF COLOR VISION IN INSECTS. *Annual Review of*
Entomology **46**:471–510. doi:10.1146/annurev.ento.46.1.471
Burke CJ, Huetteroth W, Owald D, Perisse E, Krashes MJ, Das G, Gohl D, Silies M, Certel S,
Waddell S. 2012. Layered reward signalling through octopamine and dopamine in *Drosophila*.
Nature **492**:433–437.
Caron SJC, Ruta V, Abbott LF, Axel R. 2013. Random convergence of olfactory inputs in the
Drosophila mushroom body. *Nature* **497**:113–117. doi:10.1038/nature12063
Cassenaer S, Laurent G. 2012. Conditional modulation of spike-timing-dependent plasticity for
olfactory learning. *Nature* **482**:47–52. doi:10.1038/nature10776
Cassenaer S, Laurent G. 2007. Hebbian STDP in mushroom bodies facilitates the synchronous flow
of olfactory information in locusts. *Nature* **448**:709–713. doi:10.1038/nature05973
Ce H, V R, Ey R, Sh D, Mp N, A F. 2022. Visualization of learning-induced synaptic plasticity in
output neurons of the *Drosophila* mushroom body γ-lobe. *Scientific reports* **12**. doi:10.1038/s41598-
022-14413-5
Chittka L. 2022. The Mind of a Bee. Princeton University Press. doi:10.1515/9780691236247
Chittka L, Niven J. 2009. Are bigger brains better? *Curr Biol* **19**:R995–R1008.
doi:10.1016/j.cub.2009.08.023
Chittka L, Skorupski P. 2017. Active vision: a broader comparative perspective is needed.
Clark DA, Fitzgerald JE, Ales JM, Gohl DM, Silies MA, Norcia AM, Clandinin TR. 2014a. Flies and
humans share a motion estimation strategy that exploits natural scene statistics. *Nature*
neuroscience **17**:296–303.
Clark DA, Fitzgerald JE, Ales JM, Gohl DM, Silies MA, Norcia AM, Clandinin TR. 2014b. Flies and
humans share a motion estimation strategy that exploits natural scene statistics. *Nat Neurosci*
17:296–303. doi:10.1038/nn.3600
Cognigni P, Felsenberg J, Waddell S. 2018. Do the right thing: neural network mechanisms of
memory formation, expression and update in *Drosophila*. *Current Opinion in Neurobiology* **49**:51–
58. doi:10.1016/j.conb.2017.12.002
Collett TS, Fry SN, Wehner R. 1993. Sequence learning by honeybees. *J Comp Physiol A* **172**:693–
706. doi:10.1007/BF00195395
Cope AJ, Vasilaki E, Minors D, Sabo C, Marshall JAR, Barron AB. 2018. Abstract concept learning
in a simple neural network inspired by the insect brain. *PLOS Computational Biology* **14**:e1006435.
doi:10.1371/journal.pcbi.1006435

Davidson AM, Kaushik S, Hige T. 2023. Dopamine-Dependent Plasticity Is Heterogeneously
Expressed by Presynaptic Calcium Activity across Individual Boutons of the *Drosophila* Mushroom
Body. *eNeuro* **10**. doi:10.1523/ENEURO.0275-23.2023
Dawkins MS, Woodington A. 2000. Pattern recognition and active vision in chickens. *Nature*
403:652–655. doi:10.1038/35001064
de Croon GCHE, Dupeyroux JJG, Fuller SB, Marshall JAR. 2022. Insect-inspired AI for autonomous
robots. *Science Robotics* **7**:eabl6334. doi:10.1126/scirobotics.abl6334
Doussot C, Bertrand OJN, Egelhaaf M. 2021. The Critical Role of Head Movements for Spatial
Representation During Bumblebees Learning Flight. *Frontiers in Behavioral Neuroscience* **14**.
Dyakova O, Lee Y-J, Longden KD, Kiselev VG, Nordström K. 2015. A higher order visual neuron
tuned to the spatial amplitude spectra of natural scenes. *Nature Communications* **6**:8522.
doi:10.1038/ncomms9522
Dyakova O, Müller MM, Egelhaaf M, Nordström K. 2019. Image statistics of the environment
surrounding freely behaving hoverflies. *Journal of Comparative Physiology A* **205**:373–385.
Dyakova O, Nordström K. 2017. Image statistics and their processing in insect vision. *Current*
Opinion in Insect Science, Neuroscience * Pheromones **24**:7–14. doi:10.1016/j.cois.2017.08.002
Dyer AG, Neumeyer C, Chittka L. 2005. Honeybee (*Apis mellifera*) vision can discriminate between
and recognise images of human faces. *Journal of Experimental Biology* **208**:4709–4714.
Dyer AG, Paulk AC, Reser DH. 2011. Colour processing in complex environments: insights from the
visual system of bees. *Proceedings of the Royal Society of London B: Biological Sciences* **278**:952–
959. doi:10.1098/rspb.2010.2412
Egelhaaf M, Kern R, Lindemann JP, Braun E, Geurten B. 2009. Active Vision in Blowflies:
Strategies and Mechanisms of Spatial Orientation In: Floreano D, Zufferey J-C, Srinivasan MV,
Ellington C, editors. Flying Insects and Robots. Berlin, Heidelberg: Springer Berlin Heidelberg. pp.
51–61. doi:10.1007/978-3-540-89393-6_4
Ehmer B, Gronenberg W. 2002. Segregation of visual input to the mushroom bodies in the
honeybee (*Apis mellifera*). *J of Comparative Neurology* **451**:362–373. doi:10.1002/cne.10355
Fenk LM, Avritzer SC, Weisman JL, Nair A, Randt LD, Mohren TL, Siwanowicz I, Maimon G. 2022.
Muscles that move the retina augment compound eye vision in *Drosophila*. *Nature* **612**:116–122.
doi:10.1038/s41586-022-05317-5
Fiala A, Kaun KR. 2024. What do the mushroom bodies do for the insect brain? Twenty-five years
of progress. *Learn Mem* **31**:a053827. doi:10.1101/lm.053827.123
Fisher YE. 2024. Octopamine enhances learning. *National Science Review* **11**:nwae185.
doi:10.1093/nsr/nwae185
Fisher YE, Silies M, Clandinin TR. 2015. Orientation Selectivity Sharpens Motion Detection in
Drosophila. *Neuron* **88**:390–402. doi:10.1016/j.neuron.2015.09.033
Geisler WS. 2008. Visual Perception and the Statistical Properties of Natural Scenes. *Annual*
Review of Psychology **59**:167–192. doi:10.1146/annurev.psych.58.110405.085632
Gibson JJ. 1979. The ecological approach to visual perception.
Giger AD, Srinivasan MV. 1996. Pattern recognition in honeybees: chromatic properties of
orientation analysis. *J Comp Physiol A* **178**:763–769. doi:10.1007/BF00225824
Giurfa M. 2013. Cognition with few neurons: higher-order learning in insects. *Trends in*
Neurosciences **36**:285–294. doi:10.1016/j.tins.2012.12.011
Giurfa M, Hammer M, Stach S, Stollhoff N, MüLler-Deisig N, Mizyrycki C. 1999. Pattern learning by
honeybees: conditioning procedure and recognition strategy. *Animal Behaviour* **57**:315–324.
doi:10.1006/anbe.1998.0957
Giurfa M, Zhang S, Jenett A, Menzel R, Srinivasan MV. 2001. The concepts of 'sameness' and
'difference' in an insect. *Nature* **410**:930–933. doi:10.1038/35073582
Gribakin FG. 1975. Functional morphology of the compound eye of the bee. *The compound eye and*
vision of insects **154**:176.
Guiraud M, Roper M, Chittka L. 2018. High-Speed Videography Reveals How Honeybees Can Turn
a Spatial Concept Learning Task Into a Simple Discrimination Task by Stereotyped Flight
Movements and Sequential Inspection of Pattern Elements. *Front Psychol* **9**.
doi:10.3389/fpsyg.2018.01347

Guiraud M, Roper M, Wolf S, Woodgate JL, Chittka L. 2022. Discrimination of edge orientation by
bumblebees. *Plos one* **17**:e0263198.
Guiraud M-G, Gallo V, Quinsal-Keel E, MaBouDi H. 2025a. Bumble bee visual learning: simple
solutions for complex stimuli. *Animal Behaviour* **221**:123070.
Guiraud M-G, MaBouDi H, Woodgate J, Bates OK, Rodriguez OR, Gallo V, Barron AB. 2025b. How
bumblebees manage conflicting information seen on arrival and departure from flowers. *Anim Cogn*
28:11. doi:10.1007/s10071-024-01926-x
Hammer M. 1993. An identified neuron mediates the unconditioned stimulus in associative olfactory
learning in honeybees. *Nature* **366**:59–63. doi:10.1038/366059a0
Hammer M, Menzel R. 1995. Learning and memory in the honeybee. *J Neurosci* **15**:1617–1630.
Hardie RC, Franze K. 2012. Photomechanical Responses in *Drosophila* Photoreceptors. *Science*
338:260–263. doi:10.1126/science.1222376
Hateren J van, Srinivasan MV, Wait PB. 1990. Pattern recognition in bees: orientation
discrimination. *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and*
Behavioral Physiology **167**:649–654.
Heisenberg M. 2003. Mushroom body memoir: from maps to models. *Nat Rev Neurosci* **4**:266–275.
doi:10.1038/nrn1074
Hertel H. 1983. Change of synapse frequency in certain photoreceptors of the honeybee after
chromatic deprivation. *Journal of comparative physiology* **151**:477–482.
Hertel H. 1982. The effect of spectral light deprivation on the spectral sensitivity of the honey bee.
Journal of comparative physiology **147**:365–369.
Hyvärinen A, Hurri J, Hoyer PO. 2009. Natural image statistics: A probabilistic approach to early
computational vision. Springer Science & Business Media.
James AC, Osorio D. 1996. Characterisation of columnar neurons and visual signal processing in
the medulla of the locust optic lobe by system identification techniques. *Journal of Comparative*
Physiology A **178**:183–199. doi:10.1007/BF00188161
Juston R, Kerhuel L, Franceschini N, Viollet S. 2013. Hyperacute edge and bar detection in a
bioinspired optical position sensing device. *IEEE/ASME Transactions on Mechatronics* **19**:1025–
1034.
Juusola M, Dau A, Song Z, Solanki N, Rien D, Jaciuch D, Dongre SA, Blanchard F, Polavieja GG
de, Hardie RC, Takalo J. 2017. Microsaccadic sampling of moving image information provides
Drosophila hyperacute vision. *eLife Sciences* **6**:e26117. doi:10.7554/eLife.26117
Juusola M, Hardie RC. 2001a. Light adaptation in *Drosophila* photoreceptors: I. Response dynamics
and signaling efficiency at 25 C. *The Journal of general physiology* **117**:3–25.
Juusola M, Hardie RC. 2001b. Light adaptation in *Drosophila* photoreceptors: II. Rising temperature
increases the bandwidth of reliable signaling. *The Journal of general physiology* **117**:27–42.
Juusola M, Song Z. 2017. How a fly photoreceptor samples light information in time. *The Journal of*
Physiology **595**:5427–5437.
Juusola M, Takalo J, Kemppainen J, Haghighi KR, Scales B, McManus J, Bridges A, MaBouDi H,
Chittka L. 2025. Theory of morphodynamic information processing: linking sensing to behaviour.
Vision Research **227**:108537.
Kemppainen J, Mansour N, Takalo J, Juusola M. 2022a. High-speed imaging of light-induced
photoreceptor microsaccades in compound eyes. *Communications Biology* **5**:203.
Kemppainen J, Scales B, Razban Haghighi K, Takalo J, Mansour N, McManus J, Leko G, Saari P,
Hurcomb J, Antohi A, Suuronen J-P, Blanchard F, Hardie RC, Song Z, Hampton M, Eckermann M,
Westermeier F, Frohn J, Hoekstra H, Lee C-H, Huttula M, Mokso R, Juusola M. 2022b. Binocular
mirror–symmetric microsaccadic sampling enables *Drosophila* hyperacute 3D vision. *Proc Natl Acad*
Sci USA **119**:e2109717119. doi:10.1073/pnas.2109717119
Kuang X, Poletti M, Victor JD, Rucci M. 2012. Temporal encoding of spatial information during
active visual fixation. *Curr Biol* **22**:510–514. doi:10.1016/j.cub.2012.01.050
Land MF. 1999. Motion and vision: why animals move their eyes. *J Comp Physiol A* **185**:341–352.
Land MF. 1973. Head Movement of Flies during Visually Guided Flight. *Nature* **243**:299–300.
doi:10.1038/243299a0
Land MF, Nilsson D-E. 2012. Animal eyes. Oxford University Press.

Langridge KV, Wilke C, Riabinina O, Vorobyev M, de Ibarra NH. 2018. Approach direction prior to
landing explains patterns of colour learning. *bioRxiv* 381210.
Langridge KV, Wilke C, Riabinina O, Vorobyev M, Hempel de Ibarra N. 2021. Approach Direction
Prior to Landing Explains Patterns of Colour Learning in Bees. *Frontiers in Physiology* **12**.
Lehrer M, Collett TS. 1994. Approaching and departing bees learn different cues to the distance of a
landmark. *J Comp Physiol A* **175**:171–177. doi:10.1007/BF00215113
Li L, MaBouDi H, Egertová M, Elphick MR, Chittka L, Perry CJ. 2017. A possible structural correlate
of learning performance on a colour discrimination task in the brain of the bumblebee. *Proc R Soc B*
284:20171323. doi:10.1098/rspb.2017.1323
Lin A, Yang R, Dorkenwald S, Matsliah A, Sterling AR, Schlegel P, Yu S, McKellar CE, Costa M,
Eichler K. 2024. Network statistics of the whole-brain connectome of *Drosophila*. *Nature* **634**:153–
165.
Locatelli FF, Fernandez PC, Villareal F, Muezzinoglu K, Huerta R, Galizia CG, Smith BH. 2013.
Nonassociative plasticity alters competitive interactions among mixture components in early
olfactory processing. *Eur J Neurosci* **37**:63–79. doi:10.1111/ejn.12021
Ma X, Hou X, Edgecombe GD, Strausfeld NJ. 2012. Complex brain and optic lobes in an early
Cambrian arthropod. *Nature* **490**:258–261. doi:10.1038/nature11495
MaBouDi H, Barron AB, Li S, Honkanen M, Loukola OJ, Peng F, Li W, Marshall JAR, Cope A,
Vasilaki E, Solvi C. 2021a. Non-numerical strategies used by bees to solve numerical cognition
tasks. *Proceedings of the Royal Society B: Biological Sciences* **288**:20202711.
doi:10.1098/rspb.2020.2711
MaBouDi H, Dona H, Gatto E, Loukola OJ, Buckley E, Onoufriou PD, Skorupski P, Chittka L. 2020a.
Bumblebees use sequential scanning of countable items in visual patterns to solve numerosity
tasks. *Integrative and Comparative Biology*.
MaBouDi H, Marshall JA, Dearden N, Barron AB. 2023b. How honey bees make fast and accurate
decisions. *eLife* **12**:e86176.
MaBouDi H, Marshall JAR, Barron AB. 2020b. Honeybees solve a multi-comparison ranking task by
probability matching. *Proceedings of the Royal Society B: Biological Sciences* **287**:20201525.
doi:10.1098/rspb.2020.1525
MaBouDi H, Roper M, Guiraud M, Marshall JAR, Chittka L. 2021b. Automated video tracking and
flight analysis show how bumblebees solve a pattern discrimination task using active vision. *bioRxiv*
2021.03.09.434580. doi:10.1101/2021.03.09.434580
MaBouDi H, Shimazaki H, Amari S, Soltanian-Zadeh H. 2016. Representation of higher-order
statistical structures in natural scenes via spatial phase distributions. *Vision research* **120**:61–73.
MaBouDi H, Shimazaki H, Giurfa M, Chittka L. 2017a. Olfactory learning without the mushroom
bodies: Spiking neural network models of the honeybee lateral antennal lobe tract reveal its
capacities in odour memory tasks of varied complexities. *PLoS computational biology* **13**:e1005551.
MaBouDi H, Shimazaki H, Giurfa M, Chittka L. 2017b. Olfactory learning without the mushroom
bodies: spiking neural network models of the honeybee lateral antennal lobe tract reveal its
capacities in odour memory tasks of varied complexities. *PLoS computational biology* **13**:e1005551.
MaBouDi H, Solvi C, Chittka L. 2020c. Bumblebees learn a relational rule but switch to a win-
stay/lose-switch heuristic after extensive training. *bioRxiv* 2020.05.08.085142.
doi:10.1101/2020.05.08.085142
Maddess T, Laughlin SB, Horridge GA, Levick WR. 1997. Adaptation of the motion-sensitive neuron
H1 is generated locally and governed by contrast frequency. *Proceedings of the Royal Society of
London Series B Biological Sciences* **225**:251–275. doi:10.1098/rspb.1985.0061
Manoonpong P, Patanè L, Xiong X, Brodoline I, Dupeyroux J, Viollet S, Arena P, Serres JR. 2021.
Insect-inspired robots: bridging biological and artificial systems. *Sensors* **21**:7609.
Markram H, Lübke J, Frotscher M, Sakmann B. 1997. Regulation of synaptic efficacy by
coincidence of postsynaptic APs and EPSPs. *Science* **275**:213–215.
Matsumoto Y, Matsumoto C-S, Wakuda R, Ichihara S, Mizunami M. 2015. Roles of octopamine and
dopamine in appetitive and aversive memory acquisition studied in olfactory conditioning of
maxillary palpi extension response in crickets. *Front Behav Neurosci* **9**.
doi:10.3389/fnbeh.2015.00230

Meeks JP, Holy TE. 2008. Pavlov's moth: olfactory learning and spike timing–dependent plasticity. *Nature neuroscience* **11**:1126–1127.

Menzel R. 2022. In Search for the Retrievable Memory Trace in an Insect Brain. *Front Syst Neurosci* **16**. doi:10.3389/fnsys.2022.876376

Menzel R. 2012. The honeybee as a model for understanding the basis of cognition. *Nat Rev Neurosci* **13**:758–768. doi:10.1038/nrn3357

Menzel R, Blakers M. 1976. Colour receptors in the bee eye — Morphology and spectral sensitivity. *J Comp Physiol* **108**:11–13. doi:10.1007/BF00625437

Menzel R, Giurfa M. 2006. Dimensions of cognition in an insect, the honeybee. *Behav Cogn Neurosci Rev* **5**:24–40. doi:10.1177/1534582306289522

Mohammad F, Mai Y, Ho J, Zhang X, Ott S, Stewart JC, Claridge-Chang A. 2024. Dopamine neurons that inform *Drosophila* olfactory memory have distinct, acute functions driving attraction and aversion. *PLOS Biology* **22**:e3002843. doi:10.1371/journal.pbio.3002843

Näher T, Zhang Y, Pandinelli M, Fries P. 2023. Primate Saccade Rhythmicity. *bioRxiv* 2023–09.

Nityananda V, Skorupski P, Chittka L. 2014. Can bees see at a glance? *J Exp Biol* **217**:1933–1939. doi:10.1242/jeb.101394

Odenthal L, Doussot C, Meyer S, Bertrand OJN. 2021. Analysing Head-Thorax Choreography During Free-Flights in Bumblebees. *Front Behav Neurosci* **14**. doi:10.3389/fnbeh.2020.610029

Oja E. 1982. Simplified neuron model as a principal component analyzer. *Journal of mathematical biology* **15**:267–273.

Okada R, Rybak J, Manz G, Menzel R. 2007. Learning-Related Plasticity in PE1 and Other Mushroom Body-Extrinsic Neurons in the Honeybee Brain. *J Neurosci* **27**:11736–11747. doi:10.1523/JNEUROSCI.2216-07.2007

Olshausen BA. 2003. Principles of image representation in visual cortex. *The visual neurosciences* **2**:1603–1615.

Paulk AC, Dacks AM, Phillips-Portillo J, Fellous J-M, Gronenberg W. 2009. Visual Processing in the Central Bee Brain. *J Neurosci* **29**:9987–9999. doi:10.1523/JNEUROSCI.1325-09.2009

Paulk AC, Gronenberg W. 2008. Higher order visual input to the mushroom bodies in the bee, Bombus impatiens. *Arthropod Structure & Development* **37**:443–458. doi:10.1016/j.asd.2008.03.002

Paulk AC, Phillips-Portillo J, Dacks AM, Fellous J-M, Gronenberg W. 2008. The Processing of Color, Motion, and Stimulus Timing Are Anatomically Segregated in the Bumblebee Brain. *J Neurosci* **28**:6319–6332. doi:10.1523/JNEUROSCI.1196-08.2008

Perry CJ, Barron AB. 2013. Neural mechanisms of reward in insects. *Annual review of entomology* **58**:543–562.

Roper M, Fernando C, Chittka L. 2017. Insect Bio-inspired Neural Network Provides New Evidence on How Simple Feature Detectors Can Enable Complex Visual Generalization and Stimulus Location Invariance in the Miniature Brain of Honeybees. *PLOS Computational Biology* **13**:e1005333. doi:10.1371/journal.pcbi.1005333

Rucci M, Iovin R, Poletti M, Santini F. 2007. Miniature eye movements enhance fine spatial detail. *Nature* **447**:852–855.

Rucci M, Victor JD. 2015. The unsteady eye: an information-processing stage, not a bug. *Trends Neurosci* **38**:195–206. doi:10.1016/j.tins.2015.01.005

Ruderman DL, Bialek W. 1994. Statistics of natural images: Scaling in the woods. *Phys Rev Lett* **73**:814–817. doi:10.1103/PhysRevLett.73.814

Schlegel P, Yin Y, Bates AS, Dorkenwald S, Eichler K, Brooks P, Han DS, Gkantia M, Dos Santos M, Munnelly EJ. 2024. Whole-brain annotation and multi-connectome cell typing of *Drosophila*. *Nature* **634**:139–152.

Schmalz F, El Jundi B, Rössler W, Strube-Bloss M. 2022. Categorizing visual information in subpopulations of honeybee mushroom body output neurons. *Frontiers in Physiology* **13**:866807.

Schwaerzel M, Monastirioti M, Scholz H, Friggi-Grelin F, Birman S, Heisenberg M. 2003. Dopamine and octopamine differentiate between aversive and appetitive olfactory memories in *Drosophila*. *J Neurosci* **23**:10495–10502.

Schwegmann A, Lindemann JP, Egelhaaf M. 2014. Temporal statistics of natural image sequences generated by movements with insect flight characteristics. *PLoS ONE* **9**:e110386. doi:10.1371/journal.pone.0110386

Seelig JD, Jayaraman V. 2013. Feature detection and orientation tuning in the *Drosophila* central
complex. *Nature* **503**:262–266. doi:10.1038/nature12601
Selcho M. 2024. Octopamine in the mushroom body circuitry for learning and memory. *Learn Mem*
31:a053839. doi:10.1101/lm.053839.123
Serbe E, Meier M, Leonhardt A, Borst A. 2016. Comprehensive characterization of the major
presynaptic elements to the *Drosophila* OFF motion detector. *Neuron* **89**:829–841.
Serres JR, Viollet S. 2018. Insect-inspired vision for autonomous vehicles. *Current opinion in insect*
science **30**:46–51.
Severance E, Washburn MF. 1907. The loss of associative power in words after long fixation. *The*
American Journal of Psychology 182–186.
Simoncelli EP, Olshausen BA. 2001. Natural image statistics and neural representation. *Annual*
review of neuroscience **24**:1193–1216.
Skorupski P, Döring TF, Chittka L. 2007. Photoreceptor spectral sensitivity in island and mainland
populations of the bumblebee, Bombus terrestris. *Journal of Comparative Physiology A* **193**:485–
494.
Song S, Miller KD, Abbott LF. 2000. Competitive Hebbian learning through spike-timing-dependent
synaptic plasticity. *Nature neuroscience* **3**:919–926.
Song Z, Juusola M. 2014. Refractory Sampling Links Efficiency and Costs of Sensory Encoding to
Stimulus Statistics. *Journal of Neuroscience* **34**:7216–7237. doi:10.1523/JNEUROSCI.4463-
13.2014
Spaethe J, Briscoe AD. 2004. Early duplication and functional diversification of the opsin gene
family in insects. *Molecular Biology and Evolution* **21**:1583–1594.
Spaethe J, Chittka L. 2003a. Interindividual variation of eye optics and single object resolution in
bumblebees. *Journal of Experimental Biology* **206**:3447–3453.
Spaethe J, Chittka L. 2003b. Interindividual variation of eye optics and single object resolution in
bumblebees. *Journal of Experimental Biology* **206**:3447–3453.
Spaethe J, Tautz J, Chittka L. 2001. Visual constraints in foraging bumblebees: flower size and
color affect search time and flight behavior. *Proceedings of the National Academy of Sciences*
98:3898–3903.
Srinivasan MV. 2010. Honey bees as a model for vision, perception, and cognition. *Annual review of*
entomology **55**:267–284.
Srinivasan MV. 1994. Pattern recognition in the honeybee: recent progress. *Journal of Insect*
Physiology **40**:183–194.
Srinivasan MV, Lehrer M. 1988. Spatial acuity of honeybee vision and its spectral properties. *J*
Comp Physiol **162**:159–172. doi:10.1007/BF00606081
Stach S, Benard J, Giurfa M. 2004. Local-feature assembling in visual pattern recognition and
generalization in honeybees. *Nature* **429**:758–761. doi:10.1038/nature02594
Stach S, Giurfa M. 2005. The influence of training length on generalization of visual feature
assemblies in honeybees. *Behavioural Brain Research* **161**:8–17. doi:10.1016/j.bbr.2005.02.008
Streinzer M, Brockmann A, Nagaraja N, Spaethe J. 2013. Sex and Caste-Specific Variation in
Compound Eye Morphology of Five Honeybee Species. *PLOS ONE* **8**:e57702.
doi:10.1371/journal.pone.0057702
Strube-Bloss MF, Nawrot MP, Menzel R. 2011. Mushroom Body Output Neurons Encode Odor–
Reward Associations. *J Neurosci* **31**:3129–3140. doi:10.1523/JNEUROSCI.2583-10.2011
Szyszka P, Ditzen M, Galkin A, Galizia CG, Menzel R. 2005. Sparsening and temporal sharpening
of olfactory representations in the honeybee mushroom bodies. *J Neurophysiol* **94**:3303–3313.
doi:10.1152/jn.00397.2005
Turner CH. 1911. Experiments on Pattern-Vision of the Honey Bee. *Biological Bulletin* **21**:249–264.
doi:10.2307/1536017
Van Hateren JH. 1997. Processing of natural time series of intensities by the visual system of the
blowfly. *Vision research* **37**:3407–3416.
van Hateren JH. 1992. Theoretical predictions of spatiotemporal receptive fields of fly LMCs, and
experimental validation. *Journal of Comparative Physiology A* **171**:157–170.
Varella TT, Takahashi DY, Ghazanfar AA. 2024. Active sampling as an information seeking strategy
in primate vocal interactions. *Communications Biology* **7**:1098.

Vasas V, Peng F, MaBouDi H, Chittka L. 2019. Randomly weighted receptor inputs can explain the
large diversity of colour-coding neurons in the bee visual system. *Scientific reports* **9**:1–13.
Vetter RS, Visscher PK. n.d. Influence of Age on Antennal Response of Male Honey Bees, *Apis*
mellifera, to Queen Mandibular Pheromone and Alarm Pheromone Component. *J Chem Ecol*
23:1867–1880. doi:10.1023/B:JOEC.0000006456.90528.94
Viollet S, Franceschini N. 2010. A hyperacute optical position sensor based on biomimetic retinal
micro-scanning. *Sensors and Actuators A: Physical* **160**:60–68.
Vogels TP, Sprekeler H, Zenke F, Clopath C, Gerstner W. 2011. Inhibitory Plasticity Balances
Excitation and Inhibition in Sensory Pathways and Memory Networks. *Science* **334**:1569–1573.
doi:10.1126/science.1211095
Von Frisch K. 1914. Der farbensinn und formensinn der biene. Рипол Классик.
Washburn MF. 1926. The animal mind: A text-book of comparative psychology. Macmillan.
Washburn MF. 1916. Movement and mental imagery: Outlines of a motor theory of the complexer
mental processes. Houghton Mifflin.
Webb B. 2020. Robots with insect brains. *Science* **368**:244–245.
Wehner R. 1967. Pattern recognition in bees. *Nature* **215**:1244–1248.
Werner A, Stürzl W, Zanker J. 2016. Object Recognition in Flight: How Do Bees Distinguish
between 3D Shapes? *PLoS ONE* **11**:e0147106. doi:10.1371/journal.pone.0147106
Willmore B, Tolhurst DJ. 2001. Characterizing the sparsenessof neural codes. *Network:*
Computation in Neural Systems **12**:255.
Yang E-C, Maddess T. 1997. Orientation-sensitive Neurons in the Brain of the Honey Bee (*Apis*
mellifera). *Journal of Insect Physiology* **43**:329–336. doi:10.1016/S0022-1910(96)00111-4
Yarbus AL. 2013. Eye movements and vision. Springer.
Zhang LI, Tao HW, Holt CE, Harris WA, Poo M. 1998. A critical window for cooperation and
competition among developing retinotectal synapses. *Nature* **395**:37–44.
Zhang SW, Horridge GA. 1992. Pattern recognition in bees: size of regions in spatial layout.
Philosophical Transactions of the Royal Society of London Series B: Biological Sciences **337**:65–
71.
Zheng L, de Polavieja GG, Wolfram V, Asyali MH, Hardie RC, Juusola M. 2006. Feedback Network
Controls Photoreceptor Output at the Layer of First Visual Synapses in *Drosophila*. *Journal of*
General Physiology **127**:495–510. doi:10.1085/jgp.200509470
Zimmermann MJ, Nevala NE, Yoshimatsu T, Osorio D, Nilsson D-E, Berens P, Baden T. 2018.
Zebrafish differentially process color across visual space to match natural scenes. *Current Biology*
28.
1285